

# LOG KODAIRA DIMENSION OF HOMOGENEOUS VARIETIES

MICHEL BRION AND DE-QI ZHANG

ABSTRACT. Let  $V$  be a complex algebraic variety, homogeneous under the action of a complex algebraic group. We show that the log Kodaira dimension of  $V$  is non-negative if and only if  $V$  is a semi-abelian variety.

## 1. INTRODUCTION

Throughout this short note, we work over the field of complex numbers. Given a smooth quasi-projective variety  $V$ , we recall the definition of the log Kodaira dimension  $\bar{\kappa}(V)$ . Take any projective variety  $X$  which contains  $V$  as a dense open subset. By blowing up subvarieties of the (reduced) boundary  $D := X \setminus V$ , we may assume that  $X$  is smooth and  $D$  is a simple normal crossing divisor. In this case, the pair  $(X, D)$  is called a *smooth projective compactification of  $V$* . Let  $K_X$  be a canonical divisor of  $X$ . The *log Kodaira dimension*  $\bar{\kappa} = \bar{\kappa}(X, K_X + D)$  is defined as the unique value

$$\bar{\kappa} \in \{-\infty, 0, 1, \dots, \dim V\}$$

such that for some positive constants  $\alpha, \beta$ , we have

$$\alpha m^{\bar{\kappa}} \leq h^0(X, m(K_X + D)) \leq \beta m^{\bar{\kappa}}$$

for any sufficiently large and divisible positive integer  $m$ . As  $\bar{\kappa}$  is independent of the choice of the smooth projective compactification  $(X, D)$  (cf. [4, §11.1]), we may set  $\bar{\kappa} := \bar{\kappa}(V)$ .

Also, recall that a Cartier divisor  $D$  on a projective variety  $X$  is *pseudo-effective* if its numerical equivalence class is contained in the closure of the convex cone spanned by the effective divisor classes on  $X$ .

We may now state the main result of this note:

**Theorem 1.1.** *Let  $X$  be a smooth projective variety, and  $G$  a connected algebraic group (possibly nonlinear) of automorphisms of  $X$ . Assume that  $G$  has a dense open orbit  $V$  in  $X$ , and  $D := X \setminus V$  is a simple normal crossing divisor. Then the following conditions are equivalent.*

- (1) *The log canonical divisor  $K_X + D$  is pseudo-effective.*

---

2010 *Mathematics Subject Classification.* 14J50, 14L10, 14M17, 32M12.

*Key words and phrases.* log Kodaira dimension, homogeneous variety, semi-abelian variety.

The second-named author is supported by an ARF of NUS.

- (2)  $K_X + D$  is linearly equivalent to 0.
- (3) The log Kodaira dimension  $\bar{\kappa}(V)$  is non-negative.
- (4)  $G$  is a semi-abelian variety.

We recall that a *semi-abelian variety* is an algebraic group  $G$  that lies in an exact sequence

$$1 \longrightarrow T \longrightarrow G \longrightarrow A \longrightarrow 1,$$

where  $T$  is an algebraic torus (isomorphic to a product of copies of the multiplicative group  $\mathbb{G}_m$ ), and  $A$  an abelian variety; then  $G$  is connected and commutative.

Here are a few words about the proof of Theorem 1.1, which is very simple. If  $V$  is not a semi-abelian variety, we show that the general member of some covering family of affine lines on  $G$  is not contractible to a point in  $V$ . Chevalley's structure theorem for algebraic groups is also used, so our proof is quite geometric.

From Theorem 1.1, we derive the following purely group-theoretic statement:

**Corollary 1.2.** *Let  $G$  be a connected algebraic group, and  $H \leq G$  a closed subgroup. Then  $\bar{\kappa}(G/H) \leq 0$ , with equality if and only if  $H$  contains a closed normal subgroup  $N$  of  $G$  such that  $G/N$  is a semi-abelian variety.*

## 2. PROOFS OF THEOREM 1.1 AND COROLLARY 1.2

The following result is well-known. See [5, Lemma 5.11] for a generalization to singular pairs.

**Lemma 2.1.** *Let  $Y$  be a smooth projective variety and  $A$  a simple normal crossing divisor on  $Y$ . Suppose that  $C$  is the general member of a covering family of rational curves on  $Y$ . Let  $d$  be the number of times that  $C$  meets  $A$ , i.e., the degree of the (reduced) support of the divisor  $\nu^*(A|_C)$ , where  $\nu : \mathbb{P}^1 \rightarrow C$  is the normalization. Then we have:*

- (1) If  $d \leq 2$ , then  $(K_Y + A) \cdot C \leq 0$ .
- (2) If  $d \leq 1$ , then  $(K_Y + A) \cdot C < 0$ . In particular  $K_Y + A$  is not pseudo-effective.

We now prove Theorem 1.1.

(2)  $\Rightarrow$  (3) and (3)  $\Rightarrow$  (1) follow readily from the definitions of the log Kodaira dimension and pseudo-effectivity.

(4)  $\Rightarrow$  (2). Since the semi-abelian variety  $G \subseteq \text{Aut}(X)$  is commutative and acts faithfully on  $X$ , the stabilizer of any point of  $V$  is trivial, and hence  $V \cong G$ . Also, the action of the Lie algebra of  $G$  on  $X$  yields a trivialization of the log tangent bundle  $T_X(-\log D)$  (see [8, Main Thm.] and [2, Thm. 2.5.1]). In particular, the determinant of  $T_X(-\log D)$  is trivial as well, i.e.,  $K_X + D \sim 0$ .

(1)  $\Rightarrow$  (4). Fix a point  $x_0 \in V$  and denote by  $H \leq G$  its isotropy subgroup, so that  $V$  is  $G$ -equivariantly isomorphic to the homogeneous space  $G/H$ , with both the  $G$ -actions from the left. Since  $G$  acts faithfully on  $V$ , we have:

**Claim 2.2.**  *$H$  contains no non-trivial closed normal subgroup of  $G$ .*

Next we show:

**Claim 2.3.**  *$H$  contains the image of every additive one-parameter subgroup of  $G$ .*

We prove Claim 2.3. Assume the contrary that there exists an additive one-parameter subgroup of  $G$ , i.e., a homomorphism of algebraic groups  $u : \mathbb{G}_a \rightarrow G$  (where  $\mathbb{G}_a$  denotes the additive group), with image not contained in  $H$ . Then the  $\mathbb{G}_a$ -orbit  $u(\mathbb{G}_a) \cdot x_0$  is a curve and isomorphic to  $\mathbb{G}_a/F$ , where the isotropy subgroup  $F$  is a finite subgroup of  $\mathbb{G}_a$  and hence is trivial. Thus we obtain an embedding

$$f : \mathbb{A}^1 \cong \mathbb{G}_a \cong u(\mathbb{G}_a) \cdot x_0 \hookrightarrow G/H,$$

$$t \longmapsto u(t) \cdot x$$

and hence a rational curve  $C \subset X$  which intersects the boundary  $D$  at at most one point. Also, the translates of  $C$  by  $G$  form a covering family of rational curves on  $X$ . By Lemma 2.1, it follows that  $K_X + D$  is not pseudo-effective, contradicting our assumption. This proves Claim 2.3.

We return to the proof of Theorem 1.1. By Chevalley's structure theorem (see [7, Thm. 16]),  $G$  lies in a unique extension

$$1 \longrightarrow L \longrightarrow G \longrightarrow A \longrightarrow 1,$$

where  $L$  is a connected linear algebraic group, and  $A$  an abelian variety. Consider the unipotent radical  $R_u(L)$ ; this is a closed connected normal subgroup of  $G$ . If  $R_u(L)$  is non-trivial, consider the last non-trivial term of its lower central series,  $U$ . Then  $U$  is a closed connected normal subgroup of  $G$ , isomorphic to the additive group of a finite-dimensional vector space. Hence,  $U$  is not contained in  $H$  by Claim 2.2; but then  $G$  has an additive one-parameter subgroup with image not contained in  $H$ , contradicting Claim 2.3. So  $R_u(L)$  is trivial, i.e.,  $L$  is reductive. Thus, the derived subgroup  $[L, L]$  is semi-simple, and hence generated by images of additive one-parameter subgroups. So  $[L, L] \subseteq H$  by Claim 2.3. As  $[L, L]$  is a normal subgroup of  $G$ , Claim 2.2 implies that  $[L, L]$  is trivial. So  $L$  is an algebraic torus. Hence  $G$  is a semi-abelian variety.

This completes the proof of Theorem 1.1.

Next we prove Corollary 1.2.

Denote by  $N$  the kernel of the action of  $G$  on  $G/H$ , i.e., the largest (closed) normal subgroup of  $G$  contained in  $H$ . Replacing  $G, H$  with  $G/N, H/N$ , respectively, we may assume that  $G$  acts faithfully on  $G/H$ .

By [3, Thm. 2], there exists a projective compactification  $X$  on  $G/H$  such that the natural  $G$ -action on  $G/H$  from the left extends to an algebraic action on  $X$ . Using equivariant resolution of singularities (see [6]), we may assume that  $X$  is smooth and  $X \setminus V$  is a simple normal crossing divisor. Then the desired assertion follows from Theorem 1.1. This proves Corollary 1.2.

### 3. CONCLUDING REMARKS

3.1. With the assumptions of Theorem 1.1, the action of the Lie algebra of  $G$  on  $X$  preserves  $D$ , and hence yields a linear map

$$\mathrm{op}_{X,D} : \mathrm{Lie}(G) \longrightarrow H^0(X, T_X(-\log D)),$$

where  $T_X(-\log D)$  denotes the log tangent bundle. Moreover, the pull-back of  $T_X(-\log D)$  to  $V$  is just the tangent bundle  $T_V$ , which is generated by the image of  $\mathrm{op}_{X,D}$ . It follows that the induced map

$$\wedge^n \mathrm{Lie}(G) \longrightarrow H^0(X, -(K_X + D))$$

is nonzero, where  $n := \dim(V) = \dim(X)$ . In particular,  $-(K_X + D)$  is effective. Thus, the pseudo-effectivity of  $K_X + D$  in Theorem 1.1 (1) immediately implies Theorem 1.1 (2):  $K_X + D \sim 0$ .

3.2. Theorem 1.1 is applicable to an arbitrary (not necessarily smooth) equivariant compactification  $Y \supseteq V$  as long as  $Y$  is normal, the boundary  $E := Y \setminus V$  is a divisor, and  $K_Y + E$  is  $\mathbb{Q}$ -Cartier. Indeed, by equivariant resolution of singularities, there exists a  $G$ -equivariant birational morphism  $f : X \rightarrow Y$  such that  $X$  is smooth, projective and equipped with a  $G$ -action, and  $D := X \setminus V$  (which is the inverse of  $E$ ) is a simple normal crossing divisor. By the logarithmic ramification divisor formula, we have

$$K_X + D = f^*(K_Y + E) + R_f$$

where  $R_f$  is an effective  $f$ -exceptional divisor. Thus the pairs  $(X, D)$  and  $(Y, E)$  (as compactifications of  $V$ ) have the same log Kodaira dimensions:  $\kappa(X, K_X + D) = \kappa(Y, K_Y + E)$ , and the log canonical divisor  $K_X + D$  is pseudo-effective if and only if so is  $K_Y + E$ .

In particular, if  $K_Y + E$  is pseudo-effective, then so is  $K_X + D$ , and hence both  $V \subseteq X$  and  $V \subseteq Y$  are the embeddings of the semi-abelian variety  $V$  in  $X$  and  $Y$ , respectively.

3.3. The normal projective equivariant compactifications of a given semi-abelian variety are well-understood by work of Alexeev (see [1]), where they are called semi-abelic varieties. In particular, for a semi-abelian variety  $G$  given by an extension

$$1 \rightarrow T \rightarrow G \rightarrow A \rightarrow 1$$

where  $T$  is an algebraic torus and  $A$  an abelian variety, and a normal projective equivariant compactification  $Y$  of  $G$ , the morphism  $q : G \rightarrow A$  extends to a  $T$ -invariant morphism

$$f : Y \longrightarrow A,$$

which is a fibration in projective toric varieties with the big torus  $T$ . Thus,  $f$  is the Albanese map of the variety  $Y$ .

#### REFERENCES

- [1] V. Alexeev, *Complete moduli in the presence of semiabelian group actions*, Ann. of Math. (2) **155** (2002), no. 3, 611–708.
- [2] M. Brion, *Log homogeneous varieties*, Proceedings of the XVIth Latin American Algebra Colloquium, 1–39, Bibl. Rev. Mat. Iberoamericana, Madrid, 2007.
- [3] M. Brion, *Some basic results on actions of nonaffine algebraic groups*, Symmetry and spaces, 1–20, Progr. Math. **278**, Birkhäuser, Boston, 2010.
- [4] S. Iitaka, *Algebraic Geometry – an Introduction to Birational Geometry of Algebraic Varieties*, Grad. Texts in Math. **76**, Springer 1981.
- [5] S. Keel and J. M<sup>c</sup>Kernan, *Rational curves on quasi-projective surfaces*, Mem. Amer. Math. Soc. **140** (1999), no. **669**.
- [6] J. Kollár, *Lectures on resolution of singularities*, Ann. of Math. Stud. **166**, Princeton Univ. Press, 2007.
- [7] M. Rosenlicht, *Some basic theorems on algebraic groups*, Amer. J. Math. **78** (1956), 401–443.
- [8] J. Winkelmann, *On manifolds with trivial logarithmic tangent bundle*, Osaka J. Math. **41** (2004), no. 2, 473–484.

INSTITUT FOURIER, UNIVERSITÉ DE GRENOBLE,  
 B.P. 74, 38402 SAINT-MARTIN D'HÈRES, FRANCE  
*E-mail address:* Michel.Brion@ujf-grenoble.fr

DEPARTMENT OF MATHEMATICS, NATIONAL UNIVERSITY OF SINGAPORE,  
 10 LOWER KENT RIDGE ROAD, SINGAPORE 119076  
*E-mail address:* matzdz@nus.edu.sg