

NEWTON–HOOKE/CARROLLIAN EXPANSIONS OF (A)dS AND CHERN–SIMONS GRAVITY

Joaquim Gomis¹, Axel Kleinschmidt^{2,3}, Jakob Palmkvist⁴, Patricio Salgado-Rebolledo⁵

¹*Departament de Física Quàntica i Astrofísica
and Institut de Ciències del Cosmos (ICCUB), Universitat de Barcelona
Martí i Franquès, ES-08028 Barcelona, Spain*

²*Max-Planck-Institut für Gravitationsphysik (Albert-Einstein-Institut)
Am Mühlenberg 1, DE-14476 Potsdam, Germany*

³*International Solvay Institutes
ULB-Campus Plaine CP231, BE-1050 Brussels, Belgium*

⁴*Department of Mathematical Sciences, Chalmers University of Technology
and University of Gothenburg, SE-412 96 Göteborg, Sweden*

⁵*Instituto de Física, Pontificia Universidad Católica de Valparaíso
Casilla 4059, Valparaíso, Chile*

We construct finite- and infinite-dimensional non-relativistic extensions of the Newton–Hooke and Carroll (A)dS algebras using the algebra expansion method, starting from the (anti-)de Sitter relativistic algebra in D dimensions. These algebras are also shown to be embedded in different affine Kac–Moody algebras. In the three-dimensional case, we construct Chern–Simons actions invariant under these symmetries. This leads to a sequence of non-relativistic gravity theories, where the simplest examples correspond to extended Newton–Hooke and extended (post-)Newtonian gravity together with their Carrollian counterparts.

Contents

1	Introduction	2
2	Expansions of the (A)dS algebra	4
2.1	Extended Newton–Hooke algebras	4
2.2	Extended Carroll (A)dS algebras	8
3	Newton–Hooke and Carrollian affine algebra	10
3.1	Newton–Hooke affine algebras	11
3.2	(A)dS Carrollian affine algebras	14
4	Chern–Simons non-relativistic gravities	15
4.1	Extended Newton–Hooke gravities	16
4.2	Extended Carrollian gravities	21
5	Conclusions and Outlook	24
A	Further generalisations	26
A.1	Non-relativistic expansions of the Maxwell algebra	26
A.2	Non-relativistic brane expansions of (A)dS	28

1 Introduction

Non-relativistic symmetries are usually obtained from relativistic ones by means of Lie algebra contractions. The most well-known example is the Wigner–Inönü contraction of the Poincaré algebra that leads to the Galilei algebra [1] by sending the speed of light to infinity. Lie algebra contractions necessarily preserve the number of generators but can alter the cohomology of the Lie algebra, thus allowing for different central extensions like the Bargmann extension of the Galilei algebra [2], which is crucial when taking the limit of the relativistic particle action to the non-relativistic one [3, 4].

In recent years, more general constructions of non-relativistic symmetry algebras from relativistic ones have been explored. One example is given by the method of Lie algebra expansions [5–7] applied to the Poincaré algebra [8]. This method provides an infinite sequence of non-relativistic algebras extending the Galilei algebra with an increasing number of generators, which have been used in [9, 10, 8, 11] to construct various gravitational actions. The Lie algebra expansion method can also be related to a sequence of post-Newtonian limits as shown in [12], and has also been applied to derive diverse non-relativistic symmetries in the context of (super-)gravity [13–17]. Another method is based on a Galilean free Lie algebra [18] that can be thought as the most general extension of the Galilei algebra and, upon taking quotients, has a connection to Lie algebra expansions and Kac–Moody algebras. One interesting conceptual point made in [12] is that the sequence of Lie algebras naturally comes with a generalisation of Minkowski space with more coordinates and an extension of the Minkowski metric. Using

this generalised Minkowski space it is possible to define particle actions invariant under these extended algebras that naturally incorporate post-Newtonian corrections in the non-relativistic limit.

In the present article, we study the non-relativistic symmetries obtained by Lie algebra expansion of the AdS or dS algebra in D space-time dimensions, i.e. $\mathfrak{so}(D-1, 2)$ or $\mathfrak{so}(D, 1)$. This generalises previous constructions to include a cosmological constant and generates an infinite family of algebras of Newton–Hooke [19, 20] type. By taking the limit of the cosmological constant to zero one recovers the non-relativistic algebras introduced in [10] and further studied in [8, 18, 12].

Besides the non-relativistic limit related to Galilean symmetries ($c \rightarrow \infty$) we also consider the case of ultra-relativistic Carrollian limit ($c \rightarrow 0$).¹ As is known from [24] the Carroll algebra can be understood by very specific changes in the commutation relations and associated changes in the starting point of the algebra expansion procedure. Applying the expansion procedure then produces an infinite family of Lie algebras associated with the Carrollian limit. We shall also show how our construction in both cases is related to affine Kac–Moody algebras, where the role of the expansion parameter is taken by the spectral (or loop) parameter of the affine algebra.

After the introduction of the infinite family of algebras we consider gravitational models based on them by focussing on the case of $2+1$ space-time dimensions. We construct Chern–Simons theories based on these algebras since they admit non-degenerate bilinear forms. We show that the family of algebras systematically generates non-relativistic gravity theories extended by a cosmological constant, such as extended Newton–Hooke gravity [25, 26], that reduces to extended Bargmann gravity for $\Lambda \rightarrow 0$ [27, 28], and post-Newtonian gravity [11]².

Our construction also connects to recent discussions of non-relativistic expansions of the metric as described in [29–32, 9, 10, 33]. This can be made very precise in the context of the Chern–Simons formulation and we shall elaborate on this connection in the conclusions.

In an appendix, we generalise the expansion procedure to obtain an infinite family of extensions of the non-relativistic Maxwell algebra. Moreover, we outline that our method is not only applicable non-relativistic symmetries corresponding to point particle limits, but also to extended objects where the D covariant dimensions are split into $p+1$ and $D-p-1$ directions for a p -brane [34, 35]. Similar constructions in the case without a cosmological constant have been considered previously in [36, 24, 8, 37].

Note added. While this manuscript was being finalised, the preprints [38, 39] appeared on the arXiv that have some overlap with some of our results.

¹Work on Carroll symmetries in the context of electrodynamics and brane dynamics can be found for instance in [21–23].

²In [11] this was called ‘extended gravity’ but in the light of the results of [12] this is better understood as a post-Newtonian correction.

2 Expansions of the (A)dS algebra

In this section, we will construct an infinite family of non-relativistic algebras of the Newton–Hooke and Carroll AdS type. The first members of this family will be given by the Newton–Hooke algebra in the former case and the Carroll AdS algebra in the latter (without extensions) [19, 20]. We also obtain an infinite-dimensional algebra that contains these and other intermediate cases, generalises the one obtained in [10, 8, 18], as quotients. All our algebras contain a cosmological constant as we start from the AdS algebra.

We will construct the series of non-relativistic algebras by means of Lie algebra expansions [5–7]. More precisely, we will use the semigroup expansion technique [7] with semigroup $S_E^{(N)}$, which will be defined below.

Our starting point is the (A)dS algebra in D dimensions:

$$[\tilde{J}_{AB}, \tilde{P}_C] = 2\eta_{C[B}\tilde{P}_{A]}, \quad (2.1a)$$

$$[\tilde{J}_{AB}, \tilde{J}_{CD}] = 4\eta_{[A[C}\tilde{J}_{D]B]}, \quad (2.1b)$$

$$[\tilde{P}_A, \tilde{P}_B] = -\Lambda\tilde{J}_{AB}, \quad (2.1c)$$

which denotes in a unified manner $\mathfrak{so}(D-1, 2)$ (for $\Lambda < 0$) and $\mathfrak{so}(D, 1)$ (for $\Lambda > 0$). Here, \tilde{P}_A and \tilde{J}_{AB} are the generators of spacetime translations and Lorentz transformations, respectively. Capital indices run over $A = 0, \dots, D-1$ and the Minkowski metric has been chosen to have mostly plus signature.

In order to perform a non-relativistic expansion of the (A)dS algebra, it is convenient to decompose the relativistic indices in the time and space components, $A = (0, a)$, where $a = 1, \dots, D-1$, and relabel the Lie algebra generators³ as

$$\tilde{J}_{AB} \rightarrow \{\tilde{J}_{0a} \equiv \tilde{G}_a, \tilde{J}_{ab}\}, \quad \tilde{P}_A \rightarrow \{\tilde{P}_0 \equiv \tilde{H}, \tilde{P}_a\}. \quad (2.2)$$

Then, the commutation relations (2.1) can be rewritten in the form

$$[\tilde{G}_a, \tilde{H}] = \tilde{P}_a, \quad (2.3a) \quad [\tilde{J}_{ab}, \tilde{G}_c] = 2\delta_{c[b}\tilde{G}_{a]}, \quad (2.3e)$$

$$[\tilde{G}_a, \tilde{P}_b] = \delta_{ab}\tilde{H}, \quad (2.3b) \quad [\tilde{J}_{ab}, \tilde{J}_{cd}] = 4\delta_{[a[c}\tilde{J}_{d]b]}, \quad (2.3f)$$

$$[\tilde{J}_{ab}, \tilde{P}_c] = 2\delta_{c[b}\tilde{P}_{a]}, \quad (2.3c) \quad [\tilde{P}_a, \tilde{H}] = \Lambda\tilde{G}_a, \quad (2.3g)$$

$$[\tilde{G}_a, \tilde{G}_b] = \tilde{J}_{ab}, \quad (2.3d) \quad [\tilde{P}_a, \tilde{P}_b] = -\Lambda\tilde{J}_{ab}. \quad (2.3h)$$

We note that the decomposition (2.2) is adapted to point particles in the sense that one direction—that can be thought of as the world-line direction of the particle—is singled out. In appendix A.2, we also consider the case of extended objects.

2.1 Extended Newton–Hooke algebras

In order to perform expansions of (A)dS of Newton–Hooke type, we first note that (2.3) allows for the following subspace decomposition $V = V_0 \oplus V_1$ [8]

$$V_0 = \{\tilde{H}, \tilde{J}_{ab}\}, \quad V_1 = \{\tilde{P}_a, \tilde{G}_a\}, \quad (2.4)$$

³Notice this convention is different from the one in [12] where $\tilde{J}_{a0} \equiv \tilde{G}_a$ was used.

which satisfies a \mathbb{Z}_2 -graded structure, i.e.,

$$[V_0, V_0] \subset V_0, \quad [V_0, V_1] \subset V_1, \quad [V_1, V_1] \subset V_0. \quad (2.5)$$

This decomposition is similar to a symmetric space decomposition. The homogeneous coset space in the case of AdS is $SO(D-1, 2)/(SO(D-1) \times \mathbb{R})$ and $SO(D, 1)/(SO(D-1) \times \mathbb{R})$ in the dS case.

We will consider expansions with the semigroup $S_E^{(N)} = \{\lambda_0, \dots, \lambda_{N+1}\}$, whose multiplication law is given by [7]

$$\lambda_i \cdot \lambda_j = \begin{cases} \lambda_{i+j} & \text{if } i+j \leq N, \\ \lambda_{N+1} & \text{if } i+j > N, \end{cases} \quad (2.6)$$

where λ_{N+1} acts as the zero of the semigroup, as it satisfies $\lambda_{N+1} \cdot \lambda_i = \lambda_i \cdot \lambda_{N+1} = \lambda_{N+1}$ for all λ_i . This semigroup admits the subset decomposition

$$\begin{aligned} S_0^{(N)} &= s_0^{(N)} \cup \{\lambda_{N+1}\}, & s_0^{(N)} &= \{\lambda_{2m} \mid m = 0, \dots, [\frac{N}{2}]\}, \\ S_1^{(N)} &= s_1^{(N)} \cup \{\lambda_{N+1}\}, & s_1^{(N)} &= \{\lambda_{2m+1} \mid m = 0, \dots, [\frac{N-1}{2}]\}, \end{aligned} \quad (2.7)$$

which is compatible with (2.5) in the sense that

$$S_0^{(N)} \cdot S_0^{(N)} \subset S_0^{(N)}, \quad S_0^{(N)} \cdot S_1^{(N)} \subset S_1^{(N)}, \quad S_1^{(N)} \cdot S_1^{(N)} \subset S_0^{(N)}, \quad (2.8)$$

and therefore *resonant* with the choice of V_0 and V_1 in (2.4).

Thus, a reduced resonant expanded algebra can be defined as the direct sum

$$\left\{ s_0^{(N)} \otimes V_0 \right\} \oplus \left\{ s_1^{(N)} \otimes V_1 \right\} \quad (2.9)$$

where the reduction condition in the algebra is implemented by the constraints

$$\lambda_{N+1} \otimes \tilde{J}_{AB} = 0, \quad \lambda_{N+1} \otimes \tilde{P}_A = 0, \quad (2.10)$$

which map the zero λ_{N+1} of the semigroup to the zero element in the expanded algebra. In the following we will show how, for different choices of the semigroup $S_E^{(N)}$, the reduction (2.9) lead to a non-relativistic algebras that generalise the Newton–Hooke symmetry.

The simplest example in our construction corresponds to the expansion with the semigroup $S_E^{(1)}$, which as we will see is equivalent to a non-relativistic contraction of the (A)dS algebra. In other words, this expansion gives the Newton–Hooke algebras. By setting $N = 1$ in (2.6) and (2.9), the generators of the expanded algebra are given by

$$\begin{aligned} J_{ab} &= \lambda_0 \otimes \tilde{J}_{ab}, & G_a &= \lambda_1 \otimes \tilde{G}_a, \\ H &= \lambda_0 \otimes \tilde{H}, & P_a &= \lambda_1 \otimes \tilde{P}_a. \end{aligned} \quad (2.11)$$

Using the relativistic commutation relations (2.3) together with the reduction condition (2.10) for the zero element λ_2 , the commutation relations for the generators (2.11) read

$$[G_a, H] = P_a, \quad (2.12a) \quad [J_{ab}, J_{cd}] = 4\delta_{[a[c}J_{d]b]}, \quad (2.12d)$$

$$[J_{ab}, P_c] = 2\delta_{c[b}P_{a]}, \quad (2.12b) \quad [P_a, H] = \Lambda G_a, \quad (2.12e)$$

$$[J_{ab}, G_c] = 2\delta_{c[b}G_{a]}, \quad (2.12c)$$

which corresponds to the Newton–Hooke $_{\mp}$ algebra (with $\Lambda \leq 0$) without extensions [19, 20], this algebra reduces to the Galilei symmetry for $\Lambda = 0$. Therefore the use of the semigroup $S_E^{(N)}$ will create a family of generalised Newton–Hooke algebras for all the possible values of N .

The case with $N = 2$ leads to a Newton–Hooke algebra with some extensions. Unlike the previous case, when using the semigroup $S_E^{(2)}$, the element λ_2 is no longer the zero of the semigroup. In this case the reduction condition (2.10) holds for a new semigroup zero element λ_3 , and (2.11) has to be supplemented with two extra expanded generators

$$S_{ab} = \lambda_2 \otimes \tilde{J}_{ab}, \quad M = \lambda_2 \otimes \tilde{H}. \quad (2.13)$$

Using the semigroup product law (2.6) for $N = 2$, we find that the non-vanishing commutation relations for the expanded algebra are given by the Newton–Hooke commutators (2.12) together with

$$[G_a, P_b] = \delta_{ab}M, \quad (2.14a) \quad [J_{ab}, S_{cd}] = 4\delta_{[a[c}S_{d]b]}, \quad (2.14c)$$

$$[G_a, G_b] = S_{ab}, \quad (2.14b) \quad [P_a, P_b] = -\Lambda S_{ab}. \quad (2.14d)$$

This expansion produces the Bargmann central extension M and a non-central extension S_{ab} [40]. This algebra can be understood as the generalisation of the double central extension [41–43] of the Newton–Hooke algebra to $D > 3$. In fact, in 2+1 dimensions the generator S_{ab} can be dualised to a scalar and becomes central. In that case the algebra (2.14) becomes exactly the extended Newton–Hooke algebra ($\Lambda \neq 0$) [44, 45] or the extended Bargmann algebra ($\Lambda = 0$) [46, 47].

Extended (post-)Newtonian gravity algebra

For $N = 3$ we obtain the Newton–Hooke version of the algebra found in [9] as the symmetry of post-Newtonian gravity. Moreover, when considering $N = 4$, we find an extension of the post-Newtonian gravity algebra that generalises the algebra of [11] to any dimension. Explicitly, for $N = 3$ the generators of the expanded algebra are given by (2.11), (2.13) and

$$B_a = \lambda_3 \otimes \tilde{G}_a, \quad (2.15)$$

$$T_a = \lambda_3 \otimes \tilde{P}_a.$$

The zero of the semigroup $S_E^{(3)}$ is λ_4 , and the reduction condition (2.10) together with the semigroup law and the (A)dS commutation relations (2.3) leads to the following non-vanishing commutators for the expanded algebra:

$$\begin{aligned}
[G_a, H] &= P_a, & (2.16a) & & [J_{ab}, S_{cd}] &= 4\delta_{[a[c}S_{d]b]}, & (2.16i) \\
[G_a, M] &= T_a, & (2.16b) & & [P_a, H] &= \Lambda G_a, & (2.16j) \\
[B_a, H] &= T_a, & (2.16c) & & [P_a, M] &= \Lambda B_a, & (2.16k) \\
[G_a, P_b] &= \delta_{ab}M, & (2.16d) & & [T_a, H] &= \Lambda B_a, & (2.16l) \\
[G_a, G_b] &= S_{ab}, & (2.16e) & & [P_a, P_b] &= -\Lambda S_{ab} & (2.16m) \\
[S_{ab}, G_c] &= 2\delta_{c[b}B_{a]}, & (2.16f) & & [J_{ab}, X_c] &= 2\delta_{c[b}X_{a]}, & (2.16n) \\
[S_{ab}, P_c] &= 2\delta_{c[b}T_{a]}, & (2.16g) & & X_a &= \{G_a, P_a, B_a, T_a\}. \\
[J_{ab}, J_{cd}] &= 4\delta_{[a[c}J_{d]b]}, & (2.16h) & & & &
\end{aligned}$$

In the case $\Lambda = 0$ this algebra corresponds to the one found in [9] and further studied in [8, 18, 12].

One can extend this algebra by considering $N = 4$ in the expansion prescription (2.9). In this case, (2.15) has to be supplemented with extra expanded generators given by

$$Z_{ab} = \lambda_4 \otimes \tilde{J}_{ab}, \quad Y = \lambda_4 \otimes \tilde{H}. \quad (2.17)$$

The zero element in $S_E^{(4)}$ is given by λ_5 , which means that the reduction condition (2.10) in this case sets $\lambda_5 \otimes \tilde{J}_{AB} = \lambda_5 \otimes \tilde{P}_A \equiv 0$ and the non-vanishing commutation relations of the corresponding expanded algebra are given by (2.16) together with

$$[G_a, T_b] = \delta_{ab}Y, \quad (2.18a) \quad [J_{ab}, Z_{cd}] = 4\delta_{[a[c}Z_{d]b]}, \quad (2.18d)$$

$$[B_a, P_b] = \delta_{ab}Y, \quad (2.18b) \quad [P_a, T_b] = -\Lambda Z_{ab}. \quad (2.18e)$$

$$[G_a, B_b] = Z_{ab}, \quad (2.18c) \quad (2.18f)$$

In the case $D = 2 + 1$ and $\Lambda = 0$ this algebra corresponds to the one studied in [11] and defines a central extension of the post-Newtonian gravity symmetry (2.18). For $D > 2 + 1$, however, the generator Z_{ab} is no longer central, exactly as it happens with S_{ab} in the extension of the Newton–Hooke algebra.

Notice that if some generators are central for some value N of the semigroup, they are no longer central in the $N + 1$ case. More specifically, the expansions with odd N do not possess central elements, while expansions with even values of N always yield two central elements given by $\lambda_N \otimes \tilde{J}$ and tilde $\lambda_N \otimes \tilde{H}$.

Instead of continuing with increasing values of N , in the next subsection we will construct an infinite-dimensional graded algebra using an infinite-dimensional semigroup.

Infinite-dimensional Galilean algebra

The previous analysis suggests that we can construct an infinite-dimensional graded non-relativistic algebra using the expansion procedure with an infinite-dimensional semigroup of the form [14]

$$S^{(\infty)} = \{\lambda_0, \lambda_1, \lambda_2, \dots\} \quad (2.19)$$

with multiplication law

$$\lambda_\alpha \cdot \lambda_\beta = \lambda_{\alpha+\beta}. \quad (2.20)$$

(This semigroup can be realised as powers of a formal variable ω by letting $\lambda_\alpha = \omega^\alpha$. The finite semigroups $S_E^{(N)}$ then correspond to working to order $O(\omega^{N+1})$.) As this semigroup does not have a zero element, the resonant subset decomposition that satisfies (2.8) in this case is simply given by

$$\begin{aligned} S_0^{(\infty)} &= \{\lambda_{2m} \mid m = 0, 1, 2, \dots\}, \\ S_1^{(\infty)} &= \{\lambda_{2m+1} \mid m = 0, 1, 2, \dots\}, \end{aligned} \quad (2.21)$$

and the corresponding resonant (non-reduced) expansion of (A)dS is

$$\left\{ S_0^{(\infty)} \otimes V_0 \right\} \oplus \left\{ S_1^{(\infty)} \otimes V_1 \right\}, \quad (2.22)$$

where V_0 and V_1 were defined in (2.4). Denoting the generators of the expanded algebra (2.22) as

$$\begin{aligned} J_{ab}^{(m)} &= \lambda_{2m} \otimes \tilde{J}_{ab}, & B_a^{(m)} &= \lambda_{2m+1} \otimes \tilde{G}_a, \\ H^{(m)} &= \lambda_{2m} \otimes \tilde{H}, & P_a^{(m)} &= \lambda_{2m+1} \otimes \tilde{P}_a, \end{aligned} \quad (2.23)$$

leads to the infinite-dimensional graded algebra

$$[J_{ab}^{(m)}, P_c^{(n)}] = 2\delta_{c[b} P_a^{(m+n)}], \quad (2.24a) \quad [J_{ab}^{(m)}, B_c^{(n)}] = 2\delta_{c[b} B_a^{(m+n)}], \quad (2.24e)$$

$$[J_{ab}^{(m)}, J_{cd}^{(n)}] = 4\delta_{[a[c} J_{d]b}^{(m+n)}], \quad (2.24b) \quad [B_a^{(m)}, B_b^{(n)}] = J_{ab}^{(m+n+1)}, \quad (2.24f)$$

$$[B_a^{(m)}, P_b^{(n)}] = \delta_{ab} H^{(m+n+1)}, \quad (2.24c) \quad [P_a^{(m)}, H^{(n)}] = \Lambda B_a^{(m+n)}, \quad (2.24g)$$

$$[B_a^{(m)}, H^{(n)}] = P_a^{(m+n)}, \quad (2.24d) \quad [P_a^{(m)}, P_b^{(n)}] = -\Lambda J_{ab}^{(m+n+1)}. \quad (2.24h)$$

We can get the finite-dimensional algebras presented in the previous subsections as quotients of (2.24) by suitable infinite ideals. For example, in the case of (2.16) the ideal is generated by $J_{ab}^{(m)}, B_a^{(m)}, P_a^{(m)}, H^{(m)}$ for $m > 2$.

Taking the limit $\Lambda \rightarrow 0$, we obtain as a contraction of (2.24), the infinite-dimensional extension of the Galilei algebra introduced in [10] and further studied in [18].

2.2 Extended Carroll (A)dS algebras

The (A)dS Carroll algebra corresponds to the ultra-relativistic contraction of the (A)dS algebra (2.1) [19]. It can alternatively be obtained by means of an expansion procedure when considering the following subspace decomposition for (A)dS,

$$V_0 = \{\tilde{P}_a, \tilde{J}_{ab}\}, \quad V_1 = \{\tilde{H}, \tilde{G}_b\}, \quad (2.25)$$

which is \mathbb{Z}_2 -graded. Note that here P_a and H have been interchanged with respect to the subspace decomposition used in the Newton–Hooke case (2.4). This is a special case of a general duality at the level of the translation generators between Galilean and Carrollian symmetries [24]. Using the decomposition (2.25), we can generalise the procedure outlined in the previous section to define Carrollian expansions of (A)dS, whose simplest case is the Carroll (A)dS algebra. Indeed, the non-vanishing commutation relations of the corresponding reduced resonant expanded algebra is given by

$$\left(s_0^{(N)} \times \{\tilde{P}_a, \tilde{J}_{ab}\} \right) \oplus \left(s_1^{(N)} \times \{\tilde{H}, \tilde{G}_b\} \right), \quad (2.26)$$

where $s_0^{(N)}$ and $s_1^{(N)}$ are given in (2.7), and the case $N = 1$ leads to the (A)dS Carroll algebra in D dimensions

$$[J_{ab}, P_c] = 2\delta_{c[b}P_a], \quad (2.27a) \quad [J_{ab}, G_c] = 2\delta_{c[b}G_a], \quad (2.27d)$$

$$[J_{ab}, J_{cd}] = 4\delta_{[a[c}J_{d]b}], \quad (2.27b) \quad [P_a, H] = \Lambda G_a, \quad (2.27e)$$

$$[G_a, P_b] = \delta_{ab}H, \quad (2.27c) \quad [P_a, P_b] = -\Lambda J_{ab}, \quad (2.27f)$$

where the expanded generators have been defined as

$$\begin{aligned} P_a &= \lambda_0 \otimes \tilde{P}_a, & H &= \lambda_1 \otimes \tilde{H}, \\ J_{ab} &= \lambda_0 \otimes \tilde{J}_{ab}, & G_a &= \lambda_1 \otimes \tilde{G}_a. \end{aligned} \quad (2.28)$$

Naturally, when setting $\Lambda = 0$, this allows one to obtain the Carroll algebra in D dimensions as an expansion of the Poincaré algebra. Subsequently, we can consider greater values of N to obtain extended Carroll (A)dS algebras. In the case $N = 2$ the expanded algebra has extra generators given by

$$\begin{aligned} T_a &= \lambda_2 \otimes \tilde{P}_a, \\ S_{ab} &= \lambda_2 \otimes \tilde{J}_{ab}. \end{aligned} \quad (2.29)$$

and the commutation relations are the ones of (2.27) together with

$$[G_a, H] = T_a, \quad (2.30a) \quad [S_{ab}, P_c] = 2\delta_{c[b}T_a], \quad (2.30d)$$

$$[G_a, G_b] = S_{ab}, \quad (2.30b) \quad [J_{ab}, S_{cd}] = 4\delta_{[a[c}S_{d]b}], \quad (2.30e)$$

$$[J_{ab}, T_c] = 2\delta_{c[b}T_a], \quad (2.30c) \quad [P_a, T_b] = -\Lambda S_{ab}. \quad (2.30f)$$

In the case $N = 3$, we get two extra generators,

$$\begin{aligned} M &= \lambda_3 \otimes \tilde{H}, \\ B_a &= \lambda_3 \otimes \tilde{G}_a. \end{aligned} \quad (2.31)$$

The commutation relations of this expanded algebra are (2.27), (2.30) plus

$$[J_{ab}, B_c] = 2\delta_{c[b}B_a], \quad (2.32a) \quad [S_{ab}, G_c] = 2\delta_c B_a], \quad (2.32d)$$

$$[G_a, T_b] = \delta_{ab}M, \quad (2.32b) \quad [T_a, H] = \Lambda B_a, \quad (2.32e)$$

$$[B_a, P_b] = \delta_{ab}M, \quad (2.32c) \quad [P_a, M] = \Lambda B_a. \quad (2.32f)$$

For $\Lambda = 0$, this algebra defines a Carrollian counterpart of the (post-)Newtonian symmetry introduced in [10] in the context of general relativity. In the same way, one can check that the $N = 4$ case defines a higher-dimensional Carrollian (A)dS analogue of the extended (post-)Newtonian symmetry given in [11].

Infinite-dimensional Carroll algebra

Similarly to the Newton–Hooke construction, the ultra-relativistic expansions of (A)dS form a family of algebras, which can be described in a unified fashion by considering the infinite semigroup (2.19) and the non-reduced resonant expansion

$$\left(S_0^{(\infty)} \times \left\{ \tilde{J}_{ab}, \tilde{P}_a \right\} \right) \oplus \left(S_1^{(\infty)} \times \left\{ \tilde{G}_a, \tilde{H} \right\} \right), \quad (2.33)$$

where $S_0^{(\infty)}$ and $S_1^{(\infty)}$ are given in (2.21). By defining an infinite set of expanded generators by

$$\begin{aligned} J_{ab}^{(m)} &= \lambda_{2m} \otimes \tilde{J}_{ab}, & B_a^{(m)} &= \lambda_{2m+1} \otimes \tilde{G}_a, \\ P_a^{(m)} &= \lambda_{2m} \otimes \tilde{P}_a, & H^{(m)} &= \lambda_{2m+1} \otimes \tilde{H}, \end{aligned} \quad (2.34)$$

we obtain an infinite-dimensional Carrollian expansion of the (A)dS algebra

$$[J_{ab}^{(m)}, P_c^{(n)}] = 2\delta_{c[b} P_a^{(m+n)}], \quad (2.35a) \quad [J_{ab}^{(m)}, B_c^{(n)}] = 2\delta_{c[b} B_a^{(m+n)}], \quad (2.35e)$$

$$[J_{ab}^{(m)}, J_{cd}^{(n)}] = 4\delta_{[a[c} J_{d]b}^{(m+n)}], \quad (2.35b) \quad [B_a^{(m)}, B_b^{(n)}] = J_{ab}^{(m+n+1)}. \quad (2.35f)$$

$$[B_a^{(m)}, P_b^{(n)}] = \delta_{ab} H^{(m+n)}, \quad (2.35c) \quad [P_a^{(m)}, H^{(n)}] = \Lambda B_a^{(m+n)}, \quad (2.35g)$$

$$[B_a^{(m)}, H^{(n)}] = P_a^{(m+n+1)}, \quad (2.35d) \quad [P_a^{(m)}, P_b^{(n)}] = -\Lambda J_{ab}^{(m+n)}. \quad (2.35h)$$

The different finite expansions previously constructed using the semigroup $S_E^{(N)}$ can be obtained from the infinite case by considering suitable quotients. Redefining the generators according to

$$\begin{aligned} J_{ab}^{(m)} &\rightarrow J_{ab}^{(m)}, \\ P_a^{(m)} &\rightarrow P_a^{(m-1)} \quad (m \geq 1), \\ B_a^{(m)} &\rightarrow B_a^{(m)}, \\ H^{(m)} &\rightarrow -H^{(m)} \end{aligned} \quad (2.36)$$

and, taking the limit $\Lambda \rightarrow 0$, we obtain an extension of the infinite-dimensional algebra in [10] as a contraction of the infinite-dimensional extended Carrollian algebra (2.35). A related contraction was obtained from the infinite-dimensional extended Newton–Hooke algebra (2.24) above. In the contraction of the Carroll algebra (2.35), it is extended in a semidirect sum by the additional generator $P_a^{(0)}$ (which should then rather be called $P_a^{(-1)}$ in the notation of [10]). Conversely, the algebra in [10] can be seen as an extension of the contracted infinite-dimensional extended Carroll algebra if we again redefine $H^{(m)} \rightarrow H^{(m+1)}$ and then add a generator $H^{(0)}$.

3 Newton–Hooke and Carrollian affine algebra

In this section, we will show that the infinite-dimensional Lie algebras (2.24) and (2.35) also can be obtained from the extension of $\mathfrak{so}(D-1)$ to an (untwisted or twisted) affine Kac–Moody algebra. We shall only require parabolic subalgebras of these Kac–Moody algebras and this construction can be linked to free Lie algebras in terms of quotients by Serre relations. The algebras for finite N discussed in the previous section then correspond to further quotients, similar to the constructions in [48, 18].




Figure 1: *Dynkin diagrams of $B_{r+1}^{(1)}$ (lower) and $D_{r+1}^{(1)}$ (upper). These are relevant for the infinite Newton–Hooke algebras.*

3.1 Newton–Hooke affine algebras

Consider first the complex Lie algebra $\mathfrak{g}_r = D_r$ if $d = D - 1$ is even ($d = 2r$), or $\mathfrak{g}_r = B_r$ if $d = D - 1$ is odd ($d = 2r + 1$). Thus \mathfrak{g}_r is the complexification of $\mathfrak{so}(d)$. We can extend \mathfrak{g}_r to \mathfrak{g}_{r+1} (either D_{r+1} or B_{r+1}) by adding a node labelled 0 and then further to an affine Kac–Moody algebra $\mathfrak{g}_{r+1}^{(1)}$ by adding a node labelled -1 to the Dynkin diagram of \mathfrak{g}_r . The resulting diagram, together with our labelling of the nodes, is shown in Figure 1. The corresponding Cartan matrix is

$$A_{ij} = \begin{pmatrix} 2 & 0 & -1 & \cdots & 0 & 0 & 0 \\ 0 & 2 & -1 & \cdots & 0 & 0 & 0 \\ -1 & -1 & 2 & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 2 & -1 & -1 \\ 0 & 0 & 0 & \cdots & -1 & 2 & 0 \\ 0 & 0 & 0 & \cdots & -1 & 0 & 2 \end{pmatrix} \quad (3.1)$$

for $D_{r+1}^{(1)}$ and

$$A_{ij} = \begin{pmatrix} 2 & -2 & 0 & \cdots & 0 & 0 & 0 \\ -1 & 2 & -1 & \cdots & 0 & 0 & 0 \\ 0 & -1 & 2 & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 2 & -1 & 0 \\ 0 & 0 & 0 & \cdots & -1 & 2 & -1 \\ 0 & 0 & 0 & \cdots & 0 & -2 & 2 \end{pmatrix} \quad (3.2)$$

for $B_{r+1}^{(1)}$ (where rows and columns are counted $-1, 0, 1, \dots, r$ from left to right and from top to bottom).

When we add the two nodes 0 and -1 we also add six generators e_i, f_i, h_i for $i = 0, -1$ to the generators of \mathfrak{g}_r . This gives a central extension of the loop algebra of \mathfrak{g}_{r+1} where the central element is given by

$$c = h_{-1} + h_0 + 2h_1 + 2h_2 + \dots + 2h_{r-2} + h_{r-1} + h_r. \quad (3.3)$$

The affine algebra $\mathfrak{g}_{r+1}^{(1)}$ is then obtained from it by adding also a derivation generator d satisfying $[d, e_{-1}] = e_{-1}$ and $[d, f_{-1}] = -f_{-1}$ and commuting with all the other generators.

We consider the subalgebra of $\mathfrak{g}_{r+1}^{(1)}$ generated by e_i, h_i for $i = 0, -1$ together with the generators of \mathfrak{g}_r . This Lie algebra has an $(\mathbb{N} \times \mathbb{N})$ -grading associated to these two nodes, which can be trivially extended to a $(\mathbb{Z} \times \mathbb{Z})$ -grading. It can thus be decomposed into a direct sum of subspaces, each labelled by a pair (ℓ_0, ℓ_{-1}) of non-negative integers. The subspace labelled by (ℓ_0, ℓ_{-1}) is spanned by elements formed as multibrackets of the generators, where e_0 and e_{-1} appear ℓ_0 and ℓ_{-1} times, respectively. The subalgebra at $(\ell_0, \ell_{-1}) = (0, 0)$ contains \mathfrak{g}_r but also the two additional Cartan generators h_0 and h_{-1} . By taking the linear combinations

$$\begin{aligned} h &= h_{-1} + h_1 + h_2 + \dots + h_{r-2} + \frac{1}{2}h_{r-1} + \frac{1}{2}h_r, \\ h' &= h_0 + h_1 + h_2 + \dots + h_{r-2} + \frac{1}{2}h_{r-1} + \frac{1}{2}h_r, \end{aligned} \quad (3.4)$$

(such that $c = h + h'$) we get elements that commute with \mathfrak{g}_r . Furthermore, they satisfy

$$\begin{aligned} [h, e_{-1}] &= e_{-1}, & [h', e_{-1}] &= -e_{-1}, \\ [h, e_0] &= -e_0, & [h', e_0] &= e_0. \end{aligned} \quad (3.5)$$

The $(\mathbb{Z} \times \mathbb{Z})$ -grading gives rise to a \mathbb{Z} -grading, where the single level ℓ is the sum of ℓ_0 and ℓ_{-1} . Since \mathfrak{g}_r is a subalgebra at level zero, we get representations of it at each level.

At level $\ell = 0$ we have the generators J_{ab} of (the complexification of) $\mathfrak{so}(d)$ and the two $\mathfrak{so}(d)$ scalars h and h' . At level $\ell = 1$ we have two lowest weight representations with lowest weight vectors e_{-1} and e_0 , respectively. The Dynkin labels of both corresponding highest weight representations are $[1, 0, 0, \dots, 0]$ since

$$\begin{aligned} [h_1, e_{-1}] &= -1, & [h_2, e_{-1}] &= [h_2, e_{-1}] = \dots = [h_r, e_{-1}] = 0, \\ [h_1, e_0] &= -1, & [h_2, e_0] &= [h_2, e_0] = \dots = [h_r, e_0] = 0. \end{aligned} \quad (3.6)$$

Thus they are vector representations, and we denote the corresponding generators by X_a (with lowest weight vector e_{-1}) and Y_a (with lowest weight vector e_0), where $a = 1, 2, \dots, d$ as before. It follows from (3.5) that

$$\begin{aligned} [h, X_a] &= X_a, & [h', X_a] &= -X_a, \\ [h, Y_a] &= -Y_a, & [h', Y_a] &= Y_a. \end{aligned} \quad (3.7)$$

(The parabolic subalgebra of the ‘horizontal’ algebra \mathfrak{g}_{r+1} is generated by J_{ab}, Y_a and h_0 .)

In the *free Lie algebra* generated by all X_a and Y_a at level $\ell = 1$, the subspace at level $\ell = 2$ decomposes into a direct sum of $\mathfrak{so}(d)$ modules with Dynkin labels

$$\begin{aligned} 2[1, 0, 0, \dots, 0] \wedge 2[1, 0, 0, \dots, 0] &= 2([1, 0, 0, \dots, 0] \wedge [1, 0, 0, \dots, 0]) \\ &\quad \oplus [1, 0, 0, \dots, 0] \otimes [1, 0, 0, \dots, 0] \\ &= 3[0, 1, 0, \dots, 0] \oplus [2, 0, 0, \dots, 0] \oplus [0, 0, 0, \dots, 0], \end{aligned} \quad (3.8)$$

(where \wedge denotes the antisymmetric tensor product, coming from the antisymmetry of the Lie bracket). The free Lie algebra construction continues to all positive levels [48, 18] but in order to reproduce the algebra (2.24) we need to take a quotient. This quotient leads to the subalgebra of $\mathfrak{g}_{r+1}^{(1)}$ at positive levels and the ideal that one quotients out is generated by the *Serre relations*

$$[e_{-1}, [e_{-1}, e_1]] = [e_0, [e_0, e_1]] = [e_{-1}, e_0] = 0, \quad (3.9)$$

corresponding to the representation

$$2[0, 1, 0, \dots, 0] \oplus [2, 0, 0, \dots, 0]. \quad (3.10)$$

Thus, this representation has to be removed from the antisymmetric tensor product (3.8), leaving only the direct sum

$$[0, 1, 0, \dots, 0] \oplus [0, 0, 0, \dots, 0] \quad (3.11)$$

at level $\ell = 2$. We denote the corresponding 2-form and scalar generators by J_{ab}^2 and h^2 , respectively, where the superscript indicates that they appear at level $\ell = 2$. The commutation relations among the generators at level $\ell = 1$ giving rise to these generators at level $\ell = 2$ are

$$[X_a, Y_b] = J_{ab}^2 + \delta_{ab} h^2, \quad [X_a, X_b] = [Y_a, Y_b] = 0. \quad (3.12)$$

The pattern with two vectors at odd levels and a 2-form and a scalar at even levels continues, as shown in Table 1. The set of generators at non-negative levels are⁴

$$\{h', h^{2k}, J_{ab}^{2k}, X_a^{2k+1}, Y_a^{2k+1}\} \quad (3.13)$$

for $k \geq 0$ (where again the superscript is the level ℓ), and the non-vanishing commutation relations (except for those involving h' , which we have omitted since they turn out to be irrelevant) are

$$\begin{aligned} [J_{ab}^{2k}, J_{cd}^{2k'}] &= 4\delta_{[c[b} J_{a]d]}^{2(k+k')}, \\ [J_{ab}^{2k}, X_c^{2k'+1}] &= 2\delta_{c[b} X_a]^{2(k+k')+1}, \\ [h^{2k}, X_a^{(2k'+1)}] &= X_a^{2(k+k')+1}, \\ [J_{ab}^{2k}, Y_c^{2k'+1}] &= 2\delta_{c[b} Y_a]^{2(k+k')+1}, \\ [h^{2k}, Y_a^{2k'+1}] &= -Y_a^{2(k+k')+1}, \\ [X_a^{2k+1}, Y_b^{2k'+1}] &= J_{ab}^{2(k+k'+1)} + \delta_{ab} h^{2(k+k'+1)}, \end{aligned} \quad (3.14)$$

⁴In this notation, we have singled out h' arbitrarily. As the diagrams are symmetric under the exchange of nodes 0 and -1 , we could have also exchanged the roles of h and h' .

	$\ell_{-1} = 0$	$\ell_{-1} = 1$	$\ell_{-1} = 2$	$\ell_{-1} = 3$	$\ell_{-1} = 4$	\dots
$\ell_0 = 0$	J_{ab}^0, h^0, h'	X_a^1				
$\ell_0 = 1$	Y_a^1	J_{ab}^2, h^2	X_a^3			
$\ell_0 = 2$		Y_a^3	J_{ab}^4, h^4	X_a^5		
$\ell_0 = 3$			Y_a^5	J_{ab}^6, h^6	X_a^7	
$\ell_0 = 4$				Y_a^7	J_{ab}^8, h^8	\ddots
\vdots					\ddots	\ddots

Table 1: *The non-negative levels of the affine Kac–Moody algebra $\mathfrak{g}_{r+1}^{(1)}$.*

where $J_{ab}^0 = J_{ab}$, $h^0 = h$, $X_a^1 = X_a$ and $Y_a^1 = Y_a$. If we then set

$$\begin{aligned}
J_{ab} &= J_{ab}^0, & S_{ab} &= J_{ab}^2, \\
H &= \sqrt{\Lambda} h^0, & M &= \sqrt{\Lambda} h^2, \\
G_a &= \frac{1}{\sqrt{2}}(Y_a^1 + X_a^1), & B_a &= \frac{1}{\sqrt{2}}(Y_a^3 + X_a^3), \\
P_a &= \frac{\sqrt{\Lambda}}{\sqrt{2}}(Y_a^1 - X_a^1), & T_a &= \frac{\sqrt{\Lambda}}{\sqrt{2}}(Y_a^3 - X_a^3),
\end{aligned} \tag{3.15}$$

for $\Lambda > 0$, then we recover the commutation relations (2.16). More generally, if we set

$$\begin{aligned}
J_{ab}^{(m)} &= J_{ab}^{2m}, \\
H^{(m)} &= \sqrt{\Lambda} h^{2m}, \\
B_a^{(m)} &= \frac{1}{\sqrt{2}}(Y_a^{2m+1} + X_a^{2m+1}), \\
P_a^{(m)} &= \frac{\sqrt{\Lambda}}{\sqrt{2}}(Y_a^{2m+1} - X_a^{2m+1})
\end{aligned} \tag{3.16}$$

for $\Lambda > 0$, then we recover the commutation relations (2.24). These formulas are still valid for $\Lambda < 0$ if we interpret $\sqrt{\Lambda}$ as $\pm i\sqrt{|\Lambda|}$ and we will then obtain a different real form of the complex Lie algebra $\mathfrak{g}_{r+1}^{(1)}$.

3.2 (A)dS Carrollian affine algebras

The procedure for obtaining (2.35) from an affine algebra is very similar to the previous discussion so we shall be rather brief.

We first note that due to the choice of subspaces in (2.25), the infinitely expanded algebra (2.35) consists of infinite repetitions of copies of $\{\tilde{J}_{ab}, \tilde{P}_a\}$ as even level spaces and $\{\tilde{H}, \tilde{G}_a\}$




Figure 2: *Dynkin diagrams of $B_{r+1}^{(1)}$ (upper) and $D_{r+1}^{(2)}$ (lower). These are relevant to the (A)dS Carroll algebras.*

as odd level spaces, see (2.33). Since in the Carrollian limit the commutation relations (2.27) are perfectly compatible with identifying the complexification of $V_0 = \{\tilde{J}_{ab}, \tilde{P}_a\}$ as the algebra $\mathfrak{so}(D)$ and the complexification of $V_1 = \{\tilde{H}, \tilde{G}_a\}$ as its D -dimensional vector representation, we have to look for a Kac–Moody algebra where these two spaces repeat infinitely. Thus, there must be Serre relations corresponding to an ideal such that we obtain the positive levels as a quotient of the free Lie algebra generated by the space V_1 .

The problem turns out to be identical to one already solved in [18], where the corresponding algebras were identified as the untwisted affine algebra of type $B_{r+1}^{(1)}$ (for D even and $r+1 = D/2$) and as the twisted affine algebra $D_{r+1}^{(2)}$ (for D odd and $r = (D-1)/2$). The corresponding Dynkin diagrams are reproduced in Figure 2 for convenience. By redefining the generators in (2.35) according to

$$\begin{aligned}
 J_{ab}^{(m)} &\rightarrow J_{ab}^{(m)}, \\
 P_a^{(m)} &\rightarrow \sqrt{-\Lambda} P_a^{(m-1)} \quad (m \geq 1), \\
 B_a^{(m)} &\rightarrow B_a^{(m)}, \\
 H^{(m)} &\rightarrow -\sqrt{-\Lambda} H^{(m)},
 \end{aligned} \tag{3.17}$$

in the AdS case ($\Lambda < 0$) we recover the commutation relations in (3.11) of [18]. The dS case ($\Lambda > 0$) corresponds to taking a different real form of the complex Kac–Moody algebra, interpreting $\sqrt{-\Lambda}$ as $\pm i\sqrt{\Lambda}$.

4 Chern–Simons non-relativistic gravities

In this section we want to find realisations of the previous non-relativistic symmetries. We will be interested in the construction of Chern–Simons gravities in $2+1$ dimensions, defined by the action⁵

$$S_{\text{CS}}[A] = \int \left\langle A \wedge dA + \frac{2}{3} A \wedge A \wedge A \right\rangle, \tag{4.1}$$

⁵All our integrals are over three-dimensional space-time.

where the gauge algebra is given by (2.24) and (2.35). In order to carry out this construction we shall construct invariant tensors of these algebras in the following way: in 2 + 1 dimensions, an invariant tensor for the expanded algebras of interest can be obtained following [7], by first defining the structure constants $K^\gamma_{\alpha\beta}$ of the semigroup (2.19) as

$$\lambda_\alpha \lambda_\beta = K^\gamma_{\alpha\beta} \lambda_\gamma. \quad (4.2)$$

Given the invariant tensor on $\mathfrak{so}(2, 2)$,

$$\langle \tilde{J}_{AB} \tilde{P}_C \rangle = \epsilon_{ABC} \quad (\epsilon_{012} = -1), \quad (4.3)$$

the invariant tensor on the expanded algebra can be defined as

$$\langle (\lambda_\alpha \otimes \tilde{J}_{AB}) (\lambda_\beta \otimes \tilde{P}_C) \rangle = \alpha_{(\gamma)} K^\gamma_{\alpha\beta} \epsilon_{ABC}, \quad (4.4)$$

where we have introduced an infinite set of arbitrary constants $\alpha_{(\gamma)}$.

4.1 Extended Newton–Hooke gravities

Now we will consider the infinite-dimensional extended Newton–Hooke algebra (2.24), where the definition (2.23) of the expanded generators leads to

$$\langle B_a^{(m)} P_b^{(n)} \rangle = -\alpha_{(2m+2n+2)} \epsilon_{ab}, \quad \langle J_{ab}^{(m)} H^{(n)} \rangle = -\alpha_{(2m+2n)} \epsilon_{ab}. \quad (4.5)$$

Therefore, as the constants $\alpha_{(\gamma)}$ in (4.5) are non-vanishing only for even values of γ , it is convenient to relabel them in terms of a set of constants $\mu_{(m)} = \alpha_{(2m)}$. Finally, using (2.23) and dualising J_{ab} and B_a in the form

$$J^{(m)} \equiv \frac{1}{2} \epsilon^{ab} J_{ab}^{(m)}, \quad G_a^{(m)} \equiv \epsilon_a^b B_b^{(m)}, \quad (4.6)$$

($a, b = 1, 2$) yields

$$\langle G_a^{(m)} P_b^{(n)} \rangle = \mu_{(m+n+1)} \delta_{ab}, \quad \langle J^{(m)} H^{(n)} \rangle = -\mu_{(m+n)}. \quad (4.7)$$

In terms of the dual generators (4.6), the infinite-dimensional graded algebra (2.24) takes the form

$$[J^{(m)}, P_a^{(n)}] = -\epsilon_{ab} P_b^{(m+n)}, \quad (4.8a) \quad [H^{(m)}, G_a^{(n)}] = -\epsilon_{ab} P_b^{(m+n)}, \quad (4.8e)$$

$$[G_a^{(m)}, G_b^{(n)}] = \epsilon_{ab} J^{(m+n+1)}, \quad (4.8b) \quad [H^{(m)}, P_a^{(n)}] = \Lambda \epsilon_{ab} G_b^{(m+n)}, \quad (4.8f)$$

$$[J^{(m)}, G_a^{(n)}] = -\epsilon_{ab} G_b^{(m+n)}, \quad (4.8c) \quad [P_a^{(m)}, P_b^{(n)}] = -\Lambda \epsilon_{ab} J^{(m+n+1)}. \quad (4.8g)$$

$$[G_a^{(m)}, P_b^{(n)}] = \epsilon_{ab} H^{(m+n+1)}, \quad (4.8d)$$

Action

In order to construct a Chern–Simons action, we define a connection one-form taking values on (4.8)

$$A = \sum_{m=0}^{\infty} \left(e_{(m)}^a P_a^{(m)} + \omega_{(m)}^a G_a^{(m)} + \tau_{(m)} H^{(m)} + \omega_{(m)} J^{(m)} \right). \quad (4.9)$$

The curvature 2-form $F = dA + A \wedge A$ then reads

$$F = \sum_{m=0}^{\infty} \left(F^a [P^{(m)}] P_a^{(m)} + F^a [G^{(m)}] G_a^{(m)} + F [H^{(m)}] H^{(m)} + F [J^{(m)}] J^{(m)} \right). \quad (4.10)$$

where the curvature components can be worked out using the algebra (4.8) and read explicitly

$$\begin{aligned} F^a [P^{(m)}] &= de_{(m)}^a - \sum_{n,p=0}^{\infty} \delta_m^{n+p} \epsilon_b^a \left(\omega_{(n)}^b \tau_{(p)} + e_{(n)}^b \omega_{(p)} \right) \\ F^a [G^{(m)}] &= d\omega_{(m)}^a - \sum_{n,p=0}^{\infty} \delta_m^{n+p} \epsilon_b^a \left(\omega_{(n)}^b \omega_{(p)} - \Lambda e_{(n)}^b \tau_{(p)} \right), \\ F [H^{(m)}] &= d\tau_{(m)} + \sum_{n,p=0}^{\infty} \delta_m^{n+p+1} \epsilon_{ab} e_{(n)}^a \omega_{(p)}^b, \\ F [J^{(m)}] &= d\omega_{(m)} + \frac{1}{2} \sum_{n,p=0}^{\infty} \delta_m^{n+p+1} \epsilon_{ab} \left(\omega_{(n)}^a \omega_{(p)}^b - \Lambda e_{(n)}^a e_{(p)}^b \right). \end{aligned} \quad (4.11)$$

Here and from now on the wedge product between forms are not written explicitly. Using the gauge connection (4.9) and the invariant bilinear form (4.7), the action (4.1) takes the form

$$\begin{aligned} S_{\text{CS}} &= \sum_{m,n=0}^{\infty} \mu_{(m+n+1)} \int \left(e_{(m)}^a F_a [G^{(n)}] + \omega_{(m)}^a F_a [P^{(n)}] \right) \\ &\quad - \sum_{m,n=0}^{\infty} \mu_{(m+n)} \int \left(\tau_{(m)} F [J^{(n)}] + \omega_{(m)} F [H^{(n)}] \right) \\ &\quad + \sum_{m,n,p=0}^{\infty} \mu_{(m+n+p+1)} \int \epsilon_{ab} \left(e_{(m)}^a \omega_{(n)}^b \omega_{(p)} + \frac{1}{2} \left(\omega_{(m)}^a \omega_{(n)}^b - \Lambda e_{(m)}^a e_{(n)}^b \right) \tau_{(p)} \right), \end{aligned} \quad (4.12)$$

where the different terms can be rearranged in terms of $\mu_{(n)}$ as

$$S_{\text{CS}} = \sum_{i=0}^{\infty} \mu_{(i)} S_{(i)}. \quad (4.13)$$

We shall now discuss in more detail the first three terms in the sum.

- The $\mu_{(0)}$ term can be directly read off from the second sum in (4.12). Up to total derivatives it reads

$$S_{(0)} = -2 \int \tau_{(0)} d\omega_{(0)}. \quad (4.14)$$

which corresponds to Galilean gravity in 2+1 dimensions [49].

- In order to write down the $\mu_{(1)}$ term in a familiar way, we first note that we can relate the curvatures associated to the different generators (4.11) in the form

$$\begin{aligned}
\omega_{(0)}^a F_a \left[P^{(0)} \right] &= e_{(0)}^a F_a \left[G^{(0)} \right] - \epsilon_{ab} \left(\omega_{(0)}^a \omega_{(0)}^b + \Lambda e_{(0)}^a e_{(0)}^b \right) \tau_{(0)} + \text{T.D.}, \\
\omega_{(1)} F \left[H^{(0)} \right] &= \tau_{(0)} F \left[J^{(1)} \right] - \frac{1}{2} \epsilon_{ab} \left(\omega_{(0)}^a \omega_{(0)}^b - \Lambda e_{(0)}^a e_{(0)}^b \right) \tau_{(0)} + \text{T.D.}, \\
\omega_{(0)} F \left[H^{(1)} \right] &= \tau_{(1)} F \left[J^{(0)} \right] + \epsilon_{ab} e_{(0)}^a \omega_{(0)}^b \omega_{(0)} + \text{T.D.},
\end{aligned} \tag{4.15}$$

where T.D. stands for total derivative terms. Using these relations, the action $S_{(1)}$ takes the form

$$S_{(1)} = 2 \int \left(e_{(0)}^a F_a \left[G^{(0)} \right] - \tau_{(0)} F \left[J^{(1)} \right] - \tau_{(1)} F \left[J^{(0)} \right] - \Lambda \epsilon_{ab} e_{(0)}^a e_{(0)}^b \tau_{(0)} \right). \tag{4.16}$$

This action defines extended Newton–Hooke gravity [25, 26] and generalises extended Bargmann gravity [27, 28] to include the cosmological constant. This can be seen clearly by relabeling the gauge fields as

$$\begin{aligned}
e_{(0)}^a &= e^a, & \tau_{(0)} &= \tau, & \tau_{(1)} &= m, \\
\omega_{(0)}^a &= \omega^a, & \omega_{(0)} &= \omega, & \omega_{(1)} &= s.
\end{aligned} \tag{4.17}$$

- For the $\mu_{(2)}$ term we use the relations

$$\begin{aligned}
\omega_{(0)}^a F_a \left[P^{(1)} \right] &= e_{(1)}^a F_a \left[G^{(0)} \right] - \epsilon_{ab} \left(\omega_{(0)}^a \omega_{(0)}^b \tau_{(1)} + \omega_{(0)}^a \omega_{(1)}^b \tau_{(0)} + \omega_{(0)}^a e_{(0)}^b \omega_{(1)} \right) + \text{T.D.}, \\
\omega_{(1)}^a F_a \left[P^{(0)} \right] &= e_{(0)}^a F_a \left[G^{(1)} \right] - \epsilon_{ab} \left(\omega_{(1)}^a \omega_{(0)}^b \tau_{(0)} - \omega_{(0)}^a e_{(0)}^b \omega_{(1)} \right) + \text{T.D.}, \\
\omega_{(0)} F \left[H^{(2)} \right] &= \tau_{(2)} F \left[J^{(0)} \right] + \epsilon_{ab} \left(e_{(0)}^a \omega_{(1)}^b \omega_{(0)} + e_{(1)}^a \omega_{(0)}^b \omega_{(0)} \right) + \text{T.D.}, \\
\omega_{(1)} F \left[H^{(1)} \right] &= \tau_{(1)} F \left[J^{(1)} \right] + \epsilon_{ab} \left(e_{(0)}^a \omega_{(0)}^b \omega_{(1)} - \frac{1}{2} \omega_{(0)}^a \omega_{(0)}^b \tau_{(1)} \right) + \text{T.D.}, \\
\omega_{(2)} F \left[H^{(0)} \right] &= \tau_{(0)} F \left[J^{(2)} \right] - \epsilon_{ab} \omega_{(0)}^a \omega_{(1)}^b \tau_{(0)} + \text{T.D.},
\end{aligned} \tag{4.18}$$

which allows one to express $S_{(2)}$ as

$$\begin{aligned}
S_{(2)} &= 2 \int \left(e_{(0)}^a F_a \left[G^{(1)} \right] + e_{(1)}^a F_a \left[G^{(0)} \right] \right. \\
&\quad \left. - \tau_{(0)} F \left[J^{(2)} \right] - \tau_{(1)} F \left[J^{(1)} \right] - \tau_{(2)} F \left[J^{(0)} \right] \right. \\
&\quad \left. - \Lambda \epsilon_{ab} \left(e_{(0)}^a e_{(0)}^b \tau_{(1)} + e_{(0)}^a e_{(1)}^b \tau_{(0)} + e_{(1)}^a e_{(0)}^b \tau_{(0)} \right) \right).
\end{aligned} \tag{4.19}$$

For $\Lambda = 0$, this is precisely the Lagrangian for non-relativistic three-dimensional gravity found in [11], where the gauge fields should be identified as in (4.20) together with

$$\begin{aligned}
e_{(1)}^a &= t^a, & \tau_{(2)} &= y, \\
\omega_{(1)}^a &= b^a, & \omega_{(2)} &= z.
\end{aligned} \tag{4.20}$$

In general the action $S_{(i)}$ can be written, after partial integration, in the form

$$S_{(i)} = 2 \int \left(\sum_{m,n=0}^{\infty} \left(\delta_i^{m+n+1} e_{(m)}^a F_a \left[G^{(n)} \right] + \delta_i^{m+n} \tau_{(m)} F \left[J^{(n)} \right] \right) - \Lambda \sum_{m,n,p=0}^{\infty} \delta_i^{m+n+p+1} \epsilon_{ab} e_{(m)}^a e_{(n)}^b \tau_{(p)} \right) \quad (4.21)$$

It is important to note that the infinite sum (4.13) can be consistently truncated to give

$$S_{\text{CS}} = \sum_{i=0}^{i_0+1} \mu_{(i)} S_i, \quad (4.22)$$

which gives an action for the gauge fields

$$e_{(m)}^a, \omega_{(m)}^a, \quad \text{for } m = 0, 1, \dots, i_0, \quad \tau_{(n)}, \omega_{(n)}, \quad \text{for } n = 0, 1, \dots, i_0 + 1, \quad (4.23)$$

and sets all the other gauge fields to zero. In this case, the invariant tensor (4.7) is non-degenerate for $\mu_{(i_0+1)} \neq 0$, and the truncation leads to the condition

$$\mu_{(m)} = 0, \quad \text{for } m > i_0 + 1. \quad (4.24)$$

This corresponds to the Chern–Simons action for non-relativistic gravity invariant under the expanded algebra (2.9) for $N = 2(i_0 + 1)$,

$$\left(s_0^{(2(i_0+1))} \times \{ \tilde{H}, \tilde{J} \} \right) \oplus \left(s_1^{(2(i_0+1))} \times \{ \tilde{P}_a, \tilde{G}_b \} \right). \quad (4.25)$$

This means that only the expanded algebras with even values of N give rise to Chern–Simons actions with a non-degenerate invariant bilinear form.

AdS case

Let us consider now the general non-relativistic algebra (4.8) in the case of negative cosmological constant and set $\Lambda = -\ell^{-2} < 0$. In this case it takes the form

$$[J^{(m)}, P_a^{(n)}] = -\epsilon_{ab} P_b^{(m+n)}, \quad (4.26a) \quad [H^{(m)}, G_a^{(n)}] = -\epsilon_{ab} P_b^{(m+n)}, \quad (4.26e)$$

$$[G_a^{(m)}, G_b^{(n)}] = \epsilon_{ab} J^{(m+n+1)}, \quad (4.26b) \quad [H^{(m)}, P_a^{(n)}] = -\frac{1}{\ell^2} \epsilon_{ab} G_b^{(m+n)}, \quad (4.26f)$$

$$[J^{(m)}, G_a^{(n)}] = -\epsilon_{ab} G_b^{(m+n)}, \quad (4.26c) \quad [P_a^{(m)}, P_b^{(n)}] = \frac{1}{\ell^2} \epsilon_{ab} J^{(m+n+1)}. \quad (4.26g)$$

$$[G_a^{(m)}, P_b^{(n)}] = \epsilon_{ab} H^{(m+n+1)}, \quad (4.26d)$$

By defining the change of basis

$$L_a^{\pm(m)} = \frac{1}{2} \left(G_a^{(m)} \pm \ell P_a^{(m)} \right), \quad L^{\pm(m)} = \frac{1}{2} \left(J^{(m)} \pm \ell H^{(m)} \right), \quad (4.27)$$

this algebra can be written as the direct sum $\{L_a^{+(m)}, L^{+(m)}\} \oplus \{L_a^{-(m)}, L^{-(m)}\}$, where

$$[L^{\pm(m)}, L_a^{\pm(n)}] = -\epsilon_a{}^b L_b^{\pm(m+n)}, \quad [L_a^{\pm(m)}, L_b^{\pm(n)}] = \epsilon_a{}^b L^{\pm(m+n+1)}. \quad (4.28)$$

The invariant tensor in this basis follows from (4.7) and has takes the form

$$\langle L_a^{\pm(m)} L_b^{\pm(n)} \rangle = \pm \frac{\ell}{2} \mu_{(m+n+1)} \delta_{ab}, \quad \langle L^{\pm(p)} L^{\pm(q)} \rangle = \mp \frac{\ell}{2} \mu_{(p+q)}. \quad (4.29)$$

This isomorphism implies that one can reformulate the expanded Newton–Hooke gravity action (4.12) in the form

$$S_{\text{CS}}[A] = S_{\text{CS}}[A^+] - S_{\text{CS}}[A^-], \quad (4.30)$$

where the gauge connections are given by can consider two connections of the form

$$A^\pm = \sum_m \left(\omega_{(m)}^a \pm \frac{1}{\ell} e_{(m)}^a \right) L_a^{(m)} + \sum_p \left(\omega_{(p)} \pm \frac{1}{\ell} \tau_{(p)} \right) L^{(p)}, \quad (4.31)$$

and the $\{L_a^{(m)}, L^{(m)}\}$ is now a single set of generators satisfying

$$[L^{(m)}, L_a^{(p)}] = -\epsilon_a{}^b L_b^{(m+p)}, \quad [L_a^{(m)}, L_b^{(n)}] = \epsilon_a{}^b L^{(m+n+1)}, \quad (4.32)$$

$$\langle L_a^{(m)} L_b^{(n)} \rangle = \frac{1}{2} \gamma_{(m+n+1)} \delta_{ab}, \quad \langle L^{(p)} L^{(q)} \rangle = \frac{1}{2} \gamma_{(p+q)}. \quad (4.33)$$

This is in completely analogy with the relativistic case, where three-dimensional Einstein gravity with negative cosmological constant can be reformulated as a Chern–Simons theory of the form (4.30) with chiral connections taking values in the $\mathfrak{sl}(2, \mathbb{R})$ algebra [50–54]. In fact, the algebra (4.32) can be obtained as a non-relativistic $S^{(\infty)}$ expansion of $\mathfrak{so}(2, 1) \approx \mathfrak{sl}(2, \mathbb{R})$ by defining

$$L^{(m)} = \lambda_{2m} \otimes \tilde{L}, \quad L_a^{(m)} = \lambda_{2m+1} \otimes \tilde{L}_a, \quad (4.34)$$

where $\tilde{L}_A = \{\tilde{L} \equiv \tilde{L}_0, \tilde{L}_a\}$ are the generators of the Lorentz algebra in 2+1 dimensions and satisfy $[L_A, L_B] = \epsilon^C{}_{AB} \tilde{L}_C$. Once again, quotients by suitable ideals reproduce $S_E^{(N)}$ expansions. The case $N = 1$ shows that the Newton–Hooke algebra in 2+1 dimensions is isomorphic to two copies of the Euclidean algebra in 1+1 dimensions, while the $N = 2$ case expresses the extended Newton–Hooke symmetry in three-dimensions as two copies of the centrally extended Newton–Hooke symmetry in two dimensions [45], isomorphic to the Nappi–Witten algebra [55, 16].

Exotic invariant bilinear form

As $\mathfrak{so}(2, 2)$ is semi-simple there are two independent invariant bilinear forms. Besides (4.3), a second invariant bilinear form on $\mathfrak{so}(2, 2)$ is

$$\langle J_{AB} J_{CD} \rangle = \eta_{AD} \eta_{BC} - \eta_{AC} \eta_{BD}, \quad \langle P_A P_B \rangle = -\Lambda \eta_{AB}, \quad (4.35)$$

which induces the following invariant tensor for the expanded generators (2.23)

$$\langle J_{ab}^{(m)} J_{cb}^{(n)} \rangle = \beta_{(2m+2n)} (\delta_{ad}\delta_{bc} - \delta_{ac}\delta_{bd}), \quad \langle B_a^{(m)} B_b^{(n)} \rangle = \beta_{(2m+2n+2)} \delta_{ab}, \quad (4.36a)$$

$$\langle H^{(m)} H^{(n)} \rangle = \Lambda \beta_{(2m+2n)}, \quad \langle P_a^{(m)} P_b^{(n)} \rangle = -\Lambda \beta_{(2m+2n+2)} \delta_{ab}. \quad (4.36b)$$

Defining $\beta_{(2m)} = \nu_{(m)}$ and using (4.6) this leads to

$$\langle J^{(p)} J^{(q)} \rangle = -\nu_{(p+q)}, \quad \langle G_a^{(m)} G_b^{(n)} \rangle = \nu_{(m+n+1)} \delta_{ab}, \quad (4.37a)$$

$$\langle H^{(p)} H^{(q)} \rangle = \Lambda \nu_{(p+q)}, \quad \langle P_a^{(m)} P_b^{(n)} \rangle = -\Lambda \nu_{(m+n+1)} \delta_{ab}. \quad (4.37b)$$

In this case, the action (4.1) boils down to

$$\begin{aligned} S_{\text{CS}} &= \sum_{m,n=0}^{\infty} \nu_{(m+n+1)} \int \left(\omega_{(m)}^a F_a [G^{(n)}] - \Lambda e_{(m)}^a F_a [P^{(n)}] \right) \\ &\quad - \sum_{m,n=0}^{\infty} \nu_{(m+n)} \int \left(\omega_{(m)} F [J^{(n)}] - \Lambda \tau_{(p)} F [H^{(q)}] \right) \\ &\quad + \sum_{m,n,p=0}^{\infty} \nu_{(m+n+p+1)} \int \epsilon_{ab} \left(-\Lambda e_{(m)}^a \omega_{(n)}^b \tau_{(p)} + \frac{1}{2} \left(\omega_{(m)}^a \omega_{(n)}^b - \Lambda e_{(m)}^a e_{(n)}^b \right) \omega_{(p)} \right). \end{aligned} \quad (4.38)$$

The exotic invariant tensor in the chiral basis (4.27) takes the form

$$\langle L_a^{\pm(m)} L_b^{\pm(n)} \rangle = \frac{1}{2} \nu_{(m+n+1)} \delta_{ab}, \quad \langle L^{(p)} L^{(q)} \rangle = -\frac{1}{2} \nu_{(p+q)}. \quad (4.39)$$

Therefore, the action (4.38) can be reformulated in terms of (4.31) as

$$S_{\text{CS}}[A] = S_{\text{CS}}[A^+] + S_{\text{CS}}[A^-], \quad (4.40)$$

where now the relative sign between the two Chern–Simons actions is changed compared to (4.30).

4.2 Extended Carrollian gravities

In the Carrollian case, the definition of the expanded generators with algebra given in (2.35) together with (4.4) yields the following invariant tensor

$$\langle B_a^{(m)} P_b^{(n)} \rangle = -\alpha_{(2m+2n+1)} \epsilon_{ab}, \quad \langle J_{ab}^{(m)} H^{(n)} \rangle = -\alpha_{(2m+2n+1)} \epsilon_{ab}. \quad (4.41)$$

Thus, in contrast with the Newton–Hooke case, the constants $\alpha_{(\gamma)}$ in (4.5) are non-vanishing only for odd values of γ , and we will relabel them as $\rho_{(m)} = \alpha_{(2m+1)}$. Using (2.34) and the dual generators (4.6) the invariant bilinear form in the $D = 3$ case can also be written as

$$\langle G_a^{(m)} P_b^{(n)} \rangle = \rho_{(m+n)} \delta_{ab}, \quad \langle J^{(m)} H^{(n)} \rangle = -\rho_{(m+n)}. \quad (4.42)$$

The infinite-dimensional algebra (2.35) takes the form

$$[J^{(m)}, P_a^{(n)}] = -\epsilon_{ab} P_b^{(m+n)}, \quad (4.43a) \quad [H^{(m)}, G_a^{(n)}] = -\epsilon_{ab} P_b^{(m+n+1)}, \quad (4.43e)$$

$$[G_a^{(m)}, G_b^{(n)}] = \epsilon_{ab} J^{(m+n+1)}, \quad (4.43b) \quad [H^{(m)}, P_a^{(n)}] = \Lambda \epsilon_{ab} G_b^{(m+n)}, \quad (4.43f)$$

$$[J^{(m)}, G_a^{(n)}] = -\epsilon_{ab} G_b^{(m+n)}, \quad (4.43c) \quad [P_a^{(m)}, P_b^{(n)}] = -\Lambda \epsilon_{ab} J^{(m+n)}. \quad (4.43g)$$

$$[G_a^{(m)}, P_b^{(n)}] = \epsilon_{ab} H^{(m+n)}, \quad (4.43d)$$

Action

In the Carrollian case, the connection one-form (4.9) together with the commutation relations (4.43) yields a curvature two-form (4.10) with the following components:

$$\begin{aligned} F^a [P^{(m)}] &= de_{(m)}^a - \sum_{n,p=0}^{\infty} \epsilon_b^a \left(\delta_m^{n+p+1} \omega_{(n)}^b \tau_{(p)} - \delta_m^{(n+p)} e_{(n)}^b \omega_{(p)} \right) \\ F^a [G^{(m)}] &= d\omega_{(m)}^a - \sum_{n,p=0}^{\infty} \delta_m^{n+p} \epsilon_b^a \left(\omega_{(n)}^b \omega_{(p)} - \Lambda e_{(p)}^b \tau_{(n)} \right), \\ F [H^{(m)}] &= d\tau_{(m)} + \sum_{n,p=0}^{\infty} \delta_m^{n+p} \epsilon_{ab} e_{(p)}^a \omega_{(n)}^b, \\ F [J^{(m)}] &= d\omega_{(m)} + \frac{1}{2} \sum_{n,p=0}^{\infty} \epsilon_{ab} \left(\delta_m^{n+p+1} \omega_{(n)}^a \omega_{(p)}^b - \Lambda \delta_m^{n+p} e_{(n)}^a e_{(p)}^b \right). \end{aligned} \quad (4.44)$$

The Chern–Simons action (4.1) then takes the form

$$\begin{aligned} S_{\text{CS}} &= \sum_{m,n=0}^{\infty} \rho_{(m+n)} \int \left(e_{(m)}^a F_a [G^{(n)}] + \omega_{(m)}^a F_a [P^{(n)}] - \tau_{(m)} F [J^{(n)}] - \omega_{(m)} F [H^{(n)}] \right) \\ &+ \sum_{m,n,p=0}^{\infty} \int \epsilon_{ab} \left(\rho_{(m+n+p)} \left(e_{(m)}^a \omega_{(n)}^b \omega_{(p)} - \frac{\Lambda}{2} e_{(m)}^a e_{(n)}^b \tau_{(p)} \right) + \frac{1}{2} \rho_{(m+n+p+1)} \omega_{(m)}^a \omega_{(n)}^b \tau_{(p)} \right), \end{aligned} \quad (4.45)$$

which can be also written as

$$S_{\text{CS}} = \sum_{i=0}^{\infty} \rho_{(i)} S_{(i)}, \quad (4.46)$$

where after integration by parts we can write $S_{(i)}$ as

$$S_{(i)} = 2 \int \left(\sum_{m,n=0}^{\infty} \delta_i^{m+n} \left(e_{(m)}^a F_a [G^{(n)}] - \tau_{(m)} F [J^{(n)}] \right) - \Lambda \sum_{m,n,p=0}^{\infty} \delta_i^{m+n+p} \epsilon_{ab} e_{(m)}^a e_{(n)}^b \tau_{(p)} \right). \quad (4.47)$$

The first action in this sequence is given by

$$S_{(0)} = 2 \int \left(e^a F_a [G] - \tau F [J] - \Lambda \epsilon_{ab} e^a e^b \tau \right), \quad (4.48)$$

where we have used the field redefinition (4.17). This corresponds to Carrollian AdS Chern-Simons gravity [39, 56] and reduces to three-dimensional Carroll gravity [57, 58, 43] in the vanishing cosmological constant limit.⁶

Using also (4.20), the second term of the sum reads

$$S_{(1)} = 2 \int \left(e^a F_a [B] + t^a F_a [G] - \tau F [S] - m F [J] - \Lambda \epsilon_{ab} \left(e^a e^b m + 2e^a t^b \tau \right) \right) \quad (4.49)$$

and defines a Carrollian counterpart of (post-)Newtonian gravity action [9] in three dimensions including cosmological constant. Similarly, the next actions in the sequence define further post-Carrollian corrections for three-dimensional Carroll (A)dS gravity.

It is interesting that unlike the Newton–Hooke expansion, truncations of the general Carrollian action (4.46) for arbitrary r lead to a non-degenerate invariant bilinear form for the corresponding truncation in (4.43). Also, the extended Carrollian algebras for (A)dS do not include central terms, which is in contrast with the Newton–Hooke case, where the central extensions are precisely the ones that allow us to find non-generate pairings when (4.22) holds.

Exotic invariant bilinear form

We can also consider the exotic invariant tensor for (A)dS (4.35), which leads to the following expression for pairings of the expanded generators (2.34)

$$\left\langle J_{ab}^{(m)} J_{cb}^{(n)} \right\rangle = \beta_{(2m+2n)} (\delta_{ad} \delta_{bc} - \delta_{ac} \delta_{bd}) , \quad \left\langle B_a^{(m)} B_b^{(n)} \right\rangle = \beta_{(2m+2n+2)} \delta_{ab} , \quad (4.50a)$$

$$\left\langle H^{(m)} H^{(n)} \right\rangle = \Lambda \beta_{(2m+2n+2)} , \quad \left\langle P_a^{(m)} P_b^{(n)} \right\rangle = -\Lambda \beta_{(2m+2n)} \delta_{ab} . \quad (4.50b)$$

Defining $\beta_{(2m)} = \sigma_{(m)}$ and using (4.6) this can also be written as

$$\left\langle J^{(m)} J^{(n)} \right\rangle = -\sigma_{(m+n)} , \quad \left\langle G_a^{(m)} G_b^{(n)} \right\rangle = \sigma_{(m+n+1)} \delta_{ab} , \quad (4.51a)$$

$$\left\langle H^{(m)} H^{(n)} \right\rangle = \Lambda \sigma_{(m+n+1)} , \quad \left\langle P_a^{(m)} P_b^{(n)} \right\rangle = -\Lambda \sigma_{(m+n)} \delta_{ab} . \quad (4.51b)$$

⁶The first realisations of the Carroll symmetry in gravity appeared in the study of the zero signature limit [59, 60] or strong coupling limit [61, 62] (see also [63, 64]) of general relativity, the last one being closely related to the Belinski–Khalatnikov–Lifshitz (BKL) limit of gravity [65–67]. On the other hand, the ultra-relativistic expansion of general relativity has been explored in [68], while the Carrollian limit at the level of the Einstein–Hilbert action has been studied in [49]. It is also important to mention the realisation of the Carroll symmetry in the near horizon limit of black holes [69].

In this case, the action (4.1) reduces to

$$\begin{aligned}
S_{\text{CS}} = & \sum_{m,n=0}^{\infty} \int \left(\sigma_{(m+n+1)} \left(\omega_{(m)}^a F_a \left[G^{(n)} \right] - \Lambda \tau_{(m)} F \left[H^{(n)} \right] \right) \right. \\
& \left. - \sigma_{(m+n)} \left(\omega_{(m)} F \left[J^{(n)} \right] + \Lambda e_{(m)}^a F_a \left[P^{(n)} \right] \right) \right) \\
& + \sum_{m,n,p=0}^{\infty} \epsilon_{ab} \int \left(\sigma_{(m+n+p+1)} \left[-\Lambda e_{(m)}^a \omega_{(n)}^b \tau_{(p)} + \frac{1}{2} \omega_{(m)}^a \omega_{(n)}^b \omega_{(p)} \right] \right. \\
& \left. - \frac{\Lambda}{2} \sigma_{(m+n+p)} e_{(m)}^a e_{(n)}^b \omega_{(p)} \right). \tag{4.52}
\end{aligned}$$

One can consider again finite truncations of this actions and, which produce exotic Carroll and post-Carrollian gravity theories in $2 + 1$ dimensions.

5 Conclusions and Outlook

We have studied the non-relativistic symmetries obtained by Lie algebra expansion of the AdS or dS algebra in D space-time dimensions, i.e. $\mathfrak{so}(D-1, 2)$ or $\mathfrak{so}(D, 1)$. This generalises previous constructions to include a cosmological constant and generates an infinite family of algebras of Newton–Hooke [19, 20] or Carrollian type. Subsequently we have shown how these infinite-dimensional symmetries can be embedded in different Kac-Moody algebras. Moreover, the different finite $S_E^{(N)}$ expansions contained in each family can be systematically constructed as a free algebra and correspond to suitable quotients of the infinite-dimensional case.

We have considered a gravitational model based on them by focussing on the case of $(2+1)$ -dimensional Chern–Simons theories. We show that the family of algebras generates systematically non-relativistic gravity theories extended by a cosmological constant, such as extended Bargmann gravity [28, 26] and (post-)Newtonian gravity [11].

As stated in [10] these extended Newton–Hooke symmetries encode the large c expansion of general relativity. An interesting way to see this in $2+1$ dimensions is that the Chern–Simons formulation of AdS three-dimensional gravity allows to construct the metric out of the chiral connections (4.31) in the form [70]

$$ds^2 = g_{\mu\nu} dx^\mu dx^\nu = \frac{\ell^2}{2} \langle (A^+ - A^-) \otimes (A^+ - A^-) \rangle. \tag{5.1}$$

Using (4.31) and rearranging the sum in terms of $\gamma_{(n)}$ we get

$$ds^2 = - \sum_{p,q} \gamma_{(m+n)} \tau_{(m)} \tau_{(n)} + \sum_{m,n} \gamma_{(m+n+1)} e_{(m)}^a e_{(n)}^b. \tag{5.2}$$

The flat version of this metric has been studied in [12] as an infinite extension of Minkowski space where post-Newtonian corrections to relativistic symmetries can be naturally implemented. This metric can be found in the vanishing cosmological constant limit by defining coordinates $x_{(m)}^\mu$

such that $\tau_{(m)} = \tau_{(m)\mu} dx_{(m)}^\mu$ and $e_{(m)}^a = e_{(m)\mu}^a dx_{(m)}^\mu$. Then, the Minkowskian case is obtained by setting $e_{(m)\mu}^a = \delta_\mu^a$ and $\tau_{(m)\mu} = (1, 0, 0, 0)$ for all m , which leads to the expanded metric constructed in [12].

On the other hand, keeping the gauge field components $\tau_{(m)\mu}$ and $e_{(m)\mu}^a$ arbitrary while identifying the coordinates as $x_{(m)}^\mu = x^\mu$, allows one to find the (2+1)-dimensional version of the post-Newtonian expansion of the metric used in [29–32, 10, 33]. Indeed, identifying the invariant tensor constants as powers of the speed of light in the form $\gamma_{(n)} \rightarrow c^{2(1-n)}$, and defining the spatial metrics $h_{\mu\nu} dx^\mu dx^\nu = e_{(0)}^a e_{a(0)}$ and $\Phi_{\mu\nu} dx^\mu dx^\nu = e_{(0)}^a e_{a(1)} + e_{(1)}^a e_{a(0)}$, this expression reproduces the aforementioned non-relativistic expansion:

$$g_{\mu\nu} = -c^2 \tau_\mu \tau_\nu + h_{\mu\nu} - \tau_\mu m_\nu - m_\mu \tau_\nu + c^{-2} (\Phi_{\mu\nu} - m_\mu m_\nu - \tau_\mu y_\nu - y_\mu \tau_\nu) + O(c^{-4}), \quad (5.3)$$

where we have also relabeled the metric fields according to (4.17) and (4.20). Alternatively one can introduce an expansion of the dreibein forms

$$E^0 = \sum_m \lambda_{2m} \tau_{(m)}, \quad E^a = \sum_m \lambda_{2m+1} e_{(m)}^a, \quad (5.4)$$

similar to what was done in [8]. Then, by identifying the tangent space metric with the exotic pairing for the translations (4.35), one can write [71]

$$g_{\mu\nu} = \eta_{AB} E^A E^B = \ell^2 \langle E^A P_A E^B P_B \rangle, \quad (5.5)$$

which also leads to the expansion of the metric (5.2) with $\gamma_{(n)} = \nu_{(n)}$, where $\nu_{(n)}$ appears in the pairing of the momentum generators in (4.37b).

We have also considered the Carrollian case with cosmological constant to construct a corresponding Chern–Simons theory based on the algebra (2.35). This reproduces known ultra-relativistic gravity theories in 2 + 1 dimensions and an infinite family of generalisations.

Applying the same procedure as (5.2) in the the case of expanded Carrollian (A)dS algebras (4.43) leads to a novel expansion for the metric tensor that reads

$$\begin{aligned} g_{\mu\nu} &= - \sum_{m,n} \sigma_{(m+n+1)} \tau_{(m)\mu} \tau_{(n)\nu} + \sum_{m,n} \sigma_{(m+n)} e_{(m)\mu}^a e_{(n)\nu}^b \\ &= \sigma_{(0)} h_{\mu\nu} + \sigma_{(1)} (\Phi_{\mu\nu} - \tau_\mu \tau_\nu) + \sigma_{(2)} (\Psi_{\mu\nu} - \tau_\mu m_\nu - m_\mu \tau_\nu) + \dots \end{aligned} \quad (5.6)$$

where we have also defined $\Psi_{\mu\nu} dx^\mu dx^\nu = e_{(1)}^a e_{a(1)} + e_{(0)}^a e_{a(2)} + e_{(2)}^a e_{a(0)}$. This metric can be conjectured to describe the ultra-relativistic expansion of three-dimensional Einstein gravity. To evaluate whether this expansion can be generalised to higher dimensions in the context of the Carroll limit of general relativity along the lines of [59] or [68] is an interesting question that we hope to address in the future.

There are several directions in which this research could be extended. One would be to generalise our work of symmetries of post-Newtonian correction in the flat Minkowski space [12] to curved (A)dS space and construct particle actions in the curved generalisation of the infinite-dimensional Minkowski space. On the other hand it would be interesting to analyse particle systems in the presence of a constant background field by considering the non-relativistic expansions of the Maxwell algebra according to Appendix A.1. One could also construct actions for extended objects by means of the p -brane symmetries outlined in Appendix A.2. Finally, it would be interesting to extend the analysis to the supersymmetric (A)dS case.

Acknowledgements

We acknowledge discussions with Luis Avilés, Eric Bergshoeff, Diego Hidalgo, Diederik Roest, Patricio Salgado and Jorge Zanelli. JG acknowledges the hospitality and support of the Van Swinderen Institute. JG also has been supported in part by MINECO FPA2016-76005-C2-1-P and Consolider CPAN, and by the Spanish government (MINECO/FEDER) under project MDM-2014-0369 of ICCUB (Unidad de Excelencia Marà de Maeztu). PS-R acknowledges DI-VRIEA for financial support through Proyecto Postdoctorado 2019 VRIEA-PUCV.

A Further generalisations

The procedure for obtaining non-relativistic expansions of a given relativistic algebra can be generalised to several other interesting cases. Here, we apply the general scheme to obtain non-relativistic Maxwell algebras and non-relativistic brane symmetries.

A.1 Non-relativistic expansions of the Maxwell algebra

The relativistic Maxwell algebra in D space-time dimensions is given is given by the following extension of the Poincaré algebra [72] (see also [73, 48, 74] and references therein)

$$[\tilde{J}_{AB}, \tilde{P}_C] = 2\eta_{C[B}\tilde{P}_{A]}, \quad (\text{A.1a}) \quad [\tilde{J}_{AB}, \tilde{Z}_{CD}] = 4\eta_{[A[C}\tilde{Z}_{D]B]}, \quad (\text{A.1c})$$

$$[\tilde{J}_{AB}, \tilde{J}_{CD}] = 4\eta_{[A[C}\tilde{J}_{D]B]}, \quad (\text{A.1b}) \quad [\tilde{P}_A, \tilde{P}_B] = \tilde{Z}_{AB}. \quad (\text{A.1d})$$

There is no cosmological constant present in this algebra.

As in the (A)dS case, we decompose the relativistic indices in the time and space components, $A = (0, a)$, where $a = 1, \dots, D-1$, and relabel the Lie algebra generators using (2.2) plus the corresponding relation for the new generator \tilde{Z}_{AB} given by

$$\tilde{Z}_{AB} \rightarrow \{\tilde{Z}_a \equiv \tilde{Z}_{0a}, \tilde{Z}_{ab}\}. \quad (\text{A.2})$$

In the following, we consider the infinite-dimensional semigroup $S^{(\infty)}$ given in (2.19) and derive Galilean and Carrollian expansions of the Maxwell algebra by choosing different resonant subspace decompositions.

Galilean Expansions

Galilean expansions of the Maxwell algebra are determined by the following \mathbb{Z}_2 -graded subspace decomposition

$$V_0 = \{\tilde{J}_{ab}, \tilde{H}, \tilde{Z}_{ab}\}, \quad V_1 = \{\tilde{G}_a, \tilde{P}_a, \tilde{Z}_a\}, \quad (\text{A.3})$$

and the resonant non-reduced expansion

$$\left(S_0^{(\infty)} \times \{\tilde{J}_{ab}, \tilde{H}, \tilde{Z}_{ab}\}\right) \oplus \left(S_1^{(\infty)} \times \{\tilde{G}_a, \tilde{P}_a, \tilde{Z}_a\}\right) \quad (\text{A.4})$$

where $S_0^{(\infty)}$ and $S_1^{(\infty)}$ are given in (2.21). Defining the expanded generators in the form

$$\begin{aligned} J_{ab}^{(m)} &= \lambda_{2m} \otimes \tilde{J}_{ab}, & B_a^{(m)} &= \lambda_{2m+1} \otimes \tilde{G}_a, \\ H^{(m)} &= \lambda_{2m} \otimes \tilde{H}, & P_a^{(m)} &= \lambda_{2m+1} \otimes \tilde{P}_a, \\ Z_{ab}^{(m)} &= \lambda_{2m} \otimes \tilde{Z}_{ab}, & Z_a^{(m)} &= \lambda_{2m+1} \otimes \tilde{Z}_a, \end{aligned} \quad (\text{A.5})$$

leads to the following infinite-dimensional algebra

$$[B_a^{(m)}, H^{(n)}] = P_a^{(m+n)}, \quad (\text{A.6a}) \quad [J_{ab}^{(m)}, Z_{cd}^{(n)}] = 4\delta_{[a[c}Z_{d]b}^{(m+n)}, \quad (\text{A.6g})$$

$$[B_a^{(m)}, P_b^{(n)}] = \delta_{ab}H^{(m+n+1)}, \quad (\text{A.6b}) \quad [J_{ab}^{(m)}, Z_c^{(n)}] = 2\delta_{c[b}Z_a^{(m+n)}, \quad (\text{A.6h})$$

$$[J_{ab}^{(m)}, P_c^{(n)}] = 2\delta_{c[b}P_a^{(m+n)}, \quad (\text{A.6c}) \quad [Z_{ab}^{(m)}, B_c^{(n)}] = 2\delta_{c[b}Z_a^{(m+n)}, \quad (\text{A.6i})$$

$$[B_a^{(m)}, B_b^{(n)}] = J_{ab}^{(m+n+1)}, \quad (\text{A.6d}) \quad [B_a^{(m)}, Z_b^{(n)}] = Z_{ab}^{(m+n+1)}, \quad (\text{A.6j})$$

$$[J_{ab}^{(m)}, B_c^{(n)}] = 2\delta_{c[b}B_a^{(m+n)}, \quad (\text{A.6e}) \quad [P_a^{(m)}, H^{(n)}] = -Z_a^{(m+n)}, \quad (\text{A.6k})$$

$$[J_{ab}^{(m)}, J_{cd}^{(n)}] = 4\delta_{[a[c}J_{d]b}^{(m+n)}, \quad (\text{A.6f}) \quad [P_a^{(m)}, P_b^{(n)}] = Z_{ab}^{(m+n+1)}. \quad (\text{A.6l})$$

Truncations of this infinite-dimensional algebra by different ideals lead to expansions with finite semigroups $S_E^{(N)}$ (2.6). For $N = 1$, we get the electric non-relativistic Maxwell algebra [18], while the $N = 2$ case leads to a generalisation of the exotic Maxwellian Bargmann algebra [75] for $D > 3$. For $N = 3$, the resulting algebra is a Maxwell extension of the post-Newtonian symmetry (2.24) with $\Lambda = 0$, first found in [10]. For greater values of N we obtain further post-Newtonian corrections of the electric non-relativistic Maxwell algebra.

Carrollian Expansions

Similarly, we can define Carrollian expansions of the Maxwell algebra by using the following alternative \mathbb{Z}_2 -graded subspace decomposition

$$V_0 = \{\tilde{J}_{ab}, \tilde{P}_a, \tilde{Z}_{ab}\}, \quad V_1 = \{\tilde{G}_a, \tilde{H}, \tilde{Z}_a\} \quad (\text{A.7})$$

where we have interchanged the generators H and P_a with respect to the Galilean case. This leads to the the resonant non-reduced expanded algebra

$$\left(S_0^{(\infty)} \times \{\tilde{J}_{ab}, \tilde{P}_a, \tilde{Z}_{ab}\}\right) \oplus \left(S_1^{(\infty)} \times \{\tilde{G}_a, \tilde{H}, \tilde{Z}_a\}\right) \quad (\text{A.8})$$

The expanded in this case take the form

$$\begin{aligned} J_{ab}^{(m)} &= \lambda_{2m} \otimes \tilde{J}_{ab}, & B_a^{(m)} &= \lambda_{2m+1} \otimes \tilde{G}_a \\ P_a^{(m)} &= \lambda_{2m} \otimes \tilde{P}_a, & H^{(m)} &= \lambda_{2m+1} \otimes \tilde{H} \\ Z_{ab}^{(m)} &= \lambda_{2m} \otimes \tilde{Z}_{ab}, & Z_a^{(m)} &= \lambda_{2m+1} \otimes \tilde{Z}_a, \end{aligned} \quad (\text{A.9})$$

which leads to the infinite-dimensional Carrollian Maxwell algebra

$$[B_a^{(m)}, H^{(n)}] = P_a^{(m+n+1)}, \quad (\text{A.10a}) \quad [J_{ab}^{(m)}, Z_{cd}^{(n)}] = 4\delta_{[a[c}Z_{d]b}^{(m+n)}, \quad (\text{A.10g})$$

$$[B_a^{(m)}, P_b^{(n)}] = \delta_{ab}H^{(m+n)}, \quad (\text{A.10b}) \quad [J_{ab}^{(m)}, Z_c^{(n)}] = 2\delta_{c[b}Z_a^{(m+n)}, \quad (\text{A.10h})$$

$$[J_{ab}^{(m)}, P_c^{(n)}] = 2\delta_{c[b}P_a^{(m+n)}, \quad (\text{A.10c}) \quad [Z_{ab}^{(m)}, B_c^{(n)}] = 2\delta_{c[b}Z_a^{(m+n)}, \quad (\text{A.10i})$$

$$[B_a^{(m)}, B_b^{(n)}] = J_{ab}^{(m+n+1)}, \quad (\text{A.10d}) \quad [B_a^{(m)}, Z_b^{(n)}] = Z_{ab}^{(m+n+1)}, \quad (\text{A.10j})$$

$$[J_{ab}^{(m)}, B_c^{(n)}] = 2\delta_{c[b}B_a^{(m+n)}, \quad (\text{A.10e}) \quad [P_a^{(m)}, H^{(n)}] = -Z_a^{(m+n)}, \quad (\text{A.10k})$$

$$[J_{ab}^{(m)}, J_{cd}^{(n)}] = 4\delta_{[a[c}J_{d]b}^{(m+n)}, \quad (\text{A.10f}) \quad [P_a^{(m)}, P_b^{(n)}] = Z_{ab}^{(m+n)}. \quad (\text{A.10l})$$

As before, quotients of this algebra by suitable ideals reproduce the $S_E^{(N)}$ expansions, which define Maxwell extensions of the algebras presented in Section 2.2 for $\Lambda = 0$.

A.2 Non-relativistic brane expansions of (A)dS

In order to define non-relativistic p -brane expansions of the (A)dS algebra (2.1) we decompose the relativistic indices

$$A = (\alpha, a), \quad \alpha = 0, 1, \dots, p, \quad a = p + 1, \dots, D - 1. \quad (\text{A.11})$$

This induces the following decomposition of the generators:

$$\tilde{J}_{AB} \rightarrow \{\tilde{J}_{\alpha\beta}, \tilde{J}_{\alpha a}, \tilde{J}_{ab}\}, \quad \tilde{P}_A \rightarrow \{\tilde{P}_\alpha, \tilde{P}_a\}. \quad (\text{A.12})$$

p -brane Newton–Hooke expansions

Non-relativistic brane expansions of (A)dS of Newton–Hooke type can be defined starting from the following subspace decomposition,

$$V_0 = \{\tilde{P}_\alpha, \tilde{J}_{\alpha\beta}, \tilde{J}_{ab}\}, \quad V_1 = \{\tilde{P}_a, \tilde{J}_{\alpha a}\}. \quad (\text{A.13})$$

Using (2.21) we define the following resonant expansion of (A)dS

$$\left(S_0^{(\infty)} \times \{\tilde{P}_\alpha, \tilde{J}_{\alpha\beta}, \tilde{J}_{ab}\}\right) \oplus \left(S_1^{(\infty)} \times \{\tilde{P}_a, \tilde{J}_{\alpha a}\}\right). \quad (\text{A.14})$$

which is spanned by the following expanded generators

$$\begin{aligned} H_\alpha^{(m)} &= \lambda_{2m} \otimes \tilde{P}_\alpha, & P_a^{(m)} &= \lambda_{2m+1} \otimes \tilde{P}_a, \\ J_{\alpha\beta}^{(m)} &= \lambda_{2m} \otimes \tilde{J}_{\alpha\beta}, & B_{\alpha a}^{(m)} &= \lambda_{2m+1} \otimes \tilde{J}_{\alpha a}, \\ J_{ab}^{(m)} &= \lambda_{2m} \otimes \tilde{J}_{ab}. \end{aligned} \quad (\text{A.15})$$

The algebra of these generators is given by

$$[J_{\alpha\beta}^{(m)}, H_\gamma^{(n)}] = 2\eta_{\gamma[\beta}H_{\alpha]}^{(m+n)}, \quad (\text{A.16a}) \quad [J_{\alpha\beta}^{(m)}, J_{\gamma\delta}^{(n)}] = 4\eta_{[\alpha[\gamma}J_{\delta]\beta]}^{(m+n)}, \quad (\text{A.16c})$$

$$[J_{ab}^{(m)}, P_c^{(n)}] = 2\delta_{c[b}P_a^{(m+n)}, \quad (\text{A.16b}) \quad [J_{ab}^{(m)}, J_{cd}^{(n)}] = 4\delta_{[a[c}J_{d]b]}^{(m+n)}, \quad (\text{A.16d})$$

$$[B_{\alpha a}^{(m)}, P_b^{(n)}] = \delta_{ab} H_{\alpha}^{(m+n+1)}, \quad (\text{A.16e}) \quad [P_a^{(m)}, P_b^{(n)}] = -\Lambda J_{ab}^{(m+n+1)}, \quad (\text{A.16i})$$

$$[B_{\alpha a}^{(m)}, H_{\beta}^{(n)}] = -\eta_{\alpha\beta} P_a^{(m+n)}, \quad (\text{A.16f}) \quad [J_{\alpha\beta}^{(m)}, B_{\gamma d}^{(n)}] = 2\eta_{\gamma[\beta} B_{\alpha]d}^{(m+n)}, \quad (\text{A.16j})$$

$$[H_{\alpha}^{(m)}, H_{\beta}^{(n)}] = -\Lambda B_{\alpha\beta}^{(m+n)}, \quad (\text{A.16g}) \quad [J_{ab}^{(m)}, B_{\alpha c}^{(n)}] = 2\delta_{c[b} B_{\alpha]a}^{(m+n)}, \quad (\text{A.16k})$$

$$[P_a^{(m)}, H_{\alpha}^{(n)}] = \Lambda B_{\alpha a}^{(m+n)}, \quad (\text{A.16h}) \quad [B_{\alpha a}^{(m)}, B_{\beta b}^{(n)}] = -\eta_{\alpha\beta} J_{ab}^{(m+n+1)} - \delta_{ab} J_{\alpha\beta}^{(m+n+1)}. \quad (\text{A.16l})$$

Truncations of this infinite-dimensional algebra by suitable ideals give rise to finite expansions with the semigroup $S_E^{(N)}$ for different values of N . The simplest case corresponds to $N = 1$, which is the unextended p -brane Newton–Hooke algebra in D dimensions [35, 76]. The case $N = 2$ gives a higher-dimensional p -brane generalisation of the three-dimensional extended stringy Newton–Hooke algebra [77]. As in the particle case, higher values of N give rise to post-Newtonian extensions of the brane Newton–Hooke algebra. In the case of vanishing cosmological constant, these reduce to the non-relativistic expansion of the p -brane Galilean algebra [78–80], which have been studied in [37].

p -brane Carrollian expansions

In order to formulate p -brane Carrollian expansions, we follow [24] and interchange \tilde{P}_{α} and \tilde{P}_a in the subspace decomposition (A.13), i.e.

$$V_0 = \{\tilde{P}_a, \tilde{J}_{\alpha\beta}, \tilde{J}_{ab}\}, \quad V_1 = \{\tilde{P}_{\alpha}, \tilde{J}_{\alpha a}\}, \quad (\text{A.17})$$

leading to the resonant expanded algebra

$$\left(S_0^{(\infty)} \times \{\tilde{P}_a, \tilde{J}_{\alpha\beta}, \tilde{J}_{ab}\}\right) \oplus \left(S_1^{(\infty)} \times \{\tilde{P}_{\alpha}, \tilde{J}_{\alpha a}\}\right). \quad (\text{A.18})$$

and expanded generators

$$\begin{aligned} P_a^{(m)} &= \lambda_{2m} \otimes \tilde{P}_a, & H_{\alpha}^{(m)} &= \lambda_{2m+1} \otimes \tilde{P}_{\alpha}, \\ J_{\alpha\beta}^{(m)} &= \lambda_{2m} \otimes \tilde{J}_{\alpha\beta}, & B_{\alpha a}^{(m)} &= \lambda_{2m+1} \otimes \tilde{J}_{\alpha a}, \\ J_{ab}^{(m)} &= \lambda_{2m} \otimes \tilde{J}_{ab}. \end{aligned} \quad (\text{A.19})$$

This yields the following infinite-dimensional Carrollian (A)dS algebra

$$[J_{\alpha\beta}^{(m)}, H_{\gamma}^{(n)}] = 2\eta_{\gamma[\beta} H_{\alpha]}^{(m+n)}, \quad (\text{A.20a}) \quad [H_{\alpha}^{(m)}, H_{\beta}^{(n)}] = -\Lambda B_{\alpha\beta}^{(m+n+1)}, \quad (\text{A.20g})$$

$$[J_{ab}^{(m)}, P_c^{(n)}] = 2\delta_{c[b} P_{a]}^{(m+n)}, \quad (\text{A.20b}) \quad [P_a^{(m)}, H_{\alpha}^{(n)}] = \Lambda B_{\alpha a}^{(m+n)}, \quad (\text{A.20h})$$

$$[J_{\alpha\beta}^{(m)}, J_{\gamma\delta}^{(n)}] = 4\eta_{[\alpha[\gamma} J_{\delta]\beta]}^{(m+n)}, \quad (\text{A.20c}) \quad [P_a^{(m)}, P_b^{(n)}] = -\Lambda J_{ab}^{(m+n)}, \quad (\text{A.20i})$$

$$[J_{ab}^{(m)}, J_{cd}^{(n)}] = 4\delta_{[a[c} J_{d]b]}^{(m+n)}, \quad (\text{A.20d}) \quad [J_{\alpha\beta}^{(m)}, B_{\gamma d}^{(n)}] = 2\eta_{\gamma[\beta} B_{\alpha]d}^{(m+n)}, \quad (\text{A.20j})$$

$$[B_{\alpha a}^{(m)}, P_b^{(n)}] = \delta_{ab} H_{\alpha}^{(m+n)}, \quad (\text{A.20e}) \quad [J_{ab}^{(m)}, B_{\alpha c}^{(n)}] = 2\delta_{c[b} B_{\alpha]a}^{(m+n)}, \quad (\text{A.20k})$$

$$[B_{\alpha a}^{(m)}, H_{\beta}^{(n)}] = -\eta_{\alpha\beta} P_a^{(m+n+1)}, \quad (\text{A.20f}) \quad [B_{\alpha a}^{(m)}, B_{\beta b}^{(n)}] = -\eta_{\alpha\beta} J_{ab}^{(m+n+1)} - \delta_{ab} J_{\alpha\beta}^{(m+n+1)}. \quad (\text{A.20l})$$

The $N = 1$ truncation of this infinite-dimensional symmetry corresponds to the p -brane Carroll (A)dS algebra in D dimensions [22], while $S_E^{(N)}$ expansions for greater values of N define post-Carrollian (A)dS p -brane symmetries.

References

- [1] E. Inönü and E. P. Wigner, “On the contraction of groups and their representations,” *Proc. Nat. Acad. Sci.* **39** (1953) 510–524.
- [2] V. Bargmann, “On unitary ray representations of continuous groups,” *Annals Math.* **59** (1954) 1–46.
- [3] J.-M. Lévy-Leblond, “Group-theoretical foundations of classical mechanics: the lagrangian gauge problem,” *Communications in Mathematical Physics* **12** no. 1, (1969) 64–79.
- [4] G. Marmo, G. Morandi, A. Simoni, and E. C. G. Sudarshan, “Quasiinvariance and Central Extensions,” *Phys. Rev.* **D37** (1988) 2196–2205.
- [5] M. Hatsuda and M. Sakaguchi, “Wess-Zumino term for the AdS superstring and generalized Inonu-Wigner contraction,” *Prog. Theor. Phys.* **109** (2003) 853–867, [arXiv:hep-th/0106114 \[hep-th\]](#).
- [6] J. A. de Azcarraga, J. M. Izquierdo, M. Picon, and O. Varela, “Generating Lie and gauge free differential (super)algebras by expanding Maurer-Cartan forms and Chern-Simons supergravity,” *Nucl. Phys.* **B662** (2003) 185–219, [arXiv:hep-th/0212347 \[hep-th\]](#).
- [7] F. Izaurieta, E. Rodriguez, and P. Salgado, “Expanding Lie (super)algebras through Abelian semigroups,” *J. Math. Phys.* **47** (2006) 123512, [arXiv:hep-th/0606215 \[hep-th\]](#).
- [8] E. Bergshoeff, J. M. Izquierdo, T. Ortín, and L. Romano, “Lie Algebra Expansions and Actions for Non-Relativistic Gravity,” [arXiv:1904.08304 \[hep-th\]](#).
- [9] D. Hansen, J. Hartong, and N. A. Obers, “Action Principle for Newtonian Gravity,” *Phys. Rev. Lett.* **122** no. 6, (2019) 061106, [arXiv:1807.04765 \[hep-th\]](#).
- [10] D. Hansen, J. Hartong, and N. A. Obers, “Gravity between Newton and Einstein,” *Int. J. Mod. Phys.* **D28** no. 14, (2019) 1944010, [arXiv:1904.05706 \[gr-qc\]](#).
- [11] N. Ozdemir, M. Ozkan, O. Tunca, and U. Zorba, “Three-Dimensional Extended Newtonian (Super)Gravity,” *JHEP* **05** (2019) 130, [arXiv:1903.09377 \[hep-th\]](#).
- [12] J. Gomis, A. Kleinschmidt, J. Palmkvist, and P. Salgado-Rebolledo, “Symmetries of post-Newtonian expansions,” [arXiv:1910.13560 \[hep-th\]](#).
- [13] N. González, G. Rubio, P. Salgado, and S. Salgado, “Generalized Galilean algebras and Newtonian gravity,” *Phys. Lett.* **B755** (2016) 433–438, [arXiv:1604.06313 \[hep-th\]](#).

- [14] D. M. Peñafiel and L. Ravera, “Infinite S-Expansion with Ideal Subtraction and Some Applications,” *J. Math. Phys.* **58** no. 8, (2017) 081701, [arXiv:1611.05812 \[hep-th\]](#).
- [15] J. A. de Azcárraga, D. Gútiez, and J. M. Izquierdo, “Extended $D = 3$ Bargmann supergravity from a Lie algebra expansion,” [arXiv:1904.12786 \[hep-th\]](#).
- [16] D. M. Peñafiel and P. Salgado-Rebolledo, “Non-relativistic symmetries in three space-time dimensions and the Nappi-Witten algebra,” *Phys. Lett.* **B798** (2019) 135005, [arXiv:1906.02161 \[hep-th\]](#).
- [17] P. Concha and E. Rodríguez, “Non-Relativistic Gravity Theory based on an Enlargement of the Extended Bargmann Algebra,” [arXiv:1906.00086 \[hep-th\]](#).
- [18] J. Gomis, A. Kleinschmidt, and J. Palmkvist, “Galilean free Lie algebras,” *JHEP* **09** (2019) 109, [arXiv:1907.00410 \[hep-th\]](#).
- [19] H. Bacry and J. Levy-Leblond, “Possible kinematics,” *J. Math. Phys.* **9** (1968) 1605–1614.
- [20] J.-R. Derome and J.-G. Dubois, “Hooke’s symmetries and nonrelativistic cosmological kinematics. I,” *Il Nuovo Cimento B (1971-1996)* **9** no. 2, (1972) 351–376.
- [21] C. Duval, G. W. Gibbons, P. A. Horvathy, and P. M. Zhang, “Carroll versus Newton and Galilei: two dual non-Einsteinian concepts of time,” *Class. Quant. Grav.* **31** (2014) 085016, [arXiv:1402.0657 \[gr-qc\]](#).
- [22] T. E. Clark and T. ter Veldhuis, “AdS-Carroll Branes,” *J. Math. Phys.* **57** no. 11, (2016) 112303, [arXiv:1605.05484 \[hep-th\]](#).
- [23] R. Basu and U. N. Chowdhury, “Dynamical structure of Carrollian Electrodynamics,” *JHEP* **04** (2018) 111, [arXiv:1802.09366 \[hep-th\]](#).
- [24] A. Barducci, R. Casalbuoni, and J. Gomis, “Confined dynamical systems with Carroll and Galilei symmetries,” *Phys. Rev.* **D98** no. 8, (2018) 085018, [arXiv:1804.10495 \[hep-th\]](#).
- [25] G. Papageorgiou and B. J. Schroers, “Galilean quantum gravity with cosmological constant and the extended q -Heisenberg algebra,” *JHEP* **11** (2010) 020, [arXiv:1008.0279 \[hep-th\]](#).
- [26] J. Hartong, Y. Lei, and N. A. Obers, “Nonrelativistic Chern-Simons theories and three-dimensional Hořava-Lifshitz gravity,” *Phys. Rev.* **D94** no. 6, (2016) 065027, [arXiv:1604.08054 \[hep-th\]](#).
- [27] G. Papageorgiou and B. J. Schroers, “A Chern-Simons approach to Galilean quantum gravity in 2+1 dimensions,” *JHEP* **11** (2009) 009, [arXiv:0907.2880 \[hep-th\]](#).
- [28] E. A. Bergshoeff and J. Rosseel, “Three-Dimensional Extended Bargmann Supergravity,” *Phys. Rev. Lett.* **116** no. 25, (2016) 251601, [arXiv:1604.08042 \[hep-th\]](#).

- [29] G. Dautcourt, “On the Newtonian limit of general relativity,” *Acta Physica Polonica. Series B* **21** no. 10, (1990) 755–765.
- [30] R. De Pietri, L. Lusanna, and M. Pauri, “Standard and generalized Newtonian gravities as ‘gauge’ theories of the extended Galilei group. I. The standard theory,” *Class. Quant. Grav.* **12** (1995) 219–254, [arXiv:gr-qc/9405046 \[gr-qc\]](#).
- [31] G. Dautcourt, “PostNewtonian extension of the Newton-Cartan theory,” *Class. Quant. Grav.* **14** (1997) A109–A118, [arXiv:gr-qc/9610036 \[gr-qc\]](#).
- [32] D. Van den Bleeken, “Torsional Newton-Cartan gravity from the large c expansion of general relativity,” *Class. Quant. Grav.* **34** no. 18, (2017) 185004, [arXiv:1703.03459 \[gr-qc\]](#).
- [33] D. Hansen, J. Hartong, and N. A. Obers, “Non-relativistic expansion of the Einstein-Hilbert Lagrangian,” in *15th Marcel Grossmann Meeting on Recent Developments in Theoretical and Experimental General Relativity, Astrophysics, and Relativistic Field Theories (MG15) Rome, Italy, July 1-7, 2018*. 2019. [arXiv:1905.13723 \[gr-qc\]](#).
- [34] J. Gomis, J. Gomis, and K. Kamimura, “Non-relativistic superstrings: A New soluble sector of $\text{AdS}(5) \times \text{S}^5$,” *JHEP* **12** (2005) 024, [arXiv:hep-th/0507036 \[hep-th\]](#).
- [35] J. Brugues, J. Gomis, and K. Kamimura, “Newton-Hooke algebras, non-relativistic branes and generalized pp-wave metrics,” *Phys. Rev.* **D73** (2006) 085011, [arXiv:hep-th/0603023 \[hep-th\]](#).
- [36] C. Batlle, J. Gomis, and D. Not, “Extended Galilean symmetries of non-relativistic strings,” *JHEP* **02** (2017) 049, [arXiv:1611.00026 \[hep-th\]](#).
- [37] T. Harmark, J. Hartong, L. Menculini, N. A. Obers, and G. Oling, “Relating non-relativistic string theories,” *JHEP* **11** (2019) 071, [arXiv:1907.01663 \[hep-th\]](#).
- [38] P. Concha, L. Ravera, and E. Rodríguez, “Three-dimensional exotic Newtonian gravity with cosmological constant,” [arXiv:1912.02836 \[hep-th\]](#).
- [39] F. Ali and L. Ravera, “ \mathcal{N} -extended Chern-Simons Carrollian supergravities in $2 + 1$ spacetime dimensions,” [arXiv:1912.04172 \[hep-th\]](#).
- [40] S. Bonanos and J. Gomis, “A Note on the Chevalley-Eilenberg Cohomology for the Galilei and Poincare Algebras,” *J. Phys.* **A42** (2009) 145206, [arXiv:0808.2243 \[hep-th\]](#).
- [41] A. Medina and P. Revoy, “Algèbres de lie et produit scalaire invariant,” *Annales scientifiques de l’École Normale Supérieure* **4e série, 18** no. 3, (1985) 553–561.
- [42] J. M. Figueroa-O’Farrill and S. Stanciu, “On the structure of symmetric selfdual Lie algebras,” *J. Math. Phys.* **37** (1996) 4121–4134, [arXiv:hep-th/9506152 \[hep-th\]](#).

- [43] J. Matulich, S. Prohazka, and J. Salzer, “Limits of three-dimensional gravity and metric kinematical Lie algebras in any dimension,” *JHEP* **07** (2019) 118, [arXiv:1903.09165 \[hep-th\]](#).
- [44] O. Arratia, M. A. Martin, and M. A. Olmo, “Classical Systems and Representations of (2+1) Newton-Hooke Symmetries,” *arXiv e-prints* (Mar, 1999) [math-ph/9903013](#), [arXiv:math-ph/9903013 \[math-ph\]](#).
- [45] P. D. Alvarez, J. Gomis, K. Kamimura, and M. S. Plyushchay, “(2+1)D Exotic Newton-Hooke Symmetry, Duality and Projective Phase,” *Annals Phys.* **322** (2007) 1556–1586, [arXiv:hep-th/0702014 \[hep-th\]](#).
- [46] J.-M. Levy-Leblond, “Galilei group and galilean invariance,” in *Group theory and its applications*, pp. 221–299. Elsevier, 1971.
- [47] Y. Brihaye, C. Gónner, S. Giller, and P. Kosinski, “Galilean invariance in (2+1)-dimensions,” [arXiv:hep-th/9503046 \[hep-th\]](#).
- [48] J. Gomis and A. Kleinschmidt, “On free Lie algebras and particles in electro-magnetic fields,” *JHEP* **07** (2017) 085, [arXiv:1705.05854 \[hep-th\]](#).
- [49] E. Bergshoeff, J. Gomis, B. Rollier, J. Rosseel, and T. ter Veldhuis, “Carroll versus Galilei Gravity,” *JHEP* **03** (2017) 165, [arXiv:1701.06156 \[hep-th\]](#).
- [50] A. Achúcarro and P. Townsend, “A Chern-Simons Action for Three-Dimensional anti-De Sitter Supergravity Theories,” *Phys.Lett.* **B180** (1986) 89.
- [51] E. Witten, “(2+1)-Dimensional Gravity as an Exactly Soluble System,” *Nucl. Phys.* **B311** (1988) 46.
- [52] O. Coussaert, M. Henneaux, and P. van Driel, “The Asymptotic dynamics of three-dimensional Einstein gravity with a negative cosmological constant,” *Class. Quant. Grav.* **12** (1995) 2961–2966, [arXiv:gr-qc/9506019 \[gr-qc\]](#).
- [53] M. Banados, “Three-dimensional quantum geometry and black holes,” *AIP Conf. Proc.* **484** no. 1, (1999) 147–169, [arXiv:hep-th/9901148 \[hep-th\]](#).
- [54] S. Carlip, “Conformal field theory, (2+1)-dimensional gravity, and the BTZ black hole,” *Class. Quant. Grav.* **22** (2005) R85–R124, [arXiv:gr-qc/0503022 \[gr-qc\]](#).
- [55] J. Hartong, Y. Lei, N. A. Obers, and G. Oling, “Zooming in on AdS₃/CFT₂ near a BPS bound,” *JHEP* **05** (2018) 016, [arXiv:1712.05794 \[hep-th\]](#).
- [56] L. Ravera, “AdS Carroll Chern-Simons supergravity in 2+1 dimensions and its flat limit,” [arXiv:1905.00766 \[hep-th\]](#).
- [57] J. Hartong, “Gauging the Carroll Algebra and Ultra-Relativistic Gravity,” *JHEP* **08** (2015) 069, [arXiv:1505.05011 \[hep-th\]](#).

- [58] E. Bergshoeff, D. Grumiller, S. Prohazka, and J. Rosseel, “Three-dimensional Spin-3 Theories Based on General Kinematical Algebras,” *JHEP* **01** (2017) 114, [arXiv:1612.02277 \[hep-th\]](#).
- [59] M. Henneaux, “Geometry of Zero Signature Space-times,” *Bull. Soc. Math. Belg.* **31** (1979) 47–63.
- [60] C. Teitelboim, “Quantum Mechanics of the Gravitational Field,” *Phys. Rev.* **D25** (1982) 3159.
- [61] C. J. Isham, “Some Quantum Field Theory Aspects of the Superspace Quantization of General Relativity,” *Proc. Roy. Soc. Lond.* **A351** (1976) 209–232.
- [62] M. Henneaux, M. Pilati, and C. Teitelboim, “Explicit Solution for the Zero Signature (Strong Coupling) Limit of the Propagation Amplitude in Quantum Gravity,” *Phys. Lett.* **110B** (1982) 123–128.
- [63] E. Anderson, “Strong coupled relativity without relativity,” *Gen. Rel. Grav.* **36** (2004) 255–276, [arXiv:gr-qc/0205118 \[gr-qc\]](#).
- [64] M. Niedermaier, “The gauge structure of strong coupling gravity,” *Class. Quant. Grav.* **32** (2015) 015007.
- [65] V. A. Belinski, I. M. Khalatnikov, and E. M. Lifshitz, “Oscillatory approach to a singular point in the relativistic cosmology,” *Adv. Phys.* **19** (1970) 525–573.
- [66] V. A. Belinski, I. M. Khalatnikov, and E. M. Lifshitz, “A General Solution of the Einstein Equations with a Time Singularity,” *Adv. Phys.* **31** (1982) 639–667.
- [67] T. Damour, M. Henneaux, and H. Nicolai, “Cosmological billiards,” *Class. Quant. Grav.* **20** (2003) R145–R200, [arXiv:hep-th/0212256 \[hep-th\]](#).
- [68] G. Dautcourt, “On the ultrarelativistic limit of general relativity,” *Acta Phys. Polon.* **B29** (1998) 1047–1055, [arXiv:gr-qc/9801093 \[gr-qc\]](#).
- [69] L. Donnay and C. Marteau, “Carrollian Physics at the Black Hole Horizon,” *Class. Quant. Grav.* **36** no. 16, (2019) 165002, [arXiv:1903.09654 \[hep-th\]](#).
- [70] D. Grumiller and M. Riegler, “Most general AdS₃ boundary conditions,” *JHEP* **10** (2016) 023, [arXiv:1608.01308 \[hep-th\]](#).
- [71] A. Campoleoni, “Higher Spins in $D = 2 + 1$,” *Subnucl. Ser.* **49** (2013) 385–396, [arXiv:1110.5841 \[hep-th\]](#).
- [72] R. Schrader, “The Maxwell group and the quantum theory of particles in classical homogeneous electromagnetic fields,” *Fortsch. Phys.* **20** (1972) 701–734.
- [73] S. Bonanos and J. Gomis, “Infinite Sequence of Poincare Group Extensions: Structure and Dynamics,” *J. Phys.* **A43** (2010) 015201, [arXiv:0812.4140 \[hep-th\]](#).

- [74] P. Salgado-Rebolledo, “The Maxwell group in 2+1 dimensions and its infinite-dimensional enhancements,” [arXiv:1905.09421 \[hep-th\]](#).
- [75] L. Avilés, E. Frodden, J. Gomis, D. Hidalgo, and J. Zanelli, “Non-Relativistic Maxwell Chern-Simons Gravity,” *JHEP* **05** (2018) 047, [arXiv:1802.08453 \[hep-th\]](#).
- [76] R. Andringa, E. Bergshoeff, J. Gomis, and M. de Roo, “‘Stringy’ Newton-Cartan Gravity,” *Class. Quant. Grav.* **29** (2012) 235020, [arXiv:1206.5176 \[hep-th\]](#).
- [77] L. Avilés, J. Gomis, and D. Hidalgo, “Stringy (Galilei) Newton-Hooke Chern-Simons Gravities,” [arXiv:1905.13091 \[hep-th\]](#).
- [78] E. Bergshoeff, J. Gomis, and Z. Yan, “Nonrelativistic String Theory and T-Duality,” *JHEP* **11** (2018) 133, [arXiv:1806.06071 \[hep-th\]](#).
- [79] E. A. Bergshoeff, K. T. Grosvenor, C. Şimşek, and Z. Yan, “An Action for Extended String Newton-Cartan Gravity,” *JHEP* **01** (2019) 178, [arXiv:1810.09387 \[hep-th\]](#).
- [80] E. A. Bergshoeff, J. Gomis, J. Rosseel, C. Şimşek, and Z. Yan, “String Theory and String Newton-Cartan Geometry,” *J. Phys.* **A53** no. 1, (2019) 014001, [arXiv:1907.10668 \[hep-th\]](#).