EUROPEAN ORGANIZATION FOR NUCLEAR RESEARCH (CERN)



CERN-EP-2017-325 LHCb-PAPER-2017-039 October 15, 2018

Search for the rare decay $\Lambda_c^+ ightarrow p \mu^+ \mu^-$

LHCb collaboration[†]

Abstract

A search for the flavor-changing neutral-current decay $\Lambda_c^+ \to p\mu^+\mu^-$ is reported using a data set corresponding to an integrated luminosity of 3.0 fb⁻¹ collected by the LHCb collaboration. No significant signal is observed outside of the dimuon mass regions around the ϕ and ω resonances and an upper limit is placed on the branching fraction of $\mathcal{B}(\Lambda_c^+ \to p\mu^+\mu^-) < 7.7 (9.6) \times 10^{-8}$ at 90% (95%) confidence level. A significant signal is observed in the ω dimuon mass region for the first time.

Published in Phys. Rev. D97 (2018) 091101(R)

© CERN on behalf of the LHCb collaboration, licence CC-BY-4.0.

[†]Authors are listed at the end of this paper.

The flavor-changing neutral-current (FCNC) decay $\Lambda_c^+ \to p\mu^+\mu^-$ (inclusion of the charge-conjugate processes is implied throughout) is expected to be heavily suppressed in the Standard Model (SM) by the Glashow-Iliopoulos-Maiani mechanism [1]. The branching fractions for short-distance $c \to u\ell^+\ell^-$ contributions to the transition are expected to be of $\mathcal{O}(10^{-9})$ in the SM, but can be enhanced by effects beyond the SM. However, long-distance contributions proceeding via a tree-level amplitude, with an intermediate meson resonance decaying into a dimuon pair [2,3], can increase the branching fraction up to $\mathcal{O}(10^{-6})$ [4]. The short-distance and hadronic contributions can be separated by splitting the data set into relevant regions of dimuon mass. The $\Lambda_c^+ \to p\mu^+\mu^-$ decay has been previously searched for by the BaBar collaboration [5] yielding 11.1 ± 5.0 ± 2.5 events and an upper limit on the branching fraction of 4.4×10^{-5} at 90% CL.

Similar FCNC transitions for the *b*-quark system $(b \to s\ell^+\ell^-)$ exhibit a pattern of consistent deviations from the current SM predictions both in branching fractions [6] and angular observables [7], with the combined significance reaching 4 to 5 standard deviations [8,9]. Processes involving $c \to u\ell^+\ell^-$ transitions are far less explored at both the experimental and theoretical levels, which makes such measurements desirable. Similar analyses of the *D* system have reported evidence for the long-distance contribution [10], however, the short-distance contributions have not been established [11].

In this paper we report on the search for the $\Lambda_c^+ \to p\mu^+\mu^-$ decay, using a data set corresponding to an integrated luminosity of $3.0 \,\mathrm{fb}^{-1}$ of pp collisions collected in 2011 and 2012 with the LHCb experiment. The branching fraction is measured with respect to the branching fraction of the decay $\Lambda_c^+ \to p\phi(1020)$ with $\phi(1020) \to \mu^+\mu^-$ (here and after denoted as $\Lambda_c^+ \to p\phi(\mu^+\mu^-)$) decay, which has the benefit of having the same initial and final states and consequently many sources of systematic uncertainty are expected to cancel.

The LHCb detector [12, 13] is a single-arm forward spectrometer covering the pseudorapidity range $2 < \eta < 5$, designed for the study of particles containing b or c quarks. The detector includes a high-precision tracking system consisting of a siliconstrip vertex detector surrounding the pp interaction region [14], a large-area silicon-strip detector located upstream of a dipole magnet with a bending power of about 4 Tm, and three stations of silicon-strip detectors and straw drift tubes [15] placed downstream of the magnet. The tracking system provides a measurement of momentum, of charged particles with a relative uncertainty that varies from 0.5% at low momentum to 1.0% at 200 GeV/c. The minimum distance of a track to a primary vertex (PV), the impact parameter (IP), is measured with a resolution of $(15 + 29/p_T) \mu m$, where p_T is the component of the momentum transverse to the beam, in GeV/c. Different types of charged hadrons are distinguished using information from two ring-imaging Cherenkov detectors [16]. Photons, electrons and hadrons are identified by a calorimeter system consisting of scintillating-pad and preshower detectors, an electromagnetic calorimeter and a hadronic calorimeter. Muons are identified by a system composed of alternating layers of iron and multiwire proportional chambers [17]. The online event selection is performed by a trigger [18], which consists of a hardware stage, based on information from the calorimeter and muon systems, followed by a software stage, which applies a full event reconstruction.

Samples of simulated events are used to understand the properties of the signal and normalization channels. The *pp* collisions are generated using PYTHIA [19] with a specific LHCb configuration [20]. Decays of hadronic particles are described by EVTGEN [21], in which final-state radiation is generated using PHOTOS [22]. The decay of the Λ_c^+ baryon to $p\mu^+\mu^-$ is simulated with a three-body phase-space model. The interaction of the generated particles with the detector, and its response, are implemented using the GEANT4 toolkit [23] as described in Ref. [24]. The Λ_c^+ baryons are produced in two ways at a hadron collider; as prompt Λ_c^+ or in *b*-hadron decays. The simulation contains a mixture of these two production mechanisms, according to the known Λ_c^+ and *b*-hadron production cross-sections [25, 26].

The simulated samples are used to determine the selection criteria, in particular to train a multivariate classifier that is aimed at distinguishing signal signatures in the background-dominated data set. The simulated samples are also used to calculate the efficiencies of several selection steps.

Candidate events of $\Lambda_c^+ \to p\mu^+\mu^-$ decay are reconstructed by combining a pair of charged tracks identified as muons with one identified as a proton. Candidates that pass the trigger selections are subject to further requirements consisting of kinematic and particle identification criteria and based on the response of a multivariate classifier. Each of the final-state tracks is required to be of good quality, to have $p_T > 300 \text{ MeV}/c$ and to be incompatible with originating from any of the PVs in the event. The tracks are also required to form a good-quality secondary vertex with a corresponding flight distance of at least 0.1 mm from all of the PVs in the event. The invariant mass of the dimuon system is required to be smaller than 1400 MeV/ c^2 . Three dimuon mass regions are defined:

- a region around the known ϕ mass, [985, 1055] MeV/ $c^2,$ used as a normalization channel;
- a region around the known ω mass (the ω denotes hereafter the $\omega(782)$ meson), [759, 805] MeV/ c^2 , used to isolate the $\Lambda_c^+ \to p\omega$ decay;
- a nonresonant region $(\Lambda_c^+ \to p \mu^+ \mu^-)$, with excluded ranges $\pm 40 \text{ MeV}/c^2$ around the known ω and ϕ masses.

After the preselection, the normalization channel is still dominated by combinatorial background, *i.e.* combinations of tracks that do not all originate from a genuine Λ_c^+ baryon. A boosted decision tree (BDT) is trained to reduce the combinatorial background to manageable level. The BDT is trained using the kinematic and topological variables of the Λ_c^+ candidate, related to its flight distance, decay vertex quality, $p_{\rm T}$ and impact parameter with respect to the primary vertex. In the BDT training, $\Lambda_c^+ \to p\mu^+\mu^-$ simulated events are used as a proxy for the signal and data outside the signal $p\mu^+\mu^-$ invariant mass region extending up to $\pm 300 \,{\rm MeV}/c^2$ around the known Λ_c^+ mass is used as a proxy for the background.

A k-folding technique is used to ensure the training is unbiased [27], while keeping the full available data sets for further analysis. A loose BDT cut is applied to reduce the background to the same level as the normalization channel yield.

A fit to the $p\mu^+\mu^-$ invariant-mass distribution of $\Lambda_c^+ \to p\phi(\mu^+\mu^-)$ candidates after the loose BDT requirement is shown in Fig. 1. The shape of the Λ_c^+ peak is parametrized by a Crystal Ball function [28] with parameters determined from the simulation, while the background is modeled with a first-order polynomial. The yield of the $\Lambda_c^+ \to p\phi(\mu^+\mu^-)$ decay is determined to be 395 ± 45 candidates. This sample is used for the final optimization of selection requirements. It is checked at this stage that the variables used in the signal selection are well described by simulation within the available sample size.



Figure 1: Mass distribution of $\Lambda_c^+ \to p\mu^+\mu^-$ candidates in the ϕ region after the first BDT requirement. The solid line shows the result of the fit described in the text, while the dashed line indicates the background component.

For the final selection a second BDT is trained, which includes additional variables related to Λ_c^+ -baryon decay properties and the isolation of the proton and muons in the detector. The final discrimination is performed in three dimensions: the BDT variable and two particle identification (PID) variables, the proton-identification discriminant and the muon-identification discriminant. The optimal set of BDT and PID requirements is determined by finding the best expected upper limit on the branching fraction of the signal relative to the normalization channel using the CL_s method [29] by means of Monte Carlo methods.

Several sources of background have been considered. An irreducible background due to long-distance contributions originates from $\Lambda_c^+ \to pV(\mu^+\mu^-)$ decays, with intermediate resonances indicated by V. The $\rho(770)^0$, ω and η resonances are studied, however, their contribution to the nonresonant region is expected to be negligible, because the V meson mass is well separated from the nonresonant region and/or the $\Lambda_c^+ \to pV(\mu^+\mu^-)$ branching fraction is small. Another background source considered is due to misidentification of final-state particles in hadronic D^+ , D_s^+ and Λ_c^+ decays. The expected contribution from this source has been estimated using large samples of simulated events. Given the tight PID requirements obtained from the optimization, only 2.0 ± 1.1 candidates are expected to fall into the Λ_c^+ mass window in the nonresonant region.

The ratio of branching fractions is measured using

$$\frac{\mathcal{B}(\Lambda_c^+ \to p\mu^+\mu^-)}{\mathcal{B}(\Lambda_c^+ \to p\phi)\mathcal{B}(\phi \to \mu^+\mu^-)} = \frac{\epsilon_{\text{norm}}}{\epsilon_{\text{sig}}} \times \frac{N_{\text{sig}}}{N_{\text{norm}}},\tag{1}$$

where N_{sig} (N_{norm}) is the observed yield for the signal (normalization) decay mode. The factors ϵ_{sig} and ϵ_{norm} indicate the corresponding total efficiencies for signal and normalization channels, respectively. The efficiencies are determined from the simulation.

In the case of the observation of the decay $\Lambda_c^+ \to pV$, the ratio of branching fractions is determined by

$$\frac{\mathcal{B}(\Lambda_c^+ \to pV)\mathcal{B}(V \to \mu^+\mu^-)}{\mathcal{B}(\Lambda_c^+ \to p\phi)\mathcal{B}(\phi \to \mu^+\mu^-)} = \frac{\epsilon_{\rm norm}}{\epsilon_{\rm V}} \times \frac{N_V}{N_{\rm norm}},\tag{2}$$

Uncertainty source	Value [%]	Value [%]
	$\Lambda_c^+ \rightarrow p \mu^+ \mu^-$	$\Lambda_c^+ \rightarrow pV(\mu^+\mu^-)$
	nonresonant	ω region
Size of simulation samples	4.4	10.0
BDT cut	4.8	4.8
PID cut	0.7	0.7
Total	6.5	11.1

Table 1: Systematic uncertainties on the efficiency ratio used in the determination of the branching fraction in the nonresonant and ω regions.

where $N_V(N_{\text{norm}})$ is the number of candidates observed for the $\Lambda_c^+ \to pV$ (normalization) decay mode. The factors ϵ_V and ϵ_{norm} indicate the corresponding total efficiencies for $\Lambda_c^+ \to pV$ and normalization channel, respectively.

As the final states of the signal and normalization channels are identical, many sources of systematic uncertainty cancel in the ratio of the efficiencies. There are three significant sources of systematic uncertainty. The first is related to the finite size of the simulation samples, which limits the precision on the efficiency ratio. The second is linked to residual differences between data and simulation of the BDT distribution. The third is associated to the simulation of PID and is determined from the uncertainty on the PID calibration samples. The values of the contributions are given in Table 1.

Several other sources of systematic uncertainty were considered: the trigger efficiency, the shapes used in the invariant mass fit for signal and normalization channels, the shape of the combinatorial background, and the fraction of prompt Λ_c^+ baryons and Λ_c^+ baryons from *b*-hadron decays. All of these, however, are at negligible level when compared to three dominant sources of systematic uncertainty.

The simulated $\Lambda_c^+ \to p\mu^+\mu^-$ decays have been generated according to a phase-space model for the decay products. As the exact physics model for the decay is not known, no systematic uncertainty is assigned. Instead, the weights needed to recast the result in terms of any physics model are provided in Fig. 2. The weights are described by a function of the dimuon invariant mass squared $m^2(\mu^+\mu^-)$ and the invariant mass of the proton and the negatively charged muon squared $m^2(p\mu^-)$. The weights are normalized to the average efficiency.

The distributions of the $p\mu^+\mu^-$ invariant mass for the $\Lambda_c^+ \to p\mu^+\mu^-$ candidates after final selections in the three dimuon mass ranges are presented in Fig. 3. The Λ_c^+ peak is parametrized by a Crystal Ball [28] function with parameters determined from the simulation and the background is described by a first-order polynomial. The fits are used to determine the signal yields. No significant signal is observed in the nonresonant region (Fig. 3a). The yield for the normalization channel is determined to be 96 ± 11 candidates (Fig. 3b). An accumulation of 13.2 ± 4.3 candidates at the Λ_c^+ mass is observed in the ω region (Fig. 3c). The statistical significance of the excess is determined to be 5.0σ using Wilks' theorem [30].

The distribution of the dimuon invariant mass of the Λ_c^+ candidates is shown in Fig. 4. An excess is seen at the known ω and ϕ resonance masses. The data is well described by



Figure 2: The efficiency weights for $\Lambda_c^+ \to p\mu^+\mu^-$ as a function of the dimuon invariant mass squared $m^2(\mu^+\mu^-)$ and by the invariant mass of the proton and the negatively charged muon squared $m^2(p\mu^-)$. The weights are normalized to the average efficiency.

a simple model including these resonances and a background component. The ω and ϕ peaks are parametrized as Breit-Wigner functions of relevant decay width [31] convolved with a Gaussian function to take into account the experimental resolution. The addition of a component for the $\rho(770)^0$ resonance (and its interference with the ω meson) does not improve the fit quality. It is therefore assumed that the observed candidates in the ω region are dominated by decays via the ω resonance.

As no evidence for nonresonant $\Lambda_c^+ \to p\mu^+\mu^-$ decays is found, an upper limit on the branching fractions is determined using the CL_s method. The systematic uncertainties are included in the construction of CL_s . The following upper limits are obtained at different confidence levels (CL)

$$\frac{\mathcal{B}(\Lambda_c^+ \to p\mu^+\mu^-)}{\mathcal{B}(\Lambda_c^+ \to p\phi)\mathcal{B}(\phi \to \mu^+\mu^-)} < 0.24 \ (0.28) \quad \text{ at } 90\% \ (95\%) \ \text{CL}$$

The corresponding distribution of CL_s is shown in Fig. 5. Using the values of the branching fractions for $\Lambda_c^+ \to p\phi$ and $\phi \to \mu^+\mu^-$ decays from Ref. [31] and including their uncertainties in the CL_s construction, an upper limit on the branching fraction is determined to be

$$\mathcal{B}(\Lambda_c^+ \to p \mu^+ \mu^-) < 7.7 \ (9.6) \times 10^{-8}$$
 at 90% (95%) CL.

Under the above-mentioned assumption of the $\Lambda_c^+ \to p\omega$ dominance in the ω region, the relative branching fraction with respect to the normalization channel is determined according to Eq. 2

$$\frac{\mathcal{B}(\Lambda_c^+ \to p\omega)\mathcal{B}(\omega \to \mu^+\mu^-)}{\mathcal{B}(\Lambda_c^+ \to p\phi)\mathcal{B}(\phi \to \mu^+\mu^-)} = 0.23 \pm 0.08 \text{ (stat)} \pm 0.03 \text{ (syst)}.$$

Using the relevant branching fractions from Ref. [31], the branching fraction of $\Lambda_c^+ \to p\omega$ is determined to be

$$\mathcal{B}(\Lambda_c^+ \to p\omega) = (9.4 \pm 3.2 \text{ (stat)} \pm 1.0 \text{ (syst)} \pm 2.0 \text{ (ext)}) \times 10^{-4},$$



Figure 3: Mass distribution for selected $p\mu^+\mu^-$ candidates in the three regions of the dimuon invariant mass: a) nonresonant region, b) ϕ region, c) ω region. The solid lines show the results of the fit as described in the text. The dashed lines indicate the background component.

where the first uncertainty is statistical, the second corresponds to the above-mentioned systematic effects and the third is due to the limited knowledge of the relevant branching fractions. Assuming lepton universality, the branching fraction $\mathcal{B}(\omega \to e^+e^-)$ is used instead of $\mathcal{B}(\omega \to \mu^+\mu^-)$.

In summary, a search for the $\Lambda_c^+ \to p\mu^+\mu^-$ decay is reported, using pp data collected with the LHCb experiment. The analysis is performed in three regions of dimuon mass: ϕ , ω and nonresonant. The upper limit on the nonresonant mode is improved by two orders of magnitude with respect to the previous measurement [5]. For the first time the signal is seen in the ω region with a statistical significance of 5 standard deviations.



Figure 4: Invariant mass distribution $m(\mu^+\mu^-)$ for $\Lambda_c^+ \to p\mu^+\mu^-$ candidates with mass $\pm 25 \text{ MeV}/c^2$ around the Λ_c^+ mass. The solid line shows the result of the fit, while the dashed line indicates the background component.



Figure 5: The CL_s value as a function of the $\mathcal{B}(\Lambda_c^+ \to p\mu^+\mu^-)$ branching fraction. The median expected value of an ensemble (assuming no signal component) is shown by the dashed line, with the $\pm 1\sigma$ and $\pm 2\sigma$ regions shaded. The observed distribution is shown by the solid line.

Acknowledgements

We express our gratitude to our colleagues in the CERN accelerator departments for the excellent performance of the LHC. We thank the technical and administrative staff at the LHCb institutes. We acknowledge support from CERN and from the national agencies: CAPES, CNPq, FAPERJ and FINEP (Brazil); MOST and NSFC (China); CNRS/IN2P3

(France); BMBF, DFG and MPG (Germany); INFN (Italy); NWO (The Netherlands); MNiSW and NCN (Poland); MEN/IFA (Romania); MinES and FASO (Russia); MinECo (Spain); SNSF and SER (Switzerland); NASU (Ukraine); STFC (United Kingdom); NSF (USA). We acknowledge the computing resources that are provided by CERN, IN2P3 (France), KIT and DESY (Germany), INFN (Italy), SURF (The Netherlands), PIC (Spain), GridPP (United Kingdom), RRCKI and Yandex LLC (Russia), CSCS (Switzerland), IFIN-HH (Romania), CBPF (Brazil), PL-GRID (Poland) and OSC (USA). We are indebted to the communities behind the multiple open-source software packages on which we depend. Individual groups or members have received support from AvH Foundation (Germany), EPLANET, Marie Skłodowska-Curie Actions and ERC (European Union), ANR, Labex P2IO and OCEVU, and Région Auvergne-Rhône-Alpes (France), RFBR, RSF and Yandex LLC (Russia), GVA, XuntaGal and GENCAT (Spain), Herchel Smith Fund, the Royal Society, the English-Speaking Union and the Leverhulme Trust (United Kingdom).

References

- S. Glashow, J. Iliopoulos, and L. Maiani, Weak interactions with lepton-hadron symmetry, Phys. Rev. D2 (1970) 1285.
- [2] S. Fajfer et al., The rare decays of D mesons, Nucl. Phys. Proc. Supp. 115 (2003) 93.
- [3] G. Burdman, E. Golowich, J. L. Hewett, and S. Pakvasa, Rare charm decays in the standard model and beyond, Phys. Rev. D66 (2002) 014009, arXiv:hep-ph/0112235.
- S. Fajfer and S. Prelovsek, Effects of littlest Higgs model in rare D meson decays, Phys. Rev. D73 (2006) 054026, arXiv:hep-ph/0511048.
- [5] BaBar collaboration, J. P. Lees et al., Searches for rare or forbidden semileptonic charm decays, Phys. Rev. D84 (2011) 072006, arXiv:1107.4465.
- [6] LHCb collaboration, R. Aaij et al., Measurement of the S-wave fraction in $B^0 \rightarrow K^+\pi^-\mu^+\mu^-$ decays and the $B^0 \rightarrow K^*(892)^0\mu^+\mu^-$ differential branching fraction, JHEP 11 (2016) 047, Erratum ibid. 04 (2017) 142, arXiv:1606.04731.
- [7] LHCb collaboration, R. Aaij *et al.*, Angular analysis of the $B^0 \to K^{*0}\mu^+\mu^-$ decay using $3 fb^{-1}$ of integrated luminosity, JHEP **02** (2016) 104, arXiv:1512.04442.
- [8] B. Capdevila et al., Patterns of new physics in $b \to s\ell^+\ell^-$ transitions in the light of recent data, arXiv:1704.05340.
- [9] W. Altmannshofer, C. Niehoff, P. Stangl, and D. M. Straub, Status of the $B \rightarrow K^* \mu^+ \mu^-$ anomaly after Moriond 2017, Eur. Phys. J. C77 (2017) 377, arXiv:1703.09189.
- [10] LHCb collaboration, R. Aaij *et al.*, Observation of D^0 meson decays to $\pi^+\pi^-\mu^+\mu^-$ and $K^+K^-\mu^+\mu^-$ final states, Phys. Rev. Lett. **119** (2017) 181805, arXiv:1707.08377.
- [11] LHCb collaboration, R. Aaij *et al.*, Search for $D^+_{(s)} \to \pi^+ \mu^+ \mu^-$ and $D^+_{(s)} \to \pi^- \mu^+ \mu^+$ decays, Phys. Lett. **B724** (2013) 203, arXiv:1304.6365.

- [12] LHCb collaboration, A. A. Alves Jr. et al., The LHCb detector at the LHC, JINST 3 (2008) S08005.
- [13] LHCb collaboration, R. Aaij et al., LHCb detector performance, Int. J. Mod. Phys. A30 (2015) 1530022, arXiv:1412.6352.
- [14] R. Aaij et al., Performance of the LHCb Vertex Locator, JINST 9 (2014) P09007, arXiv:1405.7808.
- [15] R. Arink et al., Performance of the LHCb Outer Tracker, JINST 9 (2014) P01002, arXiv:1311.3893.
- [16] M. Adinolfi et al., Performance of the LHCb RICH detector at the LHC, Eur. Phys. J. C73 (2013) 2431, arXiv:1211.6759.
- [17] A. A. Alves Jr. et al., Performance of the LHCb muon system, JINST 8 (2013) P02022, arXiv:1211.1346.
- [18] R. Aaij et al., The LHCb trigger and its performance in 2011, JINST 8 (2013) P04022, arXiv:1211.3055.
- T. Sjöstrand, S. Mrenna, and P. Skands, *PYTHIA 6.4 physics and manual*, JHEP 05 (2006) 026, arXiv:hep-ph/0603175; T. Sjöstrand, S. Mrenna, and P. Skands, *A brief introduction to PYTHIA 8.1*, Comput. Phys. Commun. 178 (2008) 852, arXiv:0710.3820.
- [20] I. Belyaev et al., Handling of the generation of primary events in Gauss, the LHCb simulation framework, J. Phys. Conf. Ser. 331 (2011) 032047.
- [21] D. J. Lange, The EvtGen particle decay simulation package, Nucl. Instrum. Meth. A462 (2001) 152.
- [22] P. Golonka and Z. Was, PHOTOS Monte Carlo: A precision tool for QED corrections in Z and W decays, Eur. Phys. J. C45 (2006) 97, arXiv:hep-ph/0506026.
- [23] Geant4 collaboration, J. Allison et al., Geant4 developments and applications, IEEE Trans. Nucl. Sci. 53 (2006) 270; Geant4 collaboration, S. Agostinelli et al., Geant4: A simulation toolkit, Nucl. Instrum. Meth. A506 (2003) 250.
- [24] M. Clemencic et al., The LHCb simulation application, Gauss: Design, evolution and experience, J. Phys. Conf. Ser. 331 (2011) 032023.
- [25] LHCb collaboration, R. Aaij et al., Measurement of B meson production cross-sections in proton-proton collisions at $\sqrt{s} = 7$ TeV, JHEP **08** (2013) 117, arXiv:1306.3663.
- [26] LHCb collaboration, R. Aaij *et al.*, Prompt charm production in pp collisions at $\sqrt{s} = 7 \ TeV$, Nucl. Phys. **B871** (2013) 1, arXiv:1302.2864.
- [27] A. Blum, A. Kalai, and J. Langford, Beating the hold-out: Bounds for k-fold and progressive cross-validation, in Proceedings of the Twelfth Annual Conference on Computational Learning Theory, COLT '99, (New York, NY, USA), pp. 203–208, ACM, 1999.

- [28] T. Skwarnicki, A study of the radiative cascade transitions between the Upsilon-prime and Upsilon resonances, PhD thesis, Institute of Nuclear Physics, Krakow, 1986, DESY-F31-86-02.
- [29] A. L. Read, Presentation of search results: The CL_s technique, J. Phys. G28 (2002) 2693.
- [30] S. S. Wilks, The large-sample distribution of the likelihood ratio for testing composite hypotheses, Ann. Math. Statist. **9** (1938) 60.
- [31] Particle Data Group, C. Patrignani et al., Review of particle physics, Chin. Phys. C40 (2016) 100001.

LHCb collaboration

R. Aaij⁴⁰, B. Adeva³⁹, M. Adinolfi⁴⁸, Z. Ajaltouni⁵, S. Akar⁵⁹, J. Albrecht¹⁰, F. Alessio⁴⁰, M. Alexander⁵³, A. Alfonso Albero³⁸, S. Ali⁴³, G. Alkhazov³¹, P. Alvarez Cartelle⁵⁵, A.A. Alves Jr⁵⁹, S. Amato², S. Amerio²³, Y. Amhis⁷, L. An³, L. Anderlini¹⁸, G. Andreassi⁴¹, M. Andreotti^{17,g}, J.E. Andrews⁶⁰, R.B. Appleby⁵⁶, F. Archilli⁴³, P. d'Argent¹², J. Arnau Romeu⁶, A. Artamonov³⁷, M. Artuso⁶¹, E. Aslanides⁶, M. Atzeni⁴², G. Auriemma²⁶, M. Baalouch⁵, I. Babuschkin⁵⁶, S. Bachmann¹², J.J. Back⁵⁰, A. Badalov^{38,m}, C. Baesso⁶², S. Baker⁵⁵, V. Balagura^{7,b}, W. Baldini¹⁷, A. Baranov³⁵, R.J. Barlow⁵⁶, C. Barschel⁴⁰, S. Barsuk⁷, W. Barter⁵⁶, F. Baryshnikov³², V. Batozskaya²⁹, V. Battista⁴¹, A. Bay⁴¹, L. Beaucourt⁴, J. Beddow⁵³, F. Bedeschi²⁴, I. Bediaga¹, A. Beiter⁶¹, L.J. Bel⁴³, N. Beliv⁶³, V. Bellee⁴¹, N. Belloli^{21,i}, K. Belous³⁷, I. Belyaev^{32,40}, E. Ben-Haim⁸, G. Bencivenni¹⁹, S. Benson⁴³, S. Beranek⁹, A. Berezhnoy³³, R. Bernet⁴², D. Berninghoff¹², E. Bertholet⁸, A. Bertolin²³, C. Betancourt⁴², F. Betti¹⁵, M.O. Bettler⁴⁰, M. van Beuzekom⁴³, Ia. Bezshyiko⁴², S. Bifani⁴⁷, P. Billoir⁸, A. Birnkraut¹⁰, A. Bizzeti^{18,u}, M. Bjørn⁵⁷, T. Blake⁵⁰, F. Blanc⁴¹, S. Blusk⁶¹, V. Bocci²⁶, T. Boettcher⁵⁸, A. Bondar^{36,w}, N. Bondar³¹, I. Bordyuzhin³², S. Borghi^{56,40}, M. Borisyak³⁵, M. Borsato³⁹, F. Bossu⁷, M. Boubdir⁹, T.J.V. Bowcock⁵⁴, E. Bowen⁴², C. Bozzi^{17,40}, S. Braun¹², J. Brodzicka²⁷, D. Brundu¹⁶, E. Buchanan⁴⁸, C. Burr⁵⁶, A. Bursche^{16, f}, J. Buytaert⁴⁰, W. Byczynski⁴⁰, S. Cadeddu¹⁶, H. Cai⁶⁴, R. Calabrese^{17, g}, R. Calladine⁴⁷, M. Calvi^{21,i}, M. Calvo Gomez^{38,m}, A. Camboni^{38,m}, P. Campana¹⁹, D.H. Campora Perez⁴⁰, L. Capriotti⁵⁶, A. Carbone^{15,e}, G. Carboni^{25,j}, R. Cardinale^{20,h}, A. Cardini¹⁶, P. Carniti^{21,i}, L. Carson⁵², K. Carvalho Akiba², G. Casse⁵⁴, L. Cassina²¹, M. Cattaneo⁴⁰, G. Cavallero^{20,40,h}, R. Cenci^{24,t}, D. Chamont⁷, M.G. Chapman⁴⁸, M. Charles⁸, Ph. Charpentier⁴⁰, G. Chatzikonstantinidis⁴⁷, M. Chefdeville⁴, S. Chen¹⁶, S.F. Cheung⁵⁷, S.-G. Chitic⁴⁰, V. Chobanova³⁹, M. Chrzaszcz⁴², A. Chubykin³¹, P. Ciambrone¹⁹, X. Cid Vidal³⁹, G. Ciezarek⁴⁰, P.E.L. Clarke⁵², M. Clemencic⁴⁰, H.V. Cliff⁴⁹, J. Closier⁴⁰, V. Coco⁴⁰, J. Cogan⁶, E. Cogneras⁵, V. Cogoni^{16,f}, L. Cojocariu³⁰, P. Collins⁴⁰, T. Colombo⁴⁰, A. Comerma-Montells¹², A. Contu¹⁶, G. Coombs⁴⁰, S. Coquereau³⁸, G. Corti⁴⁰, M. Corvo^{17,g}, C.M. Costa Sobral⁵⁰, B. Couturier⁴⁰, G.A. Cowan⁵², D.C. Craik⁵⁸, A. Crocombe⁵⁰, M. Cruz Torres¹, R. Currie⁵², C. D'Ambrosio⁴⁰, F. Da Cunha Marinho², C.L. Da Silva⁷³, E. Dall'Occo⁴³, J. Dalseno⁴⁸, A. Davis³, O. De Aguiar Francisco⁴⁰, K. De Bruyn⁴⁰, S. De Capua⁵⁶, M. De Cian¹², J.M. De Miranda¹, L. De Paula², M. De Serio¹⁴, d, P. De Simone¹⁹, C.T. Dean⁵³, D. Decamp⁴, L. Del Buono⁸, H.-P. Dembinski¹¹, M. Demmer¹⁰, A. Dendek²⁸, D. Derkach³⁵, O. Deschamps⁵, F. Dettori⁵⁴, B. Dey⁶⁵, A. Di Canto⁴⁰, P. Di Nezza¹⁹, H. Dijkstra⁴⁰, F. Dordei⁴⁰, M. Dorigo⁴⁰, A. Dosil Suárez³⁹, L. Douglas⁵³, A. Dovbnya⁴⁵, K. Dreimanis⁵⁴, L. Dufour⁴³, G. Dujany⁸, P. Durante⁴⁰, J.M. Durham⁷³, D. Dutta⁵⁶, R. Dzhelyadin³⁷, M. Dziewiecki¹², A. Dziurda⁴⁰, A. Dzyuba³¹, S. Easo⁵¹, U. Egede⁵⁵, V. Egorychev³², S. Eidelman^{36,w}, S. Eisenhardt⁵², U. Eitschberger¹⁰, R. Ekelhof¹⁰, L. Eklund⁵³, S. Ely⁶¹, S. Esen¹², H.M. Evans⁴⁹, T. Evans⁵⁷, A. Falabella¹⁵, N. Farley⁴⁷, S. Farry⁵⁴, D. Fazzini^{21,i}, L. Federici²⁵, D. Ferguson⁵², G. Fernandez³⁸, P. Fernandez Declara⁴⁰, A. Fernandez Prieto³⁹, F. Ferrari¹⁵, L. Ferreira Lopes⁴¹, F. Ferreira Rodrigues², M. Ferro-Luzzi⁴⁰, S. Filippov³⁴, R.A. Fini¹⁴, M. Fiorini^{17,g}, M. Firlej²⁸, C. Fitzpatrick⁴¹, T. Fiutowski²⁸, F. Fleuret^{7,b}, M. Fontana^{16,40}, F. Fontanelli^{20,h}, R. Forty⁴⁰, V. Franco Lima⁵⁴, M. Frank⁴⁰, C. Frei⁴⁰, J. Fu^{22,q}, W. Funk⁴⁰, E. Furfaro^{25,j}, C. Färber⁴⁰, E. Gabriel⁵², A. Gallas Torreira³⁹, D. Galli^{15,e}, S. Gallorini²³, S. Gambetta⁵², M. Gandelman², P. Gandini²², Y. Gao³, L.M. Garcia Martin⁷¹, J. García Pardiñas³⁹, J. Garra Tico⁴⁹, L. Garrido³⁸, D. Gascon³⁸, C. Gaspar⁴⁰, L. Gavardi¹⁰, G. Gazzoni⁵, D. Gerick¹², E. Gersabeck⁵⁶, M. Gersabeck⁵⁶, T. Gershon⁵⁰, Ph. Ghez⁴, S. Giani⁴¹, V. Gibson⁴⁹, O.G. Girard⁴¹, L. Giubega³⁰, K. Gizdov⁵², V.V. Gligorov⁸, D. Golubkov³², A. Golutvin^{55,69}, A. Gomes^{1,a}, I.V. Gorelov³³, C. Gotti^{21,i}, E. Govorkova⁴³, J.P. Grabowski¹², R. Graciani Diaz³⁸,

L.A. Granado Cardoso⁴⁰, E. Graugés³⁸, E. Graverini⁴², G. Graziani¹⁸, A. Grecu³⁰, R. Greim⁹, P. Griffith¹⁶, L. Grillo⁵⁶, L. Gruber⁴⁰, B.R. Gruberg Cazon⁵⁷, O. Grünberg⁶⁷, E. Gushchin³⁴, Yu. Guz³⁷, T. Gys⁴⁰, C. Göbel⁶², T. Hadavizadeh⁵⁷, C. Hadjivasiliou⁵, G. Haefeli⁴¹, C. Haen⁴⁰, S.C. Haines⁴⁹, B. Hamilton⁶⁰, X. Han¹², T.H. Hancock⁵⁷, S. Hansmann-Menzemer¹², N. Harnew⁵⁷, S.T. Harnew⁴⁸, C. Hasse⁴⁰, M. Hatch⁴⁰, J. He⁶³, M. Hecker⁵⁵, K. Heinicke¹⁰, A. Heister⁹, K. Hennessy⁵⁴, P. Henrard⁵, L. Henry⁷¹, E. van Herwijnen⁴⁰, M. Heß⁶⁷, A. Hicheur², D. Hill⁵⁷, P.H. Hopchev⁴¹, W. Hu⁶⁵, W. Huang⁶³, Z.C. Huard⁵⁹, W. Hulsbergen⁴³, T. Humair⁵⁵, M. Hushchyn³⁵, D. Hutchcroft⁵⁴, P. Ibis¹⁰, M. Idzik²⁸, P. Ilten⁴⁷, R. Jacobsson⁴⁰, J. Jalocha⁵⁷, E. Jans⁴³, A. Jawahery⁶⁰, M. Jezabek²⁷, F. Jiang³, M. John⁵⁷, D. Johnson⁴⁰, C.R. Jones⁴⁹, C. Joram⁴⁰, B. Jost⁴⁰, N. Jurik⁵⁷, S. Kandybei⁴⁵, M. Karacson⁴⁰, J.M. Kariuki⁴⁸, S. Karodia⁵³, N. Kazeev³⁵, M. Kecke¹², F. Keizer⁴⁹, M. Kelsey⁶¹, M. Kenzie⁴⁹, T. Ketel⁴⁴, E. Khairullin³⁵, B. Khanji¹², C. Khurewathanakul⁴¹, K.E. Kim⁶¹, T. Kirn⁹, S. Klaver¹⁹, K. Klimaszewski²⁹, T. Klimkovich¹¹, S. Koliiev⁴⁶, M. Kolpin¹², R. Kopecna¹², P. Koppenburg⁴³, A. Kosmyntseva³², S. Kotriakhova³¹, M. Kozeiha⁵, L. Kravchuk³⁴, M. Kreps⁵⁰, F. Kress⁵⁵, P. Krokovny^{36,w}, W. Krzemien²⁹, W. Kucewicz^{27,l}, M. Kucharczyk²⁷, V. Kudryavtsev^{36,w}, A.K. Kuonen⁴¹, T. Kvaratskheliya^{32,40}, D. Lacarrere⁴⁰, G. Lafferty⁵⁶, A. Lai¹⁶, G. Lanfranchi¹⁹, C. Langenbruch⁹, T. Latham⁵⁰, C. Lazzeroni⁴⁷, R. Le Gac⁶, A. Leflat^{33,40}, J. Lefrançois⁷, R. Lefèvre⁵, F. Lemaitre⁴⁰, E. Lemos Cid³⁹, O. Leroy⁶, T. Lesiak²⁷, B. Leverington¹², P.-R. Li⁶³, T. Li³, Y. Li⁷, Z. Li⁶¹, X. Liang⁶¹, T. Likhomanenko⁶⁸, R. Lindner⁴⁰, F. Lionetto⁴², V. Lisovskyi⁷, X. Liu³, D. Loh⁵⁰, A. Loi¹⁶, I. Longstaff⁵³, J.H. Lopes², D. Lucchesi^{23,o}, M. Lucio Martinez³⁹, H. Luo⁵², A. Lupato²³, E. Luppi^{17,g}, O. Lupton⁴⁰, A. Lusiani²⁴, X. Lyu⁶³, F. Machefert⁷, F. Maciuc³⁰, V. Macko⁴¹, P. Mackowiak¹⁰, S. Maddrell-Mander⁴⁸, O. Maev^{31,40}, K. Maguire⁵⁶, D. Maisuzenko³¹, M.W. Majewski²⁸, S. Malde⁵⁷, B. Malecki²⁷, A. Malinin⁶⁸, T. Maltsev^{36,w}, G. Manca^{16,f}, G. Mancinelli⁶, D. Marangotto^{22,q}, J. Maratas^{5,v}, J.F. Marchand⁴, U. Marconi¹⁵, C. Marin Benito³⁸, M. Marinangeli⁴¹, P. Marino⁴¹, J. Marks¹², G. Martellotti²⁶, M. Martin⁶, M. Martinelli⁴¹, D. Martinez Santos³⁹, F. Martinez Vidal⁷¹, A. Massafferri¹, R. Matev⁴⁰, A. Mathad⁵⁰, Z. Mathe⁴⁰, C. Matteuzzi²¹, A. Mauri⁴², E. Maurice^{7,b}, B. Maurin⁴¹, A. Mazurov⁴⁷, M. McCann^{55,40}, A. McNab⁵⁶, R. McNulty¹³, J.V. Mead⁵⁴, B. Meadows⁵⁹, C. Meaux⁶, F. Meier¹⁰, N. Meinert⁶⁷, D. Melnychuk²⁹, M. Merk⁴³, A. Merli^{22,40,q}, E. Michielin²³, D.A. Milanes⁶⁶, E. Millard⁵⁰, M.-N. Minard⁴, L. Minzoni¹⁷, D.S. Mitzel¹², A. Mogini⁸, J. Molina Rodriguez¹, T. Mombächer¹⁰, I.A. Monroy⁶⁶, S. Monteil⁵, M. Morandin²³, M.J. Morello^{24,t}, O. Morgunova⁶⁸, J. Moron²⁸, A.B. Morris⁵², R. Mountain⁶¹, F. Muheim⁵², M. Mulder⁴³, D. Müller⁴⁰, J. Müller¹⁰, K. Müller⁴², V. Müller¹⁰, P. Naik⁴⁸, T. Nakada⁴¹, R. Nandakumar⁵¹, A. Nandi⁵⁷, I. Nasteva², M. Needham⁵², N. Neri^{22,40}, S. Neubert¹², N. Neufeld⁴⁰, M. Neuner¹², T.D. Nguyen⁴¹, C. Nguyen-Mau^{41,n}, S. Nieswand⁹, R. Niet¹⁰, N. Nikitin³³, T. Nikodem¹², A. Nogay⁶⁸, D.P. O'Hanlon⁵⁰, A. Oblakowska-Mucha²⁸, V. Obraztsov³⁷, S. Ogilvy¹⁹, R. Oldeman^{16,f}, C.J.G. Onderwater⁷², A. Ossowska²⁷, J.M. Otalora Goicochea², P. Owen⁴², A. Oyanguren⁷¹, P.R. Pais⁴¹, A. Palano¹⁴, M. Palutan^{19,40}, G. Panshin⁷⁰, A. Papanestis⁵¹, M. Pappagallo⁵², L.L. Pappalardo^{17,g}, W. Parker⁶⁰, C. Parkes⁵⁶, G. Passaleva^{18,40}, A. Pastore^{14,d}, M. Patel⁵⁵, C. Patrignani^{15,e}, A. Pearce⁴⁰, A. Pellegrino⁴³, G. Penso²⁶, M. Pepe Altarelli⁴⁰, S. Perazzini⁴⁰, D. Pereima³², P. Perret⁵, L. Pescatore⁴¹, K. Petridis⁴⁸, A. Petrolini^{20,h}, A. Petrov⁶⁸, M. Petruzzo^{22,q}, E. Picatoste Olloqui³⁸, B. Pietrzyk⁴, G. Pietrzyk⁴¹, M. Pikies²⁷, D. Pinci²⁶, F. Pisani⁴⁰, A. Pistone^{20,h}, A. Piucci¹², V. Placinta³⁰, S. Playfer⁵², M. Plo Casasus³⁹, F. Polci⁸, M. Poli Lener¹⁹, A. Poluektov⁵⁰, I. Polyakov⁶¹, E. Polycarpo², G.J. Pomery⁴⁸, S. Ponce⁴⁰, A. Popov³⁷, D. Popov^{11,40}, S. Poslavskii³⁷, C. Potterat², E. Price⁴⁸, J. Prisciandaro³⁹, C. Prouve⁴⁸, V. Pugatch⁴⁶, A. Puig Navarro⁴², H. Pullen⁵⁷, G. Punzi^{24,p}, W. Qian⁵⁰, J. Qin⁶³, R. Quagliani⁸, B. Quintana⁵, B. Rachwal²⁸, J.H. Rademacker⁴⁸, M. Rama²⁴, M. Ramos Pernas³⁹, M.S. Rangel², I. Raniuk^{45,†}, F. Ratnikov^{35,x}, G. Raven⁴⁴, M. Ravonel Salzgeber⁴⁰, M. Reboud⁴, F. Redi⁴¹, S. Reichert¹⁰,

A.C. dos Reis¹, C. Remon Alepuz⁷¹, V. Renaudin⁷, S. Ricciardi⁵¹, S. Richards⁴⁸, M. Rihl⁴⁰,

K. Rinnert⁵⁴, P. Robbe⁷, A. Robert⁸, A.B. Rodrigues⁴¹, E. Rodrigues⁵⁹, J.A. Rodriguez Lopez⁶⁶, A. Rogozhnikov³⁵, S. Roiser⁴⁰, A. Rollings⁵⁷, V. Romanovskiy³⁷, A. Romero Vidal^{39,40}, M. Rotondo¹⁹, M.S. Rudolph⁶¹, T. Ruf⁴⁰, P. Ruiz Valls⁷¹, J. Ruiz Vidal⁷¹, J.J. Saborido Silva³⁹, E. Sadykhov³², N. Sagidova³¹, B. Saitta^{16, f}, V. Salustino Guimaraes⁶², C. Sanchez Mayordomo⁷¹, B. Sanmartin Sedes³⁹, R. Santacesaria²⁶, C. Santamarina Rios³⁹, M. Santimaria¹⁹, E. Santovetti^{25,j}, G. Sarpis⁵⁶, A. Sarti^{19,k}, C. Satriano^{26,s}, A. Satta²⁵, D.M. Saunders⁴⁸, D. Savrina^{32,33}, S. Schael⁹, M. Schellenberg¹⁰, M. Schiller⁵³, H. Schindler⁴⁰, M. Schmelling¹¹, T. Schmelzer¹⁰, B. Schmidt⁴⁰, O. Schneider⁴¹, A. Schopper⁴⁰, H.F. Schreiner⁵⁹, M. Schubiger⁴¹, M.H. Schune⁷, R. Schwemmer⁴⁰, B. Sciascia¹⁹, A. Sciubba^{26,k}, A. Semennikov³², E.S. Sepulveda⁸, A. Sergi⁴⁷, N. Serra⁴², J. Serrano⁶, L. Sestini²³, P. Seyfert⁴⁰, M. Shapkin³⁷, Y. Shcheglov³¹, T. Shears⁵⁴, L. Shekhtman^{36,w}, V. Shevchenko⁶⁸, B.G. Siddi¹⁷, R. Silva Coutinho⁴², L. Silva de Oliveira², G. Simi^{23,o}, S. Simone^{14,d}, M. Sirendi⁴⁹, N. Skidmore⁴⁸, T. Skwarnicki⁶¹, I.T. Smith⁵², J. Smith⁴⁹, M. Smith⁵⁵, l. Soares Lavra¹, M.D. Sokoloff⁵⁹, F.J.P. Soler⁵³, B. Souza De Paula², B. Spaan¹⁰, P. Spradlin⁵³, F. Stagni⁴⁰, M. Stahl¹², S. Stahl⁴⁰, P. Stefko⁴¹, S. Stefkova⁵⁵, O. Steinkamp⁴², S. Stemmle¹², O. Stenyakin³⁷, M. Stepanova³¹, H. Stevens¹⁰, S. Stone⁶¹, B. Storaci⁴², S. Stracka^{24,p}, M.E. Stramaglia⁴¹, M. Straticiuc³⁰, U. Straumann⁴², S. Strokov⁷⁰, J. Sun³, L. Sun⁶⁴, K. Swientek²⁸, V. Syropoulos⁴⁴, T. Szumlak²⁸, M. Szymanski⁶³, S. T'Jampens⁴, A. Tayduganov⁶, T. Tekampe¹⁰, G. Tellarini^{17,g}, F. Teubert⁴⁰, E. Thomas⁴⁰, J. van Tilburg⁴³, M.J. Tilley⁵⁵, V. Tisserand⁵, M. Tobin⁴¹, S. Tolk⁴⁹, L. Tomassetti^{17,g}, D. Tonelli²⁴, R. Tourinho Jadallah Aoude¹, E. Tournefier⁴, M. Traill⁵³, M.T. Tran⁴¹, M. Tresch⁴², A. Trisovic⁴⁹, A. Tsaregorodtsev⁶, P. Tsopelas⁴³, A. Tully⁴⁹, N. Tuning^{43,40}, A. Ukleja²⁹, A. Usachov⁷, A. Ustvuzhanin³⁵, U. Uwer¹², C. Vacca^{16,f}, A. Vagner⁷⁰, V. Vagnoni^{15,40}, A. Valassi⁴⁰, S. Valat⁴⁰, G. Valenti¹⁵, R. Vazquez Gomez⁴⁰, P. Vazquez Regueiro³⁹, S. Vecchi¹⁷, M. van Veghel⁴³, J.J. Velthuis⁴⁸, M. Veltri^{18,r}, G. Veneziano⁵⁷, A. Venkateswaran⁶¹, T.A. Verlage⁹, M. Vernet⁵, M. Vesterinen⁵⁷, J.V. Viana Barbosa⁴⁰, D. Vieira⁶³, M. Vieites Diaz³⁹, H. Viemann⁶⁷, X. Vilasis-Cardona^{38,m}, M. Vitti⁴⁹, V. Volkov³³, A. Vollhardt⁴², B. Voneki⁴⁰, A. Vorobyev³¹, V. Vorobyev^{36,w}, C. Voß⁹, J.A. de Vries⁴³, C. Vázquez Sierra⁴³, R. Waldi⁶⁷, J. Walsh²⁴, J. Wang⁶¹, Y. Wang⁶⁵, D.R. Ward⁴⁹, H.M. Wark⁵⁴, N.K. Watson⁴⁷, D. Websdale⁵⁵, A. Weiden⁴², C. Weisser⁵⁸, M. Whitehead⁴⁰, J. Wicht⁵⁰, G. Wilkinson⁵⁷, M. Wilkinson⁶¹, M. Williams⁵⁶, M. Williams⁵⁸, T. Williams⁴⁷, F.F. Wilson^{51,40}, J. Wimberley⁶⁰, M. Winn⁷, J. Wishahi¹⁰, W. Wislicki²⁹, M. Witek²⁷, G. Wormser⁷, S.A. Wotton⁴⁹, K. Wyllie⁴⁰, Y. Xie⁶⁵, M. Xu⁶⁵, Q. Xu⁶³, Z. Xu³, Z. Xu⁴, Z. Yang³, Z. Yang⁶⁰, Y. Yao⁶¹, H. Yin⁶⁵, J. Yu⁶⁵, X. Yuan⁶¹, O. Yushchenko³⁷, K.A. Zarebski⁴⁷, M. Zavertyaev^{11,c}, L. Zhang³, Y. Zhang⁷, A. Zhelezov¹², Y. Zheng⁶³, X. Zhu³, V. Zhukov^{9,33}, J.B. Zonneveld⁵², S. Zucchelli¹⁵.

¹Centro Brasileiro de Pesquisas Físicas (CBPF), Rio de Janeiro, Brazil

² Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil

³Center for High Energy Physics, Tsinghua University, Beijing, China

⁴ Univ. Grenoble Alpes, Univ. Savoie Mont Blanc, CNRS, IN2P3-LAPP, Annecy, France

⁵Clermont Université, Université Blaise Pascal, CNRS/IN2P3, LPC, Clermont-Ferrand, France

⁶Aix Marseille Univ, CNRS/IN2P3, CPPM, Marseille, France

⁷LAL, Univ. Paris-Sud, CNRS/IN2P3, Université Paris-Saclay, Orsay, France

⁸LPNHE, Université Pierre et Marie Curie, Université Paris Diderot, CNRS/IN2P3, Paris, France

⁹I. Physikalisches Institut, RWTH Aachen University, Aachen, Germany

¹⁰ Fakultät Physik, Technische Universität Dortmund, Dortmund, Germany

¹¹Max-Planck-Institut für Kernphysik (MPIK), Heidelberg, Germany

¹²Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany

¹³School of Physics, University College Dublin, Dublin, Ireland

¹⁴Sezione INFN di Bari, Bari, Italy

¹⁵Sezione INFN di Bologna, Bologna, Italy

¹⁶Sezione INFN di Cagliari, Cagliari, Italy

¹⁷ Universita e INFN, Ferrara, Ferrara, Italy

¹⁸Sezione INFN di Firenze, Firenze, Italy

¹⁹Laboratori Nazionali dell'INFN di Frascati, Frascati, Italy

²⁰Sezione INFN di Genova, Genova, Italy

²¹Sezione INFN di Milano Bicocca, Milano, Italy

²²Sezione di Milano, Milano, Italy

²³Sezione INFN di Padova, Padova, Italy

²⁴Sezione INFN di Pisa, Pisa, Italy

²⁵Sezione INFN di Roma Tor Vergata, Roma, Italy

²⁶Sezione INFN di Roma La Sapienza, Roma, Italy

²⁷Henryk Niewodniczanski Institute of Nuclear Physics Polish Academy of Sciences, Kraków, Poland

²⁸AGH - University of Science and Technology, Faculty of Physics and Applied Computer Science, Kraków, Poland

²⁹National Center for Nuclear Research (NCBJ), Warsaw, Poland

³⁰ Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest-Magurele, Romania ³¹ Petersburg Nuclear Physics Institute (PNPI), Gatchina, Russia

³²Institute of Theoretical and Experimental Physics (ITEP), Moscow, Russia

³³Institute of Nuclear Physics, Moscow State University (SINP MSU), Moscow, Russia

³⁴Institute for Nuclear Research of the Russian Academy of Sciences (INR RAS), Moscow, Russia

³⁵ Yandex School of Data Analysis, Moscow, Russia

³⁶Budker Institute of Nuclear Physics (SB RAS), Novosibirsk, Russia

³⁷Institute for High Energy Physics (IHEP), Protvino, Russia

³⁸ICCUB, Universitat de Barcelona, Barcelona, Spain

³⁹Instituto Galego de Física de Altas Enerxías (IGFAE), Universidade de Santiago de Compostela, Santiago de Compostela, Spain

⁴⁰ European Organization for Nuclear Research (CERN), Geneva, Switzerland

⁴¹Institute of Physics, Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland

⁴² Physik-Institut, Universität Zürich, Zürich, Switzerland

⁴³Nikhef National Institute for Subatomic Physics, Amsterdam, The Netherlands

⁴⁴Nikhef National Institute for Subatomic Physics and VU University Amsterdam, Amsterdam, The Netherlands

⁴⁵NSC Kharkiv Institute of Physics and Technology (NSC KIPT), Kharkiv, Ukraine

⁴⁶Institute for Nuclear Research of the National Academy of Sciences (KINR), Kyiv, Ukraine

⁴⁷ University of Birmingham, Birmingham, United Kingdom

⁴⁸H.H. Wills Physics Laboratory, University of Bristol, Bristol, United Kingdom

⁴⁹Cavendish Laboratory, University of Cambridge, Cambridge, United Kingdom

⁵⁰Department of Physics, University of Warwick, Coventry, United Kingdom

⁵¹STFC Rutherford Appleton Laboratory, Didcot, United Kingdom

⁵²School of Physics and Astronomy, University of Edinburgh, Edinburgh, United Kingdom

⁵³School of Physics and Astronomy, University of Glasgow, Glasgow, United Kingdom

⁵⁴Oliver Lodge Laboratory, University of Liverpool, Liverpool, United Kingdom

⁵⁵Imperial College London, London, United Kingdom

⁵⁶School of Physics and Astronomy, University of Manchester, Manchester, United Kingdom

⁵⁷Department of Physics, University of Oxford, Oxford, United Kingdom

⁵⁸ Massachusetts Institute of Technology, Cambridge, MA, United States

⁵⁹ University of Cincinnati, Cincinnati, OH, United States

⁶⁰ University of Maryland, College Park, MD, United States

⁶¹Syracuse University, Syracuse, NY, United States

⁶²Pontifícia Universidade Católica do Rio de Janeiro (PUC-Rio), Rio de Janeiro, Brazil, associated to ² ⁶³University of Chinese Academy of Sciences, Beijing, China, associated to ³

⁶⁴School of Physics and Technology, Wuhan University, Wuhan, China, associated to ³

⁶⁵Institute of Particle Physics, Central China Normal University, Wuhan, Hubei, China, associated to ³

⁶⁶Departamento de Fisica, Universidad Nacional de Colombia, Bogota, Colombia, associated to ⁸

⁶⁷Institut für Physik, Universität Rostock, Rostock, Germany, associated to ¹²

⁶⁸National Research Centre Kurchatov Institute, Moscow, Russia, associated to ³²

⁶⁹National University of Science and Technology MISIS, Moscow, Russia, associated to ³²

⁷⁰National Research Tomsk Polytechnic University, Tomsk, Russia, associated to ³²

⁷¹Instituto de Fisica Corpuscular, Centro Mixto Universidad de Valencia - CSIC, Valencia, Spain, associated to ³⁸

⁷² Van Swinderen Institute, University of Groningen, Groningen, The Netherlands, associated to ⁴³
 ⁷³ Los Alamos National Laboratory (LANL), Los Alamos, United States, associated to ⁶¹

^a Universidade Federal do Triângulo Mineiro (UFTM), Uberaba-MG, Brazil

^bLaboratoire Leprince-Ringuet, Palaiseau, France

^cP.N. Lebedev Physical Institute, Russian Academy of Science (LPI RAS), Moscow, Russia

^d Università di Bari, Bari, Italy

^e Università di Bologna, Bologna, Italy

^f Università di Cagliari, Cagliari, Italy

^g Università di Ferrara, Ferrara, Italy

^h Università di Genova, Genova, Italy

ⁱ Università di Milano Bicocca, Milano, Italy

^j Università di Roma Tor Vergata, Roma, Italy

^k Università di Roma La Sapienza, Roma, Italy

¹AGH - University of Science and Technology, Faculty of Computer Science, Electronics and

Telecommunications, Kraków, Poland

^mLIFAELS, La Salle, Universitat Ramon Llull, Barcelona, Spain

ⁿHanoi University of Science, Hanoi, Vietnam

^o Università di Padova, Padova, Italy

^pUniversità di Pisa, Pisa, Italy

^qUniversità degli Studi di Milano, Milano, Italy

^r Università di Urbino, Urbino, Italy

^s Università della Basilicata, Potenza, Italy

^tScuola Normale Superiore, Pisa, Italy

 u Università di Modena e Reggio Emilia, Modena, Italy

 v Iligan Institute of Technology (IIT), Iligan, Philippines

 w Novosibirsk State University, Novosibirsk, Russia

^x National Research University Higher School of Economics, Moscow, Russia

 $^{\dagger}Deceased$