

## Ground-based observations of stratospheric NO<sub>2</sub> at high and midlatitudes in Europe after the Mount Pinatubo eruption

M. Van Roozendael,<sup>1</sup> M. De Mazière,<sup>1</sup> C. Hermans,<sup>1</sup> P. C. Simon,<sup>1</sup> J.-P. Pommereau,<sup>2</sup> F. Goutail,<sup>2</sup> X. X. Tie,<sup>3</sup> G. Brasseur,<sup>3</sup> and C. Granier<sup>3</sup>

**Abstract.** Nitrogen dioxide has been monitored at the International Scientific Station at the Jungfraujoch (46°N, 8°E) since June 1990 and at Sodankyla (67°N, 27°E) since January 1990. NO<sub>2</sub> vertical column abundances are measured during the morning and evening twilights by application of the differential absorption method using the sunlight scattered at zenith in the visible range. The available time series shows a significant reduction of NO<sub>2</sub> starting in winter 1992, after the eruption of the Mount Pinatubo volcano. A maximum decrease of about 35% is observed in January 1992 at both stations. The continued series of observations shows the recovery of the NO<sub>2</sub> column until August 1995. These results are compared with two-dimensional chemical model calculations, including the effect of heterogeneous reactions on observed Pinatubo aerosols. In general, the modeled NO<sub>2</sub> columns agree qualitatively with the observations although the amplitude of the seasonal variation is underestimated, possibly due to internal limitations of the model which, for example, does not include diurnal changes. The observed and calculated NO<sub>2</sub> percent changes are in good agreement, which confirms quantitatively the impact of the heterogeneous chemistry on stratospheric NO<sub>2</sub>.

### 1. Introduction

Strong evidence exists on the basis of comparisons between model simulations and observations that heterogeneous processes occurring on the surface of stratospheric sulfuric acid aerosols (SSAs) are responsible for stratospheric ozone losses at midlatitudes. These heterogeneous processes lead to ozone depletion by converting halogens from reservoir forms to active forms, hence feeding halogen-based ozone-destroying catalytic cycles [Solomon *et al.*, 1996, and references therein].

The key chemical reactions are the hydrolysis of N<sub>2</sub>O<sub>5</sub> and ClONO<sub>2</sub>:



The conversion of nitrogen oxide into nitric acid is more efficient at high latitude during winter because the lack of solar radiation in those conditions reduces the efficiency of nitric acid photodissociation. In addition, the conversion of chlorine nitrate is strongly temperature dependent and efficient only at low temperature. Therefore reaction (2) can generally be neglected at midlatitude [Tie *et al.*, 1994].

As both N<sub>2</sub>O<sub>5</sub> and ClONO<sub>2</sub> are produced by reactions involving NO<sub>2</sub> (NO<sub>2</sub> + NO<sub>3</sub> and NO<sub>2</sub> + ClO, respectively) and HNO<sub>3</sub> is the long-lived sink for reactive nitrogen NO<sub>x</sub> (NO +

NO<sub>2</sub>), both reactions (1) and (2) decrease NO<sub>x</sub>. Hence they were already proposed as possible mechanisms for the significant NO<sub>x</sub> reduction observed between 30 and 35 km from balloon measurements performed before and after the El Chichon eruption in April 1982 [Roscoe *et al.*, 1986].

Following the major eruption of Mount Pinatubo in June 1991 which injected 20 megaton of SO<sub>2</sub> in the stratosphere [Bluth *et al.*, 1993; McCormick *et al.*, 1995] several observations have confirmed the chemical changes of the midlatitude stratosphere. Johnston *et al.* [1992] have reported 35% reduction of NO<sub>2</sub> total amount based on the longest time series of stratospheric NO<sub>2</sub> reported so far, beginning in the 1980s in Lauder (New Zealand). A similar reduction has been reported for the winter 1991–1992 by Van Roozendael *et al.* [1994] from measurements made at the International Scientific Station of the Jungfraujoch (ISSJ) in the Alps (46°N). Mills *et al.* [1993] confirmed the strong anticorrelation between the NO<sub>2</sub> total amount above Colorado and the SSAs. They also confirmed experimentally that NO<sub>2</sub> perturbation tends to saturate with increasing surface area of SSA as indicated by Prather [1992]. In situ measurements published by Fahey *et al.* [1993], including reactive nitrogen and chlorine species, confirmed the importance of SSA surface area in the production of the inactive nitrogen reservoir HNO<sub>3</sub>. Changing the partitioning between reservoir halogenated compounds and active ozone-destroying species, SSAs and associated heterogeneous reactions modify the relative importance of the destroying catalytic cycles of ozone. In addition to NO<sub>2</sub> total amount reduction, changes in nitrogen reservoirs resulting from reactions (1) and (2) have been measured. Koike *et al.* [1994] reported measured HNO<sub>3</sub> total amounts anticorrelated with NO<sub>2</sub> above Lauder (New Zealand) over the period 1992–1993. However, these observations are not confirmed by aircraft measurements at northern midlatitudes by Coffey and Mankin [1993] although they report a NO<sub>2</sub> total amount reduction of about 50% after both El Chichon and Mount Pinatubo eruptions.

<sup>1</sup>Belgian Institute for Space Aeronomy, Brussels, Belgium.

<sup>2</sup>Service d'Aéronomie du Centre National de la Recherche Scientifique, Verrières-le-Buisson, France.

<sup>3</sup>National Center for Atmospheric Research, Boulder, Colorado.

Copyright 1997 by the American Geophysical Union.

Paper number 97JD01098.  
0148-0227/97/97JD-01098\$09.00

Profile measurements of N<sub>2</sub>O<sub>5</sub> and HNO<sub>3</sub> obtained by ATMOS between 25°N and 53°S in March–April 1992 during the ATLAS 1 mission confirmed the perturbed vertical distribution of those species, especially in the tropics [Rinsland *et al.*, 1994]. These measurements confirm the role of reaction (1) in altering the odd nitrogen partitioning after the Mount Pinatubo eruption.

Several model calculations have been performed which support the experimental results. Regarding NO<sub>2</sub> total amounts, the most significant comparison has been published by Bekki and Pyle [1994] based on a coupled two-dimensional chemical-microphysical model and on measurements performed at Sodankyla (67°N) one year before and one year after the Mt. Pinatubo eruption.

This work reports on time series of ground-based visible measurements of NO<sub>2</sub> total columns covering pre- and post-Pinatubo eruption conditions at two latitudes (46°N and 67°N). The experimental results obtained over 6 years with identical instruments at both sites are compared with 2-D model simulations, including the heterogeneous processes already mentioned. The aerosol climatology used in the calculations is based on lidar measurements above the Alps (47°N) from which surface area of SSAs have been calculated [Jäger, 1995].

## 2. Instrument and Data Analysis

Measurements of the NO<sub>2</sub> vertical column amount have been performed since June 1990 at the International Scientific Station at the Jungfraujoch (46°N, 8°E) and since January 1990 at the Sodankyla Observatory (67°N, 26°E) using a Système d'Analyse par Observations Zénithales (SAOZ) [Pommereau and Goutail, 1988]. The SAOZ spectrometer and the data analysis method were described in details in previous papers [Goutail *et al.*, 1994; Van Roozendael *et al.*, 1994a]. After participating in the Network for Detection of Stratospheric Changes (NDSC) intercomparison of UV-visible instruments (May 1992, Lauder, New Zealand [Hofmann *et al.*, 1995]), this instrument has been qualified for inclusion in the NDSC.


Daily observations of the zenith-scattered sunlight are carried out at sunrise and sunset twilights in the visible region. The NO<sub>2</sub> column amounts are retrieved by the differential absorption method between 400 and 455 nm. Twilight measurements result from an average over a few spectra (between 4 and 8) recorded between 87° and 91° solar zenith angle (SZA). The conversion from slant to vertical column amounts is obtained using air mass factors (AMFs) calculated according to Solomon *et al.* [1987]. These calculations use a NO<sub>2</sub> vertical distribution measured from balloon experiment at midlatitude. It is assumed that the AMF depends on the SZA only; that is, the same set of AMFs is used throughout the year at both sites. Comparisons with integrated profiles measured with the SAOZ balloon-borne version in France and Scandinavia [Pommereau and Piquard, 1994] have shown that the uncertainty on the NO<sub>2</sub> vertical column due to this approximation is less than 10% [Goutail *et al.*, 1994]. The seasonal dependence of the error due to seasonal changes in the NO<sub>2</sub> profile is estimated less than 5% from calculations based on SAGE II data (unpublished results). Other sources of uncertainties, discussed by Van Roozendael *et al.* [1994b], are (1) the absorption cross-sections, (2) the calculation of the residual amount in the reference spectrum and (3) the meas-

urement noise. At 90° SZA the overall precision of the NO<sub>2</sub> measurements by the SAOZ instrument is limited by the AMF for large NO<sub>2</sub> columns (relative error less than 10%) and by the measurement noise for small NO<sub>2</sub> columns (absolute error of about 2×10<sup>14</sup> molecules/cm<sup>2</sup>). Much larger errors can be encountered due to contamination of the site by tropospheric NO<sub>2</sub>. Very large NO<sub>2</sub> enhancements are observed regularly at both sites mostly due to a combination of surface pollution and tropospheric multiple scattering in the presence of rain or snow showers. In the case of Jungfraujoch data, a systematic method has been developed to detect and reject pollution events [Van Roozendael *et al.*, 1994b]. The Sodankyla data set was cleaned from the largest pollution peaks by simple observation of the time series.

## 3. Results

Daily sunrise and sunset NO<sub>2</sub> total column measurements were carried out from January 1990 to August 1995 at Sodankyla (67°N) and at the Jungfraujoch Station (46°N). The 5-year NO<sub>2</sub> time series from Sodankyla is complete, but some data are missing at the Jungfraujoch, as listed below. During winter 1990/1991, morning values could not be reliably retrieved due to noise at the detector level. In October 1991 the SAOZ spectrometer was replaced by a new version of the same instrument. No data were obtained between June and October 1991 due to the first instrument being requested at another site and the delivery time of the second one. Other gaps in summer 1992 and 1994 resulted from damages to the instrument's electronics due to thunderstorms.

Figure 1 shows monthly averaged morning and evening NO<sub>2</sub> column amounts calculated at both stations in order to smooth out the day-to-day variability. Instead of using arithmetic means, monthly means were obtained by fitting a re-




**Figure 1.** Time series of monthly averaged NO<sub>2</sub> data available at (a) Sodankyla and (b) the Jungfraujoch Station. For each month an average morning and evening twilight value is plotted. The vertical dotted line indicates the day of the Mount Pinatubo eruption.

gression line to the daily columns within a month and calculating the value of this line on the 15th of the month. This method minimizes the error on the monthly averages that might arise because of missing NO<sub>2</sub> values. The error bars shown together with the monthly means represent the variability of the NO<sub>2</sub> columns within a month (1  $\sigma$  standard deviation). Except for winter data at 67°N, these variabilities are larger than the uncertainties on the measurements.

The observed NO<sub>2</sub> columns follow the known latitudinal distribution of stratospheric NO<sub>2</sub> [e.g., Noxon *et al.*, 1979, 1980]. Midlatitude data are characterized by larger diurnal changes, while the NO<sub>2</sub> winter minimum is significantly lower at high than at midlatitude. The winter minima and summer maxima in 1990 and 1995 at both sites are in reasonable agreement with other authors at northern latitudes [e.g., Coffey *et al.*, 1981]. As already reported [Van Roozendael *et al.*, 1994; Goutail *et al.*, 1994], a significant reduction of the NO<sub>2</sub> column amount starting after the eruption of Mt Pinatubo in June 1991 and reaching a maximum during winter 1991/1992 is observed at both latitudes. Similar reductions were reported at northern midlatitude [Koike *et al.*, 1993; Elokhov and Gruzdev, 1993] as well as at southern middle [Johnston *et al.*, 1992] and high latitudes [Solomon *et al.*, 1994]. The progressive recovery of the NO<sub>2</sub> column amounts between 1992 and 1995 can be easily followed in the data of Figure 1. In 1995 the NO<sub>2</sub> level is similar (even slightly larger) to the one measured in 1990.

From these results the monthly NO<sub>2</sub> column amounts in the absence of volcanic aerosols were determined, for Sodankyla, using observations before the Pinatubo eruption (i.e., from January 1990 to June 1991). In the case of Jungfraujoch data, the NO<sub>2</sub> reference was obtained from a combination of pre-Pinatubo and 1995 column measurements, the latter year being considered as free of Pinatubo aerosols (H. Jäger, private communication, 1995). The percentage deviation calculated from these reference values are shown in Figure 2 for morning and evening observations. The uncertainties on these calculated percent changes are related to the natural variability of the NO<sub>2</sub> column which limits both the precision on the NO<sub>2</sub> monthly averages and the precision in the determination of the NO<sub>2</sub> reference. Given the standard deviation of the NO<sub>2</sub> data within a month and assuming a standard deviation of the interannual variations of the same magnitude [Johnston and McKenzie, 1989], error bars shown in Figure 2 were calculated. There is a large uncertainty on the percent changes at 67°N during winter periods, which is due to the large variability of the NO<sub>2</sub> column in winter at high latitude (typically 50%). Part of this variability results from measurement noise, larger for small NO<sub>2</sub> columns, but also from the location of the station alternatively outside and inside the cold polar vortex where the NO<sub>2</sub> abundance is reduced due to (1) conversion of NO<sub>x</sub> into HNO<sub>3</sub> on Polar Stratospheric Clouds (PSCs), (2) the temperature dependence of the gas phase NO<sub>x</sub> chemistry, and (3) possible denitrification known to have occurred at several occasions in the Arctic winter. Within these uncertainties, the maximum NO<sub>2</sub> reduction of about 35% is obtained in January 1992 at both latitudes. It is followed by a progressive recovery lasting until summer 1994. There is no significant reduction measured during winters 1992/1993 and 1993/1994 due to the larger uncertainties on the measurements. As reported by Koike *et al.* [1993] and Johnston *et al.* [1992], NO<sub>2</sub> percent changes seem to be larger for morning



**Figure 2.** Relative change of the monthly averaged morning and evening NO<sub>2</sub> columns, relative to the respective unperturbed pre-Pinatubo reference value (see text), observed at (a) Sodankyla and (b) the Jungfraujoch Station. The vertical dotted line indicates the day of the Mt. Pinatubo eruption.

data than for evening ones at the midlatitude station. However, this difference might be insignificant considering the error bars. There is no difference between morning and evening percent changes measured at Sodankyla, but the amplitude of the diurnal variation is also considerably smaller than at midlatitude.


#### 4. Discussion

The observation of significant reductions of the NO<sub>2</sub> column amount was reported after the eruption of El Chichon in April 1982 [Roscoe *et al.*, 1986; Johnston and McKenzie, 1989; Coffey *et al.*, 1993] and after the eruption of Mount Pinatubo in June 1991 [Johnston *et al.*, 1992; Koike *et al.*, 1993; Elokhov and Gruzdev, 1993; Mills *et al.*, 1993; Goutail *et al.*, 1994; Solomon *et al.*, 1994; Van Roozendael *et al.*, 1994]. Most of these observations were obtained from ground-based UV-visible measurements. Two possible causes of alteration of these measurements were investigated by Perliski and Solomon [1992]. First, the dependence of air mass factors on the presence of volcanic aerosols was studied using a spherical Monte Carlo radiative transfer model. It was shown that NO<sub>2</sub> AMFs were almost insensitive to changes in scattering geometry induced by Pinatubo aerosols. Secondly, the changes in NO<sub>2</sub> photolysis rates due to changes in the radiation field, which might affect NO<sub>2</sub> vertical distribution through a modification of the partitioning between NO and NO<sub>2</sub> [e.g., Michelangeli *et al.*, 1989], were shown to have only a small effect.

The impact of stratospheric aerosols on ozone and NO<sub>2</sub> was investigated by several authors [e.g., Hofmann and Solomon, 1989; Granier and Brasseur, 1992; Bekki and Pyle, 1994; Tie *et al.*, 1994]. The measurements mentioned above

support the theory of NO<sub>2</sub> depletion due to heterogeneous conversion of N<sub>2</sub>O<sub>5</sub> to HNO<sub>3</sub> on the surface of the volcanic aerosol. This interpretation has been confirmed by HNO<sub>3</sub> measurements carried out at Lauder, New Zealand, showing a HNO<sub>3</sub> increase occurring in phase with a NO<sub>2</sub> decrease after the Pinatubo eruption [Koike *et al.*, 1994].

In Figure 3 the measured monthly averaged NO<sub>2</sub> column amounts are compared with calculations at 65° and 45°N obtained from a two-dimensional chemical model [Granier and Brasseur, 1992]. The aerosol climatology used in the model for midlatitudes was constrained by aerosol lidar measurements from Garmisch-Partenkirchen provided by H. Jäger (private communication, 1995). For high latitudes the same climatology was adjusted in phase with a simulation of the Mount Pinatubo aerosol evolution as a function of the latitude and time [Tie *et al.*, 1994]. The 2-D model does not include diurnal variations, therefore morning and evening observed values were averaged in order to facilitate the comparison. Although the mean NO<sub>2</sub> level is rather well reproduced by the calculations, the amplitude of the seasonal variation is largely underestimated, especially at midlatitude. Part of the discrepancy between model and observations can be accounted for by the temperature dependence of the NO<sub>2</sub> absorption cross sections that were measured for the SAOZ instrument at room temperature instead of stratospheric temperature. This could lead to a systematic overestimation of the NO<sub>2</sub> column by 20% [Harwood and Jones, 1994]. It is clear however that the observations does not always exceed the calculations and therefore other factors have to be searched. We believe that one other important reason for the differences between the model and the observations may be due to the fact that the



**Figure 3.** Comparison between experimental and modeled monthly averaged NO<sub>2</sub> column data at (a) Sodankyla and (b) the Jungfraujoch Station. The experimental data considered are the average between morning and evening twilight values except when only reliable evening data are available. The vertical dotted line indicates the day of the Mount Pinatubo eruption.

model does not include the diurnal cycle. The diurnally averaged model used in this work will produce at least two differences with respect to the observations. First, the morning/evening mean values determined from the measurements may not represent the diurnal averaged values calculated in the model over the full day. Second, the diurnal averaged model uses approximations that may have important effects. For example, the real loss rate for NO<sub>2</sub> by reacting with OH should be


$$\int_{\text{day-night}} [\text{OH}][\text{NO}_2].dt \quad (3)$$

while the model uses


$$\overline{[\text{OH}]_{\text{diurnal average}}} \cdot \overline{[\text{NO}_2]_{\text{diurnal average}}} \quad (4)$$

Figure 4 shows a comparison between the observed percent changes in NO<sub>2</sub> column amounts (again diurnally averaged) and the calculated ones. These latter were derived using a model calculation including background aerosol only as pre-Pinatubo reference. In the model results, the main NO<sub>2</sub> decrease starts in November 1991. The largest reduction of about 35% is obtained in January 1992 at middle and high latitudes. These results match quite well the observations both in time and in magnitude. The recovery of the NO<sub>2</sub> column is also captured by the model but, however, at a slower rate, particularly at 46°N. The agreement with the observations is better in 1992, when the aerosol layer extended up to 26 km and the aerosol density was large, than during the following years when the aerosol had partly sedimented. This may indicate a tendency of the model to better simulate conditions of high aerosol loading when the NO<sub>x</sub> reduction process is largely saturated; that is, the production of N<sub>2</sub>O<sub>5</sub> becomes the rate-limiting step in the conversion of NO<sub>x</sub> to HNO<sub>3</sub>. On the contrary, the model appears to overestimate the NO<sub>x</sub> reduction at low aerosol loading. Another disagreement, again more significant at 46°N, is the marked minimum in winter in the simulations of the NO<sub>2</sub> percent changes at the opposite of what is observed. This might relate with the overestimation of the column by the model in winter (Figure 3), possibly due, as discussed above, to the fact that the model does not include the diurnal cycle. The dip in the model results is particularly important at 67°N. However, profile measurements from balloon in Scandinavia in the winter have shown that the NO<sub>2</sub> concentration is dramatically reduced below 20 km just because of the lack of sunlight [Latelin *et al.*, 1994], and therefore it is difficult to understand how aerosol located below 20 km in 1993 and 1994 could have reduced significantly the column.

Finally, it must be mentioned that the model results also indicate that although the NO<sub>2</sub> column change is not significantly sensitive to the heterogeneous reaction rates, it is much more sensitive to the aerosol surface area used in the calculations. Some discrepancies between observations and calculated values may be due to uncertainties in the sulfate aerosol surface areas applied in the model. To illustrate this, Figure 5 displays the results of a sensitivity study of sulfate aerosol loading on the calculation. It shows that if the aerosol loading after the eruption of Mt. Pinatubo is reduced by a factor of 2, the NO<sub>2</sub> reduction becomes less important, and at 45°N, the



**Figure 4.** Comparison between experimental and modeled results for the monthly averaged relative NO<sub>2</sub> column change at (a) Sodankyla and (b) the Jungfraujoch Station. The experimental data considered are the same as in Figure 3. The vertical dotted line indicates the day of the Mount Pinatubo eruption.



**Figure 5.** Sensitivity study showing the variation in calculated relative NO<sub>2</sub> column change at (a) 67°N and (b) 46°N, when reducing by a factor of 2 the aerosol loading after the Mount Pinatubo eruption. Dotted lines in both plots were obtained with half the amount of sulfate aerosols used to calculate solid curves.

calculation is in better agreement with the observations, indicating that uncertainties in the aerosol loading could significantly affect the model results.

The maximum reduction in the NO<sub>2</sub> column determined from Sodankyla and Jungfraujoch data for the northern hemisphere is delayed by three months compared with the reduction of similar amplitude observed at southern midlatitude [Johnston *et al.*, 1992]. This delay is consistent with the different arrival times of the Pinatubo aerosols in the two hemispheres. SAGE II measurements [Trepte *et al.*, 1993] show that by the end of September 1991 the volcanic aerosol had spread to southern midlatitudes at an altitude of about 30 km. At northern midlatitudes, the aerosol cloud reached altitudes above 25 km by the end of December. These satellite observations are in agreement with ground-based lidar measurements reported at northern hemisphere sites [Jäger *et al.*, 1995].

Koike *et al.* [1993] report the largest NO<sub>2</sub> reduction occurring in April 1992 at the northern hemisphere midlatitude station of Moshiri, Japan (44°N). According to these measurements, the maximum NO<sub>2</sub> reduction above Japan appears to be delayed by three months with respect to European midlatitudes. However, the uncertainty in the NO<sub>2</sub> percent changes determined from Moshiri data might be larger because the Pinatubo-free NO<sub>2</sub> values used at that site were deduced from data obtained at 45°S (shifted by six months in order to account for the interhemispheric difference in the phase of the NO<sub>2</sub> seasonal variation). Moreover, anomalously low NO<sub>2</sub> values might have been observed in April 1992 above Japan due to transport of air masses originating in the periphery of the polar vortex [see Koike *et al.*, 1993].

## 5. Conclusion

Ground-based visible measurements of NO<sub>2</sub> abundances covering one year before and four years after the Pinatubo eruption have been carried out at the Jungfraujoch Station (46°N) and at Sodankyla (67°N). The data display a similar behavior at both latitudes, i.e., a reduction of the NO<sub>2</sub> column starting at the end of 1991, reaching a maximum of about 35% in January 1992 and then followed by a progressive recovery until summer 1994. In 1995 the NO<sub>2</sub> abundances appear to be similar to those measured before the eruption.

Measured NO<sub>2</sub> column amounts are qualitatively well reproduced by the 2-D model calculations, including heterogeneous reactions on the observed volcanic aerosols, although somewhat underestimated by the model mainly at midlatitudes. There is a good quantitative agreement between modeled and observed percent changes in NO<sub>2</sub> column. These results confirm the important effect of stratospheric sulfuric acid aerosols on the composition of the stratosphere.

**Acknowledgments.** This work has been supported by the contract EV5V-CT93-0348 from the EC, by the OSTC (contracts GC/35/002 and GC/11/004), the "Nationaal Fonds voor Wetenschappelijk Onderzoek" (FKFO 2.0115.93), and the "Fonds National de la Recherche Scientifique" (FRFC 2.4533.93). X. X. Tie and G. Brasseur are supported in part by DOE Atmospheric Chemistry Program under contract number DE-AI05-94ER619877. C. Granier is partly supported by NASA, under contract number W-18,531. The National Center for Atmospheric Research is sponsored by the National Science Foundation. SAOZ observations are conducted in cooperation with the Finnish Meteorological Institute.

## References

- Bekki, S., and J. A. Pyle, A two-dimensional modeling study of the volcanic eruption of Mount Pinatubo, *J. Geophys. Res.*, **99**, 18,861-18,869, 1994.
- Bluth, G. J. S., C. C. Schetzler, A. J. Krueger, and L. S. Walter, The contribution of explosive volcanism to global atmospheric sulphur dioxide concentrations, *Nature*, **366**, 327-239, 1993.
- Coffey, M. T., and W. G. Mankin, Observations of the loss of stratospheric NO<sub>2</sub> following volcanic eruptions, *Geophys. Res. Lett.*, **20**, 2873-2876, 1993.
- Coffey, M. T., W. G. Mankin, and A. Goldman, Simultaneous spectroscopic determination of the latitudinal, seasonal, and diurnal variability of stratospheric N<sub>2</sub>O, NO, NO<sub>2</sub>, and HNO<sub>3</sub>, *J. Geophys. Res.*, **86**, 7331-7341, 1981.
- Elokhov, A. S., and A. N. Gruzdev, Spectrometric measurements of total NO<sub>2</sub> in different regions of the globe, in *Optical Monitoring of the Environment*, SPIE, **2107**, 111-121, 1993.
- Fahey, D. W., et al., In situ measurements constraining the role of sulfate aerosols in mid-latitude ozone depletion, *Nature*, **363**, 509-514, 1993.
- Goutail, F., J.-P. Pommereau, A. Sarkissian, E. Kyro, and V. Dorokhov, Total nitrogen dioxide at the Arctic polar circle since 1990, *Geophys. Res. Lett.*, **21**, 1371-1374, 1994.
- Granier, C., and G. P. Brasseur, Impact of heterogeneous chemistry on model predictions of ozone changes, *J. Geophys. Res.*, **97**, 18,015-18,033, 1992.
- Harwood, M. H., and R. L. Jones, Temperature dependent ultraviolet-visible absorption cross-sections of NO<sub>2</sub> and N<sub>2</sub>O<sub>4</sub>: Low temperature measurements of the equilibrium constant for 2NO<sub>2</sub>=N<sub>2</sub>O<sub>4</sub>, *J. Geophys. Res.*, **99**, 22,955-22,964, 1994.
- Hofmann, D. J., and S. Solomon, Ozone destruction through heterogeneous chemistry following the eruption of El Chichon, *J. Geophys. Res.*, **94**, 5029-5041, 1989.
- Hofmann, D. J., et al., Intercomparison of UV/visible spectrometers for measurements of stratospheric NO<sub>2</sub> for the Network for the Detection of Stratospheric Change, *J. Geophys. Res.*, **100**, 16,765-16,791, 1995.
- Jäger, H., O. Uchino, T. Nagai, T. Fujimoto, V. Freudenthaler, and F. Homburg, Ground-based remote sensing of the decay of the Pinatubo eruption cloud at three northern hemisphere sites, *Geophys. Res. Lett.*, **22**, 607-610, 1995.
- Johnston, P. V., and R. L. McKenzie, NO<sub>2</sub> observations at 45°S during the decreasing phase of solar cycle 21, from 1980 to 1987, *J. Geophys. Res.*, **94**, 3473-3486, 1989.
- Johnston, P. V., R. L. McKenzie, J. G. Keys, and W. A. Matthews, Observations of depleted stratospheric NO<sub>2</sub> following the Pinatubo volcanic eruption, *Geophys. Res. Lett.*, **19**, 211-213, 1992.
- Koike, M., Y. Kondo, W. A. Matthews, P. V. Johnston, K. Yamazaki, Decrease of stratospheric NO<sub>2</sub> at 44°N caused by Pinatubo volcanic aerosols, *Geophys. Res. Lett.*, **20**, 1975-1978, 1993.
- Koike, M., N. B. Jones, W. A. Matthews, P. V. Johnston, R. L. McKenzie, D. Kinnison, and J. Rodriguez, Impact of Pinatubo aerosols on the partitioning between NO<sub>2</sub> and HNO<sub>3</sub>, *Geophys. Res. Lett.*, **21**, 597-600, 1994.
- Lateltin, E., J.-P. Pommereau, H. Le Texier, M. Pirre, and R. A. Ramarosan, Perturbation of stratospheric nitrogen dioxide by volcanic aerosol in the arctic, *Geophys. Res. Lett.*, **21**, 1411-1414, 1994.
- McCormick, M. P., L. W. Thomason, and C. R. Trepte, Atmospheric effects of the Mt Pinatubo eruption, *Nature*, **373**, 399-404, 1995.
- Michelangeli, D. V., M. Allen, and Y. Yung, El Chichon volcanic aerosols: Impact of radiative, thermal, and chemical perturbations, *J. Geophys. Res.*, **94**, 18,429-18,443, 1989.
- Mills, M. J., A. O. Langford, T. J. O'Leary, K. Arpag, H. L. Miller, M. H. Proffit, R. W. Sanders, and S. Solomon, On the relationship between stratospheric aerosols and nitrogen dioxide, *Geophys. Res. Lett.*, **20**, 1187-1190, 1993.
- Noxon, J. F., Stratospheric NO<sub>2</sub>, 2, Global behavior, *J. Geophys. Res.*, **84**, 5067-5076, 1979.
- Noxon, J. F., Correction, *J. Geophys. Res.*, **85**, 4560-4561, 1980.
- Perliski, L. M., and S. Solomon, Radiative influences of Pinatubo volcanic eruption aerosols on twilight observations of NO<sub>2</sub> column abundances, *Geophys. Res. Lett.*, **19**, 1923-1926, 1992.
- Pommereau, J.-P., and J. Piquard, Ozone and nitrogen dioxide vertical distributions by UV-visible solar occultation from balloons, *Geophys. Res. Lett.*, **21**, 1227-1230, 1994.
- Prather, M. J., Catastrophic loss of stratospheric ozone in dense volcanic clouds, *J. Geophys. Res.*, **97**, 10,187-10,191, 1992.
- Rinsland, C. P., M. R. Gunson, M. C. Abrams, L. L. Lowes, R. Zander, E. Mahieu, A. Goldman, M. K. W. Ko, J. M. Rodriguez, and N. D. Sze, Heterogeneous conversion of N<sub>2</sub>O<sub>5</sub> to HNO<sub>3</sub> in the post-Mount Pinatubo eruption stratosphere, *J. Geophys. Res.*, **99**, 8213-8219, 1994.
- Roscoe, H. K., B. J. Kerridge, L. J. Gray, R. J. Wells, and J. Pyle, Simultaneous measurements of stratospheric NO and NO<sub>2</sub> and their comparison with model predictions, *J. Geophys. Res.*, **91**, 5405-5419, 1986.
- Solomon, S., A.L. Schmeltekopf, R.W. Sanders, On the interpretation of zenith-sky absorption measurements, *J. Geophys. Res.*, **92**, 8311-8319, 1987.
- Solomon, S., R. W. Sanders, R. O. Jakoubek, K. H. Arpag, S. L. Stephens, J. G. Keys, and R. R. Garcia, Visible and near-ultraviolet spectroscopy at McMurdo Station, Antarctica. 10. Reductions of stratospheric NO<sub>2</sub> due to Pinatubo aerosols, *J. Geophys. Res.*, **99**, 3509-3516, 1994.
- Solomon, S., R. W. Portmann, R. R. Garcia, L. W. Thomason, L. R. Poole, M. P. McCormick, Role of aerosol variations in anthropogenic ozone depletion at northern midlatitudes, *J. Geophys. Res.*, **101**, 6713-6727, 1996.
- Tie, X.X., G. P. Brasseur, B. Briegleb, and C. Granier, Two-dimensional simulation of Pinatubo aerosol and its effect on stratospheric ozone, *J. Geophys. Res.*, **99**, 20,545-20,562, 1994.
- Trepte, C. R., R. E. Veiga, and M. P. McCormick, The poleward dispersal of Mount Pinatubo volcanic aerosol, *J. Geophys. Res.*, **98**, 18,563-18,573, 1993.
- Van Roozendael, M., C. Hermans, M. De Mazière, and P. C. Simon, Stratospheric NO<sub>2</sub> Observation at the Jungfrauoch Station between June 1990 and May 1992, *Geophys. Res. Lett.*, **21**, 1383-1386, 1994a.
- Van Roozendael, M., M. De Mazière and P. C. Simon, Ground-based visible measurements at the Jungfrauoch Station since 1990, *J. Quant. Spectrosc. Radiat. Transfer*, **52**, 231-240, 1994b.

G.Brasseur, C. Granier, and X. X. Tie, National Center for Atmospheric Research, Boulder CO 80307. (e-mail: brasseur@acd.ucar.edu; granier@acd.ucar.edu; xxtie@ra.cgd.ucar.edu)

F. Goutail and J.-P. Pommereau, Service d'Aéronomie du CNRS, Verrières-le-Buisson, France. (e-mail: florence.goutail@aerov.jussieu.fr; pommereau@aerov.jussieu.fr)

M. De Mazière, C. Hermans, M. Van Roozendael, and P. C. Simon, Belgian Institute for Space Aeronomy, 3 Av. Circulaire, B-1180 Brussels, Belgium. (e-mail: martine@bira-iasb.oma.be; christh@bira-iasb.oma.be; michelv@bira-iasb.oma.be; pauls@bira-iasb.oma.be)

(Received January 3, 1996; revised February 7, 1997; accepted April 10, 1997.)