COMPACTNESS OF SELF-SHRINKERS IN $\mathbb{R}^3$ WITH FIXED GENUS

AO SUN AND ZHICHAO WANG

ABSTRACT. We prove the compactness of self-shrinkers in $\mathbb{R}^3$ with bounded entropy and fixed genus. As a corollary, we show that numbers of ends of such surfaces are uniformly bounded by the entropy and genus.

1. INTRODUCTION

A hypersurface $\Sigma \subset \mathbb{R}^{n+1}$ is called a *self-shrinker* if it satisfies the equation

$$(1.1) \quad H = \frac{1}{2} \langle x, \mathbf{n} \rangle,$$

where H is the mean curvature of the hypersurface, x is the position vector on the hypersurface, and $\mathbf{n}$ is the unit normal vector of Σ at x .

Self-shrinkers are the homothetic solitons of mean curvature flow. That is, suppose Σ is a self-shrinker, then $\{\sqrt{-t}\Sigma\}_{t \in (-\infty, 0)}$ is a solution to the mean curvature flow. Moreover, by using Huisken's monotonicity formula in [26], Ilmanen [28] and White [40] proved that self-shrinkers are models for singularities of mean curvature flows.

The simplest self-shrinkers in $\mathbb{R}^3$ are the planes, the spheres, and the cylinders. Brendle [8] proved that these are the only possible self-shrinkers with genus 0. Angenent [2] constructed a genus 1 self-shrinker which is known to be the Angenent donut. For higher genus, Kapouleas-Kleene-Møller [29] and Nyugen [33] constructed a number of examples of non-compact self-shrinkers by a gluing method.

In this paper, we prove the compactness of self-shrinkers in $\mathbb{R}^3$ with fixed genus.

Theorem 1.1. *Let $\{\Sigma_i\} \subseteq \mathbb{R}^3$ be a sequence of embedded self-shrinkers with genus g and uniformly bounded entropy. Then there exist a subsequence (still denoted by $\{\Sigma_i\}$) and a self-shrinker Σ with genus g such that Σ_i converges to Σ in $C_{loc}^\infty(\mathbb{R}^3)$.*

Here entropy is defined to be the Gaussian area of self-shrinkers (see (2.3) for the precise definition).

Given $\Lambda > 0$ and a non-negative integer g , we define $\mathcal{M}_S(\Lambda, g)$ to be the space of self-shrinkers in $\mathbb{R}^3$ with genus g and entropy no more than Λ . For the sake of brevity, set

$$(1.2) \quad \mathcal{M}_S(g) = \bigcup_{\Lambda > 0} \mathcal{M}_S(\Lambda, g).$$

Then our main theorem is equivalent to say that $\mathcal{M}_S(\Lambda, g)$ is compact under $C_{loc}^\infty(\mathbb{R}^3)$ topology.

The smooth compactness theorem of $\bigcup_{k=0}^g \mathcal{M}_S(\Lambda, k)$ was first developed by Colding-Minicozzi in [19], see also some further generalizations in [10, 11]. However, it is not known

that whether the topological type of the self-shrinkers would change in the limit. For example, when passing to limit, a “hole” on a sequence of self-shrinkers may move further and further, and vanishes at infinity when passing to limit. Our theorem rules out this possibility, and improves the understanding of the structure of self-shrinkers with finite genus.

As a result, a direct corollary of our main theorem is

Corollary 1.2. *Given $\Lambda > 0$, $g > 0$, there exists $R = R(\Lambda, g) > 0$ such that for any $\Sigma \in \mathcal{M}_S(\Lambda, g)$, $\Sigma \setminus B_R(0)$ consists of finite union of topological annuli and topological disks.*

In other words, we prove that the “holes” on a self-shrinker can not stay far away from the origin of $\mathbb{R}^3$.

The classification of self-shrinkers’ ends plays an important role in our paper. Such a result has been firstly studied in [37] by L. Wang, who used a Carleman type technique to prove that two self-shrinkers’ ends which are asymptotic to the same cone must identically coincide with each other. Later in [39], L. Wang proved that two self-shrinkers’ ends which are asymptotic to the same cylinder with certain decay rate must identically coincide with each other. As a result, the asymptotic behavior of noncompact self-shrinkers at infinity is quite clear. Recently, L. Wang classified the self-shrinkers’ ends with finite topology in [38], which is the most important structure theory of self-shrinkers in $\mathbb{R}^3$.

It may be interesting to compare the structure theory of self-shrinkers with the structure theory of minimal surfaces in $\mathbb{R}^3$, cf. Colding-Minicozzi [14–17, 20], Meeks-Rosenberg [32]. The structure of minimal surfaces with finite topology and finite total curvature has been well studied, and the ends of minimal surfaces look like planes or helicoids at infinity. This is very different from the case of self-shrinkers.

The key step in [38] is to prove the multiplicity of a tangent flow of a self-shrinker equals to 1. Then the classification theorem follows from Brakke’s Regularity Theorem (see [7, 41]). Note that the argument in [38] works for all $\Sigma \in \mathcal{M}_S(\Lambda, g)$. Consequently, Σ can only have finitely many ends (see Lemma A.2). Our theorem in this paper implies that the number of ends are bounded by a constant depending only on Λ and g .

Corollary 1.3. *There exists a constant $C = C(\Lambda, g)$ such that for all $\Sigma \in \mathcal{M}_S(\Lambda, g)$, the number of ends of Σ is no more than C .*

Another application of Theorem 1.1 is the existence of entropy minimizer in $\mathcal{M}_S(g)$. It follows from the definition of entropy that the plane $\mathbb{R}^2$ minimizes the entropy among all surfaces in $\mathbb{R}^3$. Among all closed self-shrinkers in $\mathbb{R}^3$, the round sphere is the only one achieves the least entropy for all dimensions, which was proven by Colding-Ilmanen-Minicozzi-White in [21]. After that, Bernstein-Wang [5] confirmed that the cylinders have the third least entropy among all self-shrinkers in $\mathbb{R}^3$. As a corollary of our main theorem, we show the existence of an entropy minimizer with fixed genus g .

Corollary 1.4. *There exists an entropy minimizer in $\mathcal{M}_S(g)$ whenever $\mathcal{M}_S(g) \neq \emptyset$.*

We remark that, Bernstein-Wang [4] proved that the entropy of closed hypersurfaces is minimized by round spheres among all closed surfaces in $\mathbb{R}^{n+1}$ with $n \leq 6$. Later Zhu

generalized this result to all dimensions in [43]. In $\mathbb{R}^3$, Ketover-Zhou [30] gave an alternative proof using min-max theory.

This kind of minimizing problem is of great interest in the study of geometry. For example, Simon [34] proved that for a fixed genus number g (see also [3]), there is a genus g Willmore surface in $\mathbb{R}^3$ minimizes Willmore energy among all closed surfaces with genus g . In contrast, for a sequence of minimal surfaces or constant mean curvature surfaces, the topology may change when passing to a limit. Therefore in general it is hard to find an extremal surface with fixed genus.

Outline of the proof. By the compactness of $\cup_{k \leq g} \mathcal{M}_S(\Lambda, k)$ proved by Colding-Minicozzi in [19], Σ_i converges to Σ smoothly in arbitrarily large balls B_R . This fact implies that $\Sigma_i \cap B_R$ is diffeomorphic to $\Sigma \cap B_R$ when i sufficiently large. Thus we only need to analyze Σ_i outside a large ball. By Wang's result [38, Theorem 1.1], $\Sigma \setminus B_R(0)$ can be decomposed to finitely many annuli. Moreover, each connected component is either asymptotic to a regular cone or a round cylinder. We will call them *conical* or *cylindrical* ends respectively. Our purpose is to prove that the connected components of $\Sigma_i \setminus B_R$ are topological annuli or topological disks. There are two cases.

Firstly, consider that Γ is a connected component of $\Sigma \setminus B_R(0)$ which is conical. $\Sigma_i \setminus B_R(0)$ can also be decomposed into several connected components. We consider the one which is close to Γ , and denote it by Γ_i . By the monotonicity formulas of F -functional near the asymptotic cone, we can prove that Γ_i completely lies in a tubular neighborhood of Γ (see Lemma 4.1). If $|x|^2$ has no critical points in such a neighborhood, then by Morse theory we know that Γ_i has to be a topological annulus. So we only need to prove that $|x|^2$ has no critical points on Γ_i .

To prove that $|x|^2$ has no critical points on Γ_i , we argue it by contradiction. Suppose $p_i \in \Gamma_i$ is a critical point of the function $|x|^2$. Then Σ_i locally smoothly converging to Σ implies that $|p_i| \rightarrow \infty$. We can blow up these ends at these critical points, and the limit would be a translating soliton which is not a plane. Moreover it is a Bowl soliton (see Theorem 2.8). However, the monotonicity formulas show that the entropy of the translator should be bounded by the F -functional of Σ at $\partial\Gamma$ (see Proposition 2.7), which is closed to 1 since Γ is asymptotic to a regular cone. Recall that the entropy of a Bowl soliton equals to λ_1 (the entropy of S^1 ; see [24]) which is larger than 1. Thus we get a contradiction. So Γ_i is diffeomorphic to an annulus.

Next consider that Γ is a cylindrical end with direction y . The simplest case is that Γ_i has an end which is also asymptotic to a cylinder. Denote the direction of a cylindrical end of Γ_i by y_i , then $y_i \rightarrow y$ (see Lemma 4.1). For any sequence $\rho_i \rightarrow \infty$, we claim that $\Sigma_i - \rho_i y_i$ locally smoothly converges to a self-shrinking cylinder. Indeed, by monotonicity formulas, F -functional on $\Sigma_i - \rho_i y_i$ is greater or equal than λ_1 (see Proposition 2.7). On the other hand, the limit (denoted by ν) of $\Sigma_i - \rho_i y_i$ has F -functional less than or equal to the F -functional of $\Sigma - \rho y$, which converges to λ_1 as $\rho \rightarrow \infty$. Thus $\Sigma_i - \rho_i y_i$ converges to an F -stationary varifold with entropy λ_1 . Then the claim follows from Brakke's regularity theorem (see [7, 41] or Theorem 2.6) if ν is supporting on a smooth self-shrinker with multiplicity 1.

It suffices to show the smoothness of ν . Though Ilmanen [28] shows that the singularities of mean curvature flow of surfaces in $\mathbb{R}^3$ is smooth, we can not follow his argument to prove that ν supports on a smooth self-shrinker. The reason is that we need to prove the smoothness of the limits for all $\rho_i \rightarrow \infty$, and then the integral of H^2 is hard to control.

To prove ν is smooth, we show ν splits off a line and then the claim follows from the entropy bound. Indeed, the limit is also F -stationary if we take limits for $\Sigma - (\rho_i + \tau)y_i$ for each $\tau \in \mathbb{R}$. Thus $\nu - \tau y$ is also F -stationary, which implies that ν splits off a line. The entropy of ν equals to λ_1 , which implies that ν is supporting on a self-shrinking cylinder.

Consequently, $\Sigma_i - \rho_i y_i$ converges to a self-shrinker for any sequence of $\rho_i \rightarrow \infty$. Combining with the fact that Σ_i locally smoothly converges to Σ , we can prove that Γ_i is diffeomorphic to Γ (see step 4 of Proposition 5.1 for details).

When Γ_i consists of conical ends or it is compact, the idea is the same but a suitable blow up/down argument is necessary. We refer to Proposition 5.5 and 5.7 for details.

The paper is organized as follows: In Section 2, we summarize some known results in mean curvature flows and Brakke flows. Then we give an upper bound for entropy after blowing up. Using this, we show that the Bowl soliton is the only non-trivial blowing up limit. In Section 3, another version (Theorem 3.1) of Theorem 1.1 will be stated. Using this, the proof of Corollary 1.2 and 1.3 follows. The Section 4 is devoted to the first part of Theorem 3.1, which deals with the case of conical ends. In Section 5, we prove the second part of Theorem 3.1, which deals with the case of cylindrical ends. In Appendix A, we recall L. Wang's classification result of ends [38]. In Appendix B, we recall the classification result of low entropy translating solitons in [25].

Acknowledgement. We want to thank Professor Bill Minicozzi and Professor Lu Wang for the helpful discussion and comments. Z.W. would like to thank Professor Gang Tian and Professor Xin Zhou for constant encouragement. A.S. wants to thank Kyeongsu Choi for the helpful discussion. We also would like to thank Jonathan Zhu for bringing our attention to the paper of Bernstein-Wang [6].

2. PRELIMINARIES

In this section, we summarize some known results which are used in this paper. The experts should feel free to skip this section.

2.1. F -functional and monotonicity formulas. Let us recall Huisken's monotonicity formulas. Given $(y, s) \in \mathbb{R}^{n+1} \times \mathbb{R}$, let $\Phi_{y,s}$ be the backward heat kernel defined on $\mathbb{R}^{n+1} \times (-\infty, s)$:

$$(2.1) \quad \Phi_{y,s}(x, t) = (4\pi(s-t))^{-\frac{n}{2}} e^{-\frac{|x-y|^2}{4(s-t)}}.$$

We will omit the subscript if $(y, s) = (0, 0)$

For any hypersurface Σ in $\mathbb{R}^{n+1}$ and $(x_0, t_0) \in \mathbb{R}^{n+1} \times (0, +\infty)$, the F -functional is defined by

$$(2.2) \quad F_{x_0, t_0}(\Sigma) = (4\pi t_0)^{-\frac{n}{2}} \int_{\Sigma} e^{-\frac{|x-x_0|^2}{4t_0}} = \int_{\Sigma} \Phi(x - x_0, -t_0).$$

We say Σ is a *self-shrinker* in $\mathbb{R}^{n+1}$ if it satisfies the equation,

$$H = \frac{\langle x, \mathbf{n} \rangle}{2}.$$

By monotonicity formulas [18, 1.9], for a self-shrinker Σ , $V \in \mathbb{R}^{n+1}$ and $a \in \mathbb{R}$, $F_{sV, 1+as^2}(\Sigma)$ is monotonically decreasing in s whenever $1 + as^2 > 0$.

The *entropy* of Σ is defined by

$$(2.3) \quad \lambda(\Sigma) := \sup_{(x_0, t_0) \in \mathbb{R}^3 \times (0, \infty)} F_{x_0, t_0}(\Sigma).$$

In this paper, we always denote the entropy of a self-shrinking cylinder in $\mathbb{R}^3$ by λ_1 .

The entropy is a quantity characterizing all the scales of a hypersurface. This fact is suggested by the following useful proposition (see [11, 18]).

Proposition 2.1. *There is a constant C such that for any embedded hypersurface Σ in $\mathbb{R}^{n+1}$,*

$$(2.4) \quad \sup_{x \in \mathbb{R}^{n+1}} \sup_{r > 0} \frac{\text{Area}(\Sigma \cap B_r(x))}{r^n} \leq C \lambda(\Sigma),$$

and

$$(2.5) \quad \lambda(\Sigma) \leq C \sup_{x \in \mathbb{R}^{n+1}} \sup_{r > 0} \frac{\text{Area}(\Sigma \cap B_r(x))}{r^n}.$$

2.2. Brakke flows. We refer to [28; 38, §2] for the definition of Brakke flows. The following monotonicity formula is similar to the monotonicity formula of mean curvature flow:

Lemma 2.2 (Monotonicity Formulas for Brakke Flows, [28, Lemma 7]). *Let $\{\mu_t\}_{t \in (a, b)}$ be a Brakke flow satisfying*

$$\sup_{x \in \mathbb{R}^{n+1}} \sup_{R > 0} \frac{\mu_t(B_R(x))}{R^n} \leq C < +\infty.$$

Then for all $(y, s) \in \mathbb{R}^{n+1} \times (a, b)$, and all $a \leq t_1 \leq t_2 < s$,

$$\begin{aligned} & \int_{t_1}^{t_2} \int \Phi_{y, s}(x, t) \left| H - \frac{(x - y)^\perp}{2(s - t)} \right|^2 d\mu_t(x) dt \\ & \leq \int \Phi_{y, s}(x, t_1) d\mu_{t_1}(x) - \int \Phi_{y, s}(x, t_2) d\mu_{t_2}(x). \end{aligned}$$

Remark 2.3. For a Radon measure μ , we can also compute the F -functional, which is defined by

$$F_{x, t}(\mu) := \int (4\pi t)^{-\frac{n}{2}} e^{-\frac{|y-x|^2}{4t}} d\mu(y).$$

Note that if $\mu = \mathcal{H}^n \llcorner \Sigma$ for a hypersurface Σ , then $F_{x, t}(\mu) = F_{x, t}(\Sigma)$.

The following proposition will be used frequently in our argument. This result is well-known and we refer to [36] for a proof.

Proposition 2.4. *Let $\{\mu_i\}$ be a sequence of Radon measures and*

$$\limsup_{i \rightarrow \infty} \sup_{(x, t) \in \mathbb{R}^{n+1} \times (0, +\infty)} F_{x, t}(\mu_i) \leq C_0 < +\infty.$$

Suppose that $\mu_i \rightarrow \mu$. Then $\lim_{i \rightarrow \infty} F_{x, t}(\mu_i) = F_{x, t}(\mu)$.

We need the compactness theorem for Brakke flows in the following sections.

Lemma 2.5 (Compactness of Brakke flows, [27, Lemma 7.1]). *Let $\{\{\mu_t^i\}_{t \in [a,b]}\}$ be a sequence of Brakke flows so that for all bounded open $U \subset \mathbb{R}^{n+1}$,*

$$(2.6) \quad \limsup_{i \rightarrow \infty} \sup_t \mu_t^i(U) \leq C(U) < +\infty.$$

Then there is a subsequence $\{i_k\}$ and an integral Brakke flow $\{\mu_t\}_{t \in [a,b]}$, so that $\mu_t^{i_k} \rightarrow \mu_t$ in the sense of Radon measures as $k \rightarrow \infty$.

The following theorem is proven by White in [41], which is a consequence of Brakke's local regularity theorem in [7]. Our statement here is a special case of [41].

Theorem 2.6 ([41, Theorem 7.3]). *Let $\{\Sigma_t^i\}_{t \in (a,b)}$ be a sequence of mean curvature flows in $\mathbb{R}^{n+1}$ ($n \geq 2$). Suppose that the corresponding Brakke flows $\{\mu_t^i\}$ (where $\mu_t^i := \mathcal{H}^n \llcorner \Sigma_t^i$) converge to a Brakke flow μ_t as Radon measures. If μ_t is a mean curvature flow measure for $t \in (a,b)$, that is, μ_t supports on a smooth manifold Σ_t with multiplicity 1, then Σ_t^i locally smoothly converge to Σ_t for $t \in (a,b)$.*

2.3. Blow-up and blow-down arguments. In this paper, $y \in \mathbb{R}^3$ also represents a constant vector field. Denote the line in $\mathbb{R}^3$ containing y by $\mathbb{R}_y$. For $a, b \in \mathbb{R}$, set

$$(2.7) \quad I_y(a, b) = \{ty \mid t \in (a, b)\}.$$

We will use $\mathbb{R}_y$ to denote $I_y(-\infty, \infty)$.

The following proposition will be used frequently in this paper.

Proposition 2.7. *Let $\{\Sigma_i\} \subseteq \mathcal{M}_S(\Lambda, g)$ (see (1.2)) and Σ_i locally smoothly converges to Σ . Suppose $y_i \in S^2(1)$ and $y_i \rightarrow y$. For any $\rho_i \rightarrow +\infty$ and $a_i > 0$ satisfying $\rho_i a_i \rightarrow +\infty$, define*

$$\mu = \lim_{i \rightarrow \infty} \mathcal{H}^2 \llcorner a_i(\Sigma_i - \rho_i y_i).$$

Then for any $(x, t) \in \mathbb{R}^3 \times (0, +\infty)$,

$$F_{x,t}(\mu) = \lim_{i \rightarrow \infty} F_{x,t}(a_i(\Sigma_i - \rho_i y_i)) \leq \lim_{\rho \rightarrow \infty} F_{\rho y, 1}(\Sigma).$$

Proof. The proof is based on the continuity of F -functional (Proposition 2.4) and monotonicity formulas (§2.1). Indeed,

$$\begin{aligned} F_{x,t}(\mu) &= \lim_{i \rightarrow \infty} F_{x,t}(a_i(\Sigma_i - \rho_i y_i)) \\ &= \lim_{i \rightarrow \infty} F_{0,1}\left(\frac{a_i(\Sigma_i - \rho_i y_i) - x}{\sqrt{t}}\right) \\ &= \lim_{i \rightarrow \infty} F_{\rho_i(y_i + x/(\rho_i a_i)), [\rho_i^2(t/a_i^2 - 1)/\rho_i^2] + 1}(\Sigma_i) \\ &\leq \limsup_{\rho \rightarrow \infty} \limsup_{i \rightarrow \infty} F_{\rho(y_i + x/(\rho_i a_i)), [\rho^2(t/a_i^2 - 1)/\rho_i^2] + 1}(\Sigma_i) \\ &= \limsup_{\rho \rightarrow \infty} F_{\rho y, 1}(\Sigma) \quad (\text{since } \rho_i \rightarrow \infty \text{ and } a_i \rho_i \rightarrow \infty) \\ &= \lim_{\rho \rightarrow \infty} F_{\rho y, 1}(\Sigma) \quad (\text{by monotonicity formulas}). \end{aligned}$$

Thus we complete the proof. $\square$

Theorem 2.8. *Suppose $\{\Sigma_i\} \subseteq \mathcal{M}_S(\Lambda, g)$ (see (1.2)) and $p_i \in \Sigma_i$ such that $|p_i| \rightarrow \infty$. Then the blow-up sequence*

$$\tilde{\Sigma}_i = \frac{|p_i|}{2}(\Sigma_i - p_i)$$

locally smoothly converges to a Bowl soliton $\tilde{\Sigma}$ with direction $-y$, or $\mathbb{R}_y \times \mathbb{R}$ up to a subsequence. Here y is the limit point of $p_i/|p_i|$. Moreover if there is a constant $\delta > 0$ such that $|A|(p_i) \geq \delta|p_i|$, then $\tilde{\Sigma}$ is not a plane.

Proof. By Lemma A.4, we conclude that there exists a constant C such that Σ_i has uniformly curvature bound

$$|A^{\Sigma_i}| \leq C(1 + |x|),$$

where A^{Σ_i} is the second fundamental form of Σ_i . Then the rescaling sequence $\tilde{\Sigma}_i$ has a locally uniformly curvature bound, which implies that there is a subsequence of $\tilde{\Sigma}_i$ locally smoothly converging to $\tilde{\Sigma}$. Note $\tilde{\Sigma}_i$ satisfies the equation

$$\tilde{H} = \left\langle \frac{2x}{|p_i|^2}, \mathbf{n} \right\rangle + \left\langle \frac{p_i}{|p_i|}, \mathbf{n} \right\rangle.$$

Passing to a limit, then $\tilde{\Sigma}$ satisfies the equation

$$\tilde{H} = \langle y, \mathbf{n} \rangle,$$

where $y = \lim_{i \rightarrow \infty} p_i/|p_i|$. Hence the limit $\tilde{\Sigma}$ is a translating soliton.

By Proposition 2.7,

$$F_{x,t}(\tilde{\Sigma}) \leq \lim_{\rho \rightarrow \infty} F_{\rho y, 1}(\Sigma) \leq \lambda_1.$$

This entropy bound implies that $\tilde{\Sigma}$ is a Bowl soliton or $\mathbb{R}_y \times \mathbb{R}$ by Theorem B.1 (see also Choi-Haslhofer-Hershkovits [13, Theorem 1.2]).

Now we suppose that there is $\delta > 0$ such that $|A|(p_i) \geq \delta|p_i|$. Then after rescaling we conclude that $|\tilde{A}|(0) \geq \delta > 0$. Thus the limit cannot be a plane. $\square$

Remark 2.9. The results in this section show a part of the correspondence of self-shrinkers and translating solitons which have been pointed out by Ilmanen in [27, Appendix J]. We believe that there would be a more general and uniform arguments to set up this kind of correspondence.

3. MAIN RESULTS

For a subset $U \subset \mathbb{R}^3$ and $r \geq 0$, define the related conical end as

$$(3.1) \quad \mathcal{C}_{U,r} := \{tx : x \in U, t \geq r\}.$$

We also use $\mathcal{C}_U$ to if $r = 0$.

In this section, $\{\Sigma_i\}$ is always assumed to be a sequence in $\mathcal{M}_S(\Lambda, g)$. By [19], there exists $\Sigma \in \cup_{k \leq g} \mathcal{M}_S(\Lambda, k)$ such that $\Sigma_i \rightarrow \Sigma$ locally smoothly up to a subsequence (still denoted by $\{\Sigma_i\}$). By L. Wang [38] (see our statement in Theorem A.1),

$$S^2 \cap \lim_{\rho \rightarrow \infty} \rho^{-1} \Sigma$$

consists of disjoint simple closed curves and points. Therefore, there exists large R such that $\Sigma \setminus B_R(0)$ consists of self-shrinking ends $\Gamma_1, \Gamma_2, \dots, \Gamma_k$.

We say that $\mathcal{C}_{U,R}$ is a *conical neighborhood* of Γ_i if $\mathcal{C}_{U,R} \cap \Sigma = \Gamma_i$.

The key ingredient is the following theorem.

Theorem 3.1. *Let $\{\Sigma_i\}$ be a sequence of self-shrinkers in $\mathcal{M}_S(\Lambda, g)$ (see (1.2) for definition) that locally smoothly converges to Σ . Suppose that there exist an open set $U \subset S^2(1)$ and $R > 0$ such that $\mathcal{C}_{U,R} \cap \Sigma$ is a self-shrinker end Γ . Then up to a subsequence, one of the following happens:*

- (1) Γ is asymptotic to a cone and $\Sigma_i \cap \mathcal{C}_{U,R}$ is diffeomorphic to Γ ;
- (2) Γ is asymptotic to a cylinder, and $\Sigma_i \cap \mathcal{C}_U$ is diffeomorphic to Γ or a disk.

The proof of Theorem 3.1 will be postponed to §4 and §5.

In the rest of this section, we use Theorem 3.1 to prove our main results of this paper. We also need Lemma A.2 in the Appendix. Let us state it here.

Lemma 3.2 (Lemma A.2). *Let M be a self-shrinker with finite entropy and genus. Then M has finite ends.*

Now we are willing to prove our main results.

Proof of Theorem 1.1. Given a complete surface M in $\mathbb{R}^3$, denote the number of genus of M by $g(M)$.

Since Σ has finite genus, then the classification of ends in [37] and Lemma A.2 implies that we can take R large enough such that

$$g(B_R(0) \cap \Sigma) = g(\Sigma).$$

Note that Σ_i locally smoothly converges to Σ . Hence for large i ,

$$g(B_R(0) \cap \Sigma_i) = g(B_R(0) \cap \Sigma).$$

Now by Theorem 3.1, $\Sigma_i \cap \mathcal{C}_{U,R}$ is diffeomorphic to $S^1 \times \mathbb{R}$ or a disk. Hence

$$g(\Sigma_i) = g(\Sigma_i \cap B_R(0)) = g(\Sigma).$$

Thus we complete the proof. $\square$

Proof of Corollary 1.2. Assume on the contrary that $\{\Sigma_i\} \subseteq \mathcal{M}_S(\Lambda, g)$ satisfy the following: for any $R > 0$, there exists $i > 0$ such that one of the connected components of $\Sigma_i \setminus B_R(0)$ is not diffeomorphic to an annulus or a disk.

By Theorem 1.1, Σ_i locally smoothly converge to $\Sigma \in \mathcal{M}_S(\Lambda, g)$. Then there exists $R_0 > 0$, such that $\Sigma \setminus B_{R_0}(0)$ consists of self-shrinker ends. Then by Theorem 3.1, each connected component of $\Sigma_i \setminus B_R(0)$ is diffeomorphic to an annulus or a disk, which leads to a contradiction. $\square$

Given a complete surface M in $\mathbb{R}^3$, denote the number of ends of M by $E(M)$. Now we prove the uniform bound of E .

Proof of Corollary 1.3. Assume on the contrary that $\{\Sigma_i\} \subseteq \mathcal{M}_S(\Lambda, g)$ and $E(\Sigma_i) \rightarrow \infty$. By Theorem 1.1, there is a subsequence of $\{\Sigma_i\}$ (still denoted by $\{\Sigma_i\}$) locally smoothly converges to $\Sigma \in \mathcal{M}_S(\Lambda, g)$. Then by Theorem 3.1, $E(\Sigma) \geq E(\Sigma_i)$ for large i . Hence

$$E(\Sigma) \geq \liminf_{i \rightarrow \infty} E(\Sigma_i) = +\infty,$$

which contradicts Lemma A.2. □

4. ASYMPTOTIC TO A CONE

Now we are going to prove Theorem 3.1.

This section is devoted to the case that $\Sigma \cap \mathcal{C}_{U,R}$ (see (3.1) for notations) is asymptotic to a cone. The purpose is to show that $\Sigma_i \cap \mathcal{C}_{U,R}$ is also a self-shrinker's end and it is asymptotic to a cone when i is large enough. We first show that Σ_i does not intersect $\mathcal{C}_{\partial U,R}$ for large i .

Lemma 4.1. *Let $\{\Sigma_i\}$ be a sequence of self-shrinkers in $\mathcal{M}_S(\Lambda, g)$ (see (1.2) for definition) that locally smoothly converges to Σ . Suppose that $\Sigma \setminus B_R(0) \subset \mathcal{C}_{U,R}$ for some real number $R > 0$ and an open set $U \subset S^2(1)$. Then $\Sigma_i \setminus B_R(0) \subset \mathcal{C}_{U,R}$ for all large i . Particularly, $\lim_{\rho \rightarrow \infty} \rho^{-1} \Sigma_i \cap S^2(1)$ converges to a subset of $\lim_{\rho \rightarrow \infty} \rho^{-1} \Sigma \cap S^2(1)$ in the Hausdorff distance.*

Proof. Assume on the contrary that there exists $\rho_i y_i \in \Sigma_i$ (here $|y_i| = 1$) so that $\rho_i > R$ and $\rho_i y_i \notin \mathcal{C}_{U,R}$. Denote by y the limit of y_i . Then $y \notin U$. Since Σ_i locally smoothly converges to Σ , we must have $\rho_i \rightarrow \infty$.

Take $a_i \rightarrow 0$ so that

$$0 < 1 + a_i \rho_i^2 \ll 1.$$

Because Σ_i is a smooth surface, we must have

$$F_{\rho_i y_i, 1 + a_i \rho_i^2}(\Sigma_i) > 1 - 1/i.$$

By the monotonicity formulas in Section 2.1,

$$F_{\rho y, 1}(\Sigma) = \lim_{i \rightarrow \infty} F_{\rho y_i, 1 + a_i \rho^2}(\Sigma_i) \geq \liminf_{i \rightarrow \infty} F_{\rho_i y_i, 1 + a_i \rho_i^2}(\Sigma_i) \geq 1,$$

which contradicts that $\Sigma \setminus B_R(0) \subset \mathcal{C}_{U,R}$. □

In the rest of this section, we prove the first part of Theorem 3.1.

Theorem 4.2. *Let Σ_i, Σ, U and Γ be the same as in Theorem 3.1. Suppose that there is a curve $\gamma \subseteq S^2(1)$ such that*

$$\lim_{\rho \rightarrow \infty} \rho^{-1} \Sigma \cap U = \gamma,$$

i.e. $\Sigma \cap \mathcal{C}_{U,R}$ is a conical end for some $R > 0$. Then there exist $R > 0, i_0 > 0$ such that $\Sigma_i \cap \mathcal{C}_{U,R}$ is diffeomorphic to $S^1 \times [0, +\infty)$ for $i \geq i_0$.

Proof. Let us consider the function $|x|^2$ defined on Γ_i . Take R large enough such that $\Sigma \cap \mathcal{C}_{U,R}$ is diffeomorphic to $S^1 \times [0, +\infty)$.

If $|x|^2$ has no critical points in $\Sigma_i \cap \mathcal{C}_{U,R}$ for large i , Σ_i locally smoothly converging to Σ implies that the level set $\{|x| = R\} \cap \Sigma_i \cap \overline{\mathcal{C}_{U,R}}$ is S^1 when i is sufficiently large. Then by standard Morse theory we know that $\Sigma_i \cap \mathcal{C}_{U,R}$ is diffeomorphic to $S^1 \times [0, +\infty)$ for sufficiently large i .

To complete the proof, it suffices to show that $|x|^2$ has no critical point in $\Sigma_i \cap \mathcal{C}_{U,R}$ for sufficiently large i . We prove it by contradiction. Assume on the contrary that $|x|^2$ has a critical point $p_i \in \Sigma_i \cap \mathcal{C}_{U,R}$ for large i . Since Σ_i locally smoothly converges to Σ , we must have $|p_i| \rightarrow \infty$.

We claim $p_i^\perp = p_i$. Suppose $\{e_1, e_2\}$ is an orthogonal frame around p_i on $\Sigma_i \cap \mathcal{C}_{U,R}$. Since p_i is a critical point of $|x|^2$,

$$\nabla_{e_j} \langle x, x \rangle(p_i) = 0, \quad j = 1, 2.$$

Note that

$$\nabla_{e_j} \langle x, x \rangle(p_i) = 2\langle e_i, p_i \rangle,$$

which implies that $p_i^\top = 0$, hence $p_i^\perp = p_i$.

Now set $y_i = p_i/|p_i|$ and $\rho_i = |p_i|$. $|p_i| \rightarrow \infty$ implies that $\rho_i \rightarrow \infty$. Let y be a limit point of y_i . Then $y \in \lim_{\rho \rightarrow \infty} \rho^{-1} \Sigma \cap U$ and $\lim_{\rho \rightarrow \infty} F_{\rho y, 1}(\Sigma) = 1$.

Following the blow-up process in Theorem 2.8, we define $\tilde{\Sigma}_i = \frac{|p_i|}{2}(\Sigma_i - p_i)$. Then $\tilde{\Sigma}_i$ locally smoothly converges to a translating soliton, denoted by $\tilde{\Sigma}$. Moreover, $H(p_i) = 1/2|p_i^\perp| = 1/2|p_i|$ implies that $|A|(p_i) \geq 1/2|p_i|$, and Theorem 2.8 implies that $\tilde{\Sigma}$ is a Bowl soliton.

On the other hand, by Proposition 2.7,

$$F_{x,t}(\tilde{\Sigma}) \leq \limsup_{\rho \rightarrow \infty} F_{\rho y, 1}(\Sigma) \leq 1.$$

Such an upper bound leads to a contradiction to $\tilde{\Sigma}$ being a Bowl soliton. Thus we complete the proof. $\square$

5. ASYMPTOTIC TO A CYLINDER

We now prove the second part of Theorem 3.1. In this section, U is always assumed to be a subset of S^2 such that $U \cap \lim_{\rho \rightarrow \infty} \rho^{-1} \Sigma$ has only one point y . Take R large enough such that $\Sigma \cap \mathcal{C}_{U,R}$ is diffeomorphic to $S^1 \times [0, +\infty)$. It follows from Lemma 4.1 that $\lim_{\rho \rightarrow \infty} \rho^{-1} \Sigma_i \cap U$ consists of disjoint simple closed curves and points for i sufficiently large.

Given a point $x \neq 0$ in $\mathbb{R}^3$, we denote by $P(x)$ the hyperplane containing x meanwhile being perpendicular to the vector x . We will use the following notation to denote the disk of plane $P(x)$:

$$B_r^x(y) := B_r(y) \cap P(x).$$

The purpose of this section is to prove that $\Sigma_i \cap \mathcal{C}_{U,R}$ is diffeomorphic to $S^1 \times [0, +\infty)$ or a disk. The proof is divided into three cases by the types of $\mathcal{C}_U \cap \Sigma_i$ (see 3.1 for notations): $\lim_{\rho \rightarrow \infty} \rho^{-1} \Sigma_i \cap U$ is either empty or one of the connected components of $\lim_{\rho \rightarrow \infty} \rho^{-1} \Sigma_i \cap U$ is a point, or a curve.

5.1. $\Sigma_i \cap \mathcal{C}_U$ contains a cylindrical end. Firstly, we consider that $\lim_{\rho \rightarrow \infty} \rho^{-1} \Sigma_i \cap U$ always contains a connected component which is a single point y_i . Then by Lemma 4.1, $y_i \rightarrow y$ as $i \rightarrow \infty$. The goal of this section is to prove the following:

Proposition 5.1. *For sufficiently large i , $\Sigma_i \cap \mathcal{C}_{U,R}$ is diffeomorphic to $\Sigma \cap \mathcal{C}_{U,R}$.*

Proof. Let us first sketch the idea. We want to show that $\Sigma_i \cap \mathcal{C}_{U,R}$ is topologically a cylinder by contradiction. Suppose $\Sigma_i \cap \mathcal{C}_{U,R}$ is not topologically a cylinder near the point p_i , and the projection of p_i to y_i is $\rho_i y_i$. Then we can translate Σ_i by the vector $-\rho_i y_i$, in other words we consider the sequence $\Sigma_i - \rho_i y_i$. We will show that $\Sigma_i - \rho_i y_i$ converges to

a cylinder smoothly. As a result, Σ_i should be topologically a cylinder near p_i , which is a contradiction.

The proof is divided into four steps.

Step 1: For any sequence $\rho_i \rightarrow \infty$, $F_{\rho_i y_i, 1}(\Sigma_i) \rightarrow \lambda_1$.

Proof of Step 1. On one hand,

$$\liminf_{i \rightarrow \infty} F_{\rho_i y_i, 1}(\Sigma_i) \geq \liminf_{i \rightarrow \infty} \liminf_{\rho \rightarrow \infty} F_{\rho y_i, 1}(\Sigma_i) = \lambda_1.$$

On the other hand, by Proposition 2.7 (taking $a_i \equiv 1$, $x = 0$, $t = 1$),

$$\limsup_{i \rightarrow \infty} F_{\rho_i y_i, 1}(\Sigma_i) \leq \limsup_{\rho \rightarrow \infty} F_{\rho y, 1}(\Sigma) = \lambda_1.$$

Then the desired result follows. $\square$

Step 2: For any sequence $\rho_i \rightarrow \infty$, $\Sigma_i - \rho_i y_i$ locally smoothly converges to $S^1(\sqrt{2}) \times \mathbb{R}_y$ up to a subsequence.

Proof of Step 2. Define the Brakke flow for each i and $t < 0$,

$$\mu_t^i = \mathcal{H}^2 \llcorner (\sqrt{-t} \Sigma_i - \rho_i y_i).$$

By the compactness theorem of Brakke flow, there exists a Brakke flow $\{\nu_t\}_{t < 0}$ such that for each $t < 0$, $\mu_t^i \rightarrow \nu_t$ as varifolds. In the following, we show that the support of ν_{-1} is a self-shrinker.

By the monotonicity of F -functional, $\lambda(\nu_t) \leq \lambda_1$. Meanwhile, for any $t < 0$,

$$\begin{aligned} \int \Phi(x, t) d\nu_t &= \lim_{i \rightarrow \infty} \int \Phi(x, t) d\mu_t^i \\ &= \lim_{i \rightarrow \infty} \int_{\sqrt{-t} \Sigma_i - \rho_i y_i} \Phi(x, t) dx \\ &= \lim_{i \rightarrow \infty} \int_{\Sigma_i - \frac{\rho_i}{\sqrt{-t}} y_i} \Phi(x, -1) dx \\ &= \lim_{i \rightarrow \infty} F_{\frac{\rho_i}{\sqrt{-t}} y_i, 1}(\Sigma_i) \\ &= \lambda_1. \end{aligned}$$

By the monotonicity formula for Brakke flows, ν_{-1} is F -stationary, i.e.

$$\vec{H} + \frac{S(x)^\perp \cdot x}{2} = 0, \quad \nu_{-1} - a.e. \ x.$$

Here $S(x)^\perp$ is the projection operator, sending x to the normal direction of x . In other words, ν_{-1} is a self-shrinker in the sense of varifolds. Now we show that ν_{-1} is supporting on a smooth self-shrinker.

For $\tau \in \mathbb{R}$, $\rho_i + \tau \rightarrow +\infty$ as $\rho_i \rightarrow +\infty$. Then $\nu_{-1} - \tau y$, which is the limit of $\mu_{-1}^i - \tau y = \mathcal{H}^2 \llcorner (\Sigma_i - (\rho_i + \tau) y_i)$, is also F -stationary. Thus $S(x)^\perp \cdot y = 0$, which implies that ν_{-1} splits off a line (c.f. [43] Lemma 1.10). Recall that $F_{0,1}(\nu_{-1}) = \lambda_1$. We conclude that $\nu_{-1} = \mathcal{H}^2 \llcorner S^1(\sqrt{2}) \times \mathbb{R}_y$.

By the local regularity of Brakke flows [41] (see Theorem 2.6), $\Sigma_i - \rho_i y_i$ locally smoothly converges to $S^1(\sqrt{2}) \times \mathbb{R}_y$. $\square$

Step 3: For sufficiently large i , $\Sigma_i \cap \mathcal{C}_{U,R} \subseteq N_i$, where $N_i = \bigcup_{\rho=R}^{+\infty} B_2^{\rho y_i}(\rho y_i)$.

Here N_i is a part of the solid cylinder with direction y_i and radius 2.

Proof of Step 3. We prove by contradiction. Assume on the contrary that there is a subsequence of $\{\Sigma_i\}$ (still denoted by $\{\Sigma_i\}$) such that $\Sigma_i \cap \mathcal{C}_{U,R}$ does not completely lie in N_i . Then for each i , $\Sigma_i \cap \bigcup_{\rho=R}^{+\infty} \partial B_2^{\rho y_i}(\rho y_i) \neq \emptyset$. Take p_i in this set and let $\rho_i = \langle p_i, y_i \rangle$ (see Figure I).

FIGURE I. Asymptotic to a cylinder.

If ρ_i is bounded, take a subsequence of p_i (still denoted by p_i) such that $\lim \rho_i = \rho_\infty$. Then $\Sigma_i - \rho_i y_i$ locally smoothly converges to $\Sigma - \rho_\infty y$, which is also close to $S^1(\sqrt{2}) \times \mathbb{R}_y$. Hence for i sufficiently large, $\Sigma_i \cap \bigcup_{\rho=\rho_i-1}^{\rho_i+1} \partial B_2^{\rho y_i}(\rho y_i) = \emptyset$. This is a contradiction.

If ρ_i is unbounded, take a subsequence of p_i (still denoted by p_i) such that $\lim \rho_i = +\infty$. Step 2 gives that $\Sigma_i - \rho_i y_i$ locally smoothly converges to $S^1(\sqrt{2}) \times \mathbb{R}_y$ up to a subsequence (still denoted by $\{\Sigma_i\}$). However, $p_i \in \partial B_2^{\rho_i y_i}(\rho_i y_i)$ implies that $\Sigma_i - \rho_i y_i$ can not converge to $S^1(\sqrt{2}) \times \mathbb{R}_y$ locally smoothly. This is a contradiction.

Therefore, $\Sigma_i \cap \mathcal{C}_{U,R} \subseteq N_i$ for large i . □

Now we can prove our main theorem in this subsection by contradiction. Suppose on the contrary that there is a subsequence of $\Sigma_i \cap \mathcal{C}_{U,R}$ which are not diffeomorphic to $\Sigma \cap \mathcal{C}_{U,R}$ (still denoted by Σ_i). Then we can take $r_i \rightarrow \infty$ so that Σ_i is not diffeomorphic to $S^1 \times [0, 1]$ near $r_i y_i$. Applying the result in Step 2 gives a contradiction. We present the details below.

Step 4: $\Sigma_i \cap \mathcal{C}_{U,R}$ is diffeomorphic to $\Sigma \cap \mathcal{C}_{U,R}$.

Proof of Step 4. By Step 3, it suffices to show that there exists $I > 0$ such that for any $i > I$ and $r > R$, $\Sigma_i \cap \bigcup_{\rho=r}^{r+1} B_2^{\rho y_i}(\rho y_i)$ is diffeomorphic to $S^1 \times [0, 1]$.

Assume the statement is not true. Then for each i , take $r_i > R$ such that $\Sigma_i \cap \bigcup_{\rho=r_i}^{r_i+1} B_2^{\rho y_i}(\rho y_i)$ is not diffeomorphic to $S^1 \times [0, 1]$.

If r_i is bounded, take a subsequence of r_i (still denoted by r_i) such that $\lim r_i = r$. Then $\Sigma_i - r_i y_i$ locally smoothly converges to $\Sigma - r y$, which is a contradiction.

If r_i is unbounded, take a subsequence of r_i (still denoted by r_i) such that $\lim r_i = +\infty$. Because the topological type is different, $\Sigma_i - r_i y_i$ can not locally smoothly converges to $S^1(\sqrt{2}) \times \mathbb{R}_y$, which contradicts the conclusion in Step 2. □

Combining all the steps above shows the proposition. □

5.2. $\Sigma_i \cap \mathcal{C}_U$ contains a conical end. Now we consider that $\lim_{\rho \rightarrow \infty} \rho^{-1} \Sigma_i \cap U$ contains a connected component which is a simple closed curve.

Let γ_i be a connected component of $\lim_{\rho \rightarrow \infty} \rho^{-1} \Sigma_i \cap U$. By Lemma 4.1, $\gamma_i \rightarrow y$ in Hausdorff distance. Denote by $\tilde{\gamma}_i$ the curve $\mathcal{C}_{\gamma_i} \cap B_1^y(y)$. In the following of this section, $\tilde{\gamma}_i$ is also regarded as a curve in $\mathbb{R}^2 (= P(y))$.

Before we state our main proposition, we need some Lemmas. We want to show that $\{\tilde{\gamma}_i\}$ is a entropy minimizing sequence among all close plane curves.

Lemma 5.2.

$$\lim_{i \rightarrow \infty} \lambda(\tilde{\gamma}_i) = \lim_{i \rightarrow \infty} \lambda(\mathcal{C}_{\tilde{\gamma}_i}) = \lambda_1.$$

Proof. On one hand, take $s_i > 0$ and $y_i \in S^2(1)$ such that

$$\lambda(\mathcal{C}_{\tilde{\gamma}_i}) = F_{s_i y_i, 1}(\mathcal{C}_{\tilde{\gamma}_i}).$$

γ_i converges to a point implies that $s_i \rightarrow \infty$. Then Proposition 2.7 and the monotonicity formula imply that for an increasing to infinity sequence ρ_i ,

$$\begin{aligned} \lambda_1 &= \lim_{\rho \rightarrow \infty} F_{\rho y, 1}(\Sigma) \geq \lim_{i \rightarrow \infty} F_{\rho_i y_i, 1 + s_i^{-2} \rho_i^2}(\Sigma_i) \\ &\geq \lim_{i \rightarrow \infty} \lim_{\rho \rightarrow \infty} F_{\rho y_i, 1 + s_i^{-2} \rho^2}(\Sigma_i) \\ &= \lim_{i \rightarrow \infty} F_{0, 1}(\mathcal{C}_{\tilde{\gamma}_i} - s_i y_i) = \lim_{i \rightarrow \infty} \lambda(\mathcal{C}_{\tilde{\gamma}_i}). \end{aligned}$$

Therefore,

$$(5.1) \quad \lim_{i \rightarrow \infty} \lambda(\mathcal{C}_{\tilde{\gamma}_i}) \leq \lambda_1 \leq \lim_{i \rightarrow \infty} \lambda(\tilde{\gamma}_i).$$

On the other hand, take $x_i \in B_1^{y_i}(y_i)$ and $a_i \in (0, \infty)$ such that

$$\lambda(\tilde{\gamma}_i) = F_{0, 1}(a_i^{-1}(\tilde{\gamma}_i - x_i)).$$

Then a_i must converge to 0 since the diameter of $\tilde{\gamma}_i$ converges to 0. Then in any compact domain, the varifold distance between $a_i^{-1}(\mathcal{C}_{\tilde{\gamma}_i} - x_i)$ and $a_i^{-1}(\tilde{\gamma}_i - x_i) \times \mathbb{R}$ goes to zero. Since the entropy of Σ_i are uniformly bounded, so does the entropy of $\mathcal{C}_{\tilde{\gamma}_i}$ and $\tilde{\gamma}_i$. Therefore we conclude that

$$|F_{0, 1}(a_i^{-1}(\mathcal{C}_{\tilde{\gamma}_i} - x_i)) - \lambda(\tilde{\gamma}_i)| = |F_{0, 1}(a_i^{-1}(\mathcal{C}_{\tilde{\gamma}_i} - x_i)) - F_{0, 1}(a_i^{-1}(\tilde{\gamma}_i - x_i) \times \mathbb{R})| \rightarrow 0.$$

This deduces

$$\lim_{i \rightarrow \infty} \lambda(\mathcal{C}_{\tilde{\gamma}_i}) \geq \lim_{i \rightarrow \infty} \lambda(\tilde{\gamma}_i).$$

Together with (5.1), we get the desired equality. $\square$

With the entropy minimizing property, we prove that after normalizing, these curves converges to a round circle in the sense of varifolds. The argument here is the one-dimensional case of [6].

Lemma 5.3. *There exist $x_i \in \mathbb{R}^2$ and $T_i > 0$ so that $T_i^{-1/2}(\tilde{\gamma}_i - x_i)$ converges to $S^1(\sqrt{2})$ in the sense of varifolds.*

Proof. Let $\{\gamma_i(t)\}_{t \in [0, T_i]}$ be the maximal smooth curve shortening flow with $\gamma_i(0) = \gamma_i$. By the entropy bound and [22, 23], the first singularity of this flow is at (x_i, T_i) , where the flow disappears in a round point. Now let $\Gamma_i(t) = T_i^{-1/2}(\tilde{\gamma}_i((t+1)T_i) - x_i)$ for $t \in [-1, 0)$.

We can follow the argument in [6] by replacing [6, Proposition 3.2] with Proposition 5.4 to prove that $\Gamma_i(t)$ actually have uniformly curvature bound for each fixed $t \in (-1, 0)$.

By Brakke's compactness theorem, up to a subsequence $\{\Gamma_i(t)\}_{t \in [-1, 0)}$ converges to a Brakke flow $\{\nu_t\}_{t \in [-1, 0)}$ in the sense of varifolds. By the uniform curvature bounds for each fixed $t \in (-1, 0)$, ν_t is a smooth curve. Clearly, $\lambda(\nu_t) = \lambda_1$ and, by the upper semi-continuous of Gaussian density, this flow becomes singular at $(0, 0)$. Hence $\nu_{-1/2}$ must be a smooth circle with radius 1, which implies that ν_{-1} is a round circle with radius $\sqrt{2}$. $\square$

Proposition 5.4. *Suppose $\{\Gamma(t)\}_{t \in \mathbb{R}}$ is an eternal curve shortening flow in the plane $\mathbb{R}^2$ with $\sup_{t \in \mathbb{R}} \lambda(\Gamma(t)) < 2$. Then $\Gamma(t)$ is a straight line for any $t \in \mathbb{R}$.*

Proof. We follow the proof of [6, Proposition 3.2]. Again we consider the blow-down sequence of curve shortening flow $\{\rho^{-1}\Gamma(\rho^2 t)\}_{t \in \mathbb{R}}$. Then a subsequence converges to a self-shrinking Brakke flow $\{\nu_t\}_{t \in \mathbb{R}}$, with entropy less than 2. Moreover, the entropy bound and [42] suggest that the ν_{-1} must be a 1-dimensional F -stationary varifold without triple junction. Thus ν_{-1} can only be either a circle or a straight line. If it is a circle, then the smooth convergence given by Brakke's regularity theorem, $\rho^{-1}\Gamma(-\rho^2)$ is contained in $S^1(2)$ for large ρ . However, this contradicts the hypothesis that $\{\Gamma(t)\}$ is a smooth flow for all $t \in \mathbb{R}$ and then we must have that ν_{-1} is a straight line.

As such, by the monotonicity formula, $\lambda(\Gamma(t)) = 1$. Then we conclude the proposition. $\square$

Let x_i and T_i be the point and real number in Lemma 5.3. Recall that x_i is a point in $\mathbb{R}^2 = P(y) \subset \mathbb{R}^3$. Let $y_i = x_i/|x_i|$ and $\beta_i = T_i^{1/2}/|x_i|$. Note that $T_i^{-1/2}(\tilde{\gamma}_i - x_i) \times \mathbb{R}$ converges to $S^1(\sqrt{2}) \times \mathbb{R}$ in the sense of varifolds. Recall that in any fixed compact domain, the varifold distance between $T_i^{-1/2}(\mathcal{C}_{\tilde{\gamma}_i} - x_i)$ and $T_i^{-1/2}(\tilde{\gamma}_i - x_i) \times \mathbb{R}$ goes to zero. Then we conclude that $\mathcal{C}_{\gamma_i} - \beta_i^{-1}y_i$ converges to $S^1(\sqrt{2}) \times \mathbb{R}$ in the sense of varifolds.

Proposition 5.5. $\Sigma_i \cap \mathcal{C}_{U,R}$ is diffeomorphic to $\Sigma \cap \mathcal{C}_{U,R}$.

Proof. We use the similar argument as in the proof of Proposition 5.1. However, there is a difference. For the cylindrical ends case, we only need translations to bring the point where the topology of the end is bad, to a neighborhood of the origin to argue by a contradiction; for the conical ends case, we need not only translations but also dilations to bring the point, where the topology of the end is bad, back to a neighborhood of the origin and then argue by contradiction. The proof is also divided into four steps:

Step 1: For any sequence $\rho_i \rightarrow \infty$, we have

$$F_{\rho_i y_i, 1 + \beta_i^2 \rho_i^2}(\Sigma_i) \rightarrow \lambda_1.$$

Proof of Step 1. By the monotonicity of F -functional,

$$\begin{aligned} \liminf_{i \rightarrow \infty} F_{\rho_i y_i, 1 + \beta_i^2 \rho_i^2}(\Sigma_i) &\geq \liminf_{i \rightarrow \infty} \liminf_{\rho \rightarrow \infty} F_{\rho y_i, 1 + \beta_i^2 \rho^2}(\Sigma_i) \\ &= \liminf_{i \rightarrow \infty} \liminf_{\rho \rightarrow \infty} F_{0,1}((\rho \beta_i)^{-1} \Sigma_i - \beta_i^{-1} y_i) \\ &= \liminf_{i \rightarrow \infty} F_{0,1}(\mathcal{C}_{\gamma_i} - \beta_i^{-1} y_i) = \lambda_1. \end{aligned}$$

By Proposition 2.7,

$$\limsup_{i \rightarrow \infty} F_{\rho_i y_i, 1 + \beta_i^2 \rho_i^2}(\Sigma_i) \leq \liminf_{\rho \rightarrow \infty} F_{\rho y, 1}(\Sigma) = \lambda_1.$$

Then the desired result follows immediately. $\square$

Step 2: For any $\rho_i \rightarrow \infty$ with $\lim_{i \rightarrow \infty} \rho_i \beta_i < +\infty$, $\frac{\Sigma_i - \rho_i y_i}{\sqrt{1 + \beta_i^2 \rho_i^2}}$ locally smoothly converges to $S^1(\sqrt{2}) \times \mathbb{R}_y$ up to a subsequence.

Remark 5.6. Intuitively, such a sequence is “bounded” by $\mathcal{C}_{\gamma_i} - \beta_i^{-1} y_i$ and $\Sigma - \rho y$. However, both sequences locally smoothly converge to the round cylinder $S^1(\sqrt{2}) \times \mathbb{R}_y$.

Proof of Step 2. For any sequence of $\rho_i \rightarrow \infty$ with $\lim_{i \rightarrow \infty} \rho_i \beta_i < +\infty$, there exists a constant $K > 0$ so that $\rho_i \beta_i < K$ for all large i . Define the Brakke flow for $t \leq -K^2/(1+K^2)$ by

$$\mu_t^i = \mathcal{H}^2 \llbracket \left(\sqrt{\frac{1}{1 + \beta_i^2 \rho_i^2} - 1 - t} \right) \Sigma_i - \frac{\rho_i}{\sqrt{1 + \beta_i^2 \rho_i^2}} y_i \rrbracket.$$

This is the mean curvature flow so that $\mu_{-1}^i = \frac{\Sigma_i - \rho_i y_i}{\sqrt{1 + \beta_i^2 \rho_i^2}}$. Then by the compactness of Brakke flows (see Lemma 2.5), μ_t^i converges to a Brakke flow ν_t in the sense of Radon measure. We want to show that the support of ν_{-1} is a self-shrinker.

By the monotonicity of F -functional, $\lambda(\nu_t) \leq \lambda_1$. By computation,

$$(5.2) \quad \mu_t^i = \mathcal{H}^2 \llbracket \left(\sqrt{-t} \left(\frac{\Sigma_i - \tilde{\rho}_i y_i}{\sqrt{1 + \beta_i^2 \tilde{\rho}_i^2}} \right) \right) \rrbracket,$$

where $\tilde{\rho}_i = \rho_i / \sqrt{1 - (1+t)(1 + \beta_i^2 \rho_i^2)}$. Note that for any $t < -K^2/(1 + K^2)$, we have $\tilde{\rho}_i \rightarrow \infty$ since $\rho_i \rightarrow \infty$ and $\beta_i \rightarrow 0$. Therefore,

$$\begin{aligned} \int \Phi(x, t) d\nu_t &= \lim_{i \rightarrow \infty} \int \Phi(x, t) d\mu_t^i \\ &= \lim_{i \rightarrow \infty} F_{\tilde{\rho}_i y_i, 1 + \beta_i^2 \tilde{\rho}_i^2}(\Sigma_i) \\ &= \lambda_1. \end{aligned}$$

By the monotonicity formula, ν_{-1} is F -stationary, i.e.

$$\vec{H} + \frac{S(x)^\perp \cdot x}{2} = 0, \quad \nu_{-1} - a.e. \ x.$$

We now show that ν_{-1} is supporting on a smooth self-shrinker. For any $\tau \in \mathbb{R}$, set

$$\rho'_i = \rho_i + \tau \sqrt{1 + \beta_i^2 \rho_i^2} \quad \text{and} \quad s_i = \frac{1 + \beta_i^2 \rho_i'^2}{1 + \beta_i^2 \rho_i^2}.$$

It follows that $\rho'_i \rightarrow \infty$, $\lim_{i \rightarrow \infty} \beta_i \rho'_i < +\infty$ and $s_i \rightarrow 1$ as $i \rightarrow \infty$. Then $\nu_{-1} - \tau y$, the limit of $\mu_{-1}^i - \tau y_i = \mathcal{H}^2 \llbracket \left(\sqrt{s_i} \left(\frac{\Sigma_i - \rho'_i y_i}{\sqrt{1 + \beta_i^2 \rho_i'^2}} \right) \right) \rrbracket$, is also F -stationary. This shows that $S(x)^\perp \cdot y = 0$ for almost x . Therefore, ν_{-1} splits off a line with direction y . Note that $\lambda(\nu_{-1}) = \lambda_1$. Hence ν_{-1} supports on $S^1(\sqrt{2}) \times \mathbb{R}_y$ with multiplicity 1. Then by local regularity for Brakke flows

(Theorem 2.6), we conclude that $\frac{\Sigma_i - \rho_i y_i}{\sqrt{1 + \beta_i^2 \rho_i^2}}$ locally smoothly converges to $S^1(\sqrt{2}) \times \mathbb{R}_y$ up to a subsequence. $\square$

Step 3: For sufficiently large i , $\Sigma_i \cap \mathcal{C}_{U,R} \subseteq N_i$, where $N_i = \bigcup_{\rho=0}^{+\infty} B_{2\sqrt{1+\beta_i^2 \rho^2}}^{\rho y_i}(\rho y_i)$.

Proof of Step 3. Assume on the contrary that a subsequence of Σ_i is not contained in N_i . We still denote the subsequence by $\{\Sigma_i\}$. Then for each i , $\Sigma_i \cap \mathcal{C}_{U,R} \cap \partial N_i$ is non-empty. Take p_i in this intersection and set $\rho_i = \langle p_i, y_i \rangle$.

If ρ_i is bounded, take a subsequence of p_i (still denoted by p_i) such that $\lim \rho_i = \rho$. Then

$$\lim_{i \rightarrow \infty} \left| \frac{p_i - \rho_i y_i}{\sqrt{1 + \beta_i^2 \rho_i^2}} \right| = \lim_{i \rightarrow \infty} |p_i - \rho_i y_i| = 2.$$

Recall that $\Sigma_i - \rho_i y_i$ locally smoothly converges to a smooth surface. Therefore the limit surface must intersect $\mathbb{R}_y \times S^1(2)$. This contradicts the fact that $\Sigma_i - \rho_i y_i$ locally smoothly converges to $\Sigma - \rho y$.

If ρ_i is unbounded and $\rho_i \beta_i$ is bounded, take a subsequence of p_i (still denoted by p_i) such that $\lim_{i \rightarrow \infty} \rho_i = +\infty$ and $\lim_{i \rightarrow \infty} \beta_i \rho_i < +\infty$. Then $p_i \in \Sigma_i \cap \partial N_i$ also implies that

$$\lim_{i \rightarrow \infty} \left| \frac{p_i - \rho_i y_i}{\sqrt{1 + \beta_i^2 \rho_i^2}} \right| = 2,$$

which contradicts that $\frac{\Sigma_i - \rho_i y_i}{\sqrt{1 + \beta_i^2 \rho_i^2}}$ locally smoothly converges to $S^1(\sqrt{2}) \times \mathbb{R}_y$ by Step 2.

It remains to consider $\rho_i \beta_i \rightarrow \infty$ (see Figure II). Denote by $z_i = p_i / \rho_i$. Then for any $k > 0$,

$$\begin{aligned} \lim_{i \rightarrow \infty} \frac{\text{dist}(k\beta_i^{-1}z_i, \mathbb{R}_{y_i})}{\sqrt{1+k^2}} &= \lim_{i \rightarrow \infty} \frac{\text{dist}(\rho_i z_i, \mathbb{R}_{y_i})}{\sqrt{1+k^2}} \cdot \frac{k\beta_i^{-1}}{\rho_i} \\ &= \lim_{i \rightarrow \infty} \frac{\text{dist}(p_i, \mathbb{R}_{y_i})}{\sqrt{1+\rho_i^2 \beta_i^2}} \cdot \frac{\sqrt{1+\rho_i^2 \beta_i^2}}{\sqrt{1+k^2}} \cdot \frac{k}{\rho_i \beta_i} \\ &= \frac{2k}{\sqrt{1+k^2}}. \end{aligned}$$

Recall that $\frac{\Sigma_i - k\beta_i^{-1}z_i}{\sqrt{1+k^2}}$ locally smoothly converges to $S^1(\sqrt{2}) \times \mathbb{R}_y$. Thus we must have that for large i ,

$$(5.3) \quad B_{k/4}(k\beta_i^{-1}z_i) \cap \Sigma_i = \emptyset.$$

Take K large enough and fixed, then the uniform entropy bound and (5.3) give that

$$(5.4) \quad F_{K\beta_i^{-1}z_i, 1}(\Sigma_i) < 1/2.$$

Now we derive a entropy lower bound to get a contradiction. Because $\rho_i z_i \in \Sigma_i$, we always have

$$\lim_{t \rightarrow 0} F_{\rho_i z_i, t}(\Sigma_i) = 1.$$

FIGURE II. Small cones.

Then by the monotonicity formula for F -functional,

$$\begin{aligned} \lim_{t \rightarrow 0} F_{\rho_i z_i, t}(\Sigma_i) &= \lim_{t \rightarrow 0} F_{\rho_i z_i, 1 - \frac{1-t}{\rho_i^2} \cdot \rho_i^2}(\Sigma_i) \\ &\leq \limsup_{i \rightarrow \infty} \lim_{t \rightarrow 0} F_{K\beta_i^{-1} z_i, 1 - \frac{1-t}{\rho_i^2} \cdot (K\beta_i^{-1})^2}(\Sigma_i) \\ &\leq \limsup_{i \rightarrow \infty} F_{K\beta_i^{-1} z_i, 1}(\Sigma_i), \end{aligned}$$

which contradicts (5.4).

Above all, $\Sigma_i \cap \mathcal{C}_{U,R} \subseteq N_i$ for large i . □

We now prove Proposition 5.5 by contradiction. Suppose on the contrary that $\Sigma_i \cap \mathcal{C}_{U,R}$ is not diffeomorphic to $\Sigma \cap \mathcal{C}_{U,R}$ for a subsequence of Σ_i (still denoted by Σ_i). Then take r_i so that Σ_i is not diffeomorphic to Σ in a suitable neighborhood of $r_i y_i$. If r_i is uniformly bounded, then it contradicts the fact that Σ_i locally smoothly converges to Σ . If r_i is unbounded, then by taking the limit, we see a contradiction to the conclusion in Step 2.

Step 4: $\Sigma_i \cap \mathcal{C}_{U,R}$ is diffeomorphic to $\Sigma \cap \mathcal{C}_{U,R}$.

Proof of Step 4. It suffices to show that there exists $I > 0$ such that for any $i > I$ and $r > R$, $\Sigma_i \cap \bigcup_{\rho=r}^{r+1} B_{2\sqrt{1+\beta_i^2 \rho^2}}^{\rho y_i}(\rho y_i)$ is diffeomorphic to $S^1 \times [0, 1]$. Define

$$\Sigma_i(r_i; y_i) = \Sigma_i \cap \bigcup_{\rho=r}^{r+1} B_{2\sqrt{1+\beta_i^2 \rho^2}}^{\rho y_i}(\rho y_i).$$

Assume the statement is not true. Then for each i , take $r_i > R$ such that $\Sigma_i(r_i; y_i)$ is not diffeomorphic to $S^1 \times [0, 1]$.

If r_i is bounded, take a subsequence of r_i (still denoted by r_i) such that $\lim r_i = r$. Note that $\Sigma_i(r_i; y_i) - r_i y_i$ smoothly converges to $(\Sigma - ry) \cap B_2(0) \times [0, 1]$, which leads to a contradiction.

If r_i is unbounded and $r_i \beta_i$ is bounded, take a subsequence of r_i (still denoted by r_i) such that $\lim r_i = +\infty$ and $\lim r_i \beta_i < +\infty$. Then we have proved in Step 3 that $\frac{\Sigma_i - r_i y_i}{\sqrt{1+\beta_i^2 r_i^2}}$ smoothly converges to $S^1(\sqrt{2}) \times \mathbb{R}$. It follows that $\frac{\Sigma_i(r_i; y_i) - r_i y_i}{\sqrt{1+\beta_i^2 r_i^2}}$ smoothly converges to $S^1(\sqrt{2}) \times [0, 1]$, which leads to a contradiction.

It remains to consider $r_i \beta_i \rightarrow \infty$. Then the function $|x|^2$ must have critical points on $\Sigma_i(r_i; y_i)$. Without loss of generality, denote by $\rho_i z_i \in \Sigma_i(r_i; y_i)$ ($|z_i| = 1$) the critical point

of $|x|^2$. It follows that $\rho_i \beta_i \rightarrow \infty$. By Theorem 2.8 and 4.2, $\frac{\rho_i}{2}(\Sigma_i - \rho_i z_i)$ locally smoothly converges to a Bowl soliton. Thus there exists $(x, t) \in \mathbb{R}^3 \times (0, \infty)$ (see [24]) so that

$$(5.5) \quad \liminf_{i \rightarrow \infty} F_{x,t}(\rho_i(\Sigma_i - \rho_i z_i)) > \frac{1 + \lambda_1}{2}.$$

However, the monotonicity formula of F -functional gives that

$$\begin{aligned} \lim_{i \rightarrow \infty} F_{x,t}(\rho_i(\Sigma_i - \rho_i z_i)) &= \lim_{i \rightarrow \infty} F_{\rho_i(z_i + x/\rho_i^2), 1 + (t/\rho_i^4 - 1/\rho_i^2) \cdot \rho_i^2}(\Sigma_i) \\ &\leq \limsup_{k \rightarrow \infty} \limsup_{i \rightarrow \infty} F_{k\beta_i^{-1}(z_i + x/\rho_i^2), 1 + (t/\rho_i^4 - 1/\rho_i^2) \cdot (k\beta_i^{-1})^2}(\Sigma_i) \\ &= \limsup_{k \rightarrow \infty} \limsup_{i \rightarrow \infty} F_{k\beta_i^{-1}z_i, 1}(\Sigma_i) \\ &= \limsup_{k \rightarrow \infty} \limsup_{i \rightarrow \infty} F_{k\beta_i^{-1}(z_i - y_i), 1}(\Sigma_i - k\beta_i^{-1}y_i). \end{aligned}$$

A direct computation gives that, for large i ,

$$\begin{aligned} k\beta_i^{-1} \cdot |z_i - y_i| &= k\beta_i^{-1} \cdot |\rho_i z_i - \rho_i y_i|/\rho_i \leq k\beta_i^{-1} \cdot 2\text{dist}(p_i, \mathbb{R}_{y_i})/\rho_i \\ &\leq k\beta_i^{-1} \cdot 4\sqrt{1 + \beta_i^2 \rho_i^2/\rho_i} \leq 8k, \end{aligned}$$

where we used the fact that $z_i \in \Sigma_i(r_i; y_i)$. Hence we have that

$$\begin{aligned} \lim_{i \rightarrow \infty} F_{x,t}(\rho_i(\Sigma_i - \rho_i z_i)) &\leq \limsup_{k \rightarrow \infty} \limsup_{i \rightarrow \infty} F_{k\beta_i^{-1}(z_i - y_i), 1}(\Sigma_i - k\beta_i^{-1}y_i) \\ &\leq \limsup_{k \rightarrow \infty} \limsup_{i \rightarrow \infty} \sup_{z \in B_{8k}(0)} F_{z,1}(\Sigma_i - k\beta_i^{-1}y_i) \\ &\leq \limsup_{k \rightarrow \infty} \sup_{z \in B_{8k}(0)} F_{z,1}(S^1(\sqrt{2(1+k^2)} \times \mathbb{R}) = 1, \end{aligned}$$

where we used that $\frac{\Sigma_i - k\beta_i^{-1}y_i}{\sqrt{1+k^2}}$ locally smoothly converges to $S^1(\sqrt{2}) \times \mathbb{R}_y$. This entropy upper bound contradicts (5.5). Then the proof of Step 4 is finished. $\square$

We have completed the proof of the proposition. $\square$

5.3. $\Sigma_i \cap \mathcal{C}_U$ is compact. In this part, we consider that $\lim_{\rho \rightarrow \infty} \rho^{-1}\Sigma_i \cap U = \emptyset$. Then $\Sigma_i \cap \mathcal{C}_U$ is compact for all i . Take $q_i \in \Sigma_i \cap \mathcal{C}_U$ such that

$$|q_i| = \sup_{x \in \Sigma_i \cap \mathcal{C}_U} |x|.$$

Set $y_i = q_i/|q_i|$ and $d_i = |q_i|$. Then $d_i \rightarrow +\infty$ and $y_i \rightarrow y$.

Proposition 5.7. $\Sigma_i \cap \mathcal{C}_{U,R}$ is diffeomorphic to a disk.

Proof. The proof is divided into four steps:

Step 1: For any $\rho_i \rightarrow +\infty$ with $d_i(d_i - \rho_i) \rightarrow +\infty$, $F_{\rho_i y_i, 1 - \rho_i^2/d_i^2}(\Sigma_i) \rightarrow \lambda_1$.

Proof of Step 1. By Theorem 2.8, $\frac{d_i}{2}(\Sigma_i - d_i y_i)$ locally smoothly converges to a translating soliton Σ_y with translating direction $-y$. According to the entropy bound, it is a Bowl

soliton by Theorem B.1. Hence $\frac{\Sigma_y + \rho y}{\sqrt{\rho}}$ locally smoothly converges to $S^1(\sqrt{2}) \times \mathbb{R}_y$ as $\rho \rightarrow +\infty$. By the monotonicity of F -functional,

$$\begin{aligned} \liminf_{i \rightarrow \infty} F_{\rho_i y_i, 1 - \frac{\rho_i^2}{d_i^2}}(\Sigma_i) &\geq \liminf_{\rho \rightarrow \infty} \liminf_{i \rightarrow \infty} F_{(d_i - \frac{\rho}{d_i})y_i, 1 - \frac{1}{d_i^2}(d_i - \frac{\rho}{d_i})^2}(\Sigma_i) \\ &= \liminf_{\rho \rightarrow \infty} \liminf_{i \rightarrow \infty} F_{0,1}\left(\frac{d_i(\Sigma_i - d_i y_i) + \rho y_i}{\sqrt{2\rho - \rho^2/d_i^2}}\right) \\ &= \liminf_{\rho \rightarrow \infty} F_{-\frac{\rho y}{2}, \frac{\rho}{2}}(\Sigma_y) \\ &= \lambda_1. \end{aligned}$$

Here the first inequality follows from the fact that for a fixed ρ , when d_i is sufficiently large, $(d_i - \frac{\rho}{d_i}) \geq \rho_i$. By Proposition 2.7

$$\limsup_{i \rightarrow \infty} F_{\rho_i y_i, 1 - \frac{\rho_i^2}{d_i^2}}(\Sigma_i) \leq \liminf_{\rho \rightarrow \infty} F_{\rho y, 1}(\Sigma) = \lambda_1.$$

This completes the proof. $\square$

Step 2: For any $\rho_i \rightarrow +\infty$ with $d_i(d_i - \rho_i) \rightarrow +\infty$, $\frac{\Sigma_i - \rho_i y_i}{\sqrt{1 - \rho_i^2/d_i^2}}$ locally smoothly converges to $S^1(\sqrt{2}) \times \mathbb{R}_y$ up to a subsequence.

For $\rho_i \rightarrow +\infty$ with $d_i(d_i - \rho_i) \rightarrow \rho$, $\frac{\Sigma_i - \rho_i y_i}{\sqrt{1 - \rho_i^2/d_i^2}}$ locally smoothly converges to $\frac{2\Sigma_y + \rho y}{\sqrt{2\rho}}$.

Remark 5.8. Note that Theorem 2.8 gives that $\frac{d_i}{2}(\Sigma_i - d_i y_i)$ locally smoothly converges to a translating soliton Σ_y with translating direction $-y$. On the one hand, the blow-down sequence of the end of a Bowl soliton is a round cylinder. On the other hand, $\Sigma - \rho y$ also converges to a round cylinder. In the first case of Step 2, the sequence must converge to a round cylinder.

Proof of Step 2. Define the Brakke flow for $t < 0$,

$$\mu_t^i = \mathcal{H}^2 \llbracket \left(\sqrt{\frac{1}{1 - \rho_i^2/d_i^2}} - 1 - t \Sigma_i - \frac{\rho_i y_i}{\sqrt{1 - \rho_i^2/d_i^2}} \right) \rrbracket.$$

By the compactness of Brakke flows, up to a subsequence, μ_t^i converges to a Brakke flow $\{\nu_t\}_{t < 0}$ in the sense of Radon measures. Computing directly, we have

$$(5.6) \quad \mu_t^i = \mathcal{H}^2 \llbracket \left(\sqrt{-t} \left(\frac{\Sigma_i - \tilde{\rho}_i y_i}{\sqrt{1 - \tilde{\rho}_i^2/d_i^2}} \right) \right) \rrbracket,$$

where $\tilde{\rho}_i = \frac{\rho_i}{\sqrt{1 - (1+t)(1 - \rho_i^2/d_i^2)}}$. Note that for any $t < 0$ fixed,

$$\frac{\tilde{\rho}_i}{d_i} = \frac{\rho_i/d_i}{\sqrt{1 - (1+t)(1 - \rho_i^2/d_i^2)}} = \frac{\rho_i/d_i}{\sqrt{\rho_i^2/d_i^2 - (1 - \rho_i^2/d_i^2)t}}.$$

Hence $\frac{\tilde{\rho}_i}{d_i} < 1$ for sufficiently large i . Moreover,

$$\begin{aligned} 1 - \frac{\tilde{\rho}_i^2}{d_i^2} &= \frac{-t(1 - \rho_i^2/d_i^2)}{\rho_i^2/d_i^2 - (1 - \rho_i^2/d_i^2)t} \\ &> \frac{-t}{1-t} \left(1 - \frac{\rho_i^2}{d_i^2}\right). \end{aligned}$$

Since $d_i^2 - \rho_i^2 = d_i^2 - d_i\rho_i + d_i\rho_i - \rho_i^2 \geq \rho_i(d_i - \rho_i + 1) \geq \rho_i \rightarrow +\infty$, we have that $d_i^2 - \tilde{\rho}_i^2 \rightarrow +\infty$. Consequently, $d_i(d_i - \tilde{\rho}_i) \rightarrow +\infty$.

Now we can compute the F -functional as follows:

$$\begin{aligned} \int \Phi(x, t) d\nu_t &= \lim_{i \rightarrow \infty} \int \Phi(x, t) d\mu_t^i \\ &= \lim_{i \rightarrow \infty} \int_{\frac{\Sigma_i - \tilde{\rho}_i y_i}{\sqrt{1 - \tilde{\rho}_i^2/d_i^2}}} \Phi(x, -1) dx \\ &= \lim_{i \rightarrow \infty} F_{\tilde{\rho}_i y_i, 1 - \frac{\tilde{\rho}_i^2}{d_i^2}}(\Sigma_i) \\ &= \lambda_1. \end{aligned}$$

It remains to show that ν_{-1} supports on a smooth self-shrinker. Indeed, by monotonicity formulas, ν_{-1} is F -stationary, i.e.

$$\vec{H} + \frac{S(x)^\perp \cdot x}{2} = 0, \quad \nu_{-1} - a.e. \ x.$$

Now we are going to prove that ν_{-1} splits off a line. For any $\tau \in \mathbb{R}$, set

$$\rho'_i = \rho_i + \tau \sqrt{1 - \frac{\rho_i^2}{d_i^2}} \quad \text{and} \quad s_i = \frac{1 - \rho_i'^2/d_i^2}{1 - \rho_i^2/d_i^2}.$$

It follows that $d_i(d_i - \rho'_i) \rightarrow +\infty$ and $s_i \rightarrow 1$ as $i \rightarrow \infty$ if $d_i(d_i - \rho_i) \rightarrow \infty$ and $\rho_i \rightarrow \infty$. Thus $\nu_{-1} - \tau y$, the limit of $\mu_{-1}^i - \tau y_i = \mathcal{H}^2 \lfloor \sqrt{s_i} \frac{\Sigma_i - \rho'_i y_i}{\sqrt{1 - \rho_i'^2/d_i^2}} \rfloor$, is also F -stationary. This deduces that

$$\vec{H}(x + y) + \frac{S(x + y)^\perp \cdot (x + y)}{2} = 0, \nu_{-1} - a.e.$$

Therefore, $S(x)^\perp y = 0$, which implies that ν_{-1} splits off a line. Note $F_{0,1}(\nu_{-1}) = \lambda_1$. It follows that ν_{-1} is supporting on $S^1(\sqrt{2}) \times \mathbb{R}_y$ with multiplicity 1. Then by the local regularity theorem for Brakke flows, we conclude that $\frac{\Sigma_i - \rho_i y_i}{\sqrt{1 - \rho_i^2/d_i^2}}$ locally smoothly converges to $S^1(\sqrt{2}) \times \mathbb{R}$ up to a subsequence.

The second statement follows from the fact that $\frac{d_i}{2}(\Sigma_i - d_i y_i)$ locally smoothly converges to a Bowl soliton Σ_y . $\square$

Step 3: For sufficiently large i , $\Sigma_i \cap \mathcal{C}_{U,R} \subseteq N_i$, where N_i (see Figure III) is defined as

$$N_i = \bigcup_{\rho=0}^{d_i-10/d_i} B_{10\sqrt{1-\rho^2/d_i^2}}^{\rho y_i}(\rho y_i) \bigcup_{\rho=d_i-20/d_i}^{d_i} B_{10\sqrt{1+(d_i-\rho)/d_i}}^{\rho y_i}(\rho y_i).$$

FIGURE III. Boundary of N_i .

Proof of Step 3. Assume on the contrary that a subsequence of $\Sigma_i \cap \mathcal{C}_{U,R}$ is not contained in N_i . We still denote the subsequence by $\{\Sigma_i\}$. Then for each i , $\Sigma_i \cap \mathcal{C}_{U,R} \cap \partial N_i \neq \emptyset$. Take $p_i \in \Sigma_i \cap \mathcal{C}_{U,R} \cap \partial N_i$ and set $\rho_i = \langle p_i, y_i \rangle$.

If ρ_i has a bounded subsequence, take a subsequence of p_i (still denoted by p_i) such that $\lim \rho_i = \rho_\infty > R$. Then for large i , $p_i \in \partial B_{10\sqrt{1-\rho_i^2/d_i^2}}^{\rho_i y_i}(\rho_i y_i)$, which implies that

$$(5.7) \quad \left| \frac{p_i - \rho_i y_i}{\sqrt{1 - \rho_i^2/d_i^2}} \right| = 10,$$

Recall that $\Sigma_i - \rho_i y_i$ locally smoothly converges to $\Sigma - \rho_\infty y$. Then (5.7) contradicts the fact that $\Sigma - \rho y$ is close to $S^1(\sqrt{2}) \times \mathbb{R}_y$ near the origin for $\rho > R$.

If $\rho_i \rightarrow \infty$ and $d_i(d_i - \rho_i)$ is unbounded, take a subsequence of p_i (still denoted by p_i) such that $\lim d_i(d_i - \rho_i) = +\infty$. Then again $p_i \in \partial B_{10\sqrt{1-\rho_i^2/d_i^2}}^{\rho_i y_i}(\rho_i y_i)$, which implies that

$$\left| \frac{p_i - \rho_i y_i}{\sqrt{1 - \rho_i^2/d_i^2}} \right| = 10.$$

This contradicts the fact that $\frac{\Sigma_i - \rho_i y_i}{\sqrt{1 - \rho_i^2/d_i^2}}$ locally smoothly converges to $S^1(\sqrt{2}) \times \mathbb{R}_y$.

It remains to consider the case that $\rho_i \rightarrow \infty$ and $d_i(d_i - \rho_i)$ is bounded. Note that the boundedness of $d_i(d_i - \rho_i)$ implies that $d_i^2 - \rho_i^2 = (d_i + \rho_i)(d_i - \rho_i)$ is also bounded. Let τ be a limit of $d_i(d_i - \rho_i)$ (up to a subsequence). Then

$$d_i(\Sigma_i - \rho_i y_i) = d_i(\Sigma_i - d_i y_i) + d_i(d_i - \rho_i) y_i \rightarrow 2\Sigma_y + \tau y,$$

where Σ_y is the Bowl soliton with direction $-y$. Denote by p the limit of $d_i(p_i - \rho_i y_i)$. Then

$$(5.8) \quad p \in \{x \in \mathbb{R}^3 : \langle x, y \rangle = 0\} \cap (2\Sigma_y + \tau y).$$

Let $u : [0, \infty) \rightarrow \mathbb{R}$ be the graph function of $\Sigma_y \cap (\mathbb{R}_y \times \mathbb{R}_p)$ (see Figure IV). Then u satisfies (see [1, (8.7)])

$$\frac{u''}{1 + u'^2} + \frac{u'}{r} = \frac{1}{2},$$

FIGURE IV. Bowl soliton.

and $u(0) = u'(0) = 0$. Since the Bowl soliton is convex, $u'' \geq 0$ everywhere. Hence $u'' + u'/r \geq 1/2$, which implies that $u \geq r^2/8$. Then a direct computation and (5.8) give that

$$(5.9) \quad |p| \leq 4\sqrt{\tau}.$$

Now we are going to estimate $|p|$ by the choice of p_i to get a contradiction.

Claim 1. For ρ_i satisfying $\lim_{i \rightarrow \infty} d_i(d_i - \rho_i) < \infty$, we always have

$$|p_i - \rho_i y_i| \geq 10\sqrt{1 + d_i(d_i - \rho_i)}/d_i.$$

Proof. If $\rho_i \leq d_i - 20/d_i$ for large i , then we always have

$$|p_i - \rho_i y_i| = 10\sqrt{1 - \rho^2/d_i^2}.$$

Recall that the boundedness of $d_i(d_i - \rho_i)$ implies that $\rho_i > d_i - 1$ for large i and

$$(5.10) \quad 10\sqrt{1 - \rho^2/d_i^2} \geq 10\sqrt{1 + d_i(d_i - \rho)}/d_i$$

is equivalent to $\rho^2 - d_i \rho + 1 \leq 0$. Hence, the claim holds true in this case.

We now consider $\rho_i > d_i - 20/d_i$. In this case, the claim follows from the definition of N_i . $\square$

Claim 1 gives that

$$|p| = \lim_{i \rightarrow \infty} d_i \cdot |p_i - \rho_i y_i| \geq \lim_{i \rightarrow \infty} 10\sqrt{1 + d_i(d_i - \rho)} = 10\sqrt{1 + \tau}.$$

which leads to a contradiction to (5.9).

Above all, we must have $(\Sigma_i \cap \mathcal{C}_{U,R}) \subseteq N_i$. $\square$

Now we argue by contradiction to study the topology of $\Sigma_i \cap \mathcal{C}_{U,R} = \Sigma_i \cap N_i$ outside a small neighborhood of $d_i y_i$. We want to prove that the ends $\Sigma_i \cap \mathcal{C}_{U,R}$ outside a small

neighborhood of $d_i y_i$ are topological cylinders. Let us first sketch the ideas. Roughly speaking, suppose that $r_i y_i$ are the bad points, in the sense that they prevent the parts of ends being cylinder, then

- if r_i is bounded, it contradicts the fact that Σ_i locally smoothly converges to Σ ;
- if r_i is unbounded and $d_i(d_i - r_i)$ is unbounded, we apply Step 2 to get a contradiction;
- if r_i is unbounded and $d_i(d_i - r_i)$ is bounded, we use Theorem 2.8 to get a contradiction.

The following are the precise arguments.

Step 4: $\Sigma_i \cap \mathcal{C}_{U,R} \setminus \mathcal{C}_{U,d_i-20/d_i}$ is diffeomorphic to $\Sigma \cap \mathcal{C}_{U,R} \setminus \mathcal{C}_{U,d_i-20/d_i}$.

Proof of Step 4. It suffices to show that there exists $I > 0$ such that for any $i > I$ and $r_i \in (R, d_i - 20/d_i)$, $\Sigma_i \cap \bigcup_{\rho=r_i}^{r_i+\sqrt{1-r_i^2/d_i^2}} B_{\sqrt{1-\rho^2/d_i^2}}^{\rho y_i}(\rho y_i)$ is diffeomorphic to $S^1 \times [0, 1]$. Set

$$\Sigma_i(r_i) := \Sigma_i \cap \bigcup_{\rho=r_i}^{r_i+\sqrt{1-r_i^2/d_i^2}} B_{\sqrt{1-\rho^2/d_i^2}}^{\rho y_i}(\rho y_i).$$

We argue by contradiction. Assume the statement is not true. Then for each i , take $r_i \in (R, d_i - 20/d_i)$ such that $\Sigma_i(r_i)$ is not diffeomorphic to $S^1 \times [0, 1]$.

If r_i is bounded, let $r > 0$ so that a subsequence of r_i (still denoted by r_i) satisfies $\lim r_i = r$. Then $\Sigma_i(r_i)$ locally smoothly converges to $(\Sigma - ry) \cap B_2(0) \times [0, 1]$, which leads to a contradiction.

If r_i is unbounded and $d_i(d_i - r_i)$ is unbounded, take a subsequence of r_i (still denoted by r_i) such that $\lim r_i = +\infty$ and $\lim d_i(d_i - r_i) = +\infty$. Then by Step 2, $\frac{\Sigma_i - r_i y_i}{\sqrt{1-r_i^2/d_i^2}}$ locally smoothly converges to $S^1(\sqrt{2}) \times \mathbb{R}_y$. Therefore, $\frac{\Sigma_i(r_i) - r_i y_i}{\sqrt{1-r_i^2/d_i^2}}$ locally smoothly converges to $S^1(\sqrt{2}) \times [0, 1]$, which leads to a contradiction.

If r_i is unbounded but $d_i(d_i - r_i)$ is bounded, take a subsequence of r_i (still denoted by r_i) such that $\lim r_i = +\infty$ and $\lim d_i(d_i - r_i) = \alpha$. Then by assumptions, $\alpha \geq 20$. Note that

$$\lim_{i \rightarrow \infty} d_i \sqrt{1 - r_i^2/d_i^2} = \lim_{i \rightarrow \infty} \sqrt{(d_i + r_i)(d_i - r_i)} = \sqrt{2\alpha}.$$

Together with Theorem 2.8, it follows that $\frac{\Sigma_i(r_i) - r_i y_i}{\sqrt{1-r_i^2/d_i^2}}$ locally smoothly converges to $\frac{2\Sigma_y + \alpha y}{\sqrt{2\alpha}} \cap (B_2(0) \times [0, 1])$, which leads to a contradiction. $\square$

We now proceed the proof of Proposition 5.7. Step 2 suggests that $\Sigma_i \cap \mathcal{C}_{U,d_i-20/d_i}$ is diffeomorphic to a bowl soliton, i.e. a disk; Step 4 suggests that $\Sigma_i \cap \mathcal{C}_{U,R} \setminus \mathcal{C}_{U,d_i-20/d_i}$ is diffeomorphic to a cylinder. Step 3 suggests that $\Sigma_i \cap \mathcal{C}_{U,R}$ is just the union of these two parts. Now the proposition follows from these steps. $\square$

Proof of Theorem 3.1. The desired results follow from Theorem 4.2 and Proposition 5.1, 5.5 and 5.7. $\square$

APPENDIX A. CLASSIFICATION OF SELF-SHRINKER ENDS WITH FINITE GENUS

In [38], L. Wang classified properly embedded self-shrinker ends with finite topology. The key step is to prove the multiplicity of the tangent flow is 1. Though L. Wang stated the result for finite topology, the argument works for all self-shrinkers with finite genus. We state it here and sketch the proof for completeness.

Theorem A.1 ([38, Theorem 1.1]). *Let $M \in \mathcal{M}_S(\Lambda, g)$ and $y \in S^2(1)$ such that*

$$\lim_{\rho \rightarrow \infty} F_{\rho y, 1}(M) \geq 1.$$

Then $\Sigma - \rho y$ locally smoothly converges to $\mathbb{R}_y \times \mathbb{R}$ or $\mathbb{R}_y \times S^1(\sqrt{2})$.

Proof. In this proof, we will use the same notations as in [38].

The theorem follows from [38, §5] if we prove [38, Theorem 1.3] under the assumptions of finite genus instead of finite topology. Then there exists $\rho_0 > 0$ so that $M \setminus B_{\rho_0}(0)$ does not contain two simple closed curves with intersection number 1. Now take a connected component of $M \setminus B_{\rho_0}(0)$, still denoted by M . Then there exists $\rho_1 > 0$ so that $\partial M \subset B_{\rho_1}(0)$ and ∂M lie in a connected component of $M \cap B_{\rho_1}(0)$.

Note that [38, Proposition 4.1] holds true under the assumption of finite genus. Hence the desired results follow from [38, Theorem 1.3]. We now sketch the proof by pointing out the necessary modifications.

Without any changes, L. Wang's argument [38, Proof of Theorem 1.3] shows that given i sufficiently large, for any $k \geq i$ and any connected component

$$M^{i,k} \subset M \cap (I_y(\tau_k, \tau_{k+1}) \times B_{R_i}^2),$$

the two sets

$$\partial M^{i,k} \cap \{x \in \mathbb{R}^3 : x \cdot y = \tau_k\} \text{ and } \partial M^{i,k} \cap \{x \in \mathbb{R}^3 : x \cdot y = \tau_{k+1}\}$$

are union of the same number of elements of $\{\gamma_j^{i,k}\}$ and $\{\gamma_j^{i,k+1}\}$, respectively. Hence the connected components of $M \cap B_{R_i}^3(\tau_i y)$ extend to infinity for all i sufficiently large.

Then similar to [38, Proof of Theorem 1.3], it suffices to show that, for sufficiently large i , the connected components of $M \cap B_{R_i}^3(\tau_i y)$ remain disjoint in the solid half-cylinder $I_y(\tau_i, +\infty) \times B_{R_i}^2$.

Suppose not, then there exist $i > 0$ and $k_l \rightarrow +\infty$ so that two elements of $\{\gamma_j^{i,k_l}\}_{j=1}^L$ lie in the boundary of a connected component of $M \cap (I_y(\tau_{k_l}, \tau_{k_l+1}) \times B_{R_i}^2)$. Hence there exist k_0 sufficiently large and a connected component of $M \cap (I_y(\tau_{k_0}, +\infty) \times B_{R_i}^2)$, denoted by N^{i,k_0} , which is connected at infinity, so that the boundary of a connected component of $N^{i,k_0} \cap (I_y(\tau_{k_0}, \tau_{k_0+1}) \times B_{R_i}^2)$ contains two of $\{\gamma_j^{i,k_0+1}\}_{j=1}^L$. Denote such a component by M^{i,k_0} . Then we can write

$$\partial M^{i,k_0} \cap \{x \in \mathbb{R}^3 : x \cdot y = \tau_{k_0+1}\} = \bigcup_{l=1}^m \gamma_{j_l}^{i,k_0+1},$$

where $m \geq 2$.

Then the following argument is similar to [38, Proof of Theorem 1.3]. Indeed, we can take $\{z_l\}_{l=1}^2, \{\gamma_l\}_{l=1}^2, \gamma$ like L. Wang did in [38, Proof of Theorem 1.3]. As N^{i,k_0} is connected at infinity, there exist points $z'_1 \in \gamma_1$ and $z'_2 \in \gamma_2$ so that they can be jointed by a smooth

embedded curve γ' in N^{i,k_0} outside a large enough ball so that $\gamma' \cap \gamma_{j_1}^{i,k_0+1} = \emptyset$. Denote the part of γ connecting z'_1 and z'_2 that contains γ_0 by γ'' . Thus, by a smoothing process, $\overline{\gamma' \cup \gamma''}$ can be made a simple closed smooth curve in M that transversally intersects $\gamma_{j_1}^{i,k_0+1}$ only at z_1 . Then from the curve $\gamma_{j_1}^{i,k_0+1}$ we will construct, in the following paragraphs, a simple closed curve that transversally intersects $\overline{\gamma' \cup \gamma''}$ only at one point. This leads to a contradiction.

If $\Sigma^{\tau_{k_0+1}} = \mathbb{R}_y \times \mathbb{S}^1$, then $\gamma_{j_1}^{i,k_0+1}$ is closed and we are done.

If $\Sigma^{\tau_{k_0+1}} = \mathbb{R}_y \times \mathbb{R}$, then by using the maximum principle for self-shrinkers [38, Lemma 4.2], L. Wang has proved that there is a curve $\tilde{\gamma}_l$ in M jointing z_l to ∂M for $l = 1, 2$ which is disjoint with γ_0 . Note that ∂M lies on a connected component of $M \cap B_{\rho_1}(0)$. Hence z_1 and z_2 can be jointed by a curve, denoted by $\gamma_{1,2}$, which is disjoint with γ_0 or $\overline{\gamma' \cup \gamma''}$. Thus, by a smooth process again, $\overline{\gamma_{1,2} \cup \gamma_0}$ can be made the desired simple closed smooth curve. $\square$

Here is an application of Theorem A.1.

Lemma A.2. *Let M be a self-shrinker with finite entropy and genus. Then M has finitely many ends.*

Proof. Assume on the contrary that M has finite entropy and finite genus, but infinitely many ends.

Case 1: There are infinitely many cylindrical ends.

Suppose that M has cylindrical ends with direction y_i and $|y_i| = 1$. Assume that $y_i \rightarrow y$. Then

$$F_{\rho y, 1}(M) = \lim_{i \rightarrow \infty} F_{\rho y_i, 1}(M) \geq \lim_{i \rightarrow \infty} \lim_{r \rightarrow \infty} F_{r y_i, 1}(M) = \lambda_1.$$

By [38], $M - \rho y$ locally smoothly converges to $\mathbb{R}_y \times S^1(\sqrt{2})$ with multiplicity 1. Thus for any $R > 2$, there exists $L_0 > 0$ such that

$$(A.1) \quad M \cap \{x \in \mathbb{R}^3 : \langle x, y \rangle \geq L_0, \text{dist}(x, I_y(L_0, +\infty)) \in (2, R)\} = \emptyset.$$

Take i large enough such that $L_0|y_i - y| < 1$. Consider a smooth function $f \in C^\infty(0, +\infty)$ given by

$$f(\rho) = \rho^2 - \langle \rho y_i, y \rangle^2.$$

Note that $f(L_0) < 1$ and $\lim_{\rho \rightarrow \infty} f(\rho) = +\infty$. Hence there exists $L_1 > L_0$ such that $f(L_1) = R^2/4$. Together with (A.1), we have $(M - L_1 y_i) \cap B_{R/4}(0) = \emptyset$. This deduces that $F_{L_1 y_i, 1}(M) \leq C/R$, which contradicts

$$F_{L_1 y_i, 1}(M) \geq \lim_{\rho \rightarrow \infty} F_{\rho y_i, 1}(M) = \lambda_1.$$

Case 2: There are infinitely many conical ends.

Suppose that M admits infinitely many conical ends with links $\{\gamma_i\}$. Take $y_i \in \gamma_i$ and let $y = \lim_{i \rightarrow \infty} y_i$. Then

$$F_{\rho y, 1}(M) = \lim_{i \rightarrow \infty} F_{\rho y_i, 1}(M) \geq \lim_{i \rightarrow \infty} \lim_{r \rightarrow \infty} F_{r y_i, 1}(M) = 1.$$

By [38], $\Sigma - \rho y$ locally smoothly converges to $\mathbb{R}_y \times S^1(\sqrt{2})$ or $\mathbb{R}_y \times \mathbb{R}$. By the similar argument as in Case 1, the limit surface can not be a cylinder. Then it remains to rule out

the case that $M - \rho y$ locally smoothly converges to $\mathbb{R}_y \times \mathbb{R}$. Without loss of generality, we assume that $\text{dist}(y, \gamma_i) = \text{dist}(y, y_i)$, where $\text{dist}(\cdot, \cdot)$ is the distance function in $\mathbb{R}^3$.

Take $R \gg 1$. Then applying local regularity theorem [41, Theorem 3.1], we have that:

Lemma A.3. *There exists $\epsilon > 0$ so that, for any $x \in S^2(1)$, $\alpha > 0$ with*

$$(A.2) \quad \lim_{\rho \rightarrow \infty} F_{\rho x, 1}(M) = 1 \text{ and } 1 \leq F_{\alpha x, 1}(M) < 1 + \epsilon,$$

then $M \cap B_{3R}(\alpha x)$ can be written as a graph, with C^2 norm $\ll 1$, over a hyperplane P containing $\mathbb{R}_x$. Particularly,

$$d_H(P \cap B_{3R}(\alpha x), M \cap B_{3R}(\alpha x)) \ll 1,$$

where d_H is the Hausdorff distance.

Proof of Lemma A.3. The proof here is similar to Step 1 and 2 in Proposition 5.1, and we leave it to readers. $\square$

Claim 2. *There exists $\rho_1 > 0$ so that $F_{x, 1}(M) < 1 + \epsilon$ for all $x \in B_R(\rho y)$ and $\rho > \rho_1$.*

Proof of Claim 2. Since $\lim_{\rho \rightarrow \infty} F_{\rho y, 1}(M) = 1$, then we can take ρ_0 and r_0 so that $F_{x, 1}(M) < 1 + \epsilon$ for $x \in B_{r_0}(\rho_0 y)$. Let $\rho_1 = \rho_0 \cdot R/r_0$. Then monotonicity of $F_{\rho x, 1}(M)$ as a function of ρ suggests that such ρ_1 is the desired constant. $\square$

Now take k large enough so that $\rho_1 \text{dist}(y_k, y) < 1$. Set $\rho_2 = 10/\text{dist}(y_k, y)$. It follows that $\text{dist}(\rho_2 y, \mathcal{C}_{\gamma_k}) > 9$. Note that $x = y$ and $\alpha = \rho_2$ satisfy (A.2). By Lemma A.3, there exists a hyperplane P containing $\mathbb{R}_y$ so that $M \cap B_{3R}(\rho_2 y)$ can be written as a graph over P with

$$d_H(P \cap B_{3R}(\rho_2 y), M \cap B_{3R}(\rho_2 y)) \ll 1.$$

We now prove that:

Claim 3. $\mathcal{C}_{\gamma_k}$ closes to P in $B_R(\rho_2 y)$.

Proof of Claim 3. Since $\rho_2 y_k \in B_R(\rho_2 y)$, then $\mathcal{C}_{\gamma_k} \cap B_R(\rho_2 y) \neq \emptyset$. By Claim 2, for all $z \in \mathcal{C}_{\gamma_k} \cap B_R(\rho_2 y)$, we have that $x = z/|z|$ and $\alpha = |z|$ satisfy (A.2). By Lemma A.3, $\text{dist}(z, M \cap B_{3R}(z)) \ll 1$ for all $z \in \mathcal{C}_{\gamma_k} \cap B_R(\rho_2 y)$. By triangle inequalities,

$$\begin{aligned} \text{dist}(z, P) &\leq \text{dist}(z, M \cap B_R(z)) + d_H(M \cap B_R(z), P \cap B_R(z)) \\ &\leq \text{dist}(z, M \cap B_R(z)) + d_H(M \cap B_{3R}(\rho_2 y), P \cap B_{3R}(\rho_2 y)) \ll 1. \end{aligned}$$

$\square$

There are two cases, depending on whether $\rho_2 \gamma_k$ is completely contained in the ball $B_R(\rho_2 y)$. Let us first consider the case that $\rho_2 \gamma_k$ is not completely contained in $B_R(\rho_2 y)$. Let γ' be the connected component of $\rho_2 \gamma_k \cap B_R(\rho_2 y)$ containing $\rho_2 y_k$. Then by Claim 3 and the fact that $\rho_2 y_k$ is the closest point of γ' to $\rho_2 y$, γ' must be a curve close to half of $P \setminus \mathbb{R}_y$; see Figure V. Denote by γ'_1 and γ'_2 the two connected components of $\gamma' \setminus \{\rho_2 y_k\}$. Taking $z_1 \in \rho_2 \gamma_k$ so that $|z_1 - \rho_2 y_k| = R/2$, then we have

$$\begin{aligned} F_{0,1}(\mathcal{C}_{\gamma_k} - z_1) &\geq F_{0,1}(B_{R/4}(z_1) \cap \mathcal{C}_{\gamma'_1} - z_1) + F_{0,1}(B_{R/4}(z_1) \cap \mathcal{C}_{\gamma'_2} - z_1) \\ &\geq 1 - C/R + 1 - C/R > 3/2. \end{aligned}$$

Therefore,

$$\lim_{\rho \rightarrow \infty} F_{\rho z_1 / \rho_2, 1 + \rho^2 / \rho_2^2}(M) = F_{0,1}(\lim_{\rho \rightarrow \infty} \rho^{-1} M - z_1) > 3/2,$$

FIGURE V. $\rho_2 \gamma_k$ closes to a line.

which implies that

$$F_{z_1,2}(M) = F_{\rho_2 z_1 / \rho_2, 1 + \rho_2^2 / \rho_2^2}(M) \geq \lim_{\rho \rightarrow \infty} F_{\rho z_1 / \rho_2, 1 + \rho^2 / \rho_2^2}(M) > 3/2.$$

This gives a contradiction.

To complete the proof, it suffices to consider the case that $\rho_2 \gamma_k$ is contained in $B_R(\rho_2 y)$. Let $x_1, x_2 \in \rho_2 \gamma_k$ so that $|x_1 - x_2| = \text{diam}(\rho_2 \gamma_k)$ (see Figure VI). Set $\rho_3 = 2R/|x_1 - x_2|$. Then $x = x_1/|x_1|$ and $\alpha = \rho_3|x_1|$ satisfy (A.2). Hence by Lemma A.3, there exists a hyperplane P_1 containing $\mathbb{R}_{x_1}$ so that $M \cap B_{3R}(\rho_3 x_1)$ can be written as a graph over P_1 with

$$d_H(P_1 \cap B_{3R}(\rho_3 x_1), M \cap B_{3R}(\rho_3 x_1)) \ll 1.$$

FIGURE VI. $\rho_3 \gamma_k$ closes to a line.

Claim 4. $\mathcal{C}_{\gamma_k}$ closes to P_1 in $B_R(\rho_3 x_1)$.

Proof of Claim 3. Since $\rho_2 y_k \in B_R(\rho_2 y)$, then by Claim 2, $F_{z,1}(M) < 1 + \epsilon$ for all $z \in \rho_2 \gamma_k$. By monotonicity formulas, for any $z \in \rho_3 \rho_2 \gamma_k$, $x = z/|z|$ and $\alpha = |z|$ satisfy (A.2), i.e.

$$1 \leq F_{z,1}(M) < 1 + \epsilon.$$

Then Lemma A.3 gives that $\text{dist}(z, M \cap B_{3R}(z)) \ll 1$ for all $z \in \rho_3 \rho_2 \gamma_k$. By triangle inequalities,

$$\begin{aligned} \text{dist}(z, P_1) &\leq \text{dist}(z, M \cap B_R(z)) + d_H(M \cap B_R(z), P_1 \cap B_R(z)) \\ &\leq \text{dist}(z, M \cap B_R(z)) + d_H(M \cap B_{3R}(\rho_3 x_1), P_1 \cap B_{3R}(\rho_3 x_1)) \ll 1. \end{aligned}$$

□

To proceed the argument, we take $z_2 \in \rho_3 \rho_2 \gamma_k$ so that $|z_2 - \rho_3 x_1| = R$. Denote by γ_1'' and γ_2'' the two components of $\rho_2 \rho_3 \gamma_k \setminus \{\rho_3 x_1, \rho_3 x_2\}$. Then we have

$$\begin{aligned} F_{0,1}(\mathcal{C}_{\gamma_k} - z_2) &\geq F_{0,1}(B_R(z_2) \cap \mathcal{C}_{\gamma_1''} - z_2) + F_{0,1}(B_R(z_2) \cap \mathcal{C}_{\gamma_2''} - z_2) \\ &\geq 1 - C/R + 1 - C/R > 3/2. \end{aligned}$$

Therefore,

$$\lim_{\rho \rightarrow \infty} F_{\rho z_2 / \rho_2 \rho_3, 1 + \rho^2 / \rho_2^2 \rho_3^2}(M) = F_{0,1}(\mathcal{C}_{\gamma_k} - z_2) > 3/2,$$

which implies that

$$F_{z_2, 2}(M) = F_{\rho_2 \rho_3 z_2 / \rho_2 \rho_3, 1 + \rho_2^2 \rho_3^2 / \rho_2^2 \rho_3^2}(M) \geq \lim_{\rho \rightarrow \infty} F_{\rho z_2 / \rho_2 \rho_3, 1 + \rho^2 / \rho_2^2 \rho_3^2}(M) > 3/2.$$

This gives a contradiction. □

The following linear bound of $|A|$ is proven by Song [35]. Here we give a uniform bound for $M \in \mathcal{M}_S(\Lambda, g)$, which is used in Theorem 2.8.

Lemma A.4 (cf. [35, Theorem 19; 38, Appendix A.1]). *There exists a constant $C = C(\Lambda, g)$ such that for any self-shrinker $M \in \mathcal{M}_S(\Lambda, g)$,*

$$|A|(x) \leq C(|x| + 1).$$

Proof of Lemma A.4. Assume on the contrary that there exist a sequence of self-shrinkers $\{\Sigma_i\}$ with entropy bound Λ and genus bound g , and $x_k \in \Sigma_k$ such that

$$|A|(x_k) \geq k(|x_k| + 1).$$

By Colding-Minicozzi [19], Σ_k locally smoothly converges to a self-shrinker, denoted by Σ , with $\lambda(\Sigma) \leq \Lambda$ and $g(\Sigma) \leq g$. Without loss of generality, $\{x_k\}$ is assumed to be unbounded. Take $y_i \in B_{1/|x_i|}(x_i)$ such that

$$(|x_i|^{-1} - |y_i - x_i|)|A(y_i)| = \sup_{y \in B_{1/|x_i|}(x_i)} (|x_i|^{-1} - |y - x_i|)|A(y)|.$$

Set $r_i = (|x_i|^{-1} - |y_i - x_i|)/2$. Then $|A(y_k)|r_k \geq k/2$ and for $y \in B_{r_k}(y_k) \cap \Sigma_k$,

$$2r_k|A(y_k)| \geq (|x_k|^{-1} - |y - x_k|)|A(y)| \geq r_k|A(y)|.$$

Thus for $y \in B_{k/2}(0) \cap |A(y_k)|(\Sigma_k - y_k)$, $|A(y)| \leq 2$ and $|H(y)| \leq 1/k$. Let $k \rightarrow \infty$, $|A(y_k)|(\Sigma_k - y_k)$ locally smoothly converges to a minimal surface Γ .

Let y be a limit point of $y_k/|y_k|$. Then by Proposition 2.7, for any $(z, t) \in \mathbb{R}^3 \times (0, +\infty)$,

$$F_{z, t^2}(\Gamma) \leq \lim_{\rho \rightarrow \infty} F_{\rho y, 1}(\Sigma) \leq \lambda_1,$$

which implies that Γ has only one end. Thus Σ is a plane, which contradicts $|A_\Gamma(0)| = 1$. □

APPENDIX B. TRANSLATING SOLITONS WITH LOW ENTROPY

For completeness of this paper, we sketch the proof of classification of translating solitons with low entropy in $\mathbb{R}^3$. The argument is based on the proof by Choi-Haslhofer-Hershkovits [13, Theorem 1.2] and Hershkovits [25, Theorem 3]. We refer to [24] the computation of the entropy of translating solitons.

Theorem B.1. *Let $M \subseteq \mathbb{R}^3$ be a translating soliton with entropy less than 2. Then M is a plane or a Bowl soliton.*

Proof. Let y be the translating vector field of M , i.e.

$$H = -\langle y, \mathbf{n} \rangle.$$

For each $r > 0$, define the blow-down Brakke flow by

$$\mu_t^r = \mathcal{H}^2 \llcorner \left(\frac{M - ry}{\sqrt{1+r}} + \sqrt{1+r}(1+t)y \right).$$

By the compactness of Brakke flows (Lemma 2.5), as $r \rightarrow +\infty$, there exist a sequence $r_i \rightarrow \infty$ and a Brakke flow $\{\nu_t\}$ such that $\mu_t^{r_i} \rightarrow \nu_t$ in the sense of Radon measures for all $t < 0$.

We are going to show that ν_{-1} is F -stationary. Indeed, by computing directly,

$$\mu_t^r = \mathcal{H}^2 \llcorner \left(\sqrt{-t} \left(\frac{M - \tilde{r}y}{\sqrt{1+\tilde{r}}} \right) \right),$$

where $\tilde{r} = r - (1+t)(1+r)$. Note that for $t < 0$ fixed, $\tilde{r} \rightarrow +\infty$ as $r \rightarrow +\infty$.

Note that M is a translating soliton with direction y . Then by the monotonicity of F -functional, $F_{ry,1+r}(M)$ is monotonically increasing (see [18, (1.9)]). Denote by λ the limiting value. Then $\lambda < 2$ by the assumption of entropy bound.

Now computing the F -functional for the limiting flow gives

$$\int \Phi(x, t) d\nu_t = \lim_{r \rightarrow \infty} \int \Phi(x, t) d\mu_t^r = \lim_{r \rightarrow +\infty} F_{\tilde{r}y, 1+\tilde{r}}(\Sigma) = \lambda.$$

By the monotonicity formulas (see Lemma 2.2), ν_{-1} is F -stationary.

Now we can show that ν_{-1} is supporting on a smooth self-shrinker. Namely, by the same argument, for all $\tau \in \mathbb{R}$, the related varifold of the limit of $\mu_{-1}^{r_i+\tau\sqrt{1+r_i}}$ is also F -stationary. Recall that

$$\mu_{-1}^{r_i+\tau\sqrt{1+r_i}} = \mathcal{H}^2 \llcorner \sqrt{s_i} \left(\frac{M - r_i y}{\sqrt{1+r_i}} - \tau y \right),$$

where $s_i = \frac{\sqrt{1+r_i}}{\sqrt{1+r_i}+\tau}$. Note that $s_i \rightarrow 1$ as $r_i \rightarrow +\infty$. We conclude that $\nu_{-1} - \tau y$ is also stationary. This deduces that $y^\perp = 0$. It follows that ν_{-1} splits off a line.

Now ν_{-1} is a F -stationary cycle with entropy less than 2. It follows that ν_{-1} is supporting on a plane or a self-shrinker cylinder.

In the first case, M is a plane by Brakke local regularity or the argument in [6, Proposition 3.2]. In the second case, M is a Bowl soliton by Hershkovits [25, Theorem 3].

For completeness of this paper, we sketch the proof given by Hershkovits [25]. Firstly, using the Neck Improvement Theorem [9, Theorem 4.4], the end can be written as a graph

(see [25, §2] for details) of a function g over $\mathbb{R}^2 \setminus B_R(0)$ satisfying

$$g = \frac{1}{2}(x_1^2 + x_2^2) - \frac{1}{2} \log(x_1^2 + x_2^2) + O\left(\frac{1}{\sqrt{x_1^2 + x_2^2}}\right).$$

Then the claim follows from [31, Theorem A] once we prove M has only one end.

We now show that M admits only one end. We first claim that $|A|$ is bounded by Choi-Schoen's estimates [12].

Claim 5. *There exists $C = C(M)$ such that $|A|(x) \leq C$ for any $x \in M$, where A is the second fundamental form of M .*

Proof of Claim 5. Assume on the contrary that $x_i \in M$ satisfying $|A|(x_i) > i$. Take $q_i \in M$ such that

$$\max_{x \in B_1(p) \cap M} (1 - |x - x_i|)|A|(x) = (1 - |z_i - x_i|)|A|(z_i),$$

where $B_r(p)$ is the ball in $\mathbb{R}^3$.

Set $r_i = (1 - |z_i - x_i|)/2$. Then $r_i|A|(z_i) \rightarrow \infty$ and for any $x \in B_{r_i}(z_i)$, we have $|A|(x) \leq 2|A|(z_i)$. Hence $|A|(z_i)(M - z_i)$ locally smoothly converges to a hypersurface $\Gamma \subseteq \mathbb{R}^3$. Recall M is a translating soliton and then $|H| \leq 1$. It follows that Γ is a minimal surface. By Proposition 2.7, the entropy of Γ is less than 2, which implies that Γ is a plane. This contradicts the fact of $|A_\Gamma|(0) = 1$.

Thus we have proved Claim 5. $\square$

As a result of claim 5, there exists a constant $\delta > 0$ such that $F_{x,1}(M) \geq \delta$ for any $x \in M$.

To proceed our argument, assume on the contrary that $p_i = (x_{1,i}, x_{2,i}, y_i) \rightarrow \infty$ lies in M but not in the end given by the graph of g . Recall that $\frac{M-ry}{\sqrt{1+r}}$ locally smoothly converges to $S^1(\sqrt{2}) \times \mathbb{R}_y$, which yields that for any $k > 0$ and large i , $y_i < \frac{1}{2k}(x_{1,i}^2 + x_{2,i}^2)$.

Claim 6. *There exists $K > 0$ such that $Ky_i \leq -(x_{1,i}^2 + x_{2,i}^2)$ for all large i .*

Proof of Claim 6. Assume on the contrary that for any fixed $k > 0$, we have $ky_i \geq -(x_{1,i}^2 + x_{2,i}^2)$ for large i . Take $s_{i,k} > 0$ such that $p_i + s_{i,k}y \in \{ky = x_1^2 + x_2^2\}$ and set $r_i = (x_{1,i}^2 + x_{2,i}^2)/k$ (see Figure VII).

Because $\frac{M-ry}{\sqrt{1+r}}$ locally smoothly converges to $S^1(\sqrt{2}) \times \mathbb{R}_y$, we have $B_{\sqrt{x_{1,i}^2 + x_{2,i}^2}/2}(p_i + s_{i,k}y) \cap M = \emptyset$. Note that $y_i < \frac{1}{2k}(x_{1,i}^2 + x_{2,i}^2)$. Clearly,

$$x_{1,i}^2 + x_{2,i}^2 \leq 2ks_{i,k} \leq 4(x_{1,i}^2 + x_{2,i}^2).$$

Therefore,

$$B_{\sqrt{k}/4}(0) \cap \frac{M - p_i - s_{i,k}y}{\sqrt{1 + s_{i,k}}} = \emptyset,$$

which implies that

$$\lim_{k \rightarrow \infty} \lim_{i \rightarrow \infty} F_{s_{i,k}y, 1+s_{i,k}}(M - p_i) = 0.$$

Note that $M - p_i$ is also a translating soliton with direction y . Hence $F_{sy, 1+s}(M - p_i)$ is monotonically increasing (see [18, (1.9)]). This deduces that

$$\delta \leq \liminf_{i \rightarrow \infty} F_{0,1}(M - p_i) \leq \lim_{k \rightarrow \infty} \lim_{i \rightarrow \infty} F_{s_{i,k}y, 1+s_{i,k}}(M - p_i) = 0,$$

FIGURE VII. $p_i \rightarrow -\infty$.

which leads to a contradiction. $\square$

Now set $p_i^0 = (x_{1,i}, x_{2,i}, 0)$. By Claim 6, $Ky_i < -|p_i^0|^2$. In particular, $|y_i| \rightarrow \infty$ and $p_i^0 / \sqrt{1 + |y_i|}$ is bounded. Note that $\frac{M}{\sqrt{1+r}} = \mu_0^r$ vanishes as $r \rightarrow \infty$. Hence

$$\lim_{i \rightarrow \infty} F_{0,1+|y_i|}(M - p_i^0) = 0,$$

;see also Claim 6). On the other hand,

$$F_{0,1}(M - p_i) \leq F_{|y_i|y,1+|y_i|}(M - p_i) = F_{0,1+|y_i|}(M - p_i^0),$$

which leads to a contradiction.

Then we finish the proof. $\square$

REFERENCES

- [1] Sigurd Angenent, Panagiota Daskalopoulos, and Natasa Sesum, *Unique asymptotics of ancient convex mean curvature flow solutions*, J. Differential Geom. **111** (2019), no. 3, 381–455. MR3934596
- [2] Sigurd B. Angenent, *Shrinking doughnuts*, Nonlinear diffusion equations and their equilibrium states, 3 (Gregynog, 1989), 1992, pp. 21–38. MR1167827
- [3] Matthias Bauer, *Existence of minimizing Willmore surfaces of prescribed genus*, Int. Math. Res. Not. **10** (2003), 553–576. MR1941840, MR1941840
- [4] Jacob Bernstein and Lu Wang, *A sharp lower bound for the entropy of closed hypersurfaces up to dimension six*, Invent. Math. **206** (2016), no. 3, 601–627. MR3573969
- [5] ———, *A topological property of asymptotically conical self-shrinkers of small entropy*, Duke Math. J. **166** (2017), no. 3, 403–435. MR3606722
- [6] ———, *Hausdorff stability of the round two-sphere under small perturbations of the entropy*, Math. Res. Lett. **25** (2018), no. 2, 347–365. MR3826825
- [7] Kenneth A. Brakke, *The motion of a surface by its mean curvature*, Mathematical Notes, vol. 20, Princeton University Press, Princeton, N.J., 1978. MR485012
- [8] Simon Brendle, *Embedded self-similar shrinkers of genus 0*, Ann. of Math. (2) **183** (2016), no. 2, 715–728. MR3450486

- [9] Simon Brendle and Kyeongsu Choi, *Uniqueness of convex ancient solutions to mean curvature flow in $\mathbb{R}^3$* , Invent. Math. **217** (2019), no. 1, 35–76. MR3958790
- [10] Xu Cheng, Tito Mejia, and Detang Zhou, *Stability and compactness for complete f -minimal surfaces*, Trans. Amer. Math. Soc. **367** (2015), no. 6, 4041–4059. MR3324919
- [11] Xu Cheng and Detang Zhou, *Volume estimate about shrinkers*, Proc. Amer. Math. Soc. **141** (2013), no. 2, 687–696. MR2996973
- [12] Hyeon In Choi and Richard Schoen, *The space of minimal embeddings of a surface into a three-dimensional manifold of positive Ricci curvature*, Invent. Math. **81** (1985), no. 3, 387–394. MR807063
- [13] Kyeongsu Choi, Robert Haslhofer, and O Hershkovits, *Ancient low entropy flows, mean convex neighborhoods, and uniqueness*, ArXiv e-prints (October 2018), available at arXiv:math/1810.08467.
- [14] Tobias H. Colding and William P. Minicozzi II, *The space of embedded minimal surfaces of fixed genus in a 3-manifold. I. Estimates off the axis for disks*, Ann. of Math. (2) **160** (2004), no. 1, 27–68. MR2119717
- [15] ———, *The space of embedded minimal surfaces of fixed genus in a 3-manifold. II. Multi-valued graphs in disks*, Ann. of Math. (2) **160** (2004), no. 1, 69–92. MR2119718
- [16] ———, *The space of embedded minimal surfaces of fixed genus in a 3-manifold. III. Planar domains*, Ann. of Math. (2) **160** (2004), no. 2, 523–572. MR2123932
- [17] ———, *The space of embedded minimal surfaces of fixed genus in a 3-manifold. IV. Locally simply connected*, Ann. of Math. (2) **160** (2004), no. 2, 573–615. MR2123933
- [18] ———, *Generic mean curvature flow I: generic singularities*, Ann. of Math. (2) **175** (2012), no. 2, 755–833. MR2993752
- [19] ———, *Smooth compactness of self-shrinkers*, Comment. Math. Helv. **87** (2012), no. 2, 463–475. MR2914856
- [20] ———, *The space of embedded minimal surfaces of fixed genus in a 3-manifold V; fixed genus*, Ann. of Math. (2) **181** (2015), no. 1, 1–153. MR3272923
- [21] Tobias Holck Colding, Tom Ilmanen, William P. Minicozzi II, and Brian White, *The round sphere minimizes entropy among closed self-shrinkers*, J. Differential Geom. **95** (2013), no. 1, 53–69. MR3128979
- [22] M. Gage, *The heat equation shrinking convex plane curves*, J. Differential Geom. **23** (1986), no. 1, 69–96. MR840401
- [23] Matthew A. Grayson, *The heat equation shrinks embedded plane curves to round points*, J. Differential Geom. **26** (1987), no. 2, 285–314. MR906392
- [24] Qiang Guang, *Volume growth, entropy and stability for translating solitons*, Comm. Anal. Geom. **27** (2019), no. 1, 47–72. MR3951020
- [25] Or Hershkovits, *Translators asymptotic to cylinders*, to appear in Crelle’s Journal, available at arXiv:math/1805.10553.
- [26] Gerhard Huisken, *Asymptotic behavior for singularities of the mean curvature flow*, J. Differential Geom. **31** (1990), no. 1, 285–299. MR1030675
- [27] Tom Ilmanen, *Elliptic regularization and partial regularity for motion by mean curvature*, Mem. Amer. Math. Soc. **108** (1994), no. 520, x+90. MR1196160
- [28] ———, *Singularities of mean curvature flow of surfaces*, preprint (1995).
- [29] Nikolaos Kapouleas, Stephen James Kleene, and Niels Martin Møller, *Mean curvature self-shrinkers of high genus: non-compact examples*, J. Reine Angew. Math. **739** (2018), 1–39. MR3808256
- [30] Daniel Ketover and Xin Zhou, *Entropy of closed surfaces and min-max theory*, J. Differential Geom. **110** (2018), no. 1, 31–71. MR3851744
- [31] Francisco Martín, Andreas Savas-Halilaj, and Knut Smoczyk, *On the topology of translating solitons of the mean curvature flow*, Calc. Var. Partial Differential Equations **54** (2015), no. 3, 2853–2882. MR3412395
- [32] William H. Meeks III and Harold Rosenberg, *The uniqueness of the helicoid*, Ann. of Math. (2) **161** (2005), no. 2, 727–758. MR2153399
- [33] Xuan Hien Nguyen, *Construction of complete embedded self-similar surfaces under mean curvature flow, Part III*, Duke Math. J. **163** (2014), no. 11, 2023–2056. MR3263027
- [34] Leon Simon, *Existence of surfaces minimizing the Willmore functional*, Comm. Anal. Geom. **1** (1993), no. 2, 281–326. MR1243525
- [35] Antoine Song, *A maximum principle for self-shrinkers and some consequences*, ArXiv e-prints (December 2014), available at arXiv:math/1412.4755.

- [36] Ao Sun, *Local Entropy and Generic Multiplicity One Singularities of Mean Curvature Flow of Surfaces*, ArXiv e-prints (October 2018), available at arXiv:math/1810.08114.
- [37] Lu Wang, *Uniqueness of self-similar shrinkers with asymptotically conical ends*, J. Amer. Math. Soc. **27** (2014), no. 3, 613–638. MR3194490
- [38] ———, *Asymptotic structure of self-shrinkers*, ArXiv e-prints (October 2016), available at arXiv:math/1610.04904.
- [39] ———, *Uniqueness of self-similar shrinkers with asymptotically cylindrical ends*, J. Reine Angew. Math. **715** (2016), 207–230. MR3507924
- [40] Brian White, *Partial regularity of mean-convex hypersurfaces flowing by mean curvature*, Internat. Math. Res. Notices **4** (1994), 1073–7928. MR1266114
- [41] ———, *A local regularity theorem for mean curvature flow*, Ann. of Math. (2) **161** (2005), no. 3, 1487–1519. MR2180405
- [42] ———, *Currents and flat chains associated to varifolds, with an application to mean curvature flow*, Duke Math. J. **148** (2009), no. 1, 41–62. MR2515099
- [43] Jonathan J. Zhu, *On the entropy of closed hypersurfaces and singular self-shrinkers*, to appear in JDG, available at arXiv:math/1607.07760.

DEPARTMENT OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, 77 MASSACHUSETTS AVENUE, CAMBRIDGE, MA 02139, USA

E-mail address: aosun@mit.edu

MAX-PLANCK INSTITUTE FOR MATHEMATICS, VIVATSGASSE 7, 53111 BONN, GERMANY

E-mail address: wangzhichaonk@gmail.com