


Current Biology

Symbiont Digestive Range Reflects Host Plant Breadth in Herbivorous Beetles

Graphical Abstract



Authors

Hassan Salem, Roy Kirsch, Yannick Pauchet, ..., Donald Windsor, Takema Fukatsu, Nicole M. Gerardo

Correspondence

hassan.salem@tuebingen.mpg.de

In Brief

Tortoise leaf beetles rely on the symbiotic bacterium, *Stammera*, to digest foliage rich in pectin. Salem et al. reveal that *Stammera* varies in the pectinases it encodes and supplements. *Stammera* encoding a more dynamic digestive range allows its host to overcome a greater diversity of plant polysaccharides, corresponding to a wider ecological distribution.

Highlights

- *Stammera* genomes are structurally conserved across Cassidinae species
- Symbiont pectinases complement the host's endogenous cellulases and xylanases
- *Stammera* differentially encode pectinases in their reduced genomes
- Symbiont pectinolytic range reflects host plant breadth



Article

Symbiont Digestive Range Reflects Host Plant Breadth in Herbivorous Beetles

Hassan Salem,^{1,2,3,8,*} Roy Kirsch,⁴ Yannick Pauchet,⁴ Aileen Berasategui,¹ Kayoko Fukumori,⁵ Minoru Moriyama,⁵ Michael Cripps,⁶ Donald Windsor,⁷ Takema Fukatsu,⁵ and Nicole M. Gerardo¹

¹Department of Biology, Emory University, Atlanta, GA 30322, USA

²National Museum of Natural History, Smithsonian Institution, Washington, DC 20560, USA

³Mutualisms Research Group, Max Planck Institute for Developmental Biology, Tübingen 72076, Germany

⁴Department of Entomology, Max Planck Institute for Chemical Ecology, Jena 07745, Germany

⁵National Institute for Advanced Industrial Science and Technology, Tsukuba 305-8566, Japan

⁶AgResearch, Lincoln Research Centre, Lincoln 7608, New Zealand

⁷Smithsonian Tropical Research Institute, Panama City 0843-03092, Panama

⁸Lead Contact

*Correspondence: hassan.salem@tuebingen.mpg.de

<https://doi.org/10.1016/j.cub.2020.05.043>

SUMMARY

Numerous adaptations are gained in light of a symbiotic lifestyle. Here, we investigated the obligate partnership between tortoise leaf beetles (Chrysomelidae: Cassidinae) and their pectinolytic *Stammera* symbionts to detail how changes to the bacterium's streamlined metabolic range can shape the digestive physiology and ecological opportunity of its herbivorous host. Comparative genomics of 13 *Stammera* strains revealed high functional conservation, highlighted by the universal presence of polygalacturonase, a primary pectinase targeting nature's most abundant pectic class, homogalacturonan (HG). Despite this conservation, we unexpectedly discovered a disparate distribution for rhamnogalacturonan lyase, a secondary pectinase hydrolyzing the pectic heteropolymer, rhamnogalacturonan I (RG-I). Consistent with the annotation of rhamnogalacturonan lyase in *Stammera*, cassidines are able to depolymerize RG-I relative to beetles whose symbionts lack the gene. Given the omnipresence of HG and RG-I in foliage, *Stammera* that encode pectinases targeting both substrates allow their hosts to overcome a greater diversity of plant cell wall polysaccharides and maximize access to the nutritionally rich cytosol. Possibly facilitated by their symbionts' expanded digestive range, cassidines additionally endowed with rhamnogalacturonan lyase appear to utilize a broader diversity of angiosperms than those beetles whose symbionts solely supplement polygalacturonase. Our findings highlight how symbiont metabolic diversity, in concert with host adaptations, may serve as a potential source of evolutionary innovations for herbivorous lineages.

INTRODUCTION

Beetles display a remarkable degree of adaptability to exploit a correspondingly diverse flora [1, 2]. As half of all beetles are herbivorous, traits mediating phytophagy are central to the evolutionary success of Earth's most speciose animal group [3]. Recent genomic insights into beetle diversification revealed that the two independent origins of specialized herbivory within the order coincided with the co-option of microbial enzymes to more efficiently digest complex plant polymers, such as cellulose, hemicellulose, and pectin [4–7]. By endowing phytophagous beetles with the catalytic tools to breach the plant cell wall, these enzymes are implicated in the evolution of correspondingly varied and specialized plant-feeding habits [8, 9], fueling the diversification of major herbivorous lineages [4].

Although horizontal gene transfer from bacteria and fungi allowed phytophagous beetles to endogenously maintain a battery of key plant-cell-wall-degrading enzymes [4, 7, 8], novel metabolic

features can also be integrated through symbiosis. The pectinolytic phenotype of the tortoise leaf beetle *Cassida rubiginosa* (Coleoptera: Chrysomelidae: Cassidinae) is entirely outsourced to its obligate γ -proteobacterial symbiont, *Candidatus Stammera capleta* (henceforth *Stammera*) [10]. Possessing a highly reduced genome (271 Kb), the symbiont's limited metabolism is streamlined to produce and export pectinolytic enzymes that enable *C. rubiginosa* to process foliage rich in recalcitrant pectins [10, 11]. To ensure that future host generations are endowed with *Stammera*, each egg is maternally provisioned at the anterior pole with a symbiont-harboring caplet as a vehicle for vertical transmission [10]. Infection by *Stammera* is initiated when an emerging larva consumes the caplet. As the sole source of pectinases for its host, the experimental removal of *Stammera* diminishes the digestive capacity of the host beetle, resulting in stunted growth and low survivorship [10].

Our initial characterization of the symbiosis revealed that the thistle-feeding tortoise leaf beetle *C. rubiginosa* relies on two classes of pectinases in its partnership with *Stammera* [10]: a



Table 1. List of Cassidine Leaf Beetles Included in This Study, Their Host Plants, and the Digestive Enzymes Encoded by Their *Stammera* Symbionts

Beetle Species	Tribe	Location	Host Plant Clade, Order	<i>Stammera</i> Digestive Enzymes		
				Polygalacturonase (GH 28)	Rhamnogalacturonan Lyase (PL 4)	α -Glucuronidase (GH 67)
<i>Cassida versicolor</i>	Cassidini	Japan	Fabids, Rosales [21]	+	+	
<i>Cassida viridis</i>	Cassidini	Germany	Lamiid, Lamiales [22]	+	+	
<i>Cassida vibex</i>	Cassidini	Germany	Campanulids, Asterales [23]	+	+	
<i>Cassida rubiginosa</i>	Cassidini	New Zealand	Campanulids, Asterales [24, 25]	+	+	
<i>Parachirida semiannulata</i>	Cassidini	Panama	Lamiid, Solanales [18]	+	+	
<i>Ischnocodia annulus</i>	Cassidini	Panama	Lamiid, Boraginales [18]	+		+
<i>Stolas discoides</i>	Mesomphaliini	Panama	Lamiid, Solanales [18]	+		+
<i>Chelymorpha alternans</i>	Mesomphaliini	Panama	Lamiid, Solanales [18]	+		+
<i>Acromis sparsa</i>	Mesomphaliini	Panama	Lamiid, Solanales [20]	+		
<i>Cistudinella foveolata</i>	Ischyrosomychini	Panama	Lamiid, Boraginales [18, 26]	+		+
<i>Discomorpha panamensis</i>	Omozerini	Panama	Lamiid, Boraginales [18, 26]	+		+
<i>Agroiconota porpinqua</i>	Cassidini	Panama	Lamiid, Solanales [18]	+		+
<i>Charidotella sexpunctata</i>	Cassidini	USA	Lamiid, Solanales [27]	+		+

Symbol (+) indicates the presence of an encoding gene within the symbiont's genome. GH, glycoside hydrolase; PL, polysaccharide lyase.

polygalacturonase that cleaves homogalacturonan (HG), the homopolymeric backbone of pectin, and a rhamnogalacturonan lyase that targets the heteropolymeric pectic region, rhamnogalacturonan I (RG-I). As the most plentiful polysaccharide of the primary plant cell wall [12], pectin is instrumental for embedding cellulose and xylan into a carbohydrate matrix that ensures the physical integrity of the cell and contributes to adhesion and signal transduction [13]. Although HG is typically more abundant than RG-I, both polysaccharidic sequences are universally present in the primary cell wall and comprise up to 95% of all pectic content in plant tissues [13, 14]. Because pectin degradation facilitates, and typically precedes, the downstream breakdown of other primary cell wall components like cellulose and hemicellulose [15, 16], symbiont-encoded pectinases are critical for *C. rubiginosa* to maximize the dietary value of ingested leaves.

The Cassidinae subfamily includes more than 6,000 species that are all herbivorous [17]. Despite a cosmopolitan distribution and their specializations to exploit a diverse range of angiosperms [17–20], our understanding of the metabolic features behind the origin and radiation of this highly speciose insect lineage is limited, especially in the context of its symbiosis with *Stammera*. Reconciling an extensive record of life history traits for tortoise leaf beetles with comparative genomics, transcriptomics, and biochemical assays, we explore the metabolic features of both host and symbiont to (1) characterize the range of digestive enzymes defining the partnership, (2) demonstrate how variation in symbiotic factors can shape the digestive physiology of the insect host, and (3) highlight how this variation might facilitate niche expansion and adaptation across a diverse group of herbivorous beetles.

RESULTS AND DISCUSSION

Stammera Genomes Are Structurally and Metabolically Conserved

DNA sequencing of the symbiotic organs associated with the foregut of 13 cassidine species, representing four taxonomic tribes and collected from four continents, revealed that all investigated tortoise leaf beetle species maintain a symbiosis with *Stammera* (Table 1). In light of the phylogenetic congruence between Cassidinae beetles and *Stammera* (Figure 1), we applied the reconciliation tool Jane 4 [28] to test for co-speciation. Our analysis outlined nine co-speciation events between host and symbiont (Table S1). Three events were attributed to duplication and host switching, and two were labeled as loss events. No other evolutionary events, such as duplication or failure to diverge, were revealed. Although the general pattern of co-cladogenesis (Figure 1) is in line with cassidines relying on egg caplets to vertically transmit *Stammera* [10], limited horizontal exchange of the symbiont cannot be ruled out. This is consistent with the coevolutionary dynamics reported for many insect groups that extracellularly transfer their symbionts [29], including the hemipteran families Acanthosomatidae [30], Plataspidae [31], and Urostylididae [32].

The assembled genomes of all 13 *Stammera* strains are drastically reduced in size (215–310 Kb) and highly AT-rich (83%–89%; Figure 1; Table S2). In addition to a circular chromosome, each *Stammera* possesses 1 to 2 plasmids that range in size from 3 to 5 Kb (Table S2). Encoding 232–296 putative protein-coding open reading frames (ORFs) with an average size of 912 bp, the resulting coding percentage is between 89% and 92%. All strains possess operons for the three structural




Figure 1. Tortoise Leaf Beetles (Chrysomelidae: Cassidinae) in Co-cladogenesis with Their *Stammera* Symbionts

(A) Cassidinae beetles.

(B) Maximum-likelihood (ML) phylogeny of the host is based on concatenated sequences of 18S and 28S ribosomal RNA and mitochondrial 16S rRNA genes. Symbiont phylogeny is based on ML analysis of a concatenated nucleotide alignment for 30 genes shared by 13 *Stammera* genomes. Diagrams of *Stammera*'s genomes are represented by circles with diameters corresponding to size. Genes coding for rRNA (red), metabolism (blue), and cellular processes (green) are indicated on each circle. Node coloration reflects bootstrap support. Photographs of Cassidinae species from left to right are as follows: *Cassida rubiginosa*; *Ischnocodia annulus*; *Chelymorpha alternans*; *Acromis sparsa*; *Discomorpha panamensis*; and *Charidotella sexpunctata*.

See also Table S1.

ribosomal RNA genes, as well as tRNA genes for all 20 amino acids (Table S2).

Similar to other bacterial symbionts possessing tiny genomes [33], *Stammera* retained gene families involved in the core processes of replication, transcription, and translation (Figures 2 and S1A). Our annotation also revealed that genes related to protein folding and stability are shared throughout, including the *GroES-GroEL* chaperonin complex and the bacterial heat shock protein 70 (Hsp70) (Figure S1A). Suggestive of an important role in the biology of bacterial symbionts [33], such chaperones are among the most highly expressed enzymes within the bacteriomes of aphids [34], cicadas [35], tsetse flies [36], sharpshooters [37], and leaf beetles [11]. In contrast, pathways for the synthesis of essential amino acids, cofactors, and lipids

are missing, as are genes encoding signal transduction, cell surface structures, and motility (Figure 2; Data S1).

Energy production and the recovery of reducing equivalents in *Stammera* are achieved through the oxidation of sugars to pyruvate via glycolysis, followed by the fermentation of pyruvate to lactate through the activity of lactate dehydrogenase [11]. The combination of glycolysis and obligate fermentation appears to be a common feature across this γ -proteobacterial lineage (Figure S1B), presumably mediated by the microbe's extracellular localization and relaxed selection to maintain the citric acid cycle. This represents a notable departure from the respiratory metabolism fueling many intracellular symbionts [11], where the citric acid cycle serves as a precursor to produce key micronutrients (e.g., essential amino acids) for their insect hosts



Figure 2. High Metabolic Conservation Identified in the Genomes of *Stammera* from 13 Cassidinae Species

Relative abundance of Clusters of Orthologous Groups (COG)-annotated functional groups is represented as a heatmap. Symbiont phylogeny, as in Figure 1, is based on a ML analysis of a concatenated nucleotide alignment for 30 genes shared by 13 *Stammera* genomes. See also Figure S1, Table S2, and Data S1.

[11, 38, 39]. In the four instances we could not annotate lactate dehydrogenase in *Stammera* (hosts: *Acromis sparsa*; *Chelymorpha alternans*; *Stolas discoides*; and *Cistudinella foveolata*), we also observed an incomplete glycolysis pathway, as well as a missing ATP synthase complex (Figure S1B). Given the monophyly of these four strains within the symbiont’s phylogeny (Figures 1 and S1B), it appears that the glycolytic and fermentative pathways were lost once, pointing toward the host presumably supplementing *Stammera* with the necessary energy equivalents as described for other microbes with highly reduced genomes, such as *Carsonella ruddii* in psyllids [39].

Other notable variation across *Stammera* genomes involves genes coding for peptidoglycan biosynthesis (Figure S1B). Although host-restricted symbionts can experience gene losses in all functional categories, pathways underlying production of cell envelope components are commonly depleted [40], as shown in endosymbionts like *Buchnera* in aphids [41], *Hodgkinia* in cicadas [42], and *Carsonella* in psyllids [39]. The differential distribution of these pathways in *Stammera* may indicate some

variation in the degree of metabolic dependence and the level of host control over microbial growth and proliferation within the symbiotic organs of cassidine beetles.

Variation in Symbiont-Derived Digestive Enzymes

Our first sequencing and annotation of a *Stammera* genome associated with *C. rubiginosa* revealed the presence of polygalacturonase and rhamnogalacturonan lyase as two pectinases mediating digestion of HG and RG-I, respectively [10]. Polygalacturonase appears foundational to the Cassidinae-*Stammera* symbiosis, evidenced by its distribution across all 13 symbiont genomes (Table 1). As an endo-active enzyme belonging to the glycoside hydrolase family 28 (GH 28), polygalacturonase hydrolyzes the backbone of HG at random intervals to generate galacturonic acid products of various sizes [7, 10].

In contrast to the conservation of polygalacturonase across all 13 genomes, rhamnogalacturonan lyase is encoded by a subset of *Stammera* strains (Table 1). The symbionts supplementing rhamnogalacturonan lyase partner with beetles distributed




Figure 3. Enzymatic Synergy between Host and Symbiont to Deconstruct the Primary Cell Wall

(A) Host- and *Stammera*-encoded plant-cell-wall-degrading enzymes (PCWDEs) as inferred from transcriptome profiling of seven representative cassidine species. Source of PCWDEs is designated by color, where gray denotes an endogenous beetle gene and green denotes a *Stammera*-encoded enzyme.

(B) Predicted mode of action of the annotated PCWDEs across cellulose, xylan, and pectin.

GalA, galacturonic acid; GH, glycoside hydrolase; GlcA, glucuronic acid; Glu, glucose; PL, polysaccharide lyase; Rha, rhamnose; Xyl, xylose. See also Tables S3 and S4.

across Panama, Germany, Japan, and New Zealand (Table 1). These beetles feed on phylogenetically diverse angiosperms, including plants within the lamiid, campanulid, and fabid eudicot clades (Table 1). Belonging to the polysaccharide lyase family 4 (PL 4), rhamnogalacturonan lyase cleaves the α -(1,4) glycosidic bonds between rhamnose and galacturonic acid of the RG-I backbone through a β -elimination reaction [43].

Furthermore, genome annotation revealed a third class of plant-cell-wall-degrading enzymes in *Stammera*: an α -glucuronidase (GH 67; Table 1). In contrast to the aforementioned pectinases, α -glucuronidase is predicted to cleave the glycosidic bonds linking the xylose backbone to glucuronic acid through a single displacement mechanism [44]. As noted for rhamnogalacturonan lyase, α -glucuronidase is not universally present in all *Stammera* strains (Table 1). But unlike rhamnogalacturonan lyase, the distribution of α -glucuronidase in our survey is restricted to symbionts associated with New World tropical beetle species specializing on host plants within the lamiid eudicot clade (Table 1). The binary distribution of rhamnogalacturonan lyase and α -glucuronidase in this symbiosis is striking (each occurs in the absence of the other), highlighting how *Stammera* can supplement different sets of digestive enzymes to its leaf beetle hosts.

Deconstructing the Plant Cell Wall Together: A Complementary Role for Host Enzymes

As with many phytophagous beetles [4, 5, 45–47], the tortoise leaf beetle *C. rubiginosa* maintains a range of cellulases and

xylanases endogenously [10]. This is attributed to a string of horizontal gene transfer events that coincided with an ancestral transition from fungivory to herbivory in the Coleoptera, currently dated at 250 mya [5, 8]. Acquired from a range of bacterial and fungal donors, many of these enzymes underwent a number of duplication events followed by functional diversification [4]. By evolving complementary catalytic activities, they allowed beetles to process cellulose and hemicellulose more efficiently [45–47]. Expanding on our previous transcriptional profiling, we sequenced six additional gut transcriptomes of cassidine beetles to characterize the range of endogenous plant-cell-wall-degrading enzymes in the subfamily and detail the degree of predicted synergy between host and microbe toward deconstructing the complex polysaccharidic networks dominating plant leaves. Conserved across the seven species of cassidines transcriptionally profiled here are four endogenous carbohydrate-active enzymatic families, including cellulases belonging to the GH families 9, 45, and 48, as well as a single xylanase belonging to the family GH 10 (Figure 3; Table S5). Genes encoding these enzymatic families are enriched in most chrysomelid genomes [4, 48].

Heterologous expression of endogenous GHs in phytophagous beetles revealed that GH 45 is functionally active and confers a cellulolytic phenotype, which is in line with descriptions from wider herbivorous taxa, such as nematodes [49], mollusks [50], and rotifers [51]. In contrast, functional characterization of GH 9 failed to confirm the enzyme's predicted cellulolytic activity

in the Chrysomelidae [45, 46], suggesting that it may have lost its enzymatic potential altogether. The low expression levels of GH 9 in leaf beetle guts relative to other cellulases support the latter hypothesis.

For the endogenous xylanase (Figure 3), GH 10 was previously annotated in the genome of the coffee berry borer, and heterologous expression demonstrated the enzyme's ability to effectively depolymerize the xylan backbone [47]. The optimal activity dynamics of the xylanase was shown to resemble conditions similar to the physiological medium within the beetle's digestive tract [47], allowing the insect to subsist solely on the coffee seed's endosperm, a diet composed primarily of hemicellulose polysaccharides.

Displaying a division of labor that appears to reflect no catalytic overlap (Figure 3), host- and *Stammera*-encoded enzymes contribute to the processing of cellulose, xyloglucan, and pectin: the three most abundant classes of polysaccharides within leaves. However, relative to the conserved battery of digestive enzymes encoded in the genomes of most phytophagous beetles [4], cassidines notably do not encode their own pectinases (Figure 3). Beyond the endogenous cellulases and xylanases, as well as the symbiotic polygalacturonase, rhamnogalacturonan lyase, and α -glucuronidase detected in the reduced genomes of *Stammera*, no additional enzymes involved in the processing of the plant cell wall could be annotated in our transcriptomic survey of the beetles. As our annotations are informed by characterized digestive enzymes spanning public databases, we cannot exclude the possibility that novel catalytic features are additionally encoded.

Two Pectinases Are Better Than One

Given the conservation of host-encoded cellulases and xylanases and variation of plant-cell-wall-degrading enzymes encoded by *Stammera* (Figure 3), we set out to compare the digestive physiology of two leaf beetle species possessing metabolically representative symbionts: *Cassida rubiginosa* and *Chelymorpha alternans*. *Stammera* in *C. rubiginosa* contributes polygalacturonase and rhamnogalacturonan lyase, although *Ch. alternans* harbors a symbiont encoding for polygalacturonase and α -glucuronidase (Table 1; Figure 3).

To confirm *in silico* predictions that *C. rubiginosa* should depolymerize a greater diversity of pectic substrates than *Ch. alternans*, we qualitatively analyzed the breakdown products of HG and RG-I by thin-layer chromatography (TLC). Because both leaf beetle species possess symbionts that encode for polygalacturonase, gut extracts tested against HG revealed that the insects can monomerize the homopolymer into galacturonic acid (Figure 4A). Although the monosaccharide is the sole product accumulating in *C. rubiginosa*, di-galacturonic acid, tri-galacturonic acid, and larger oligosaccharides also appear in assays featuring *Ch. alternans* (Figure 4A). Similarly, and consistent with earlier observations [10], *C. rubiginosa* is also able to depolymerize RG-I, evidenced by the accumulation of monogalacturonic acid as a breakdown product (Figure 4B). This is also the case for other cassidines whose symbionts encode rhamnogalacturonan lyase, including *Cassida vibex* and *Cassida viridis* (Figure S2). In contrast, monogalacturonic acid is not a breakdown product in *Ch. alternans* (Figure 4B). Instead, we observe the accumulation of a larger, uncharacterized molecule, which

also appears in *C. rubiginosa* (Figure 4), *C. vibex*, and *C. viridis* (Figure S2). Overall, relative to beetles only supplemented with *Stammera*'s polygalacturonase, the combination of symbiont-encoded primary and secondary pectinases endows a subset of cassidines with a more proficient pectinolytic phenotype across the two polysaccharidic sequences that form the pectic backbone (Figure 4).

The exact mode of action of rhamnogalacturonan lyases in *Stammera* is of particular interest. Its activity in phytopathogens and plants involves cleaving the α -(1,4) glycosidic bond between rhamnose and galacturonic acid at random intervals, producing oligomers of varying sizes at the non-reducing end [14]. Although we do not observe the additional buildup of larger oligosaccharides in our assays, the catalytic phenotype against RG-I is lost following aposymbiosis in *C. rubiginosa* [10], implicating rhamnogalacturonan lyase in this process, either through direct activity or in synergy with other digestive enzymes.

Despite possessing a more limited pectinolytic metabolism relative to *C. rubiginosa*, we hypothesized *Ch. alternans* to display a greater xylanolytic range, owing to its symbiont encoding α -glucuronidase instead of rhamnogalacturonan lyase (Table 1; Figure 3). Testing for activity against xylotriose linked to glucuronic acid, both *Ch. alternans* and *C. rubiginosa* display an ability to depolymerize the hemicellulose's backbone into monomers and oligomers of xylose (Figure 5). This is consistent with the annotation of the endogenous xylanase GH 10 in both species (Figure 3) and the aforementioned catalytic capacity of the enzyme in other phytophagous beetles [47]. Surprisingly, we could not characterize any activity in either species against the glycoside linkage between xylose and glucuronic acid (Figure 5), given the absence of the latter as a breakdown product in the TLC. This is especially notable for *Ch. alternans* in light of the annotation of an α -glucuronidase in its symbiont's genome (Table 1; Figure 3). This indicates that the enzyme does not upgrade *Ch. alternans*' digestive physiology against the hemicellulose. The loss of α -glucuronidase from the symbiont of *Acromis sparsa* suggests that, in at least one lamiiid-feeding Cassidinae species, the enzyme may no longer be adaptive.

Reflecting the dominant abundance of HG in angiosperm leaves [12, 13] and the pivotal role polygalacturonases played in the evolution of obligate herbivory in beetles [7], we highlight the conserved distribution of the primary pectinase in *Stammera* (Table 1) and emphasize its importance as a foundational enzyme for the stability of this partnership. Predicted to upgrade the nutritional ecology of herbivorous beetles by facilitating the exploitation of young plant tissues rich in HG (e.g., leaves, stems, seeds, and fruits), the acquisition of polygalacturonases following horizontal gene transfer [4] or symbiosis (Table 1; Figure 3) underscores their adaptive significance for phytophagous taxa [7].

Symbiont Pectinolytic Range Reflects Host Ecological Breadth

As the second most abundant polysaccharide in the plant cell wall, RG-I constitutes 20%–35% of the overall pectic content in plant tissues [13, 14]. Omnipresent in eudicots, an exceedingly diverse and abundant group of angiosperm plants, RG-I contributes to the functionality of the primary cell wall by enhancing its rigidity and firmness [52]. Because RG-I is covalently linked to




Figure 4. Polygalacturonase and Rhamnogalacturonan Lyase Endow a More Dynamic Pectinolytic Phenotype to Cassidine Leaf Beetles
(A and B) Thin-layer chromatogram (TLC) illustrating the breakdown products of pectinase activity against (A) HG and (B) RG-I using gut contents from *Cassida rubiginosa* and *Chelymormpha alternans*. GC, gut content; S, substrate; Std, standard; GalA, mono-galacturonic acid; GalA-2, di-galacturonic acid; GalA-3, tri-galacturonic acid; Rha, rhamnose. Dashed box highlights GalA differences across treatments. See also Figure S2.

HG, cleavage of the heteropolymeric backbone compromises the stability and mechanical properties of the pectin network [52, 53]. RG-I-degrading enzymes, like rhamnogalacturonan lyase, are typically found in the genomes of plants and their pathogens [54]. Activity of the secondary pectinase in plant roots, leaves, and fruits is tied to developmental modifications requiring plasticity during growth, ripening, and senescing [54]. In phytopathogens like *Dickeya* and *Bacillus*, rhamnogalacturonan lyase activity constitutes a pathogenicity factor by allowing the microbes to break through the primary plant cell wall to initiate intracellular infection [55, 56].

Given that nitrogen is the major nutrient limitation influencing insects exploiting living plant material [57], a more efficient plant-cell-wall-degrading phenotype is predicted to mediate specialization on host plants increasingly divergent in tissue form, composition, and complexity by facilitating access to cytosolic content rich in these compounds [4, 7, 9]. This is evident in the convergent evolution of phytophagy across the Coleoptera, where the independent integration of gene sets encoding for plant-cell-wall-degrading enzymes coincided with the rapid radiation of beetles to exploit

angiosperms during the Jurassic [4]. An analogous example for how the integration of novel digestive enzymes can influence ecological range, the evolution of entomopathogenicity in the fungus *Metarhizium* is tied to the horizontal acquisition of a core set of enzymes that facilitate the breakdown of insect cuticular components [58, 59]. Through the complementary activities of horizontally acquired lipid carriers and proteases, *Metarhizium* is able to penetrate the insect exoskeleton by efficiently digesting epicuticular lipids and procuticular proteins [58, 59]. In characterizing the impact of horizontal gene transfer on the pathogenic breadth of *Metarhizium*, generalist species were found to share several genes, including an endoprotease, which lacked homologs in specialist endophytes. Strikingly, experimental transfer of these genes through genome transformation resulted in *Metarhizium* specialists expanding their host range [59].

Coinciding with an extended range of universal plant cell wall polysaccharides that a folivore can metabolize, cassidines whose symbionts additionally provision rhamnogalacturonan lyase appear to have radiated onto host plants spanning a wider phylogenetic breadth (Table 1; Figure 6). Although *Stammera*




Figure 5. α -Glucuronidase Does Not Upgrade the Xylanolytic Capacity of Cassidine Leaf Beetles

TLC illustrating the breakdown products of xylanase activity assays against xylotriase linked to glucuronic acid using gut contents from *Cassida rubiginosa* and *Chelymorpha alternans*. Xyl, mono-xylose; Xyl-2, di-xylose; Xyl-3, tri-xylose. Dashed box highlights the absence of GlcA buildup across both treatments.

***Stammera* digestive enzymes**

Polygalacturonase (GH 28)

α -glucuronidase (GH 67)

Rhamnogalacturonan lyase (PL 4)

	GC + S	GC	S	GC + S	GC	S	Std	Std	Std
<i>Cassida rubiginosa</i>									
<i>Chelymorpha alternans</i>									

annotated to only express polygalacturonase are restricted to associating with cassidines feeding on plants within the lamiid eudicot clade, beetles symbiotically supplemented with both polygalacturonase and rhamnogalacturonan lyase evolved to specialize on angiosperms across three distinct eudicot clades (lamiids, campanulids, and fabids; Figure 6), which are separated by ~125 Ma of evolution [60].

Radiations to exploit novel ecological niches are rarely a consequence of just a single adaptation but rather an amalgam of several complementary traits, where the evolution of one may facilitate the rise of others that, collectively, drive the origin of novel adaptive forms [61]. Among beetles, the ability to exploit and coevolve with an increasing diversity of angiosperms is attributed to the development of grinding mandibular structures to fragment plant tissues [62], coupled with the acquisition and expansion of enzymatic classes to process toxic secondary compounds and recalcitrant polysaccharides [4]. Consistent with the latter, an upgraded digestive capacity by *Stammera* appears to not only reflect a broader phylogenetic host plant coverage but also an expanded geographical distribution that extends beyond the tropical origins of the Cassidinae [17]. Beetles symbiotically endowed with rhamnogalacturonan lyase and polygalacturonase exploit host plants adapted for temperate climates, such as thistles [23–25], mints [22], and Maleae trees [21],

in contrast to the tropical or subtropical distributions of lamiid host plants for cassidines relying solely on polygalacturonase [18–20, 27]. This biogeographical pattern is notable, given that land plants enrich for RG-I in response to drier growth conditions [63, 64]. Because it serves as an important anti-desiccant underlying plant cells' response to turgor changes, plants upregulate the synthesis of RG-I and its arabinan side chains during episodes of osmotic stress [65–69]. Where rainfall and humidity are key determinants of plant adaptation and distribution [70], rhamnogalacturonan lyase may allow beetles to more efficiently exploit temperate plants responding to sharper gradients of osmotic stress than those encountered in the tropics. In outlining an ecological context for the possible adaptive role of rhamnogalacturonan lyase in cassidines, we do not exclude the likelihood that complementary preadaptations may have also contributed to their expanded host plant range.

Following at least two independent horizontal gene transfer events [4], rhamnogalacturonan lyase is endogenously maintained by a subset of beetles belonging to the Buprestidae and Curculionidae families [4, 71–73]. Acquired horizontally from Actinobacteria (Buprestidae) and Proteobacteria (Curculionidae), the secondary pectinase is encoded by beetles specializing on host plants adapted for temperate or arid biomes [4], including pine [74], ash [75], date [76, 77], and citrus trees [78]. In parallel, a symbiosis with *Stammera* may have allowed certain cassidines to do the same. A similar interaction was recently described between cochineal scale insects and their β -proteobacterial symbiont, *Dactylopiibacterium* [79]. By encoding and expressing rhamnogalacturonan lyase and other pectinases, *Dactylopiibacterium* appears to be involved in the digestive capacity of cochineals subsisting exclusively on desert cactuses [79]. Thus, whether endogenously maintained or symbiotically supplemented, the secondary pectinase may be critical for insect herbivores overcoming host plants adapted for drier climates.

We note that our current study of this pectinolytic symbiosis does not extend to monocot-feeding, leaf-mining beetles. Previously a separate subfamily under the Hispinae, this group was recently reclassified as part of the Cassidinae [17]. Microbial




Figure 6. Cassidines Endowed with *Stammera*'s Polygalacturonase and Rhamnogalacturonan Lyase Have Radiated onto Host Plants Spanning a Wide Phylogenetic Range

Representative enzymatic combinations encoded by *Stammera* and the corresponding host plant range of cassidines. Eudicot phylogeny is adapted from Chase et al. [26].

profiling of these monocot-feeding taxa revealed the absence of *Stammera* and host symbiotic organs in *Cephaloleia* species [80], as well as in the palm pest *Octodonta nipae* [81]. This indicates that the symbiosis with *Stammera* is not shared by all members of the expanded subfamily. Toward understanding when *Stammera* was initially acquired, a broader sampling of monocot-feeding taxa is necessary and may offer new insights into the evolutionary transition and adaptation to eudicots within the Cassidinae.

The obligate symbioses insects form with micro-organisms serve as some of the most striking examples of beneficial partnerships in nature [82]. Endowing many key adaptations, these interactions are cited in spurring evolutionary innovation by expanding the availability of resources that may be evolutionarily exploited [83, 84]. Offsetting the loss of endogenous pectinases in cassidines, we demonstrate how polygalacturonase is a defining and consistent feature of the symbiosis with *Stammera*. Contrasting a conserved catalytic capacity against HG, cassidines additionally supplemented with rhamnogalacturonan lyase display a more-proficient digestive phenotype against RG-I. As the pectic heteropolymer is dynamically regulated by eudicots to contend with abiotic stress throughout development, we offer a framework for how a symbiont may modulate plant-herbivore interactions by expanding the range of universal plant cell wall polymers that its host can readily overcome.

STAR★METHODS

Detailed methods are provided in the online version of this paper and include the following:

- KEY RESOURCES TABLE
- RESOURCE AVAILABILITY
 - Lead Contact
 - Materials Availability
 - Data and Code Availability
- EXPERIMENTAL MODEL AND SUBJECT DETAILS

● METHOD DETAILS

- Symbiont genome sequencing, assembly, and annotation
- Phylogenetic reconstruction
- Transcriptome sequencing, assembly, and annotation
- Enzymatic assays

SUPPLEMENTAL INFORMATION

Supplemental Information can be found online at <https://doi.org/10.1016/j.cub.2020.05.043>.

ACKNOWLEDGMENTS

We thank the Department of Bioorganic Chemistry at the Max Planck Institute for Chemical Ecology for contributing live *Chelymorpha alternans* beetles, Tobin Hammer for scientific matchmaking, and Ryuichi Koga for insightful discussions. Financial support from the Alexander von Humboldt Foundation, Smithsonian Institution, National Science Foundation, Max Planck Society, and the Wissenschaftskolleg zu Berlin is gratefully acknowledged. Specimen images from Udo Schmidt (<https://www.kaefer-der-welt.de>) are greatly appreciated, as are illustrations from BioRender (<https://biorender.com>).

AUTHOR CONTRIBUTIONS

H.S. and N.M.G. conceived of the study. H.S., N.M.G., D.W., R.K., Y.P., T.F., and A.B. designed the experiments. H.S., A.B., K.F., and M.M. performed genome sequencing, assembly, and analysis. R.K. and Y.P. carried out transcriptome sequencing, assembly, and analysis. R.K. and Y.P. performed the enzymatic assays. M.C. contributed insect specimens for initial assays. H.S. wrote the manuscript. All authors edited and commented on the paper.

DECLARATION OF INTERESTS

The authors declare no competing interests.

Received: January 19, 2020

Revised: April 5, 2020

Accepted: May 12, 2020

Published: June 4, 2020

REFERENCES

- Mitter, C., Farrell, B., and Wiegmann, B. (1988). The phylogenetic study of adaptive zones: has phytophagy promoted insect diversification? *Am. Nat.* **132**, 107–128.
- Farrell, B.D. (1998). "Inordinate fondness" explained: why are there so many beetles? *Science* **281**, 555–559.
- Barraclough, T.G., Barclay, M.V.L., and Vogler, A.P. (1998). Species richness: does flower power explain beetle-mania? *Curr. Biol.* **8**, R843–R845.
- McKenna, D.D., Shin, S., Ahrens, D., Balke, M., Beza-Beza, C., Clarke, D.J., Donath, A., Escalona, H.E., Friedrich, F., Letsch, H., et al. (2019). The evolution and genomic basis of beetle diversity. *Proc. Natl. Acad. Sci. USA* **116**, 24729–24737.
- Busch, A., Danchin, E.G.J., and Pauchet, Y. (2019). Functional diversification of horizontally acquired glycoside hydrolase family 45 (GH45) proteins in Phytophaga beetles. *BMC Evol. Biol.* **19**, 100.
- Acuña, R., Padilla, B.E., Flórez-Ramos, C.P., Rubio, J.D., Herrera, J.C., Benavides, P., Lee, S.-J., Yeats, T.H., Egan, A.N., Doyle, J.J., and Rose, J.K. (2012). Adaptive horizontal transfer of a bacterial gene to an invasive insect pest of coffee. *Proc. Natl. Acad. Sci. USA* **109**, 4197–4202.
- Kirsch, R., Gramzow, L., Theißen, G., Siegfried, B.D., French-Constant, R.H., Heckel, D.G., and Pauchet, Y. (2014). Horizontal gene transfer and functional diversification of plant cell wall degrading polygalacturonases: Key events in the evolution of herbivory in beetles. *Insect Biochem. Mol. Biol.* **52**, 33–50.
- Wybouw, N., Pauchet, Y., Heckel, D.G., and Van Leeuwen, T. (2016). Horizontal gene transfer contributes to the evolution of arthropod herbivory. *Genome Biol. Evol.* **8**, 1785–1801.
- Calderón-Cortés, N., Quesada, M., Watanabe, H., Cano-Camacho, H., and Oyama, K. (2012). Endogenous plant cell wall digestion: a key mechanism in insect evolution. *Annu. Rev. Ecol. Evol. Syst.* **43**, 45–71.
- Salem, H., Bauer, E., Kirsch, R., Berasategui, A., Cripps, M., Weiss, B., Koga, R., Fukumori, K., Vogel, H., Fukatsu, T., and Kaltenpoth, M. (2017). Drastic genome reduction in an herbivore's pectinolytic symbiont. *Cell* **171**, 1520–1531.e13.
- Bauer, E., Kaltenpoth, M., and Salem, H. (2020). Minimal fermentative metabolism fuels extracellular symbiont in a leaf beetle. *ISME J.* **14**, 866–870.
- Burton, R.A., Gidley, M.J., and Fincher, G.B. (2010). Heterogeneity in the chemistry, structure and function of plant cell walls. *Nat. Chem. Biol.* **6**, 724–732.
- Mohnen, D. (2008). Pectin structure and biosynthesis. *Curr. Opin. Plant Biol.* **11**, 266–277.
- Ochoa-Jiménez, V.-A., Berumen-Varela, G., Fernández-Valle, R., Ernesto, M., and Hernández, T. (2018). Rhamnogalacturonan lyase: a pectin modification enzyme of higher plants. *Emir. J. Food Agric.* **30**, 910–917.
- De Lorenzo, G., and Ferrari, S. (2002). Polygalacturonase-inhibiting proteins in defense against phytopathogenic fungi. *Curr. Opin. Plant Biol.* **5**, 295–299.
- Blackman, L.M., Cullerle, D.P., Torreña, P., Taylor, J., and Hardham, A.R. (2015). RNA-seq analysis of the expression of genes encoding cell wall degrading enzymes during infection of lupin (*Lupinus angustifolius*) by *Phytophthora parasitica*. *PLoS ONE* **10**, e0136899.
- Chaboo, C.S. (2007). Biology and phylogeny of the Cassidinae Gyllenhal sensu lato (tortoise and leaf-mining beetles) (Coleoptera, Chrysomelidae). *Bull. Am. Mus. Nat. Hist.* **305**, 1–250.
- Windsor, D.M., Riley, E.G., and Stockwell, H.P. (1992). An introduction to the biology and systematics of Panamanian tortoise beetles (Coleoptera: Chrysomelidae: Cassidinae). In *Insects of Panama and Mesoamerica: Selected Studies*, D. Quintero, and A. Aiello, eds. (Oxford University), pp. 372–391.
- Morrison, C.R., and Windsor, D.M. (2018). The life history of *Chelymophra alternans* (Coleoptera: Chrysomelidae: Cassidinae) in Panamá. *Ann. Entomol. Soc. Am.* **111**, 31–41.
- Windsor, D.M. (1987). Natural history of a subsocial tortoise beetle, *Acromis sparsa* Boheman (Chrysomelidae, Cassidinae) in Panama. *Psyche J. Entomol.* **94**, 19861.
- Kimoto, S., and Takizawa, H. (1994). Leaf Beetles (Chrysomelidae) of Japan (Tokai University).
- Marshall, S.A., and Paiero, S.M. (2016). *Cassida viridis* (Coleoptera: Chrysomelidae), a Palaearctic leaf beetle newly recorded from North America. *Can. Entomol.* **148**, 140–142.
- Fujiyama, N., Togashi, K., Kikuta, S., and Katakura, H. (2011). Distribution and host specificity of the thistle-feeding tortoise beetle *Cassida vibex* (Coleoptera : Chrysomelidae) in southwestern Hokkaido, northern Japan. *Entomol. Sci.* **14**, 271–277.
- Cripps, M.G., Jackman, S.D., Roquet, C., van Koten, C., Rostás, M., Bourdôt, G.W., et al. (2016). Evolution of Specialization of *Cassida rubiginosa* on *Cirsium arvense* (Compositae, Cardueae). *Front. Plant Sci.* **7**, 1261.
- Ward, R.H., and Pienkowski, R.L. (1978). Biology of *Cassida rubiginosa* a thistle-feeding shield beetle. *Ann. Entomol. Soc. Am.* **71**, 585–591.
- The Angiosperm Phylogeny Group. (2016). An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG IV. *Bot. J. Linn. Soc.* **181**, 1–20.
- Barrows, E.M. (1979). Life cycles, mating, and color change in tortoise beetles (Coleoptera:Chrysomelidae:Cassidinae). *Coleopt. Bull.* **33**, 9–16.
- Conow, C., Fielder, D., Ovidia, Y., and Libeskind-Hadas, R. (2010). Jane: a new tool for the cophylogeny reconstruction problem. *Algorithms Mol. Biol.* **5**, 16.
- Salem, H., Florez, L., Gerardo, N., and Kaltenpoth, M. (2015). An out-of-body experience: the extracellular dimension for the transmission of mutualistic bacteria in insects. *Proc. Biol. Sci.* **282**, 20142957.
- Kikuchi, Y., Hosokawa, T., Nikoh, N., Meng, X.-Y., Kamagata, Y., and Fukatsu, T. (2009). Host-symbiont co-speciation and reductive genome evolution in gut symbiotic bacteria of acanthosomatid stinkbugs. *BMC Biol.* **7**, 2.
- Hosokawa, T., Kikuchi, Y., Nikoh, N., Shimada, M., and Fukatsu, T. (2006). Strict host-symbiont cospeciation and reductive genome evolution in insect gut bacteria. *PLoS Biol.* **4**, e337.
- Kaiwa, N., Hosokawa, T., Nikoh, N., Tanahashi, M., Moriyama, M., Meng, X.-Y., Maeda, T., Yamaguchi, K., Shigenobu, S., Ito, M., and Fukatsu, T. (2014). Symbiont-supplemented maternal investment underpinning host's ecological adaptation. *Curr. Biol.* **24**, 2465–2470.
- McCutcheon, J.P., and Moran, N.A. (2011). Extreme genome reduction in symbiotic bacteria. *Nat. Rev. Microbiol.* **10**, 13–26.
- Poliakov, A., Russell, C.W., Ponnala, L., Hoops, H.J., Sun, Q., Douglas, A.E., and van Wijk, K.J. (2011). Large-scale label-free quantitative proteomics of the pea aphid-*Buchnera* symbiosis. *Mol. Cell. Proteomics* **10**, M110.007039.
- McCutcheon, J.P., McDonald, B.R., and Moran, N.A. (2009). Convergent evolution of metabolic roles in bacterial co-symbionts of insects. *Proc. Natl. Acad. Sci. USA* **106**, 15394–15399.
- Haines, L.R., Haddow, J.D., Aksoy, S., Gooding, R.H., and Pearson, T.W. (2002). The major protein in the midgut of teneral *Glossina morsitans morsitans* is a molecular chaperone from the endosymbiotic bacterium *Wigglesworthia glossinidia*. *Insect Biochem. Mol. Biol.* **32**, 1429–1438.
- Bennett, G.M., and Chong, R.A. (2017). Genome-wide transcriptional dynamics in the companion bacterial symbionts of the glassy-winged sharpshooter (Cicadellidae: Homalodisca vitripennis) reveal differential gene expression in bacteria occupying multiple host organs. *G3* **7**, 3073–3082.
- Pelley, J.W. (2007). 7 - Citric acid cycle, electron transport chain, and oxidative phosphorylation. In *Elsevier's Integrated Biochemistry*, J.W. Pelley, ed. (Mosby), pp. 55–63.
- Tamames, J., Gil, R., Latorre, A., Peretó, J., Silva, F.J., and Moya, A. (2007). The frontier between cell and organelle: genome analysis of Candidatus *Carsonella ruddii*. *BMC Evol. Biol.* **7**, 181.

40. Bennett, G.M., and Moran, N.A. (2015). Heritable symbiosis: the advantages and perils of an evolutionary rabbit hole. *Proc. Natl. Acad. Sci. USA* *112*, 10169–10176.
41. Gil, R., Sabater-Muñoz, B., Latorre, A., Silva, F.J., and Moya, A. (2002). Extreme genome reduction in *Buchnera* spp.: toward the minimal genome needed for symbiotic life. *Proc. Natl. Acad. Sci. USA* *99*, 4454–4458.
42. Van Leuven, J.T., Meister, R.C., Simon, C., and McCutcheon, J.P. (2014). Sympatric speciation in a bacterial endosymbiont results in two genomes with the functionality of one. *Cell* *158*, 1270–1280.
43. Azadi, P., O'Neill, M.A., Bergmann, C., Darvill, A.G., and Albersheim, P. (1995). The backbone of the pectic polysaccharide rhamnogalacturonan I is cleaved by an endohydrolase and an endolyase. *Glycobiology* *5*, 783–789.
44. Biely, P., de Vries, R.P., Vrsanská, M., and Visser, J. (2000). Inverting character of α -glucuronidase A from *Aspergillus tubingensis*. *Biochim. Biophys. Acta* *1474*, 360–364.
45. McKenna, D.D., Scully, E.D., Pauchet, Y., Hoover, K., Kirsch, R., Geib, S.M., et al. (2016). Genome of the Asian longhorned beetle (*Anoplophora glabripennis*), a globally significant invasive species, reveals key functional and evolutionary innovations at the beetle-plant interface. *Genome Biol.* *17*, 227.
46. Busch, A., Kunert, G., Wielsch, N., and Pauchet, Y. (2018). Cellulose degradation in *Gastrophysa viridula* (Coleoptera: Chrysomelidae): functional characterization of two CAZymes belonging to glycoside hydrolase family 45 reveals a novel enzymatic activity. *Insect Mol. Biol.* *27*, 633–650.
47. Padilla-Hurtado, B., Flórez-Ramos, C., Aguilera-Gálvez, C., Medina-Olaya, J., Ramírez-Sanjuan, A., Rubio-Gómez, J., et al. (2012). Cloning and expression of an endo-1,4- β -xylanase from the coffee berry borer, *Hypothenemus hampei*. *BMC Res. Notes* *5*, 23.
48. Schoville, S.D., Chen, Y.H., Andersson, M.N., Benoit, J.B., Bhandari, A., Bowsher, J.H., et al. (2018). A model species for agricultural pest genomics: the genome of the Colorado potato beetle, *Leptinotarsa decemlineata* (Coleoptera: Chrysomelidae). *Sci. Rep.* *8*, 1931.
49. Kikuchi, T., Jones, J.T., Aikawa, T., Kosaka, H., and Ogura, N. (2004). A family of glycosyl hydrolase family 45 cellulases from the pine wood nematode *Bursaphelenchus xylophilus*. *FEBS Lett.* *572*, 201–205.
50. Rahman, M.M., Inoue, A., and Ojima, T. (2014). Characterization of a GHF45 cellulase, AKEG21, from the common sea hare *Aplysia kurodai*. *Front Chem.* *2*, 60.
51. Szydlowski, L., Boschetti, C., Crisp, A., Barbosa, E.G.G., and Tunnacliffe, A. (2015). Multiple horizontally acquired genes from fungal and prokaryotic donors encode cellulolytic enzymes in the bdelloid rotifer *Adineta ricciae*. *Gene* *566*, 125–137.
52. Yapo, B.M. (2011). Rhamnogalacturonan-I: a structurally puzzling and functionally versatile polysaccharide from plant cell walls and mucilages. *Polym. Rev.* *51*, 391–413.
53. Vincken, J.-P., Schols, H.A., Oomen, R.J.F.J., McCann, M.C., Ulvskov, P., Voragen, A.G.J., and Visser, R.G.F. (2003). If homogalacturonan were a side chain of rhamnogalacturonan I. Implications for cell wall architecture. *Plant Physiol.* *132*, 1781–1789.
54. Silva, I.R., Jers, C., Meyer, A.S., and Mikkelsen, J.D. (2016). Rhamnogalacturonan I modifying enzymes: an update. *N. Biotechnol.* *33*, 41–54.
55. Laatu, M., and Condemine, G. (2003). Rhamnogalacturonate lyase RhiE is secreted by the out system in *Erwinia chrysanthemi*. *J. Bacteriol.* *185*, 1642–1649.
56. Silva, I.R., Larsen, D.M., Meyer, A.S., and Mikkelsen, J.D. (2011). Identification, expression, and characterization of a novel bacterial RGI lyase enzyme for the production of bio-functional fibers. *Enzyme Microb. Technol.* *49*, 160–166.
57. Mattson, W.J., Jr. (1980). Herbivory in relation to plant nitrogen content. *Annu. Rev. Ecol. Syst.* *11*, 119–161.
58. Zhao, H., Xu, C., Lu, H.-L., Chen, X., St Leger, R.J., and Fang, W. (2014). Host-to-pathogen gene transfer facilitated infection of insects by a pathogenic fungus. *PLoS Pathog.* *10*, e1004009.
59. Zhang, Q., Chen, X., Xu, C., Zhao, H., Zhang, X., Zeng, G., Qian, Y., Liu, R., Guo, N., Mi, W., et al. (2019). Horizontal gene transfer allowed the emergence of broad host range entomopathogens. *Proc. Natl. Acad. Sci. USA* *116*, 7982–7989.
60. Zeng, L., Zhang, N., Zhang, Q., Endress, P.K., Huang, J., and Ma, H. (2017). Resolution of deep eudicot phylogeny and their temporal diversification using nuclear genes from transcriptomic and genomic datasets. *New Phytol.* *214*, 1338–1354.
61. Stroud, J.T., and Losos, J.B. (2016). Ecological opportunity and adaptive radiation. *Annu. Rev. Ecol. Syst.* *47*, 507–532.
62. Acorn, J.H., and Ball, G.E. (1991). The mandibles of some adult ground beetles: structure, function, and the evolution of herbivory (Coleoptera: Carabidae). *Can. J. Zool.* *69*, 638–650.
63. Tenhaken, R. (2015). Cell wall remodeling under abiotic stress. *Front. Plant Sci.* *5*, 771.
64. Le Gall, H., Philippe, F., Domon, J.-M., Gillet, F., Pelloux, J., and Rayon, C. (2015). Cell wall metabolism in response to abiotic stress. *Plants (Basel)* *4*, 112–166.
65. Leucci, M.R., Lenucci, M.S., Piro, G., and Dalessandro, G. (2008). Water stress and cell wall polysaccharides in the apical root zone of wheat cultivars varying in drought tolerance. *J. Plant Physiol.* *165*, 1168–1180.
66. Moore, J.P., Nguema-Ona, E., Chevalier, L., Lindsey, G.G., Brandt, W.F., Lerouge, P., et al. (2006). Response of the leaf cell wall to desiccation in the resurrection plant *Myrothamnus flabellifolius*. *Plant Physiol.* *141*, 651–662.
67. Moore, J.P., Farrant, J.M., and Driouch, A. (2008). A role for pectin-associated arabinans in maintaining the flexibility of the plant cell wall during water deficit stress. *Plant Signal. Behav.* *3*, 102–104.
68. Jones, L., Milne, J.L., Ashford, D., and McQueen-Mason, S.J. (2003). Cell wall arabinan is essential for guard cell function. *Proc. Natl. Acad. Sci. USA* *100*, 11783–11788.
69. Piro, G., Leucci, M.R., Waldron, K., and Dalessandro, G. (2003). Exposure to water stress causes changes in the biosynthesis of cell wall polysaccharides in roots of wheat cultivars varying in drought tolerance. *Plant Sci.* *165*, 559–569.
70. Holdridge, L.R. (1967). *Life Zone Ecology* (Tropical Science Center).
71. Keeling, C.I., Yuen, M.M.S., Liao, N.Y., Docking, T.R., Chan, S.K., Taylor, G.A., et al. (2013). Draft genome of the mountain pine beetle, *Dendroctonus ponderosae* Hopkins, a major forest pest. *Genome Biol.* *14*, R27.
72. Vega, F.E., Brown, S.M., Chen, H., Shen, E., Nair, M.B., Ceja-Navarro, J.A., et al. (2015). Draft genome of the most devastating insect pest of coffee worldwide: the coffee berry borer, *Hypothenemus hampei*. *Sci. Rep.* *5*, 12525.
73. Wang, L., Zhang, X.-W., Pan, L.-L., Liu, W.-F., Wang, D.-P., Zhang, G.-Y., et al. (2013). A large-scale gene discovery for the red palm weevil *Rhynchophorus ferrugineus* (Coleoptera: Curculionidae). *Insect Sci.* *20*, 689–702.
74. Erbilgin, N., Ma, C., Whitehouse, C., Shan, B., Najjar, A., and Evenden, M. (2014). Chemical similarity between historical and novel host plants promotes range and host expansion of the mountain pine beetle in a naïve host ecosystem. *New Phytol.* *201*, 940–950.
75. Anulewicz, A.C., McCullough, D.G., Cappaert, D.L., and Poland, T.M. (2008). Host range of the emerald ash borer (*Agrilus planipennis* Fairmaire) (Coleoptera: Buprestidae) in North America: results of multiple-choice field experiments. *Environ. Entomol.* *37*, 230–241.
76. Kontodimas, D.C., Milonas, P.G., Vassiliou, V., Thymakis, N., and Economou, D. (2006). The occurrence of *Rhynchophorus ferrugineus* in Greece and Cyprus and the risk against the native greek palm tree Phoenix theophrasti. *Entomol. Hell.* *16*, 11–15.
77. Soroker, V., Blumberg, D., Haberman, A., Hamburger-Rishard, M., Reneh, S., Talebaev, S., Anshelevich, L., and Harari, A.R. (2005). Current status of

- red palm weevil infestation in date palm plantations in Israel. *Phytoparasitica* 33, 97–106.
78. CISR (2019). Diaprepes root weevil. https://cizr.ucr.edu/diaprepes_root_weevil.html.
79. Bustamante-Brito, R., Vera-Ponce de León, A., Rosenblueth, M., Martínez-Romero, J.C., and Martínez-Romero, E. (2019). Metatranscriptomic analysis of the bacterial symbiont *Dactylopiibacterium carminicum* from the carmine cochineal *Dactylopius coccus* (Hemiptera: Coccoidea: Dactylopiidae). *Life* 9, 4.
80. Blankenchip, C.L., Michels, D.E., Braker, H.E., and Goffredi, S.K. (2018). Diet breadth and exploitation of exotic plants shift the core microbiome of *Cephaloleia*, a group of tropical herbivorous beetles. *PeerJ* 6, e4793.
81. Ali, H., Muhammad, A., Sanda, N.B., Huang, Y., and Hou, Y. (2019). Pyrosequencing uncovers a shift in bacterial communities across life stages of *Octodonta nipae* (Coleoptera: Chrysomelidae). *Front. Microbiol.* 10, 466.
82. Douglas, A.E. (2015). Multiorganismal insects: diversity and function of resident microorganisms. *Annu. Rev. Entomol.* 60, 17–34.
83. Chomicki, G., Weber, M., Antonelli, A., Bascompte, J., and Kiers, E.T. (2019). The impact of mutualisms on species richness. *Trends Ecol. Evol.* 34, 698–711.
84. Sudakran, S., Kost, C., and Kaltenpoth, M. (2017). Symbiont acquisition and replacement as a source of ecological innovation. *Trends Microbiol.* 25, 375–390.
85. Bolger, A.M., Lohse, M., and Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30, 2114–2120.
86. Miller, M.A., Pfeiffer, W., and Schwartz, T. (2010). Creating the CIPRES Science Gateway for inference of large phylogenetic trees. In 2010 Gateway Computing Environments Workshop (GCE) (IEEE), pp. 1–8.
87. Langmead, B., and Salzberg, S.L. (2012). Fast gapped-read alignment with Bowtie 2. *Nat. Methods* 9, 357–359.
88. Bankevich, A., Nurk, S., Antipov, D., Gurevich, A.A., Dvorkin, M., Kulikov, A.S., Lesin, V.M., Nikolenko, S.I., Pham, S., Pribelski, A.D., et al. (2012). SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J. Comput. Biol.* 19, 455–477.
89. Boetzer, M., and Pirovano, W. (2012). Toward almost closed genomes with GapFiller. *Genome Biol.* 13, R56.
90. Aziz, R.K., Bartels, D., Best, A.A., DeJongh, M., Disz, T., Edwards, R.A., Formisano, K., Gerdes, S., Glass, E.M., Kubal, M., et al. (2008). The RAST Server: rapid annotations using subsystems technology. *BMC Genomics* 9, 75.
91. Edgar, R.C. (2004). MUSCLE: a multiple sequence alignment method with reduced time and space complexity. *BMC Bioinformatics* 5, 113.
92. Lanfear, R., Frandsen, P.B., Wright, A.M., Senfeld, T., and Calcott, B. (2017). PartitionFinder 2: new methods for selecting partitioned models of evolution for molecular and morphological phylogenetic analyses. *Mol. Biol. Evol.* 34, 772–773.
93. Arkin, A.P., Cottingham, R.W., Henry, C.S., Harris, N.L., Stevens, R.L., Maslov, S., Dehal, P., Ware, D., Perez, F., Canon, S., et al. (2018). KBase: The United States Department of Energy Systems Biology Knowledgebase. *Nat. Biotechnol.* 36, 566–569.
94. Bazinet, A.L., Zwickl, D.J., and Cummings, M.P. (2014). A gateway for phylogenetic analysis powered by grid computing featuring GARLI 2.0. *Syst. Biol.* 63, 812–818.
95. Peters, R.S., Krogmann, L., Mayer, C., Donath, A., Gunkel, S., Meusemann, K., Kozlov, A., Podsiadlowski, L., Petersen, M., Lanfear, R., et al. (2017). Evolutionary history of the hymenoptera. *Curr. Biol.* 27, 1013–1018.

STAR★METHODS

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Bacterial and Virus Strains		
<i>Candidatus</i> <i>Stammera capleta</i>	This study and [10]	NCBI Taxonomy ID: 2608262
Biological Samples		
<i>Cassida versicolor</i> collected in Tsukuba, Japan	This paper	N/A
<i>Cassida viridis</i> collected in Jena, Germany	This paper	N/A
<i>Cassida vibex</i> collected in Jena, Germany	This paper	N/A
<i>Cassida rubiginosa</i> collected in Jena, Germany	This paper	N/A
<i>Parachirida semiannulata</i> collected in Gamboa, Panama	This paper	N/A
<i>Ischnocodia annulus</i> collected in Gamboa, Panama	This paper	N/A
<i>Stolas discoides</i> collected in Gamboa, Panama	This paper	N/A
<i>Chelymorpha alternans</i> collected in Gamboa, Panama	This paper	N/A
<i>Acromis sparsa</i> collected in Gamboa, Panama	This paper	N/A
<i>Cistudinella foveolata</i> collected in Gamboa, Panama	This paper	N/A
<i>Discomorpha panamensis</i> collected in Gamboa, Panama	This paper	N/A
<i>Agroiconota porpinqua</i> collected in Gamboa, Panama	This paper	N/A
<i>Charidotella sexpunctata</i> collected in Gamboa, Panama	This paper	N/A
Chemicals, Peptides, and Recombinant Proteins		
RNAlater	Sigma-Aldrich	Cat#R0901
TURBO DNase	Thermo Fisher Scientific	Cat#AM2238
TLC plates Silica gel 60	Merck	Cat#116835
rhamnogalacturonan I	Megazyme	Cat#P-RHAM1
Polygalacturonic acid	Megazyme	Cat#P-PGACT
Galacturonic acid	Santa Cruz	Cat#91510-62-2
di-galacturonic acid	Sigma-Aldrich	Cat#D4288
Tri-galacturonic acid	Sigma-Aldrich	Cat#T7407
rhamnose	Sigma-Aldrich	Cat#83650
Xylose with glucuronic acid decoration	Megazyme	Cat# O-XUX
Xylose	Sigma-Aldrich	Cat#47253
di-xylose	Megazyme	Cat#O-XBI
Tri-xylose	Megazyme	Cat#O-XTR
Glucuronic acid	Sigma-Aldrich	Cat#G5269
Critical Commercial Assays		
innuPrep DNA/RNA Mini kit	Analytik Jena	Cat# 845-KS-2080050
RNeasy MinElute Clean up Kit	Quiagen	Cat# 74204
RNA 6000 Nano LabChip kit	Agilent	Cat#5067-1511
QIAGEN RNA Extraction kit	QIAGEN	Cat# 74104
QIAGEN RNeasy Blood & Tissue Kit	QIAGEN	Cat# 69506
Taq DNA polymerase	VWR	Cat# 89167-762
Deposited Data		
Transcriptome sequencing Bioproject ID	This paper	NCBI:PRJNA561700
Genome sequencing Bioproject ID	This paper	NCBI:PRJNA561424

(Continued on next page)

Continued

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Experimental Models: Organisms/Strains		
<i>Cassida viridis</i> collected in Jena, Germany	This paper	N/A
<i>Cassida vibex</i> collected in Jena, Germany	This paper	N/A
<i>Cassida rubiginosa</i> collected in Jena, Germany	This paper	N/A
<i>Chelymormpha alternans</i> collected in Gamboa, Panama	This paper	N/A
Software and Algorithms		
CLC Genomics Workbench v11.0	Quiagen	Cat#832001
Trimmomatic v0.36	[85]	http://www.usadellab.org/cms/index.php?page=trimmomatic
Trinity v2.5.1	[86]	https://github.com/trinityrnaseq/trinityrnaseq/releases/tag/Trinity-v2.5.1
Bowtie2	[87]	http://bowtie-bio.sourceforge.net/bowtie2/index.shtml
Spades	[88]	http://bioinf.spbau.ru/spades
GapFiller	[89]	https://www.baseclear.com/genomics/bioinformatics/basetools/gapfiller
RAST	[90]	http://rast.nmpdr.org
MUSCLE	[91]	http://phylogeny.lirmm.fr/phylo.cgi/one_task.cgi?task_type=muscle
PartitionFinder 2	[92]	http://www.robertlanfear.com/partitionfinder/
Jane 4	[28]	https://www.cs.hmc.edu/~hadas/jane/

RESOURCE AVAILABILITY**Lead Contact**

Further information and requests for resources (e.g., available reference specimens), peptic substrates and commercial reagents should be directed to and will be fulfilled by the Lead Contact, Hassan Salem (hassan.salem@tuebingen.mpg.de).

Materials Availability

This study did not generate new unique reagents.

Data and Code Availability

Genomic and transcriptomic sequencing data generated in this study are available at the National Center for Biotechnology Information under BioProjects PRJNA561424 and PRJNA561700, respectively.

BioProjects can be accessed at:

PRJNA561424: <https://www.ncbi.nlm.nih.gov/bioproject/PRJNA561424>

PRJNA561700: <https://www.ncbi.nlm.nih.gov/bioproject/PRJNA561700>

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Adult Cassidinae specimens were collected in Germany, Panama, the United States of America, Japan and New Zealand between 2016 and 2018. The insects were submerged in 70% ethanol until they were processed ahead of DNA extraction. Ten adults were included per sample. Representative specimens were identified based on morphological characters and their identity was confirmed by partial sequencing of the mitochondrial 16S rRNA gene, as well as the nuclear 18S and 28S rRNA genes. For enzymatic assays, *Cassida rubiginosa*, *Cassida vibex* and *Cassida viridis* were collected from leaves of *Cirsium arvense*, *Cirsium oleraceum* and *Mentha arvensis* in Jena, Germany; whereas *Chelymormpha alternans* are continuously reared at the Max Planck Institute for Chemical Ecology on *Ipomoea batatas*.

METHOD DETAILS**Symbiont genome sequencing, assembly, and annotation**

Symbiotic organs were dissected from ten individuals of each of the 13 cassidine species, and DNA was purified using the QIAGEN DNeasy Blood & Tissue Kit (Hilden, Germany) according to the manufacturer's instructions. Sequencing was performed on a HiSeq 2500 Sequencing System from Illumina (<https://www.illumina.com/systems/sequencing-platforms/hiseq-2500.html>), utilizing the

paired-end 150 bp technology and at a depth of ~40 million reads. Adaptor sequences were trimmed with Trimmomatic v0.03 using the default parameters [85]. To filter beetle-, plant-, and *Wolbachia*-associated sequences, the quality-controlled reads were mapped using Bowtie2 [87] to the publicly available sequences of closely related species. Assembly was performed using Spades under default parameters [88]. This resulted in ~40,000 contigs per assembly. After assembly, the contigs were screened for GC content and taxonomic identity to Enterobacteriaceae via BLAST to filter out remaining contaminant sequences. Contigs were then further assembled into scaffolds using GapFiller [89]. Resulting genomic scaffolds were subsequently connected and circularized by PCR and Sanger sequencing where necessary. The final genome sequences were automatically annotated using RAST [90] (as implemented in KBase [93]) and according to the genetic code 4 (TGA encoding tryptophan) to computationally translate the predicted protein-coding genes (Data S1).

Phylogenetic reconstruction

The phylogenetic relationships among the Cassidinae were reconstructed using a mitochondrial (16S rRNA) and two nuclear genes (18S and 28S rRNA) for all host species. Toward inferring the relationships among the different *Stammera* strains, a set of reference alignments based on highly conserved Clusters of Orthologous Groups (COG) families was used to identify 30 protein-coding genes shared by all taxa. Host and symbiont genes were aligned using MUSCLE [91] ahead of concatenation. PartitionFinder 2 [92] was used to select the best partitioning scheme. Applying the greedy algorithm resulted in 3 and 28 partitions for host and symbiont alignments, respectively (Table S5). Maximum likelihood (ML) tree inference was performed using GARLI 2.0 [94] as implemented in Cypres Science Gateway [86]. The identified partitioning schemes were used in the ML analyses, and each subset was assigned the model suggested by PartitionFinder 2. For each analysis, branch support was estimated following 1,000 bootstrap replicates.

The phylogenetic reconciliation tool Jane 4 [28] was applied to describe the coevolutionary association between Cassidinae beetles and *Stammera*, using default cost settings. This analysis leverages five evolutionary scenarios to characterize host-symbiont interactions: cospeciation (joint speciation with the host lineage); duplication (both symbionts are kept in the same host); duplication and host switch (symbionts are duplicated and transferred from one host species to another); losses (loss of symbiont); and failure to diverge. Accounting for possible vector combinations, our optimal reconstructions reported a total vector cost of 8. The statistical significance of the total cost was tested using the null distribution of cost values based on 1,000 randomly generated trees. Following these randomizations, 100% had a total vector cost value higher than that of our predicted results, indicating that the obtained reconstructions were not randomly attributed.

Transcriptome sequencing, assembly, and annotation

Dissected beetle midguts from six representative cassidines were preserved in RNAlater. Preserved midguts were extracted, snap-frozen in liquid nitrogen and stored at -80°C . Total RNA was extracted from three beetles using the innuPrep DNA/RNA Mini kit (Analytik Jena, Jena, Germany) following the manufacturer's instructions. Genomic DNA contamination was removed by DNase treatment (TURBO DNase, Invitrogen, Carlsbad, CAL, USA) for 30 min at 37°C . Total RNA samples were further purified by using the RNeasy MinElute Clean up Kit (QIAGEN, Hilden, Germany) following the manufacturer's protocol, except for elution to 20 μL volume. The integrity and quality of the RNA samples were determined using the RNA 6000 Nano LabChip kit on an Agilent 2100 Bioanalyzer (both Agilent Technologies, Santa Clara, CAL, USA) according to the manufacturer's instructions.

RNA-Seq was performed at the Max Planck Genome Center (Cologne, Germany). There, poly(A)⁺ enriched RNA was fragmented to an average size of 300–350 nucleotides and a TruSeq compatible, directional library was prepared for each sample. Libraries were tagged using dual indexing and were multiplexed on the same sequencing lane. Sequencing was carried out on a HiSeq3000 sequencing platform (Illumina, CA, USA) using paired-end (2 × 150 bp) reads. Quality control measures, including the filtering of high-quality reads based on fastq file scores, the removal of reads containing primer/adaptor sequences and trimming of the read length, were carried out using CLC Genomics Workbench v11.0 (QIAGEN, Hilden, Germany). Multiple assemblies were performed for each sample whereby an increasing number of randomly selected read pairs were included per assembly (Table S4).

The RNA-Seq data for *Chelymorpha alternans* were generated in a similar way, except that the library was tagged using mono indexing and was multiplexed with other beetle-derived libraries on the same sequencing lane. Illumina-specific adapters, tags and low-quality bases were removed using Trimmomatic v0.36. The resulting clean reads were assembled using Trinity v2.5.1 using default parameters. We faced problems of cross-contamination of the sequencing data after de-multiplexing, which made subsequent analyses difficult. We cured these RNA-Seq data from cross-contamination using the protocol described by Peters et al. [95]. Briefly, we compared the *C. alternans* transcriptome assembly with all other assemblies sequenced in the same run using BLASTN. The coverage depth of BLASTN identified transcripts that shared nucleotide sequence identity of at least 98% over a length of at least 180 bp was then compared. If the relative coverage of two transcripts originating from two different assemblies differed >2-fold, the transcript with the lower relative coverage was assumed to be a contaminant and was removed from the corresponding assembly. Plant cell wall degrading enzymes (PCWDEs) were annotated in the assembled transcriptomes by BLAST-searches using representative PCWDE sequences from insects, nematodes, bacteria, and fungi, and as previously described in Kirsch et al. [7] and Busch et al. [46].

Enzymatic assays

Qualitative analysis of breakdown products was performed by thin layer chromatography (TLC) of 20 μL enzyme assays set up as follows: 14 μL of crude gut extract and pooled from *Cassida rubiginosa*, *Cassida vibex*, *Cassida viridis* or *Chelymorpha alternans*

(three beetles each) were incubated with either 0.2% polygalacturonic acid, rhamnogalacturonan I or xylotri-ose linked with glucuronic acid in 20 mM citrate/phosphate buffer pH 5.0 at 40°C for 16 h. The whole assay volumes were used for TLC afterward. Samples were applied to TLC plates (Silica gel 60, 20 cm, Merck) in 4 μ L steps, and enzymatic breakdown products were separated using the following mobile phase: ethyl acetate:glacial acetic acid:formic acid:water (9:3:1:4) for approximately 60 min. After drying, carbohydrates were stained by spraying the plates with a solution containing 0.2% (w/v) orcinol in methanol:sulfuric acid (9:1), followed by a short heating until spots appeared. The reference standard either contained 2 μ g each of galacturonic, di-galacturonic, tri-galacturonic acid and rhamnose, or 2 μ g each of xylose, di-xylose, tri-xylose and glucuronic acid.