

ASYMPTOTICS OF CLASSICAL SPIN NETWORKS

STAVROS GAROUFALIDIS AND ROLAND VAN DER VEEN

With an appendix by Don Zagier

ABSTRACT. A spin network is a cubic ribbon graph labeled by representations of $SU(2)$. Spin networks are important in various areas of Mathematics (3-dimensional Quantum Topology), Physics (Angular Momentum, Classical and Quantum Gravity) and Chemistry (Atomic Spectroscopy). The evaluation of a spin network is an integer number. The main results of our paper are: (a) an existence theorem for the asymptotics of evaluations of arbitrary spin networks (using the theory of G -functions), (b) a rationality property of the generating series of all evaluations with a fixed underlying graph (using the combinatorics of the chromatic evaluation of a spin network), (c) rigorous effective computations of our results for some $6j$ -symbols using the Wilf-Zeilberger theory, and (d) a complete analysis of the regular Cube $12j$ spin network (including a non-rigorous guess of its Stokes constants), in the appendix.

CONTENTS

1. Introduction	2
1.1. Spin networks in mathematics, physics and chemistry	2
1.2. The evaluation of a spin network	2
1.3. Three fundamental problems	3
1.4. A solution to Problem 1.2	3
1.5. A partial solution to Problem 1.3	5
1.6. A conjecture regarding Problem 1.4	5
1.7. Acknowledgement	5
2. Evaluation of spin networks	5
2.1. Evaluation of spin networks by recoupling	6
2.2. Generating series and chromatic evaluation	8
2.3. Chromatic evaluation	9
2.4. Proof of Theorem 2.9	12
3. Asymptotic expansions	12
3.1. Sequences of Nilsson type	12
3.2. G -functions	13
3.3. Hypergeometric terms	13
3.4. Integral representation of spin network evaluations	14
4. Examples and a conjecture on growth rates	15
4.1. The $6j$ -symbol and the tetrahedron	15
4.2. A conjecture on growth rates	17
5. Challenges and future directions	18
Appendix A. Asymptotics of the regular cube	19

Date: December 19, 2011.

S.G. was supported in part by NSF. R.V. was supported in part by the Netherlands Organisation for Scientific Research (NWO).

1991 *Mathematics Classification.* Primary 57N10. Secondary 57M25.

Key words and phrases: Spin networks, ribbon graphs, recoupling, $6j$ -symbols, Racah coefficients, angular momentum, asymptotics, G -functions, Kauffman bracket, Jones polynomial, Wilf-Zeilberger method, enumerative combinatorics, Nilsson.

Appendix B. Further comments on the asymptotics of the regular cube	22
References	22

1. INTRODUCTION

1.1. Spin networks in mathematics, physics and chemistry. A (classical) *spin network* (Γ, γ) consists of a *cubic ribbon graph* Γ (i.e., an abstract trivalent graph with a cyclic ordering of the edges at each vertex) and a coloring γ of its set of edges by natural numbers. According to Penrose, spin networks correspond to a diagrammatic description of tensors of representations of $SU(2)$. Here a color k on an edge indicates the $k + 1$ dimensional irreducible representation of $SU(2)$, and their evaluation is a contraction of the above tensors. Spin networks originated in work by Racah and Wigner in atomic spectroscopy in the late forties [Rac46, Rac42a, Rac42b, Rac49, Wig41]. Exact or asymptotic evaluations of spin network is a useful and interesting topic studied by Ponzano-Regge, Biedenharn-Louck and many others; see [BL81a, BL81b, PR68, VMK88]. In the past three decades, spin networks have been used in relation to classical and quantum gravity and angular momentum in 3-dimensions; see [EPR08, Pen71a, Pen71b, RS95]. In mathematics, q -deformations of spin networks (so called quantum spin networks) appeared in the eighties in the work of Kirillov-Reshetikhin [KR89]. *Quantum spin networks* are *knotted framed trivalent graphs* embedded in 3-space with a cyclic ordering of the edges near every vertex, and their evaluations are rational functions of a variable q . The quantum theta and $6j$ -symbols are the building blocks for topological invariants of closed 3-manifolds in the work of Turaev-Viro [TV92, Tur94]. Quantum spin networks are closely related to a famous invariant of knotted 3-dimensional objects, the celebrated *Jones polynomial*, [Jon87]. A thorough discussion of quantum spin networks and their relation to the Jones polynomial and the *Kauffman bracket* is given in [KL94] and [CFS95]. Recent papers on asymptotics of spin networks in physics and mathematics include: [AHH⁺09], [LY11] and [CM11]. Aside from the appearances of spin networks in the above mentioned areas, their evaluations and their asymptotics lead to challenging questions even for simple networks such as the cube, discussed in detail in the appendix. Some examples of spin networks that will be discussed in the paper are shown in Figure 1.




Figure 1. From left to right: The theta, the tetrahedron or $6j$ -symbol, the Cube, the 5-sided prism and the complete bipartite graph $K_{3,3}$ or $9j$ -symbol. The cyclic order of the edges around each vertex is counterclockwise. The left three spin networks are admissible, and the right two are not.


1.2. The evaluation of a spin network.

Definition 1.1. (a) We say a spin network is *admissible* when the sum of the three labels a, b, c around every vertex is even and a, b, c satisfy the triangle inequalities: $|a - b| \leq c \leq a + b$.

(b) The Penrose evaluation $\langle \Gamma, \gamma \rangle^P$ of a spin network (Γ, γ) is defined to be zero if it is not admissible. If it is admissible, its evaluation is given by the following algorithm.

- Use the cyclic ordering to thicken the vertices into disks and the edges into untwisted bands.

- Replace the vertices and edges by the linear combinations of arcs as follows:

(1) 

- Finally the resulting linear combination of closed loops is evaluated by assigning the value $(-2)^n$ to a term containing n loops.

In the above definition the summation is over all permutations σ of the a arcs at an edge colored a . The Penrose evaluation $\langle \Gamma, \gamma \rangle^P$ of a spin network is always an integer. Note that the admissibility condition is equivalent to saying that the strands can be connected at each vertex as in Figure 1. Note also that cubic ribbon graphs Γ are allowed to have multiple edges, loops and several connected components including components that contain no vertices. In addition, Γ is allowed to be non-planar (contrary to the requirement of many authors [Wes98, Mou79, KL94]), as long as one fixes a cyclic ordering of the edges at each vertex. The latter condition is implicit in [Pen71a]. It turns out that changing the cyclic ordering at a vertex of a spin network changes its evaluation by single sign; see Lemma 2.1 below.

1.3. Three fundamental problems. It is easy to see that if (Γ, γ) is an admissible spin network and n is a natural number, then $(\Gamma, n\gamma)$ is also admissible. A fundamental problem is to study the asymptotic behavior of the sequence of evaluations $\langle \Gamma, n\gamma \rangle^P$ when n is large. This problem actually consists of separate parts. Fix an admissible spin network (Γ, γ) .

Problem 1.2. Prove the existence of an asymptotic expansion of the sequence $\langle \Gamma, n\gamma \rangle^P$ when n is large.

Problem 1.3. Compute the asymptotic expansion of the sequence $\langle \Gamma, n\gamma \rangle^P$ to all orders in n effectively.

Problem 1.4. Identify the terms in the asymptotic expansion of $\langle \Gamma, n\gamma \rangle^P$ with geometric invariants of the spin network.

These problems are motivated by the belief that the quantum mechanics of particles with large spin will approximate the classical theory. To the best of our knowledge, the literature for Problem 1.2 is relatively new and short and concerns only thetas and $6j$ -symbols with certain labellings. For Problem 1.3, it should be noted that even for the $6j$ -symbols not much is known about the subleading terms in the asymptotic expansion. Some terms are found in [DL09] but no general algorithm is given. As for the geometric interpretation in Problem 1.4 again there is a well known conjecture in the case of the $6j$ -symbol [PR68]. Roberts used geometric quantization techniques to prove this conjecture on the leading asymptotic behavior of $6j$ -symbols in the so-called Euclidean case [Rob99, Rob02]. Some results on the $9j$ -symbol have been found in [HL10]. Finally more general interpretation for the leading order asymptotics appears in [CM11] however this assumes a hypothesis that has not been shown to hold in cases other than the $6j$ -symbol. Problems 1.2–1.4 can also be viewed as the classical analogue of the problem of understanding the asymptotics of quantum spin networks and quantum invariants. Even less is known in the quantum case but see [GvdV11] and a well known conjecture in this context is the volume conjecture [Kas97], [MM01].

1.4. A solution to Problem 1.2. In this paper we give a complete solution to Problem 1.2 in full generality. A convenient role is played by the following normalization of the spin network evaluation. This normalization was introduced independently in [Cos09] in the q -deformed case.

Definition 1.5. We define the *standard normalization* of a spin network evaluation to be

(2)
$$\langle \Gamma, \gamma \rangle = \frac{1}{\mathcal{I}!} \langle \Gamma, \gamma \rangle^P$$

where $\mathcal{I}!$ is defined to be the product

$$(3) \quad \mathcal{I}! = \prod_{v \in V(\Gamma)} \left(\frac{-a_v + b_v + c_v}{2} \right)! \left(\frac{a_v - b_v + c_v}{2} \right)! \left(\frac{a_v + b_v - c_v}{2} \right)!$$

where a_v, b_v, c_v are the colors of the edges emanating from vertex v , and $V(\Gamma)$ is the set of vertices of Γ .

The standard normalization has a number of useful properties (see Theorem 1.7 below) that can be stated conveniently in terms of a generating function that we now define. If we fix a cubic ribbon graph Γ one can consider many spin network evaluations, one for each admissible labeling γ of Γ . We organize these in a generating function by taking a formal variable for every edge and encoding γ in the exponents of monomials in these variables.

Definition 1.6. Given a cubic ribbon graph Γ define a formal power series in the variables $z = (z_e)_{e \in E(\Gamma)}$ by

$$F_\Gamma(z) = \sum_{\gamma \geq 0} \langle \Gamma, \gamma \rangle z^\gamma$$

where $z^\gamma = \prod_{e \in E(\Gamma)} z_e^{\gamma(e)}$, and $E(\Gamma)$ denote the set of edges of Γ .

By virtue of our use of the standard normalization we can prove the following theorem about our generating function F_Γ .

Theorem 1.7. (1) For all spin networks (Γ, γ) , the standard evaluation $\langle \Gamma, \gamma \rangle$ is an integer.
(2) The sequence $\langle \Gamma, n\gamma \rangle$ is exponentially bounded.
(3) For any cubic ribbon graph Γ the generating series F_Γ is rational function explicitly defined in terms of Γ .

To illustrate the last part of the theorem let us mention the special case in which Γ is planar with the counterclockwise orientation. In this case a result from [Wes98] that states that

$$F_\Gamma(z) = \frac{1}{P_\Gamma(z)^2}$$

where $P_\Gamma(z) = \sum_{c \in C_\Gamma} z^c$ and C_Γ is the set of 2-regular subgraphs of Γ . Our theorem generalizes this result to arbitrary Γ , the precise statement can be found in Theorem 2.9. The next result gives a complete answer to Problem 1.2. To state it, we need to recall a useful type of sequence; see [Gar09, Gar11].

Definition 1.8. We say that a sequence (a_n) is of *Nilsson type* if it has an *asymptotic expansion* of the form

$$(4) \quad a_n \sim \sum_{\lambda, \alpha, \beta} \lambda^n n^\alpha (\log n)^\beta S_{\lambda, \alpha, \beta} h_{\lambda, \alpha, \beta}(1/n)$$

where

- the summation is over a finite set of triples (λ, α, β) ,
- the *growth rates* λ are algebraic numbers of equal magnitude,
- the *exponents* α are rational and the *nilpotency exponents* β are natural numbers,
- the *Stokes constants* $S_{\lambda, \alpha, \beta}$ are complex numbers,
- the $h_{\lambda, \alpha, \beta}(x)$ are formal power series with coefficients in a number field K such that the coefficient of x^n is bounded by $C^n n!$ for some $C > 0$ and the constant coefficient is 1.

Note that the set of sequences of Nilsson type is closed under addition and pointwise multiplication. Using the theory of G -functions, (discussed in Section 3.2), we prove:

Theorem 1.9. For any spin network (Γ, γ) the sequence $\langle \Gamma, n\gamma \rangle$ is of Nilsson type.

1.5. **A partial solution to Problem 1.3.** Regarding Problem 1.3, we introduce a new method (the Wilf-Zeilberger theory) which

- computes a linear recursion for the sequence $\langle \Gamma, n\gamma \rangle$,
- given a linear recursion, effectively computes the corresponding triples (λ, α, β) , the number field K and any number of terms of the power series $h_{\lambda, \alpha, \beta}(x) \in 1 + xK[[x]]$ in Definition 1.8,
- numerically computes the Stokes constants $S_{\lambda, \alpha, \beta}$.

Given this information, one may guess exact values of the Stokes constants. In some cases, we obtain alternative exact computation of the Stokes constants, too. As an illustration of the theorem we will present computations of the asymptotic expansions of three representative $6j$ -symbols up to high order using the Wilf-Zeilberger method in Section 4.1. In the appendix we will present additional numerical results on the asymptotic expansion of the case of the cube spin network. About 20 more examples of spin network evaluations (including the s -sided prisms for $s = 2, \dots, 7$ and the twisted s -sided prisms for $s = 2, \dots, 5$) have been computed, and the data is available from the first author upon request.

1.6. **A conjecture regarding Problem 1.4.** The example of the cube spin network also provides evidence for the following conjecture on the growth rates λ in the Nilsson type expansion. The conjecture connects the growth rates of suitable spin networks to the total mean curvature of a related Euclidean polyhedron. Let P be a convex polyhedron in three dimensional Euclidean space. Denote by $M(P)$ the total mean curvature of P . Recall that $M(P) = \frac{1}{2} \sum_e \ell_e \phi_e$, where ϕ_e is the exterior dihedral angle at edge e and ℓ_e is the length of the edge.

Conjecture 1.10. Let (Γ, γ) be a planar spin network such that the dual of Γ is realized as the 1-skeleton of a convex Euclidean polyhedron P with edge lengths given by γ . The numbers $e^{\pm iM(P)}$ are growth rates in the asymptotic expansion of the unitary evaluations of $\langle \Gamma, n\gamma \rangle^U$.

In the conjecture we are using the so called unitary evaluation of a spin network defined in Section 4. This evaluation differs from the standard one by an explicit factor ensuring that it is still of Nilsson type.

After this work was completed, an approach to Problems 1.2–1.4 was proposed by Costantino-Marche, [CM11] using generating functions. Their approach requires certain nondegeneracy conditions, and in particular does not give a solution to Problem 1.2 or Problem 1.3 for the regular cube spin network, see the Appendix.

1.7. **Acknowledgement.** The results were conceived in a workshop in Aarhus, Denmark, and presented in HaNoi, Vietnam and Strasbourg, France in the summer of 2007. The authors wish to thank the organizers for their hospitality. S.G. wishes to thank C. Koutschan, D. Zeilberger and D. Zagier for many enlightening conversations.

2. EVALUATION OF SPIN NETWORKS

In this section we treat two ways of calculating the evaluation of a spin network. The first is by recoupling theory and leads to practical but non-canonical formulas for the evaluations as multi-sums. The second way is the method of chromatic evaluation. This leads to the proof of the generating function result, Theorem 1.7 announced above.

We start by recording some elementary facts about spin network evaluations. First of all our definition of the standard evaluation assumes that there are no edges without vertices. By definition we will add an $(a, a, 0)$ colored vertex to any a -colored component that has none. This makes sense because of Part (a) of the following.

Lemma 2.1. Let (Γ, γ) be a spin network and consider the standard evaluation.

- (a) Inserting a vertex colored $(0, a, a)$ in the interior of an edge of Γ colored a does not change the standard evaluation of the spin network.
- (b) Changing the cyclic ordering at a vertex whose edges are colored a, b, c changes the evaluation by a sign $(-1)^{(a(a-1)+b(b-1)+c(c-1))/2}$

Proof. (a) The chosen normalization introduces an extra factor $1/a!$ for the new vertex labeled $(0, a, a)$, while it follows from the definition that one also inserts an extraneous summation over permutations in the pre-existing edge labeled a . Since

$$\sum_{\sigma \in S_a} \sum_{\tau \in S_a} \text{sign}(\sigma)\sigma \text{sign}(\tau)\tau = a! \sum_{\sigma \in S_a} \text{sign}(\sigma)\sigma$$

the evaluation is unchanged.

(b) Changing the cyclic order at a vertex with edge labels a, b, c has the following effect. The alternating sum at each of the adjacent edges is multiplied by the permutation that turns the arcs in the edge by 180 degrees. This element has sign $a(a-1)/2$ in S_a . \square

As a consequence of part (a) of the above lemma, an edge labeled 0 in a spin network can be removed without affecting the evaluation. There is an alternative *bracket normalization* $\langle \Gamma, \gamma \rangle^B$ of the evaluation of a spin network (Γ, γ) which agrees with a specialization of the *Jones polynomial* or *Kauffman bracket*.

Definition 2.2. The *bracket normalization* of a spin network (Γ, γ) is defined by

$$(5) \quad \langle \Gamma, \gamma \rangle^B = \frac{1}{\mathcal{E}!} \langle \Gamma, \gamma \rangle^P$$

where

$$(6) \quad \mathcal{E}! = \prod_{e \in E(\Gamma)} \gamma(e)!$$

This normalization has the property that it coincides with the Kauffman bracket (Jones polynomial) of a quantum spin network evaluated at $A = -1$ [KL94]. However, $\langle \Gamma, \gamma \rangle^B$ is not necessarily an integer number, and the analogous generating series does not satisfy the rationality property of Theorem 1.7.

2.1. Evaluation of spin networks by recoupling. In this subsection we describe a way of evaluating spin networks by recoupling. We will reduce the evaluation of spin networks to multi-dimensional sums of $6j$ and theta-symbols. The value of the $6j$ and theta-symbols is given by the following lemma of [KL94] and [Wes98], using our normalization. The choice of letters in labeling the $6j$ -symbol is traditional following for example [KL94].

Lemma 2.3. (a) Let (\triangleleft, γ) denote a tetrahedron labeled and oriented as in Figure 1 with $\gamma = (a, b, c, d, e, f)$. Its standard evaluation is given by

$$(7) \quad \langle \triangleleft, \gamma \rangle = \sum_{k=\max T_i}^{\min S_j} (-1)^k (k+1) \binom{k}{S_1 - k, S_2 - k, S_3 - k, k - T_1, k - T_2, k - T_3, k - T_4}$$

where, as usual

$$\binom{a}{a_1, a_2, \dots, a_r} = \frac{a!}{a_1! \dots a_r!}$$

denotes the multinomial coefficient when $a_1 + \dots + a_r = a$, and S_i are the half sums of the labels in the three quadrangular curves in the tetrahedron and T_j are the half sums of the three edges emanating from a given vertex. In other words, the S_i and T_j are given by

$$(8) \quad S_1 = \frac{1}{2}(a + d + b + c) \quad S_2 = \frac{1}{2}(a + d + e + f) \quad S_3 = \frac{1}{2}(b + c + e + f)$$

$$(9) \quad T_1 = \frac{1}{2}(a + b + e) \quad T_2 = \frac{1}{2}(a + c + f) \quad T_3 = \frac{1}{2}(c + d + e) \quad T_4 = \frac{1}{2}(b + d + f).$$

(b) Let (Θ, γ) denote the Θ spin network of Figure 1 admissibly colored by $\gamma = (a, b, c)$. Then we have

$$(10) \quad \langle \Theta, \gamma \rangle = (-1)^{\frac{a+b+c}{2}} \left(\frac{a+b+c}{2} + 1 \right) \binom{\frac{a+b+c}{2}}{\frac{-a+b+c}{2}, \frac{a-b+c}{2}, \frac{a+b-c}{2}}.$$

Finally note that the evaluation of an n -labeled unknot is equal to $(-1)^n(n+1)$. Recoupling is a way to modify a spin network locally, while preserving its evaluation. This is done as in Figure 2. The top-most formula is called the recoupling formula and follows from the recoupling formula in [KL94], using our conventions. The other two pictures in the figure show the bubble formula and the triangle formula. The




Figure 2. The recoupling formula (top), the bubble formula (left) and the triangle formula (right). The sum is over all k for which the network is admissible, and $\delta_{k,l}$ is the *Kronecker delta function*.

bubble formula shown on the left of Figure 2 serves to remove all bigon faces. Likewise the triangle formula can be used to remove triangles. To see why the recoupling and bubble formulas suffice to write any spin network as a multi sum of products of $6j$ -symbols divided by thetas we argue as follows. Applying the recoupling formula to a cycle in the graph reduces its length by one. Keep going until you get a multiple edge which can then be removed by the bubble formula. Although the triangle formula follows quickly from the bubble formula and the recoupling formula it is important enough to state on its own. For example the triangle formula shows that the evaluation of the class of *triangular networks* is especially simple. The triangular networks are the planar graphs that can be obtained from the tetrahedron by repeatedly replacing a vertex by a triangle. By the triangle formula the evaluation of any triangular network is simply a product of $6j$ -symbols divided by thetas. No extraneous summation will be introduced. To illustrate how recoupling theory works, let us evaluate the regular s -sided prism and $K_{3,3}$. Consider the s -sided prism network as shown in Equation (11) (for $s = 5$) where every edge is colored by the integer n . In the figure we have left out most of the labels n for clarity. By convention unlabeled edges are colored by n . Performing the recoupling move on every inward pointing edge we transform the prism into a string of bubbles that is readily evaluated.

$$(11) \quad \begin{array}{c} \text{Prism} \\ \text{with } n \text{ labels} \end{array} = \sum_{k \text{ admissible}} \left(\frac{\text{Bubble with } k \text{ labels}}{\text{Bubble with } k \text{ labels}} \right)^s \text{String of bubbles} = \sum_{k \text{ admissible}} \left(\frac{\text{Bubble with } k \text{ labels}}{\text{Bubble with } k \text{ labels}} \right)^s \text{Loop with } k \text{ label}$$

Observing that if n is odd the network is not admissible (and thus evaluates to zero), and denoting the tetrahedron and the theta with one edge colored by k and the others by n by $S(n, k)$ and $\theta(n, k)$ we conclude the following formula for the n -colored s -sided prism.

Proposition 2.4. If $n = 2N$ is even we have

$$\langle \text{Prism}_s, 2N \rangle = \sum_{j=0}^{2N} (2j+1) \left(\frac{S(2N, 2j)}{\theta(2N, 2j)} \right)^s$$

and if n is odd we have $\langle \text{Prism}_s, n \rangle = 0$.

For small values of s the prism can be evaluated in a more straight forward way, thus providing some well known identities of $6j$ -symbols. Namely when $s = 1$ we get zero, when $s = 2$ we find some thetas and when $s = 3$ we have by the triangle formula a product of two $6j$ -symbols thus giving a special case of the Biedenharn-Elliott identity [KL94]. For $s = 4$ we find a formula for the regular cube, that will be used in the appendix. We know of no easier expression for the evaluation in this case. A similar computation for $K_{3,3}$ cyclically ordered as a plane hexagon with its three diagonals gives the following.

Proposition 2.5. If $n = 2N$ is even we have

$$\langle K_{3,3}, 2N \rangle = \sum_{j=0}^{2N} (-1)^j (2j+1) \left(\frac{S(2N, 2j)}{\theta(2N, 2j)} \right)^3$$

and if n is odd we have $\langle K_{3,3}, n \rangle = 0$.

Note the similarity between Prism_3 and $K_{3,3}$. The only difference is the sign that comes up in the calculation when one needs to change the cyclic order. The extra sign makes $\langle K_{3,3}, 2N \rangle = 0$ for all odd N . This is because changing the cyclic ordering at a vertex takes the graph into itself, while it produces a sign $(-1)^N$ when all edges are colored $2N$.

2.2. Generating series and chromatic evaluation. Recall the generating function $F_\Gamma(z)$ for all spin network evaluations with the same underlying graph Γ from Definition 1.6. We are using variables $z = (z_e)_{e \in E(\Gamma)}$, one for each edge, and abbreviate monomials $\prod_{e \in E(\Gamma)} z_e^{\gamma(e)}$ as z^γ . Our goal is to express F_Γ explicitly in terms of Γ . To do so we need a couple of definitions.

Definition 2.6. Given a cubic ribbon graph Γ define a *cycle* to be a (possibly disconnected) 2-regular subgraph of Γ . The set of all cycles is denoted by C_Γ .


In terms of the cycles we define a polynomial and a quadratic form.

Definition 2.7. Given a cubic ribbon graph Γ and $X \subset C_\Gamma$ we define

$$(12) \quad P_{\Gamma, X}(z) = \sum_{c \in C_\Gamma} \epsilon_X(c) \prod_{e \in c} z_e$$

where $\epsilon_X(c) = -1$ (resp. 1) when $c \in X$ (resp. $c \notin X$). Also define the function Q_Γ on the subsets of C_Γ as follows. Let $Q_\Gamma(X)$ be the number of unordered pairs $\{c, c'\} \subset X$ with the property that c and c' intersect in an odd number of places when drawn on the thickening of Γ .

Note that the cyclic orientation of Γ defines a unique thickening. Also, the polynomials $P_{\Gamma, X}$ all have constant coefficient 1. We will call $P_\Gamma = P_{\Gamma, \emptyset}$ the *cycle polynomial* of Γ . It is independent of the cyclic orientation of Γ . Notice how the other $P_{\Gamma, X}$ only differ from the cycle polynomial in the signs of the individual monomials. It is interesting to note that the cycle polynomial determines the cubic graph, up to a well-determined ambiguity. This follows from a classic theorem of Whitney, which we quote for the benefit of the reader. Recall that a graph Γ is 2-connected (resp. 3-connected) if it remains connected after removing any one (resp. any two) vertices of Γ . A *Whitney flip* is the following move on a graph (where R and L contain at least two edges):



A Whitney flip is the graph-theoretic analogue of a *knot mutation* (see [Ada94]) and can only be applied to graphs which are not 3-connected.

Proposition 2.8. [Whi33] (a) Let Γ_1, Γ_2 be two 2-connected cubic graphs with same cycle polynomial. Then Γ_2 is obtained from Γ_1 by a sequence of Whitney flips.
 (b) Let Γ be a 3-connected cubic graph. The cycle polynomial P_Γ uniquely determines Γ .

Proof. This follows from a more general theorem of Whitney [Whi33] that states the following. Two finite 2-connected graphs with a bijection on the set of vertices that preserves the set of cycles are related by a sequence of Whitney flips. This also works for non-cubic graphs if we define a *cycle* C of a finite graph Γ to be a subgraph of Γ with the same vertex set as Γ such that every vertex of C has even valency. In case Γ is a cubic graph, a cycle of Γ in the above sense exactly coincides with Definition 2.6. Thus, part (a) follows.

Part (b) follows from (a) and the fact that 3-connected graphs cannot be Whitney flipped. \square

With the definitions in place we can finally state the precise version of the last part of Theorem 1.7:

Theorem 2.9. *For every cubic ribbon graph Γ we have*

$$(13) \quad F_\Gamma(z) = \sum_{X \subset C_\Gamma} \frac{a_X}{P_{\Gamma, X}^2} \in \mathbb{Z}[[z]] \cap \mathbb{Q}(z).$$

where the coefficients are given by

$$a_X = \frac{1}{2^{|C_\Gamma|}} \sum_{Y \subset C_\Gamma} (-1)^{Q_\Gamma(Y) + |X \cap Y|}$$

Corollary 2.10. For every spin network (Γ, γ) , the evaluation $\langle \Gamma, \gamma \rangle$ is an integer number and $\langle \Gamma, n\gamma \rangle$ is exponentially bounded.

In particular Theorem 1.7 reduces to Theorem 2.9 above. To see how the particular case of planar spin networks comes about we note that

Corollary 2.11. When Γ is planar with the counterclockwise orientation, then all cycles intersect an even number of times so $(-1)^{Q(X)} = 1$ and hence only a_\emptyset is non-zero. It follows that

$$F_\Gamma = \frac{1}{P_{\Gamma, \emptyset}^2}$$

recovering an earlier theorem by Westbury [Wes98].

The proof of this theorem uses the *chromatic evaluation method* which goes back to [Pen71b]. Our proof builds on earlier work by [Wes98] and [KL94] on planar spin networks and will be given in the next subsection.

2.3. Chromatic evaluation.

Definition 2.12. For $N \in \mathbb{Z}$ define the evaluation $\langle \Gamma, \gamma \rangle_N^P$ just as in Definition 1.1 except that the value of the a loop is now N instead of -2 .

Note that by definition $\langle \Gamma, \gamma \rangle_{-2}^P = \langle \Gamma, \gamma \rangle^P$. However for positive N the evaluations are easier to work with. Let V be an N -dimensional vector space with basis $B = \{b_1, \dots, b_N\}$. For definiteness, let us state that by a loop we mean an immersion of the circle. Recall that in the definition of the evaluation, the first step is to replace a diagram of the graph Γ by a linear combination of closed loops. We will call this collection of embedded loops the expanded diagram of Γ . Instead of directly evaluating each closed loop as N we can also view the expanded diagram as the composition (and linear combination) of copies of three types of maps:

$$\begin{array}{lll} \cup : V \otimes V \rightarrow \mathbb{C} & \cap : \mathbb{C} \rightarrow V \otimes V & \times : V \otimes V \rightarrow V \otimes V \\ e_i \otimes e_j \mapsto \delta_{i,j} & 1 \mapsto \sum_i e_i \otimes e_i & e_i \otimes e_j \mapsto e_j \otimes e_i \end{array}$$

Here the direction of composition is upwards. Composing all the maps gives a linear map from the ground field \mathbb{C} to itself, which is multiplication by the scalar $\langle \Gamma, \gamma \rangle_N^P$. This interpretation of the invariant coupled with a counting argument leads us to an expression of the evaluation in terms of cycle configurations that we will define now.

Definition 2.13. Define a *cycle configuration* to be a function $L : C_\Gamma \rightarrow \mathbb{N}$ such that $L(\emptyset) = 0$. A cycle configuration L defines a coloring $\gamma(L)$ as follows

$$\gamma(L)(e) = \sum_{c \in C_\Gamma : e \in c} L(c) \quad \text{and} \quad |L| = \sum_{c \in C_\Gamma} L(c) \quad \text{and} \quad L! = \prod_{c \in C_\Gamma} L(c)!$$

Finally define the quadratic form Q_Γ on cycle configurations as $Q_\Gamma(L) = \sum_{\{c,d\}} L(c)L(d)$, where $\{c,d\} \subset C$ runs over the unordered pairs of cycles that intersect in an odd number of places.

Viewing a (non-empty) subset $X \subset C_\Gamma$ as the cycle configuration that is 1 on X and 0 elsewhere, this definition of Q_Γ coincides with the one given in Definition 2.7. We can now state and prove the main lemma that expresses the evaluation for positive N in terms of cycle configurations.

Lemma 2.14. For positive integers N we have

$$\langle \Gamma, \gamma \rangle_N^P = \sum_{L: \gamma(L)=\gamma} (-1)^{Q_\Gamma(L)} \binom{N}{L} \mathcal{I}!$$

Here $\binom{N}{L}$ is defined as $\frac{N(N-1)\dots(N-|L|+1)}{L!}$ and recall $\mathcal{I}!$ is the normalization factor from Definition 1.5.

Proof. We view $\langle \Gamma, \gamma \rangle_N^P$ as a linear combination of the composition of maps consisting of the three elementary maps as explained above. This can be made more precise by orienting the edges of Γ and defining Cups to be the set of the cups (local minima) in the expanded diagram of Γ . Now define the product of symmetric groups $S_\gamma = \prod_{e \in E(\Gamma)} S_{\gamma(e)}$. For a function $f : \text{Cups} \rightarrow B$ and $\sigma \in S_\gamma$, define $\langle f, \sigma \rangle$ to be 1 if f is constant along the loops we get by connecting the cups using the crossings prescribed by σ . In all other cases $\langle f, \sigma \rangle = 0$. Composing the elementary maps shows directly that the evaluation can be expressed as:

$$\langle \Gamma, \gamma \rangle_N^P = \sum_{\sigma \in S_\gamma} \text{sgn}(\sigma) \sum_{f: \text{Cups} \rightarrow B} \langle f, \sigma \rangle$$

Call a pair (f, σ) good if $\langle f, \sigma \rangle = 1$, no loops self intersect and given any two distinct loops ℓ, ℓ' sharing an edge of Γ we have $f(\ell) \neq f(\ell')$. Reversing the order of summation in the above equation we see that all but the good pairs (f, σ) cancel out. A good pair (f, σ) determines a unique cycle configuration $L_{f,\sigma}$ as follows. Define $L_{f,\sigma}(\emptyset) = 0$ and for non-empty cycles c we define $L_{f,\sigma}(c)$ to be the number of $b \in B$ such that $f^{-1}(b)$ traces out c . Notice that for any $b \in B$ the fiber $f^{-1}(b)$ consists of a set of loops that trace out a cycle c without hitting any edge of Γ twice.

Two good pairs that determine the same cycle configuration must have the same sign $\text{sgn}(\sigma)$. To see this we interpret the sign as $(-1)^{\# \text{ internal crossings}}$, where a crossing in the expanded diagram is *internal* if it comes from a choice of σ (not the crossings between distinct edges in a projection of Γ). Changing σ without changing the cycles that are to be traced out will always change the number of (internal) crossings by an even amount. It therefore makes sense to define the sign of a cycle configuration as $(-1)^{Q(L)}$. To summarize what we have done so far we can rewrite the evaluation as follows

$$\langle \Gamma, \gamma \rangle_N^P = \sum_{L: \gamma(L)=\gamma} (-1)^{Q(L)} |G'(L)|$$

where $G'(L)$ is the set of all good pairs (f, σ) satisfying $L_{f,\sigma} = L$. Ordering the basis B and the $|L|$ loops determined by L shows that $|G'(L)| = \frac{N!}{(N-|L|)!} |G(L)|$. Here $G(L)$ is the subset of $G'(L)$ in which the order of the basis elements agrees with the order of the loops. Elements of $G(L)$ are uniquely determined by σ and L so we will suppress f in the notation. We should however keep in mind that using f we are able to distinguish all the loops as we follow along the edges of Γ . In particular we know which arcs at the vertices of Γ trace out which cycles. To finish the proof we need to show that

$$|G(L)| = \frac{\mathcal{I}!}{L!}$$

Intuitively this can be understood as the number of ways to arrange the loops at the vertices, corrected by the symmetry of the cycle configuration. We will however give a careful proof of this fact below.

We will proceed to count elements of the set $G(L)$ by defining a transitive action of a group of order $\mathcal{I}!$ on $G(L)$ with stabilizer of size $L!$. To this end call a pair of half edges at a common vertex of Γ an *angle* and denote the set of all angles by \mathcal{A} . Looking at the expanded diagram of Γ we define $|a|$ to be the number of arcs passing through the angle a . By the cyclic orientation on Γ the angles are automatically oriented. Finally define $S_A = \prod_{a \in \mathcal{A}} S_{|a|}$ and notice that as promised $|S_A| = \mathcal{I}!$. The action of S_A on $G(L)$ is defined as follows. At every angle a we take a permutation T_a and its inverse and push T_a into the permutation at the edge of Γ it points to and T_a^{-1} into the edge the angle points away from. So more formally an element $T = (T_a)_{a \in \mathcal{A}} \in S_A$ acts on $\sigma = (\sigma_e)_{e \in E} \in G(L)$ by sending σ_e to $T_a T_b^{-1} \sigma_e T_c T_d^{-1}$. Here a, b, c, d are the angles adjacent to e , such that b and d are pointing away from e , see Figure 3. The action is well defined since the pairs (T_a, T_b) and (T_c, T_d) commute and the order of multiplication is determined by the orientation of the edge e . Also the action does not change the cycle configuration, just the way the loops are shaped by the permutations so it is indeed an action on $G(L)$. To prove that the action is transitive note that L and




Figure 3. On the left we see an edge e of Γ and its adjacent angles a, b, c, d together with their orientations. In the example on the right the permutation $\sigma_e = (1354)$ is drawn in the middle, and the numbers of strands in the four angles are given by $|a| = 1, |b| = 4, |c| = 3, |d| = 2$. The action of T is indicated by the white boxes.

f determine how the arcs making up the angles are connected. Any $\sigma' \in G(L)$ is therefore related to our σ by $\sigma'_e = \tau_{e,a} \tau_{e,b} \sigma_e \tau_{e,c} \tau_{e,d}$, for some $\tau_{e,j} \in S_{|j|}$ permuting the strands at the angle j . Let e, e' be the edges adjacent to a . Since σ and σ' both belong to the same cycle configuration L , the product $\tau_{e,a} \tau_{e',a}$ can only permute loops that trace the same cycle. The loops do not self intersect so the permutations of the loops along any (component of a) cycle multiply to 1. This means that one can modify the permutations τ so that they become inverses along each angle, without changing the product of the two permutations adjacent to any edge. Carrying out these modifications for each cycle gives the required T . It remains to show that the stabilizer of the action consists of $L!$ elements. So suppose we are given a $T \in S_A$ such that at every edge $\sigma_e = T_a T_b^{-1} \sigma_e T_c T_d^{-1}$. First of all the T_j can only permute arcs belonging to loops in the same cycle. Indeed, the T_j on one side of σ_e have to be canceled by the T 's on the other side. This is only possible if the permuted strands connect the same pair of angles at the edge. By definition of the action the permutations also have to cancel along each angle. This allows us to follow all the non-trivial T_j 's around, showing that the permuted loops must trace the same cycle. By the same follow along argument, permuting the arcs in a cycle at a single edge determines how they are to be permuted along the whole cycle. Therefore the cardinality of the stabilizer is $L!$. \square

Since the evaluation $\langle \Gamma, \gamma \rangle_N^P$ is a polynomial in N , the conclusion of the previous Lemma 2.14 actually holds for all N , in particular $N = -2$. We record this for future use as the following corollary, where we have switched back to the standard normalization.

Corollary 2.15.

$$\langle \Gamma, \gamma \rangle = \sum_{L: \gamma(L)=\gamma} (-1)^{Q_\Gamma(L)} \binom{-2}{L}$$

2.4. Proof of Theorem 2.9. To find a generating function for these evaluations, we first need to expand the sign $(-1)^{Q_\Gamma(L)}$ in terms of characters. That is we use Fourier analysis on the group $(\mathbb{Z}/2\mathbb{Z})^{|C_\Gamma|}$. We often write elements of this group as subsets of C_Γ . For every fixed $X \subset C_\Gamma$ we have a character $(-1)^{X(L)} = (-1)^{\sum_{x \in X} L(x)}$. In case L is a cycle configuration we extend the character to a Dirichlet character. So for some coefficients a_X we have

$$(-1)^{Q_\Gamma(L)} = \sum_{X \subset C_\Gamma} a_X (-1)^{X(L)}$$

Taking inner products, the coefficients are given by

$$a_X = \frac{1}{2^{|C_\Gamma|}} \sum_{Y \subset C_\Gamma} (-1)^{Q_\Gamma(Y) + |X \cap Y|}$$

To rewrite the generating function let us introduce variable for each cycle: $w = (w_c)_{c \in C_\Gamma}$ and set $w_c = \prod_{e \in c} z_e$. If $\gamma(L) = L$ then the color of an edge is the sum of the number of cycles (with multiplicity) containing that edge, hence

$$(-1)^{X(L)} z^\gamma = (-1)^{X(L)} w^L = \prod_{c \in C_\Gamma} (\epsilon_X(c) w_c)^{L(c)}$$

where $\epsilon_X(c)$ is the function that is 1 if $c \notin X$ and -1 if $c \in X$. Recall the cycle polynomial $P_{\Gamma, X}(z) = \sum_{c \in C_\Gamma} \epsilon_X(c) w_c$ so we can compute

$$\sum_{\gamma} \sum_{L: \gamma(L)=\gamma} \binom{N}{L} (-1)^{X(L)} z^\gamma = \sum_L \binom{N}{L} \prod_{c \in C_\Gamma} (\epsilon_X(c) w_c)^{L(c)} = (1 + \sum_{\emptyset \neq c \in C_\Gamma} \epsilon_X(c) w_c)^N = P_{\Gamma, X}^N$$

Now setting $N = -2$ everywhere and applying Corollary 2.15 we can finish the proof

$$F_\Gamma(z) = \sum_{\gamma} \langle \Gamma, \gamma \rangle z^\gamma = \sum_{X \subset C_\Gamma} a_X P_{\Gamma, X}^{-2}$$

□

3. ASYMPTOTIC EXPANSIONS

The goal of this section is to prove Theorem 1.9 that provides a Nilsson type asymptotic expansion for spin network evaluations. The general idea is to define the (single variable) generating function

Definition 3.1. Let (Γ, γ) be a spin network. The single variable generating function $F_{\Gamma, \gamma}$ is the formal power series

$$(14) \quad F_{\Gamma, \gamma}(z) = \sum_z \langle \Gamma, n\gamma \rangle z^n$$

Our goal is to show that this function is a G -function (defined in Section 3.2 below). It then follows from the theory of G -functions that the sequence $\langle \Gamma, n\gamma \rangle$ is of Nilsson type. Before doing so we first make some comments on Nilsson type sequences in general.

3.1. Sequences of Nilsson type. Recall from Definition 1.8 that a Nilsson type sequence (a_n) has the following asymptotic expansion as $n \rightarrow \infty$

$$a_n \sim \sum_{\lambda, \alpha, \beta} \lambda^n n^\alpha (\log n)^\beta S_{\lambda, \alpha, \beta} h_{\lambda, \alpha, \beta}(1/n)$$

The meaning of this expansion is entirely analogous to the more familiar special case where there is only one growth rate:

$$a_n \sim \lambda^n n^\alpha (\log n)^\beta \sum_{k=0}^{\infty} \frac{\mu_k}{n^k}$$

(which goes back to Poincaré, [Olv97]). In this case the meaning is that for every $r \in \mathbb{N}$ we have

$$\lim_{n \rightarrow \infty} n^r \left(a_n \lambda^{-n} n^{-\alpha} (\log n)^{-\beta} - \sum_{k=0}^{r-1} \frac{\mu_k}{n^k} \right) = \mu_r.$$

The general case is similar but to express it we would need more notation see [Gar11]. It can be shown that a Nilsson type sequence has a unique asymptotic expansion [Gar11]. With respect to point wise addition and multiplication the Nilsson type sequences form a \mathbb{C} -algebra. Formally multiplying the asymptotic expansions will produce the asymptotic expansion of the product sequence. When adding sequences of Nilsson type, the asymptotic expansion of the sum sequence is only the sum of the asymptotic expansions if all the growth rates have the same absolute value. Otherwise only the terms with maximal growth rate occur. The field K generated by the coefficients of the power series $h_{\lambda, \alpha, \beta}$ in a Nilsson type expansion a number field. See for example Section 4. An important source of Nilsson type sequences are G -functions that we will introduce next.

3.2. G -functions. In this section we recall the notion of a G -function, introduced by Siegel ([Sie29]) in relation to transcendence problems in number theory. Many of their arithmetic and algebraic properties were established by André in [And00]. G -functions appear naturally in Geometry (as Variations of Mixed Hodge Structures), in Arithmetic and most recently in Enumerative Combinatorics. For a detailed discussion, see [And00, Gar09] and references therein.

Definition 3.2. We say that a series $G(z) = \sum_{n=0}^{\infty} a_n z^n$ is a G -function if

- (a) the coefficients a_n are algebraic numbers,
- (b) there exists a constant $C > 0$ so that for every $n \geq 1$ the absolute value of every conjugate of a_n is less than or equal to C^n ,
- (c) the common denominator of the algebraic numbers a_0, \dots, a_n is less than or equal to C^n ,
- (d) $G(z)$ is holonomic, i.e., it satisfies a linear differential equation with coefficients polynomials in z .

For the purposes of this paper the most important property of G -functions is expressed in the following lemma.

Lemma 3.3. [Gar09, Prop.2.5] The sequence of Taylor coefficients of a G -function at $z = 0$ is a sequence of Nilsson type.

With the help of Lemma 3.3 we can now reduce the proof of Theorem 1.9 to the following lemma

Lemma 3.4. For any admissible spin network (Γ, γ) the generating function $F_{\Gamma, \gamma}(z)$ is a G -function.

In the next subsection we will give a proof of Lemma 3.4.

3.3. Hypergeometric terms. In this subsection we prove Lemma 3.4 (and hence Theorem 1.9) by showing that the standard evaluations of spin networks are a certain type of hypergeometric multisums that we will first describe in general.

Definition 3.5. An r -dimensional *balanced hypergeometric datum* \mathfrak{t} (in short, *balanced datum*) in variables (n, k) , where $n \in \mathbb{N}$ and $k = (k_1, \dots, k_r) \in \mathbb{N}^r$, is

- a finite list $\{(\epsilon_j, A_j(n, k)) \mid j \in J\}$ where $A_j : \mathbb{N}^{r+1} \rightarrow \mathbb{Z}$ is an affine linear form in (n, k) and $\epsilon_j \in \{-1, 1\}$ for all $j \in J$.
- a vector (C_0, \dots, C_r) of algebraic numbers

that satisfies the *balancing condition*

$$(15) \quad \sum_{j=1}^J \epsilon_j A_j = \text{constant}$$

and moreover, the set

$$(16) \quad P_{\mathfrak{t}} = \{x \in \mathbb{R}_{\geq 0}^r \mid A_j(1, x) \geq 0 \text{ for all } j \in J\}$$

is a *compact* rational convex polytope.

A balanced datum \mathfrak{t} gives rise to a balanced term $\mathfrak{t}(n, k)$ (defined for $n \in \mathbb{N}$ and $k \in \mathbb{Z}^r \cap nP_{\mathfrak{t}}$), to a sequence $(a_{\mathfrak{t}, n})$ and to a generating series $G_{\mathfrak{t}}(z)$ defined by:

$$(17) \quad \mathfrak{t}(n, k) = C_0^n \prod_{i=1}^r C_i^{k_i} \prod_{j=1}^J A_j(n, k)!^{\epsilon_j}$$

$$(18) \quad a_{\mathfrak{t}, n} = \sum_{k \in \mathbb{Z}^r \cap nP_{\mathfrak{t}}} \mathfrak{t}(n, k)$$

$$(19) \quad G_{\mathfrak{t}}(z) = \sum_{n=0}^{\infty} a_{\mathfrak{t}, n} z^n$$

We will call the sequences $(a_{\mathfrak{t}, n})$ balanced multisums. The connection between balanced multisum sequences and their asymptotics was given in [Gar09] using the theory of G -functions. More precisely,

Lemma 3.6. [Gar09, Thm.2] If \mathfrak{t} is a balanced datum, then the corresponding series $G_{\mathfrak{t}}(z)$ is a G -function.

Using this Lemma we can now easily prove Lemma 3.4.

Proof. (of Lemma 3.4) Using the recoupling formulae from Section 2.1 we can write $\langle \Gamma, \gamma \rangle$ as a multi-dimensional sum of products of $6j$ -symbols, theta-symbols and unknots (i.e., $1j$ -symbols) with a denominator consisting of theta-symbols. It follows from Equations (7) and (10) that the $6j$ -symbols (resp. theta-symbols) are balanced 1-dimensional (resp. 0-dimensional) sums, thus the ratio of the product of the theta-symbols by the product of the theta-symbols is a balanced multi-dimensional sum. The unknots can be written as $(-1)^k (k+1)!/k!$ and are therefore balanced as well. It is easy to check that admissibility guarantees that the multi-dimensional sum has finite range. \square

Beware that the term $\mathfrak{t}(n, k)$ constructed in the above proof is neither unique nor canonical in any sense.

3.4. Integral representation of spin network evaluations. In this final subsection we comment on the connection between Lemma 3.4 and Theorem 1.7 on the rationality of the multivariate generating function. The idea is that the single variable generating function $F_{\Gamma, \gamma}$ is a diagonal of the multivariate generating function F_{Γ} , where the diagonal is defined as follows.

Definition 3.7. Given a power series $f(x_1, \dots, x_r) \in \mathbb{Q}[[x_1, \dots, x_r]]$ and an exponent $J = (j_1, \dots, j_r) \in \mathbb{N}_+^r$, we define the J -diagonal of f by

$$(20) \quad (\Delta_J f)(z) = \sum_{n=0}^{\infty} [x^{nJ}](f) z^n \in \mathbb{Q}[[z]]$$

where $[x^{nJ}](f)$ denotes the coefficient of $x_1^{nj_1} \dots x_r^{nj_r}$ in f .

For every spin network (Γ, γ) we have

$$(21) \quad F_{\Gamma, \gamma} = \Delta_{\gamma} F_{\Gamma}$$

Consequently, the G -function $F_{\Gamma, \gamma}(z)$ is the diagonal of a rational function, and thus it comes from geometry in the following sense. Fix a power series $f(x_1, \dots, x_r) \in \mathbb{Q}[[x_1, \dots, x_r]]$ convergent at the origin and an exponent $J = (j_1, \dots, j_r) \in \mathbb{N}_+^r$ and consider the diagonal $(\Delta_J f)(z) \in \mathbb{Q}[[z]]$ as in Definition 3.7. Let \mathcal{C} denote a small real r -dimensional torus around the origin. Then we have the following.

Lemma 3.8. With the above assumptions,

$$(22) \quad (\Delta_J f)(z) = \frac{1}{(2\pi i)^r} \int_{\mathcal{C}} \frac{f(x_1, \dots, x_r)}{x_1^{j_1} \dots x_r^{j_r} - z} dx_1 \wedge \dots \wedge dx_r.$$

Proof. With the notation of Definition 20, an application of Cauchy's theorem gives for every natural number n

$$[x^{nJ}](f) = \frac{1}{(2\pi i)^r} \int_{\mathcal{C}} \frac{f(x_1, \dots, x_r)}{(x_1^{j_1} \dots x_r^{j_r})^{n+1}} dx_1 \wedge \dots \wedge dx_r.$$

Multiplying by z^n and summing up for n and interchanging summation and integration concludes the proof. \square

If in addition $f(x_1, \dots, x_r)$ is a rational function, then the singularities of the analytic continuation of the right hand-side of (22) can be analyzed by deforming the integration cycle \mathcal{C} and studying the corresponding variation of Mixed Hodge Structure as in [BK08]. Such G -functions come from geometry; see [And00, BK08].

4. EXAMPLES AND A CONJECTURE ON GROWTH RATES

In this section we illustrate the result of Theorem 1.9 on the asymptotic expansions in the case of the $6j$ -symbol. We also review the well known geometric interpretation of the leading asymptotics in this case. Finally we formulate a conjecture on the geometric meaning of the growth rates in the asymptotic expansion of more general spin networks. To discuss the geometric aspects of the asymptotics of spin networks it is convenient to introduce one more normalization of spin network evaluations

Definition 4.1. We define the *unitary normalization* $\langle \Gamma, \gamma \rangle^U$ of a spin network evaluation (Γ, γ) to be

$$\langle \Gamma, \gamma \rangle^U = \frac{1}{\Theta(\gamma)} \langle \Gamma, \gamma \rangle$$

where

$$\Theta(\gamma) = \prod_{v \in V(\Gamma)} \sqrt{|\langle \Theta, a_v, b_v, c_v \rangle|}$$

and a_v, b_v, c_v are the colors of the edges at vertex v .

Since the asymptotics of the normalization factor $\Theta(\gamma)$ is of Nilsson type by Stirling's formula [Olv97], we see that $\langle \Gamma, n\gamma \rangle^U$ is still of Nilsson type.

4.1. The $6j$ -symbol and the tetrahedron. The special case of the tetrahedral spin network or $6j$ -symbol motivates much of the questions we asked in the introduction. There is a well known interpretation of the leading asymptotics in terms of a metric tetrahedron T dual to Γ such that the length of a (dual) edge e is given by $\gamma(e)$ [PR68]. Provided the $6j$ symbol is admissible, such a tetrahedron T can always be found uniquely in either $\mathbb{R}^3, \mathbb{R}^2$ or Minkowski space $\mathbb{R}^{2,1}$ [Blu70, Ch.8]. We say the $6j$ -symbol is Euclidean, Plane or Minkowskian depending on the type of T . The type is determined by the sign of the Cayley-Menger determinant of T . Let us be more specific in the Euclidean case. Denote by ϕ_e the exterior dihedral angle of T at edge e .

Theorem 4.2. *Let (Γ, γ) be a Euclidean $6j$ symbol. The sequence $\langle \Gamma, n\gamma \rangle^U$ is of Nilsson type with growth rates, Stokes constants and powers of n and $\log n$ are:*

$$\lambda_{\pm} = e^{i \sum_j \frac{\gamma(e_j) \phi_j}{2}} \quad S_{\pm} = \frac{e^{i \sum_j \frac{\phi_j}{2} + \frac{i\pi}{4}}}{\sqrt{6\pi \text{Vol}(T)}} \quad \alpha = \frac{3}{2} \quad \beta = 0$$

These formulae have been proven in [Rob99]. By analytically continuing the Euclidean formula for dihedral angles in terms of edge lengths, the results can be extended to the Minkowskian case. This will be postponed to a future publication. The Plane case must be different since the volume vanishes in this case. Also any interpretation of the terms in the asymptotic expansion of the $6j$ -symbol beyond the ones just given is very much an open problem [DL09]. This warrants a detailed and exact investigation of the asymptotics of three

representative $6j$ -symbols using the Wilf-Zeilberger method. In this method we compute a recursion for the sequence from which all terms in the asymptotic expansion except for the Stokes constants may be computed.

We have chosen the simplest examples of a Euclidean $6j$ -symbol, a Plane one and a Minkowskian $6j$ -symbol. Their colorings are given by

$$\gamma_{\text{Euclidean}} = (2, 2, 2, 2, 2, 2), \quad \gamma_{\text{Plane}} = (3, 4, 4, 3, 5, 5), \quad \gamma_{\text{Minkowskian}} = (4, 4, 4, 4, 6, 6).$$

Using the unitary evaluation (Definition 4.1) we thus consider the sequences (a_n) , (b_n) and (c_n)

$$\begin{aligned} a_n &:= \langle \triangle, n \gamma_{\text{Euclidean}} \rangle^U = \frac{n!^6}{(3n+1)!^2} \sum_{k=3n}^{4n} \frac{(-1)^k (k+1)!}{(k-3n)!^4 (4n-k)!^3} \\ b_n &:= \langle \triangle, n \gamma_{\text{Plane}} \rangle^U = \frac{n!^2 (2n)!^2 (3n)!^2}{(6n+1)!^2} \sum_{k=6n}^{7n} \frac{(-1)^k (k+1)!}{(k-6n)!^4 (7n-k)! (8n-k)! (9n-k)!} \\ c_n &:= \langle \triangle, n \gamma_{\text{Minkowskian}} \rangle^U = \frac{n!^2 (3n)!^4}{(7n+1)!^2} \sum_{k=7n}^{8n} \frac{(-1)^k (k+1)!}{(k-7n)!^4 (8n-k)! (10n-k)!^2} \end{aligned}$$

In what follows we denote by $\det(C)$ the Cayley-Menger determinant and by K the field generated by the coefficients of the power series $h_{\lambda, \alpha, \beta}$ in the asymptotic expansion. The command

`<< zb.m`

loads the package of [PR] into *Mathematica*. The command

`teucl[n_, k_] := n!^6 / (3n+1)!^2 (-1)^k (k+1)! / ((4n-k)!^3 (k-3n)!^4)`

defines the summand of the sequence (a_n) , and the command

`Zb[teucl[n, k], {k, 3n, 4n}, n, 2]`

computes the following second order linear recursion relation for the sequence (a_n)

$$\begin{aligned} &-9(1+n)(2+3n)^2(4+3n)^2(451+460n+115n^2)a[n] + \\ &(3+2n)(319212+1427658n+2578232n^2+2423109n^3+1255139n^4+340515n^5+37835n^6)a[1+n] - \\ &9(2+n)(5+3n)^2(7+3n)^2(106+230n+115n^2)a[2+n] = 0 \end{aligned}$$

This linear recursion has two formal power series solutions of the form

$$\begin{aligned} a_{\pm, n} &= \frac{1}{n^{3/2}} \Lambda_{\pm}^n \left(1 + \frac{-432 \pm 31i\sqrt{2}}{576n} + \frac{109847 \mp 22320i\sqrt{2}}{331776n^2} + \frac{-18649008 \pm 4914305i\sqrt{2}}{573308928n^3} \right. \\ &+ \frac{14721750481 \pm 45578388960i\sqrt{2}}{660451885056n^4} + \frac{-83614134803760 \pm 7532932167923i\sqrt{2}}{380420285792256n^5} \\ &\left. + \frac{-31784729861796581 \mp 212040612888146640i\sqrt{2}}{657366253849018368n^6} + O\left(\frac{1}{n^7}\right) \right) \end{aligned}$$

where

$$\Lambda_{\pm} = \frac{329 \mp 460i\sqrt{2}}{729} = e^{\mp i6 \arccos(1/3)}$$

are two complex numbers of absolute value 1. Notice the growth rates indeed match the interpretation in terms of dihedral angles of the regular tetrahedron predicted in Theorem 4.2. The coefficients of the formal power series $a_{\pm, n}$ are in the number field $K = \mathbb{Q}(\sqrt{-2})$ and the Cayley-Menger determinant is $\det(C) = 2^5$. More is actually true. Namely, the sequence (a_n) generates two new sequences $(\mu_{+, n})$ and $(\mu_{-, n})$ defined by

$$a_{\pm, n} = \frac{1}{n^{3/2}} \Lambda_{\pm}^n \sum_{l=0}^{\infty} \frac{\mu_{\pm, l}}{n^l}$$

where $\mu_{\pm, 0} = 1$. Each of the sequences $(\mu_{\pm, n})$ are factorially divergent. However, the generating series $\sum_{n=0}^{\infty} z^n \mu_{\pm, n+1}/n!$ are G -functions (as follows from [And00]), and the sequences $(\mu_{\pm, n+1}/n!)$ are of Nilsson

type, with exponential growth rates $\Lambda_{\pm} - \Lambda_{\mp}$. The asymptotics of each sequence $(\mu_{\pm, n+1}/n!)$ gives rise to finitely many new sequences, and so on. All those sequences span a finite dimensional vector space, canonically attached to the sequence (a_n) . This is an instance of resurgence, and is explained in detail in [GM10, Sec.4]. The second order recursion relation for the Plane and the Minkowskian examples has lengthy coefficients, and leads to the following sequences $(b_{\pm, n})$ and $(c_{\pm, n})$

$$\begin{aligned}
b_{+, n} &= \frac{1}{n^{4/3}} \Lambda_+^n \left(1 - \frac{1}{3n} + \frac{3713}{46656n^2} - \frac{25427}{2239488n^3} + \frac{9063361}{17414258688n^4} - \frac{109895165}{104485552128n^5} \right. \\
&\quad \left. + \frac{1927530983327}{2437438960041984n^6} + \dots \right) \\
b_{-, n} &= \frac{1}{n^{5/3}} \Lambda_-^n \left(1 - \frac{37}{96n} + \frac{3883}{46656n^2} - \frac{13129}{4478976n^3} - \frac{5700973}{8707129344n^4} - \frac{14855978561}{3343537668096n^5} \right. \\
&\quad \left. + \frac{2862335448661}{2437438960041984n^6} + \dots \right) \\
c_{\pm, n} &= \frac{1}{n^{3/2}} \Lambda_{\pm}^n \left(1 + \frac{336 \mp 1369\sqrt{2}}{4032n} + \frac{1769489 \mp 831792\sqrt{2}}{1806336n^2} + \frac{67925105712 \mp 66827896993\sqrt{2}}{21849440256n^3} \right. \\
&\quad + \frac{5075437500833257 \mp 2589265090380768\sqrt{2}}{176193886224384n^4} \\
&\quad + \frac{100978405759997442992 \mp 98904713360431641651\sqrt{2}}{552544027199668224n^5} \\
&\quad \left. + \frac{685103512739058526758457 \mp 349782631602887151717776\sqrt{2}}{247539724185451364352n^6} + \dots \right)
\end{aligned}$$

where in the Plane case we have

$$\Lambda_- = \Lambda_+ = -1, \quad K = \mathbb{Q}, \quad \det(C) = 0$$

and in the Minkowskian case we have

$$\begin{aligned}
\Lambda_+ &= \frac{696321931873 - 111529584108\sqrt{2}}{678223072849} = 0.794127\dots \\
\Lambda_- &= \frac{696321931873 + 111529584108\sqrt{2}}{678223072849} = 1.25924\dots \\
K &= \mathbb{Q}(\sqrt{2}), \quad \det(C) = -2^5 3^4.
\end{aligned}$$

Again as in the Euclidean case the growth rates may be interpreted in terms of dihedral angles.

4.2. A conjecture on growth rates. In the special case where (Γ, γ) is an admissible tetrahedron, we have seen a geometric interpretation for the growth rates of $\langle \Gamma, n\gamma \rangle^U$. We can reformulate this more concisely using mean curvature. Recall that for a convex Euclidean polyhedron P in \mathbb{R}^3 the mean curvature is defined by $M(P) = \frac{1}{2} \sum_e \ell_e \phi_e$, where ϕ_e is the exterior dihedral angle at edge e and ℓ_e is the length of the edge. So in the case of the tetrahedron Theorem 4.2 says that if there exists a Euclidean tetrahedron T whose 1-skeleton is dual to Γ and whose edge lengths are given by γ , then the growth rates are given by: $\{e^{\pm iM(T)}\}$. We would like to conjecture that the growth rates of a spin network always include a growth rate corresponding to the mean curvature of the dual polyhedron whenever these terms make sense. For simplicity we formulate the conjecture for planar spin networks only.

Conjecture 4.3. Let (Γ, γ) be a planar spin network with the counterclockwise orientation. Suppose that Γ is 3-connected and that its dual can be viewed as the 1-skeleton of a convex Euclidean polyhedron P whose edge lengths are given by γ . The set of growth rates of the Nilsson type sequence $\langle \Gamma, n\gamma \rangle^U$ contains $e^{\pm iM(P)}$.

By Cauchy's theorem the dual polyhedron P is determined up to isometry by its 1-skeleton and its edge lengths, i.e. by (Γ, γ) . This follows from the fact that P has only triangular faces and is convex.

As a first test of the conjecture we show that it behaves well under the triangle formula on spin networks defined in Section 2.1. In particular this will verify the conjecture for all triangular networks as defined in Section 2.1. Let (Γ, γ) and (Γ', γ') be two spin networks that both satisfy the hypotheses of Conjecture 4.3 and denote their dual polyhedra by P and P' . Furthermore, suppose that (Γ', γ') is obtained from (Γ, γ) by replacing a vertex $v \in \Gamma$ by a triangle. Dually this implies that P can be produced by cutting off a tetrahedron from P' .

Lemma 4.4. If Conjecture 4.3 is true for (Γ, γ) then it is also true for (Γ', γ') .

Proof. Let the labels around the vertex v be a, b, c and call the labels of the edges of new triangle A, B, C as in Figure 2 (lower right) and denote by (\triangle, ψ) the tetrahedron spin network with labels a, b, c, A, B, C that shows up in the triangle formula. This formula shows that

$$\langle \Gamma', n\gamma' \rangle^U = (-1)^{\frac{n(a+b+c)}{2}} \langle \triangle, n\psi \rangle^U \langle \Gamma, n\gamma \rangle^U$$

since the theta only contributes to a sign in the unitary evaluation. We already know Conjecture 4.3 holds for tetrahedra with Euclidean duals, including (\triangle, ψ) . Let us call the dual Euclidean tetrahedron T . Multiplying the asymptotic expansions on the right hand side we see that the growth rates will include

$$(-1)^{\frac{(a+b+c)}{2}} e^{i(M(P)+M(T))} = e^{iM(P')}$$

To see why the equality holds note that we can dissect P' into P and T along the triangle with labels a, b, c that is dual to the vertex v . The minus sign coming from the theta accounts for the fact that we are working with exterior dihedral angles and these add an additional factor of π when comparing the angles in a dissection. \square

The Euclidean volume also appears in the asymptotic expansion of the tetrahedral spin network, as part of the Stokes constants, see Section 4.1. However this does not generalize well to larger networks since the volumes do not add under the triangle formula. In the appendix we will see a less trivial confirmation of the above conjecture for the cube spin network.

5. CHALLENGES AND FUTURE DIRECTIONS

In this section we list some challenges and future directions. Our first problem concerns a bound on the unitary evaluations.

Problem 5.1. Show that the unitary evaluation of a spin network (Γ, γ) satisfies

$$|\langle \Gamma, \gamma \rangle^U| \leq 1$$

This problem may be solved using unitarity and locality in a way similar to the proof that the Reshetikhin-Turaev invariants of a closed 3-manifold grow at most polynomially with respect to the level; see [Gar98, Thm.2.2]. Our next problem is a version of the Volume Conjecture for classical spin networks with all edges colored by 2. Problem 5.1 also suggests that the growth rates must be ≤ 1 . In case $\gamma = 2n$ more seems to be true.

Problem 5.2. The growth rates of the sequence $\langle \Gamma, 2n \rangle^U$ are on the unit circle.

A positive solution to this problem is known for the following ribbon graphs: the Θ , the tetrahedron, the 3-faced prism and more generally for the infinite family of drums; see [Abd10]. More generally one may pose the following

Problem 5.3. Give a geometric meaning to the set of growth rates of a spin network.

We have formulated Conjecture 4.3 as a partial answer to this question but that concerns only a single special growth rate among many. Along the same lines one may ask for an interpretation of the rest of the asymptotic expansion. Looking at the case of the $6j$ -symbol it seems reasonable to consider the number field $K_{\Gamma, \gamma}$ generated by the coefficients of the power series $h_{\lambda, \alpha, \beta}$ in the Nilsson type asymptotic expansion of $\langle \Gamma, n\gamma \rangle^U$.

Problem 5.4. Give a geometric interpretation of the number field $K_{\Gamma,\gamma}$ of a spin network (Γ, γ) .

Also the Stokes constant might have a geometric meaning as it does in the case of the tetrahedron.

Problem 5.5. Give a geometric meaning to the Stokes constants of the sequence $\langle \Gamma, n\gamma \rangle^U$.

The next problem is a computational challenge to all the known asymptotic methods, and shows their practical limitations.

Problem 5.6. Compute the asymptotics of the evaluation $\langle K_{3,3}, 2n \rangle$ (given explicitly in Proposition 2.5) and $\langle \text{Cube}, 2n \rangle$.

The next problem is formulated by looking at the examples from Section 4.1.

Problem 5.7. Prove that for every coloring γ of the tetrahedron spin network (\triangle, γ) , the sequence $\langle \triangle, n\gamma \rangle$ satisfies a second order recursion relation with coefficients polynomials in n . Can you compute the coefficients of this recursion from γ alone?

Let us end this section with a remark. The main results of our paper can be extended to evaluations of spin networks corresponding to higher rank Lie groups. This will be discussed in a later publication.

APPENDIX A. ASYMPTOTICS OF THE REGULAR CUBE

by Don Zagier

We give the asymptotic expansion of the standard evaluation a_n of the 1-skeleton of the 3-dimensional cube, with all edges colored by $2n$. Proposition 2.4 implies that (a_n) is given by

$$(23) \quad a_n = \sum_{k=0}^{2n} (2k+1) a_{n,k}^4$$

with

$$a_{n,k} = \sum_j (-1)^j \binom{k}{j-3n}^2 \binom{2n-k}{4n-j} \binom{j+1}{2n+k+1},$$

making it clear that the numbers (a_n) are integral and positive. The first few values of a_n are given by

$$\begin{aligned} a_0 &= 1, \\ a_1 &= 6144, \\ a_2 &= 505197000, \\ a_3 &= 7741440000000, \\ a_4 &= 1393762029660000000, \\ a_5 &= 3685480142898164744060928, \\ a_6 &= 1038107879077276408534853271552, \\ a_7 &= 29722354754825775224492840550400000, \\ a_8 &= 104193297934159421485149830847575156250000, \\ a_9 &= 3557731603525300009641567861059837904000000000, \\ a_{10} &= 12357485751601160513255660198337121351402277161410240. \end{aligned}$$

We first look for a recursion of the form

$$(24) \quad \sum_{j=0}^J c_j(n) a_{n+j} = 0$$

with J not too large and $c_j(n)$ polynomials of n of some not too large degree d . Using the first few hundred values of (a_n) , we find experimentally a recursion of this form with $J = 4$, $d = 61$ and with $c_j(n)$ given by

$$\begin{aligned} c_0(n) &= 3^{16} (2n+7) (3n+2)^8 (3n+4)^8 (3n+5)^7 (3n+7)^7 (3n+8) (3n+10) (4n+3) (4n+5) P_0(n+3), \\ c_1(n) &= -2 \cdot 3^8 (n+1)^5 (2n+3) (2n+7) (3n+5)^7 (3n+7)^7 (3n+8) (3n+10) P_1(n), \\ c_2(n) &= -2 \cdot 3^4 (n+1)^5 (n+2)^7 (2n+5) (3n+8) (3n+10) P_2(n), \\ c_3(n) &= -2 \cdot (n+1)^5 (n+2)^7 (n+3)^9 (2n+3) (2n+7) P_1(-n-5), \\ c_4(n) &= (n+1)^5 (n+2)^7 (n+3)^9 (n+4)^{11} (2n+3) (4n+15) (4n+17) P_0(n+2), \end{aligned}$$

where P_0 , P_1 and P_2 are irreducible polynomials (normalized to have integral coefficients with no common factor) with leading terms

$$\begin{aligned} P_0(n) &= 2^{11} 3^7 5^5 7 \cdot 23^5 47 (n^{26} + O(n^{24})), \\ P_1(n) &= 2^{15} 3^7 5^5 7^3 23^5 47^3 (n^{38} + 94n^{37} + O(n^{36})), \\ P_2(n) &= 2^{15} 3^{13} 5^5 7 \cdot 19 \cdot 23^5 47 \cdot 71 \cdot 73 (n^{46} + 115n^{45} + O(n^{44})) \end{aligned}$$

as $n \rightarrow \infty$, and with the polynomial P_0 being even. The full values are given at the end of the appendix. To analyze the asymptotics of the solutions of the above recursion, we will use the standard Frobenius theory (see e.g. [Mil06, Olv97, Was87, WZ85]). If C_j denotes the top coefficient of the polynomial $c_j(n)$, then we find that $\sum_{j=0}^4 C_j \lambda^j$ factors as $(\lambda - 3^{12})^2 (\lambda - (1 + \sqrt{-2})^{24}) (\lambda - (1 - \sqrt{-2})^{24})$ and that the indicial equation of the root 3^{12} has a double root at $-9/2$, while the indicial equations of the roots $(1 \pm \sqrt{-2})^{24}$ both have root -4 . This implies that (a_n) has an asymptotic expansion

$$(25) \quad a_n \sim S_0 \frac{3^{12n}}{n^4} \left((\log n + c) M_1\left(\frac{1}{n}\right) + M_2\left(\frac{1}{n}\right) \right) + \Re \left(S_1 \frac{(1 + \sqrt{-2})^{24n}}{n^{9/2}} M_3\left(\frac{1}{n}\right) \right)$$

for some constants $S_0, c \in \mathbb{R}$, $S_1 \in \mathbb{C}$ and power series $M_1(x)$, $M_2(x) \in \mathbb{Q}[[x]]$, $M_3(x) \in \mathbb{Q}[\sqrt{-2}][[x]]$, normalized by requiring that M_1 and M_3 have constant term 1. Notice that the three roots 3^{12} , $(1 \pm \sqrt{-2})^{24}$ have the same absolute value, so that the different terms of this expansion all have the same order of magnitude up to powers of n . Using the acceleration method¹ described at [Zag01, p.954] and in Section 4 of [GM08], applied to the values of a_n for $n = 1000, \dots, 1050$, we find the numerical values of the constants S_j and c and the first few coefficients of the power series M_i . The former are then recognized as

$$(26) \quad S_0 = \frac{3^5}{24\pi^6}, \quad c = \frac{7}{4} \log 2 + \log 3 + \gamma, \quad S_1 = \frac{(1+i)(1+\sqrt{-2})^{12}}{2^{31/4} \pi^{11/2}},$$

and the latter as

$$\begin{aligned} M_1(x) &= 1 - \frac{14}{9}x + \frac{419}{324}x^2 - \frac{5659}{8748}x^3 + \frac{84769}{629856}x^4 \dots, \\ M_2(x) &= \frac{1}{2}x - \frac{689}{864}x^2 + \frac{4771}{7776}x^3 - \frac{3799441}{22394880}x^4 + \dots, \\ M_3(x) &= 1 - \frac{2080 - 43\sqrt{-2}}{1152}x + \frac{1985023 - 114208\sqrt{-2}}{1327104}x^2 + \dots \end{aligned}$$

The acceleration method can give many more terms, but it is easier to simply substitute the Ansatz (25) into the recursion for the a_n , thus obtaining as many terms as desired. The approximation works very well in practice, e.g., the maximal relative error between a_n and the right-hand side of (25) with 50 terms of the power series $M_j(1/n)$ is about one part in 10^{105} for n between 900 and 1000. To first order, the above formulas say that the asymptotics of a_n are given by

$$a_n \sim 3^{12n+5} \frac{\log(2^{7/4}3n) + \gamma + O(1/\sqrt{n})}{\pi^6 (2n)^4}.$$

¹This method is equivalent to the Richardson transform, explained in detail in [GIKM, Sec.5.2], and also in [BO99].

We end by giving the complete values of the polynomials $P_j(n)$ that appear in the recursion relation:

$$\begin{aligned}
P_0(n) = & 29639019676089600000 n^{26} - 150687090646682256000 n^{24} + 306650022810104871540 n^{22} \\
& - 331831776907297971277 n^{20} + 219414205267920364521 n^{18} - 96826696589802950226 n^{16} \\
& + 29683042452642732342 n^{14} - 6233837158945489065 n^{12} + 868763697226715493 n^{10} - 77173811768742984 n^8 \\
& + 4094153904684504 n^6 - 111886799053248 n^4 + 797085625600 n^2 + 17508556800 ,
\end{aligned}$$

$$\begin{aligned}
P_1(n) = & 51330514060153830297600000 n^{38} + 4825068321654460047974400000 n^{37} \\
& + 219957552931873414824036864000 n^{36} + 6478694077195677171946040064000 n^{35} \\
& + 138596018058877517667573746466240 n^{34} + 2295022658488679405177615124025920 n^{33} \\
& + 30614929984046498162519595722728508 n^{32} + 338072087836667419737764233439922530 n^{31} \\
& + 3151590998989517431768295237323718623 n^{30} + 25169023605885819585932158912744414906 n^{29} \\
& + 174146716308878486922546565722791225448 n^{28} + 1053195250756920493731804102357697945572 n^{27} \\
& + 5606361750518381240594997946959656401095 n^{26} + 26414736794861925209673962053754850002124 n^{25} \\
& + 110642699366898526542975775645886257667832 n^{24} + 413447345600050228521136991970449404260966 n^{23} \\
& + 1381980145537336658418260176761712507602933 n^{22} + 4140268295003002172648827386155584658850114 n^{21} \\
& + 11132423733718852590472537735822272877436592 n^{20} + 26885849594024541421613060268809580068669016 n^{19} \\
& + 58334970199614352499186715966601305101299773 n^{18} + 113675657006866049496120543251199160823538984 n^{17} \\
& + 198774677991170902825182509342222362759066932 n^{16} + 311443610656870193242629490576780944439111836 n^{15} \\
& + 436334419544283264503767964716530868856380648 n^{14} + 545097336381579864877890591441121830311242864 n^{13} \\
& + 605044458481431111735014250352650750979996544 n^{12} + 594007313574579683774145689072368182659197376 n^{11} \\
& + 512886129222805060276163096656047277821413760 n^{10} + 386709368743514690018446501443764021880730368 n^9 \\
& + 252342374226937131766477472379392715246649344 n^8 + 140888462647571785365811030970706748098760704 n^7 \\
& + 66312852042204808325857346405562441033596928 n^6 + 25796733254687036537088890539848097231134720 n^5 \\
& + 8069974595385074631605661061376909102284800 n^4 + 1950251347843211319569463651786279088128000 n^3 \\
& + 341555150844826683309630400427989401600000 n^2 + 38554163497112285346472887524366745600000 n \\
& + 2104728968892765569954334578933760000000 ,
\end{aligned}$$

$$\begin{aligned}
P_2(n) = & 34044436942851501889228800000 n^{46} + 3915110248427922717261312000000 n^{45} \\
& + 219662706791565782848926392832000 n^{44} + 8013067972307054904678991211520000 n^{43} \\
& + 213693214418431619419298087215485120 n^{42} + 4441366173640824819720536004281937600 n^{41} \\
& + 74894384718928871397218606262165524844 n^{40} + 1053306557874097263334396534371665684400 n^{39} \\
& + 12603933341218324699775023159567066766967 n^{38} + 130270526199929371052793324163523141276865 n^{37} \\
& + 1176678761157600477488177183321806479261195 n^{36} + 9375125430920826953262133708501549064882175 n^{35} \\
& + 66383537873561799570407105177080643368970738 n^{34} + 420307309950545627790271278025374052312931480 n^{33} \\
& + 2391581100396727961084015883784259869439400176 n^{32} + 12280714697778331715472758947513061657620945580 n^{31} \\
& + 57105561410677181109714147899130291409292461050 n^{30} + 241147151880727945569092590718565623964677498750 n^{29} \\
& + 926919151084739302148924528551196602048329414090 n^{28} + 3249130587428846232389342739110375794038903230050 n^{27} \\
& + 10401395584522433137223729045847274251204938831280 n^{26} + 30443078183785042173235936392545801110737278600700 n^{25} \\
& + 81523471295041101121702148739822166982527381249680 n^{24} + 199828567976168135158731196334605133971159536918300 n^{23} \\
& + 448394647578816808473555576099796146381767002557015 n^{22} + 920880328621349858197546061838422939683875830250825 n^{21} \\
& + 1730060383317082970163176773077295571528776228823811 n^{20} + 2970768265449411986606322605605748766920711781281175 n^{19} \\
& + 4657049644272293081174713334539070013552116895070038 n^{18} + 6654448413726919814684796853369712557012807014275460 n^{17} \\
& + 8650083204510826813891208265741376798744454496660220 n^{16} + 10204350936081623227151423121609083529329974289414800 n^{15} \\
& + 10892578929911563608834673775127900025242488747892352 n^{14} + 10483723584809889429623667272318319527058201296732320 n^{13} \\
& + 9059104715506400545606048153659029802430897952414784 n^{12} + 6992047378850409545281115023066707881078577900939520 n^{11} \\
& + 4790327226399433431900479655636468590589446522365440 n^{10} + 2891085435841974142824330005531725765542027596096000 n^9 \\
& + 1522663617068026467557018527073299284644425562419200 n^8 + 691604038146644417153727841870625107738881053184000 n^7 \\
& + 266822516520374901619621632554668192771563340800000 n^6 + 85697330246713152918864338753344668302045030400000 n^5 \\
& + 22287663074878779648397028175676823501117952000000 n^4 + 450792626594235022750902319082697990289920000000 n^3 \\
& + 66504589562492929383059851236268484198400000000 n^2 + 6363528251174783559928067350587667660800000000 n \\
& + 296294973650466076876070759346376704000000000 .
\end{aligned}$$

APPENDIX B. FURTHER COMMENTS ON THE ASYMPTOTICS OF THE REGULAR CUBE

The guessed recursion relation of the sequence (a_n) from the previous section agrees with the result of the independent guessing program `Guess` of Kauers; [Kau09a, Kau09b]. The recursion for (a_n) was verified for $n = 0, \dots, 2996$, where the height (i.e., the number of digits) of a_{3000} is 17162. On the other hand, the coefficients of the polynomials $c_k(n)$ are integers with a much smaller height 73. In addition, the root λ_1 of the characteristic polynomial can be written in the form:

$$\lambda_1 = (1 + i\sqrt{2})^{24} = 3^{12} e^{12i \arccos(-1/3)}$$

where $e^{12i \arccos(-1/3)}$ is the exponentiated total mean curvature of the regular Euclidean octahedron (dual to the regular Euclidean cube) with unit sides. This confirms Conjecture 4.3 on the asymptotics of evaluations of classical spin networks. The factor 3^{12} comes from the fact that we are considering the standard normalization and not the unitary one as is done in the Conjecture.

The asymptotic expansion (25) is clearly of Nilsson type, with the presence of logarithms, and Stokes constants which are no longer algebraic, up to rational powers of π . This makes it unlikely that stationary phase type methods will be able to obtain the asymptotic expansion for the regular cube.

REFERENCES

- [Abd10] Abdelmalek Abdesselam, *On the volume conjecture for classical spin networks*, arXiv/1009.2811, 2010.
- [Ada94] Colin C. Adams, *The knot book*, W. H. Freeman and Company, New York, 1994, An elementary introduction to the mathematical theory of knots.
- [AHH⁺09] Vincenzo Aquilanti, Hal M. Haggard, Austin Hedeman, Nadir Jeevanjee, Robert G. Littlejohn, and Liang Yu, *Semiclassical mechanics of the Wigner 6j-symbol*, arXiv/0904.1734, 2009.
- [And00] Yves André, *Séries Gevrey de type arithmétique. I. Théorèmes de pureté et de dualité*, Ann. of Math. (2) **151** (2000), no. 2, 705–740.

- [BK08] Spencer Bloch and Dirk Kreimer, *Mixed Hodge structures and renormalization in physics*, Commun. Number Theory Phys. **2** (2008), no. 4, 637–718.
- [BL81a] L. C. Biedenharn and J. D. Louck, *Angular momentum in quantum physics*, Encyclopedia of Mathematics and its Applications, vol. 8, Addison-Wesley Publishing Co., Reading, Mass., 1981, Theory and application, With a foreword by Peter A. Carruthers.
- [BL81b] Lawrence C. Biedenharn and James D. Louck, *The Racah-Wigner algebra in quantum theory*, Encyclopedia of Mathematics and its Applications, vol. 9, Addison-Wesley Publishing Co., Reading, Mass., 1981, With a foreword by Peter A. Carruthers, With an introduction by George W. Mackey.
- [Blu70] Leonard M. Blumenthal, *Theory and applications of distance geometry*, Second edition, Chelsea Publishing Co., New York, 1970.
- [BO99] Carl M. Bender and Steven A. Orszag, *Advanced mathematical methods for scientists and engineers. I*, Springer-Verlag, New York, 1999, Asymptotic methods and perturbation theory, Reprint of the 1978 original.
- [CFS95] J. Scott Carter, Daniel E. Flath, and Masahico Saito, *The classical and quantum 6j-symbols*, Mathematical Notes, vol. 43, Princeton University Press, Princeton, NJ, 1995.
- [CM11] Francesco Costantino and Julien Marché, *Generating series and asymptotics of classical spin networks*, arXiv:1103.5644, 2011.
- [Cos09] Francesco Costantino, *Integrality of kauffman brackets of trivalent graphs*, arXiv:0908.0542, 2009.
- [DL09] Maïé Dupuis and Etera R. Livine, *Pushing the asymptotics of the 6j-symbol further*, Phys. Rev. D **80** (2009), no. 2, 024035.
- [EPR08] Jonathan Engle, Roberto Pereira, and Carlo Rovelli, *Flipped spinfoam vertex and loop gravity*, Nuclear Phys. B **798** (2008), no. 1-2, 251–290.
- [Gar98] Stavros Garoufalidis, *Applications of quantum invariants in low-dimensional topology*, Topology **37** (1998), no. 1, 219–224.
- [Gar09] ———, *G-functions and multiset versus holonomic sequences*, Adv. Math. **220** (2009), no. 6, 1945–1955.
- [Gar11] ———, *What is a sequence of Nilsson type?*, Interactions between hyperbolic geometry, quantum topology and number theory, Contemp. Math., vol. 541, Amer. Math. Soc., Providence, RI, 2011, pp. 145–157.
- [GIKM] Stavros Garoufalidis, A. Its, Andrei Kapaev, and Marcos Mariño, *Asymptotics of the instantons of painlevé i* , arXiv:1002.3634, Preprint 2010.
- [GM08] Daniel B. Grünberg and Pieter Moree, *Sequences of enumerative geometry: congruences and asymptotics*, Experiment. Math. **17** (2008), no. 4, 409–426, With an appendix by Don Zagier.
- [GM10] Stavros Garoufalidis and Marcos Mariño, *Universality and asymptotics of graph counting problems in non-orientable surfaces*, J. Combin. Theory Ser. A **117** (2010), no. 6, 715–740.
- [GvdV11] Stavros Garoufalidis and Roland van der Veen, *Asymptotics of quantum spin networks at a fixed root of unity*, Mathematische Annalen (2011), 1–26, 10.1007/s00208-011-0662-3.
- [HL10] H. M. Haggard and R. G. Littlejohn, *Asymptotics of the wigner 9j-symbol*, Classical and Quantum Gravity **27** (2010), 135010.
- [Jon87] V. F. R. Jones, *Hecke algebra representations of braid groups and link polynomials*, Ann. of Math. (2) **126** (1987), no. 2, 335–388.
- [Kas97] Rinat Kashaev, *The hyperbolic volume of knots from the quantum dilogarithm*, Lett. Math. Phys **39** (1997), 296–275.
- [Kau09a] Manuel Kauers, *Guessing handbook*, Tech. Report 09-07, RISC Report Series, Johannes Kepler University Linz, Austria, 2009.
- [Kau09b] ———, *Mathematica software*, 2009, <http://www.risc.uni-linz.ac.at/research/combinat/software/Guess/>.
- [KL94] Louis H. Kauffman and Sóstenes L. Lins, *Temperley-Lieb recoupling theory and invariants of 3-manifolds*, Annals of Mathematics Studies, vol. 134, Princeton University Press, Princeton, NJ, 1994.
- [KR89] A. N. Kirillov and N. Yu. Reshetikhin, *Representations of the algebra $U_q(\mathfrak{sl}(2))$, q -orthogonal polynomials and invariants of links*, Infinite-dimensional Lie algebras and groups (Luminy-Marseille, 1988), Adv. Ser. Math. Phys., vol. 7, World Sci. Publ., Teaneck, NJ, 1989, pp. 285–339.
- [LY11] Robert G. Littlejohn and Liang Yu, *Semiclassical analysis of the wigner 9j-symbol with small and large angular momenta*, arXiv/1104.1499, To appear in Phys. Rev. A, 2011.
- [Mil06] Peter D. Miller, *Applied asymptotic analysis*, Graduate Studies in Mathematics, vol. 75, American Mathematical Society, Providence, RI, 2006.
- [MM01] Jun Murakami and Hitoshi Murakami, *The colored jones polynomials and the simplicial volume of a knot*, Acta Math. **186** (2001), 85–104.
- [Mou79] John P. Moussouris, *The chromatic evaluation of strand networks*, Advances in Twistor Theory (Ward Huston, ed.), Research Notes in Mathematics, Pitman, 1979, pp. 308–312.
- [Olv97] Frank W. J. Olver, *Asymptotics and special functions*, AKP Classics, A K Peters Ltd., Wellesley, MA, 1997, Reprint of the 1974 original [Academic Press, New York; MR0435697 (55 #8655)].
- [Pen71a] Roger Penrose, *Angular momentum: An approach to combinatorial space time*, Quantum Theory and Beyond (Ted Bastin, ed.), Research Notes in Mathematics, Cambridge University Press, 1971, pp. 151–180.
- [Pen71b] ———, *Applications of negative dimensional tensors*, Combinatorial Mathematics and its Applications (Proc. Conf., Oxford, 1969), Academic Press, London, 1971, pp. 221–244.

- [PR] Peter Paule and Axel Riese, *qZeil Mathematica software*, <http://www.risc.uni-linz.ac.at>.
- [PR68] G. Ponzano and T. Regge, *Semiclassical limit of racah coefficients*, Spectroscopic and group theoretical methods in physics, Research Notes in Mathematics, North Holland, 1968, pp. 1–58.
- [Rac42a] Giulio Racah, *Theory of complex spectra II*, Physical Rev. (2) **61** (1942), 186–197.
- [Rac42b] ———, *Theory of complex spectra III*, Physical Rev. (2) **62** (1942), 438–462.
- [Rac46] ———, *Theory of complex spectra I*, Physical Rev. (2) **61** (1946), 186–197.
- [Rac49] ———, *Theory of complex spectra IV*, Physical Rev. (2) **76** (1949), 1352–1365.
- [Rob99] Justin Roberts, *Classical 6j-symbols and the tetrahedron*, Geom. Topol. **3** (1999), 21–66 (electronic).
- [Rob02] ———, *Asymptotics and 6j-symbols*, Invariants of knots and 3-manifolds (Kyoto, 2001), Geom. Topol. Monogr., vol. 4, Geom. Topol. Publ., Coventry, 2002, pp. 245–261 (electronic).
- [RS95] Carlo Rovelli and Lee Smolin, *Discreteness of area and volume in quantum gravity*, Nuclear Phys. B **442** (1995), no. 3, 593–619.
- [Sie29] C.L. Siegel, *Über einige anwendungen diophantischer approximationen*, Tech. Report 1, Abh. Preuss. Akad. Wiss., 1929, 1–70. Reprinted in *Gesammelte Abhandlungen*, vol. 1, no 16 (1966) 209–266.
- [Tur94] V. G. Turaev, *Quantum invariants of knots and 3-manifolds*, de Gruyter Studies in Mathematics, vol. 18, Walter de Gruyter & Co., Berlin, 1994.
- [TV92] V. G. Turaev and O. Ya. Viro, *State sum invariants of 3-manifolds and quantum 6j-symbols*, Topology **31** (1992), no. 4, 865–902.
- [VMK88] D. A. Varshalovich, A. N. Moskalev, and V. K. Khersonskii, *Quantum theory of angular momentum*, World Scientific Publishing Co. Inc., Teaneck, NJ, 1988, Irreducible tensors, spherical harmonics, vector coupling coefficients, $3n_j$ symbols, Translated from the Russian.
- [Was87] Wolfgang Wasow, *Asymptotic expansions for ordinary differential equations*, Dover Publications Inc., New York, 1987, Reprint of the 1976 edition.
- [Wes98] Bruce W. Westbury, *A generating function for spin network evaluations*, Knot theory (Warsaw, 1995), Banach Center Publ., vol. 42, Polish Acad. Sci., Warsaw, 1998, pp. 447–456.
- [Whi33] Hassler Whitney, *2-isomorphic graphs*, Amer. J. Math. **55** (1933), 245–254.
- [Wig41] Eugene P. Wigner, *On representations of certain finite groups*, Amer. J. Math. **63** (1941), 57–63.
- [WZ85] Jet Wimp and Doron Zeilberger, *Resurrecting the asymptotics of linear recurrences*, J. Math. Anal. Appl. **111** (1985), no. 1, 162–176.
- [Zag01] Don Zagier, *Vassiliev invariants and a strange identity related to the Dedekind eta-function*, Topology **40** (2001), no. 5, 945–960.

SCHOOL OF MATHEMATICS, GEORGIA INSTITUTE OF TECHNOLOGY, ATLANTA, GA 30332-0160, USA
<http://www.math.gatech.edu/~stavros>
E-mail address: stavros@math.gatech.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, BERKELEY, BERKELEY, CA 94720-3840, USA
<http://www.math.berkeley.edu/~roland>
E-mail address: roland@math.berkeley.edu