

IDEMPOTENT CHARACTERS AND EQUIVARIANTLY MULTIPLICATIVE SPLITTINGS OF K-THEORY

BENJAMIN BÖHME

ABSTRACT. We classify the primitive idempotents of the p -local complex representation ring of a finite group G in terms of the cyclic subgroups of order prime to p and show that they all come from idempotents of the Burnside ring. Our results hold without adjoining roots of unity or inverting the order of G , thus extending classical structure theorems. We then derive explicit group-theoretic obstructions for tensor induction to be compatible with the resulting idempotent splitting of the representation ring Mackey functor.

Our main motivation is an application in homotopy theory: we conclude that the idempotent summands of G -equivariant topological K -theory and the corresponding summands of the G -equivariant sphere spectrum admit exactly the same flavors of equivariant commutative ring structures, made precise in terms of Hill-Hopkins-Ravenel norm maps.

1. INTRODUCTION

The purpose of this paper is twofold: We first classify the primitive idempotents in the real and complex representation rings $RO(G)$ and $RU(G)$ of a finite group G and their local variants, as summarized in §1.1, extending various classical results. We then study the compatibility of tensor induction with the splittings of $RO(G)$ and $RU(G)$ into idempotent summands, and as a consequence obtain an explicit description of the G -equivariant commutative ring spectrum structures occurring as idempotent summands of real and complex G -equivariant topological K -theory. See §1.2 for a summary of these results.

We begin with some motivation. Multiplicative induction is a familiar tool in representation theory and group cohomology. In the wake of Hill, Hopkins and Ravenel's ground-breaking solution to the Kervaire invariant one problem [HHR16], it has also received much interest in equivariant homotopy theory. Starting from the observation that localization can destroy some of the structure of an equivariant commutative

2000 *Mathematics Subject Classification.* 19L47; 19A22, 20C15, 55P43, 55P60, 55P91, 55S91.

Key words and phrases. Equivariant stable homotopy theory, Hill-Hopkins-Ravenel norm, equivariant commutative ring spectrum, topological K -theory, representation ring, idempotent, multiplicative induction, Tambara functor.

ring spectrum, Hill and Hopkins [HH14] gave a necessary and sufficient criterion (cf. Proposition 4.4) for the localization

$$R[x^{-1}] := \operatorname{hocolim} \left(R \xrightarrow{x} S^{-V} \wedge R \xrightarrow{x} S^{-(V \oplus V)} \wedge R \xrightarrow{x} \dots \right)$$

of a G - E_∞ ring spectrum R at an element $x \in \pi_V^G(R)$ to admit a G - E_∞ ring structure. The critical part is that $R[x^{-1}]$ might not admit Hill-Hopkins-Ravenel *norm maps*

$$N_K^H : G_+ \wedge_H \bigwedge_{H/K} \operatorname{Res}_K^G(R) \rightarrow R$$

for all nested subgroups $K \leq H \leq G$. Subsequently, more general notions of equivariant commutative ring spectra equipped with incomplete collections of norm maps, called N_∞ *ring spectra*, were studied by Blumberg and Hill in [BH15], [BH18] and [BH20].

Interesting examples of equivariant localizations arise from primitive¹ idempotent elements $e \in \pi_0^G(R)$. These induce a decomposition of the homotopy Mackey functor $\underline{\pi}_*(R)$ into indecomposable summands (also called *blocks*) of the form

$$e \cdot \underline{\pi}_*(R) \cong \underline{\pi}_*(R)[e^{-1}]$$

and hence yield a block decomposition of R as a wedge of G -spectra $R[e^{-1}]$. One can now ask about the possible N_∞ ring structures on these blocks. Hill and Hopkins' aforementioned criterion involves checking relations involving multiplicative induction in $\pi_0^G(R)$, which in general are hard to access.

Problem 1.1. Determine the nested subgroups $K \leq H \leq G$ such that

- (1) the norm map N_K^H for R descends to a well-defined norm map²

$$\tilde{N}_K^H : G_+ \wedge_H \bigwedge_{H/K} \operatorname{Res}_K^G(R[e^{-1}]) \rightarrow R[e^{-1}]$$

on the block of R defined by the primitive idempotent $e \in \pi_0^G(R)$

- (2) the induced norm operation on homotopy groups $N_K^H : \pi_0^K(R) \rightarrow \pi_0^H(R)$ descends to a well-defined norm operation

$$\tilde{N}_K^H : \pi_0^K(R[e^{-1}]) \rightarrow \pi_0^H(R[e^{-1}]).$$

In the prequel [Böh19], the author gave an explicit group-theoretical answer in the fundamental example of the G -equivariant sphere spectrum S . It built on an analysis of

¹An idempotent is *primitive* if it cannot be written as a sum of non-zero idempotents.

²Throughout the paper, we write \tilde{N} for the norms of a localization to distinguish them from the norms of the original object.

multiplicative induction in the Burnside ring $A(G)$ and Segal's identification $\pi_0^G(\mathbb{S}) \cong A(G)$ [Seg71].

In the present paper, we present a complete solution to Problem 1.1 for G -equivariant complex topological K-theory KU_G and its real analogue KO_G . The homotopy groups

$$\pi_0^G(KU_G) \cong RU(G), \quad \pi_0^G(KO_G) \cong RO(G)$$

identify with the complex and real representation ring $RU(G)$ and $RO(G)$, respectively, see e.g. [Seg68, §2].

1.1. Primitive idempotents in representation rings. Dress' classification of primitive idempotents in the Burnside ring and its local variants [Dre69] was the starting point for the investigation of the idempotent splittings of $A(G)$ and \mathbb{S} in [Böh19]. Given a collection P of prime numbers, write $A(G)_{(P)} := A(G) \otimes \mathbb{Z}_{(P)}$ for the P -local Burnside ring, where $\mathbb{Z}_{(P)} := \mathbb{Z}[p^{-1} \mid p \notin P]$. Dress showed that the primitive idempotent elements $e_L \in A(G)_{(P)}$ are in canonical bijection with the conjugacy classes of P -perfect subgroups $L \leq G$. See § 2.1 for further details.

It is known that the complex representation ring $RU(G)$ has no idempotents other than zero or one, see [Ser77, §11.4, Corollary]. We extend this result to a classification of the primitive idempotents in the P -local representation ring $RU(G)_{(P)} := RU(G) \otimes \mathbb{Z}_{(P)}$ as follows. Consider the "linearization" map

$$\text{lin}: A(G)_{(P)} \rightarrow RU(G)_{(P)}$$

given by sending a finite G -set to its associated permutation representation.

Theorem 1.2. *The assignment $C \mapsto \text{lin}(e_C)$ defines a bijection between the conjugacy classes of cyclic subgroups $C \leq G$ of order not divisible by any prime in P and the primitive idempotent elements of the ring $RU(G)_{(P)}$. Here, $e_C \in A(G)_{(P)}$ denotes Dress' idempotent associated to C , see Theorem 2.5.*

Theorem 1.2 is an instance of the phenomenon that one passes from the Burnside ring to the representation ring by restricting attention to cyclic subgroups. The proof is given in §2.

Remark 1.3. Theorem 1.2 extends classical work in the following way: Building on work by Solomon [Sol67], Gluck [Glu81a] studies the idempotents $\text{lin}(e_C)$ and their character values in the rational and the p -local case for a single prime p , but does not show that they are primitive. He also observes that Dress' idempotent $e_L \in A(G)_{(P)}$ is in the kernel of the linearization map if L is not a cyclic group; we prove this in the general P -local case in Corollary 2.7.

We record an immediate consequence of Theorem 1.2. Write $RO(G)_{(P)}$ for the P -local real representation ring and $RQ(G) \otimes \mathbb{Z}_{(P)}$ for the ring of $\mathbb{Z}_{(P)}$ -linear combinations of G -representations over the rational numbers. It is well-known that these embed into $RU(G)_{(P)}$ as subrings.

Corollary 1.4. *The primitive idempotents of $RU(G)_{(P)}$ all lie in the subrings $RO(G)_{(P)}$ and $RQ(G) \otimes \mathbb{Z}_{(P)}$. Hence, they are precisely the primitive idempotents of these subrings.*

In the special case of $RQ(G) \otimes \mathbb{Q}$, this result appeared as [Sol67, Thm. 3].

1.2. Multiplicativity of idempotent summands. We now turn to the multiplicative properties of the idempotent splittings of the complex and real representation rings and equivariant K -theory spectra. Since the block $RU(G)_{(P)}[lin(e_C)^{-1}]$ agrees with the $A(G)_{(P)}$ -module localization

$$RU(G)_{(P)}[e_C^{-1}] \cong RU(G)_{(P)} \otimes_{A(G)_{(P)}} A(G)_{(P)}[e_C^{-1}],$$

we obtain an identification

$$(KU_G)_{(P)}[lin(e_C)^{-1}] \simeq (KU_G)_{(P)} \wedge \mathbb{S}_{(P)}[e_C^{-1}]$$

of the blocks of P -local G -equivariant K -theory with an e_C -localization in genuine G -spectra. By Corollary 1.4, the same is true for $RO(G)_{(P)}$ and $(KO_G)_{(P)}$. This enables us to reduce the solution to Problem 1.1 for equivariant K -theory to the one for the sphere given in the prequel [Böh19]. The resulting classification of the maximal N_∞ ring structures of the idempotent summands of $(KU_G)_{(P)}$ can be summarized as follows:

Theorem 1.5. *Let $C \leq G$ be a cyclic group of order not divisible by any prime in P and let e_C be the corresponding primitive idempotent in $A(G)_{(P)}$. Let $K \leq H \leq G$ be nested subgroups. Then the following are equivalent:*

- (a) *The G -spectrum $\mathbb{S}_{(P)}[e_C^{-1}]$ inherits a norm map \tilde{N}_K^H from the norm map N_K^H of $\mathbb{S}_{(P)}$.*
- (b) *The G -spectrum $(KU_G)_{(P)}[e_C^{-1}]$ inherits a norm map \tilde{N}_K^H from that of $(KU_G)_{(P)}$.*
- (c) *The Mackey functor $A(-)_{(P)}[e_C^{-1}]$ inherits a norm map \tilde{N}_K^H from that of $A(-)_{(P)}$.*
- (d) *The Mackey functor $RU(-)_{(P)}[e_C^{-1}]$ inherits a norm map \tilde{N}_K^H from that of $RU(-)_{(P)}$.*
- (e) *Any subgroup $C' \leq H$ conjugate in G to C lies in K .*

All of the above holds with $(KU_G)_{(P)}$ and $RU(-)_{(P)}$ replaced by their real variants $(KO_G)_{(P)}$ and $RO(-)_{(P)}$.

The equivalence of (a), (c) and (e) was already proven in [Böh19]. Theorem 1.5 is made more precise in Theorem 3.4, Theorem 3.8 and Corollary 4.3 in terms of Blumberg and Hill's framework of incomplete Tambara functors and N_∞ operads.

Remark 1.6. If H does not contain a group conjugate in G to C , then the norms \tilde{N}_K^H exist for trivial reasons: It can be seen from Theorem 2.5 that the restriction of e_C to H vanishes, and so $A(H)_{(p)}[e_C^{-1}]$ and $RU(H)_{(p)}[e_C^{-1}]$ must be zero. In other cases, these groups are always non-zero.

An immediate consequence of Theorem 1.5 is the following:

Corollary 1.7. *The summand $(KU_G)_{(p)}[e_C^{-1}]$ is a G - E_∞ ring spectrum if and only if $C \leq G$ is the trivial group. The same is true for real K-theory.*

1.3. Organization. In §2, we recall Dress' work on idempotents in the Burnside ring and give a proof of Theorem 1.2. The algebraic and homotopical parts of Theorem 1.5 are discussed in §3 and §4, respectively.

1.4. Acknowledgements. The present work was part of the author's PhD project at the University of Copenhagen; a previous version of the article was included in his PhD thesis [Böh18]. The author would like to thank his PhD advisor Jesper Grodal, his PhD committee consisting of Andrew Blumberg, John Greenlees and Lars Hesselholt, as well as Markus Hausmann, Joshua Hunt, Malte Leip, Riccardo Pengo, David Sprehn and an anonymous referee for many helpful discussions and suggestions. This research was supported by the Danish National Research Foundation through the Centre for Symmetry and Deformation (DNRF92).

2. IDEMPOTENT ELEMENTS IN REPRESENTATION RINGS

The goal of this section is to prove Theorem 1.2. In §2.1, we show how some parts of the theorem follow easily from the classification of idempotents in the Burnside ring. The difficult part is to prove that the images of the Burnside ring idempotents are indeed primitive. We recall Atiyah's description [Ati61] of the prime ideal spectrum $\text{Spec}(RU(G)_{(p)} \otimes \mathcal{O}_{\mathbb{F}})$ in §2.2, where $\mathcal{O}_{\mathbb{F}}$ is obtained from \mathbb{Z} by adjoining sufficiently many roots of unity, classify the idempotents of $RU(G)_{(p)} \otimes \mathcal{O}_{\mathbb{F}}$ in §2.3, and deduce the primitivity part of Theorem 1.2 in §2.4. In the rational and in the p -local case, it is possible to prove the primitivity in an easier way, as we explain in §2.5.

2.1. Idempotents in the Burnside ring. We recall Dress' classification of idempotents of $A(G)_{(p)}$ [Dre69] and prove parts of Theorem 1.2.

Notation 2.1. Recall that by the Chinese remainder theorem applied to $\langle g \rangle$, each $g \in G$ can be written uniquely as a product $g_P \cdot g_{P^\perp}$ of powers of g , where g_P is of order divisible only by primes in P , and g_{P^\perp} is of order prime to P . The elements g_P and g_{P^\perp} are called the P -part and P -prime part of g , respectively.

Definition 2.2. The P -residual subgroup of a group H is the unique minimal normal subgroup $O^P(H)$ such that the quotient is a solvable P -group, i.e. a solvable group of order only divisible by primes in P . H is called P -perfect if $O^P(H) = H$.

Lemma 2.3. For cyclic groups, we have $O^P(g) = \langle g_{P^\perp} \rangle$. In particular, a cyclic group is P -perfect if and only if its order is not divisible by any element of P . \square

Definition 2.4. For a subgroup $H \leq G$, the mark homomorphism $\phi^H: A(G)_{(P)} \rightarrow \mathbb{Z}_{(P)}$ is extended additively from the assignment $X \mapsto |X^H|$ for finite G -sets X .

Theorem 2.5 ([Dre69], Prop. 2). There is a canonical bijection between the conjugacy classes of P -perfect subgroups $L \leq G$ and the set of primitive idempotent elements of $A(G)_{(P)}$. It sends L to the element $e_L \in A(G)_{(P)}$ whose marks $\phi^H(e_L)$ at a subgroup $H \leq G$ are one if $O^P(H)$ and L are conjugate in G , and zero otherwise.

Write $\chi(V)(g)$ for the value of the character of $V \in RU(G)_{(P)}$ at the element $g \in G$. The linearization map $\text{lin}: A(G)_{(P)} \rightarrow RU(G)_{(P)}$ satisfies the following simple identity:

Lemma 2.6. For $X \in A(G)_{(P)}$, we have $\chi(\text{lin}(X))(g) = \phi^{(g)}(X) \in \mathbb{Z}$. \square

Corollary 2.7. Let $L \leq G$ be a P -perfect subgroup. Then the virtual representation $\text{lin}(e_L)$ has character values

$$\chi(\text{lin}(e_L))(g) = \phi^{(g)}(e_L) = \begin{cases} 1 & \text{if } \langle g_{P^\perp} \rangle \sim_G L \\ 0 & \text{otherwise.} \end{cases}$$

In particular, $\text{lin}(e_L)$ is zero if L is not cyclic. The elements $\text{lin}(e_C)$ are mutually orthogonal idempotents summing to one, where C ranges over a set of representatives for the conjugacy classes of cyclic P -perfect subgroups.

Proof. The statement follows from Theorem 2.5, Lemma 2.6, and Lemma 2.3. \square

This proves all the statements of Theorem 1.2 except for the primitivity of the idempotents $\text{lin}(e_C)$. Note that the rational case ($P = \emptyset$) of Corollary 2.7 is stated in [Glu81a, Theorem] and goes back to a similar result by Solomon [Sol67, Thm. 3].

The following observation is not part of the proof of Theorem 1.2, but we record it for later reference.

Lemma 2.8. The P -local Burnside ring splits as

$$A(G)_{(P)} \cong e_{\text{cyc}} \cdot A(G)_{(P)} \times e_{\text{ker}} \cdot A(G)_{(P)}$$

where e_{cyc} (respectively e_{ker}) is defined to be the sum of all primitive idempotents e_L with L cyclic (respectively non-cyclic). Moreover, the summand $e_{\text{ker}} \cdot A(G)_{(P)}$ is precisely the kernel of the linearization map $\text{lin}: A(G)_{(P)} \rightarrow R(G)_{(P)}$.

Proof. The first part follows from Theorem 2.5 by writing $1 = e_{cyc} + e_{ker}$. Lemma 2.6 implies that the kernel of lin consists of those virtual G -sets whose marks vanish at all cyclic subgroups. By Corollary 2.7, these are precisely the elements of the ideal $e_{ker} \cdot A(G)_{(P)}$. \square

2.2. Prime ideals in the splitting field case. Let $\exp(G)$ be the exponent³ of G and write \mathbb{F} for the $\exp(G)$ -th cyclotomic extension of \mathbb{Q} with ring of integers $\mathcal{O}_{\mathbb{F}}$ and Galois group $\Gamma := Gal(\mathbb{F} : \mathbb{Q})$. All characters of G -representations over the complex numbers take values in $\mathcal{O}_{\mathbb{F}}$, and therefore can be viewed as class functions $G/\sim \rightarrow \mathcal{O}_{\mathbb{F}}$, where G/\sim is the set of conjugacy classes of G . When working P -locally, the elements of $RU(G)_{(P)}$ are $\mathbb{Z}_{(P)}$ -linear combinations of irreducible representations of G over the complex numbers, hence their characters take values in $\mathcal{O}_{\mathbb{F},(P)} := \mathcal{O}_{\mathbb{F}} \otimes \mathbb{Z}_{(P)}$.

Notation 2.9. Any element $V \in RU(G)_{(P)} \otimes \mathcal{O}_{\mathbb{F}}$ can be written as an $\mathcal{O}_{\mathbb{F},(P)}$ -linear combination $V = \sum_i \lambda_i \cdot V_i$ of irreducible G -representations V_i . We write

$$\hat{\chi}(V)(g) := \sum_i \lambda_i \cdot \chi(V_i)(g)$$

for the value of the $\mathcal{O}_{\mathbb{F},(P)}$ -linear character of V at $g \in G$.

Characters are multiplicative in tensor products of representations, hence $\hat{\chi}(-)(g)$ defines a ring homomorphism $RU(G)_{(P)} \otimes \mathcal{O}_{\mathbb{F}} \rightarrow \mathcal{O}_{\mathbb{F},(P)}$. Atiyah [Ati61] described the structure of the prime ideal spectrum $\text{Spec}(RU(G) \otimes \mathcal{O}_{\mathbb{F}})$ in terms of these maps. His proof applies without changes to the open subscheme $\text{Spec}(RU(G)_{(P)} \otimes \mathcal{O}_{\mathbb{F}})$ cut out by P -localization.

Proposition 2.10 (Cf. [Ati61], Prop. 6.4). *The topological space $\text{Spec}(RU(G)_{(P)} \otimes \mathcal{O}_{\mathbb{F}})$ can be described as follows:*

(1) *Every prime ideal of $RU(G)_{(P)} \otimes \mathcal{O}_{\mathbb{F}}$ is of the form*

$$Q(\mathfrak{p}, g) := (\hat{\chi}(-)(g))^{-1}(\mathfrak{p}) = \{V \in RU(G)_{(P)} \otimes \mathcal{O}_{\mathbb{F}} \mid \hat{\chi}(V)(g) \in \mathfrak{p}\}$$

for some element $g \in G$ and some prime ideal $\mathfrak{p} \trianglelefteq \mathcal{O}_{\mathbb{F},(P)}$.

(2) *Let $\mathfrak{p}, \mathfrak{q} \trianglelefteq \mathcal{O}_{\mathbb{F},(P)}$ be prime ideals such that $\mathbb{Z} \cap \mathfrak{q} = q\mathbb{Z}$ for a prime $q \in \mathbb{Z}$. There is an inclusion $Q(\mathfrak{p}, g) \subseteq Q(\mathfrak{q}, h)$ if and only if \mathfrak{p} is contained in \mathfrak{q} and g_{q^\perp} is conjugate in G to h_{q^\perp} .*

(3) *The prime ideals $Q(\mathfrak{p}, g)$ with $\mathfrak{p} = (0)$ are minimal and the ones with $\mathfrak{p} \neq (0)$ are maximal. In particular, the Krull dimension of $RU(G)_{(P)} \otimes \mathcal{O}_{\mathbb{F}}$ is one.*

³The exponent of a finite group is the least common multiple of the orders of all group elements.

2.3. Idempotents in the splitting field case. We can deduce a classification of the idempotent elements of $RU(G)_{(P)} \otimes \mathcal{O}_{\mathbb{F}}$ from Proposition 2.10. Our proof is inspired by Dress' approach [Dre69, Prop. 2] to the idempotents in the Burnside ring.

Theorem 2.11. *The map*

$$G \rightarrow \pi_0(\text{Spec}(RU(G)_{(P)} \otimes \mathcal{O}_{\mathbb{F}}))$$

that sends $x \in G$ to the connected component of $Q(0, x)$ induces a bijection between the set of conjugacy classes of P -prime elements⁴ of G and the set of connected components of $\text{Spec}(RU(G)_{(P)} \otimes \mathcal{O}_{\mathbb{F}})$. In particular, the prime ideal spectrum of $RU(G) \otimes \mathcal{O}_{\mathbb{F}}$ is connected.

This follows directly from:

Proposition 2.12. *For any (not necessarily P -prime) elements $x, y \in G$, the prime ideals $Q(\mathfrak{p}, x)$ and $Q(\mathfrak{q}, y)$ lie in the same connected component of $\text{Spec}(RU(G) \otimes \mathcal{O}_{\mathbb{F}})$ if and only if x_{P^\perp} and y_{P^\perp} are conjugate in G .*

Proof. First observe that for $\mathfrak{p} \neq (0)$, the height one ideal $Q(\mathfrak{p}, x)$ lies in the closure of the height zero ideal $Q((0), x)$, so without loss of generality we may assume that $\mathfrak{p} = \mathfrak{q} = (0)$. Since $RU(G)_{(P)} \otimes \mathcal{O}_{\mathbb{F}}$ has Krull dimension one, two points $Q((0), x)$ and $Q((0), y)$ lie in the same component if and only if there is a zig-zag of inclusions of prime ideals

$$\begin{array}{ccccc} & & Q(\mathfrak{p}_0, x_0) = Q(\mathfrak{p}_0, x_1) & & \dots \\ & \nearrow & & \nwarrow & \nearrow \quad \nwarrow \\ Q((0), x_0) & & & & Q((0), x_1) & & Q((0), x_r) \end{array}$$

for some elements $x = x_0, x_1, \dots, x_r = y \in G$ and some prime ideals $\mathfrak{p}_i \trianglelefteq \mathcal{O}_{\mathbb{F},(P)}$. By part (2) of Theorem 2.10, we have an equality $Q(\mathfrak{p}_i, x_i) = Q(\mathfrak{p}_i, x_{i+1})$ if and only if $(x_i)_{\mathfrak{p}_i^\perp}$ is conjugate in G to $(x_{i+1})_{\mathfrak{p}_i^\perp}$, where \mathfrak{p}_i is given by $\mathbb{Z} \cap \mathfrak{p}_i = \mathfrak{p}_i \mathbb{Z}$.

For the "only if" part of the proposition, given a zig-zag as above, it follows that

$$x_{P^\perp} = ((x_0)_{\mathfrak{p}_0^\perp})_{P^\perp} \sim_G ((x_1)_{\mathfrak{p}_0^\perp})_{P^\perp} = ((x_1)_{\mathfrak{p}_1^\perp})_{P^\perp} \sim_G \dots \sim_G ((x_r)_{\mathfrak{p}_{r-1}^\perp})_{P^\perp} = y_{P^\perp}$$

where \sim_G indicates being conjugate in G .

For the "if" part, assume that $x_{P^\perp} \sim_G y_{P^\perp}$. Since the prime ideals $Q((0), g)$ only depend on the conjugacy class of g , it follows that $Q((0), x_{P^\perp})$ and $Q((0), y_{P^\perp})$ agree. Thus, it suffices to show that for any $g \in G$, the prime ideals $Q((0), g)$ and $Q((0), g_{P^\perp})$ lie in the same component. We will construct an explicit zig-zag as above. Let

⁴A P -prime element is an element x such that $x_{P^\perp} = x$.

p_0, p_1, \dots, p_r be all primes in P that divide the order of G . By the going-up theorem, we can find prime ideals $\mathfrak{p}_i \trianglelefteq \mathcal{O}_{\mathbb{F},(P)}$ such that $\mathfrak{p}_i \cap \mathbb{Z} = p_i \mathbb{Z}$. Then g_{P^\perp} may be computed as

$$g_{P^\perp} = (\cdots (g_{p_0^\perp})_{p_1^\perp} \cdots)_{p_r^\perp}.$$

Inductively, define $g_0 := g$ and $g_i := (g_{i-1})_{p_{i-1}^\perp}$ so that we have $(g_i)_{p_i^\perp} = (g_{i+1})_{p_i^\perp}$. Then these choices of elements g_i and prime ideals \mathfrak{p}_i give rise to a zig-zag between $Q((0), g)$ and $Q((0), g_{P^\perp})$, which completes the proof. \square

Corollary 2.13. *The conjugacy classes of P -prime elements (x) of G are in canonical bijection with the primitive idempotents e_x of $RU(G)_{(P)} \otimes \mathcal{O}_{\mathbb{F}}$. The character of the element e_x is given as follows:*

$$\hat{\chi}(e_x)(g) = \begin{cases} 1 & \text{if } g_{P^\perp} \sim_G x \\ 0 & \text{otherwise} \end{cases}$$

Proof. It is a standard fact of algebraic geometry that for any commutative ring R , the subsets $V \subseteq \text{Spec}(R)$ that are both open and closed are in canonical bijection with the idempotent elements of R , by assigning to V the global section⁵ which is constant one on V and constant zero on the complement of V . Under this identification, the primitive idempotents correspond to the minimal non-empty open and closed subsets. The latter agree with the connected components of $\text{Spec}(RU(G)_{(P)} \otimes \mathcal{O}_{\mathbb{F}})$ since there are only finitely many of them. The first claim now follows from Theorem 2.11. For the description of characters, note that $\hat{\chi}(e_x)(g) = 1$ if and only if the corresponding global section e_x evaluates to one at the point $Q((0), g)$ if and only if $Q((0), x)$ and $Q((0), g)$ are in the same connected component. \square

Remark 2.14. Roquette [Roq52] shows that the classification given in Corollary 2.13 also holds for the primitive idempotents in the p -adic representation ring after adjoining all e -th roots of unity.

Remark 2.15. Using Schur's orthogonality relations, it follows from Corollary 2.13 that e_x is given explicitly as

$$e_x = \frac{1}{|C_G(x)|} \sum_V \chi(V)(x^{-1}) \cdot V$$

where V runs over a system of representatives of the irreducible representations of G and $C_G(x)$ denotes the centralizer of x in G . This observation goes back at least to Brauer [Bra47, (7)]. The coefficients can also be expressed in terms of Möbius functions, see [Sol67, Thm. 4], [Glu81b, Prop.] and [Yos83, §3].

⁵Here we use that by definition, the global sections of $\text{Spec}(R)$ agree with the ring R .

2.4. Idempotents of $RU(G)_{(P)}$. Recall that $\Gamma \cong (\mathbb{Z}/\exp(G))^\times$ denotes the Galois group of the cyclotomic extension \mathbb{F}/\mathbb{Q} . The left Γ -action on \mathbb{F} restricts to an action on $\mathcal{O}_{\mathbb{F}}$. Let Γ act on $RU(G)_{(P)} \otimes \mathcal{O}_{\mathbb{F}}$ via its action on the right factor. Then clearly $RU(G)_{(P)} = (RU(G)_{(P)} \otimes \mathcal{O}_{\mathbb{F}})^\Gamma$. The group Γ then acts from the right on $\text{Spec}(RU(G)_{(P)} \otimes \mathcal{O}_{\mathbb{F}})$ and we have $\text{Spec}(RU(G)_{(P)}) \cong (\text{Spec}(RU(G)_{(P)} \otimes \mathcal{O}_{\mathbb{F}}))/\Gamma$, cf. [Ser77, §11.4, Exerc. 11.4]. We will now describe these Γ -orbits in terms of the prime ideals $Q(\mathfrak{p}, x)$.

First recall that the left Γ -action on $\mathcal{O}_{\mathbb{F}}$ induces a right Γ -action on $\text{Spec}(\mathcal{O}_{\mathbb{F}})$ that is given by $\mathfrak{p} \cdot \gamma = \gamma^{-1}(\mathfrak{p})$.

Definition 2.16 ([Ser77], §12.4). Define a right Γ -action on the underlying set of G as follows: If $\gamma \in \Gamma$ corresponds to the unit $m \in (\mathbb{Z}/\exp(G))^\times$, let $g \cdot \gamma := g^{m^{-1}}$, where m^{-1} is (any integer representing) the inverse of m in the group Γ .

This action is well-defined since the order of $g \in G$ divides $\exp(G)$. Moreover, it is compatible with conjugation in G . We can describe the Γ -orbits in G easily:

Lemma 2.17 ([Ser77], §13.1, Cor.). *Two elements $x, y \in G$ lie in the same Γ -orbit if and only if they generate the same cyclic subgroup of G .*

Proof. Let n divide $\exp(G)$. Then $\Gamma \cong (\mathbb{Z}/\exp(G))^\times$ permutes the generators of $\mathbb{Z}/\exp(G)$ transitively, and the same is true for the generators of the cyclic group \mathbb{Z}/n , viewed as a subgroup of $\mathbb{Z}/\exp(G)$. \square

Proposition 2.18. *The left Γ -action on $RU(G)_{(P)} \otimes \mathcal{O}_{\mathbb{F}}$ induces a right Γ -action on the space $\text{Spec}(RU(G)_{(P)} \otimes \mathcal{O}_{\mathbb{F}})$ which coincides with the action defined by $Q(\mathfrak{p}, x) \cdot \gamma = Q(\mathfrak{p} \cdot \gamma, x \cdot \gamma)$.*

Proof. As in Notation 2.9, write $\sum_i \lambda_i \cdot V_i$ for a generic element of $RU(G)_{(P)} \otimes \mathcal{O}_{\mathbb{F}}$. Then

$$\begin{aligned} Q(\mathfrak{p}, x) \cdot \gamma &= \gamma^{-1}(Q(\mathfrak{p}, g)) = \left\{ \sum_i \lambda_i \cdot V_i \mid \sum_i \gamma(\lambda_i) \cdot \chi(V_i)(g) \in \mathfrak{p} \right\} \\ &= \left\{ \sum_i \lambda_i \cdot V_i \mid \sum_i \lambda_i \cdot \gamma^{-1}(\chi(V_i)(g)) \in \mathfrak{p} \cdot \gamma \right\} \\ &= \left\{ \sum_i \lambda_i \cdot V_i \mid \sum_i \lambda_i \cdot \chi(V_i)(g \cdot \gamma) \in \mathfrak{p} \cdot \gamma \right\} \\ &= Q(\mathfrak{p} \cdot \gamma, g \cdot \gamma) \end{aligned} \quad \square$$

Corollary 2.19. *The map*

$$G \rightarrow \pi_0(\text{Spec}(RU(G)_{(P)}))$$

that sends an element x to the component of the orbit $Q((0), x) \cdot \Gamma$ induces a bijection between the Γ -orbits of conjugacy classes of P -prime elements $x \in G$ and the set of components of the topological space $\text{Spec}(RU(G)_{(P)})$. In particular, the spectrum of $RU(G)$ is connected.

Corollary 2.20. *There is a canonical bijection between the Γ -orbits of conjugacy classes of P -prime elements $x \in G$ and the primitive idempotents in $RU(G)_{(P)}$. The idempotent $e_{x,\Gamma}$ associated to the orbit of (x) has character given by*

$$\chi(e_{x,\Gamma})(g) = \begin{cases} 1 & \text{if } g_{P^\perp} \sim_G x \cdot \gamma \text{ for some } \gamma \in \Gamma \\ 0 & \text{otherwise} \end{cases}$$

Proof. This follows from Corollary 2.19 in the same way that Corollary 2.13 follows from Theorem 2.11, see the proof of Corollary 2.13. \square

Remark 2.21. In particular, we have $e_{x,\Gamma} = \sum_{\gamma \in \Gamma} e_{x \cdot \gamma}$ in $RU(G)_{(P)}$. A simple calculation shows that $e_{x \cdot \gamma} = \gamma^{-1}(e_x)$ in $RU(G)_{(P)} \otimes \mathcal{O}_{\mathbb{F}}$. Therefore $e_{x,\Gamma} = \text{tr}_{\mathbb{F}/\mathbb{Q}}(e_x)$ is the field trace of e_x . We will not use this fact.

By Lemma 2.17, we can write $e'_{\langle x \rangle} := e_{x,\Gamma}$ and rephrase Corollary 2.20 in terms of cyclic subgroups. At this point, there is no dependence on the field extension \mathbb{F}/\mathbb{Q} anymore.

Corollary 2.22. *There is a canonical bijection between the conjugacy classes of cyclic P -perfect subgroups $C \in G$ and the primitive idempotents in $RU(G)_{(P)}$. The primitive idempotent e'_C has character given by*

$$\chi(e'_C)(g) = \begin{cases} 1 & \text{if } \langle g_{P^\perp} \rangle \sim_G C \\ 0 & \text{otherwise.} \end{cases}$$

In particular, the character of e'_C agrees with that of $\text{lin}(e_C)$ given in Corollary 2.7 and hence we have $e'_C = \text{lin}(e_C)$.

Theorem 1.2 follows.

Remark 2.23. It is clear from Corollary 2.22 that the primitive idempotents of $RU(G)_{(P)}$ only depend on those primes $p \in P$ that divide the order of G .

2.5. Quick proofs of special cases. In the rational and p -local case, we can give short ad-hoc proofs of the primitivity of the elements $\text{lin}(e_C)$ stated as part of Theorem 1.2.

Lemma 2.24. *Let $x, y \in G$ generate the same subgroup. If all character values of the virtual representation $V \in RU(G)_{(P)}$ lie in $\mathbb{Z}_{(P)}$, then $\chi(V)(x) = \chi(V)(y)$.*

Proof. By Lemma 2.17, we can find $\gamma \in \Gamma$ such that $y = x \cdot \gamma$. Then

$$\chi(V)(y) = \chi(V)(x \cdot \gamma) = \gamma^{-1}(\chi(V)(x)) = \chi(V)(x)$$

because $\chi(V)(x) \in \mathbb{Z}_{(P)} = (\mathcal{O}_{\mathbb{F},(P)})^\Gamma$. \square

Corollary 2.25. *For any cyclic $C \leq G$, the idempotent $\text{lin}(e_C) \in RU(G) \otimes \mathbb{Q}$ is primitive.*

Proof. Recall that the character of $\text{lin}(e_C)$ is one on elements that generate subgroups conjugate to C and zero otherwise. But Lemma 2.24 shows that any integer-valued character must be constant on the set where $\text{lin}(e_C)$ is one, hence $\text{lin}(e_C)$ cannot decompose as a sum of idempotents. \square

For the p -local case, we need another lemma. It was used in Atiyah's proof of Theorem 2.10.

Lemma 2.26 ([Ati61], proof of Lemma 6.3). *Let $V \in \text{RU}(G)_{(p)}$ and let \mathfrak{p} be a prime of $\mathcal{O}_{\mathbb{F},(p)} = \mathcal{O}_{\mathbb{F}} \otimes \mathbb{Z}_{(p)}$. Then $\chi(V)(g) \equiv \chi(V)(g_{p^\perp}) \pmod{\mathfrak{p}}$.*

Proof. Without loss of generality, we may assume that G is cyclic and V one-dimensional, hence its character is multiplicative. Write $g = g_{p^\perp} \cdot h$ where the order of h is p^r , then $(\chi(V)(h))^{p^r} = 1$. But $\mathcal{O}_{\mathbb{F},(p)}/\mathfrak{p}$ is a finite field of characteristic p , so

$$\chi(V)(h) \equiv 1 \pmod{\mathfrak{p}},$$

and consequently

$$\chi(V)(g) \equiv \chi(V)(g_{p^\perp}) \cdot \chi(V)(h) \equiv \chi(V)(g_{p^\perp}) \pmod{\mathfrak{p}}. \quad \square$$

Definition 2.27. For $C \leq G$ cyclic of order prime to p , let

$$S_C := \{g \in G \mid \langle g_{p^\perp} \rangle \sim_G C\}.$$

Combining Lemma 2.24 and Lemma 2.26 gives:

Corollary 2.28. *Let \mathfrak{p} be any prime ideal in $\mathcal{O}_{\mathbb{F},(p)}$. If all character values of $V \in \text{RU}(G)_{(p)}$ lie in $\mathbb{Z}_{(p)}$, then the character of V is constant modulo \mathfrak{p} on the set S_C .*

Finally, a proof similar to that of Corollary 2.25 shows:

Corollary 2.29. *For any cyclic p -perfect $C \leq G$, the idempotent $\text{lin}(e_C) \in \text{RU}(G)_{(p)}$ is primitive.*

Lemma 2.26 does not hold in the general P -local case, as the next example shows. However, it follows from Theorem 1.2 that the statement becomes true under the additional assumption that the character of V be zero outside of S_C . We do not know how to use this assumption to give a quick proof of the primitivity of the elements $\text{lin}(e_C)$ that applies to all choices of P .

Example 2.30. Let $G = C_2 \times C_3$ be the cyclic group of order 6 and $P = \{2, 3\}$. Write $\mathbb{1}$ for the trivial representation and let $V \in \text{RU}(G)_{(p)}$ be given as the tensor product

of the sign representation of C_2 with the sum of the two non-trivial irreducible C_3 -representations. Let $g \in G$ be a generator and observe that $g_{p^\perp} = 1$. However,

$$\chi(V - \mathbb{1})(g) = 0 \not\equiv 1 = \chi(V - \mathbb{1})(1) \pmod{\mathfrak{p}}$$

for any prime ideal \mathfrak{p} of $\mathcal{O}_{\mathbb{F},(P)}$.

3. IDEMPOTENT SPLITTINGS OF REPRESENTATION RINGS

As before, let P be a fixed collection of prime numbers, and let $R(G)_{(P)}$ denote one of the rings $RO(G)_{(P)}$ or $RU(G)_{(P)}$. The goal of this section is to describe the multiplicativity of the idempotent splitting

$$R(-)_{(P)} \cong \prod_{(C)} R(-)_{(P)}[e_C^{-1}].$$

We start by briefly recalling the notion of an (incomplete) Tambara functor in §3.1. In §3.2, we study the multiplicativity of the idempotent splitting of $R(-)_{(P)}$: we characterize the norms which are compatible with e_C -localization in Theorem 3.4 and describe the incomplete Tambara functor structure of each idempotent summand in Theorem 3.8. It is then easy to read off the structure that is preserved by the entire splitting, as we explain in §3.3.

3.1. Incomplete Tambara functors. Recall that many naturally arising Mackey functors have additional multiplicative structure.

Definition 3.1. A *Green functor* is a Mackey functor \underline{R} equipped with commutative ring structures on the values $\underline{R}(H)$ for all $H \leq G$ such that all restriction maps $R_K^H: \underline{R}(H) \rightarrow \underline{R}(K)$ become ring homomorphisms and all transfer maps $T_K^H: \underline{R}(K) \rightarrow \underline{R}(H)$ are morphisms of $\underline{R}(H)$ -modules.

Often, Green functors come equipped with additional *multiplicative transfer maps* or *norms* $N_K^H: \underline{R}(K) \rightarrow \underline{R}(H)$ for all subgroup inclusions $K \leq H \leq G$, satisfying a number of compatibility relations for norms, additive transfers and restrictions. Tambara [Tam93] axiomatized the structure of these objects and called them *TNR-functors*; nowadays they are referred to as *Tambara functors*.

Blumberg and Hill [BH18] introduced the more general notion of an (incomplete) \mathcal{I} -Tambara functor that only admits a partial collection of norms for certain subgroup inclusions $K \leq H \leq G$, parametrized by well-behaved collections \mathcal{I} of *admissible H -sets* H/K . These *indexing systems* form a poset under inclusion. Thus, \mathcal{I} -Tambara functors for varying \mathcal{I} interpolate between the notion of a Green functor (which doesn't necessarily admit any norms) and that of a Tambara functor (which admits all possible norms). We refer to the above sources for precise definitions and further details.

Example 3.2. The Mackey functors defined by the Burnside ring $A(-)$ and the representation rings $R(-)$ are examples of Tambara functors. The multiplicative norms N_K^H in the Burnside ring are induced by the co-induction functor $\text{map}_K(H, -)$ from finite K -sets to finite H -sets. Those of the representation ring are induced by tensor induction of representations. The linearization maps $\text{lin}: A(-) \rightarrow R(-)$ are maps of Tambara functors.

More generally, Brun [Bru07] showed that the zeroth equivariant homotopy groups of a G - E_∞ ring spectrum naturally form a Tambara functor.

Remark 3.3. Note that the $RO(G)$ -graded homotopy groups of a G - E_∞ ring spectrum form a *graded Tambara functor* in the sense of Angeltveit-Bohmann [AB18].

3.2. Multiplicativity of the idempotent summands. Observe that the canonical localization maps $R(-)_{(P)} \rightarrow R(-)_{(P)}[e_C^{-1}]$ are levelwise ring homomorphisms that are compatible with the Mackey functor structure, hence the idempotent splitting of $R(G)_{(P)}$ induces a splitting of the underlying Green functor of $R(-)_{(P)}$. Our next goal is to describe the idempotent summands $R(-)_{(P)}[e_C^{-1}]$ by proving the equivalence of the statements (c), (d) and (e) of Theorem 1.5. For convenience of the reader, we record this in the following theorem.

Theorem 3.4. *Let $C \leq G$ be a cyclic P -perfect subgroup and let $e_C \in A(G)_{(P)}$ be the corresponding primitive idempotent element. Fix subgroups $K \leq H \leq G$. Then the following are equivalent:*

(c) *The norm map $N_K^H: A(K)_{(P)} \rightarrow A(H)_{(P)}$ descends to a well-defined map of multiplicative monoids*

$$\tilde{N}_K^H: A(K)_{(P)}[e_C^{-1}] \rightarrow A(H)_{(P)}[e_C^{-1}].$$

(d) *The norm map $N_K^H: R(K)_{(P)} \rightarrow R(H)_{(P)}$ descends to a well-defined map of multiplicative monoids*

$$\tilde{N}_K^H: R(K)_{(P)}[e_C^{-1}] \rightarrow R(H)_{(P)}[e_C^{-1}].$$

(e) *Any subgroup $C' \leq H$ conjugate in G to C lies in K .*

It was proven in [Böh19, Thm. 4.1] that the statements (c) and (e) are equivalent, so Theorem 3.4 reduces to showing that (c) and (d) are equivalent.

We recall the following fact due to Blumberg and Hill (see [Böh19, Thm. 2.33] for an elementary proof):

Theorem 3.5 ([BH18], Thm. 5.25). *Let \underline{R} be an \mathcal{I} -Tambara functor structured by an indexing system \mathcal{I} . Let $x \in \underline{R}(G)$. Then the orbit-wise localization $\underline{R}[x^{-1}]$ is a localization in the*

category of \mathcal{I} -Tambara functors if and only if for all admissible sets H/K of \mathcal{I} , the element $N_K^H R_K^G(x)$ divides a power of $R_H^G(x)$.

If the element x is idempotent, then checking the above division relation amounts to checking an equation:

Lemma 3.6. *Let $e, e' \in R$ be idempotents in a commutative ring. Then e divides e' if and only if $e \cdot e' = e'$.*

Proof. Assume that e divides e' . Then $e' \in eR$, hence $e \cdot e' = e'$, since multiplication by e is projection onto the idempotent summand eR of R . The other direction is obvious. \square

Proof of Theorem 3.4. We only need to show the equivalence (c) \Leftrightarrow (d). By Theorem 3.5 and Lemma 3.6, the statement (c) (respectively (d)) holds if and only if the equation

$$N_K^H R_K^G(x) \cdot R_H^G(x) = R_H^G(x)$$

holds in $A(H)_{(P)}$ for $x = e_C$ (respectively in $R(H)_{(P)}$ for $x = \text{lin}(e_C)$). The linearization map $\text{lin}: A(-)_{(P)} \rightarrow R(-)_{(P)}$ is a map of Tambara functors, hence preserves norms, restrictions and multiplication. By Lemma 2.8, lin is injective on the ideal summand $e_{\text{cyc}} \cdot A(G)_{(P)}$ and that summand contains the element e_C . It follows that the above equation holds for $x = e_C$ if and only if it holds for $x = \text{lin}(e_C)$. \square

We can use the language of incomplete Tambara functors [BH18] to describe the algebraic structure of $R(G)_{(P)}[e_C^{-1}]$ in terms of certain indexing systems.

Proposition 3.7 ([Böh19], Prop. 4.16). *Let $L \leq G$ be P -perfect. There is an indexing system \mathcal{I}_L given as follows: for all $H \leq G$, $\mathcal{I}_L(H)$ is the full subcategory of finite H -sets spanned by all coproducts of the orbits H/K such that the groups $K \leq H \leq G$ satisfy the following condition: Any subgroup $L' \leq H$ conjugate in G to L lies in K .*

Note that this condition on H/K is a generalized version of condition (e) of Theorem 1.5 with C and C' replaced by P -perfect subgroups L and L' that are not necessarily cyclic.

Theorem 3.8. *Let $C \leq G$ be a cyclic P -perfect subgroup, and denote by $R(-)_{(P)}$ one of the Tambara functors $RU(-)_{(P)}$ or $RO(-)_{(P)}$. Then the following hold:*

- i) *The Green functor $R(-)_{(P)}[e_C^{-1}]$ admits the structure of an \mathcal{I}_C -Tambara functor under $R(-)_{(P)}$.*
- ii) *The indexing system \mathcal{I}_C is maximal among the indexing systems that satisfy i).*

iii) The canonical map $R(-)_{(P)} \rightarrow R(-)_{(P)}[e_C^{-1}]$ is an e_C -localization in the category of \mathcal{I}_C -Tambara functors. \square

We record two easy consequences of our characterization of norm maps in the idempotent summands.

Corollary 3.9. *The summand $R(-)_{(P)}[e_C^{-1}]$ is a Tambara functor (i.e., has a complete set of norms) if and only if C is the trivial group. \square*

Corollary 3.10. *The subgroup C is normal in G if and only if the summand $R(-)_{(P)}[e_C^{-1}]$ admits all norms of the form \tilde{N}_K^H such that K contains a subgroup conjugate in G to C . \square*

3.3. Multiplicativity of the idempotent splittings. We can now describe the multiplicativity of the idempotent splitting of $R(-)_{(P)}$ in terms of the indexing system

$$\mathcal{I}_{cyc} := \bigcap_{(C)} \mathcal{I}_C$$

arising as the intersection of the indexing systems \mathcal{I}_C defined in Prop. 3.7.

Proposition 3.11. *The localization maps $R(-)_{(P)} \rightarrow R(-)_{(P)}[e_C^{-1}]$ assemble into an isomorphism of \mathcal{I}_{cyc} -Tambara functors*

$$R(-)_{(P)} \rightarrow \prod_{(C) \leq G} R(-)_{(P)}[e_C^{-1}]$$

where the product is taken over conjugacy classes of cyclic P -perfect subgroups. Moreover, \mathcal{I}_{cyc} is maximal among all indexing sets with this property.

Proof. Each of the canonical maps $R(-)_{(P)} \rightarrow R(-)_{(P)}[e_C^{-1}]$ is a map of \mathcal{I}_C -Tambara functors by 3.8, hence their product is a map of \mathcal{I}_{cyc} -Tambara functors. It is a level-wise isomorphism by construction. The maximality also follows from Theorem 3.8: it implies that \mathcal{I}_{cyc} is maximal among the indexing systems \mathcal{J} such that each summand $R(-)_{(P)}[e_C^{-1}]$ is a \mathcal{J} -Tambara functor. \square

The admissible sets of \mathcal{I}_{cyc} can be characterized as follows.

Lemma 3.12 ([Böh19], Lemma 4.23). *Let $K \leq H \leq G$, then H/K is an admissible set for \mathcal{I}_{cyc} if and only if for all cyclic P -perfect $C \leq H$, C is contained in K .*

4. IDEMPOTENT SPLITTINGS OF EQUIVARIANT K-THEORY

Let K_G denote one of the genuine G -spectra KU_G or KO_G , i.e., either complex or real equivariant K -theory. We will determine the multiplicativity of the P -local idempotent splitting

$$(K_G)_{(P)} \simeq \prod_{(C)} (K_G)_{(P)}[e_C^{-1}],$$

i.e., we will explicitly describe the maximal N_∞ algebra structure on each of the factors, as well as the maximal N_∞ algebra structure preserved by the splitting. Recall that as a consequence of Theorem 1.2, the blocks of $(K_G)_{(P)}$ are given as the e_C -localizations

$$(K_G)_{(P)} \wedge \mathbb{S}_{(P)}[e_C^{-1}]$$

of $(K_G)_{(P)}$ in the category of G -spectra.

4.1. Preliminaries. The N_∞ operads of [BH15] structure G -equivariant ring spectra with incomplete sets of norm maps parametrized by their associated indexing systems. According to [GW18, Thm. 4.7, Prop. 4.10], any given indexing system can be realized as the indexing system of a Σ -cofibrant⁶ N_∞ operad. Similar existence results were given in [Rub17, Thm. 2.16] and [BP17, Cor. IV].

Notation 4.1. For each conjugacy class of cyclic P -perfect subgroups $C \leq G$, let \mathcal{O}_C be a Σ -cofibrant N_∞ operad whose associated indexing system is \mathcal{I}_C . Let \mathcal{O}_{cyc} be a Σ -cofibrant N_∞ operad whose associated indexing system is \mathcal{I}_{cyc} .

Note that by definition, an N_∞ operad \mathcal{P} is a certain operad in the category of unbased G -spaces. By the usual abuse of notation, we refer to an algebra over the operad $\Sigma_+^\infty \mathcal{P}$ in G -spectra as a \mathcal{P} -algebra.

Remark 4.2. For any choice of the operad \mathcal{O}_C , both \mathbb{S} and K_C are naturally algebras over \mathcal{O}_C : both spectra can be modelled as strictly commutative monoids in orthogonal G -spectra, and hence admit an action by \mathcal{O}_C that factors through the action of the commutative operad.

4.2. Multiplicativity of the idempotent summands. We are now ready to state our main homotopical result.

Theorem 4.3. *Let $C \leq G$ be a cyclic P -perfect subgroup. Then:*

- i) The G -spectrum $(K_G)_{(P)}[e_C^{-1}]$ is an \mathcal{O}_C -algebra under $(K_G)_{(P)}$.*
- ii) The operad \mathcal{O}_C is maximal among the N_∞ -operads that satisfy i).*
- iii) The canonical map $(K_G)_{(P)} \rightarrow (K_G)_{(P)}[e_C^{-1}]$ is an e_C -localization in the category of \mathcal{O}_C -algebras in G -spectra.*

The key to the proof is the following preservation result for N_∞ algebras given in [Böh19]. It extends previous work of Hill and Hopkins [HH14] and uses a result of Gutiérrez and White [GW18, Cor. 7.10].

⁶An operad \mathcal{O} in G -spaces is Σ -cofibrant if each space $\mathcal{O}(n)$ is of the homotopy type of a $(G \times \Sigma_n)$ -CW complex.

Proposition 4.4 ([Böh19], Prop. 2.32). *Let \mathcal{P} be a Σ -cofibrant N_∞ operad. Fix $x \in \pi_0^G(\mathbf{S}_{(P)})$. Then the Bousfield localization L_x given by smashing with*

$$\mathbf{S}_{(P)}[x^{-1}] = \text{hocolim} \left(\mathbf{S}_{(P)} \xrightarrow{x} \mathbf{S}_{(P)} \xrightarrow{x} \dots \right)$$

preserves⁷ \mathcal{P} -algebras in P -local G -spectra if and only if for all $H \leq G$ and all transitive admissible H -sets H/K , the element $N_K^H R_K^G(x)$ divides a power of $R_H^G(x)$ in the ring $\pi_0^H(\mathbf{S}_{(P)})$.

Proof of Theorem 4.3. Ad i): We know from Theorems 3.5 and 3.4 that for each of the admissible sets of \mathcal{I}_C , hence of \mathcal{O}_C , the division relation of Prop. 4.4 holds, so $(K_G)_{(P)}[e_C^{-1}]$ is an \mathcal{O}_C -algebra under $(K_G)_{(P)}$.

Ad ii): Assume that \mathcal{P} is an element strictly greater than \mathcal{O}_C in the poset of (homotopy types of) N_∞ operads. Then any norm that comes from \mathcal{P} but not from \mathcal{O}_C induces a corresponding norm on homotopy groups that does not correspond to an admissible set of \mathcal{I}_C , thus contradicting the maximality statement included in Theorem 3.8.

Ad iii): It is an e_C -localization in G -spectra and a map of \mathcal{O}_C -algebras. \square

We obtain the homotopical analogue of Corollary 3.9, stated as Corollary 1.7 in the introduction. There is also a homotopical version of Corollary 3.10:

Corollary 4.5. *The group C is normal in G if and only if $(K_G)_{(P)}[e_C^{-1}]$ admits all norm maps of the form \tilde{N}_K^H such that K and H both contain a subgroup conjugate in G to C . \square*

4.3. Multiplicativity of the idempotent splitting. We can also describe the multiplicativity of the entire idempotent splitting:

Corollary 4.6. *Let \mathcal{O}_{cyc} be a Σ -cofibrant N_∞ operad realizing the indexing system $\mathcal{I}_{cyc} = \bigcap_{(C)} \mathcal{I}_C$. Then the idempotent splitting*

$$(K_G)_{(P)} \simeq \prod_{(C)} (K_G)_{(P)}[e_C^{-1}]$$

is an equivalence of \mathcal{O}_{cyc} -algebras. Here, the product is taken over all conjugacy classes of cyclic P -perfect subgroups of G .

REFERENCES

- [AB18] V. Angeltveit and A. M. Bohmann. Graded Tambara functors. *J. Pure Appl. Algebra*, 222(12):4126–4150, 2018.
- [Ati61] M. F. Atiyah. Characters and cohomology of finite groups. *Inst. Hautes Études Sci. Publ. Math.*, 9:23–64, 1961.
- [BH15] A. J. Blumberg and M. A. Hill. Operadic multiplications in equivariant spectra, norms, and transfers. *Adv. Math.*, 285:658–708, 2015.

⁷ in the sense of [GW18, Def. 7.3]

- [BH18] A. J. Blumberg and M. A. Hill. Incomplete Tambara functors. *Algebr. Geom. Topol.*, 18(2):723–766, 2018.
- [BH20] A. J. Blumberg and M. A. Hill. G -symmetric monoidal categories of modules over equivariant commutative ring spectra. *Tunis. J. Math.*, 2(2):237–286, 2020.
- [Böh18] B. Böhme. *Equivariant multiplications and idempotent splittings of G -spectra*. PhD thesis, University of Copenhagen, 2018.
- [Böh19] B. Böhme. Multiplicativity of the idempotent splittings of the Burnside ring and the G -sphere spectrum. *Adv. Math.*, 347:904–939, 2019.
- [BP17] P. Bonventre and L. A. Pereira. Genuine equivariant operads. [arXiv:1707.02226v1](https://arxiv.org/abs/1707.02226v1), 2017.
- [Bra47] R. Brauer. On Artin’s L -series with generalized group characters. *Ann. of Math. (2)*, 48:502–514, 1947.
- [Bru07] M. Brun. Witt vectors and equivariant ring spectra applied to cobordism. *Proc. Lond. Math. Soc. (3)*, 94:351–385, 2007.
- [Dre69] A. Dress. A characterisation of solvable groups. *Math. Z.*, 110:213–217, 1969.
- [Glu81a] D. Gluck. A character table bound for the Schur index. *Illinois J. Math.*, 25(4):649–653, 1981.
- [Glu81b] D. Gluck. Idempotent formula for the Burnside ring algebra with applications to the p -subgroup simplicial complex. *Illinois J. Math.*, 25(1):63–67, 1981.
- [GW18] J. J. Gutiérrez and D. White. Encoding equivariant commutativity via operads. *Algebr. Geom. Topol.*, 18(5):2919–2962, 2018.
- [HH14] M. A. Hill and M. J. Hopkins. Equivariant multiplicative closure. *Algebraic topology: applications and new directions, Contemp. Math.*, 620, 2014. Amer. Math. Soc., Providence, RI.
- [HHR16] M. A. Hill, M. J. Hopkins, and D. C. Ravenel. On the nonexistence of elements of Kervaire invariant one. *Ann. of Math. (2)*, 184(1):1–262, 2016.
- [Roq52] P. Roquette. Arithmetische Untersuchung des Charakterringes einer endlichen Gruppe. Mit Anwendungen auf die Bestimmung des minimalen Darstellungskörpers einer endlichen Gruppe und in der Theorie der Artinschen L -Funktionen. *J. Reine Angew. Math.*, 190, 1952.
- [Rub17] J. Rubin. Combinatorial N_∞ operads. [arXiv:1705.03585v2](https://arxiv.org/abs/1705.03585v2), 2017.
- [Seg68] G. Segal. Equivariant K -theory. *Inst. Hautes Études Sci. Publ. Math.*, 34:129–151, 1968.
- [Seg71] G. Segal. Equivariant stable homotopy theory. *Actes, Congrès Intern. Math., Tome 2 (1970)*, pages 59–63, Paris, 1971.
- [Ser77] J.-P. Serre. *Linear representations of finite groups*. Graduate Texts in Mathematics, 42. Springer-Verlag, New York-Heidelberg, 1977. Translated from the second French edition by Leonard L. Scott.
- [Sol67] L. Solomon. The Burnside algebra of a finite group. *J. Combinatorial Theory*, 2:603–615, 1967.
- [Tam93] D. Tambara. On multiplicative transfer. *Comm. Algebra*, 21(4):1393–1420, 1993.
- [Yos83] T. Yoshida. Idempotents of Burnside Rings and Dress Induction Theorem. *J. Algebra*, 80:90–105, 1983.

MAX PLANCK INSTITUTE FOR MATHEMATICS, VIVATSGASSE 7, 53111 BONN, GERMANY

E-mail address: boehme@mpim-bonn.mpg.de