



# Diatom metabarcoding and microscopic analyses from sediment samples at Lake Nam Co, Tibet: The effect of sample-size and bioinformatics on the identified communities

Wengang Kang<sup>a</sup>, Sten Anslan<sup>b,\*</sup>, Nicole Börner<sup>a</sup>, Anja Schwarz<sup>a</sup>, Robin Schmidt<sup>b</sup>, Sven Künzel<sup>c</sup>, Patrick Rioual<sup>d,f</sup>, Paula Echeverría-Galindo<sup>a</sup>, Miguel Vences<sup>b</sup>, Junbo Wang<sup>e</sup>, Antje Schwalb<sup>a</sup>

<sup>a</sup> Institute of Geosystems and Bioindication, Technische Universität Braunschweig, Braunschweig, Germany

<sup>b</sup> Zoological Institute, Technische Universität Braunschweig, Braunschweig, Germany

<sup>c</sup> Department of Evolutionary Genetics, Max Planck Institute for Evolutionary Biology, Plön, Germany

<sup>d</sup> Key Laboratory of Cenozoic Geology & Environment, Institute of Geology and Geophysics, Chinese Academy of Sciences, Beijing, China

<sup>e</sup> Key Laboratory of Tibetan Environment Changes and Land Surface Processes (TEL)/Nam Co Observation and Research Station (NAMORS), Institute of Tibetan Plateau Research, Chinese Academy of Sciences, Beijing 100101, China

<sup>f</sup> CAS Center for Excellence in Life and Paleoenvironment, Beijing, China

## ARTICLE INFO

### Keywords:

Diatoms  
Metabarcoding  
Tibetan Plateau  
sedDNA  
Bioinformatics

## ABSTRACT

Diatoms (Bacillariophyceae) are characterized by silicified cell walls that favor their long-term preservation in sediments, therefore widely used as bioindicators of present and past water conditions. Alongside with traditional morphological analyses, metabarcoding has become a valuable tool to study the community structures of various organisms, including diatoms. Here, we test whether the quantity of sediment sample used for DNA extraction affects the results obtained from high-throughput sequencing (metabarcoding) of the diatom *rbcL* region by isolating DNA from 10 g and 0.5 g (wet weight) of lake surface sediment samples. Because bioinformatics processing of metabarcoding data may affect the outcome, we also tested the consistency of the results from three different pipelines: 1) ESVs (exact sequence variants) pipeline; 2) clustering sequences at 95% sequence identity to form OTUs (operational taxonomic units; 95% OTUs); and 3) 97% OTUs pipeline. Additionally, the agreement between metabarcoding data and morphological inventories of corresponding samples were compared. Our results demonstrate highly uniform patterns between the diatom *rbcL* amplicons from 10 g and 0.5 g of sedimentary DNA (sedDNA) extracts (HTS 10 and HTS 0.5, respectively). Furthermore, after the careful curation of the sequencing data, metabarcoding results were highly consistent among the data sets produced by different bioinformatics pipelines. Comparing results from metabarcoding and microscopy, we identified some taxonomic mismatches: morphological analyses identified 59 diatom genera, whereas metabarcoding 49 to 54 genera. These mismatches are related to incompleteness of the sequence databases, but also to inconsistencies in diatom taxonomy in general and potential dissolution effects of diatom valves caused by high alkalinity of the investigated lake waters. Nevertheless, multivariate community analysis revealed consistent results between data sets identified by microscopy and metabarcoding – water depth and conductivity as the most significant variables in driving diatom communities in Lake Nam Co – further confirming that metabarcoding is a viable method for identifying diatom-environment relationships.

## 1. Introduction

Diatoms (Bacillariophyceae) rank among the most important components of aquatic food webs and play an important role in carbon

fixation (Mann, 1999). Because of the fast response and narrow optima of some taxa for multiple environmental variables, diatoms are excellent indicators of ecosystem health (Van Dam et al., 1994; Battarbee et al., 2001), and may provide early warning signals for aquatic ecosystem

\* Corresponding author.

E-mail address: [sten.anslan@gmail.com](mailto:sten.anslan@gmail.com) (S. Anslan).

<https://doi.org/10.1016/j.ecolind.2020.107070>

Received 8 May 2020; Received in revised form 3 October 2020; Accepted 9 October 2020


Available online 22 October 2020

1470-160X/© 2020 The Author(s).

Published by Elsevier Ltd.

This is an open access article under the CC BY-NC-ND license

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>).




**Fig. 1.** Geographic map (A) highlighting the study area (Lake Nam Co) on the Tibetan Plateau. The bathymetric map (B) shows the major tributaries, ponds and glacier within the Nam Co catchment, location of Nam Co Monitoring and Research Station (NAMORS, red star) and sampling sites (green dots; the numbers from 1 to 23 corresponds to the sampling sites described in Table S1).

changes in face of environmental pressures such as nutrient enrichment (Wang et al., 2012) or heavy metal contamination (Chen et al., 2015). Silicified cell walls of diatoms favor their long-term preservation in the sediments, therefore they are widely applied to assess the water quality and long-term lake responses to environmental changes (Liu et al., 2017; Kang et al., 2019). The standard methods for assessing diatom communities rely on counting and identifying their silicified cell walls (valves) using mostly light microscopy (e.g. European-Committee-for-Standardization, 2014). But with the rapid development and continuously decreasing costs of high-throughput sequencing (HTS) technologies, the metabarcoding approach, allowing simultaneous identification of multiple species from environmental samples, is becoming a standard tool for fast biodiversity assessment (Ruppert et al., 2019). Because morphological analyses of diatoms (and other microorganisms) are labor-intensive, require expertise and are prone to inter-investigator variation, metabarcoding, referred to as 'Biomonitoring 2.0' (Baird and Hajibabaei, 2012), may have the potential to outperform the traditional, low throughput, monitoring methods.

Metabarcoding-based biodiversity studies, however, may face various difficulties, starting from DNA extraction to data processing in complex bioinformatics pipelines (Sinha et al., 2017; Anslan et al., 2018; Hardge et al., 2018). Therefore, the suitability of metabarcoding approach for assessing diatom communities have been the research focus for several studies. Although the DNA barcoding library for accurate species level identification is still incomplete for many organisms, metabarcoding can be a promising tool for biomonitoring diatom community assemblages as it has been shown to produce comparable results with traditional morphological analyses (Zimmermann et al., 2015; Apothéoz-Perret-Gentil et al., 2017; Vasselon et al., 2017b; Keck et al., 2018; Rimet et al., 2018a, 2018b; Rivera et al., 2018). The majority of diatom metabarcoding studies are applied to biofilms of epilithic diatom species from rivers and lakes, with the goal of assessing current-state water quality. Because diatom silicified valves are usually well preserved in sediments, they also constitute important indicators for inferring paleo-environmental conditions such as water pH, nutrient dynamics, and temperature (Douglas et al., 2010). However, only few studies have estimated the suitability of metabarcoding for identifying diatom communities directly from sediment samples and have assessed

its consistency with microscopy (Dulias et al., 2017; Piredda et al., 2017). Although morphological and metabarcoding data sets from these studies have demonstrated highly correlated results, it is not clear how this pattern is related to the quantity of sediment used for DNA extraction or affected by the use of different bioinformatics pipelines. The quantity of sediment used strongly depends on the approach taken for DNA extraction; it is common to use DNA isolation kits which allow input of 'large' quantities (usually up to 10 g) of environmental sample, to potentially capture the complete community represented in the sample. However, DNA extraction methods, such as the 'universal' Power Soil Kit (Hermans et al., 2018) processes small amount of material using less chemicals. Therefore, it costs only a fractional amount and may represent attractive alternatives for DNA metabarcoding of large numbers of samples. Multiple publicly available tools exist for bioinformatics processing of large sets of sequencing data, amongst which QIIME (Caporaso et al., 2010) and mothur (Schloss et al., 2009) are the most commonly used, but some studies have highlighted that an inappropriate choice of software and settings may heavily affect the final results (Majaneva et al., 2015; Anslan et al., 2018). Also for diatom communities, recent studies have suggested that the choice of bioinformatics pipelines may affect the outcome of metabarcoding studies (Tapolczai et al., 2019a; Rivera et al., 2020).

Here, we explore whether the characterization of diatom community structure via metabarcoding is dependent on the quantity of sediment used for DNA extraction and bioinformatics pipeline. Two most commonly used DNA isolation kits for environmental samples, PowerMax Soil and Power Soil (Qiagen), are utilized by applying those to 10 g and 0.5 g (wet weight) of surface sediment samples, respectively. We further tested the consistency of the metabarcoding results obtained via three different bioinformatics pipelines by applying exact sequence variants (ESV) and two OTU clustering approaches. In addition, we assess how the metabarcoding data sets (from 10 g vs. 0.5 g of sediments) compare with the morphological analyses of diatoms from the same samples, and how these datasets relate with environmental variables. Diatom communities were studied from Lake Nam Co, a saline lake on the Tibetan Plateau, and from nearby ponds and tributaries. The preservation of fragile valves of some diatom taxa may be poor in saline, high pH lakes (Flower, 1993) and to the best of our knowledge this is the



**Fig. 2.** Illustration of the study design. HTS 10 and HTS 0.5 represent 10 g and 0.5 g (wet weight) treatments for metabarcoding. Sequencing data from metabarcoding were subjected to three different bioinformatics pipelines (ESVs, 95% and 97% OTUs). Morphological analyses of the diatoms included light microscopy (analyses of 400 valves) and scanning electron microscopy (SEM).

first study of exploring the consistency of microscopy and metabarcoding in the saline Tibetan lakes.

## 2. Materials and methods

### 2.1. Study area

Nam Co is a dimictic and endorheic lake at high altitude ( $\sim 4,730$  m a.s.l.) in the central Tibetan Plateau (Fig. 1; Wang et al., 2010). The lake has surface area of  $2020 \text{ km}^2$  and  $10,789 \text{ km}^2$  catchment area making it as the third largest lake in TP, with a maximum water depth of 98.9 m in the central part (Zhang et al., 2013; Wang et al., 2020). Nam Co locates in the monsoon-influenced transition zone between semi-humid and semi-arid areas (Wang et al., 2020). The annual mean air temperature between 2006 and 2017 at Nam Co Monitoring and Research Station (NAMORS) was  $-0.6^\circ\text{C}$  and the annual mean precipitation was 406 mm (range from 291 to 568 mm) (Anslan et al., 2020). The large glacial areas in the southwestern catchment ( $\sim 700 \text{ km}^2$ ) and melting water generates rivers to the lake. Nam Co is a saline and alkaline with characterized by pH of 7.8–9.5 and a conductivity of  $1920 \mu\text{S mm}^{-1}$  (Keil et al., 2010).

### 2.2. Sample collection

Surface sediment samples were collected in Nam Co using an Ekman-Birge bottom sampler (Hydro-Bios) from water depths ranging between 0.2 m and 56 m (Table S1). A spatula was used to sample about top 2 cm layer of sediments. Additional samples from shallow water sites such as rivers and lagoons were collected randomly from sandy substrate by using a hand-shovel and scraping the upper 2 cm. Approximate wet weight of a sample was 200 g, which was mixed, split in half, and these two batches were then transferred to Whirl-Pak bags for metabarcoding and morphological identification of diatoms. Water temperature, pH and conductivity were measured at each sampling site using a multi parameter probe WTW 3630 (Table S1). Water anions and cations were measured using ion chromatography (IC) and inductively coupled plasma optical emission spectrometry (ICP-OES), respectively, at the Institute of Geographical Sciences, Freie Universität Berlin (Table S1).

In the field, laboratory samples for the morphological identification of diatoms were stored at  $4^\circ\text{C}$ . Samples for metabarcoding analyses were sieved through 2 mm sieves to remove coarser sediment components. Tap water was used for sieving, and therefore also as negative extraction (and PCR) control for metabarcoding analysis. Approximately, 50 g of sediment were divided between three 50 ml tubes and filled with 96% ethanol (4:1 ethanol:sediment ratio). All used

equipment was bleached (10% sodium hypochlorite solution) after each step to avoid cross-contamination. All samples were stored and transported in a freezer ( $-20^\circ\text{C}$ ). The study design is illustrated in Fig. 2.

### 2.3. Molecular analysis

Sediment samples were centrifuged at 4000 rpm for 10 min, supernatant was removed and subsamples were mixed. Wet samples were weighted to 10 g (in the following referred to as metabarcoding treatment HTS 10) and 0.5 g (HTS 0.5), and sedDNA was extracted using DNeasy PowerMax Soil Kit and DNeasy PowerSoil Kit (Qiagen, Germany), respectively (three samples that were processed with PowerMax Kit had  $< 10$  g input, see Table S1). Except for the amount of chemicals, these kits use identical chemistry and protocols. To enhance the cell lysis, we modified the initial step by adding Proteinase K (10 mg/ml) and 1 M DTT (dithiothreitol) together with the C1 solution from the extraction kits. For the PowerMax Kit (10 g of sediments) 60  $\mu\text{l}$  of Proteinase K and 100  $\mu\text{l}$  of DTT, and for the PowerSoil Kit (0.5 g of sediments), 4  $\mu\text{l}$  of Proteinase K and 25  $\mu\text{l}$  of DTT was added, respectively; following overnight incubation at  $56^\circ\text{C}$ . For potentially higher DNA yield, the elution was performed twice by adding half of the recommended amount of the buffer onto a spin column membrane and incubated at room temperature for 3 min. The rest of the steps were performed following manufacturer's instructions.

PCRs were performed using uniquely tagged primers rbcL-646F (5'-ATG CGT TGG AGA GAR CGT TTC-3') and rbcL-998R (5'-GAT CAC CTT CTA ATT TAC CWA CAA CTG-3'), which amplify 331 base pairs (bp) of the large subunit of the ribulose-bisphosphate carboxylase/oxygenase (rbcL) gene (Kelly et al., 2018) (Table S2). We also tested the primers of Diat\_rbcL\_708F and R3 (Vasselon et al., 2017b), which amplify a shorter fragment (312 bp) of the same region. However, the PCR results were visually superior for rbcL-646F and rbcL-998R primer pair (data not shown), thus here, we decided to proceed only with the latter primers. The 25  $\mu\text{l}$  PCR mix consisted of 5  $\mu\text{l}$  of Hot Start FirePol Master Mix (Solis BioDyne, Estonia), 0.5  $\mu\text{l}$  forward and reverse primer (10  $\mu\text{M}$ ), and 1–3  $\mu\text{l}$  of template DNA. The rest of the volume was filled with nuclease-free water. PCR conditions were as follows:  $95^\circ\text{C}$  for 15 min (hot start), 32–35 cycles of  $95^\circ\text{C}$  for 20 s,  $55^\circ\text{C}$  for 45 s,  $72^\circ\text{C}$  for 60 s, and final extension at  $72^\circ\text{C}$  for 5 min. Three replicate PCRs were performed per sample, following sample pooling and checking the yield of PCR products during gel electrophoresis by pipetting 5  $\mu\text{l}$  PCR product on 1% agarose gel. Amplicons per sample were pooled as based on their relative quantity and purified using Favor-Prep™ Gel/PCR Purification Kit (Favorgen-Biotech Corp., Austria), following the manufacturer's

instructions. Steps of DNA extraction, PCR and sequencing included both negative and positive controls. Sample preparations, as well as DNA isolations, were conducted under laminar flow clean bench, using 30 min UV sterilization prior analyses. Sequencing was performed on the Illumina MiSeq (2×250) using MiSeq Reagent Kit v2.

## 2.4. Bioinformatics

Three different bioinformatics workflows (pipelines) were used to process raw paired-end Illumina data: 1) ESVs (exact sequence variants) pipeline as implemented in DADA2 (Callahan et al., 2016); 2) 95% OTUs (operational taxonomic units) pipeline, where OTUs are clustered at 95% sequence identity; and 3) a pipeline based on OTUs clustered at 97% identity. The processing of sequencing data to generate ESVs and 95% OTUs followed the workflows as described in Tapolczai et al. (2019a) and Rivera et al. (2020), respectively, except that taxonomy assignment of the representative sequences was performed using blastn algorithm (instead of Naïve Bayesian classifier) (Camacho et al., 2009) with  $e\text{-value} = 0.001$ , word size = 7, reward = 1, penalty = -1, gap open = 1 and gap extend = 2 (against R-Syst v.7.2 diatom database (Rimet et al., 2016)). Based on our positive and negative controls, the 95% OTUs data set was further filtered to exclude low occurrence ( $\leq 3$ ) reads per OTU per sample to alleviate to 'tag-switching' error. The latter was not performed for the ESVs data set as no sequences were observed in the negative controls and no 'read-leakage' from the positive control sample. Singleton ESVs/OTUs were discarded from the data sets (i.e. ESVs/OTUs that had only one read across samples).

To generate 97% OTUs, raw paired-end Illumina sequencing data was processed in PipeCraft (Anslan et al., 2017), which incorporates all the following tools (except LULU). Reads were assembled and quality filtered using vsearch (fastq\_minoverlen 15, maxdiffs 45, fastq\_minmergelen 200, fastq\_maxee 1, fastq\_maxns 0, fastq\_truncqual 5, fastq\_allowmergestagger) (Rognes et al., 2016). Chimera filtering step included vsearch uchime\_denovo algorithm with options id 0.97 and abskew 2. The filtered sequences were clustered using UPARSE (Edgar, 2013) with 97% similarity threshold and minimum cluster size of 2 (i.e. singletons excluded). The obtained OTU table was further curated with post-clustering algorithms as implemented in LULU (Frøslev et al., 2017) to merge consistently co-occurring 'daughter' OTUs (minimum\_ratio = 1, minimum\_ratio\_type = "avg", minimum\_relative\_cooccurrence = 0.8, minimum\_match = 96.97). Potential tag-switching errors were also corrected based on negative and positive controls based on relative abundances of sequences in the control samples (Taberlet et al., 2018). To account for unequal sequencing depth, we rarefied samples to common depth of 7000 sequences using the mothur software (Schloss et al., 2009). The latter removed five samples from the data set. Representative sequences per OTU were compared against R-Syst v.7.2 diatom database using blastn as stated above.

The used primers (rbcl-646F and rbcl-998R) amplify DNA also from other algae and bacteria (especially Proteobacteria and Cyanobacteria). To exclude the non-target taxa, only OTUs that demonstrated the match coverage and identity of  $\geq 90\%$  against a reference database, were considered as diatom OTUs and included in the final tables produced by each pipeline. According to additional blastn search against NCBI database (Geer et al., 2010), the above threshold was confirmed to include only diatom taxa into the final OTU table. OTUs with lower thresholds to reference diatom sequences (in R-Syst) were often more closely related (based on  $e\text{-value}$ , sequence similarity and coverage) to other micro-algae (e.g. taxa from Mischozoales, Tribonematales, Eustigmatophyceae), thus excluded from the downstream analyses.

Because of the uncertainty of the most adequate species-level sequence similarity threshold for diatoms, the taxonomic composition comparisons between metabarcoding treatments (HTS 10 and HTS 0.5) and microscopy was performed on genus level. Reliable genus level classification of the OTUs in the HTS data set was here defined when sequence similarity and coverage was  $\geq 95\%$  against a reference

sequence in the R-Syst database. OTUs that displayed lower values against the R-Syst database sequences were blastn-searched against the NCBI database to check for additional genus-level annotations. Synonym names for genera were also explored in the case of mismatches between microscopy and metabarcoding data sets.

## 2.5. Morphological analysis

Sediment samples for morphological diatom analyses were treated and examined using standard methods (Battarbee et al., 2001). Specifically, approximately 0.1 g of dry sediment (freeze dried) for each sample was treated with 30% hydrogen peroxide ( $\text{H}_2\text{O}_2$ ) and 10% hydrochloric acid (HCl) for 2–3 h to remove organic matter and carbonates, respectively. The resulting slurries were then washed repeatedly with distilled water until a neutral pH was reached and stewed onto glass coverslips to dry at room temperature. The dry samples were then fixed onto microscope slides with Naphrax®, a highly refractive mountant. At least 400 valves were identified to species or genus level (range from 400 to 407) and enumerated along transects using a light microscope (Leica DM6B, magnification  $\times 1000$ ) with oil immersion objective. Diatom identification and taxonomy were based primarily on general floras such as Krammer and Lange-Bertalot (1986–1991), Lange-Bertalot et al. (2017), Mohan et al. (2018) for *Lindavia biswashanti*, and Pavlov et al. (2013) for species belonging to the *Hippodonta* genus. Scanning electron microscope (SEM) photographs were taken using a Zeiss MERLIN instrument at the Institute of Geology and Geophysics, CAS, Beijing.

## 2.6. Statistics

Differences in the ESVs/OTUs/morphospecies richness and correlation analysis between the three treatments (HTS 10, HTS 0.5 and microscopy) were tested using Kruskal-Wallis Analysis of Variance (ANOVA) by Ranks and Spearman Rank Order Correlations in STATISTICA 7 (StatSoft, 2004). Differences in diatom community assemblies were tested using Permutational Analysis of Variance (PERMANOVA, with 9999 permutations) and Principal Co-ordinate Analysis (PCoA) of Bray-Curtis similarity matrices of log transformed data (performed in PRIMER v6; Clarke and Gorley, 2006). The values of the PCoA first axis were selected as a response variable for Random Forest analysis to identify most important environmental variables for diatom community assembly. The latter was performed in R (R-Core-Team, 2015), using the package 'randomForest' (Liaw and Wiener, 2002). The variables included for Random Forest analyses were maximum depth, conductivity, and concentration of zinc (Zn), calcium (Ca), silica (Si), chromium (Cr), nitrate ( $\text{NO}_3$ ), copper (Cu), manganese (Mn), iron (Fe), strontium (Sr), vanadium (V) (Table S1). These variables were selected based on correlation matrix (Fig. S1; conducted using 'correlplot' package (Wei et al., 2017) in R) to exclude highly correlated ones. Variables, selected by Random Forest model per treatment, were used in marginal test analysis (with 9999 permutations), in PRIMER. Only variables with P-values  $< 0.05$  were displayed as vectors in the PCoA ordination plots.

Bray-Curtis similarity matrices of Hellinger-transformed data per treatment were compared with Mantel tests (with 9999 permutations, method="spear") to assess the correlations between sample similarities as implemented in the 'vegan' package (Oksanen et al., 2015). In addition, Procrustes analyses were used to compare the similarity of metaMDS ordinations between different treatments (in 'vegan'). To examine the presence of diatom OTUs that were consistently detected either by the HTS 10 or the HTS 0.5 treatment, an indicator species analysis (with 9999 permutations) was performed using the 'indicspecies' package (De Caceres et al., 2016).

The analyses for comparing metabarcoding treatments (HTS 10 and HTS 0.5) were performed for 20 corresponding samples. Because of smaller sample size for the microscopy data, analyses for HTS 10 vs.



**Table 1**

Number of ESVs/OTUs/morphospecies per sample for metabarcoding and microscopy data. TOTAL11 denotes the sum of ESVs/OTUs/species for 11 comparable samples of all data. TOTAL 20 denotes the sum of ESVs/OTUs for 20 comparable samples of metabarcoding data. N.a denotes 'not available' data; thus the last three samples were not included to HTS 10 vs. HTS 0.5 comparisons, and the analyses of microscopy vs. HTS treatments were conducted using corresponding samples with available data.

Sample	ESVs (HTS10/ HTS0.5)	95% OTUs (HTS10/ HTS0.5)	97% OTUs (HTS10/ HTS0.5)	Microscopy
1	134/125	73/76	53/49	32
2	107/39	63/32	53/23	41
3	127/112	71/69	55/50	42
4	273/261	145/137	120/128	n.a
5	60/88	45/59	36/48	n.a
6	101/133	70/77	52/60	n.a
7	38/28	27/26	18/14	28
8	219/273	117/136	103/118	n.a
9	221/251	116/122	103/102	n.a
10	273/272	131/133	115/120	n.a
11	182/169	111/100	101/91	n.a
12	137/139	79/81	67/64	n.a
13	82/81	53/57	47/48	n.a
14	187/193	101/100	82/79	41
15	41/69	32/46	20/28	38
16	102/29	64/21	54/13	34
17	185/175	105/98	81/82	43
18	73/87	53/66	47/52	46
19	139/110	86/78	64/53	41
20	53/68	37/44	29/35	57
21	92/n.a	59/n.a	44/n.a	42
22	50/n.a	41/n.a	34/n.a	44
23	11/n.a	64/n.a	46/n.a	28
TOTAL 11	474/433	175/178	165/161	189
TOTAL 20	1011/1018	297/307	306/318	n.a

microscopy and HTS 0.5 vs. microscopy were performed with 14 and 11 samples, respectively (see Table S1).

### 3. Results

#### 3.1. Richness and composition

The numbers of ESVs/OTUs/morphospecies per sample for metabarcoding and microscopy data are summarized in Table 1. The numbers of ESVs/OTUs per sample between HTS 10 and HTS 0.5 treatments demonstrated a strong positive correlation (Spearman  $R > 0.863$ ; Fig. 3a-c; Fig. S2), and no significant differences in the (ESV/OTU) richness (Fig. 3d). Accordingly, the intra-pipeline comparisons of HTS 10 and HTS 0.5 data (20 comparable samples; Table 1) demonstrated high proportions of shared ESVs/OTUs (63.9–87%; Fig. 4). For the genus level comparisons across metabarcoding data sets, ESVs and OTUs were annotated to genus level only when the similarity and coverage of the representative read of ESV/OTU was  $\geq 95\%$  against a reference sequence. Similarly, a large proportion of genera were shared between HTS treatments (87.7–92.2%; Fig. 4). Interestingly, inter-pipeline comparisons revealed that higher proportion of genera was shared between HTS 0.5 treatments compared with HTS 10 (87.7% vs. 76.4%; Fig. S3). Compared to data generated with the OTUs pipelines, the data from the ESVs pipeline harbored a higher number of different genera (51 vs. 47 for HTS 10 and 50 vs. 47 for HTS 0.5; Fig. S3). For the HTS 10 data, the unique genera (3 genera; i.e. genera that were identified only in the corresponding data set) of the ESVs data set represented a total of less than 0.01% of sequences (Table S3). For the HTS 0.5 data, the unique

genera (2 genera) of the ESVs data set represented also a total of less than 0.01% of sequences (Table S3). The data set of 95% and 97% OTUs did not contain any unique genera (Fig. S3; Table S3).


Morphological examination of the sediment samples recovered a total of 189 diatom taxa from 11 surface sediment samples (Table 1), which included 59 genera (Fig. 5; Table S3). Unlike the per sample richness correlations observed between the metabarcoding treatments (Fig. 3a-c), correlations were not obvious between richness values from the microscopy and metabarcoding data ( $P > 0.398$  for all cases; Fig. S4). Across treatments, detected species richness by microscopy differed significantly only from the ESVs data (Fig. 3d). The detected composition of genera by microscopy were compared with metabarcoding data, which harbored 54 different genera for the ESVs data, 50 and 49 genera for the 95% and 97% OTU data, respectively. The genus level comparisons (among 11 corresponding samples) revealed that 50.7–54.3% of genera were shared between microscopy and metabarcoding treatments (Fig. 5). Compared with the metabarcoding inventories, the microscopy data set harbored larger proportion of unique genera (Fig. 5). From these, the majority were represented in low abundances in the microscopy data set ( $< 9$  counted valves per sample). However, counts of the valves assigned to *Pseudostaurosira*, one of the most abundant genera that were completely missing from metabarcoding data, was 519 (11.77%) across the microscopy data set.

Comparison of the relative abundance of valves and sequences of the matching genera between microscopy and metabarcoding data, revealed overall significant positive correlations (Spearman  $R > 0.317$  and  $P < 0.023$ ; except for 97% OTUs HTS 10 vs. microscopy data, where  $P = 0.067$ ; Fig. S5). The outstanding exceptions were *Pantocsekiella* and *Achnanthisidium*, which had high relative abundance in microscopy, but low abundance in metabarcoding data (Fig. S5). *Vice versa*, *Staurosira* and *Aulacoseira* were found to have high relative abundance in metabarcoding data, but low in microscopy data (Fig. S5).


#### 3.2. Community analyses

Mantel test and Procrustes analyses revealed that the inter-sample community similarities between metabarcoding treatments were greatly correlated (Fig. 6; Table 2). For all cases, the Mantel correlation (i.e. Mantel  $R$ ) between HTS 10 and HTS 0.5 was higher than 0.851 and  $P < 0.001$  (Fig. 6). Among intra-pipeline treatments (HTS 10 vs. HTS 0.5), 95% OTUs exhibited highest correlation (Mantel  $R = 0.969$ ; Table 2; Fig. 6). Procrustes correlations, however, revealed highest values between ESVs treatments (HTS 10 vs. HTS 0.5; Procrustes correlation = 0.969,  $P < 0.001$ ; Table 1), but a much lower value between 95% OTUs treatments (Procrustes correlation = 0.688; Table 1). The high community similarity among the metabarcoding data sets was also demonstrated by PERMANOVA analyses using treatment as fixed variable ( $P > 0.999$  for all cases). Moreover, no group-specific indicator OTUs were assigned to neither HTS 10 nor HTS 0.5 treatments by indicator species analysis for any of the metabarcoding data sets.


The inter-sample community similarities (Mantel correlations) and Procrustes analyses between microscopy and HTS data also demonstrated highly correlated patterns (microscopy vs. all HTS treatments: Mantel  $R \geq 0.800$ ,  $P < 0.001$ ; Procrustes correlations  $\geq 0.681$ ,  $P \leq 0.001$ ; Fig. 7, Table 1). Compared with other treatments, the 95% OTUs data set demonstrated slightly higher Mantel correlations with microscopy data (Table 1), whereas the highest Procrustes correlations were found for the ESVs HTS 0.5 data set (Table 1). Based on Random Forest analysis, the diatom assemblages in all treatments were most strongly affected by conductivity, water depth, Si, Ca, Sr, Mn and Fe, however, with different orders in variable importance (Fig. S6). Marginal tests (the significance of an individual variable when considered alone and ignoring all other variables) showed consistent patterns for the most



**Fig. 3.** a-c) scatterplots between per-sample richness of taxonomic units from HTS 10 and HTS 0.5 treatments. d) box plot for the number of ESVs/OTUs/morphospecies (log transformed) per treatment. Different letters above the whiskers indicate significant differences according to Kruskal-Wallis pairwise test.



**Fig. 4.** Venn diagrams of the diatom genera relations between HTS 10 and HTS 0.5 treatments (20 corresponding samples).



**Fig. 5.** Venn diagrams of the diatom genera relations between treatments of HTS 10, HTS 0.5 and microscopy (11 corresponding samples). Upper diagrams demonstrate the distribution of ESVs or OTUs in the corresponding treatments; lower diagrams show the distribution of diatom genera in the corresponding treatments.

important variables for all treatments (Table S4). The highly correlated community structures were also demonstrated in the ordination plots (Fig. 8).

#### 4. Discussion

In the current study, regardless of sample size (10 g, 0.5 g) and bioinformatics pipeline (ESVs, 97% OTUs, 95% OTUs) applied, we found highly comparable diatom community patterns between metabarcoding data and microscopic inventories from lake sediment samples. This was especially pronounced between the metabarcoding treatments, i.e., amplicons from 10 g and 0.5 g of DNA extracts (HTS 10 and HTS 0.5, respectively). Although the per-sample diatom richness was not significantly correlated between microscopy and metabarcoding data, community analyses indicated consistent patterns of environmental variables shaping the diatom community structures, irrespective of sample-size and bioinformatics pipeline used. This further demonstrates that metabarcoding is a viable method to describe diatom-environment relationships.

##### 4.1. Sample size

Although the DNA extraction method may have a strong impact on taxa recovery in metabarcoding studies (Schiebelhut et al., 2017; Sinha et al., 2017), previous comparisons of different DNA isolation kits for diatoms have demonstrated comparable patterns for diversity and community assembly (Vasselon et al., 2017a). The present study also shows that richness and community structure of diatoms are highly correlated between metabarcoding data of 10 g vs. 0.5 g of sediment samples. Interestingly, the correlations of relative abundances of matching diatom genera between metabarcoding and microscopy data sets, resulted in higher correlation values for the HTS 0.5 data (Fig. S5). However, other studies comparing metabarcoding results from DNA extracts of various amounts of substrate have reported contrasting results. For example, Penton et al. (2016) reported significant effects of sample size (in terms of input quantity) on the community structure of fungi and bacteria from soil. Higher diversity estimates were associated with 10 g of soil DNA extracts compared with 5 g, 1 g and 0.25 g. Studying meiofaunal communities from marine sediment samples, Brannock and Halanach (2015) found that different extraction quantities did not result in significantly different diversity estimates, however, the OTU community compositions were different. Exploring various eukaryotes from sediment samples, Nascimento et al. (2018) also found significantly different diversity metrics and community compositions for various sample sizes. They suggested that larger volumes of sediment are

necessary to capture the representative metazoan communities compared to the non-metazoan eukaryotes. Therefore, the choice of a quantity of sediment for DNA extraction may depend on the expected distribution of the target groups in the substrate, where the detection of more patchily distributed metazoan communities requires larger quantities of sediment for analyses. Although it could be hypothesized that sample size may affect also the recovery of some microbial groups, here we demonstrate that this was not the case for diatoms from lake sediment samples (when comparing sample size on 10 g vs. 0.5 g). The benefits of using only up to 0.5 g of the sample include a more time- and cost-effective DNA extraction procedure, and the possibility to conduct meaningful analyses when only a limited amount of sediment is available.

##### 4.2. Bioinformatics

Here, three different bioinformatics pipelines were used to analyze metabarcoding data. Combination of analyses among metabarcoding data sets and between microscopy vs. metabarcoding data demonstrated highly correlated patterns (Table 2, Figs. 3–7). Although the comparisons between microscopy and metabarcoding data demonstrated highest Mantel correlation of the former with the 95% OTUs data set, Procrustes correlation was highest with the ESVs data set (Table 2). Thus, the identification of the best performing pipeline, in terms of consistency with microscopic inventories, may depend on the applied statistical method. Nevertheless, considering the genus level comparisons from metabarcoding, the ESVs data set had the slightly higher number of genus level identifications, and therefore a (marginally) higher match proportion with microscopy data. But interestingly, the ESVs data set harbored also one diatom taxon that is considered to be strictly marine, *Trachyneis* sp. (Fig. 4; Table S3). ESV assigned to the latter taxon, however, consisted of only three reads across the whole data set. It is uncertain whether this low abundance ESV assigned to *Trachyneis* sp. represents remaining sequencing errors or real occurrences, but raises caution in evaluating data quality based on highest taxonomic richness alone. In any case, the differences between bioinformatics workflows in this study were minor and indicated highly similar signals among all of them.

Although this study displayed an overall consistency of results obtained by different pipelines, several studies have demonstrated the influence of bioinformatics in metabarcoding studies targeting other organisms (Majaneva et al., 2015; Sinha et al., 2017; Anslan et al., 2018; Pauvert et al., 2019). The impacts may originate from inadequate error filtering processes (Edgar, 2017), inaccurate taxonomic annotation (Anslan et al., 2018) or inappropriate clustering methods in the specific




Fig. 6. Mantel correlation plots between HTS 10 and HTS 0.5 treatments.



**Table 2**

Mantel test and Procrustes analyses results between all treatments. HTS 10 and HTS 0.5 represent metabarcoding data from 10 g and 0.5 g of sediments, respectively.

Treatment 1	Treatment 2	Mantel R	Mantel P value	Procrustes correlation	Procrustes P value
95% OTUs HTS 10	95% OTUs HTS 0.5	0.9688	<0.001	0.688	<0.001
95% OTUs HTS 0.5	ESVs HTS 0.5	0.936	<0.001	0.933	<0.001
97% OTUs HTS 0.5	95% OTUs HTS 0.5	0.96	<0.001	0.831	<0.001
97% OTUs HTS 0.5	ESVs HTS 0.5	0.902	<0.001	0.888	<0.001
ESVs HTS 10	ESVs HTS 0.5	0.951	<0.001	0.969	<0.001
ESVs HTS 10	95% OTUs HTS 0.5	0.918	<0.001	0.929	<0.001
ESVs HTS 10	97% OTUs HTS 0.5	0.875	<0.001	0.821	<0.001
95% OTUs HTS 10	ESVs HTS 10	0.949	<0.001	0.777	<0.001
95% OTUs HTS 10	97% OTUs HTS 0.5	0.935	<0.001	0.689	<0.001
95% OTUs HTS 10	ESVs HTS 0.5	0.914	<0.001	0.705	<0.001
97% OTUs HTS 10	95% OTUs HTS 10	0.9686	<0.001	0.827	<0.001
97% OTUs HTS 10	97% OTUs HTS 0.5	0.937	<0.001	0.586	0.0015
97% OTUs HTS 10	95% OTUs HTS 0.5	0.934	<0.001	0.766	<0.001
97% OTUs HTS 10	ESVs HTS 10	0.901	<0.001	0.756	<0.001
97% OTUs HTS 10	ESVs HTS 0.5	0.852	<0.001	0.71	<0.001
Microscopy	ESVs HTS 0.5	0.865	<0.001	0.818	0.003
Microscopy	ESVs HTS 10	0.800	<0.001	0.777	0.001
Microscopy	95% OTUs HTS 0.5	0.889	<0.001	0.813	0.002
Microscopy	95% OTUs HTS 10	0.893	<0.001	0.681	0.007
Microscopy	97% OTUs HTS 0.5	0.879	<0.001	0.788	0.007
Microscopy	97% OTUs HTS 10	0.835	<0.001	0.684	0.008

case of diatoms (Tapolczai et al., 2019b). Towards standardization of analyzing short (312 bp) rbcL metabarcoding data of diatoms for bio-monitoring purposes, the studies of Tapolczai et al. (2019a) and Rivera et al. (2020) found that individual sequence units (ISU) approach tend to outperform operational taxonomic units (OTU) based approaches. Nevertheless, furthest neighbor OTU clustering and ESVs approach showed to perform equally well (Rivera et al., 2020). Our study also resulted in highly similar results across three applied pipelines (ESVs vs. two OTU approaches), which demonstrates that the appropriate filtering of erroneous sequences and critical taxonomic assignment of the target taxa may be a key step, with the potential of mitigating the otherwise considerable effect of bioinformatics.

#### 4.3. Taxonomic composition

Because different data sets in this study included ESVs, OTUs or morphospecies, the direct comparisons of their taxonomic unity were performed at the genus level. In this study, metabarcoding results from 10 g and 0.5 g of sedDNA extracts exhibited highly concurring taxonomic composition, with only few mismatched taxa, which were represented by a low relative abundance of reads. In accordance with the community analyses, this indicates sample-size independent patterns when detecting diatoms via metabarcoding from lake sediments.

However, comparisons between microscopy and metabarcoding data resulted in a higher number of mismatched taxa, ranging from 20 to 23 genera being detected only by microscopy (Fig. 5). Not completely matching identifications from microscopy vs. metabarcoding have been reported in several previous diatom-related studies (e.g. Visco et al., 2015; Rivera et al., 2018; Tapolczai et al., 2019b), with the possible reasons discussed within. One of the main reasons of such mismatches is the incompleteness of the reference sequence databases, which consists of a limited number of annotated taxa. For example, *Sichuaniella lacustris*, discovered only by morphological analyses, is the unique representative of the genus *Sichuaniella*, which was originally described from Sichuan Province on the southeastern Tibetan Plateau (Li et al., 2013) and has no genetic information in the public databases. Therefore, the identity of this species in the metabarcoding data set cannot be confirmed. Additionally, there are no reference sequences for genera such as *Platessa*, *Odontidium* and *Gomphosinica* in the public databases. *Gomphosinica* has been separated from *Gomphonema* and described as a new genus based only on their morphological differences (Kocielek et al., 2015). Thus, *Gomphosinica* in the microscopy data set could potentially be represented as *Gomphonema* in the metabarcoding data set.

The inter-investigator variation depending on changes in diatom taxonomy and the use of synonym names could add additional layers for the mismatches between microscopy and metabarcoding data. In this study, it is difficult to consistently separate *Staurosirella* and *Pseudostaurosira* (missing from metabarcoding data) from *Staurosira* (present in metabarcoding data) under the light microscope and even with support from SEM images. Although *Pseudostaurosira* was one of the most abundant genera in the microscopy data, it was missing from the metabarcoding inventories, whereas the relative abundance of *Staurosira* was high in latter data sets (Table S3). Medlin et al. (2012) have pointed out that the molecular separation of *Pseudostaurosira* and *Staurosirella* from *Staurosira* is arguable. On the other hand, in the few studies that have attempted to merge morphological- and molecular-based phylogenies of the Fragilariaceae, the morphological characterization is often poorly done (Morales et al., 2019). We speculate that morphologically identified *Pseudostaurosira* (especially *Pseudostaurosira brevistriata*) corresponds to *Staurosira* in the metabarcoding data, as their presence-absence patterns in our sediment samples correlates well (Table S3). Furthermore, identification of *Pseudostaurosira* in the metabarcoding data sets was also limited due to a fact that almost all originally named *Pseudostaurosira* were re-assigned to *Staurosira* in the curated R-Syst diatom database (Rimet et al., 2016).

The majority of other missing genera from metabarcoding data sets were represented in very low abundances in the microscopy data. Similarly, Kermarrec et al. (2013) reported that morphologically identified low abundance taxa (<1% from 450 valve counts) were often not detected in the DNA metabarcoding data set. These low abundance taxa may indicate the transport of diatom valves with highly degraded DNA from other locations (thus non-detectable with herein used primers). On the other hand, environmental DNA could be carried along large distances (Deiner et al., 2014), which also could contribute to the observed 'extra' diatom taxa in metabarcoding data sets, which were not detected via microscopy. These issues could contribute to the observed pattern of uncorrelated per-sample diatom taxonomic unit richness between microscopy and metabarcoding. Moreover, some of the diatom taxa with fragile and weakly silicified valves, such as *Cylindrotheca*, *Entomoneis*, *Fistulifera*, *Reimeria*, *Seminavis*, that were detected only in the metabarcoding data sets, might be sensitive to the chemical treatment (e.g. HCl and H<sub>2</sub>O<sub>2</sub>) during sample preparation for microscopy. Based on personal observations of water samples from Nam Co, we have confirmed the presence of several *Entomoneis* and *Fistulifera* species (data not shown), which further supports the assumption that valves of fragile diatom species may be more prone to dissolution and therefore undetectable in sediment samples. Thus, incompleteness of the reference databases, together with the continuously changing diatoms




Fig. 7. Mantel correlation plots between microscopy data and metabarcoding treatments (HTS 10 and HTS0.5).

classification system and DNA transportation characterizations contribute to at least some extent to the issue of non-matching taxa between microscopy and metabarcoding results.

#### 4.4. Perspectives

Assemblages of diatom communities reflect environmental parameters (Dixit et al., 1992), and therefore they are widely used as paleo-ecological indicators of lake ecosystems (Douglas et al., 2010). We found that the diatom communities within sediment samples from Nam Co can be related to the same environmental variables for both, the morphological and metabarcoding data sets, which is in accordance with the study by Dulas et al. (2017). This suggests that inferring (paleo-) environmental characteristics (via, for example, diatom-based transfer functions) would produce similar results using either method, where the high-throughput nature of metabarcoding analyses, however, enables simultaneous processing of much larger numbers of samples in a time-effective manner (Stoof-Leichsenring et al., 2014; Dulas et al., 2017). Although not tested here, the additional ‘fine-tuning’ of the metabarcoding data with e.g. quantification correction factors or including phylogeny of the OTUs has been suggested to further improve the (biomass) correlations between microscopy and metabarcoding results (Vasselon et al., 2018; Mortágua et al., 2019) as well as boost the applicability of the latter for biomonitoring purposes (Keck et al., 2018). Because of the incompleteness of available DNA barcode databases, taxonomy-independent methods for molecular taxa are another promising advancement towards the applicability of metabarcoding in environmental surveys (Apothélos-Perret-Gentil et al., 2017; Tapolczai


et al., 2019b). Moreover, when the preservation of diatom valves is poor, as for example in saline, high pH lakes with low sediment accumulation rates (Flower, 1993), DNA may still preserve in sediments as for example has been demonstrated by studying ‘non-fossilizing’ phytoplankton by means of sedimentary ancient DNA (Li et al., 2016). The strong similarity of our metabarcoding results from 10 g and 0.5 g of DNA extracts implies that ‘small’ DNA isolation kits (for ~ 0.5 g) may serve as an alternative approach when the amount of sediments is limited in sedimentary ancient DNA (sedaDNA) studies.

#### CRedit authorship contribution statement

**Wengang Kang:** Investigation, Formal analysis, Writing - original draft. **Sten Anslan:** Conceptualization, Formal analysis, Visualization, Writing - original draft. **Nicole Börner:** Investigation, Writing - review & editing. **Anja Schwarz:** Investigation, Writing - review & editing. **Robin Schmidt:** Investigation. **Sven Künzel:** Investigation. **Patrick Rioual:** Methodology, Writing - review & editing. **Paula Echeverría-Galindo:** Writing - review & editing. **Miguel Vences:** Writing - review & editing. **Junbo Wang:** Writing - review & editing. **Antje Schwalb:** Funding acquisition, Supervision, Writing - review & editing.

#### Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.



**Fig. 8.** Ordination plots of diatom communities for all treatments. a-d) plots for corresponding 11 samples between microscopy and HTS 0.5 data; e-h) plots for corresponding 14 samples between microscopy and HTS 10 data. Vectors on the plots denote most important variables as based on Random Forest modelling and DistLM marginal tests. Length of the vector represents the importance of variable.

### Acknowledgements

We thank Jinlei Kai and Magnus Ole Asmussen for help during fieldwork, and Philipp Hoelzmann for water chemistry analysis, and Eike Reinosch for geographical map in Fig. 1. We are indebted to the team of the NAMORS station of the Institute of Tibetan Plateau Research of the Chinese Academy of Sciences, in particular Guangjian Wu, and to the entire team of the International Research Training Group “Geoecosystems in transition on the Tibetan Plateau” (TransTiP), funded by the Deutsche Forschungsgemeinschaft (DFG-GRK 2309), for logistic support and fruitful discussion. We also thank the Institute of Tibetan Plateau Research, Chinese Academy of Sciences by supplying the research permission obtained from the Tibet Autonomous Region Government.

### Data accessibility

Illumina data sets have been deposited in the Sequence Read Archive (SRA) under BioProject PRJNA558871.

### Authors' contributions

W.K., N.B. and A. Schwarz performed sampling. R.S. and S.A. performed molecular analyses, S.K. ran high-throughput sequencing, S.A. processed the sequence data. W.K. and P.R. worked on morphological analyses of the diatoms. S.A. and W.K. analyzed the data and performed statistical analyzes. A. Schwalb developed the program (DFG-GRK 2309) that funded this research (DFG grant 317513741). S.A. prepared the manuscript with the contributions from all authors. All authors approved the final version.

### Funding

Funding was provided by the Deutsche Forschungsgemeinschaft

(DFG) via the International Research Training Group GRK 2309 “Geoecosystems in transition on the Tibetan Plateau (TransTiP)”.

### Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.ecolind.2020.107070>.

### References

- Anslan, S., Bahram, M., Hiiesalu, I., Tedersoo, L., 2017. PipeCraft: flexible open-source toolkit for bioinformatics analysis of custom high-throughput amplicon sequencing data. *Mol. Ecol. Resour.* 17 (6), e234–e240.
- Anslan, S., Nilsson, R.H., Wurzbacher, C., Baldrian, P., Tedersoo, L., Bahram, M., 2018. Great differences in performance and outcome of high-throughput sequencing data analysis platforms for fungal metabarcoding. *Mycosyst. Res.* 39, 29–40.
- Anslan, S., Azizi, R.M., Buckel, J., Echeverria G.P., Kai, J., Kang, W., Keys, L., Maurischat, P., Nieberding, F., Reinosch, E., Tang, H., Tran, T.V., Wang, Y., Schwalb, A., 2020. Reviews and syntheses: How do abiotic and biotic processes respond to climatic variations in the Nam Co catchment (Tibetan Plateau)? *Biogeosciences* 17, 1261–1279.
- Apothéoz-Perret-Gentil, L., Cordonier, A., Straub, F., Iseli, J., Esling, P., Pawlowski, J., 2017. Taxonomy-free molecular diatom index for high-throughput eDNA biomonitoring. *Mol. Ecol. Resour.* 17 (6), 1231–1242.
- Baird, D.J., Hajibabaei, M., 2012. Biomonitoring 2.0: a new paradigm in ecosystem assessment made possible by next-generation DNA sequencing. *Molecular Ecology* 21, 2039–2044.
- Battarbee, R., Carvalho, L., Jones, V., Flower, R., Cameron, N., Bennion, H., Juggins, S., 2001. *Tracking Environmental Change Using Lake Sediments: Terrestrial, Algal and Siliceous Indicators*. Kluwer Academic Publishers.
- Brannock, P.M., Halanich, K.M., 2015. Meiofaunal community analysis by high-throughput sequencing: comparison of extraction, quality filtering, and clustering methods. *Mar. Geomorphol.* 23, 67–75.
- Callahan, B.J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson, A.J.A., Holmes, S.P., 2016. DADA2: High-resolution sample inference from Illumina amplicon data. *Nat. Methods* 13 (7), 581–583.
- Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., Madden, T.L., 2009. BLAST+: architecture and applications. *BMC Bioinf.* 10 (1), 421. <https://doi.org/10.1186/1471-2105-10-421>.
- Caporaso, J.G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F.D., Costello, E.K., Fierer, N., Peña, A.G., Goodrich, J.K., Gordon, J.I., Huttley, G.A., Kelley, S.T., Knights, D., Koenig, J.E., Ley, R.E., Lozupone, C.A., McDonald, D., Muegge, B.D.,

- Pirrung, M., Reeder, J., Sevinsky, J.R., Turnbaugh, P.J., Walters, W.A., Widmann, J., Yatsunenko, T., Zaneveld, J., Knight, R., 2010. QIIME allows analysis of high-throughput community sequencing data. *Nat. Methods* 7 (5), 335–336.
- Chen, G., Shi, H., Tao, J., Chen, L.I., Liu, Y., Lei, G., Liu, X., Smol, J.P., 2015. Industrial arsenic contamination causes catastrophic changes in freshwater ecosystems. *Sci. Rep.* 5 (1) <https://doi.org/10.1038/srep17419>.
- Clarke, K., Gorley, R., 2006. PRIMER V6: User manual/tutorial. Primer-E Ltd., Plymouth, p. 192.
- De Caceres, M., Jansen, F., De Caceres, M.M., 2016. Package 'indicspecies'. Relationship between species and groups of sites. R package version 1.7.6.
- Deiner, K., Altermatt, F., 2014. Transport distance of invertebrate environmental DNA in a natural river. *PLoS ONE* 9, e88786.
- Dixit, S.S., Smol, J.P., Kingston, J.C., Charles, D.F., 1992. Diatoms - Powerful indicators of environmental change. *Environ. Sci. Technol.* 26, 22–33.
- Douglas, M.S.V., Smol, J.P., Smol, J.P., Stoermer, E.F., 2010. In: *The Diatoms: Applications for the Environmental and Earth Sciences*. Cambridge University Press, Cambridge, pp. 249–266. <https://doi.org/10.1017/CBO9780511763175.014>.
- Dulias, K., Stoof-Leichsenring, K.R., Pestryakova, L.A., Herzschuh, U., 2017. Sedimentary DNA versus morphology in the analysis of diatom-environment relationships. *J. Paleolimnol.* 57 (1), 51–66.
- Edgar, R.C., 2013. UPARSE: highly accurate OTU sequences from microbial amplicon reads. *Nat. Methods* 10 (10), 996–998.
- Edgar, R.C., 2017. Accuracy of microbial community diversity estimated by closed-and open-reference OTUs. *PeerJ* 5, e3889.
- European-Committee-for-Standardization, 2014. European Standard 13946. Water quality-Guidance for the routine sampling and preparation of benthic diatoms from rivers and lakes.
- Flower, R.J., 1993. Diatom preservation: experiments and observations on dissolution and breakage in modern and fossil material, Twelfth International Diatom Symposium. Springer, pp. 473–484.
- Frøstlev, T.G., Kjoller, R., Bruun, H.H., Ejrnæs, R., Brunbjerg, A.K., Pietroni, C., Hansen, A. J., 2017. Algorithm for post-clustering curation of DNA amplicon data yields reliable biodiversity estimates. *Nat. Commun.* 8 (1) <https://doi.org/10.1038/s41467-017-01312-x>.
- Geer, L.Y., Marchler-Bauer, A., Geer, R.C., Han, L., He, J., He, S., Liu, C., Shi, W., Bryant, S.H., 2009. The NCBI biosystems database. *Nucleic Acids Research* 38, D492–D496.
- Hardge, K., Neuhaus, S., Kilias, E.S., Wolf, C., Mettief, K., Frickenhaus, S., 2018. Impact of sequence processing and taxonomic classification approaches on eukaryotic community structure from environmental samples with emphasis on diatoms. *Mol. Ecol. Resour.* 18 (2), 204–216.
- Hermans, S.M., Buckley, H.L., Lear, G., 2018. Optimal extraction methods for the simultaneous analysis of DNA from diverse organisms and sample types. *Mol. Ecol. Resour.* 18 (3), 557–569.
- Kang, W., Chen, G., Wang, J., Huang, L., Wang, L., Li, R., Hu, K., Liu, Y., Tao, J., Blais, J. M., Smol, J.P., 2019. Assessing the impact of long-term changes in climate and atmospheric deposition on a shallow alpine lake from southeast Tibet. *Sci. Total Environ.* 650, 713–724.
- Keck, F., Vasselton, V., Rimet, F., Bouchez, A., Kahlert, M., 2018. Boosting DNA metabarcoding for biomonitoring with phylogenetic estimation of operational taxonomic units' ecological profiles. *Mol. Ecol. Resour.* 18 (6), 1299–1309.
- Keil, A., Berking, J., Mügler, I., Schütt, B., Schwalb, A., Steeb, P., 2010. Hydrological and geomorphological basin and catchment characteristics of Lake Nam Co, South-Central Tibet. *Quat. Int.* 218 (1–2), 118–130.
- Kelly, M., Boonham, N., Juggins, S., Killie, P., Mann, D., Pass, D., Sapp, M., Sato, S., Glover, R., 2018. A DNA based Diatom Metabarcoding Approach for Water Framework Directive Classification of Rivers.
- Kermerrec, L., Franc, A., Rimet, F., Chaumeil, P., Humbert, J.F., Bouchez, A., 2013. Next-generation sequencing to inventory taxonomic diversity in eukaryotic communities: a test for freshwater diatoms. *Mol. Ecol. Resour.* 13 (4), 607–619.
- Kocielek, J., You, Q., Wang, Q., Liu, Q., 2015. A consideration of some interesting freshwater gomphonemoid diatoms from North America and China, and the description of Gomphosinica gen. nov. *Nova Hedwigia* 144, 175–198.
- Krammer, K., Lange-Bertalot, H., 1986–1991. *Bacillariophyceae*, vol. 2 (1–4). Süßwasserflora von Mitteleuropa, Gustav Fischer Verlag, Stuttgart/Jena.
- Lange-Bertalot, H., Hofmann, G., Werum, M., Cantonati, M., Kelly, M.G., 2017. Freshwater benthic diatoms of Central Europe: over 800 common species used in ecological assessment. Koeltz Botanical Books Schmitten-Oberreifenberg.
- Li, G., Dong, H., Hou, W., Wang, S., Jiang, H., Yang, J., Wu, G., 2016. Temporal Succession of Ancient Phytoplankton Community in Qinghai Lake and Implication for Paleo-environmental Change. *Sci. Rep.* 6 (1) <https://doi.org/10.1038/srep19769>.
- Li, Y., Lange-Bertalot, H., Metzeltin, D., 2013. Sichuanella (Li Yanling, Lange-Bertalot et Metzeltin) nom. nov. - a new name for Sichuanella (Li Yanling et al). *Diatoms of Europe: Diatoms of the European inland waters and comparable habitats*. Vol 7, 5–12.
- Liaw, A., Wiener, M., 2002. Classification and regression by randomForest. *R news* 2, 18–22.
- Majaneva, M., Hyytiäinen, K., Varvio, S.L., Nagai, S., Blomster, J., 2015. Bioinformatic amplicon read processing strategies strongly affect eukaryotic diversity and the taxonomic composition of communities. *PLoS ONE* 10, e0130035.
- Liu, J., Rühland, K.M., Chen, J., Xu, Y., Chen, S., Chen, Q., Huang, W., Xu, Q., Chen, F., Smol, J.P., 2017. Aerosol-weakened summer monsoons decrease lake fertilization on the Chinese Loess Plateau. *Nature Clim. Change* 7 (3), 190–194.
- Mann, D.G., 1999. The species concept in diatoms. *Phycologia* 38 (6), 437–495.
- Medlin, L., Yang, I., Sato, S., 2012. Evolution of the diatoms. VII. Four gene phylogeny assesses the validity of selected araphid genera. *Nova Hedwigia* 141, 505–513.
- Mohan, J., Stone, J.R., Nicholson, K., Neumann, K., Dowling, C., SHARMA, S., 2018. *Lindavia biswashanti*, a new diatom species (Bacillariophyta) from Gokyo Cho, Himalayan Range, Nepal. *Phytotaxa* 364, 101–107.
- Morales, E.A., Wetzel, C.E., Novais, M.H., Buczkó, K., Morais, M.M., Ector, L., 2019. Morphological reconsideration of the araphid genus *Pseudostaurisira* (Bacillariophyceae), a revision of *Gedaniella*, *Popovskayella* and *Serratifera*, and a description of a new Nanofrustulum species. *Pl. Ecol. Evol.* 152 (2), 262–284.
- Mortágua, A., Vasselton, V., Oliveira, R., Elias, C., Chardon, C., Bouchez, A., Rimet, F., João Feio, M., Almeida, S.F.P., 2019. Applicability of DNA metabarcoding approach in the bioassessment of Portuguese rivers using diatoms. *Ecol. Ind.* 106, 105470. <https://doi.org/10.1016/j.ecolind.2019.105470>.
- Nascimento, F.J.A., Lallias, D., Bik, H.M., Creer, S., 2018. Sample size effects on the assessment of eukaryotic diversity and community structure in aquatic sediments using high-throughput sequencing. *Sci. Rep.* 8 (1) <https://doi.org/10.1038/s41598-018-30179-1>.
- Oksanen, J., Blanchet, F.G., Kindt, R., Legendre, P., Minchin, P.R., O'hara, R., Simpson, G.L., Solymos, P., Stevens, M., Wagner, H., 2015. *vegan: Community Ecology Package*. R package version 2.4-2. <http://CRAN.R-project.org/package=vegan>.
- Pauvert, Charlie, Bué, Marc, Laval, Valérie, Edel-Hermann, Véronique, Fauchery, Laure, Gautier, Angélique, Lesur, Isabelle, Vallance, Jessica, Vacher, Corinne, 2019. Bioinformatics matters: the accuracy of plant and soil fungal community data is highly dependent on the metabarcoding pipeline. *Fungal Ecology* 41, 23–33.
- Pavlov, A., Levkov, Z., Williams, D.M., Edlund, M.B., 2013. Observations on *Hippodonta* (Bacillariophyceae) in selected ancient lakes. *Phytotaxa* 90, 1–53.
- Penton, C.R., Gupta, V.V., Yu, J., Tiedje, J.M., 2016. Size matters: assessing optimum soil sample size for fungal and bacterial community structure analyses using high throughput sequencing of rRNA gene amplicons. *Front. Microbiol.* 7, 824.
- Piredda, Roberta, Sarno, Diana, Lange, Carina B., Tomasino, Maria Paola, Zingone, Adriana, Montresor, Marina, 2017. Diatom resting stages in surface sediments: a pilot study comparing next generation sequencing and serial dilution cultures. *Cryptogamie, Algologie* 38 (1), 31–46.
- R-Core-Team, 2015. R Foundation for Statistical Computing; 2015. R: A language and environment for statistical computing.
- Rimet, Frederic, Abarca, Nelida, Bouchez, Agnes, Kusber, Wolf-Henning, Jahn, Regine, Kahlert, Maria, Keck, Francois, Kelly, Martyn G., Mann, David, G., Piuze, Andre, Trobajo, Rosa, Tapolczai, Kalman, Vasselton, Valentin, Zimmermann, Jonas, 2018a. The potential of High-Throughput Sequencing (HTS) of natural samples as a source of primary taxonomic information for reference libraries of diatom barcodes. *Fottea* 18 (1), 37–54.
- Rimet, F., Chaumeil, P., Keck, F., Kermerrec, L., Vasselton, V., Kahlert, M., Franc, A., Bouchez, A., 2016. R-Syst: diatom: an open-access and curated barcode database for diatoms and freshwater monitoring. *Database* 2016.
- Rimet, Frédéric, Vasselton, Valentin, A.-Keszte, Barbara, Bouchez, Agnès, 2018b. Do we similarly assess diversity with microscopy and high-throughput sequencing? Case of microalgae in lakes. *Org. Divers. Evol.* 18 (1), 51–62.
- Rivera, Sinziana F., Vasselton, Valentin, Bouchez, Agnès, Rimet, Frédéric, 2020. Diatom metabarcoding applied to large scale monitoring networks: optimization of bioinformatics strategies using Mothur software. *Ecol. Ind.* 109, 105775. <https://doi.org/10.1016/j.ecolind.2019.105775>.
- Rivera, S.F., Vasselton, V., Jacquet, S., Bouchez, A., Ariztegui, D., Rimet, F., 2018. Metabarcoding of lake benthic diatoms: from structure assemblages to ecological assessment. *Hydrobiologia* 807 (1), 37–51.
- Rognes, T., Flouri, T., Nichols, B., Quince, C., Mahé, F., 2016. VSEARCH: a versatile open source tool for metagenomics. *PeerJ* 4, e2584.
- Ruppert, Krista M., Kline, Richard J., Rahman, Md Saydur, 2019. Past, present, and future perspectives of environmental DNA (eDNA) metabarcoding: a systematic review in methods, monitoring, and applications of global eDNA. *Global Ecol. Conserv.* 17, e00547. <https://doi.org/10.1016/j.gecco.2019.e00547>.
- Schiebelhut, Lauren M., Abboud, Sarah S., Gómez Daglio, Liza E., Swift, Holly F., Dawson, Michael N., 2017. A comparison of DNA extraction methods for high-throughput DNA analyses. *Mol. Ecol. Resour.* 17 (4), 721–729.
- Schloss, Patrick D., Westcott, Sarah L., Ryabin, Thomas, Hall, Justine R., Hartmann, Martin, Hollister, Emily B., Lesniewski, Ryan A., Oakley, Brian B., Parks, Donovan H., Robinson, Courtney J., Sahl, Jason W., Stres, Blaz, Thallinger, Gerhard G., Van Horn, David J., Weber, Carolyn F., 2009. Introducing mothur: open-source, platform-independent, community-supported software for describing and comparing microbial communities. *AEM* 75 (23), 7537–7541.
- Sinha, Rashmi, Abu-Ali, Galeb, Vogtmann, Emily, Fodor, Anthony A., Ren, Boyu, Amir, Ammon, Schwager, Emma, Crabtree, Jonathan, Ma, Siyuan, Abnet, Christian C., Knight, Rob, White, Owen, Huttenhower, Curtis, 2017. Assessment of variation in microbial community amplicon sequencing by the Microbiome Quality Control (MBQC) project consortium. *Nat. Biotechnol.* 35 (11), 1077–1086.
- StatSoft, I., 2004. STATISTICA (data analysis software system), version 7. [www.statsoft.com](http://www.statsoft.com).
- Stoof-Leichsenring, Kathleen R., Bernhardt, Nadine, Pestryakova, Luidmila A., Epp, Laura S., Herzschuh, Ulrike, Tiedemann, Ralph, 2014. A combined paleolimnological/genetic analysis of diatoms reveals divergent evolutionary lineages of *Staurisira* and *Staurisirella* (Bacillariophyta) in Siberian lake sediments along a latitudinal transect. *J. Paleolimnol.* 52 (1–2), 77–93.
- Taberlet, P., Bonin, A., Coissac, E., Zinger, L., 2018. *Environmental DNA: For biodiversity research and monitoring*. Oxford University Press.
- Tapolczai, K., Keck, F., Bouchez, A., Rimet, F., Vasselton, V., 2019a. Diatom DNA metabarcoding for biomonitoring: strategies to avoid major taxonomical and bioinformatical biases limiting molecular indices capacities. *Frontiers in Ecology and Evolution* 7, 409.



- Tapolczai, Kálmán, Vasselon, Valentin, Bouchez, Agnès, Stenger-Kovács, Csilla, Padišák, Judit, Rimet, Frédéric, 2019b. The impact of OTU sequence similarity threshold on diatom-based bioassessment: a case study of the rivers of Mayotte (France, Indian Ocean). *Ecol. Evol.* 9 (1), 166–179.
- Dam, Herman, Mertens, Adrienne, Sinkeldam, Jos, 1994. A coded checklist and ecological indicator values of freshwater diatoms from The Netherlands. *Netherlands J. Aquatic Ecol.* 28 (1), 117–133.
- Vasselon, Valentin, Bouchez, Agnès, Rimet, Frédéric, Jacquet, Stéphan, Trobajo, Rosa, Corniquel, Méline, Tapolczai, Kálmán, Domaizon, Isabelle, Mahon, Andrew, 2018. Avoiding quantification bias in metabarcoding: application of a cell biovolume correction factor in diatom molecular biomonitoring. *Methods Ecol. Evol.* 9 (4), 1060–1069.
- Vasselon, Valentin, Domaizon, Isabelle, Rimet, Frédéric, Kahlert, Maria, Bouchez, Agnès, 2017a. Application of high-throughput sequencing (HTS) metabarcoding to diatom biomonitoring: Do DNA extraction methods matter? *Freshwater Sci.* 36 (1), 162–177.
- Vasselon, Valentin, Rimet, Frédéric, Tapolczai, Kálmán, Bouchez, Agnès, 2017b. Assessing ecological status with diatoms DNA metabarcoding: scaling-up on a WFD monitoring network (Mayotte island, France). *Ecol. Ind.* 82, 1–12.
- Visco, Joana Amorim, Apothéloz-Perret-Gentil, Laure, Cordonier, Arielle, Esling, Philippe, Pillet, Loïc, Pawlowski, Jan, 2015. Environmental monitoring: inferring the diatom index from next-generation sequencing data. *Environ. Sci. Technol.* 49 (13), 7597–7605.
- Wang, J., Zhu, L., Wang, Y., Ju, J., Xie, M., Daut, G., 2010. Comparisons between the chemical compositions of lake water, inflowing river water, and the lake sediment in Nam Co, central Tibetan Plateau, China and their controlling mechanisms. *J. Great Lakes Res.* 36, 587–595.
- Wang, Junbo, Huang, Lei, Ju, Jianting, Daut, Gerhard, Ma, Qingfeng, Zhu, Liping, Haberzettl, Torsten, Baade, Jussi, Mäusbacher, Roland, Hamilton, Andrew, Graves, Kelly, Olsthoorn, Jason, Laval, Bernard E., 2020. Seasonal stratification of a deep, high-altitude, dimictic lake: Nam Co, Tibetan Plateau. *J. Hydrol.* 584, 124668. <https://doi.org/10.1016/j.jhydrol.2020.124668>.
- Wang, Rong, Dearing, John A., Langdon, Peter G., Zhang, Enlou, Yang, Xiangdong, Dakos, Vasilis, Scheffer, Marten, 2012. Flickering gives early warning signals of a critical transition to a eutrophic lake state. *Nature* 492 (7429), 419–422.
- Zhang, Guoqing, Yao, Tandong, Xie, Hongjie, Kang, Shichang, Lei, Yanbin, 2013. Increased mass over the Tibetan Plateau: From lakes or glaciers? *Geophys. Res. Lett.* 40 (10), 2125–2130.
- Zimmermann, Jonas, Glöckner, Gernot, Jahn, Regine, Enke, Neela, Gemeinholzer, Birgit, 2015. Metabarcoding vs. morphological identification to assess diatom diversity in environmental studies. *Mol. Ecol. Resour.* 15 (3), 526–542.
- Wei, T., Simko, V., Levy, M., Xie, Y., Jin, Y., Zelma, J., 2017. Package ‘corrplot’. *Statisticians* 56, e24.