MULTICHANNEL ENERGY ANALYSIS OF THE ION FLUX⁺) FROM A THETA PINCH

by

G. Becker
INSTITUT FÜR PLASMAPHYSIK
Garching bei München, Germany

Conventional diagnostic methods for ion energies in hot theta pinch plasmas use the neutron flux or the laser light scattered in the forward direction. None of these gives the detailed energy distribution of the ions. Determination of the spectrum from the axial end losses has already been reported by [1].

The new 10-channel energy spectrometer used here measures the time resolved energy distribution in the flux $\parallel$ to the axis from each discharge ($E = 1 - 10$ keV). Preliminary diagnostic investigations and the measurements were made on the Isar III theta pinch [2] with the standard parameters 10 mTorr D_2 , $B_{\text{max}} = 75$ kG, and 30 cm long straight coil. The experimental set-up is shown in Fig. 1.

The particle beam of small divergence passes through a metal collimator and is energy dispersed in the field of a planar capacitor. Behind each of the 10 exit slits there is a fast detector system. The energy resolution is found to be dependent only on the slit assembly and not on space charges in the capacitor.

Fig. 1

Figure 2 shows the deuteron flux for 1 - 7 keV from one discharge (channel width 200 eV). In addition, the neutron flux and the pinch field are displayed as a time reference.

Fig. 2

Fig. 3

The distance A (Fig. 1) was varied originally to investigate the influence of the coil field on the distribution. This, however, proved to be unimportant. Figure 3 shows the maximum ion currents versus A. Although the solid angle Ω_0 , in which particles can be detected, decreases with growing A, the measured flux increases. This is inconsistent with the existing conception of direct measurement of deuterons as well as the fact that the decrease for large A does not go faster than $\propto \Omega_0$. Behind the long path through neutral gas the measured flux can only exist if equilibrium between D atoms (90 %) and deuterons (10 %) builds up as a result of charge exchange cycles in D_2 . The deuteron flux diverges at the limit of the coil field (B_{lim}) in the aperture, and so the capacitor is bypassed. Additional experiments confirmed that the measured flux increases with the neutral gas length in front of B_{lim} . Deuterons can only be detected, if they pass B_{lim} as D atoms without being deflected, and if they are reionized by one more charge exchange in the field-free region. Changes of energy and direction due to charge exchange cycles can be neglected here. $I = I(A, E)$ in Fig. 3 can be calculated from the change of deuteron flux at the coil end ($I_0(E)$) due to charge exchange and electron-impact ionization. The reduction of I_0 depends only slightly on E in the range 4 - 10 keV, but is much greater for low E (1 keV).

In Fig. 4 measured flux spectra $\frac{dn}{dE}(E)$ are plotted at several times of one discharge. They are similar to the averaged results of [1]. The dip in $\frac{dn}{dE}$ at low energies, interpreted as a lack of slow ions in [1], is now explained in terms of the above processes.

For $E \gg 3$ keV $\frac{dn}{dE} \propto \exp \left\{ -\frac{E}{E} \right\}$. $\bar{E}$ is compared with kT from the neutron flux in the case of a 3-dim. Maxwellian.

10 mTorr $\bar{E} \approx 1-1.3$ keV	$kT \approx 1$ keV
20 mTorr $\bar{E} \approx 900$ eV	$kT \approx 0.9-1$ keV
30 mTorr $\bar{E} \approx 600$ eV	$kT \approx 500-600$ eV

The first pulse 300 nsec after the beginning of the main discharge ($t = 0$) is ascribed to the separation of the divergent plasma ends (Fig. 2). The onset of the main signal at $t = t_1 \approx 1$ μsec is unexpectedly steep. Mach-Zehnder measurements of the end losses into 2π show that about 30 % of the plasma has already been lost up to $t = t_1$, when $\frac{v_{\perp}}{v_{\parallel}}$ is large (anisotropic outflow, $l_p/v_{\parallel} < t_c$ ion relaxation time; l_p plasma length). The starting point and t_1 can be determined independently by means of the 10-channel energy spectrometer. l_p is found to equal the coil length fairly well. t_1 shows an n and m-dependence contrary to that of t_c given by the theory. The sudden onset of losses into Ω_0 observed in all channels at $t = t_1$ suggests anomalously fast relaxation, which may be caused by mirror instability. Relaxation must take place within a few 100 nsec because the outflow is isotropic for $t > t_1$. This follows from a comparison of the total particle flux into 2π with the end loss into Ω_0 , calculated from the absolute E-integrated flux spectrum. At $t = 1.6$ μsec one measures, for example:

Total end loss into 2π	: ≈ 2.5 kA
E-integrated loss into $\Omega_0 = 2 \cdot 10^{-6}$ rad ²	: ≈ 1 mA
Loss into Ω_0 for $E = 7 \pm 0.1$ keV	: max. ≈ 2 μA.

The extremely high spectrometer detection efficiency allows analysis of late end losses despite the fact that Ω_0 is only $2 \cdot 10^{-6}$ rad² and the channel width only 200 eV. The end loss into 2π corresponds to about 1 kA at the detection limit of interferometry for $l_p \approx 30$ cm. Ion analysis is possible, however, down to $I_0 \approx 10^{-7}$ A.

After the instability at t_1 relatively few energetic ions are seen in Ω_0 . Thus, there is no fast relaxation for these ions. A late anomaly (possibly a loss-cone instability) affects particularly $E \geq 7$ keV ions (2nd main signal), but is also to be seen in the decay of the signals of low energy ions that have already become rare at this time. The increase of the fraction of energetic ions in the flux with time corresponds to a longer confinement of these ions. The discussed results with straight coil are obtained with a mirror ratio $R = 1.6$ as well.

[1] R.L. Bingham, L.M. Goldman, R.W. Kilb; Plasma Phys. and Contr. Nucl. Fus. Res., I.A.E.A., I, 301 (1966)

[2] A. Heiss, H. Herold, E. Unsöld, Proc. APS Top. Conf., Los Alamos (1967), LA-3770, p.D7-1.

⁺) This work was performed as part of the joint research program of the Institut für Plasmaphysik, Garching, and Euratom.