

CENTER OF MASS AND KÄHLER STRUCTURES

SCOTT O. WILSON AND MAHMOUD ZEINALIAN

ABSTRACT. There is a sequence of positive numbers δ_{2n} , such that for any connected $2n$ -dimensional Riemannian manifold M , there are two mutually exclusive possibilities:

- (1) There is a complex structure on M making it into a Kähler manifold.
- (2) For any almost complex structure J compatible with the metric, at every point $p \in M$, there is a smooth loop γ at p such that

$$\text{dist}(J_p, \text{hol}_\gamma^{-1} J_p \text{hol}_\gamma) > \delta_{2n}.$$

1. INTRODUCTION

Kähler manifolds possess a tremendous amount of interesting structure, and therefore have several equivalent characterizations. It has been a focus of much research to determine conditions under which manifolds do (or do not) admit a Kähler structure. This short note shows that the holonomy action on the space of almost hermitian structures determines two mutually exclusive cases, according to whether there is a structure that is *nearly preserved* at some point, by proving that any manifold with a nearly preserved almost hermitian structure at some point in fact admits a Kähler structure. The novel idea is to use a center of mass argument, averaging a given almost complex structure at a point over the holonomy action, and parallel transporting the result to obtain a global Kähler structure.

Due to the plethora of topological implications for having a Kähler structure, one may deduce several interesting consequences that do not mention the Kähler condition at all. For example, we deduce by [DGMS] that any almost hermitian manifold which has a non-trivial Massey product (or more generally is not formal) also has a holonomy action on the space almost hermitian structures which is bounded away from the identity in a way that is precisely quantifiable.

The authors would like to thank Dennis Sullivan for conversations about this.

2010 *Mathematics Subject Classification.* 53C55, 32Q15, 53C29.

Key words and phrases. Kähler, almost hermitian, holonomy, center of mass.

2. MAIN RESULT

For a $2n$ -dimensional real vector space V with an inner product g , let

$$\mathbb{J}(V, g) = \{J : V \rightarrow V \mid J^2 = -id, g(Ju, Jv) = g(u, v)\}$$

denote the space of metric almost complex structures on V .

Lemma 1. $\mathbb{J}(V, g)$ is a compact smooth manifold. The tangent space at a point J is

$$T_J \mathbb{J}(V, g) = \{\phi : V \rightarrow V \mid \phi J = -J\phi \text{ and } \phi^\dagger = -\phi\}$$

and has a bilinear form $\tilde{g}(\phi, \psi) = \text{tr}(\phi\psi^\dagger)$, for all $\phi, \psi \in T_J \mathbb{J}(V, g)$, where ψ^\dagger denotes the adjoint of ψ . This makes $\mathbb{J}(V, g)$ into a Riemannian manifold on which $O(V)$ acts transitively by isometries.

Proof. $\mathbb{J}(V, g)$ can be identified as the $O(V)$ -homogenous space $O(V)/U(V)$ where $U(V)$ is defined using any fixed J on V , and the action of $O(V)$ on $\mathbb{J}(V, g)$ is given by conjugation. Therefore $\mathbb{J}(V, g)$ is a smooth manifold. The induced action of $O(V)$ on $T_J \mathbb{J}(V, g)$ is also given by conjugation, so that trace is invariant. \square

Lemma 2. For any two metric vector spaces (V, g) and (W, h) of the same dimension, the Riemannian manifolds $\mathbb{J}(V, g)$ and $\mathbb{J}(W, h)$ are isometric. In particular, for a vector space V and two metrics g and g' , the Riemannian manifolds $\mathbb{J}(V, g)$ and $\mathbb{J}(V, g')$ are isometric.

Proof. The Gramm-Schmidt process ensures that there is a linear isometry $f : (V, g) \rightarrow (W, h)$. Conjugation by this isometry gives the desired Riemannian isometry. \square

For an argument below, we will require convex balls in $\mathbb{J}(V, g)$ for which there is a well defined notion of center of mass. For any Riemannian manifold M , balls of radius r are convex if

$$r = \min \left\{ \frac{\text{inj } M}{2}, \frac{\pi}{2\sqrt{\epsilon}} \right\},$$

where $\text{inj } M$ denotes the injectivity radius of M , and ϵ is a finite positive upper bound on the sectional curvature (c.f. [Ch] Prop IX.6.1). Also, for any such ϵ , we have the following theorem.

Theorem 3 (Karcher, [K]). Let $f : X \rightarrow M$ be a measurable map from a probability space (X, m) to a Riemannian manifold M . If $f(X)$ is contained in a convex subset B of M , with diameter less than or equal to $\pi/2\sqrt{\epsilon}$, then there is a unique center of mass in B , defined by the minimum of

$$E(y) = \frac{1}{2} \int_X d^2(f(x), y) m(x).$$

Additionally, the center of mass is natural with respect to isometries ([K], 1.4.1). We refer the reader to [Ch] Prop IX.7.1 for an exposition of center of mass.

Definition 4. For $n > 1$ let

$$\delta_{2n} = \min\left\{\frac{\text{inj } \mathbb{J}(V, g)}{2}, \frac{\pi}{4\sqrt{\epsilon}}\right\}$$

where V is a real vector space of dimension $2n$ with any metric g . Here we choose, once and for all, a finite positive upper bound ϵ on the sectional curvature of $\mathbb{J}(V, g)$ at one point, which by homogeneity works for all points. By Lemma 2, δ_{2n} depends only on the dimension of V .

An interesting question (that we will not address here) is whether there is a positive lower bound for the set of all least such δ_{2n} , independent of n .

Definition 5. A complex structure $J \in \mathbb{J}(V, g)$ is said to be nearly preserved by a subgroup $H \subset O(V)$ if the orbit $HJ = \{\phi^{-1}J\phi \mid \phi \in H\}$ lies inside the ball $B(J, \delta_{2n}) \subset \mathbb{J}(V, g)$ with center J and radius δ_{2n} .

Recall that any closed subgroup H of $O(V)$ is a compact Lie group, admitting a bi-invariant Haar measure, which is unique up to a constant. Therefore any such H has a unique probability measure (of total mass equal to one).

Proposition 6. Let $J \in \mathbb{J}(V, g)$ be nearly preserved by a closed subgroup H of $O(V)$. Then there is a $J' \in \mathbb{J}(V, g)$ such that $HJ' = \{J'\}$.

Proof. Consider the orbit $HJ \subset B(J, \delta_{2n}) \subset \mathbb{J}(V, g)$. By assumption, $B(J, \delta_{2n})$ is a convex ball about J . Consider the mapping $H \rightarrow HJ \subset \mathbb{J}(V, g)$, from the probability space H onto its orbit. By Definition 4 and Theorem 3, the set HJ has a unique center of mass J' in S . Since the orbit of the action of H on the set HJ is itself, and the center of mass is unique and natural with respect to isometries given by the H -action, H fixes J' . \square

Given a point $p \in M$, let $H_p = \text{Im}(\rho_p)$ be the image subgroup of the holonomy homomorphism $\rho_p : \Omega_p(M) \rightarrow O(T_pV)$, which is known to be closed by the Ambrose-Singer theorem. The following proposition is a standard result which we include for completeness.

Proposition 7. Let (M, g) be a connected Riemannian manifold and assume $J_p \in \mathbb{J}(T_pM, g)$ is invariant under the action of $H_p = \text{Im}(\rho_p)$. Then M admits a unique almost complex structure $J : TM \rightarrow TM$ agreeing with J_p and making (M, g, J) into a Kähler manifold.

Proof. For any $q \in M$ define $J_q : T_qM \rightarrow T_qM$ by $P_\lambda \circ J_p \circ P_\lambda^{-1}$, where $P_\lambda : T_pM \rightarrow T_qM$ is the Riemannian parallel transport along any smooth path λ in M from p to q . Since M is connected, such a path λ always exists and J_q is independent of the choice of the path because $H_p J_p = \{J_p\}$. Thus, we have defined a smooth complex structure $J : TM \rightarrow TM$, and by way of construction, J is compatible with all parallel transports, and therefore $\nabla J = 0$. Since the Levi-Cevita connection ∇ is torsion free, and J is still Hermitian, J is integrable; see Corollary 3.5 of [KN]. Therefore, (M, g, J) is a Kähler manifold. \square

Corollary 8. *With (M, g) as above, if there is a $p \in M$ and a $J_p \in \mathbb{J}(T_p M, g)$ that is nearly preserved by $H_p = \text{Im}(\rho_p)$, then there is a $J' : TM \rightarrow TM$ making (M, g, J') into a Kähler manifold.*

Proof. This follows from Proposition 6 and Proposition 7. □

Theorem 9. *There is a sequence of positive numbers $\delta_{2n} \in \mathbb{R}$ such that for any connected $2n$ -dimensional Riemannian manifold M , one of the two following mutually exclusive properties hold:*

- (1) *There is a complex structure on M making it into a Kähler manifold.*
- (2) *For any almost complex structure J compatible with the metric, at every point $p \in M$, there is a smooth loop γ at p such that*

$$\text{dist}(J_p, \text{hol}_\gamma^{-1} J_p \text{hol}_\gamma) > \delta_{2n}.$$

Proof. If 1) is true then 2) is clearly false, and the converse follows from the previous corollary. □

REFERENCES

- [Ch] I. Chavel, *Riemannian Geometry: A Modern Introduction*, (Cambridge Studies in Advanced Mathematics), 2nd edition.
- [DGMS] P. Deligne, P. Griffiths, J. Morgan, D. Sullivan, “Real homotopy theory of Kähler manifolds,” *Invent. Math.* 29 (1975), no. 3, 245-274.
- [K] Karcher, H. “Riemannian center of mass and mollifier smoothing,” *Comm. Pure Appl. Math.* 30, (1977), 509-541.
- [KN] S. Kobayashi, K. Nomizu, *Foundations of Differential Geometry*, Vol II, Wiley-Interscience (1996).

SCOTT O. WILSON, DEPARTMENT OF MATHEMATICS, QUEENS COLLEGE, CITY UNIVERSITY OF NEW YORK, 65-30 KISSENA BLVD., FLUSHING, NY 11367

E-mail address: `scott.wilson@qc.cuny.edu`

MAHMOUD ZEINALIAN, DEPARTMENT OF MATHEMATICS, LIU POST, LONG ISLAND UNIVERSITY, 720 NORTHERN BOULEVARD, BROOKVILLE, NY 11548, USA

E-mail address: `mzeinalian@liu.edu`