ON TAU-FUNCTIONS FOR THE KDV HIERARCHY

BORIS DUBROVIN[†], DI YANG, AND DON ZAGIER

ABSTRACT. For an *arbitrary* solution to the KdV hierarchy, the generating series of logarithmic derivatives of the tau-function of the solution can be expressed by the basic matrix resolvent via algebraic manipulations. Based on this we develop in this paper two new formulae for the generating series by introducing a pair of wave functions of the solution. Applications to the Witten–Kontsevich tau-function, to the generalized Brézin–Gross–Witten (BGW) tau-function, as well as to a modular deformation of the generalized BGW tau-function which we call the Lamé tau-function are also given.

Mathematics Subject Classification (2010). 37K10; 53D45, 14N35, 05A15, 33E15.

Keywords. KdV hierarchy, tau-function, pair of wave functions, matrix resolvent, generating series.

CONTENTS

1. Introduction	1
2. Review of the matrix resolvent approach to the KdV hierarchy	11
3. From a pair of wave functions to the n -point functions	14
4. Generating series of the generalized BGW correlators	19
5. Generating series of the Lamé partial correlation functions	24
6. Further remarks	26
References	29

1. INTRODUCTION

To make this article accessible also to non-specialists, we recall the definitions of all of the main objects studied (KdV hierarchy, tau-function, matrix resolvents, wave functions, $\dots$), in some cases with definitions slightly different from the most standard ones. Experts can simply skip this material.

1.1. The Korteweg–de Vries hierarchy. The Korteweg–de Vries (KdV) equation

$$u_t = u u_x + \frac{1}{12} u_{xxx}, \quad (1)$$

discovered in the study of water waves in the 19th century, extends to a *hierarchy* of PDEs,

$$\frac{\partial u}{\partial t_k} = \frac{u^k}{k!} u_x + Q_k(u, u_x, u_{xx}, \dots) \quad (k \geq 0), \quad (2)$$

where u is now a function of the multivariable $\mathbf{t} = (t_0 = x, t_1 = t, t_2, t_3, \dots)$ and the Q_k are specific polynomials with $Q_0 = 0$ and $Q_1 = u_{xxx}/12$. We recall a simple construction of this hierarchy. Denote by $\mathcal{A}$ the polynomial algebra $\mathbb{Q}[u_0, u_1, u_2, \dots]$ with the grading $\deg u_i = i$, and by ∂ the derivation on $\mathcal{A}$ that satisfies $\partial(u_i) = u_{i+1}$. For arbitrary elements $Q_0, Q_1, Q_2, \dots$ in $\mathcal{A}$ we can define uniquely a family of derivations D_k ($k \geq 0$) on $\mathcal{A}$ by

$$D_0 = \partial, \quad [D_k, D_0] = 0, \quad D_k(u_0) = \frac{u_0^k}{k!} u_1 + Q_k. \quad (3)$$

[†]Deceased on March 19, 2019.

It turns out that if we require that $Q_k \in \mathcal{A}^{\geq 2}$ with $Q_0 = 0$ and $Q_1 = u_3/12$ and that all D_k commute with D_1 , then the polynomials Q_k exist and are unique. The first few values are

$$\begin{aligned} Q_0 &= 0, & Q_1 &= \frac{1}{12}u_3, & Q_2 &= \frac{1}{12}(2u_1u_2 + u_0u_3) + \frac{1}{240}u_5, \\ Q_3 &= \frac{1}{24}(u_1^3 + 4u_0u_1u_2 + u_0^2u_3) + \frac{1}{240}(5u_2u_3 + 3u_1u_4 + u_0u_5) + \frac{1}{6720}u_7, \end{aligned}$$

and all the D_k commute. (See [28, 58]. A new proof of the existence and pairwise commutativity is given in Section 2.) We call these unique derivations D_k the KdV derivations. The formal system (3) called the abstract KdV hierarchy leads to the compatible system of PDEs (2) if $u = u(\mathbf{t})$ and we identify u_i and D_k with $\partial_x^i(u)$ and $\partial/\partial t_k$, respectively.

Let V be a ring of functions of x closed under ∂_x . (Usually V will be $\mathbb{C}[[x]]$ or $\mathbb{C}((x))$.) For given initial data $f(x) \in V$, one can solve the KdV hierarchy (2) to get a unique solution $u = u(\mathbf{t})$ in $V[[\mathbf{t}_{>0}]]$ with $u(x, 0, 0, \dots) = f(x)$. This gives a 1-1 correspondence:

$$\{\text{solution } u \text{ of (2) in } V[[\mathbf{t}_{>0}]]\} \longleftrightarrow \{\text{initial data } f\} = V. \quad (4)$$

Below we give three cases of particular interest. More examples can be found in [13, 55].

Example 1. $f(x) = x$. The corresponding unique solution u in $\mathbb{C}[[x, \mathbf{t}_{>0}]]$ governs [57, 66] the integrals

$$\int_{\overline{\mathcal{M}}_{g,n}} \psi_1^{p_1} \cdots \psi_n^{p_n}, \quad (5)$$

where $\overline{\mathcal{M}}_{g,n}$ denotes the Deligne–Mumford moduli space of stable algebraic curves of genus g with n distinct marked points and ψ_j denotes the so-called ψ -class (see Section 4 for the precise definition). This solution is often called the *Witten–Kontsevich solution*, denoted by u_{WK} .

Example 2. $f(x) = \frac{C}{(x-1)^2}$, $C \in \mathbb{C}$. The corresponding solution u in $\mathbb{C}[[x, \mathbf{t}_{>0}]]$ will be called the *generalized BGW solution* [2, 61], denoted by $u_{\Theta(C)}$. For $C = 1/8$, $u_{\Theta(C)}$ governs [62] the integrals

$$\int_{\overline{\mathcal{M}}_{g,n}} \Theta_{g,n} \psi_1^{p_1} \cdots \psi_n^{p_n}, \quad (6)$$

called the n -point Θ -class intersection numbers. Here $\Theta_{g,n}$ denotes the Norbury Θ -class [62] whose precise definition will be recalled in Section 4. We call $u_{\Theta(1/8)}$ the *BGW–Norbury solution*.

Example 3. $f(x) = C \wp(x; \tau)$, $C \in \mathbb{C}$, $\tau \in \mathfrak{H} = \text{upper half plane}$. Here we can take V to be the ring $\mathbb{C}[g_2, g_3, \wp, \wp'] / (\wp'^2 - 4\wp^3 + g_2\wp + g_3)$ with $g_2 = 60G_4$ and $g_3 = 140G_6$, where $G_{2k} = \sum_{(m,n) \in \mathbb{Z}^2 \setminus (0,0)} \frac{1}{(m+n\tau)^{2k}}$, $k \geq 2$. The corresponding unique solution u in $V[[\mathbf{t}_{>0}]]$ is a modular deformation of $u_{\Theta(C)}$, denoted by u_{elliptic} and discussed in more detail in Section 5. We call u_{elliptic} a *Lamé solution* of the KdV hierarchy.

1.2. The tau-function and the matrix resolvent approach. Let $(D_k)_{k \geq 0}$ be the KdV derivations defined above. By a *tau-structure* for the abstract KdV hierarchy we mean a collection of elements $\Omega_{p,q} \in \mathcal{A}$ ($p, q \geq 0$) satisfying

$$\Omega_{0,0} = u_0, \quad \Omega_{p,q} = \Omega_{q,p}, \quad D_r(\Omega_{p,q}) = D_q(\Omega_{p,r}), \quad \forall p, q, r \geq 0. \quad (7)$$

(A more general abstract tau-structure allows $\Omega_{0,0}$ to be some other element of $\mathcal{A}$ with non-degenerate 0th-order term, but this will not be studied in this paper.) Since the D_k commute, the elements

$$\Omega_{p_1, \dots, p_n} := D_{p_1} \cdots D_{p_{n-2}}(\Omega_{p_{n-1}, p_n}) \quad (n \geq 3, p_1, \dots, p_n \geq 0) \quad (8)$$

are symmetric in their indices. One can show that the tau-structure exists and is unique up to replacing $\Omega_{p,q}$ by $\Omega_{p,q} + c_{p,q}$, where $c_{p,q} = c_{q,p}$ are constants. By the *canonical* tau-structure we will mean the unique choice for which all $\Omega_{p,q}$ vanish when all $u_i = 0$, the first few values being

$$\Omega_{0,1} = \frac{u_0^2}{2} + \frac{u_2}{12}, \quad \Omega_{0,2} = \frac{u_0^3}{6} + \frac{u_1^2}{24} + \frac{u_0u_2}{12} + \frac{u_4}{240}, \quad \Omega_{1,1} = \frac{u_0^3}{3} + \frac{u_1^2}{24} + \frac{u_0u_2}{6} + \frac{u_4}{144}, \quad \dots$$

If $u = u(\mathbf{t})$ is a solution of (2) and we write $\Omega_{p,q}(\mathbf{t})$ as the image of $\Omega_{p,q}$ under $u_i \mapsto \partial_x^i(u)$, then (7) implies that there exists a function $\tau = \tau(\mathbf{t})$, a so-called *tau-function* of the solution u , such that

$$\Omega_{p,q}(\mathbf{t}) = \frac{\partial^2 \log \tau(\mathbf{t})}{\partial t_p \partial t_q} \quad (p, q \geq 0). \quad (9)$$

The symmetry in (8) is then obvious, since the image $\Omega_{p_1, \dots, p_n}(\mathbf{t})$ of $\Omega_{p_1, \dots, p_n}$ under $u_i \mapsto \partial_x^i(u)$ is

$$\Omega_{p_1, \dots, p_n}(\mathbf{t}) = \frac{\partial^n \log \tau(\mathbf{t})}{\partial t_{p_1} \cdots \partial t_{p_n}} \quad (n \geq 2, p_1, \dots, p_n \geq 0). \quad (10)$$

Denote $\Omega_p(\mathbf{t}) := \partial_{t_p}(\log \tau(\mathbf{t}))$. These logarithmic derivatives are called *n-point correlation functions* of u and the specializations $\Omega_{p_1, \dots, p_n}(x) := \Omega_{p_1, \dots, p_n}(x, \mathbf{0})$ are called *n-point partial correlation functions* of u , while the evaluation of $\Omega_{p_1, \dots, p_n}(x)$ at a particular value of x will be called an *n-point correlator* of u . If we choose the canonical tau-structure as defined above, we will call τ the tau-function of u , although it is still defined up to multiplying the exponential of an arbitrary linear function of $\mathbf{t}$. This definition of the tau-function τ is known to agree with other literature ([26, 42]).

The key fact for this paper is that the canonical tau-structure for the abstract KdV hierarchy can be given by an explicit generating series using the so-called matrix resolvent (MR) approach [13, 15]. This goes as follows. Let $\mathcal{L}$ (“matrix Lax operator”) be the operator $\partial + \Lambda(\lambda) + q$, where $\Lambda(\lambda) = \begin{pmatrix} 0 & 1 \\ \lambda & 0 \end{pmatrix}$, $q = \begin{pmatrix} 0 & 0 \\ -2u_0 & 0 \end{pmatrix}$. Let $\mathcal{S} = \text{sl}_2(\mathbb{C})((\lambda^{-1}))$ be the space of $\text{sl}_2(\mathbb{C})$ -valued formal Laurent series in λ^{-1} . The *principal grading* on $\mathcal{A} \otimes \mathcal{S}$ is defined by $\deg E = 1$, $\deg F = -1$, $\deg \lambda = 2$, $\deg u_i = 0$, where $E = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$, $F = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$ are the Weyl generators. The **basic matrix resolvent** of $\mathcal{L}$ is defined as the unique element $R(\lambda) = R(\lambda; u_0, u_1, \dots) \in \mathcal{S} \otimes \mathcal{A}$ satisfying

$$[\mathcal{L}, R(\lambda)] = 0, \quad R(\lambda) = \Lambda(\lambda) + \text{l.o.t.}, \quad \text{Tr } R(\lambda)^2 = 2\lambda, \quad (11)$$

where “l.o.t.” means lower order terms with respect to the principal gradation. The proof of existence and uniqueness of $R(\lambda)$ can be found in [15] or in Section 2 of the present paper. The upper right entry $R(\lambda)_{12}$ of the basic matrix resolvent which we denote by $b(\lambda)$ plays an important role. The following proposition, originally proved in [13], will be re-proved in Section 2 in a more straightforward way.

Proposition 1 ([13, 15, 71]). (i) The differential polynomials $\Omega_{p,q} \in \mathcal{A}$ defined by the generating series

$$\frac{\text{tr}(R(\lambda)R(\mu))}{(\lambda - \mu)^2} - \frac{\lambda + \mu}{(\lambda - \mu)^2} = \sum_{p,q \geq 0} \frac{(2p+1)!!(2q+1)!!}{\lambda^{p+1}\mu^{q+1}} \Omega_{p,q} \quad (12)$$

form a tau-structure for the abstract KdV hierarchy, vanishing at $u_0 = u_1 = \dots = 0$.

(ii) For any integer $n \geq 3$ and the corresponding $\Omega_{p_1, \dots, p_n}$ as defined in (8) we have

$$\sum_{p_1, \dots, p_n \geq 0} \Omega_{p_1, \dots, p_n} \prod_{j=1}^n \frac{(2p_j+1)!!}{\lambda_j^{p_j+1}} = - \sum_{\sigma \in S_n/C_n} \frac{\text{tr}(R(\lambda_{\sigma(1)}) \cdots R(\lambda_{\sigma(n)}))}{\prod_{i=1}^n (\lambda_{\sigma(i+1)} - \lambda_{\sigma(i)})}, \quad (13)$$

where S_n denotes the symmetry group, C_n the cyclic group, and it is understood that $\sigma(n+1) = \sigma(1)$.

Just as with $\Omega_{p,q}$ and $\Omega_{p,q}(\mathbf{t})$, we can evaluate the abstract basic matrix resolvent $R(\lambda)$ for a solution $u(\mathbf{t})$ of the KdV hierarchy by substituting $\partial_x^i(u)$ for u_i . The resulting matrix-valued function $R(\lambda, \mathbf{t})$ is called the *basic matrix resolvent* of u . Denote for short $R(\lambda, x) = R(\lambda, x, \mathbf{0})$.

Corollary 1 ([13]). For any $n \geq 2$, the following formulas hold true:

$$\sum_{p_1, \dots, p_n} \Omega_{p_1, \dots, p_n}(\mathbf{t}) \prod_{j=1}^n \frac{(2p_j+1)!!}{\lambda_j^{p_j+1}} = - \sum_{\sigma \in S_n/C_n} \frac{\text{tr } R(\lambda_{\sigma(1)}, \mathbf{t}) \cdots R(\lambda_{\sigma(n)}, \mathbf{t})}{\prod_{i=1}^n (\lambda_{\sigma(i+1)} - \lambda_{\sigma(i)})} - \frac{\lambda_1 + \lambda_2}{(\lambda_1 - \lambda_2)^2} \delta_{n2}, \quad (14)$$

$$\sum_{p_1, \dots, p_n} \Omega_{p_1, \dots, p_n}(x) \prod_{j=1}^n \frac{(2p_j+1)!!}{\lambda_j^{p_j+1}} = - \sum_{\sigma \in S_n/C_n} \frac{\text{tr } R(\lambda_{\sigma(1)}, x) \cdots R(\lambda_{\sigma(n)}, x)}{\prod_{i=1}^n (\lambda_{\sigma(i+1)} - \lambda_{\sigma(i)})} - \frac{\lambda_1 + \lambda_2}{(\lambda_1 - \lambda_2)^2} \delta_{n2}. \quad (15)$$

We note that $R(\lambda, x)$ is equal to $R(\lambda)$ with u_i replaced by $\partial_x^i(f)$, where f is the initial data of the solution. Therefore, formula (15) produces all higher order logarithmic derivatives of the tau-function of u at $\mathbf{t}_{>0} = \mathbf{0}$ with the knowledge of the initial data f .

Remark 1. Tau-structures are among the most important notions in the theory of integrable systems (see e.g. [4, 22, 26, 28, 33, 42, 50, 52, 63, 64]). For the KdV hierarchy, it is known that there are several different but equivalent definitions for the canonical one [13, 28]; however, one can also define essentially different tau-structures by using normal Miura transformations [35, 42] for instance.

1.3. From wave functions to n -point correlation functions. A key ingredient in this paper (and also in the forthcoming paper [41]) will be a pair (ψ, ψ^*) , consisting of a wave function ψ and the dual wave function ψ^* associated with ψ . We explain this briefly here and in detail in Section 3.

We start with the time-independent case. Let $f(x)$ be an element of V , and L the linear Schrödinger operator $\partial_x^2 + 2f(x)$. By a *wave function* of f we will mean an element $\psi = \psi(z, x)$ in the module $\tilde{V}((z^{-1})) e^{xz}$ satisfying the equation $L(\psi) = z^2\psi$ of the form $\psi = (1 + \phi_1(x)/z + \phi_2(x)/z^2 + \dots) e^{xz}$, where $\tilde{V}$ is any ring with $V \subseteq \partial_x(\tilde{V}) \subseteq \tilde{V}$. The *dual wave function*¹ ψ^* of f associated with ψ is then defined as the unique element in $\tilde{V}((z^{-1})) e^{-xz}$ satisfying $L(\psi^*) = z^2\psi^*$ of the form $\psi^* = (1 + \phi_1^*(x)/z + \phi_2^*(x)/z^2 + \dots) e^{-xz}$ for which $\partial_x^i(\psi)\psi^*$ has residue 0 at $z = \infty$ for all $i \geq 0$. The pair (ψ, ψ^*) consisting of a wave function ψ of f and the unique dual wave function ψ^* associated with ψ is called a *pair of wave functions* of f . Given f , the wave function ψ of f is unique up to multiplication by an arbitrary element in $1 + z^{-1}\mathbb{C}((z^{-1}))$, but $\psi\psi^*$ is unique and coincides with $b(z^2, x)$, the upper right entry of the matrix $R(z^2, x)$ defined above, as is proved in Lemma 7.

We now proceed to the time-dependent case. Let $L = \partial^2 + 2u_0$ be a linear operator on $\mathcal{A}$, called the Lax operator for the KdV hierarchy, and introduce a sequence of differential operators A_k defined by $A_k = \frac{1}{(2k+1)!!} (L^{\frac{2k+1}{2}})_+$ ($k \geq 0$). Here $(\cdot)_+$ means taking the differential part of a pseudo-differential operator ([28]). Let $u = u(\mathbf{t})$ be an arbitrary solution to the KdV hierarchy (2). By a *wave function* of u we will mean an element $\psi = \psi(z, \mathbf{t})$ in $\tilde{V}[[\mathbf{t}_{>0}]]((z^{-1})) e^{\sum_{k=0}^{\infty} t_k z^{2k+1}/(2k+1)!!}$ of the form $\psi = (1 + \phi_1(\mathbf{t})/z + \phi_2(\mathbf{t})/z^2 + \dots) e^{\sum_{k=0}^{\infty} t_k z^{2k+1}/(2k+1)!!}$ satisfying the equations

$$L(\psi) = z^2\psi, \quad \partial_{t_k}(\psi) = A_k(\psi). \quad (16)$$

Define the *dual wave function* $\psi^* = \psi^*(z, \mathbf{t})$ of u associated with ψ as the unique element in $\tilde{V}[[\mathbf{t}_{>0}]]((z^{-1})) e^{-\sum_{k=0}^{\infty} t_k z^{2k+1}/(2k+1)!!}$ of the form $\psi^* = (1 + \phi_1^*(\mathbf{t})/z + \phi_2^*(\mathbf{t})/z^2 + \dots) e^{-\sum_{k=0}^{\infty} t_k z^{2k+1}/(2k+1)!!}$ satisfying

$$L(\psi^*) = z^2\psi^*, \quad -\partial_{t_k}(\psi^*) = A_k(\psi^*) \quad (17)$$

and that $\partial_x^i(\psi(z, \mathbf{t}))\psi^*(z, \mathbf{t})$ has residue 0 at $z = \infty$ for all $i \geq 0$. We say that a wave function ψ of u and the dual wave function ψ^* of u associated with ψ form a *pair of wave functions* of u . The existence of a pair of wave functions of u , which is known, will be shown in Section 3. Given a solution $u = u(\mathbf{t})$, the wave function ψ of u is unique only up to multiplication by an arbitrary element in $1 + z^{-1}\mathbb{C}((z^{-1}))$, but the product $\psi\psi^*$ is unique and coincides with $b(z^2, \mathbf{t})$, the upper right entry of $R(z^2, \mathbf{t})$, where ψ^* is the dual wave function of u associated with ψ .

Let $u = u(\mathbf{t})$ be an arbitrary solution of (2) in $V[[\mathbf{t}_{>0}]]$. The basic matrix resolvent $R(z^2, \mathbf{t})$ of u can be expressed in terms of a pair of wave functions of u (Lemma 8, Section 3). This enables us to give a new formula for the generating series of n -point correlation functions $\Omega_{p_1, \dots, p_n}(\mathbf{t})$ of u . Let $(\psi(z, \mathbf{t}), \psi^*(z, \mathbf{t}))$ be a pair of wave functions of u . Define

$$D(z, w, \mathbf{t}) := \frac{\psi(z, \mathbf{t}) \psi_x^*(w, \mathbf{t}) - \psi^*(w, \mathbf{t}) \psi_x(z, \mathbf{t})}{w^2 - z^2}. \quad (18)$$

¹A dual wave function satisfies $L^*\psi^* = \lambda\psi^*$, where L^* is the formal adjoint operator of L . For the KdV hierarchy, $L^* = L$ and a dual wave function ψ^* is also a wave function.

Theorem 1. *For any fixed $n \geq 2$ the following formula holds true:*

$$\sum_{p_1, \dots, p_n} \Omega_{p_1, \dots, p_n}(\mathbf{t}) \prod_{j=1}^n \frac{(2p_j + 1)!!}{z_j^{2p_j+2}} = - \sum_{\sigma \in S_n/C_n} \prod_{i=1}^n D(z_{\sigma(i)}, z_{\sigma(i+1)}, \mathbf{t}) - \frac{\delta_{n2}}{(z_1 - z_2)^2}. \quad (19)$$

We note that under the gauge freedom of the second type $\psi(z, \mathbf{t}) \mapsto g(z)\psi(z, \mathbf{t}) = \tilde{\psi}(z, \mathbf{t})$, the dual wave function ψ^* associated with ψ is mapped to $\psi^*(z, x)/g(z) = \tilde{\psi}^*(z, x)$. Therefore we have

$$\tilde{D}(z, w, \mathbf{t}) := \frac{\tilde{\psi}(z, \mathbf{t}) \tilde{\psi}_x^*(w, \mathbf{t}) - \tilde{\psi}^*(w, \mathbf{t}) \tilde{\psi}_x(z, \mathbf{t})}{w^2 - z^2} = \frac{g(z)}{g(w)} D(z, w, \mathbf{t}).$$

However, products of the factors of the form $g(z)/g(w)$ cancel in each sum of the right hand side of the formula (19) and therefore remain unchanged under the gauge freedom of the second type. This argument agrees with the fact that the n -point correlation functions $\Omega_{p_1, \dots, p_n}(\mathbf{t})$ of u (for $n \geq 2$) are the evaluations of certain elements of $\mathcal{A}$ with u_i replaced by $\partial_x^i(u)$.

For an element $f(x) \in V$ and a solution $u(\mathbf{t}) \in V[[\mathbf{t}_{>0}]]$ to the KdV hierarchy related via the 1-1 correspondence (4), we have $\psi(z, x, \mathbf{0}) \equiv \psi(z, x)$. For a given f , one can find an explicit recursion for solving $\psi(z, x)$ (see Section 3). However, we do not know an efficient way of solving $\psi(z, \mathbf{t})$. This is because the recursive procedure of solving $\psi(z, \mathbf{t})$ requires the knowledge of $u(\mathbf{t})$, but this requires first solving the KdV hierarchy. However, due to (19), the knowledge of $\psi(z, x)$ gives rise to a construction of the logarithm of the tau-function $\tau(\mathbf{t})$ of u (since $u = \partial_x^2(\log \tau)$, (19) also constructs u) in a simple way. More precisely, if we specialize (19) to $\mathbf{t} = (x, \mathbf{0})$, then it gives a formula $\Omega_{p_1, \dots, p_n}(x)$ for all $n \geq 2$ and $p_i \geq 0$ and hence, in view of (10), gives the entire Taylor series of $\log \tau(\mathbf{t})$ (which, we recall, is only defined up to the addition of a linear term). This is important because in our concrete computations for Examples 1–3 of Section 1.1, we can compute $\psi(z, x)$ and $\psi^*(z, x)$, and hence the function $D(z, w, x)$, explicitly.

Remark 2. For $V = \mathbb{C}[[x]]$, one could use alternatively the Sato Grassmannian approach [13, 26, 63, 72] to prove the identity (19). It would be interesting to investigate if the identity (19) could also be proved, say for $V = \mathbb{C}((x))$, by using the approach of Bergère and Eynard with the employment of topological recursion (loop equation) [1, 8, 10, 24, 29, 45, 59, 69], or by using matrix models together with appropriate Riemann–Hilbert problems (isomonodromic deformations) [12, 16, 17, 27, 29, 37, 39, 49, 57, 60, 66], or by using OPE from appropriate vertex algebras [5, 26, 47, 72]; we expect that at least for the initial data $f(x) \in V$ having the *bispectral* property defined by Duistermaat and Grünbaum [43] (see also the M -bispectrality given in Section 6.2) this might be possible. We also note that the matrix resolvent method and some of the above-mentioned methods extend to new situations (see e.g. [14, 15, 37, 38, 39, 40], [3], [8, 9, 11, 59] and [16, 17, 18, 51]).

We now formulate a theorem that is equivalent to Theorem 1 but involves a function $K(z, w, \mathbf{t})$ which, unlike $D(z, w, \mathbf{t})$, depends only on the solution $u(\mathbf{t})$ (i.e. it is independent of the choice of a pair of wave functions of $u(\mathbf{t})$). The function $K(z, w, \mathbf{t})$ is defined by

$$K(z, w, \mathbf{t}) := \frac{b(z^2, \mathbf{t}) b_x(w^2, \mathbf{t}) - b(w^2, \mathbf{t}) b_x(z^2, \mathbf{t})}{2(w^2 - z^2)} - \frac{w b(z^2, \mathbf{t}) + z b(w^2, \mathbf{t})}{w^2 - z^2}. \quad (20)$$

Theorem 2. *For any fixed $n \geq 2$ the following formula holds true:*

$$\sum_{p_1, \dots, p_n} \Omega_{p_1, \dots, p_n}(\mathbf{t}) \prod_{j=1}^n \frac{(2p_j + 1)!!}{z_j^{2p_j+2}} = - \frac{1}{\prod_{i=1}^n b(z_i^2, \mathbf{t})} \sum_{\sigma \in S_n/C_n} \prod_{i=1}^n K(z_{\sigma(i)}, z_{\sigma(i+1)}, \mathbf{t}) - \frac{\delta_{n2}}{(z_1 - z_2)^2}. \quad (21)$$

Remark 3. Both the formula (19) and the formula (21) are generalized to the Toda lattice hierarchy in [68]. For example, the following theorem is proved in [68].

Theorem ([68]). *Fix $k \geq 2$ an integer. For an arbitrary solution (v, w) to the Toda lattice hierarchy associated with the Lax operator $L_{\text{Toda}} = \Lambda + v + w\Lambda^{-1}$, let $\psi_1(\lambda, n, \mathbf{t}), \psi_2(\lambda, n, \mathbf{t})$ be a pair of wave functions of the solution (v, w) . Here Λ denotes the shift operator $\Lambda : f(n) \mapsto f(n+1)$. Define*

$$D(\lambda, \mu, n, \mathbf{t}) = \frac{\psi_1(\lambda, n, \mathbf{t}) \psi_2(\mu, n-1, \mathbf{t}) - \psi_1(\lambda, n-1, \mathbf{t}) \psi_2(\mu, n, \mathbf{t})}{\lambda - \mu}.$$

Then the following formula holds true:

$$\sum_{i_1, \dots, i_k \geq 0} \frac{\Omega_{i_1, \dots, i_k}^{\text{Toda}}(n, \mathbf{t})}{\lambda_1^{i_1+2} \dots \lambda_k^{i_k+2}} = (-1)^{k-1} \frac{e^{kq(n-1, \mathbf{t})}}{\prod_{j=1}^k \lambda_j} \sum_{\sigma \in S_k / C_k} \prod_{j=1}^k D(\lambda_{\sigma(j)}, \lambda_{\sigma(j+1)}, n, \mathbf{t}) - \frac{\delta_{k,2}}{(\lambda_1 - \lambda_2)^2}, \quad (22)$$

where $\Omega_{i_1, \dots, i_k}^{\text{Toda}}(n, \mathbf{t})$ denote the k -point correlation functions of (v, w) , and the $q(n, \mathbf{t})$ is defined via

$$w(n, \mathbf{t}) = e^{q(n-1, \mathbf{t}) - q(n, \mathbf{t})}, \quad \frac{\partial q(n, \mathbf{t})}{\partial t_i} = -S_i(n, \mathbf{t}), \quad i \geq 0.$$

See [68] for the precise definitions of the functions $S_i(n, \mathbf{t})$, $\Omega_{i_1, \dots, i_k}^{\text{Toda}}(n, \mathbf{t})$, and of a pair of wave functions for the Toda lattice hierarchy.

We observe that all the following three functions

$$K(z, w, \mathbf{t}) - \frac{b(w^2, \mathbf{t})}{z - w}, \quad K(z, w, \mathbf{t}) - \frac{b(z^2, \mathbf{t})}{z - w}, \quad K(z, w, \mathbf{t}) - \frac{1}{2} \frac{b(z^2, \mathbf{t}) + b(w^2, \mathbf{t})}{z - w}$$

belong to the ring $V[[\mathbf{t}_{>0}]][[z^{-1}, w^{-1}]]$. We also remark that if we define K as the right hand side of (20) with $b(z^2, \mathbf{t})$ and $b(w^2, \mathbf{t})$ replaced by $b(z^2)$ and $b(w^2)$, respectively, then it easily follows that the three functions $K - b(z^2)/(z - w)$, $K - b(w^2)/(z - w)$ and $K - \frac{1}{2}(b(z^2) + b(w^2))/(z - w)$ all belong to $\mathcal{A}[[z^{-1}, w^{-1}]]$. Moreover, the identity (21) is then also true in the **abstract sense** (similarly to Proposition 1). We have for example

$$\begin{aligned} K - \frac{1}{2} \frac{b(z^2) + b(w^2)}{z - w} &= \frac{u_0}{2} \left(\frac{1}{zw^2} - \frac{1}{z^2w} \right) - \frac{u_1}{2} \frac{1}{z^2w^2} + \frac{6u_0^2 + u_2}{8} \left(\frac{1}{zw^4} - \frac{1}{z^2w^3} + \frac{1}{z^3w^2} - \frac{1}{z^4w} \right) \\ &\quad - \frac{12u_0u_1 + u_3}{8} \left(\frac{1}{z^2w^4} + \frac{1}{z^4w^2} \right) + \dots \end{aligned}$$

Proposition 2. *Let $\psi(z, \mathbf{t})$ be any wave function of $u(\mathbf{t})$. The functions K and D satisfy the relation*

$$K(z, w, \mathbf{t}) = b(z^2, \mathbf{t}) \frac{\psi(w, \mathbf{t})}{\psi(z, \mathbf{t})} D(z, w, \mathbf{t}) = b(w^2, \mathbf{t}) \frac{\psi^*(z, \mathbf{t})}{\psi^*(w, \mathbf{t})} D(z, w, \mathbf{t}) = \psi^*(z, \mathbf{t}) \psi(w, \mathbf{t}) D(z, w, \mathbf{t}). \quad (23)$$

The proofs of Proposition 2 and Theorem 2 will be given in Section 3.

Let us now consider the case $V = \mathbb{C}[[x]]$, namely, consider arbitrary solutions of the KdV hierarchy in the ring $\mathbb{C}[[t_0, t_1, t_2, \dots]]$, where we recall that $t_0 = x$. Define $D(z, w)$ and $K(z, w)$ as follows:

$$D(z, w) := D(z, w, \mathbf{0}) = \frac{\psi(z, \mathbf{0}) \psi_x^*(w, \mathbf{0}) - \psi^*(w, \mathbf{0}) \psi_x(z, \mathbf{0})}{w^2 - z^2}, \quad K(z, w) := K(z, w, \mathbf{0}). \quad (24)$$

Then Proposition 2 immediately implies the following corollary.

Corollary 2. *For $V = \mathbb{C}[[x]]$, we have*

$$D(z, w) - \frac{1}{z - w} \in \mathbb{C}[[z^{-1}, w^{-1}]], \quad (25)$$

$$K(z, w) - \frac{b(z^2, \mathbf{0})}{z - w}, \quad K(z, w) - \frac{b(w^2, \mathbf{0})}{z - w}, \quad K(z, w) - \frac{1}{2} \frac{b(z^2, \mathbf{0}) + b(w^2, \mathbf{0})}{z - w} \in \mathbb{C}[[z^{-1}, w^{-1}]]. \quad (26)$$

Note that for this case, pairs of wave functions of the solution correspond to certain points of the Sato Grassmannian. In such correspondences, the coefficients of $D(z, w) - 1/(z - w)$ coincide with the affine coordinates [44] of the point of the Sato Grassmannian. Based on this observation we find that for $V = \mathbb{C}[[x]]$ the identity (19) evaluated at $\mathbf{t} = \mathbf{0}$ is equivalent to a formula obtained by J. Zhou [72] from the fermionic picture (see the Theorem 5.3 of [72]). The affine coordinates can alternatively be used to compute the Plücker coordinates in the expansion of tau-function (as opposed to its logarithm) (cf. [6, 44, 63, 70, 72]). Geometric meaning (in Sato Grassmannian) of the coefficients of the power series in (26) remains an interesting open question, which we plan to do in a subsequent publication.

1.4. Applications. We now discuss each of the three examples from Section 1.1 in turn.

Example 1: Intersection numbers of ψ -classes. An explicit formula for the generating series of (5) in terms of the basic matrix resolvent was derived in [13]; see also (128) and (129). A particular pair of wave functions $(\psi(z, x), \psi^*(z, x))$ of $f = x$ were given in [13] (cf. also [21, 54]):

$$\psi(z, x) = \frac{\sqrt{z}}{(z^2 - 2x)^{\frac{1}{4}}} e^{\frac{1}{3}z^3 - \frac{1}{3}(z^2 - 2x)^{\frac{3}{2}}} \sum_{k \geq 0} \frac{(-1)^k}{288^k} \frac{(6k)!}{(3k)!(2k)!} (z^2 - 2x)^{-\frac{3k}{2}}, \quad (27)$$

$$\psi^*(z, x) = \frac{\sqrt{z}}{(z^2 - 2x)^{\frac{1}{4}}} e^{-\frac{z^3}{3} + \frac{1}{3}(z^2 - 2x)^{\frac{3}{2}}} \sum_{k \geq 0} \frac{1}{288^k} \frac{(6k)!}{(3k)!(2k)!} (z^2 - 2x)^{-\frac{3k}{2}}. \quad (28)$$

They can be viewed as the asymptotic expansion of the following two analytic functions, respectively, as $z \rightarrow \infty$ within appropriate sectors [13, 19]:

$$\sqrt{2\pi z} e^{\frac{z^3}{3}} 2^{\frac{1}{3}} \text{Ai}(\xi), \quad e^{\frac{\pi i}{6}} \sqrt{2\pi z} e^{-\frac{z^3}{3}} 2^{\frac{1}{3}} \text{Ai}(\omega \xi),$$

where $\xi = 2^{-\frac{2}{3}}(z^2 - 2x)$, $\omega = e^{2\pi i/3}$. Restricting the formal series (27)–(28) to $x = 0$ we obtain

$$\psi(z, 0) = \sum_{k=0}^{\infty} \frac{(-1)^k}{288^k} \frac{(6k)!}{(3k)!(2k)!} z^{-3k}, \quad \psi^*(z, 0) = \psi(-z, 0), \quad (29)$$

$$\psi_x(z, 0) = \sum_{k=0}^{\infty} \frac{1+6k}{1-6k} \frac{(-1)^k}{288^k} \frac{(6k)!}{(3k)!(2k)!} z^{-3k+1}, \quad \psi_x^*(z, 0) = \psi_x(-z, 0). \quad (30)$$

These series appeared in the Faber–Zagier formula [46] on relations in the tautological ring of the moduli space of curves. Now consider the function $D(z, w) = D(z, w, \mathbf{0})$. According to (25) we know that this function has a pole only at $z = w$. The first few terms for $D(z, w) - 1/(z - w)$ are given by

$$\begin{aligned} D(z, w) - \frac{1}{z - w} &= \frac{5}{24} z^{-1} w^{-3} - \frac{7}{24} z^{-2} w^{-2} + \frac{5}{24} z^{-3} w^{-1} - \frac{455}{1152} z^{-2} w^{-5} \\ &\quad + \frac{385}{1152} z^{-3} w^{-4} - \frac{385}{1152} z^{-4} w^{-3} + \frac{455}{1152} z^{-5} w^{-2} + \dots \end{aligned} \quad (31)$$

Theorem 1 then implies the following slightly simplified version of a formula of J. Zhou:

Theorem (Zhou [72]). *For $n \geq 2$, the generating series of the n -point intersection numbers of ψ -classes has the expression*

$$\sum_{g \geq 0} \sum_{p_1, \dots, p_n \geq 0} \int_{\mathcal{M}_{g,n}} \psi_1^{p_1} \dots \psi_n^{p_n} \prod_{j=1}^n \frac{(2p_j + 1)!!}{z_j^{2p_j+2}} = - \sum_{\sigma \in S_n/C_n} \prod_{i=1}^n D(z_{\sigma(i)}, z_{\sigma(i+1)}) - \frac{\delta_{n2}}{(z_1 - z_2)^2}. \quad (32)$$

We observe that an explicit formula for the coefficients in (31) was obtained by J. Zhou [70] by solving Virasoro constraints using the *fermionic* method. Later the explicit formula was re-proved by F. Balogh and one of the authors of the present paper [6] using the Sato Grassmannian approach [63].

The Laurent series of the corresponding function $K(z, w)$ begins

$$K(z, w) - \frac{1}{2} \frac{b(z^2, \mathbf{0}) + b(w^2, \mathbf{0})}{z - w} = -\frac{1}{2} \frac{1}{z^2 w^2} + \frac{5}{16} \left(\frac{1}{z w^6} - \frac{1}{z^2 w^5} + \frac{1}{z^3 w^4} - \frac{1}{z^4 w^3} + \frac{1}{z^5 w^2} - \frac{1}{z^6 w} \right) + \cdots \quad (33)$$

Example 2: Theta-classes. For the BGW-Norbury solution $u_{\Theta(1/8)}$ introduced in Example 2 (with $C = 1/8$), denote by $\Omega_{p_1, \dots, p_n}^{\Theta(1/8)}(\mathbf{t})$ the n -point correlation functions of $u_{\Theta(1/8)}$. According to Norbury [62] $\Omega_{p_1, \dots, p_n}^{\Theta(1/8)}(\mathbf{t})$ for $n \geq 2$ are related to the Θ -class intersection numbers by

$$\Omega_{p_1, \dots, p_n}^{\Theta(1/8)}(\mathbf{0}) = \int_{\overline{\mathcal{M}}_{g,n}} \Theta_{g,n} \psi_1^{p_1} \cdots \psi_n^{p_n} \quad (g = p_1 + \cdots + p_n + 1). \quad (34)$$

(The expression on the right vanishes for $g \neq p_1 + \cdots + p_n + 1$.) For any $n \geq 1$, denote by $F(\lambda_1, \dots, \lambda_n)$ the following generating series of the n -point Θ -class intersection numbers

$$F(\lambda_1, \dots, \lambda_n) = \sum_{p_1, \dots, p_n} \prod_{j=1}^n \frac{(2p_j + 1)!!}{\lambda_j^{p_j+1}} \int_{\overline{\mathcal{M}}_{1+p_1+\cdots+p_n,n}} \Theta_{1+p_1+\cdots+p_n,n} \psi_1^{p_1} \cdots \psi_n^{p_n}. \quad (35)$$

Using the relationship (34) and Corollary 1 we will prove:

Theorem 3. *For fixed $n \geq 2$, the generating series $F(\lambda_1, \dots, \lambda_n)$ has the expression*

$$F(\lambda_1, \dots, \lambda_n) = - \sum_{\sigma \in S_n / C_n} \frac{\text{tr}(M(\lambda_{\sigma(1)}) \cdots M(\lambda_{\sigma(n)}))}{\prod_{i=1}^n (\lambda_{\sigma(i+1)} - \lambda_{\sigma(i)})} - \delta_{n,2} \frac{\lambda_1 + \lambda_2}{(\lambda_1 - \lambda_2)^2}, \quad (36)$$

where $M(\lambda) = \begin{pmatrix} 0 & 0 \\ \lambda & 0 \end{pmatrix} + \sum_{k \geq 0} \left[\frac{(2k-1)!!}{2^k} \right]^3 \begin{pmatrix} k & 1 \\ -\frac{8k^3+12k^2+4k+1}{8(k+1)} & -k \end{pmatrix} \frac{\lambda^{-k}}{k!}.$

The string type equation (see in Section 4) relates the 2-point Θ -class intersection numbers to 1-point intersection numbers, from which we deduce:

Corollary 3. *The 1-point Θ -class intersection numbers have the expression*

$$\int_{\overline{\mathcal{M}}_{g,1}} \Theta_{g,1} \psi_1^{g-1} = \frac{(2g-1)!!^2}{8^g g! (2g-1)}, \quad \forall g \geq 1. \quad (37)$$

For a general C in Example 2, let $\tau_{\Theta(C)}(\mathbf{t})$ be the tau-function of $u_{\Theta(C)}$, $R(\lambda, \mathbf{t})$ the basic matrix resolvent of $u_{\Theta(C)}$, and $\Omega_{p_1, \dots, p_n}^{\Theta(C)}(\mathbf{t})$ the n -point correlation functions of $u_{\Theta(C)}$. We will call $\Omega_{p_1, \dots, p_n}^{\Theta(C)}(\mathbf{0})$ the n -point generalized BGW correlators. For $n \geq 2$, $\Omega_{p_1, \dots, p_n}^{\Theta(C)}(\mathbf{0})$ are uniquely determined by $u_{\Theta(C)}$ and so by f , where $f = \frac{C}{(x-1)^2}$. To determine $\Omega_p^{\Theta(C)}(\mathbf{0})$, we impose a string-type equation for $\tau_{\Theta(C)}$

$$\sum_{i \geq 0} (1+2i) t_i \frac{\partial \log \tau_{\Theta(C)}}{\partial t_i} + C = \frac{\partial \log \tau_{\Theta(C)}}{\partial t_0}. \quad (38)$$

Since $\sum_{i \geq 0} (1+2i) t_i \frac{\partial}{\partial t_i}$ gives the infinitesimal scaling symmetry of the KdV hierarchy and since eq. (38) implies that $u_{\Theta(C)}(x, \mathbf{0}) = C/(x-1)^2$, the tau-function $\tau_{\Theta(C)}$ of $u_{\Theta(C)}$ satisfying the additional constraint (38) exists. (From this way, $\tau_{\Theta(C)}$ is determined up to a multiplicative constant.) A matrix model for $\tau_{\Theta(C)}$ was obtained by A. Alexandrov [2]. The function $\tau_{\Theta(C)}$ is called the *generalized BGW tau-function*. As before denote $R(\lambda, x) = R(\lambda, x, \mathbf{0})$, $b(\lambda, x) = R(\lambda, x)_{12}$.

Theorem 4. For any fixed $C \in \mathbb{C}$, the formal power series $b(\lambda, x)$ of λ^{-1} only depends on the variable $\zeta := \lambda(x-1)^2$. Explicitly, the matrix-valued formal series $R = R(\lambda, x)$ has the explicit expression

$$R = \begin{pmatrix} -\frac{C\lambda^{\frac{1}{2}}}{\zeta^{\frac{3}{2}}} G_{\frac{3}{2}}(\zeta) & G_{\frac{1}{2}}(\zeta) \\ \frac{\lambda}{\zeta} \left((\zeta - 2C) G_{\frac{1}{2}}(\zeta) - \frac{3C}{\zeta} G_{\frac{3}{2}}(\zeta) - \frac{6C(C+1)}{\zeta^2} G_{\frac{5}{2}}(\zeta) \right) & \frac{C\lambda^{\frac{1}{2}}}{\zeta^{\frac{3}{2}}} G_{\frac{3}{2}}(\zeta) \end{pmatrix}, \quad (39)$$

where $\zeta^{3/2} := \lambda^{3/2}(x-1)^3$ and we have set

$$\Delta := 1 - 8C, \quad G_\alpha(\zeta) := {}_3F_0\left(\alpha, \alpha + \frac{\sqrt{\Delta}}{2}, \alpha - \frac{\sqrt{\Delta}}{2}; ; \frac{1}{\zeta}\right). \quad (40)$$

The proof is in Section 4. It is interesting to mention that if $-C$ is a triangular number, i.e. $C = -\frac{1}{2}p(p+1)$ ($p \geq 0$) then $b(\lambda, x)$ truncates to a polynomial of λ^{-1} .

Corollary 4. The 1-point generalized BGW correlators have the explicit expression

$$\Omega_p^{\Theta(C)}(\mathbf{0}) = \frac{1}{(p+1)!(1+2p)} \prod_{i=0}^p \left(C + \frac{i(i+1)}{2} \right), \quad j \geq 0. \quad (41)$$

We proceed to applying Theorem 1 to the computation of generalized BGW correlators.

Theorem 5. Let $\alpha = \sqrt{\frac{1}{4} - 2C}$, and for $k \geq 0$ define $a_k(\alpha) \in \mathbb{Q}[\alpha^2] = \mathbb{Q}[C]$ by

$$a_k(\alpha) = \frac{(\alpha - k + \frac{1}{2})_{2k}}{2^k k!} = \frac{(-1)^k}{k!} \prod_{j=1}^k \left(C + \binom{j}{2} \right) = (-1)^k (2k-1) \Omega_{k-1}^{\Theta(C)}(\mathbf{0}).$$

Then the two functions

$$\psi(z, x) = \sum_{k \geq 0} \frac{a_k(\alpha)}{z^k (1-x)^k} e^{zx}, \quad \psi^*(z, x) = \psi(-z, x) \quad (42)$$

form a pair of wave functions of $f = C/(x-1)^2$. Moreover, if $D_{\Theta(C)}(z, w) = D(z, w)$ is the corresponding function as defined in (24), then for any $n \geq 2$ we have the generating function

$$\sum_{p_1, \dots, p_n} \Omega_{p_1, \dots, p_n}^{\Theta(C)}(\mathbf{0}) \prod_{j=1}^n \frac{(2p_j+1)!!}{z_j^{2p_j+2}} = - \sum_{\sigma \in S_n/C_n} \prod_{i=1}^n D_{\Theta(C)}(z_{\sigma(i)}, z_{\sigma(i+1)}) - \frac{\delta_{n2}}{(z_1 - z_2)^2}. \quad (43)$$

According to equation (25) we know that $D(z, w)$ has an expansion of the form

$$D_{\Theta(C)}(z, w) = \frac{1}{z-w} + \sum_{m, n \geq 0} \frac{A_{mn}(\alpha)}{z^m (-w)^n} \quad (44)$$

with coefficients $A_{mn}(\alpha) \in \mathbb{Q}[\alpha]$. These coefficients are given explicitly by the following proposition.

Proposition 3. The coefficients A_{mn} can be given either by the recursion and boundary conditions

$$A_{m, n+1}(\alpha) - A_{m+1, n}(\alpha) = \frac{m-n}{m+n} a_m(\alpha) a_n(\alpha), \quad A_{m, 0}(\alpha) = A_{0, n}(\alpha) = 0 \quad (45)$$

or else explicitly by any of the three formulas

$$A_{mn}(\alpha) = \sum_{\substack{r \geq m, s \geq 0 \\ r+s=m+n-1}} \frac{r-s}{r+s} a_r(\alpha) a_s(\alpha) = \sum_{\substack{r \geq n, s \geq 0 \\ r+s=m+n-1}} \frac{r-s}{r+s} a_r(\alpha) a_s(\alpha) \quad (46)$$

$$= 2 \frac{m! n! a_m(\alpha) a_n(\alpha)}{(m+n-1)(m+n-1)!} \sum_{k \in \mathbb{Z}} (-1)^k \frac{\binom{m+n-1}{m+k} \binom{m+n-1}{n+k}}{\alpha - k - \frac{1}{2}}. \quad (47)$$

Observe that the summation in (47) is finite (since the k_{th} term vanishes unless $|k + 1/2| < m, n$) and is even in α (as one sees by sending α to $-\alpha$ and k to $-k - 1$), and that either formula (46) or (47) implies that $A_{mn}(\alpha)/a_m(\alpha)$ for fixed m and n is a polynomial of degree $n - 1$ in α^2 .

The Laurent series of the corresponding function $K(z, w)$ (defined in (24)) begins

$$\begin{aligned} K(z, w) &= \frac{1}{2} \frac{b(z^2, \mathbf{0}) + b(w^2, \mathbf{0})}{z - w} \\ &= \frac{C}{2} \left(\frac{1}{zw^2} - \frac{1}{z^2w} \right) - \frac{C}{z^2w^2} + \frac{3C(1+C)}{4} \left(\frac{1}{zw^4} - \frac{1}{z^2w^3} + \frac{1}{z^3w^2} - \frac{1}{z^4w} \right) + \cdots \end{aligned} \quad (48)$$

We also note that, similarly as above, the pair of wave functions ψ, ψ^* of the above theorem can be viewed as the asymptotic expansions of the following two analytic functions, respectively, as z goes to ∞ within a certain sector of the complex z -plane:

$$\sqrt{\frac{2\xi}{\pi}} e^z K_\alpha(\xi), \quad \sqrt{2\pi\xi} e^{-z} I_\alpha(\xi), \quad (49)$$

where I_α and K_α denote the modified Bessel functions and $\xi = z(1 - x)$. Finally we note that the function $D_{\Theta(C)}(z, w)$ in terms of these analytic functions reads

$$D_{\Theta(C)}(z, w) = 2e^{z-w} \sqrt{zw} \frac{K_\alpha(z) w I'_\alpha(w) - I_\alpha(w) z K'_\alpha(z)}{z^2 - w^2}, \quad (50)$$

which can be recognized as a Bessel kernel (cf. [7, 65]).

Example 3: Lamé solutions. Let τ_{elliptic} be the tau-function of the Lamé solution $u_{\text{elliptic}}(\mathbf{t})$ introduced above, which as we will see gives an elliptic generalization of $\tau_{\Theta(C)}$. Denote by $R(\lambda, \mathbf{t})$ the basic matrix resolvent of u_{elliptic} , and denote $R(\lambda, x) = R(\lambda, x, \mathbf{0})$ and $b(\lambda, x) = R(\lambda, x)_{12}$. If we restrict to the particular case when $-C$ is a triangular number, i.e. $C = -\frac{p(p+1)}{2}$ for some integer $p \geq 0$, then $u_{\text{elliptic}}(\mathbf{t})$ is a special *finite-gap* solution [36, 32] of the KdV hierarchy. Denote by

$$y^2 = S_p(\lambda) \quad (51)$$

the corresponding *spectral curve* [36, 32], where $S_p(\lambda)$ is a polynomial of λ of degree $1 + 2p$ with leading coefficient 1. The first few S_p are $S_0 = \lambda$, $S_1 = \lambda^3 - \frac{g_2}{4}\lambda - \frac{g_3}{4}$, $S_2 = (\lambda^2 - 3g_2)(\lambda^3 - \frac{9}{4}g_2\lambda + \frac{27}{4}g_3)$. The following theorem will be deduced from results of [36, 32, 34] in Section 5.

Theorem 6. *In the case that $C = -\frac{p(p+1)}{2}$ ($p \geq 0$), the product $\sqrt{\frac{S_p(\lambda)}{\lambda}} \phi(\lambda, x)$ is a degree p polynomial in λ with leading coefficient 1 and a degree p polynomial in $\wp = \wp(x; \tau)$. For $p = 1$,*

$$b = 2\sqrt{\lambda} \frac{\lambda - \wp}{\sqrt{4\lambda^3 - g_2\lambda - g_3}};$$

For $p = 2$,

$$b = \frac{1}{2}\sqrt{\lambda} \frac{4\lambda^2 - 12\wp\lambda + 36\wp^2 - 9g_2}{\sqrt{(\lambda^2 - 3g_2)(4\lambda^3 - 9g_2\lambda + 27g_3)}};$$

For $p = 3$,

$$b = \frac{4\lambda^3 - 24\wp\lambda^2 + 60(3\wp^2 - g_2)\lambda - 900\wp^3 + 225g_2\wp + 225g_3}{\sqrt{16\lambda^6 - 504g_2\lambda^4 + 2376g_3\lambda^3 + 4185g_2^2\lambda^2 - 36450g_2g_3\lambda - 3375g_2^3 + 91125g_3^2}}.$$

Organization of the paper. In Section 2 we review the MR approach. In Section 3 we prove Theorem 1. In Section 4 we prove Theorems 3, 4, 5, and compute explicitly some generalized BGW correlators. In Section 5 we compute Lamé correlators. Further remarks are given in Section 6.

Acknowledgements. We would like to express our thanks to Paul Norbury for helpful suggestions and to the anonymous referee for many constructive comments. D.Y. is grateful to Youjin Zhang for his advice and encouragement and to Chang-Shou Lin and Alexander Alexandrov for their interest. Part of the work of D.Y. was done in MPIM, Bonn while he was a postdoc; he acknowledges MPIM for excellent working conditions and financial supports.

Note added. After this paper was finished we learned that the identities (36)–(37), (39)–(41), which had been presented together with their proofs by one of the authors [67], had also been found by M. Bertola and G. Ruzza (see [17], whose arXiv version appeared slightly before the arXiv version of this paper), but with a quite different method using matrix models together with the construction of appropriate Riemann–Hilbert problems.

2. REVIEW OF THE MATRIX RESOLVENT APPROACH TO THE KDV HIERARCHY

We continue with more details about the MR approach to the study of tau-functions.

2.1. Fundamental lemma. Recall from Introduction some of our notations $\mathcal{A} = \mathbb{Q}[u_0, u_1, u_2, \dots]$, $\Lambda(\lambda) = \begin{pmatrix} 0 & 1 \\ \lambda & 0 \end{pmatrix}$, $q = \begin{pmatrix} 0 & 0 \\ -2u_0 & 0 \end{pmatrix}$, $\mathcal{L} = \partial + \Lambda + q$, and $\mathcal{S} = \mathfrak{sl}_2(\mathbb{C})((\lambda^{-1}))$. Denote by $(\cdot|\cdot)$ the *normalized* Cartan–Killing form. We have $(A|B) = \text{Tr } AB$, $\forall A, B \in \mathfrak{sl}_2(\mathbb{C})$. This bilinear form naturally extends to $\mathcal{S}$. According to B. Kostant [56] and V. Kac [53], the space $\mathcal{S}$ admits the decomposition

$$\mathcal{S} = \text{Ker ad}_\Lambda \oplus \text{Im ad}_\Lambda, \quad \text{Ker ad}_\Lambda \perp \text{Im ad}_\Lambda, \quad (52)$$

$$\text{Ker ad}_\Lambda = \left\{ \sum_{\ell \leq m} c_\ell \Lambda_{1+2\ell} \mid m \in \mathbb{Z}, c_\ell \in \mathbb{C}, c_m \neq 0 \right\}, \quad (53)$$

where $\Lambda_{1+2\ell} := \Lambda \lambda^\ell$. It is easy to see that $\deg \Lambda_{1+2\ell} = 1 + 2\ell$ (recall that $\deg$ denotes the principal gradation) and that $(\Lambda|\Lambda) = 2\lambda$.

Lemma 1 (Drinfeld–Sokolov [31]). *There exists a unique pair (U, H) such that*

$$U \in \mathcal{A} \otimes (\text{Im ad}_\Lambda)^{<0}, \quad H \in \mathcal{A} \otimes (\text{Ker ad}_\Lambda)^{<0}, \quad (54)$$

$$e^{-\text{ad}_U} \mathcal{L} = \partial + \Lambda + H. \quad (55)$$

The proof, using (52)–(53) and the mathematical induction, can be found in [31, 15].

2.2. Matrix resolvent recursive relations. Let us prove that equations in (11) has a unique solution R . The first equation of (11) reads $\partial(R) + [\Lambda + q, R] = 0$. Write

$$R = \begin{pmatrix} a & b \\ c & -a \end{pmatrix}. \quad (56)$$

Then it follows that

$$a = \frac{1}{2} \partial(b), \quad (57)$$

$$c = (\lambda - 2u_0)b - \frac{1}{2} \partial^2(b), \quad (58)$$

$$\partial^3(b) - 4(\lambda - 2u_0)\partial(b) + 4u_1b = 0. \quad (59)$$

Substituting (57)–(58) into the 3rd equation of (11) we find

$$b\partial^2(b) - \frac{1}{2}(\partial(b))^2 - 2(\lambda - 2u_0)b^2 = -2\lambda. \quad (60)$$

Applying ∂ to both sides of this equation yields (59). This shows compatibility between the 1st and the 3rd equations in (11). The 2nd equation of (11) is also compatible with the 3rd one. This gives

the existence of R . Let us now prove the uniqueness. According to the 2nd equation in (11) we have

$$b = \sum_{k \geq -1} \frac{b_k}{\lambda^{k+1}}, \quad b_k \in \mathcal{A}, \quad b_{-1} = 1. \quad (61)$$

Then (59) and (60) imply that b_k satisfies the following recursive relations: for $k \geq 0$,

$$\partial(b_k) = 2u_0 \partial(b_{k-1}) + u_1 b_{k-1} + \frac{1}{4} \partial^3(b_{k-1}), \quad (62)$$

$$b_k = \sum_{\substack{k_1, k_2 \geq -1 \\ k_1 + k_2 = k-2}} \left[\left(\frac{b_{k_1} \partial^2(b_{k_2})}{4} - \frac{\partial(b_{k_1}) \partial(b_{k_2})}{8} \right) + u_0 b_{k_1} b_{k_2} \right] - \frac{1}{2} \sum_{\substack{k_1 \geq 0, k_2 \geq -1 \\ k_1 + k_2 = k-2}} b_{k_1} b_{k_2+1}, \quad (63)$$

which will be referred as the *matrix resolvent recursive relations*. (One recognizes that (62) is the Lenard–Magri recursion.) Formula (63) uniquely determines $b_0, b_1, b_2, \dots$ with the first few being

$$b_0 = u_0, \quad b_1 = \frac{3}{2} u_0^2 + \frac{1}{4} u_2. \quad (64)$$

We note that the formal series $b = b(\lambda)$ determined here coincides with b in the Introduction.

Lemma 2. *The basic matrix resolvent can be expressed as $R = e^{\text{ad}_U}(\Lambda)$.*

Proof. Applying e^{ad_U} to $[\partial + \Lambda + H, \Lambda] \equiv 0$ gives $[\mathcal{L}, e^{\text{ad}_U}(\Lambda)] = 0$. Since $\deg U < 0$, the matrix $R := e^{\text{ad}_U}(\Lambda)$ satisfies the 2nd equation in (11). Finally, $\text{Tr } R^2 = (\Lambda|\Lambda) = 2\lambda$. The lemma is proved. $\square$

Definition 1. Define a family of derivations D_k ($k \geq 0$) by means of (3) with

$$Q_k := \frac{\partial(b_k)}{(2k+1)!!} - \frac{u_0^k}{k!} u_1. \quad (65)$$

2.3. From matrix resolvent to tau-structure. The basic matrix resolvent R has the expression

$$R = \begin{pmatrix} \frac{1}{2} \sum_{i \geq -1} \partial(b_i) \lambda^{-1-i} & \sum_{i \geq -1} b_i \lambda^{-1-i} \\ \sum_{i \geq -1} \lambda^{-1-i} \left((\lambda - 2u) b_i - \frac{\partial^2(b_i)}{2} \right) & -\frac{1}{2} \sum_{i \geq -1} \partial(b_i) \lambda^{-1-i} \end{pmatrix}, \quad (66)$$

where we recall that $b_k \in \mathcal{A}$ ($k \geq -1$) are uniquely determined by (63) with $b_{-1} = 1$. Introduce

$$V_k := -\frac{1}{(2k+1)!!} (\lambda^k R(\lambda))_+ + \frac{1}{(2k+1)!!} \begin{pmatrix} 0 & 0 \\ b_k & 0 \end{pmatrix}, \quad k \geq 0. \quad (67)$$

Lemma 3. *The derivations D_k ($k \geq 0$) defined in Definition 1 satisfy*

$$[D_k - V_k, \mathcal{L}] = 0, \quad k \geq 0. \quad (68)$$

Proof. Since D_k commutes with $D_0 = \partial$, equation (68) can be written equivalently as

$$D_k(q) = [V_k, \mathcal{L}].$$

Noting that

$$V_k = \frac{1}{(1+2k)!!} \begin{pmatrix} -\frac{1}{2} \sum_{i=-1}^{k-1} \partial(b_i) \lambda^{k-1-i} & -\sum_{i=-1}^{k-1} b_i \lambda^{k-1-i} \\ \sum_{i=-1}^{k-1} \lambda^{k-1-i} \left(\frac{\partial^2(b_i)}{2} - (\lambda - 2u_0) b_i \right) & \frac{1}{2} \sum_{i=-1}^{k-1} \partial(b_i) \lambda^{k-1-i} \end{pmatrix}$$

we have by a direct calculation that

$$[V_k, \mathcal{L}] = [V_k, \partial + \Lambda + q] = \frac{1}{(1+2k)!!} \begin{pmatrix} 0 & 0 \\ -2 \partial(b_k) & 0 \end{pmatrix}.$$

Here (62) has been used for the simplification. The lemma is proved. $\square$

Definition 2. Define $P_k = e^{\text{ad}_U}(\Lambda_{1+2k})$, $k = 0, 1, 2, 3, \dots$

Lemma 4. *The following formula holds true*

$$D_k(P_\ell) = [V_k, P_\ell], \quad \forall k, \ell \geq 0. \quad (69)$$

Proof. Applying $e^{-\text{ad}_U}$ to the both sides of (68) yields

$$[D_k + S_k, \partial + \Lambda + H] = 0,$$

where $S_k := \sum_{i \geq 0} \frac{(-1)^i}{(i+1)!} \text{ad}_U^i(D_k(U)) - e^{-\text{ad}_U}(V_k)$. Clearly, S_k belongs to $\mathcal{A}_u \otimes \mathcal{S}$. Decompose $S_k = S_k^{\text{Ker}} + S_k^{\text{Im}}$ with $S_k^{\text{Ker}} \in \mathcal{A} \otimes \text{Ker}_{\text{ad}_\Lambda}$, $S_k^{\text{Im}} \in \mathcal{A} \otimes \text{Im}_{\text{ad}_\Lambda}$. We have

$$\partial(S_k^{\text{Ker}}) = D_k(H), \quad (70)$$

$$\partial(S_k^{\text{Im}}) = [S_k^{\text{Im}}, \Lambda + H]. \quad (71)$$

Equation (71) implies that S_k^{Im} must vanish. Otherwise, write $S_k^{\text{Im}} = \sum_{j \leq m} S_k^{\text{Im}, [j]}$ for some m with $S_k^{\text{Im}, [m]}$ is not zero. Then since $\deg H < 0$, the highest (principal) degree equation of (71) reads $[S_k^{\text{Im}, [m]}, \Lambda] = 0$. This implies that $S_k^{\text{Im}, [m]} = 0$, which produces a contradiction. Therefore S_k belongs to $\mathcal{A} \otimes \text{Ker}_{\text{ad}_\Lambda}$. This implies the identity

$$[D_k + S_k, \Lambda_{1+2\ell}] = 0.$$

Applying e^{ad_U} to this identity yields (69). $\square$

Let us proceed to the tau-structure. We first verify that $\Omega_{p,q}$ are well-defined from eq. (12). Indeed,

$$R(\mu) = R(\lambda) + R'(\lambda)(\mu - \lambda) + (\mu - \lambda)^2 \partial_\lambda \left(\frac{R(\lambda) - R(\mu)}{\lambda - \mu} \right),$$

so we have

$$\text{RHS of (12)} = \frac{2\lambda}{(\lambda - \mu)^2} - \frac{\text{tr}(R(\lambda)R'(\lambda))}{\lambda - \mu} + \text{tr} \left(R(\lambda) \partial_\lambda \left(\frac{R(\lambda) - R(\mu)}{\lambda - \mu} \right) \right) - \frac{\lambda + \mu}{(\lambda - \mu)^2}.$$

It is obvious from the property $\text{tr } R(\lambda)^2 = 2\lambda$ that $\text{tr}(R(\lambda)R'(\lambda)) = 1$. Therefore $\text{RHS of (12)} = \text{tr} \left(R(\lambda) \partial_\lambda \left(\frac{R(\lambda) - R(\mu)}{\lambda - \mu} \right) \right) \in \mathcal{A}[[\lambda^{-1}, \mu^{-1}]] \lambda^{-1} \mu^{-1}$. This finishes the verification.

Introduce $\nabla(\lambda) = - \sum_{k \geq 0} \frac{(1+2k)!!}{\lambda^{1+k}} D_k$.

Lemma 5. *The following identity holds true:*

$$\nabla(\lambda)(R(\mu)) = \frac{[R(\lambda), R(\mu)]}{\lambda - \mu} - [Q(\lambda), R(\mu)], \quad Q(\lambda) := \begin{pmatrix} 0 & 0 \\ b(\lambda) & 0 \end{pmatrix}. \quad (72)$$

Proof. Using equation (69) with $\ell = 0$ we have

$$\begin{aligned} \nabla(\lambda)(R(\mu)) &= - \sum_{k \geq 0} \frac{1}{\lambda^{1+k}} \left[-(\mu^k R(\mu))_+ + \begin{pmatrix} 0 & 0 \\ b_k & 0 \end{pmatrix}, R(\mu) \right] \\ &= \frac{1}{2\pi i} \oint_{|\mu| < |\rho| < |\lambda|} d\rho \frac{[R(\rho), R(\mu)]}{(\lambda - \rho)(\rho - \mu)} - \left[\begin{pmatrix} 0 & 0 \\ b(\lambda) - 1 & 0 \end{pmatrix}, R(\mu) \right] \\ &= \frac{[R(\lambda), R(\mu)]}{\lambda - \mu} - [\text{Coef}_{\lambda^1} R(\lambda), R(\mu)] - \left[\begin{pmatrix} 0 & 0 \\ b(\lambda) - 1 & 0 \end{pmatrix}, R(\mu) \right]. \end{aligned}$$

The lemma is proved. $\square$

Proof of Proposition 1. The fact that $\Omega_{0,0} = u$ can be obtained from a straightforward residue computation. Since the RHS of (12) is invariant with respect to $\lambda \leftrightarrow \mu$, we have $\Omega_{p,q} = \Omega_{q,p}$. Applying $\nabla(\nu)$ to the both sides of (12) we obtain

$$\begin{aligned} \sum_{p,q,r \geq 0} D_r(\Omega_{p,q}) \frac{(1+2p)!!(1+2q)!!(1+2r)!!}{\lambda^{p+1} \mu^{q+1} \nu^{r+1}} &= \frac{\text{Tr}[R(\nu), R(\lambda)]R(\mu)}{(\lambda - \mu)^2(\nu - \lambda)} + \frac{\text{Tr} R(\lambda) [R(\nu), R(\mu)]}{(\lambda - \mu)^2(\nu - \mu)} \\ &\quad - \frac{\text{Tr}[Q(\nu), R(\lambda)]R(\mu)}{(\lambda - \mu)^2} - \frac{\text{Tr} R(\lambda) [Q(\nu), R(\mu)]}{(\lambda - \mu)^2} \\ &= - \frac{\text{Tr}[R(\nu), R(\lambda)]R(\mu)}{(\lambda - \mu)(\mu - \nu)(\nu - \lambda)}. \end{aligned}$$

The RHS of this identity is invariant with respect to the permutations of λ, μ, ν , so is the LHS. This implies that $D_r(\Omega_{p,q}) = D_p(\Omega_{q,r}) = D_q(\Omega_{r,p})$. For $n = 3$, the statement is already proven in the above proof of Proposition 1. Formula (13) can be proved by induction; details can be found in [13, 15] and are omitted here. The proposition is proved. $\square$

Corollary 5. *The D_k ($k \geq 0$) (defined in Definition 1) commute pairwise. Moreover, we have $Q_0 = 0$, $Q_1 = u_3/12$, and $Q_k \in \mathcal{A}^{\geq 2}$.*

Proof. We have for all $k, \ell \geq 0$,

$$D_k D_\ell(\Omega_{0;0}) = D_k D_0(\Omega_{\ell;0}) = D_0 D_k(\Omega_{\ell;0}) = D_0 D_0(\Omega_{k;\ell}) = D_\ell D_k(\Omega_{0;0}).$$

Therefore $[D_k, D_\ell] = 0$. The fact that $Q_0 = 0$ and $Q_1 = u_3/12$ can be easily seen from (64) and (65). Define $B_k = b_k|_{u_i=0, i \geq 1}$. Then $B_{-1} = 1$. Equation (63) implies that

$$B_k = \sum_{\substack{k_1, k_2 \geq -1 \\ k_1 + k_2 = k-2}} u_0 B_{k_1} B_{k_2} - \frac{1}{2} \sum_{\substack{k_1 \geq 0, k_2 \geq -1 \\ k_1 + k_2 = k-2}} B_{k_1} B_{k_2+1}, \quad k \geq 0.$$

It follows from this recursion that $B_k = \frac{(1+2k)!!}{(k+1)!} u_0^{k+1}$ for all $k \geq 0$, which implies $Q_k \in \mathcal{A}^{\geq 2}$. The corollary is proved. $\square$

Corollary 5 and the uniqueness statement proved in [58] imply that the D_k defined in Definition 1 coincide with the ones defined in Section 1.1. Our definition of tau-function (see Section 1.2) of any solution also coincides with the one in the literature [25, 26, 42], which can be seen by observing that

$$\Omega_{p,q}^{[0]} = \frac{u^{p+q+1}}{p! q! (p+q+1)},$$

where $\Omega_{p,q}^{[0]}$ denotes the degree zero part of $\Omega_{p,q}$.

3. FROM A PAIR OF WAVE FUNCTIONS TO THE n -POINT FUNCTIONS

Denote by $L = \partial^2 + 2u_0 : \mathcal{A} \rightarrow \mathcal{A}$ the Lax operator for the KdV hierarchy. Let us first briefly recall the pseudo-differential operator approach [28] to the study of the KdV equation. Notations will be the same as in Section 1.3. Denote $A_k = \frac{1}{(2k+1)!!} (L^{\frac{2k+1}{2}})_+ (k \geq 0)$. We recall that A_k is a differential operator with coefficients in $\mathcal{A}$. Define a family of derivations D_k by $D_k(u_0) := [A_k, L/2]$. It is straightforward to show (by using the properties of pseudo-differential operators) that these D_k commute pairwise with $D_0 = \partial$, $D_1(u_0) = u_0 u_1 + u_3/12$ and $D_k(u_0) = u_0^k u_1/k! + \dots$, where the dots denotes terms in $\mathcal{A}^{\geq 2}$. So these D_k coincide with the ones defined in Section 1.1, and the KdV hierarchy can be written equivalently as

$$\frac{\partial L}{\partial t_k} = [A_k, L] \quad (73)$$

with ∂ replaced by ∂_x and u_i by $\partial_x^i(u)$. In the rest of this section, we give a way of constructing a pair of wave functions of an arbitrary solution, and apply it as well as Corollary 1 to prove Theorem 1.

3.1. Dressing operators and pairs of wave functions. Similarly as in Section 1.3, we start with the *time-independent* case. Namely, consider $L = \partial^2 + 2f(x)$ with arbitrarily given fixed $f(x) \in V$. Here we recall that V is a ring of functions of x closed under ∂_x . A pseudo-differential operator Φ of the form $\Phi = \sum_{m \geq 0} \phi_m(x) \partial_x^{-m}$ with $\phi_0(x) \equiv 1$ is called a *dressing operator* of L if

$$\Phi(x) \circ \partial_x^2 \circ \Phi(x)^{-1} = L.$$

Here $\phi_m(x)$ ($m \geq 1$) will live in a certain ring $\tilde{V}$ satisfying $V \subseteq \partial_x(\tilde{V}) \subseteq \tilde{V}$.

If $\Phi(x)$ exists then the freedom of $\Phi(x)$ is given by an operator of the form $\sum_{m \geq 0} g_m \partial_x^{-m}$ for arbitrary constants g_m with $g_0 = 1$ through the right composition

$$\Phi(x) \mapsto \Phi(x) \circ \sum_{m \geq 0} g_m \partial_x^{-m}. \quad (74)$$

Recall that for pseudo-differential operators [28], most operations (e.g., composition) only used differentiations in x , namely, for $a \in \mathbb{C} \setminus \mathbb{N}$ we usually think of ∂_x^a as a formal symbol and do not actually need to consider how it acts on a function of x . But, following [28], for all $p(x) \in \tilde{V}$ we can define $\partial_x^{-m}(p(x) e^{xz})$ for any $m \in \mathbb{C}$ as $\sum_{l \geq 0} \binom{-m}{l} \partial_x^l(p(x)) z^{-m-l} e^{xz}$. If $\Phi = \Phi(x) = \sum_{m \geq 0} \phi_m(x) \partial_x^{-m}$ is a dressing operator of L , then it is straightforward to check that $\psi := \Phi(x)(e^{xz})$ is in $\tilde{V}((z^{-1})) e^{xz}$ and is a wave function of f (cf. Section 1.3). Explicitly, $\psi = \sum_{m \geq 0} \phi_m(x) z^{-m} e^{xz}$. On the contrary, if $\psi = \sum_{m \geq 0} \phi_m(x) z^{-m} e^{xz}$ is a wave function of f , then $\Phi := \sum_{m \geq 0} \phi_m(x) \partial_x^{-m}$ is a dressing operator of L . This gives the 1-1 correspondence between the dressing operators of L and the wave functions of f . Also, the freedom of Φ generated by the right composition (74) matches to $\psi \mapsto g(z)\psi$ with $g(z) = \sum_{m \geq 0} g_m z^{-m}$.

Let us give a proof of the existence of a wave function. Write

$$\psi(z, x) = e^{\beta(z, x)} e^{xz}, \quad \beta(z, x) := \sum_{j \geq 1} \beta_j(x) z^{-j}. \quad (75)$$

The defining equation $L(\psi) = z^2 \psi$ for the wave function ψ written in terms of β is given by

$$\beta_{xx} + \beta_x^2 + 2z\beta_x + 2f = 0. \quad (76)$$

Substituting the expression (75) of β in this equation we obtain

$$\beta'_j(x) = -\frac{1}{2} \beta''_{j-1}(x) - \frac{1}{2} \sum_{\substack{j_1, j_2 \geq 1 \\ j_1 + j_2 = j-1}} \beta'_{j_1}(x) \beta'_{j_2}(x) - f \delta_{j,1}, \quad (77)$$

where $\beta_0(x)$ is defined to as the 0-function and $' = d/dx$. Clearly, the recursion (77) uniquely determines $\beta'_j(x)$ for all $j \geq 1$ and $\beta'_j(x) \in V$. Since $V \subseteq \partial_x(\tilde{V})$, we conclude that the solution ψ of the form (75) to the equation $L(\psi) = z^2 \psi$ exists with $\beta_j(x) \in \tilde{V}$. Expanding the exponential $e^{\beta(z, x)}$ into Taylor series of z^{-1} we obtain finally the required form $\psi = \sum_{m \geq 0} \phi_m(x) z^{-m} e^{xz}$ with $\phi_0(x) \equiv 1$ and $\phi_m(x) \in \tilde{V}$ for a wave function.

Now let ψ be a wave function of f and Φ the (unique) dressing operator of L associated to ψ . Define a particular element ψ^* in $\tilde{V}((z^{-1})) e^{-xz}$ by

$$\psi^* = \psi^*(z, x) := (\Phi^{-1})^*(e^{-xz}), \quad (78)$$

where $(\Phi^{-1})^*$ denotes the formal adjoint operator of Φ^{-1} . For any $y_1 = a_1 e^{xz} \in \tilde{V}((z^{-1})) e^{xz}$ and $\tilde{y}_1 = b_1 e^{xz} \in \tilde{V}((z^{-1})) e^{-xz}$, define $y_1 \tilde{y}_1 := a_1 b_1 \in \tilde{V}((z^{-1}))$.

Lemma 6. *The elements ψ and ψ^* form a pair of wave functions of f .*

Proof. It is easy to check that ψ^* is indeed of the form $\psi^* = (1 + \phi_1^*(x)/z + \phi_2^*(x)/z^2 + \dots) e^{-xz}$ with $\phi^*(x) \in \tilde{V}$ and satisfies $L(\psi^*) = z^2 \psi^*$. We are left to show that for all $i \geq 0$,

$$\text{res}_{z=\infty} \partial_x^i (\psi(z, x)) \psi^*(z, x) dz = 0.$$

Note that for any two pseudo-differential operators P, Q we have

$$\text{res}_{z=\infty} P(e^{xz}) Q(e^{-xz}) dz = \text{res}_{\partial_x} P \circ Q^*. \quad (79)$$

Taking $P = \partial_x^i \circ \Phi$ and $Q = \Phi^{-1}$ in this identity we find that

$$\text{res}_{z=\infty} \partial_x^i (\psi(z, x)) \psi^*(z, x) dz = \text{res}_{z=\infty} \partial_x^i \circ \Phi(e^{xz}) (\Phi^{-1})^*(e^{-xz}) dz = \text{res}_{\partial_x} \partial_x^i \circ \Phi \circ \Phi^{-1} = 0.$$

The lemma is proved. $\square$

Hence we have proved the existence of a pair of wave functions of f .

Observe that the above proof is revertible. Namely, the zero-residue condition ensures the uniqueness of the dual wave function associated with ψ . It must be the ψ^* defined by (78).

Lemma 7. *We have $\psi_x \psi^* - \psi \psi_x^* = 2z$ and $\psi(z, x) \psi^*(z, x) = b(z^2, x)$, with b as in (56).*

Proof. Denote $W = \psi_x \psi^* - \psi \psi_x^*$. We have $W_x = 0$. It follows that W must have the form $W = 2z + \sum_{k \geq 0} s_k z^{-k}$, where s_k are constants. For any $i \geq -1$,

$$\begin{aligned} \text{res}_{z=\infty} z^i W dz &= \text{res}_{z=\infty} \left(\partial_x \circ \Phi \circ \partial_x^i (e^{xz}) (\Phi^{-1})^* (e^{-xz}) - (-1)^i \Phi(e^{xz}) \partial_x \circ (\Phi^{-1})^* \circ \partial_x^i (e^{-xz}) \right) dz \\ &= \text{res}_{\partial_x} \left(\partial_x \circ \Phi \circ \partial_x^i \circ \Phi^{-1} + \Phi \circ \partial_x^i \circ \Phi^{-1} \circ \partial_x \right) = \text{res}_{\partial_x} \left(\partial_x \circ L^{\frac{i}{2}} + L^{\frac{i}{2}} \circ \partial_x \right) = 0. \end{aligned}$$

The last equality uses the fact that L is a differential operator and that $(L^*)^{1/2} = -L^{1/2}$. Therefore all s_k vanish. The first identity is proved. To prove the second equality we observe that both the LHS and the RHS satisfies the same differential equation (60) (with u_0 replaced by f and ∂ by ∂_x). Moreover, they are both formal power series in z^{-1} with the leading term 1. Then the uniqueness statement implies the second identity. The lemma is proved. $\square$

We note that the identity $\psi(z, x) \psi^*(z, x) = b(z^2, x)$ can be used as an alternative and possibly useful criterion for a pair of wave functions. We should also notice that for an arbitrary solution $u(\mathbf{t})$ there is a particular subclass of pairs of wave functions of $u(\mathbf{t})$ with the additional property $\psi(-z, x) = \psi^*(z, x)$, which was often assumed in the literature.

Let us now proceed to the *time-dependent* case, i.e. to prove the existence of a pair of wave functions of an arbitrary solution of the KdV hierarchy in the ring $V[[\mathbf{t}_{>0}]]$. We recall that for the operators $L = \partial^2 + 2u_0$ and A_k , once a solution $u = u(\mathbf{t})$ is taken, we will replace ∂ by ∂_x and u_i by $\partial_x^i(u)$.

Proposition 4. *Let $u = u(\mathbf{t})$ be a solution to the KdV hierarchy (2) in the ring $V[[\mathbf{t}_{>0}]]$. There exists a pseudo-differential operator Φ of the form*

$$\Phi = \Phi(\mathbf{t}) = \sum_{m \geq 0} \phi_m(\mathbf{t}) \partial_x^{-m}, \quad \phi_0(\mathbf{t}) \equiv 1 \quad (80)$$

such that

$$L = \Phi \circ \partial_x^2 \circ \Phi^{-1}, \quad (81)$$

$$\partial_{t_k}(\Phi) = -\frac{1}{(2k+1)!!} (\Phi \circ \partial_x^{2k+1} \circ \Phi^{-1})_- \circ \Phi, \quad \forall k \geq 0. \quad (82)$$

Here, $t_0 = x$ as usual and $\phi_m(\mathbf{t}) \in \tilde{V}[[\mathbf{t}_{>0}]]$ ($m \geq 1$).

Proof. Let $f(x) := u(x, \mathbf{0})$ be the initial data of the solution. According to the above time-independent theory, we can take a pseudo-differential operator $\Phi(x)$ of the form $\Phi(x) = \sum_{m=0}^{\infty} \phi_m(x) \partial_x^{-m}$ with $\phi_0(x) \equiv 1$ and $\phi_m(x) \in \tilde{V}$ ($m \geq 1$), such that $\partial_x^2 + 2f(x) = \Phi(x) \circ \partial_x^2 \circ \Phi(x)^{-1}$.

Now consider the initial value problem for $\Phi(\mathbf{t})$ given by equations (81) and (82) with the initial data $\Phi(x, \mathbf{0}) = \Phi(x)$. First let us check the compatibility between (81) and the $k = 0$ case of equations (82). We notice that $(\Phi \circ \partial_x \circ \Phi^{-1})_- = (L^{1/2})_- = L^{1/2} - (L^{1/2})_+ = \Phi \circ \partial_x \circ \Phi^{-1} - \partial_x$. Therefore,

$$\partial_{t_0}(\Phi) = -\Phi \circ \partial_x + \partial_x \circ \Phi = \partial_x(\Phi).$$

This is compatible with our convention $t_0 = x$. Secondly, we will prove the compatibility between (81) and equations (82) with $k \geq 1$. Indeed, we have

$$\begin{aligned} \frac{\partial L}{\partial t_k} &= \partial_{t_k}(\Phi) \circ \partial_x^2 \circ \Phi^{-1} - \Phi \circ \partial_x^2 \circ \Phi^{-1} \circ \partial_{t_k}(\Phi) \circ \Phi^{-1} \\ &= -\frac{1}{(2k+1)!!} \left(L^{\frac{2k+1}{2}} \right)_- \circ L + \frac{1}{(2k+1)!!} L \circ \left(L^{\frac{2k+1}{2}} \right)_- = [A_k, L]. \end{aligned}$$

This is true as u is a solution of the KdV hierarchy (cf. (73)). Finally, we check the compatibility between all the equations of (82). Indeed,

$$\begin{aligned} &\partial_{t_\ell} \partial_{t_k}(\Phi) - \partial_{t_k} \partial_{t_\ell}(\Phi) \\ &= \frac{\left(\left[\left(L^{\frac{2\ell+1}{2}} \right)_-, L^{\frac{2k+1}{2}} \right] - \left[\left(L^{\frac{2k+1}{2}} \right)_-, L^{\frac{2\ell+1}{2}} \right] + \left[\left(L^{\frac{2k+1}{2}} \right)_-, \left(L^{\frac{2\ell+1}{2}} \right)_- \right] \right)}{(2k+1)!! (2\ell+1)!!} \circ \Phi \\ &= \frac{\left(\left[\left(L^{\frac{2\ell+1}{2}} \right)_-, L^{\frac{2k+1}{2}} \right] - \left[L^{\frac{2k+1}{2}}, \left(L^{\frac{2\ell+1}{2}} \right)_+ \right] \right)}{(2k+1)!! (2\ell+1)!!} \circ \Phi = \frac{\left(\left[L^{\frac{2\ell+1}{2}}, L^{\frac{2k+1}{2}} \right] \right)_-}{(2k+1)!! (2\ell+1)!!} \circ \Phi = 0. \end{aligned}$$

The proposition is proved. $\square$

We call $\Phi(\mathbf{t})$ in Proposition 4 a *dressing operator* of $L = \partial_x^2 + 2u(\mathbf{t})$. It is clear from the proof of Proposition 4 that the freedom of the dressing operator $\Phi(\mathbf{t})$ is characterized by that of $\Phi(x)$, i.e., by a sequence of arbitrary constants $g_1, g_2, \dots$ through

$$\Phi \mapsto \Phi \circ \sum_{m=0}^{\infty} g_m \partial_x^{-m}, \quad g_0 = 1.$$

Similarly as in the time-independent case we will use the dressing operator to prove the existence of a pair of wave functions of u . As in (80), write $\Phi(\mathbf{t}) = \sum_{m \geq 0} \phi_m(\mathbf{t}) \partial_x^{-m}$ with $\phi_0(x) = 1$. Put

$Q = e^{\sum_{k \geq 0} \frac{z^{2k+1}}{(2k+1)!!} t_k}$. Define for any $p(\mathbf{t}) \in \tilde{V}[[\mathbf{t}_{>0}]]$ and for all $m \in \mathbb{Z}$, $\partial_x^{-m} \left(p(\mathbf{t}) e^{\sum_{k \geq 0} \frac{z^{2k+1}}{(2k+1)!!} t_k} \right) := \sum_{l \geq 0} \binom{-m}{l} \partial_x^l (p(\mathbf{t})) z^{-m-l} e^{\sum_{k \geq 0} \frac{z^{2k+1}}{(2k+1)!!} t_k}$. Then one can check that the following element

$$\psi(z, \mathbf{t}) := \Phi(\mathbf{t})(Q) = \left(1 + \frac{\phi_1(\mathbf{t})}{z} + \frac{\phi_2(\mathbf{t})}{z^2} + \dots \right) e^{\sum_{k \geq 0} \frac{z^{2k+1}}{(2k+1)!!} t_k} \quad (83)$$

is a wave function of u . For any $p(\mathbf{t}) \in \tilde{V}[[\mathbf{t}_{>0}]]$ and $m \in \mathbb{Z}$, define $\partial_x^{-m} \left(p(\mathbf{t}) e^{-\sum_{k \geq 0} \frac{z^{2k+1}}{(2k+1)!!} t_k} \right) := \sum_{l \geq 0} \binom{-m}{l} \partial_x^l (p(\mathbf{t})) (-z)^{-m-l} e^{-\sum_{k \geq 0} \frac{z^{2k+1}}{(2k+1)!!} t_k}$. For any $y_1 = a_1 e^{\sum_{k \geq 0} \frac{z^{2k+1}}{(2k+1)!!} t_k}$, $\tilde{y}_1 = \tilde{a}_1 e^{-\sum_{k \geq 0} \frac{z^{2k+1}}{(2k+1)!!} t_k}$ with a_1 and $\tilde{a}_1$ in $\tilde{V}[[\mathbf{t}_{>0}]]((z^{-1}))$, we define $y_1 \tilde{y}_1 := a_1 \tilde{b}_1 \in \tilde{V}[[\mathbf{t}_{>0}]]((z^{-1}))$. One can then verify that the following element

$$\psi^*(z, \mathbf{t}) := (\Phi(\mathbf{t})^{-1})^*(Q^{-1}) = \left(1 + \frac{\phi_1^*(\mathbf{t})}{z} + \frac{\phi_2^*(\mathbf{t})}{z^2} + \dots \right) e^{-\sum_{k \geq 0} \frac{z^{2k+1}}{(2k+1)!!} t_k} \quad (84)$$

is the dual wave function of u associated with ψ . Hence we have proved the existence of a pair of wave functions of u .

3.2. Proof of Theorem 1. Let $u = u(\mathbf{t})$ be an arbitrary solution to the KdV hierarchy (2) and (ψ, ψ^*) a pair of wave functions of u . We first prove a useful lemma.

Lemma 8. *Define*

$$\Psi(z, \mathbf{t}) = \begin{pmatrix} \psi(z, \mathbf{t}) & \psi^*(z, \mathbf{t}) \\ -\psi_x(z, \mathbf{t}) & -\psi_x^*(z, \mathbf{t}) \end{pmatrix}. \quad (85)$$

Then we have

$$\det \Psi(z, \mathbf{t}) \equiv 2z \quad (86)$$

and

$$b(\lambda, \mathbf{t}) = \psi(z, \mathbf{t}) \psi^*(z, \mathbf{t}), \quad (87)$$

where $\lambda = z^2$. Moreover, the matrix resolvent $R(\lambda, \mathbf{t})$ of u is given by

$$R(\lambda, \mathbf{t}) \equiv -\Psi(z, \mathbf{t}) \begin{pmatrix} z & 0 \\ 0 & -z \end{pmatrix} \Psi^{-1}(z, \mathbf{t}). \quad (88)$$

Proof. The proof for (86) and (87) is almost identical with that for Lemma 7, so we omit the details. Let us now prove (88). We have

$$-\Psi(z, \mathbf{t}) \begin{pmatrix} z & 0 \\ 0 & -z \end{pmatrix} \Psi^{-1}(z, \mathbf{t}) = \begin{pmatrix} \frac{1}{2}(\psi\psi^*)_x & \psi\psi^* \\ -\psi_x\psi_x^* & -\frac{1}{2}(\psi\psi^*)_x \end{pmatrix}. \quad (89)$$

So (87) shows the $(1, 2)$ -entry identity of (88). The $(1, 1)$ -entry identity and $(2, 2)$ -entry identity are also true due to (57). It remains to show

$$-\psi_x\psi_x^* = (\lambda - 2u)b - \frac{1}{2}\partial_x^2(b).$$

Indeed,

$$\text{RHS} = (\lambda - 2u)\psi\psi^* - \frac{1}{2}(\psi_{xx}\psi + 2\psi_x\psi_x^* + \psi\psi_{xx}^*) = \text{LHS}.$$

The lemma is proved. $\square$

According to equation (88) we can write the basic matrix resolvent $R(z^2, \mathbf{t})$ of u in terms of ψ, ψ^* as

$$R(z^2, \mathbf{t}) = z + \begin{pmatrix} \psi \\ -\psi_x \end{pmatrix} \begin{pmatrix} \psi_x^* & \psi^* \end{pmatrix}. \quad (90)$$

For simplicity, we will often denote $R(\lambda) = R(\lambda, \mathbf{t})$. Through a direct calculation we have

$$\text{tr } R(z_1^2) R(z_2^2) = (\psi_x^*(z_1)\psi(z_2) - \psi^*(z_1)\psi_x(z_2))(\psi_x^*(z_2)\psi(z_1) - \psi^*(z_2)\psi_x(z_1)) - 2z_1z_2. \quad (91)$$

Hence

$$\sum_{p_1, p_2} \Omega_{p_1, p_2}(\mathbf{t}) \frac{(2p_1 + 1)!! (2p_2 + 1)!!}{z_1^{2p_1+2} z_2^{2p_2+2}} = -D(z_1, z_2; \mathbf{t}) D(z_2, z_1; \mathbf{t}) - \frac{1}{(z_1 - z_2)^2}. \quad (92)$$

This shows formula (19) for $n = 2$. For $n \geq 3$, we are going to use a formula given in [37]:

$$\begin{aligned} & \sum_{\sigma \in S_n / C_n} \frac{\text{tr } R(\lambda_{\sigma(1)}) \cdots R(\lambda_{\sigma(n)})}{\prod_{i=1}^n (\lambda_{\sigma(i+1)} - \lambda_{\sigma(i)})} \\ &= \sum_{\sigma \in S_{n-2}} \frac{\langle R(\lambda_n), \text{ad}_{R(\lambda_{\sigma(1)})} \cdots \text{ad}_{R(\lambda_{\sigma(n-2)})} R(\lambda_{n-1}) \rangle}{(\lambda_{n-1} - \lambda_{\sigma(n-2)})(\lambda_n - \lambda_{n-1})(\lambda_{\sigma(1)} - \lambda_n) \prod_{i=1}^{n-3} (\lambda_{\sigma(i+1)} - \lambda_{\sigma(i)})}, \end{aligned}$$

where $\text{ad}_a b := [a, b]$ and $\langle a, b \rangle := \text{tr } ab$. Here we have abbreviated $R_i(\lambda) = R_i(\lambda, \mathbf{t})$. This formula tells that the contributions to the n -point generating series coming from the term z in (90) are zero. Therefore the theorem is proved by using the same argument as in [40]. $\square$

3.3. Proof of Theorem 2. We will first prove Proposition 2. It follows easily from Lemma 8 that the four functions $\psi, \psi^*, \psi_x, \psi_x^*$ satisfy the following three relations:

$$\begin{aligned}\psi(z, \mathbf{t}) \psi^*(z, \mathbf{t}) &= b(z^2, \mathbf{t}), \\ \psi_x(z, \mathbf{t}) \psi^*(z, \mathbf{t}) - \psi(z, \mathbf{t}) \psi_x^*(z, \mathbf{t}) &= 2z, \\ \psi_x(z, \mathbf{t}) \psi^*(z, \mathbf{t}) + \psi(z, \mathbf{t}) \psi_x^*(z, \mathbf{t}) &= b_x(z^2, \mathbf{t}).\end{aligned}$$

Solving this system we obtain

$$\psi^*(z, \mathbf{t}) = \frac{b(z^2, \mathbf{t})}{\psi(z, \mathbf{t})}, \quad \psi_x(z, \mathbf{t}) = \psi(z, \mathbf{t}) \frac{b_x(z^2, \mathbf{t}) + 2z}{2b(z^2, \mathbf{t})}, \quad \psi_x^*(z, \mathbf{t}) = \frac{b_x(z^2, \mathbf{t}) - 2z}{2\psi(z, \mathbf{t})}.$$

Substituting these expressions into (18), we obtain the first equality of (23). Proposition 2 is then proved. Theorem 2 follows easily from Theorem 1 and the first equality of (23), as the factors of the form $\psi(z, \mathbf{t})/\psi(w, \mathbf{t})$ cancel in each product of the sum of the right hand side of (19). $\square$

As the remark given in the introduction (right after Theorem 2), the abstract version of Theorem 2 follows immediately. We note that it is not difficult to give a direct proof of the abstract version of Theorem 2 by using the definition of K (cf. equation (20)), Lemma 5 and Proposition 1; we leave it as an exercise for interested readers.

4. GENERATING SERIES OF THE GENERALIZED BGW CORRELATORS

Denote by $\pi : \overline{\mathcal{M}}_{g,n+1} \rightarrow \overline{\mathcal{M}}_{g,n}$ the forgetful map forgetting the last marked point, and $\rho : \overline{\mathcal{M}}_{g-1,n+2} \rightarrow \overline{\mathcal{M}}_{g,n}$ and $\phi_{h,I} : \overline{\mathcal{M}}_{h,|I|+1} \times \overline{\mathcal{M}}_{g-h,|J|+1} \rightarrow \overline{\mathcal{M}}_{g,n}$, $I \sqcup J = \{1, \dots, n\}$ the gluing maps. Norbury [62] introduced a collection of cohomology classes $\{\Theta_{g,n} \in H^*(\overline{\mathcal{M}}_{g,n})\}_{2g-2+n>0}$ satisfying

$$\text{i) } \Theta_{g,n} \text{ is of pure degree,} \quad (93)$$

$$\text{ii) } \rho^* \Theta_{g,n} = \Theta_{g-1,n+2}, \quad \phi_{h,I}^* \Theta_{g,n} = \pi_1^* \Theta_{h,|I|+1} \cdot \pi_2^* \Theta_{g-h,|J|+1}, \quad (94)$$

$$\text{iii) } \Theta_{g,n+1} = \psi_{n+1} \cdot \pi^* \Theta_{g,n}, \quad (95)$$

$$\text{iv) } \Theta_{1,1} = 3\psi_1. \quad (96)$$

Norbury proved that such $\Theta_{g,n}$ exists and it must satisfy $\Theta_{g,n} \in H^{4g-4+2n}(\overline{\mathcal{M}}_{g,n})$. Define $Z_\Theta = Z_\Theta(\mathbf{t})$ as the following generating series of intersection numbers (called the partition function)

$$Z_\Theta(\mathbf{t}) = \exp \left(\sum_{g,n \geq 0} \frac{1}{n!} \sum_{p_1, \dots, p_n \geq 0} \int_{\overline{\mathcal{M}}_{g,n}} \Theta_{g,n} \psi_1^{p_1} \dots \psi_n^{p_n} t_{p_1} \dots t_{p_n} \right). \quad (97)$$

Define $u_\Theta = u_\Theta(\mathbf{t}) := \frac{\partial^2 \log Z_\Theta(\mathbf{t})}{\partial t_0^2}$. The integrals $\int_{\overline{\mathcal{M}}_{g,n}} \Theta_{g,n} \psi_1^{p_1} \dots \psi_n^{p_n}$, called the n -point Θ -class intersection numbers (aka the n -point Θ -class correlators), are independent of choice of $\Theta_{g,n}$. They vanish unless $p_1 + \dots + p_n = g - 1$. Therefore u_Θ belongs to $\mathbb{Q}[[t_0, t_1, t_2, \dots]]$.

Norbury's Theorem ([62]). *The formal power series u_Θ is a solution of the KdV hierarchy (2) with*

$$u_\Theta(t_0 = x, \mathbf{0}) = \frac{1}{8(x-1)^2}. \quad (98)$$

Moreover, Z_Θ is the tau-function of the solution u_Θ satisfying the string type equation

$$\sum_{i \geq 0} (1 + 2i) t_i \frac{\partial Z_\Theta}{\partial t_i} + \frac{1}{8} Z_\Theta = \frac{\partial Z_\Theta}{\partial t_0}. \quad (99)$$

Norbury's Theorem tells that $u_\Theta = u_{\Theta(1/8)}$. The latter is defined in Example 2 of Introduction. So $\tau_{\Theta(1/8)}$ and Z_Θ can only differ by the exponential of a linear function. Noting that equation (99) coincides with equation (38) with $C = 1/8$, so the linear function can only be a constant, which can easily be normalized as zero. We conclude that $Z_\Theta = \tau_{\Theta(1/8)}$. The latter is often called the BGW tau-function [20, 48, 2, 30, 61], originally studied in matrix models. The goal of this section is to prove Theorems 3, 4, 5, and give some explicit computations.

4.1. The essential second kind topological ODE of A_1 -type. The topological ODE of $\mathfrak{g}$ -type with $\mathfrak{g}$ being a simple Lie algebra was introduced and studied in [14, 15], which will be called the first kind topological ODE. To prove Theorem 4, let us introduce an ODE associated to $\mathfrak{sl}_2(\mathbb{C})$

$$2\zeta^3 \rho''' + 3\zeta^2 \rho'' - 2\zeta(\zeta - 2C)\rho' - 2C\rho = 0, \quad ' := \frac{d}{d\zeta}, \quad (100)$$

where C is an arbitrary parameter. It will be used for computing the Θ -class intersection numbers in full genera ($C = 1/8$) and the generalized BGW correlators. As one motivation, note that the matrix $R = R(\lambda, x)$ appearing in Theorem 4 is uniquely determined by the given $f(x) = \frac{C}{(1-x)^2}$, and equation (100) is nothing but a kind of transformation of the matrix resolvent recursive relation (see Section 2.2) for the corresponding $b = b(\lambda, x)$, which has two independent variables, into a recursive relation for the coefficients of the one-variable function ρ . We call (100) the *essential second kind topological ODE of A_1 -type*. See Section 6.1 for more details.

Proposition 5. *For any fixed $C \in \mathbb{C}$, there exists a unique series ρ in $\mathbb{C}[[z^{-1}]]$ satisfying equation (100) as well as the initial condition*

$$\rho(\infty) = 1. \quad (101)$$

Moreover, $\rho \in \mathbb{Q}[C][[\zeta^{-1}]]$ and it satisfies the following nonlinear ODE

$$\zeta \rho'^2 + \left(1 - \frac{2C}{\zeta}\right) \rho^2 - \rho \rho' - 2\zeta \rho \rho'' = 1. \quad (102)$$

Proof. Write

$$\rho = \sum_{k \geq -1} (2k+1)!! \frac{\rho_k}{\zeta^{k+1}} \quad (103)$$

with $\rho_{-1} = 1$. Substituting this expression in (100) we obtain

$$\rho_k = \frac{C + \frac{k(k+1)}{2}}{k+1} \rho_{k-1}, \quad k \geq 0. \quad (104)$$

This implies the uniqueness statement of the proposition and $\rho \in \mathbb{Q}[C][[\zeta^{-1}]]$. To show (102), note that

$$\frac{d}{d\zeta} \left(\zeta \rho'^2 + \left(1 - 2C/\zeta\right) \rho^2 - \rho \rho' - 2\zeta \rho \rho'' \right) = 0.$$

Therefore, $\zeta \rho'^2 + \left(1 - 2C/\zeta\right) \rho^2 - \rho \rho' - 2\zeta \rho \rho'' \equiv C_1$, where C_1 is a constant independent of ζ . The fact that $C_1 \equiv 1$ can be deduced from (103). The proposition is proved. $\square$

We now apply Proposition 5 to derive the explicit expression of the basic matrix resolvent of the solution of the KdV hierarchy characterized by the initial data as in Example 2.

Proof of Theorem 4. Let $\rho = \rho(\zeta)$ be the unique element in Proposition 5. According to Proposition 5, $\rho = \rho(\zeta)$ satisfies equations (100)–(102). Define $\tilde{b}(\lambda, x) = \rho(\lambda(x-1)^2)$. Then it is easy to check that $\tilde{b}(\lambda, x)$ satisfies (59) and (60) with u replaced by $\frac{C}{(x-1)^2}$ and ∂ by ∂_x , and it has the form (61). Hence

the uniqueness statement in the definition of the basic matrix resolvent (see Section 2.2) implies $b(\lambda, x) = \tilde{b}(\lambda, x)$. The above equation (104) yields an explicit expression for ρ :

$$\rho = 1 + \sum_{k \geq 0} \frac{(2k+1)!!}{(k+1)!} \frac{\prod_{i=0}^k (C + \frac{i(i+1)}{2})}{\zeta^{k+1}} = {}_3F_0\left(\frac{1}{2}, \frac{1}{2} + \alpha, \frac{1}{2} - \alpha; ; \frac{1}{\zeta}\right). \quad (105)$$

(Recall that $2\alpha = \sqrt{1-8C}$.) This gives the $(1, 2)$ -entry of R . Other entries can be obtained by using (57) and (58). $\square$

As already pointed out after Theorem 4, if $-C$ is a triangular number, i.e., $C = -\frac{p(p+1)}{2}$ for some $p \in \mathbb{N}$, then the formal series $\rho = \rho(\zeta)$ truncates to the polynomial $\rho = {}_3F_0(\frac{1}{2}, 1+p, -p; ; \frac{1}{\zeta})$ in ζ^{-1} . This corresponds to the rational limit of the p -soliton solution to the KdV hierarchy.

For $n \geq 2$, using Corollary 1 one immediately obtains an explicit formula of the generating series of the n -point generalized BGW correlators (any C) in terms of the matrix R given in Theorem 4.

Proof of Theorem 3. Following from Norbury's Theorem, Corollary 1, Theorem 4 with $C = 1/8$. $\square$

Proof of Corollary 4. Note that equation (38) implies

$$(1+2p)\Omega_p^{\Theta(C)}(0) = \Omega_{0,p}^{\Theta(C)}(0), \quad p \geq 0.$$

Formula (41) is then a consequence of (105) (as it is an easy exercise that $\Omega_{0,k} = b_k/(2k+1)!!$). $\square$

Proof of Corollary 3. According to Norbury's Theorem, formula (37) is a special case of (41). $\square$

4.2. Proof of Theorem 5 and Proposition 3. It is straightforward to check that the function $W := \sqrt{2\xi/\pi} e^z K_\alpha(\xi)$ with $\xi = z(1-x)$ satisfies the differential equation $L(W) = z^2 W$, where $L = \partial_x^2 + 2C/(1-x)^2$. Moreover, as $z \rightarrow \infty$ within an appropriate sector, the asymptotic behavior of W coincides with ψ defined in (42). Similarly, the analytic function $W^* := \sqrt{2\pi\xi} e^{-z} I_\alpha(\xi)$ satisfies $L(W^*) = z^2 W^*$ and has asymptotics coinciding with ψ^* . It remains to show the zero-residue condition. This can be proved by noticing that

$$\begin{aligned} \psi(z, x) \psi^*(z, x) &= \sum_{k \geq 0} \frac{a_k(\alpha)}{z^k (1-x)^k} \sum_{\ell \geq 0} (-1)^\ell \frac{a_\ell(\alpha)}{z^\ell (1-x)^\ell} \\ &= {}_3F_0\left(\frac{1}{2}, \frac{1}{2} + \alpha, \frac{1}{2} - \alpha; ; \frac{1}{z^2(1-x)^2}\right) = b(z^2, x). \end{aligned} \quad (106)$$

This proves that ψ and ψ^* are a pair of wave functions as claimed, and equation (43) then follows from Theorem 1 with $\mathbf{t} = 0$. This completes the proof of Theorem 5.

For Proposition 3 we first observe that the recursion and boundary conditions for $A_{mn}(\alpha)$ given in (45) are obviously equivalent to either of the closed formulas in (46), so it suffices to prove the former. For convenience we write simply a_k and A_{mn} for $a_k(\alpha)$ and $A_{mn}(\alpha)$ and set

$$A(z) := \psi(z, 0) = \sum_{k=0}^{\infty} \frac{a_k}{z^k}, \quad B(z) := \psi_x(z, 0) = z + \sum_{k=0}^{\infty} \frac{b_k}{z^k}$$

with $b_k = a_{k+1} + ka_k$. (Avoid from confusion with the notations a_k, b_k in other sections.) One checks easily that $(A(z), B(z))$ satisfies the first-order differential system

$$\theta_z \begin{pmatrix} A(z) \\ B(z) \end{pmatrix} = \begin{pmatrix} z & -1 \\ 2C - z^2 & 1 + z \end{pmatrix} \begin{pmatrix} A(z) \\ B(z) \end{pmatrix}, \quad (107)$$

where $2C = \frac{1}{4} - \alpha^2$ and $\theta_z := z \frac{d}{dz}$. (This implies a second-order differential equation for A equivalent to the equation $LW = z^2 W$ given above.) Setting $y = -w$, we find the two generating functions

$$\sum_{m,n \geq 0} \frac{(m+n)(A_{m+1,n} - A_{m,n+1})}{z^m y^n} = (\theta_z + \theta_y) \left(\frac{A(z)B(y) - B(z)A(y)}{z+y} \right)$$

(this follows from the definition of the A_{mn} and the fact that the Euler operator $\theta_z + \theta_y$ annihilates the degree zero function $\frac{z-y}{z+y}$) and

$$\sum_{m,n \geq 0} \frac{(n-m)a_m a_n}{z^m y^n} = (\theta_z - \theta_y)(A(z)A(y)).$$

Using the differential equations (107) we find that the right hand sides of both of these formulas equal $(z-y)A(z)A(y) + A(z)B(y) - B(z)A(y)$, proving equation (45). For (47), we notice that the first of equations (46) shows that the polynomial $A_{mn}(\alpha)$ is divisible by $a_m(\alpha)$ (because each a_r with $r \geq m$ is divisible by a_m), and since $a_n(\alpha)$ has only simple zeros, this shows that the quotient $\tilde{A}_{mn} := A_{mn}/a_m a_n$ has only simple poles at half-integral values of α . Since $\tilde{A}_{mn}$ is also small at infinity, it has a partial fraction decomposition as $\sum_k c_{mn}(k)/(\alpha - k - \frac{1}{2})$ for some coefficients $c_{mn}(k)$. That these coefficients have the values given in (47) can be proved by comparing residues, using either the recursion (45) or one of the closed formulas (46), together with a simple binomial coefficient identity. The details are left to the reader. $\square$

4.3. Computations.

4.3.1. *Some Θ -class intersection numbers.* According to Norbury's Theorem, we have

$$\Omega_{p_1, \dots, p_n}^{\Theta(1/8)}(0) = \sum_{g=0}^{\infty} \int_{\mathcal{M}_{g,n}} \Theta_{g,n} \psi_1^{p_1} \dots \psi_n^{p_n}.$$

Here $n \geq 1$. Note that the degree-dimension matching reads

$$p_1 + \dots + p_n + 2g - 2 + n = 3g - 3 + n \quad \Leftrightarrow \quad p_1 + \dots + p_n = g - 1.$$

So, actually, $\Omega_{p_1, \dots, p_n}^{\Theta(1/8)}(0) = \int_{\overline{\mathcal{M}}_{1+p_1+\dots+p_n,n}} \Theta_{1+p_1+\dots+p_n,n} \psi_1^{p_1} \dots \psi_n^{p_n}$ as we have already given in (34). We are going to compute some Θ -class intersection numbers of the form $\int_{\overline{\mathcal{M}}_{1+nb,n}} \Theta_{1+nb,n} \psi_1^b \dots \psi_n^b$, denoted by $\Omega_{b^n}^{\Theta(1/8)}(0)$ for short. Using an algorithm designed in [37] and using Theorem 3, one can compute these Θ -class intersection numbers in relatively high genera. For example we have

$$\Omega_{1^{11}}^{\Theta(1/8)}(0) = \frac{3727154672771403705644393643825}{67108864},$$

$$\Omega_{2^8}^{\Theta(1/8)}(0) = \frac{10497097022517857530944569189202112366380675}{36028797018963968},$$

$$\Omega_{3^6}^{\Theta(1/8)}(0) = \frac{291143373745168297982109927833062542748609508458550221626925092375}{604462909807314587353088}.$$

Several more Θ -class correlators are in Table 1. One observes that $\Omega_{0^n}^{\Theta(1/8)}(0) = \frac{(n-1)!}{8}$.

	$b = 0$	$b = 1$	$b = 2$	$b = 3$
$n = 1$	$\frac{1}{8}$	$\frac{3}{128}$	$\frac{15}{1024}$	$\frac{525}{32768}$
$n = 2$	$\frac{1}{8}$	$\frac{63}{512}$	$\frac{125565}{131072}$	$\frac{178066035}{8388608}$
$n = 3$	$\frac{1}{4}$	$\frac{7221}{2048}$	$\frac{8160299505}{8388608}$	$\frac{5357097499513095}{4294967296}$
$n = 4$	$\frac{3}{4}$	$\frac{4825971}{16384}$	$\frac{6118287865593075}{1073741824}$	$\frac{3673662570422147820860595}{4398046511104}$
$n = 5$	3	$\frac{3540311739}{65536}$	$\frac{2089963670900974355205}{17179869184}$	$\frac{7614423907504732590945890803999875}{2251799813685248}$
$n = 6$	15	$\frac{1209901485555}{65536}$	$\frac{31867458860062839143669852025}{4398046511104}$	$\frac{32942281960173069977596091564715863342175375}{576460752303423488}$

TABLE 1. $\Omega_{b^n}^{\Theta(1/8)}$, $b = 0, 1, 2, 3$.

The algorithm designed in [37] also produces explicit full genera formulas for the Θ -class correlators:

$$\begin{aligned}
\sum_{p \geq 0} \frac{(1+2p)!!}{\lambda^{p+1}} \Omega_{0,p}^{\Theta(1/8)}(0) &= b - 1 \\
- 512 \, 3!! \, 5!! \sum_{i \geq 0} \frac{(1+2p)!!}{\lambda^{p+1}} \Omega_{1,2,p}^{\Theta(1/8)}(0) &= -1024 \lambda^3 a + 256 \lambda^2 (b - a) + \lambda (848b - 616a - 128c) \\
&\quad + 3321b - 2187a - 432c \\
1024 \, 3!!^3 \sum_{p \geq 0} \frac{(1+2p)!!}{\lambda^{p+1}} \Omega_{1,1,p}^{\Theta(1/8)}(0) &= 2048 \lambda^4 b + \lambda^3 (256b - 2048c) - \lambda^2 (1536a + 96b + 768c) \\
&\quad + \lambda (1728a - 8820b - 864c) + 26352a - 46989b + 4284c
\end{aligned}$$

In the above formulas, $a = \frac{1}{8\lambda} {}_3F_0\left(\frac{3}{2}, \frac{1}{2}, \frac{1}{2}; \frac{1}{\lambda}\right)$, $b = {}_3F_0\left(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}; \frac{1}{\lambda}\right)$, and

$$c = \left(\lambda - \frac{1}{4}\right) {}_3F_0\left(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}; \frac{1}{\lambda}\right) - \frac{3}{8\lambda} {}_3F_0\left(\frac{3}{2}, \frac{1}{2}, \frac{1}{2}; \frac{1}{\lambda}\right) - \frac{27}{32\lambda^2} {}_3F_0\left(\frac{5}{2}, \frac{1}{2}, \frac{1}{2}; \frac{1}{\lambda}\right).$$

4.3.2. *Correlators with parameter C .* We list some correlators of the form $\Omega_{b^n}^{\Theta(C)}(0)$ in Tables 2–3.

	$b = 0$	$b = 1$
$n = 1$	C	$\frac{1}{6}C(C+1)$
$n = 2$	C	$\frac{1}{6}C(C+1)(2C+5)$
$n = 3$	$2C$	$\frac{1}{3}C(C+1)(2C+7)(3C+10)$
$n = 4$	$6C$	$C(C+1)(22C^3 + 292C^2 + 1320C + 1925)$
$n = 5$	$24C$	$C(C+1)(364C^4 + 8028C^3 + 69089C^2 + 261625C + 350350)$
$n = 6$	$120C$	$C(C+1)(8160C^5 + 272480C^4 + 3843730C^3 + 27340910C^2 + 93831500C + 119119000)$

TABLE 2. $\Omega_{b^n}^{\Theta(C)}(0)$, $b = 0, 1$.

$n = 1$	$\frac{1}{30}C(C+1)(C+3)$
$n = 2$	$\frac{1}{60}C(C+1)(C+3)(3C^2+38C+126)$
$n = 3$	$\frac{1}{60}C(C+1)(C+3)(15C^4+550C^3+8011C^2+52521C+126126)$
$n = 4$	$\frac{1}{120}C(C+1)(C+3)(285C^6+21120C^5+701455C^4+12823420C^3+131525532C^2+698301072C+1466593128)$
$n = 5$	$\frac{1}{30}C(C+1)(C+3)(1035C^8+131220C^7+7929500C^6+286890460C^5+6581287505C^4+95511193020C^3+838324176858C^2+3995785717308C+7792009289064)$
$n = 6$	$\frac{1}{72}C(C+1)(C+3)(49329C^{10}+9683190C^9+941488056C^8+57517664804C^7+2370818604241C^6+67214920642718C^5+1301029520426886C^4+16691838842700000C^3+133819015248860760C^2+596987475819494760C+1110408075747354384)$

TABLE 3. $\Omega_{2^n}^{\Theta(C)}(0)$.

We list first few explicit formulas for certain correlators in full genera:

$$\begin{aligned}
\sum_{i \geq 0} \frac{(1+2p)!!}{\lambda^{p+1}} \Omega_{0,p}^{\Theta(C)}(0) &= b - 1 \\
3!! \sum_{i \geq 0} \frac{(1+2p)!!}{\lambda^{p+1}} \Omega_{1,p}^{\Theta(C)}(0) &= \lambda(2b-3) + c - Cb \\
5!! \sum_{i \geq 0} \frac{(1+2p)!!}{\lambda^{i+1}} \Omega_{2,p}^{\Theta(C)}(0) &= \lambda^2(3b-5) + 2\lambda(c-Cb) + 2Ca - \frac{C(C+3)}{2}b + Cc \\
&- \frac{3!!^2}{2} \sum_{p \geq 0} \frac{(1+2p)!!}{\lambda^{i+1}} \Omega_{1,1,p}^{\Theta(C)}(0) = -\lambda^2a + 2C\lambda(b-a) - C(C+3)a + C(2C+3)b - Cc \\
&- 2 \cdot 5!!^2 \sum_{i \geq 0} \frac{(1+2p)!!}{\lambda^{p+1}} \Omega_{2,2,p}^{\Theta(C)}(0) = -4\lambda^4a + 8C\lambda^3(b-a) + 4C\lambda^2((8C+9)b - c - (4C+6)a) \\
&\quad + C\lambda((36C^2+216C+180)b - (12C^2+132C+120)a - 24(C+1)c) \\
&\quad + 12C(3C^3+49C^2+151C+105)b - 18C(5+7C+2C^2)c - 9C(C^3+30C^2+99C+70)a
\end{aligned}$$

with $a = \frac{C}{\lambda}G_{\frac{3}{2}}(\lambda)$, $b = G_{\frac{1}{2}}(\lambda)$, and $c = (\lambda-2C)G_{\frac{1}{2}}(\lambda) - \frac{3C}{\lambda}G_{\frac{3}{2}}(\lambda) - \frac{6C(C+1)}{\lambda^2}G_{\frac{5}{2}}(\lambda)$, where we recall that $G_\alpha(\zeta)$ is defined in (40).

5. GENERATING SERIES OF THE LAMÉ PARTIAL CORRELATION FUNCTIONS

In this section, we study the Lamé tau-function τ_{elliptic} .

5.1. An explicit recursion for the basic matrix resolvent. Recall that the Lamé solution u_{elliptic} of the KdV hierarchy is the unique solution in the ring $V[[t_{>0}]]$, where $V = \mathbb{C}[g_2, g_3, \wp, \wp']/(\wp'^2 - 4\wp^3 + g_2\wp + g_3)$, having the initial data $f = C\wp(x; \tau)$. Let $R(\lambda, x)$ and $b(\lambda, x)$ denote the functions as in Section 1.4 (Example 3). Replacing ∂ with ∂_x and with u with $f = C\wp(x; \tau)$ in (60) we obtain the following nonlinear ODE for $b(\lambda, x)$

$$b_{xx}b - \frac{1}{2}b_x^2 - 2(\lambda - 2C\wp(x; \tau))b^2 = -2\lambda. \quad (108)$$

We note again that $b = b(\lambda, x)$ is the unique power series solution in λ^{-1} , having the form $b = \sum_{k \geq -1} b_k/\lambda^{k+1}$ with $b_{-1} = 1$, to equation (108). Denote $X = \wp(x; \tau)$, $Y = \wp'(x; \tau)$. Recall that

$$Y^2 = 4X^3 - g_2X - g_3 = 4(X - e_1)(X - e_2)(X - e_3)$$

with g_2 and g_3 as in Section 1.1. In the (X, λ) -coordinates, equation (108) has the equivalent expression

$$\left(X^3 - \frac{g_2}{4}X - \frac{g_3}{4}\right)(2bb'' - b'^2) + \left(3X^2 - \frac{g_2}{4}\right)bb' + (2CX - \lambda)b^2 + \lambda = 0. \quad (109)$$

Here $' = \frac{\partial}{\partial X}$. It follows that b_k must have the form $b_k = P_{k+1}$ with $P_0 = 1$ and P_m ($m \geq 1$) being polynomials in X of degree m whose coefficients are polynomials of g_2, g_3, C . Equation (109) uniquely determines P_k , $k \geq 1$ in the following recursive way:

$$\begin{aligned} P_k = & \sum_{i=0}^{k-1} \left[CXP_iP_{k-i-1} + \left(\frac{3}{2}X^2 - g_2\right)P_iP'_{k-i-1} + \left(X^3 - \frac{1}{4}g_2X - \frac{1}{4}g_3\right)\left(P_iP''_{k-i-1} - \frac{1}{2}P'_iP'_{k-i-1}\right) \right] \\ & - \frac{1}{2} \sum_{i=1}^{k-1} P_iP_{k-i}. \end{aligned} \quad (110)$$

We do not have a closed expression of b for a general value of C .

5.2. Proof of Theorem 6. Define $R^{\text{sp}}(\lambda, x) := \sqrt{\frac{S_p(\lambda)}{\lambda}} R(\lambda, x)$. We have

$$\begin{aligned} [\mathcal{L}, R^{\text{sp}}] &= 0, \\ \det R^{\text{sp}} &= -S_p(\lambda). \end{aligned}$$

One identifies R^{sp} with the matrix resolvent in [34]. This shows that $\sqrt{\frac{S_p(\lambda)}{\lambda}} b(\lambda, x)$ is a polynomial in λ of degree p with leading coefficient 1. The fact that $\sqrt{\frac{S_p(\lambda)}{\lambda}} b(\lambda, x)$ is also a degree p polynomial of $X = \wp$ can be deduced from homogeneity. Indeed, if we introduce a gradation by $\text{wt } \lambda = 2$, $\text{wt } X = 2$, $\text{wt } g_2 = 4$, $\text{wt } g_3 = 6$, then we find that $b(\lambda, x)$ is homogeneous of degree 0. Note that $S_p(\lambda)$ is homogeneous of degree $4p + 2$, i.e. $\text{wt } S_p(\lambda) = 4p + 2$. The theorem is proved. $\square$

5.3. Computations. The first few partial correlation functions of u_{elliptic} are

$$\begin{aligned} \Omega_{0,0}(x) &= C\wp(x), \quad \Omega_{0,0,0}(x) = C\wp'(x), \quad \Omega_{0,0,0,0}(x) = 6C\wp(x)^2 - \frac{Cg_2}{2}, \\ \Omega_{1,1}(x) &= C\left(\frac{(C+1)(2C+5)}{6}\wp(x)^3 - \frac{(C+1)g_2}{8}\wp(x) - \frac{C+2}{24}g_3\right), \\ \Omega_{1,1,1}(x) &= \frac{C(C+1)}{24}\wp'(x)\left(4(2C+7)(3C+10)\wp(x)^3 - (11C+28)g_2\wp(x) - 2(C+5)g_3\right). \end{aligned}$$

Considering the Laurent series expansion of $\wp(x)$ at $x = 0$, we obtain

$$\begin{aligned} \Omega_{0,0}(x) &= C\left(\frac{1}{x^2} + \frac{g_2}{20}x^2 + \frac{g_3}{28}x^4 + \frac{g_2^2}{1200}x^6 + \frac{3g_2g_3}{6160}x^8 + \frac{49g_2^3 + 750g_3^2}{7644000}x^{10} + \dots\right), \\ \Omega_{0,0,0}(x) &= C\left(\frac{2}{x^3} - \frac{g_2}{10}x - \frac{g_3}{7}x^3 - \frac{g_2^2}{200}x^5 - \frac{3g_2g_3}{770}x^7 - \frac{49g_2^3 + 750g_3^2}{764400}x^9 - \dots\right), \\ \Omega_{1,1}(x) &= \frac{C(C+1)(2C+5)}{6x^6} + \frac{C^2(C+1)}{20x^2}g_2 + \frac{C(6C^2+14C+1)}{168}g_3 \\ &\quad + \frac{C(C+1)(8C+5)}{2400}g_2^2x^2 + \dots, \\ \Omega_{1,1,1}(x) &= \frac{C(C+1)(2C+7)(3C+10)}{6x^9} + \frac{C(C+1)(9C^2+34C+35)}{60x^5}g_2 \\ &\quad + \frac{C(C+1)(18C^2+109C+140)}{168x^3}g_3 + \frac{C(C+1)(24C^2+109C+140)}{2400x}g_2^2 + \dots. \end{aligned}$$

Since each of these expressions is modular of some weight with respect to the action $\tau \mapsto \frac{a\tau+b}{c\tau+d}$, $x \mapsto \frac{x}{c\tau+d}$ of the full modular group, we call these Lamé partial correlation functions *modular deformations* of the generalized BGW partial correlation functions (after a shift of x by 1). The partial correlation functions also allow us to consider other analytic in τ aspects. For example, using

$$\wp(x) = \frac{\pi^2}{3}(2 + 3 \cot^2(\pi x)) + 16 q^2 \pi^2 \sin^2(\pi x) + 4 q^4 (5 + 4 \cos(2\pi x)) \sin^2(\pi x) + \dots$$

one can obtain the $q \rightarrow 0$ expansion ($q = e^{\pi i \tau}$) of the Lamé partial correlation functions. The above modular deformations could be interpreted in an alternative way. Namely, we first switch on the periods $(2\omega, 2\omega')$ of the Weierstrass $\wp$ -function. (Previously we have taken $\omega = \frac{1}{2}$ and $\omega' = \frac{\tau}{2}$). It is easy to see that the form of the partial correlation functions do not change when the initial data is given by $f(x) \equiv C \wp(x; 2\omega, 2\omega')$. Denote $\tau = \frac{\omega'}{\omega}$, $q = e^{\pi i \tau}$. Then we have as $\omega \rightarrow \infty$,

$$\wp(x) = \frac{1}{x^2} + \frac{\pi^4 x^2 (1 + 240 q^2 + 2160 q^4 + \dots)}{240 \omega^4} + \frac{\pi^6 x^4 (1 - 504 q^2 - 16632 q^4 + \dots)}{6048 \omega^6} + O(\omega^{-7}).$$

Hence the modular deformation of a generalized BGW partial correlation function can also be viewed as an $\omega \rightarrow \infty$ asymptotic.

Remark 4. For $C = 1/8$, we expect the existence of a deformation of the Norbury class such that the partition function is equal to the Lamé tau-function. For other values of C , we also expect the existence of cohomology classes on $\overline{\mathcal{M}}_{g,n}$ giving rise to the Lamé tau-function.

Using Corollary 1 we can obtain formulas like

$$\begin{aligned} -\frac{3!!^2}{2} \sum_{i \geq 0} \frac{(1+2i)!!}{\lambda^{i+1}} \Omega_{1,1,i}(x) &= -\lambda^2 a + C \lambda (\wp'(x) b - 2\wp(x) a) \\ &+ \frac{C}{4} (g_2 - 4(C+3)\wp(x)^2) a + \frac{C(2C+3)}{2} \wp(x) \wp'(x) b - \frac{C}{2} \wp'(x) c, \end{aligned} \quad (111)$$

where a, b, c are the components of $R(\lambda, x)$.

6. FURTHER REMARKS

6.1. The second kind topological ODE of A_1 -type. Consider the solution of the KdV hierarchy with initial data $f = C/(1-x)^2$. Let $R = R(\lambda, x)$ denote the basic matrix resolvent of this solution evaluated at $\mathbf{t}_{>0} = \mathbf{0}$ and $t_0 = x$. Define $M(\lambda, x) = \frac{1}{\lambda^{\frac{3}{2}}} \lambda^{\frac{\sigma_3}{4}} R(\lambda, x) \lambda^{-\frac{\sigma_3}{4}}$. According to Theorem 4 we have $M(\lambda, x) = M(\zeta)$ with $\zeta = \lambda(x-1)^2$. Moreover, $M = M(\zeta)$ satisfies

$$\frac{dM}{d\zeta} + \frac{1}{2} \left[\begin{pmatrix} 0 & 1/\sqrt{\zeta} \\ 1/\sqrt{\zeta} - 2C/\zeta & 0 \end{pmatrix}, M \right] = 0. \quad (112)$$

We call (112) the *second kind topological ODE of A_1 -type*. Note that as $\zeta \rightarrow \infty$, we have

$$M(\zeta) \sim \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} + O(1/\sqrt{\zeta}).$$

Equation (112) together with this boundary condition uniquely determines M . This M further satisfies $\det M(\zeta) = -1$. The second kind topological ODE (112) can be written equivalently as

$$\frac{dM}{dz} + \left[\begin{pmatrix} 0 & 1 \\ 1 - \frac{2C}{z^2} & 0 \end{pmatrix}, M \right] = 0, \quad (113)$$

where $\zeta = z^2$. More details about the second kind topological ODE and its generalization to an arbitrary simple Lie algebra will be given in a subsequent publication.

6.2. M -bispectrality. Let us consider $V = \mathbb{C}((x))$, the ring of Laurent series of x . Let $u = u(\mathbf{t})$ be a solution of the KdV hierarchy (2) in $V[[\mathbf{t}_{>0}]]$, $f(x) = u(x, \mathbf{0})$ the initial value. Let us recall a notion of bispectral solutions [40] defined by using $R(\lambda, x)$. Denote $\sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$.

Definition 3. The solution u is called M -bispectral if there exist three scalar functions $g(\lambda) \neq 0$, $g_0(\lambda) \neq 0$, $h(\lambda, x)$, and an $\mathfrak{sl}_2(\mathbb{C})$ -valued function $\tilde{R}(h)$ such that

$$R(\lambda, x) = g(\lambda) g_0(\lambda)^{\sigma_3} \tilde{R}(h(\lambda, x)) g_0(\lambda)^{-\sigma_3}. \quad (114)$$

Theorem 7. The solution u of the KdV hierarchy is M -bispectral iff

$$\text{either } f(x) = (x - A)C, \quad \text{or } f(x) = \frac{C}{(x - A)^2} + B, \quad (115)$$

where A, B, C are constants.

Proof. Denote by $a(\lambda, x), b(\lambda, x), c(\lambda, x)$ the entries of $R(\lambda, x)$. They must have the form

$$a(\lambda, x) = g(\lambda) \rho_1(h(\lambda, x)), \quad (116)$$

$$b(\lambda, x) = q(\lambda) g(\lambda) \rho_2(h(\lambda, x)), \quad (117)$$

$$c(\lambda, x) = q(\lambda)^{-1} g(\lambda) \rho_3(h(\lambda, x)) \quad (118)$$

for some functions $g(\lambda) \neq 0$, $q(\lambda) = g_0(\lambda)^2 \neq 0$, $\rho_1(h)$, $\rho_2(h) \neq 0$, $\rho_3(h)$. Substituting these expressions in equation (57) we obtain

$$\frac{1}{2} q(\lambda) g(\lambda) \rho_2'(h(\lambda, x)) h_x = g(\lambda) \rho_1(h(\lambda, x)).$$

This implies existence of three functions of one-variable W_1, W_2, P satisfying

$$h(\lambda, x) = P(x W_1(\lambda) + W_2(\lambda)). \quad (119)$$

Hence we can assume that

$$h(\lambda, x) = x W_1(\lambda) + W_2(\lambda), \quad W_1(\lambda) \neq 0. \quad (120)$$

It further implies that

$$\frac{1}{2} q(\lambda) W_1(\lambda) = \frac{\rho_1(h(\lambda, x))}{\rho_2'(h(\lambda, x))} = C_0,$$

where C_0 is a constant. Using equations (118) and (58) we have

$$(\lambda - 2f(x)) \rho_2(h(\lambda, x)) - \frac{W_1(\lambda)^2}{2} \rho_2''(h(\lambda, x)) = \frac{W_1(\lambda)^2}{4C_0^2} \rho_3(h(\lambda, x)).$$

Therefore

$$\lambda - 2f(x) = \frac{W_1(\lambda)^2}{2} \frac{\rho_2''(h(\lambda, x))}{\rho_2(h(\lambda, x))} + \frac{W_1(\lambda)^2}{4C_0^2} \frac{\rho_3(h(\lambda, x))}{\rho_2(h(\lambda, x))}. \quad (121)$$

Differentiating both sides w.r.t. λ we obtain

$$1 = \partial_\lambda \left(\frac{W_1(\lambda)^2}{2} \frac{\rho_2''(h(\lambda, x))}{\rho_2(h(\lambda, x))} + \frac{W_1(\lambda)^2}{4C_0^2} \frac{\rho_3(h(\lambda, x))}{\rho_2(h(\lambda, x))} \right). \quad (122)$$

Denote

$$G(h) := \frac{1}{2} \frac{\rho_2''(h)}{\rho_2(h)} + \frac{1}{4C_0^2} \frac{\rho_3(h)}{\rho_2(h)}.$$

We have

$$1 = \partial_\lambda (W_1(\lambda)^2 G(h(\lambda, x))) = 2W_1 W_1' G + (x W_1' + W_2') W_1^2 G'.$$

Differentiating both sides with respect to x we find

$$0 = 3W_1^2 W_1' G' + (x W_1' + W_2') W_1^3 G''.$$

This leads to two possibilities:

$$\text{i) } W_1' = 0 : \quad W_2' G'' = 0, \quad (123)$$

$$\text{ii) } W_1' \neq 0 : \quad \left(G'' \neq 0 \text{ and } x = -\frac{3W_1' G' + W_2' W_1 G''}{W_1' W_1 G''} \right) \quad \text{or} \quad (G'' = G' = 0). \quad (124)$$

For the case i), we have $W_1 \equiv C_1$ for some non-zero constant C_1 . So

$$1 = W_2' G' C_1^2.$$

This implies that $W_2' \neq 0$ and so $G'' \equiv 0$. Hence $G(h) = C_2 h + C_4$ and $W_2 = \frac{\lambda}{C_2 C_1^2} + C_3$ for some constants C_2, C_3, C_4 . Substituting these expressions in (121) we find

$$u = -\frac{1}{2} C_1^3 C_2 x - \frac{1}{2} C_1^2 C_2 C_3 - \frac{1}{2} C_1^2 C_4.$$

For the case ii), we have

$$1 = -3 \left(\frac{G'}{G''} \right)', \quad \text{or} \quad G = \text{Const}.$$

So

$$G(h) = \frac{C_1}{(h + C_2)^2} + C_3$$

for some constants C_1, C_2, C_3 . Then we find that

$$W_1 = \left(\frac{\lambda + 2C_3 C_4}{C_3} \right)^{\frac{1}{2}}, \quad (125)$$

$$W_2 = C_5 (\lambda + 2C_3 C_4)^{\frac{1}{2}} - C_2, \quad (126)$$

$$f = -\frac{C_1}{2(x + \sqrt{C_3} C_5)^2} - C_3 C_4 \quad (127)$$

for some constants C_4, C_5 . We have proved that u must have the form in (115).

We are left to prove the solvability of existence of $g_0, g, \tilde{R}, h$ when f is taken to be one of (115). For $f(x) = (x - A)C$, we can take for instance

$$g_0 = 1, \quad g = \frac{\sqrt{\lambda}}{2^{\frac{1}{3}} C^{\frac{1}{3}}}, \quad W_1 = -2^{\frac{1}{3}} C^{\frac{1}{3}}, \quad W_2 = \frac{\lambda}{2^{\frac{2}{3}} C^{\frac{2}{3}}},$$

and we have

$$b = \frac{\sqrt{\lambda}}{2^{\frac{1}{3}} C^{\frac{1}{3}}} \sum_{g=0}^{\infty} \frac{(6g-1)!!}{96^g g!} \zeta^{-3g-\frac{1}{2}}, \quad \zeta := -(x - A) 2^{\frac{1}{3}} C^{\frac{1}{3}} + \frac{\lambda}{2^{\frac{2}{3}} C^{\frac{2}{3}}}.$$

For $f(x) = \frac{C}{(x-A)^2} + B$, we can take for instance

$$g_0 = \frac{1}{(\lambda - 2B)^{\frac{1}{4}}}, \quad g = \sqrt{\lambda}, \quad W_1 = \sqrt{\lambda - 2B}, \quad W_2 = -A \sqrt{\lambda - 2B},$$

and we have

$$b = \frac{\sqrt{\lambda}}{\sqrt{\lambda - 2B}} {}_3F_0 \left(\frac{1}{2}, \frac{1 + \sqrt{1 - 8C}}{2}, \frac{1 - \sqrt{1 - 8C}}{2}; ; \frac{1}{(x - A)\sqrt{\lambda - 2B}} \right).$$

The theorem is proved. $\square$

Remark 5. For the first case of (115), when (A, C) is taken as $(0, 1)$, the above theorem recovers the following formula of [13, 15, 71]:

$$\sum_{g, p_1, \dots, p_n \geq 0} \int_{\mathcal{M}_{g,n}} \psi_1^{p_1} \cdots \psi_n^{p_n} \prod_{j=1}^n \frac{(2p_j + 1)!!}{\lambda_j^{p_j+1}} = - \sum_{\sigma \in S_n/C_n} \frac{\text{tr } M(\lambda_{\sigma(1)}) \cdots M(\lambda_{\sigma(n)})}{\prod_{i=1}^n (\lambda_{\sigma(i+1)} - \lambda_{\sigma(i)})} - \delta_{n2} \frac{\lambda_1 + \lambda_2}{(\lambda_1 - \lambda_2)^2}, \quad (128)$$

where $M(\lambda)$ is a 2 by 2 matrix given explicitly by

$$M(\lambda) = \begin{pmatrix} \frac{1}{2} \sum_{g=1}^{\infty} \frac{(6g-5)!!}{2^{4g-1} (g-1)!} \lambda^{-3g+2} & \sum_{g=0}^{\infty} \frac{(6g-1)!!}{2^{4g} g!} \lambda^{-3g} \\ \sum_{g=0}^{\infty} \frac{1+6g}{1-6g} \frac{(6g-1)!!}{2^{4g} g!} \lambda^{-3g+1} & -\frac{1}{2} \sum_{g=1}^{\infty} \frac{(6g-5)!!}{2^{4g-1} (g-1)!} \lambda^{-3g+2} \end{pmatrix}. \quad (129)$$

We note that $\lambda^{-\frac{1}{2}} M(\lambda)$ satisfies an ODE in λ , *the first kind topological ODE of A_1 -type* [14, 15].

Remark 6. For the first case of (115), when $(A, C) = (q, 1)$, the corresponding KdV tau-function gives the partition function of stable quasi-map invariants [23] of a point.²

Remark 7. For the second case of (115), when (A, B, C) is take as $(1, 0, \frac{1}{8})$, the above theorem recovers the formula (39).

The following lemma due to Norbury has a similar flavour with Theorem 7.

Lemma 9 ([62]). *A collection of cohomology classes $\{\Theta_{g,n} \in H^*(\overline{\mathcal{M}}_{g,n})\}_{2g-2+n>0}$ satisfying the properties i) and ii) (see equations (93) and (94)) must satisfy*

$$\text{either } \deg \Theta_{g,n} \equiv 0, \quad \text{or } \deg \Theta_{g,n} \equiv 2g - 2 + n.$$

Proof Denote $d(g, n) = \deg \Theta_{g,n}$. Using the first equation of (94) we obtain

$$d(g, n) = d(g-1, n+2)$$

which implies the existence of a function H of one variable such that $d(g, n) = H(2g - 2 + n)$. Now using the second equation of (94) we find

$$H(x+y) = H(x) + H(y).$$

Therefore there exists an integer constant Q such that $d(g, n) = (2g - 2 + n) Q$. Noting that

$$0 \leq d(g, n) \leq 3g - 3 + n,$$

we have either $Q = 0$ or $Q = 1$. □

REFERENCES

- [1] Adler, M., van Moerbeke, P. (1992). A matrix integral solution to two-dimensional W_p -gravity. *Comm. Math. Phys.*, **147**, 25–56.
- [2] Alexandrov, A. (2018). Cut-and-join description of generalized Brezin–Gross–Witten model. *Adv. Theor. Math. Phys.*, **22**, 1347–1399.
- [3] Alexandrov, A. Matrix model for the stationary sector of Gromov–Witten theory of $\mathbf{P}^1$. arXiv:2001.08556.
- [4] Babelon, O., Bernard, D., Talon, M. (2003). *Introduction to classical integrable systems*. Cambridge Univ. Press.
- [5] Belavin, A., Dubrovin, B., Mukhametzhanov, B. (2014). Minimal Liouville gravity correlation numbers from Douglas string equation. *J. High Energy Phys.*, **156**, front matter+50 pp.
- [6] Balogh, F., Yang, D. (2017). Geometric interpretation of Zhou’s explicit formula for the Witten–Kontsevich tau function. *Lett. Math. Phys.*, **107**, 1837–1857.
- [7] Basor, E.L., Tracy, C.A. (1993). Variance calculations and the Bessel kernel. *Journal of Statistical Physics*, **73**, 415–421.
- [8] Belliard, R., Eynard, B., Marchal, O. (2017). Integrable differential systems of topological type and reconstruction by the topological recursion. *Ann. Henri Poincaré*, **18**, 3193–3248.

²We are grateful to Bumsig Kim for sharing with us his knowledge about this interesting point.

- [9] Belliard, R., Eynard, B., Marchal, O. (2018). Loop equations from differential systems on curves. *Ann. Henri Poincaré*, **19**, 141–161.
- [10] Bergère, M., Eynard, B. Determinantal formulae and loop equations. arXiv:0901.3273.
- [11] Bergère, M., Borot, G., Eynard, B. (2015). Rational differential systems, loop equations, and application to the qth reductions of KP. *Ann. Henri Poincaré*, **16**, 2713–2782.
- [12] Bertola, M., Cafasso, M. (2017). The Kontsevich matrix integral: convergence to the Painlevé hierarchy and Stokes’ phenomenon. *Comm. Math. Phys.*, **352**, 585–619.
- [13] Bertola, M., Dubrovin, B., Yang, D. (2016). Correlation functions of the KdV hierarchy and applications to intersection numbers over $\overline{\mathcal{M}}_{g,n}$. *Physica D: Nonlinear Phenomena*, **327**, 30–57.
- [14] Bertola, M., Dubrovin, B., Yang, D. (2018). Simple Lie algebras and topological ODEs. *IMRN*, **2018**, 1368–1410.
- [15] Bertola, M., Dubrovin, B., Yang, D. Simple Lie algebras, Drinfeld–Sokolov hierarchies, and multi-point correlation functions. *Moscow Mathematical Journal* (to appear). arXiv:1610.07534.
- [16] Bertola, M., Ruzza, G. (2019). The Kontsevich–Penner matrix integral, isomonodromic tau functions and open intersection numbers. *Annales Henri Poincaré*, **20**, 393–443.
- [17] Bertola, M., Ruzza, G. (2019). Brezin–Gross–Witten tau function and isomonodromic deformations. *Commun. Number Theory Phys.*, **13**, 827–883.
- [18] Bertola, M., Ruzza, G. Matrix models for stationary Gromov–Witten invariants of the Riemann sphere. arXiv:2001.10466.
- [19] Bertola, M., Yang, D. (2015). The partition function of the extended r -reduced Kadomtsev–Petviashvili hierarchy. *Journal of Physics A: Mathematical and Theoretical*, **48**, 195205.
- [20] Brézin, E., Gross, D.J. (1980). The external field problem in the large N limit of QCD. *Phys. Lett., B*, **97**, 120–124.
- [21] Buryak, A. (2016). Open intersection numbers and the wave function of the KdV hierarchy. *Mosc. Math. J.* **16**, 27–44.
- [22] Buryak, A., Dubrovin, B., Guéré, J., Rossi, P. (2018). Tau-structure for the double ramification hierarchies. *Comm. Math. Phys.* **363**, 191–260.
- [23] Ciocan-Fontanine, I., Kim, B. (2010). Moduli stacks of stable toric quasimaps. *Advances in Mathematics*, **225**, 3022–3051.
- [24] Chekhov, L., Eynard, B., Orantin, N. (2006). Free energy topological expansion for the 2-matrix model. *J. High Energy Phys.* **2006**, 053, 31pp.
- [25] Date, E., Jimbo, M., Kashiwara, M., Miwa, T. (1981). Transformation groups for soliton equations. In *Publications of RIMS*, **18** (3), 1077–1110.
- [26] Date, E., Kashiwara, M., Jimbo, M., Miwa, T. (1983). Transformation groups for soliton equations. *Nonlinear integrable systems—classical theory and quantum theory* (Kyoto, 1981), 39–119, World Sci. Publishing, Singapore.
- [27] Deift, P. *Polynomials, Orthogonal, Random Matrices: A Riemann–Hilbert Approach*. American Mathematical Society, Providence, RI, 1999.
- [28] Dickey, L.A. (2003). *Soliton Equations and Hamiltonian Systems*, 2nd edition. World Scientific.
- [29] Dijkgraaf, R., Verlinde, H., Verlinde, E. (1991). Loop equations and Virasoro constraints in nonperturbative two-dimensional quantum gravity. *Nuclear Phys. B*, **348**, 435–456.
- [30] Do, N., Norbury, P. (2018). Topological recursion on the Bessel curve. *Communications in Number Theory and Physics*, **12**, 53–73.
- [31] Drinfeld, V.G., Sokolov, V.V. (1985). Lie algebras and equations of Korteweg–de Vries type, *J. Math. Sci.*, **30**, 1975–2036. Translated from *Itogi Nauki i Tekhniki, Seriya Sovremennye Problemy Matematiki (Noveishie Dostizheniya)*, **24** (1984), 81–180.
- [32] Dubrovin, B. (1975). Periodic problem for the Korteweg–de Vries equation in the class of finite band potentials. *Functional Analysis and its Applications*, **9**, 215–223.
- [33] Dubrovin, B. (1996). Geometry of 2D topological field theories. In “Integrable Systems and Quantum Groups” (Montecatini Terme, 1993). Editors: Francaviglia, M., Greco, S. Springer Lecture Notes in Math. **1620**, 120–348.
- [34] Dubrovin, B. (2009). Integrable Systems and Riemann Surfaces. Lecture Notes (preliminary version). Available online: http://people.sissa.it/~dubrovin/rsnleq_web.pdf.
- [35] Dubrovin, B., Liu S.-Q., Yang, D., Zhang, Y. (2016). Hodge integrals and tau-symmetric integrable hierarchies of Hamiltonian evolutionary PDEs. *Advances in Mathematics*, **293**, 382–435.
- [36] Dubrovin, B., Novikov, S.P. (1974). Periodic and conditionally periodic analogs of the many-soliton solutions of the Korteweg–de Vries equation. *Zh. Eksper. Teoret. Fiz.*, **67**, 2131–2144.
- [37] Dubrovin, B., Yang, D. (2016). Generating series for GUE correlators. *Lett. Math. Phys.*, **107**, 1971–2012.
- [38] Dubrovin, B., Yang, D. (2019). On Gromov–Witten invariants of $\mathbb{P}^1$. *Mathematical Research Letters*, **26**, 729–748.
- [39] Dubrovin, B., Yang, D. (2020). Matrix resolvent and the discrete KdV hierarchy. *Comm. Math. Phys.* **377**, 1823–1852.
- [40] Dubrovin, B., Yang, D., Zagier, D. (2018). Gromov–Witten invariants of the Riemann sphere. *Pure Appl. Math. Q.*, **16**, 153–190.

- [41] Dubrovin, B., Yang, D., Zagier, D. Geometry and arithmetic of integrable hierarchies of KdV type. I. Integrality. arXiv:2101.10924.
- [42] Dubrovin, B., Zhang, Y. Normal forms of hierarchies of integrable PDEs, Frobenius manifolds and Gromov-Witten invariants. arXiv preprint math/0108160.
- [43] Duistermaat, J.J., Grünbaum, F.A. (1986). Differential equations in the spectral parameter. *Comm. Math. Phys.*, **103**, 177–240.
- [44] Ènolskii, V.Z., Harnad, J. (2011). Schur function expansions of KP τ -functions associated to algebraic curves. *Uspekhi Mat. Nauk*, **66**, 137–178; translation in *Russian Math. Surveys*, **66** (2011), 767–807.
- [45] Eynard, B., Orantin, N. (2007). Invariants of algebraic curves and topological expansion. *Commun. Number Theory Phys.* **1**, 347–452.
- [46] Faber, C. (1999). A conjectural description of the tautological ring of the moduli space of curves, *Moduli of curves and abelian varieties*, Aspects Math., E33, Friedr. Vieweg, Braunschweig, 109–129.
- [47] Frenkel, I., Zhu, Y. (1992). Vertex operator algebras associated to representations of affine and Virasoro algebras. *Duke Math. J.*, **66**, 123–168.
- [48] Gross, D.J., Witten, E. (1980). Possible third-order phase transition in the large-N lattice gauge theory. *Physical Review D*, **21**, 446–453.
- [49] Harer, J., Zagier, D. (1986). The Euler characteristic of the moduli space of curves. *Invent. Math.*, **85**, 457–485.
- [50] Hirota, R. (1971). Exact solution of the Korteweg–de Vries equation for multiple collisions of solitons. *Physical Review Letters*, **27**, 1192–1194.
- [51] Gissonni, M., Grava, T., Ruzza, G. (2020). Laguerre Ensemble: Correlators, Hurwitz Numbers and Hodge Integrals. *Annales Henri Poincaré*, **21**, 3285–3339.
- [52] Jimbo, M., Miwa, T., Ueno, K. (1981). Monodromy preserving deformation of linear ordinary differential equations with rational coefficients I. General theory and τ -function. *Phys. D*, **2**, 306–352.
- [53] Kac, V.G. (1978). Infinite-dimensional algebras, Dedekind’s η -function, classical Möbius function and the very strange formula. *Advances in Mathematics*, **30**, 85–136.
- [54] Kac, V., Schwarz, A. (1991). Geometric interpretation of the partition function of 2D gravity. *Phys. Lett. B*, **257**, 329–334.
- [55] Kaufmann, R., Manin, Y., Zagier, D. (1996). Higher Weil–Petersson volumes of moduli spaces of stable-pointed curves. *Comm. Math. Phys.*, **181**, 763–787.
- [56] Kostant, B. (1959). The Principal Three-Dimensional Subgroup and the Betti Numbers of a Complex Simple Lie Group. *American Journal of Mathematics*, **81**, 973–1032.
- [57] Kontsevich, M. (1992). Intersection theory on the moduli space of curves and the matrix Airy function. *Comm. Math. Phys.*, **147**, 1–23.
- [58] Liu, S.-Q., Zhang, Y. (2006). On quasi-triviality and integrability of a class of scalar evolutionary PDEs. *Journal of Geometry and Physics*, **57**, 101–119.
- [59] Marchal, O. (2018). WKB solutions of difference equations and reconstruction by the topological recursion. *Non-linearity* **31**, 226–262.
- [60] Mehta, M.L. *Random Matrices*, 2nd edition. Academic Press, New York, 1991.
- [61] Mironov, A., Morozov, A., Semenoff, G.W. (1996). Unitary matrix integrals in the framework of the generalized Kontsevich model. 1. Brezin-Gross-Witten Model. *International Journal of Modern Physics A*, **11**, 5031–5080.
- [62] Norbury, P. A new cohomology class on the moduli space of curves. arXiv:1712.03662.
- [63] Sato, M. (1981). Soliton Equations as Dynamical Systems on a Infinite Dimensional Grassmann Manifolds (Random Systems and Dynamical Systems). *RIMS Kokyuroku*, **439**, 30–46.
- [64] Segal, G., Wilson, G. (1985). Loop groups and equations of KdV type. *Inst. Hautes Études Sci. Publ. Math. No.* **61**, 5–65.
- [65] Tracy, C.A., Widom, H. (1994). Level spacing distributions and the Bessel kernel. *Comm. Math. Phys.*, **161**, 289–309.
- [66] Witten, E. (1991). Two-dimensional gravity and intersection theory on moduli space. *Surveys in Differential Geometry* (Cambridge, MA, 1990), (pp. 243–310), Lehigh Univ., Bethlehem, PA.
- [67] Yang, D. On the matrix-resolvent approach to tau-functions. Talk given at the IBS-CGP workshop “Integrable systems and applications”, Pohang, 2018-May.
- [68] Yang, D. (2020). On tau-functions for the Toda lattice hierarchy. *Lett. Math. Phys.* **110**, 555–583.
- [69] Zhou, J. (2013). Topological recursions of Eynard-Orantin type for intersection numbers on moduli spaces of curves. *Lett. Math. Phys.* **103**, 1191–1206.
- [70] Zhou, J. Explicit formula for Witten-Kontsevich tau-function. arXiv:1306.5429.
- [71] Zhou, J. On Absolute N-Point Function Associated with Gelfand-Dickey Polynomials. Preprint (2015).
- [72] Zhou, J. Emergent geometry and mirror symmetry of a point. arXiv:1507.01679.

Boris Dubrovin

SISSA, via Bonomea 265, Trieste 34136, Italy

Di Yang

School of Mathematical Sciences, University of Science and Technology of China,
Jinzhai Road 96, Hefei 230026, P.R. China

diyang@ustc.edu.cn

Don Zagier

Max-Planck-Institut für Mathematik, Vivatsgasse 7, Bonn 53111, Germany,
and International Centre for Theoretical Physics, Strada Costiera 11, Trieste 34014, Italy
dbz@mpim-bonn.mpg.de