

Search for lensing signatures in the gravitational-wave observations from the first half of LIGO-Virgo's third observing run

THE LIGO SCIENTIFIC COLLABORATION AND THE VIRGO COLLABORATION

(Dated: 13 May 2021)

ABSTRACT

We search for signatures of gravitational lensing in the gravitational-wave signals from compact binary coalescences detected by Advanced LIGO and Advanced Virgo during O3a, the first half of their third observing run. We study: 1) the expected rate of lensing at current detector sensitivity and the implications of a non-observation of strong lensing or a stochastic gravitational-wave background on the merger-rate density at high redshift; 2) how the interpretation of individual high-mass events would change if they were found to be lensed; 3) the possibility of multiple images due to strong lensing by galaxies or galaxy clusters; and 4) possible wave-optics effects due to point-mass microlenses. Several pairs of signals in the multiple-image analysis show similar parameters and, in this sense, are nominally consistent with the strong lensing hypothesis. However, taking into account population priors, selection effects, and the prior odds against lensing, these events do not provide sufficient evidence for lensing. Overall, we find no compelling evidence for lensing in the observed gravitational-wave signals from any of these analyses.

1. INTRODUCTION

Gravitational lensing occurs when a massive object bends spacetime in a way that focuses light rays toward an observer (see [Bartelmann 2010](#) for a review). Lensing observations are widespread in electromagnetic astrophysics and have been used for, among other purposes, making a compelling case for dark matter ([Clowe et al. 2004](#); [Markevitch et al. 2004](#)), discovering exoplanets ([Bond et al. 2004](#)), and uncovering massive objects and structures that are too faint to be detected directly ([Coe et al. 2013](#)).

Similarly to light, when gravitational waves (GWs) travel near a galaxy or a galaxy cluster, their trajectories curve, resulting in gravitational lensing ([Ohanian 1974](#); [Thorne 1982](#); [Deguchi & Watson 1986](#); [Wang et al. 1996](#); [Nakamura 1998](#); [Takahashi & Nakamura 2003](#)) which changes the GW amplitude without affecting the signal's frequency evolution ([Deguchi & Watson 1986](#); [Wang et al. 1996](#); [Nakamura 1998](#); [Takahashi & Nakamura 2003](#); [Dai & Venumadhav 2017](#); [Ezquiaga et al. 2021](#)). Strong lensing, in particular, can also produce multiple images observed at the GW detectors as repeated events separated by a time delay of minutes to months for galaxies ([Ng et al. 2018](#); [Li et al. 2018](#); [Oguri 2018](#)), and up to years for galaxy clusters ([Smith et al. 2018, 2017, 2019](#); [Robertson et al. 2020](#); [Rydzanowski et al. 2020](#)). The detection of such strongly lensed GWs has been forecast within this decade ([Ng et al. 2018](#); [Li et al. 2018](#); [Oguri 2018](#)), at design sensitivity of Advanced LIGO and Advanced Virgo, assuming that binary black holes (BBHs) trace the star-formation rate density. In addition, if GWs propagate near smaller lenses such as stars or compact objects, microlensing may induce observable beating patterns in the waveform ([Cao](#)

[et al. 2014](#); [Jung & Shin 2019](#); [Lai et al. 2018](#); [Christian et al. 2018](#); [Diego et al. 2019](#); [Diego 2020](#); [Pagano et al. 2020](#)). Indeed, lensing can induce a plethora of effects on GWs.

If observed, GW lensing could enable numerous scientific pursuits, such as localization of merging black holes to sub-arcsecond precision ([Hannuksela et al. 2020](#)), precision cosmography studies ([Sereno et al. 2011](#); [Liao et al. 2017](#); [Cao et al. 2019](#); [Li et al. 2019b](#); [Hannuksela et al. 2020](#)), precise tests of the speed of gravity ([Baker & Trodden 2017](#); [Fan et al. 2017](#)), tests of the GW's polarization content ([Goyal et al. 2021](#)), and detecting intermediate-mass or primordial black holes ([Lai et al. 2018](#); [Diego 2020](#); [Oguri & Takahashi 2020](#)).

Here we perform a comprehensive lensing analysis of data from the first half of the third LIGO–Virgo observing run, called O3a for short, focusing on compact binary coalescence (CBC) signals. We begin by outlining the expected rate of strongly lensed events. Strong lensing is rare, but magnified signals enable us to probe a larger comoving volume, thus potentially giving us access to more sources ([Dai et al. 2017](#); [Ng et al. 2018](#); [Smith et al. 2018](#); [Li et al. 2018](#); [Oguri 2018](#); [Smith et al. 2017, 2019](#); [Robertson et al. 2020](#); [Rydzanowski et al. 2020](#)). We forecast the lensed event rates using standard lens and black hole population models (Sec. 3). These expected rates are subject to some astrophysical uncertainty but are vital to interpreting our search results in later sections.

The rate of lensing can also be inferred from the stochastic GW background (SGWB) ([Buscicchio et al. 2020a](#); [Mukherjee et al. 2021](#); [Buscicchio et al. 2020b](#)). Thus, we use the non-observation of strong lensing and the stochastic background to constrain the BBH merger-rate density and the rate of lensing at high redshifts.

In addition, lensing magnification biases the inferred GW luminosity distance and source mass measurements, which could contribute to an apparently high-mass population of binaries (Dai et al. 2017; Broadhurst et al. 2018; Oguri 2018; Hannuksela et al. 2019; Broadhurst et al. 2020a). Therefore, we analyze several LIGO–Virgo detections with unusually high masses under the alternative interpretation that they are lensed signals from lower-mass sources (Sec. 4).

We then move on to search for signatures of lensing-induced multiple images, which should appear as repeated similar signals, magnified and with waveform differences determined by the image type (Dai & Venumadhav 2017; Ezquiaga et al. 2021), separated in time by minutes to months (or even years). Consequently, if an event pair is strongly lensed, we expect to infer consistent parameters for both events (Haris et al. 2018; Hannuksela et al. 2019).

We search for these multiple images by first comparing the posterior overlap between pairs of events occurring during the O3a period as reported in Abbott et al. (2020d) (Sec. 5.1). After identifying a list of candidates from the posterior-overlap analysis, we follow these up with more computationally expensive but more accurate joint-parameter estimation (PE) procedures (Sec. 5.2). Next, we perform a targeted search for previously undetected counterpart images of known events in Sec. 5.3, images that could have fallen below the threshold of previous wide-parameter space CBC searches (as discussed in Li et al. 2019a; McIsaac et al. 2020; Dai et al. 2020). Finally, we search for microlensing induced by point-mass lenses in the intermediate and low mass range, including wave-optics effects (Sec. 6).

Several searches for GW lensing signatures have already been performed in the data from the first two observing runs O1 and O2 (Hannuksela et al. 2019; Li et al. 2019a; McIsaac et al. 2020; Pang et al. 2020; Liu et al. 2021; Dai et al. 2020), including strong lensing and microlensing effects. We will discuss these previous studies in the appropriate sections. Given the growing interest in GW lensing and the existing forecasts, an analysis of the most recent GW observations for lensing effects is now timely.

Results of all analyses in this paper and associated data products can be found in LIGO Scientific Collaboration and Virgo Collaboration (2021). GW strain data and posterior samples for all events from GWTC-2 are available (GWOSC 2020) from the Gravitational Wave Open Science Center (Abbott et al. 2021a).

2. DATA AND EVENTS CONSIDERED

The analyses presented here use data taken during O3a by the Advanced LIGO (Aasi et al. 2015) and Advanced Virgo (Acernese et al. 2015) detectors. O3a extended from 2019 April 1 to 2019 October 1. Various instrumental upgrades have led to more sensitive data, with median binary

neutron star (BNS) inspiral ranges (Allen et al. 2012a) increased by a factor of 1.64 in LIGO Hanford, 1.53 in LIGO Livingston, and 1.73 in Virgo compared to O2 (Abbott et al. 2020d). The duty factor for at least one detector being online was 97%; for any two detectors being online at the same time it was 82%; and for all three detectors together it was 45%. Further details regarding instrument performance and data quality for O3a are available in Abbott et al. (2020d); Buikema et al. (2020).

The LIGO and Virgo detectors used a photon recoil based calibration (Karki et al. 2016; Cahillane et al. 2017; Viets et al. 2018) resulting in a complex-valued, frequency-dependent detector response with typical errors in magnitude of 7% and 4 degrees in phase (Sun et al. 2020; Acernese et al. 2018) in the calibrated O3a strain data.

Transient noise sources, referred to as glitches, contaminate the data and can affect the confidence of candidate detections. Times affected by glitches are identified so that searches for GW events can exclude (veto) these periods of poor data quality (Abbott et al. 2016a, 2020a; Davis et al. 2021; Nguyen et al. 2021; Fiori et al. 2020). In addition, several known noise sources are subtracted from the data using information from witness auxiliary sensors (Driggers et al. 2019; Davis et al. 2019).

Candidate events, including those reported in Abbott et al. (2020d) and the new candidates found by the searches for sub-threshold counterpart images in Sec. 5.3 of this paper, have undergone a validation process to evaluate if instrumental artifacts could affect the analysis; this process is described in detail in Sec. 5.5 of Davis et al. (2021). This process can also identify data quality issues that need further mitigation for individual events, such as subtraction of glitches (Cornish & Littenberg 2015) and non-stationary noise couplings (Vajente et al. 2020), before executing PE algorithms. See Table V of Abbott et al. (2020d) for the list of events requiring such mitigation.

The GWTC-2 catalog (Abbott et al. 2020d) contains 39 events from O3a (in addition to the 11 previous events from O1 and O2) with a false-alarm rate (FAR) below two per year, with an expected contamination fraction of less than 10%. We neglect the potential contamination in this analysis. These events were identified by three search pipelines: one minimally modeled transient search cWB (Klimenko et al. 2004, 2005, 2006, 2011, 2016) and the two matched-filter searches GstLAL (Sachdev et al. 2019; Hanna et al. 2020; Messick et al. 2017) and PyCBC (Allen et al. 2012b; Allen 2005; Dal Canton et al. 2014; Usman et al. 2016; Nitz et al. 2017). Their parameters were estimated through Bayesian inference using the LALINFERENCE (Veitch et al. 2015) and Bilby (Ashton et al. 2019; Smith et al. 2020; Romero-Shaw et al. 2020) packages. Both the matched-filter searches and PE use a variety of CBC waveform models which gener-

ally combine knowledge from post-Newtonian theory, the effective-one-body formalism and numerical relativity (for general introductions to these approaches, see [Blanchet 2014](#); [Damour & Nagar 2016](#); [Palenzuela 2020](#); [Schmidt 2020](#) and references therein). The analyses in this paper rely on the same methods, and the specific waveform models and analysis packages used are described in each section.

Most of the 39 events from O3a are most probably BBHs, while three (GW190425, GW190426.152155, and GW190814) have component masses below $3 M_{\odot}$ ([Abbott et al. 2020b,d,e](#)), thus potentially containing a neutron star. We consider these 39 events in most of the analyses in this paper, except in the magnification analysis (Sec. 4), which concerns only six of the more unusual events, and the microlensing analysis (Sec. 6), which focuses on the 36 clear BBH events only.

Specifically, we use the following input data sets for each analysis. The magnification analysis in Sec. 4 and posterior-overlap analysis in Sec. 5.1 start from the Bayesian inference posterior samples released with GWTC-2 ([GWOSC 2020](#)). The joint-PE analyses in Sec. 5.2 and microlensing analysis in Sec. 6 reanalyze the strain data in short segments around the event times, available from the same data release, with data selection and noise mitigation choices matching those of the PE analyses in [Abbott et al. \(2020d\)](#). In addition, the searches for sub-threshold counterpart images in Sec. 5.3 cover the whole O3a strain data set, using the same data quality veto choices as in [Abbott et al. \(2020d\)](#) but a strain data set consistent with the PE analyses: the final calibration version of LIGO data ([Sun et al. 2020](#)) with additional noise subtraction ([Vajente et al. 2020](#)).

3. LENSING STATISTICS

In this section, we first forecast the number of detectable strongly lensed events (Sec. 3.1). Then, we infer upper limits on the rate of strongly lensed events using two different methods; the first uses only the non-detection of resolvable strongly lensed BBH events (Sec. 3.2), while the second leverages additionally the non-observation of the SGWB (Sec. 3.3) ([Callister et al. 2020](#); [Abbott et al. 2021b](#)). Since the background would originate from higher redshifts, this second method complements the first method.

Throughout this section, we model the mass distribution of BBHs following the results for the POWER LAW + PEAK MODEL of [Abbott et al. \(2020f\)](#). We consider two distinct models of the BBH merger rate density. Model A brackets most of the population synthesis results ([Eldridge et al. 2019](#); [Neijssel et al. 2019](#); [Boco et al. 2019](#); [Santoliquido et al. 2021](#)) corresponding to Population I and II stars while Model B assumes the Madau–Dickinson ansatz ([Madau & Dickinson 2014](#)) where the rate peaks at a particular redshift. For consistency with previous analyses (e.g., [Abbott et al. 2021b](#)),

we take the Hubble constant from Planck 2015 observations to be $H_0 = 67.9 \text{ km s}^{-1} \text{ Mpc}^{-1}$ ([Ade et al. 2016](#)). Detailed discussion on both models and their respective parametrization is given in Appendix A. The obtained rates are subject to uncertainty because of their dependence on the merger rate density, which is model-dependent and only partially constrained. They are nevertheless vital to interpreting our search results in later sections (see Sec. 5).

3.1. Strong lensing rate

We predict the rate of lensing using the standard methods outlined in the literature ([Ng et al. 2018](#); [Li et al. 2018](#); [Oguri 2018](#)), at galaxy and galaxy-cluster lens mass scales. To model the lens population, we need to choose a density profile and a mass function. We adopt the singular isothermal sphere (SIS) density profile for both galaxies and clusters. Moreover, we use the velocity dispersion function (VDF) from the Sloan Digital Sky Survey (SDSS; [Choi et al. 2007](#)) for galaxies and the halo mass function from [Tinker et al. \(2008\)](#) for clusters which have been used in other lensing studies as well (e.g., [Oguri & Marshall 2010](#); [Robertson et al. 2020](#)). The SIS profile can well describe galaxies. However, the mass distribution of clusters tends to be more complicated. Nevertheless, [Robertson et al. \(2020\)](#) have demonstrated that the SIS model can reproduce the lensing rate predictions from a study of numerically simulated cluster lenses. Thus, we adopt the same model. Under the SIS model, we obtain two images with different magnifications and arrival times. The rate of strong lensing is

$$\mathcal{R}_{\text{lens}} = \int \frac{dN(M_h, z_l)}{dM_h} \frac{dD_c}{dz_l} \frac{\mathcal{R}_m(z_m)}{1+z_m} \frac{dV_c}{dz_m} \sigma(M_h, z_l, z_m, \rho, \rho_c) \times p(\rho|z_m) d\rho dz_m dz_l dM_h, \quad (1)$$

where $dN(M_h, z_l)/dM_h$ is the differential comoving number density of lensing halos in a halo mass shell at lens redshift z_l , D_c and V_c are the comoving distance and volume, respectively, at a given redshift, $\mathcal{R}_m(z_m)$ is the total merger rate density at redshift z_m , $p(\rho | z_m)$ is the distribution of signal-to-noise ratio (SNR) at a given redshift, and σ is the lensing cross-section (Appendix A). Throughout this section and in Sec. 3.3 we choose a network SNR threshold $\rho_c = 8$ as a point estimator of the detectability of GW signals. We find it to be consistent with the search results in [Abbott et al. \(2020d\)](#) and in Sec. 5.3, and we estimate its impact to be subdominant with respect to other source of uncertainties.

In Table 1, we show our estimates of the relative rate of lensing assuming different models (Models A and B) for the merger rate density. The results are shown separately for galaxy-scale (G) and cluster-scale (C) lenses. Furthermore, these rates are calculated for events that are doubly lensed and for two cases: when only a single event (i.e., the brighter

Table 1. Expected fractional rates of observable lensed double events at current LIGO–Virgo sensitivity.

Merger Rate Density Model	Galaxies		Galaxy Clusters	
	R_D	R_S	R_D	R_S
A	$0.8\text{--}4.3 \times 10^{-4}$	$2.6\text{--}8.6 \times 10^{-4}$	$0.4\text{--}1.7 \times 10^{-4}$	$1.3\text{--}3.7 \times 10^{-4}$
B	$0.5\text{--}14.0 \times 10^{-4}$	$1.2\text{--}24.0 \times 10^{-4}$	$0.4\text{--}7.0 \times 10^{-4}$	$0.8\text{--}11.0 \times 10^{-4}$

NOTE—This table lists the relative rates of lensed double events expected to be observed by LIGO–Virgo at the current sensitivity where both of the lensed events are detected (R_D) and only one of the lensed events is detected (R_S). For Model A, the range corresponds to the bracketing function (see Appendix A) and for Model B, the rates encompass a 90 per cent credible interval. We show the rate of lensing by galaxies ($\sigma_{\text{vd}} = 10\text{--}300 \text{ km s}^{-1}$) and galaxy clusters ($\log_{10}(M_{\text{halo}}/M_{\odot}) \sim 14\text{--}16$) separately. Besides their usage for forecasts, the fraction of lensed events allows us to interpret the prior probability of the strong lensing hypothesis, which we require to identify lensed events confidently.

one) is detected (S), and when both of the doubly lensed events are detected (D). The expected fractional rate of lensing (lensed to unlensed rate), which will be necessary for the multi-image analyses (Sec. 5), ranges from $\mathcal{O}(10^{-3}\text{--}10^{-4})$, depending on the merger rate density assumed. We estimate the fractional rate of observed double (single) events for galaxy-scale lenses in the range $0.8\text{--}4.3 \times 10^{-4}$ ($2.6\text{--}8.6 \times 10^{-4}$) when using Model A for the merger rate density. Similarly, for cluster-scale lenses, the fractional rate is estimated to be in the range of $0.4\text{--}1.7 \times 10^{-4}$ ($1.3\text{--}3.7 \times 10^{-4}$) much rarer than the rates at galaxy scales. These estimates suggest that observing a lensed double image is unlikely at the current sensitivity of the LIGO–Virgo network of detectors. Nevertheless, at design sensitivity and with future upgrades, standard forecasts suggest that the possibility of observing such events might become significant (Ng et al. 2018; Li et al. 2018; Oguri 2018). The main uncertainty in the rate estimates derives from the uncertainties in the merger-rate density at high redshift.

Depending on the specific distribution of lenses and the source population, the time delays between images can change. Models favoring galaxy lensing produce minutes to perhaps months of time delay, while galaxy cluster lensing can produce time delays up to even years. However, the time delay distribution for galaxy cluster lenses is more difficult to model accurately, owing to the more complex lensing morphology.

Since the merger rate density at high redshift is observationally constrained only by the absence of the SGWB, these rates are subject to uncertainty. Nevertheless, standard theoretical models will still produce useful forecasts. We will later refer to these rate estimates in the relevant sections (see Sec. 5).

3.2. Implications from the non-observation of strongly lensed events

Motivated by the absence of evidence for strong lensing (Sec. 5), we assume that no strong lensing has occurred, in order to constrain the merger rate density at high redshift. We use the standard constraints on the merger rate density at low redshift from the LIGO–Virgo population studies (Abbott et al. 2020f). We assume the Madau–Dickinson form for the merger rate density (Model B). This model’s free parameters include




Figure 1. Merger rate density as a function of redshift based on the GWTC-2 results without lensing constraints (blue) and with lensing (red) included in LIGO–Virgo detections. We show the results for galaxy-scale lenses (G) and cluster-scale lenses (C) separately. Furthermore, S (or D) correspond to doubly lensed events where single (or double) events are detected. Because lensed detections occur at higher redshifts than unlensed events, their non-observation can be used to constrain mergers at higher redshifts.

the local merger rate density, the merger rate density peak, and the power-law slope. As illustrated in Fig. 1, we find that the non-observation of lensing sets independent constraints on the merger rate density at high redshift, where current limits are obtained through SGWB non-observation (Abbott et al. 2021b).

3.3. Constraints from stochastic background

We can also constrain the redshift evolution of the merger rate density from the reported non-observation of the SGWB from BBHs (Callister et al. 2020; Abbott et al. 2021b). This, in turn, provides constraints on the relative abundance of distant mergers, which are more likely to undergo lensing. Thus, the non-observation of the SGWB can inform the estimate of the probability of observing lensed BBH mergers (Buscicchio et al. 2020a; Mukherjee et al. 2021).

Following Buscicchio et al. (2020a), we forecast constraints on the merger rate density in O3 using up-to-date constraints




Figure 2. Survival function for the lensing probability of detectable BBH mergers, constrained by the non-observation of the SGWB. The solid line shows the value obtained from the median BBH merger rate density posterior. The shaded region corresponds to the 90% credible interval. Fewer than 1 in 10^3 events are expected to be lensed with magnification $\mu > 2$, on average. Significantly higher magnifications (e.g., $\mu > 30$) are suppressed by a further factor of 10.

on the mass distribution and redshift evolution of BBH mergers obtained from the latest detections (Abbott et al. 2019a,b, 2020d,f), as well as those inferred from current upper limits on the SGWB, given its non-observation (Abbott et al. 2021b).

While the measured parameters for each merger (redshifts, source masses) are potentially biased by lensing, as discussed in Sec. 4, we express all quantities as functions of non-biased merger redshift z_m and chirp mass \mathcal{M} (Buscicchio et al. 2020a) for consistency with other sections. We model the differential lensing probability following Dai et al. (2017). The differential merger rate in a redshift and magnification shell is

$$\frac{d^2\mathcal{R}}{dz_m d \ln \mu} = \frac{dP(\mu | z_m)}{d \ln \mu} \frac{4\pi D_c^2(z_m)}{H_0(1+z_m)E(z_m)} \times \int dm_1 dm_2 \frac{d^3\mathcal{R}_m(z_m)}{dm_1 dm_2 dz_m} p(\rho > \rho_c | m_1, m_2, z_m, \mu), \quad (2)$$

where $d^3\mathcal{R}_m(z_m)/dm_1 dm_2 dz_m$ is the differential merger rate density; $p(\rho > \rho_c | m_1, m_2, z_m, \mu)$ provides the probability of observing mergers with source masses m_1, m_2 , redshift z_m , and magnified by a factor μ above a fixed network SNR threshold $\rho_c = 8$, integrated over the population distribution of source parameters; the factor $4\pi D_c^2(z_m)/[H_0(1+z_m)E(z_m)]$ gives the comoving volume of a redshift shell in an expanding Universe (taking into account the redshifted rate definition with respect to the source frame); and $dP(\mu | z_m)/(d \ln \mu)$ is the lensing probability. We then integrate the differential merger rate (Eq. 2) over redshift and magnifications in $[\mu, \mu_{\max}]$ to obtain the survival function for μ at any redshift. We show the result in Fig. 2. We find the observation of lensed events to be unlikely, with the fractional rate at $\mu > 2$ being $3.3^{+0.9}_{-0.7} \times 10^{-4}$. More significantly magnified events are even more suppressed,

with a rate of $2.7^{+0.2}_{-0.1} \times 10^{-5}$ at $\mu > 30$. These estimates suggest that most binary mergers that we observe are not strongly lensed. However, as projected in Buscicchio et al. (2020a), at design sensitivity, the same probability might be significantly enhanced, as a widened horizon will probe the merger rate density deeper in redshift.

Comparing the above predictions with those obtained for single lensed detections with Model B in Sec. 3.1, we observe decreases of about a factor of 5 for the lensing rate. These decreases are due to the different assumptions on the underlying lensing model.

4. ANALYZING HIGH-MASS EVENTS

If a GW signal is strongly lensed, it will receive a magnification μ defined such that the GW amplitude increases by a factor $|\mu|^{1/2}$ relative to an unlensed signal. The luminosity distance inferred from the GW observation will be degenerate with the magnification such that the inferred luminosity distance

$$D_L^{\text{inferred}} = \frac{D_L}{\sqrt{|\mu|}}. \quad (3)$$

Because of this degeneracy, lensing biases the inferred redshift and thus the source masses. Consequently, the binary appears to be closer than it truly is, and it appears to be more massive than it truly is.

Broadhurst et al. (2018, 2020a,b) argued that some of the relatively high-mass LIGO–Virgo events could be strongly lensed GWs from the lower-mass stellar black hole population observed in the electromagnetic bands. However, the expected strong lensing rates and the current estimates of the merger-rate density disfavor this interpretation (Dai et al. 2017; Ng et al. 2018; Li et al. 2018; Oguri 2018; Hannuksela et al. 2019; Buscicchio et al. 2020a,b) compared to the standard interpretation of a genuine unlensed high-mass population (Abbott et al. 2019a; Roulet et al. 2020; Abbott et al. 2020f; Kimball et al. 2020). Hence, in the absence of more direct evidence, such as identifying multiple images within LIGO–Virgo data (Sec. 5), it is difficult to support the lensing hypothesis purely based on magnification considerations. Nevertheless, it is informative to analyze the degree to which the lensed interpretation would change our understanding of the observed sources.

Under the strong lensing hypothesis \mathcal{H}_{SL} , the GW would originate from a well-known, intrinsically lower-mass population, and the LIGO–Virgo observations have been biased by lensing. Using such a mass prior, we infer the required magnification and corrected redshift and component masses under \mathcal{H}_{SL} . The posterior distribution of the parameters is (Pang et al. 2020)

$$p(\mu, \vartheta | d, \mathcal{H}_{\text{SL}}) \propto p(d | \vartheta) p(\vartheta | \mu, \mathcal{H}_{\text{SL}}) p(\mu | \mathcal{H}_{\text{SL}}), \quad (4)$$

where we distinguish the *apparent* parameters of the waveform received at the detector ϑ , which differ from the intrinsic

Table 2. Inferred properties of selected O3a events under the lensing magnification hypothesis.

Event name	m_1 [M_\odot]	m_2 [M_\odot]	z	μ	τ
GW190425	$1.42^{+0.16}_{-0.12}$	$1.27^{+0.12}_{-0.15}$	$0.3^{+0.1}_{-0.1}$	68^{+163}_{-44}	$6^{+8}_{-4} \times 10^{-6}$
GW190426.152155	$1.89^{+0.40}_{-0.55}$	$0.90^{+0.25}_{-0.40}$	$1.3^{+0.5}_{-0.2}$	497^{+452}_{-272}	$2^{+2}_{-0} \times 10^{-4}$

Event name	m_1^{50} (m_1^{65}) [M_\odot]	m_2^{50} (m_2^{65}) [M_\odot]	z^{50} (z^{65})	μ^{50} (μ^{65})	τ^{50} (τ^{65})
GW190521	43^{+6}_{-16} (55^{+9}_{-22})	36^{+10}_{-15} (45^{+13}_{-19})	$2.5^{+2.1}_{-0.7}$ ($1.8^{+1.7}_{-0.5}$)	13^{+55}_{-8} (6^{+28}_{-4})	9^{+10}_{-3} (5^{+9}_{-2}) $\times 10^{-4}$
GW190602.175927	42^{+7}_{-17} (48^{+14}_{-19})	31^{+13}_{-16} (33^{+15}_{-16})	$1.4^{+1.5}_{-0.5}$ ($1.1^{+1.4}_{-0.4}$)	10^{+65}_{-7} (6^{+46}_{-4})	2^{+8}_{-1} (2^{+7}_{-1}) $\times 10^{-4}$
GW190706.222641	39^{+10}_{-15} (42^{+17}_{-17})	29^{+12}_{-13} (29^{+13}_{-13})	$1.7^{+1.8}_{-0.5}$ ($1.6^{+1.7}_{-0.6}$)	5^{+26}_{-3} (4^{+22}_{-2})	5^{+9}_{-3} (4^{+9}_{-2}) $\times 10^{-4}$

NOTE—Under the hypothesis that the listed events are lensed signals from intrinsically lower-mass binary populations with $\mu > 2$, this table lists the favored source masses, redshifts, magnifications, and estimated optical depths for the BNS and neutron star–black hole (NSBH) (top) and BBH (bottom) high-mass events. For the BBHs, two sets of numbers are given for different assumptions about the edge of the pair instability supernova (PISN) mass gap (a cut at $50 M_\odot$ and $65 M_\odot$). For the BNSs, we presume that they originate from the Galactic BNS population. To interpret the heavy BBHs as lensed signals originating from the assumed lower-mass population, they should be magnified at a moderate magnification $\mu \sim O(10)$ at $z \sim 1$ to 2 . The BNS and NSBH events would require extreme magnifications.

parameters θ due to bias by lensing magnification. Therefore, we can compute the magnification posterior and other parameters by simply re-weighting existing posteriors.

Studies along these lines were already done for the GW190425 BNS event by [Pang et al. \(2020\)](#) and for the GW190521 BBH event in [Abbott et al. \(2020g\)](#). Here we extend the approach to cover additional interesting O3a events, focusing on two cases: (i) the (apparently) most massive observed BBHs, and (ii) sources with an (apparent) heavy neutron star component. In the BBH case, we take the prior over component masses, m_1 and m_2 , and redshift, z of the source $p(m_1, m_2, z)$ from the power-law BBH population model used in [Abbott et al. \(2019a\)](#) for O1 and O2 observations, with a mass power-law index $\alpha = 1$, mass ratio power-law index $\beta_q = 0$, and minimum component mass $m_{\min} = 5 M_\odot$, and assume an absence of BBHs above the pair instability supernova (PISN) mass gap. As in the previous GW190521 study ([Abbott et al. 2020g](#)), we consider two different values to account for uncertainties on the edge of the PISN gap, $m_{\max} = (50, 65) M_\odot$. Such a simple model is adequate for this analysis because our analysis results are most sensitive to the mass cut (highest masses allowed by the prior) and less sensitive to the specific shape of the mass distribution. For events with an apparent heavy neutron star component, we assume a Galactic BNS prior following a total mass with a $2.69 M_\odot$ mean and $0.12 M_\odot$ standard deviation ([Farrow et al. 2019](#)). In both cases, the magnification could explain the apparent high mass of the events from the LIGO–Virgo observations.

We assume that the redshift prior $p(z) \propto \tau(z) dV_c/dz$, where the optical depth of lensing by galaxies or galaxy clusters $\tau(z) \propto D_c(z)^3$ ([Haris et al. 2018](#)). The redshift dependence of the optical depth is approximately the same for both galaxies and galaxy clusters, while the overall scaling can

change ([Fukugita & Turner 1991](#)). We use the lensing prior $p(\mu|\mathcal{H}_{\text{SL}}) \propto \mu^{-3}$ ([Blandford & Narayan 1986](#)) with a lower limit $\mu > 2$ appropriate to strong lensing ([Ng et al. 2018](#)). This prior is appropriate when we are in the high-magnification, strong lensing limit, i.e. assuming that the observed masses are highly biased. We do not consider weak lensing, which does not produce multiple images and has no known prediction that we could test.

We analyze all O3a BBH events with the primary mass above $50 M_\odot$ at 90% probability using the Bayesian inference posterior samples released with GWTC-2 ([GWOSC 2020](#); [Abbott et al. 2020d](#)). Moreover, we analyze GW190425, a high-mass BNS ([Abbott et al. 2020b](#)), and GW190426.152155, a low-significance potential neutron star–black hole (NSBH) event ([Abbott et al. 2020d](#)) which was investigated as a possible lensed BNS event ([Smith et al. 2019](#)). We use the results for the IMRPHENOMPv2 waveform ([Hannam et al. 2014](#); [Bohé et al. 2016](#)) for most of the events. For GW190521, where higher-order multipole moments are important to include in the analysis ([Abbott et al. 2020g](#)), we adopt the NRSUR7DQ4 waveform ([Varma et al. 2019](#)) results as in [Abbott et al. \(2020h\)](#). Furthermore, for GW190425 ([Abbott et al. 2020b](#)), we use the IMRPHENOMPv2_NRTIDAL ([Dietrich et al. 2019](#)) low-spin samples. Results are summarized in Table 2.

To interpret the heavy BBHs as lensed signals originating from the assumed lower-mass population, they should be magnified at a moderate magnification $\mu \sim 10$ at $z \sim 1$ – 2 . Depending on the lens model, this magnification may imply a moderate chance of an observable multi-image counterpart as events closer to the caustic curves experience more substantial magnifications. Consequently, they often produce events with similar magnification ratios and shorter time delays (comparable magnifications and shorter time delays can be derived from the lens’s symmetry, although if weak or

microlensing is present, the magnifications can differ even in the high-magnification limit). However, we could not identify any multi-image counterparts for any of the high-mass events in our multiple image search (Sec. 5).

The BNS and NSBH events, on the other hand, would require extreme magnifications (68_{-44}^{+163} and 497_{-272}^{+452} , respectively) to be consistent with the Galactic BNS distribution. At these magnifications, we would expect the source to be close to a caustic, and therefore it may be possible that the presence of microlenses would produce observable effects (Diego et al. 2019; Diego 2020; Pagano et al. 2020). Moreover, the event would likely be multiply imaged (Blandford & Narayan 1986; Oguri 2018). A more detailed follow-up study to quantify the likelihood of multiple images and microlensing could produce more stringent evidence for the lensing hypothesis for these events. We will briefly comment on these events in the context of multi-image and microlensing results in the sections that follow.

At this stage, we cannot set robust constraints on the lensing hypothesis based on the magnification alone. Moreover, as detailed in the following section, we have also not found any other clear evidence to indicate that these GW events are lensed. The prior lensing rate and the optical depth disfavor the lensing hypothesis for most standard binary population and lens models, as discussed in Sec. 3. However, if other BBH formation channels exist that produce an extensive number of mergers at high redshift, the lensing rates can change. In the future, more quantitative constraints could be set by connecting the inferred magnifications with lens modeling to make predictions for the appearance of multiple images or microlensing effects.

5. SEARCH FOR MULTIPLE IMAGES

In addition to magnification, strong lensing can produce multiple images of a single astrophysical event. These multiple images appear at the GW detectors as repeated events. The images will differ in their arrival time and amplitude (Wang et al. 1996; Haris et al. 2018; Hannuksela et al. 2019; Li et al. 2019a; McIsaac et al. 2020). The sky location is the same within the localization accuracy of GW detectors, given that the typical angular separations are of the order of arcseconds. Additionally, lensing can invert or Hilbert transform the image (Dai & Venumadhav 2017; Ezquiaga et al. 2021), introducing a frequency-independent phase shift. This transformation depends on the image type, set by the lensing time delay at the image position: Type-I, II, and III correspond to a time-delay minimum, saddle point, and maximum, respectively (Ezquiaga et al. 2021).

The multiply imaged waveforms $\{\tilde{h}_j^L\}$ of a single signal \tilde{h} then satisfy (Dai & Venumadhav 2017; Ezquiaga et al. 2021)

$$\tilde{h}_j^L(f; \theta, \mu_j, \Delta t_j, \Delta \phi_j) = \sqrt{|\mu_j|} \tilde{h}(f; \theta, \Delta t_j) \exp(i \text{sign}(f) \Delta \phi_j), \quad (5)$$

where $\sqrt{|\mu_j|}$ is the lensing magnification experienced by the image j and $\Delta \phi_j = -\pi n_j/2$ is the Morse phase, with index $n_j = 0, 1, 2$ for Type-I, II, and III images. $\tilde{h}(f; \theta, \Delta t_j)$ is the original (unlensed) waveform before lensing, but evaluated as arriving with a time delay Δt_j . The multi-image hypothesis then states that most parameters measured from the different lensed images of the same event are consistent.

The relative importance of different parameters for the overall consistency under the multi-image hypothesis will vary for different events. For example, the sky localization match will have greater relevance for well-localized, high-SNR events. Similarly, the overlap in measured chirp mass $(1+z)\mathcal{M} = (1+z)(m_1 m_2)^{3/5}/(m_1 + m_2)^{1/5}$ will be more significant when the uncertainty in that parameter is lower, although in this case the underlying astrophysical mass distribution will play a key role. The similarities in other parameters such as mass ratios or spins will be more important when they depart from the more common astrophysical expectations. Evidence of strong lensing could also be acquired with a single Type-II (saddle point) image if the induced waveform distortions in the presence of higher modes, precession, or eccentricity are observed (Ezquiaga et al. 2021). Such evidence is unlikely to be observed without next-generation detectors (Wang et al. 2021).

In this section, we perform three distinct but related analyses. First, we test the lensed multi-image hypothesis by analyzing, for all pairs of O3a events from GWTC-2, the overlap of posterior distributions previously inferred for the individual events. This allows us to set ranking statistics to identify an initial set of candidates for lensed multiple images. We perform a more detailed joint-PE analysis for these most promising pairs, considering all potential correlations in the full parameter space and the image type. This joint analysis provides a more solid determination of the lensing probability for a given GW pair. Finally, we search for additional sub-threshold candidates that could be multiply imaged counterparts to the previously considered events: some counterpart images can have lower relative magnification compared with the primary image and/or fall in times of worse detector sensitivity or antenna patterns, and hence may not have passed the detection threshold of the original broad searches.

Previous studies have also searched for multiple images in the O1–O2 catalog GWTC-1 (Hannuksela et al. 2019; Broadhurst et al. 2019; Li et al. 2019a; McIsaac et al. 2020; Dai et al. 2020; Liu et al. 2021). The first search for GW lensing signatures in O1 and O2 focused on the posterior overlap of the masses, spins, binary orientation and sky positions (Hannuksela et al. 2019) and the consistency of time delays with expectations for galaxy lenses, but found no conclusive evidence of lensing. The search did uncover a candidate pair GW170104–GW170814 with a relatively high Bayes factor of $\gtrsim 200$. Still, this study disfavored the candidate due to its

long time delay and the low prior probability of lensing. In parallel, [Broadhurst et al. \(2019\)](#) suggested that the candidate pair GW170809–GW170814 could be lensed, but this claim is disfavored by more comprehensive analyses ([Hannuksela et al. 2019](#); [Liu et al. 2021](#)). Both [Li et al. \(2019a\)](#) and [McIsaac et al. \(2020\)](#) performed searches for sub-threshold counterparts to the GWTC-1 events, identifying some marginal candidates but finding no conclusive evidence of lensing. More recently, [Dai et al. \(2020\)](#) and [Liu et al. \(2021\)](#) searched for lensed GW signals including the analysis of the lensing image type, which can be described through the Morse phases, $\Delta\phi_j$ in Eq. (5). These analyses have revisited the pair GW170104–GW170814 and demonstrated that the Morse phase is consistent with the lensed expectation but would require Type-III (time-delay maximum) images, which are rare from an observational standpoint. [Dai et al. \(2020\)](#) also pointed out that a sub-threshold trigger, designated by them as GWC170620, is also consistent with coming from the same source. However, the required number and type of images for this lens system make the interpretation unlikely given current astrophysical expectations. Also, two same-day O3a event pairs (on 2019 May 21 and 2019 August 28) have already been considered elsewhere, but were both ruled out due to vanishing localization overlap ([Singer et al. 2019](#); [Abbott et al. 2020g](#)).

5.1. Posterior-overlap analysis

As a consequence of degeneracies in the measurements of parameters, the lensing magnification can be absorbed into the luminosity distance (Sec. 4), the time delay can be absorbed into the time of coalescence, and, when the radiation is dominated by $\ell = |m| = 2$ multipole moments, the phase shifts introduced by lensing (the Morse phases) can be absorbed into the phase of coalescence. The multi-image hypothesis then states that all other parameters except the arrival time, luminosity distance, and coalescence phase are the same between lensed events, and thus there should be extensive overlap in their posterior distributions, even if those have been inferred without taking lensing into account.

Therefore, we use the consistency of GW signals detected by LIGO and Virgo to identify potential lensed pairs. Following [Haris et al. \(2018\)](#), we define a ranking statistic $\mathcal{B}^{\text{overlap}}$ to distinguish candidate lensed pairs from unrelated signals:

$$\mathcal{B}^{\text{overlap}} = \int d\Theta \frac{p(\Theta|d_1) p(\Theta|d_2)}{p(\Theta)}, \quad (6)$$

where the parameters Θ include the redshifted masses $(1+z)m_{1,2}$, the dimensionless spin magnitudes $\chi_{1,2}$, the cosine of spin tilt angles $\theta_{1,2}$, the sky location $(\alpha, \sin\delta)$, and the cosine of orbital inclination θ_{JN} , but they do not include the full 15-dimensional set of parameters Θ to ensure the accuracy of the kernel density estimators (KDEs) that we use to approximate the posterior distributions $p(\Theta|d_{1,2})$ for each event

when evaluating Eq. (6). Here, $p(\Theta)$ denotes the prior on Θ . We use posterior samples (GWOSC 2020) obtained using the LALINFERENCE software package ([Veitch et al. 2015](#)) with the IMRPHENOMPv2 waveform model ([Hannam et al. 2014](#); [Bohé et al. 2016](#)) for most of the events. However, for GW190521, we use NRSUR7DQ4 ([Varma et al. 2019](#)) posteriors, and for GW190412 and GW190814 we use IMRPHENOMPv3HM ([Khan et al. 2020](#)) posteriors. The prior $p(\Theta)$ is chosen to be uniform in all parameters. The component mass priors have the bound $(2-200 M_\odot)$. Equation (6) then quantifies how consistent a given event pair is with being lensed. In our analysis, we omit the BNS event GW190425 ([Abbott et al. 2020b](#)) because it was detected at relatively low redshift, and hence we expect the probability of it being lensed to be very small.

In addition to the consistency of the frequency profile of the signals (as measured by the posterior overlap), the expected time delays Δt between lensed images follow a different distribution than for pairs of unrelated events. Following [Haris et al. \(2018\)](#), we define

$$\mathcal{R}^{\text{gal}} = \frac{p(\Delta t|\mathcal{H}_{\text{SL}})}{p(\Delta t|\mathcal{H}_{\text{U}})}, \quad (7)$$

where $p(\Delta t|\mathcal{H}_{\text{SL}})$ and $p(\Delta t|\mathcal{H}_{\text{U}})$ are the prior probabilities of the time delay Δt under the strongly lensed and unlensed hypotheses, respectively. Here $p(\Delta t|\mathcal{H}_{\text{U}})$ is obtained by assuming that the GW events follow a Poisson process. We use a numerical fit to the time-delay distribution $p(\Delta t|\mathcal{H}_{\text{SL}})$ obtained in Sec. 3 for the SIS galaxy lens model, with a merger rate density given by R_{min} in Eq. (A1). Equation (7) provides another ranking statistic to test the lensing hypothesis, based on the time delay, though subject to some astrophysical uncertainties (see discussion in Sec. 3). The time-delay distribution does not include galaxy cluster lenses, which may be responsible for long time delays of several months or more. We also do not model detector downtime, but we expect the different contributions to the time delay to average out across a longer time period.

To estimate the significance of the combined ranking statistic, $\log_{10}(\mathcal{B}^{\text{overlap}} \times \mathcal{R}^{\text{gal}})$ computed for O3a event pairs, we perform an injection campaign. For the injection campaign, we sample component masses $m_{1,2}$ from a power-law distribution ([Abbott et al. 2016b](#)) in the range $(10-50 M_\odot)$. We assume that the redshift distribution follows population synthesis simulations of isolated binary evolution ([Belczynski et al. 2008, 2010](#); [Dominik et al. 2013](#); [Marchant et al. 2018](#); [Eldridge et al. 2019](#); [Neijssel et al. 2019](#); [Boco et al. 2019](#); [Santoliquido et al. 2021](#)); in particular, for illustration purposes, we show results using the redshift evolution from [Belczynski et al. \(2016a,b\)](#), but for the local universe that we look at ($z < 2$), other models produce qualitatively similar results. All other parameters are sampled from uninformative prior




Figure 3. Scatter plot of the ranking statistics $\log_{10} \mathcal{B}^{\text{overlap}}$ and $\log_{10} \mathcal{R}^{\text{gal}}$ for a subset of event pairs that have both $\mathcal{B}^{\text{overlap}} > 50$ and $\mathcal{R}^{\text{gal}} > 0.01$. The dashed lines denote the significance levels of the combined ranking statistics (in terms of Gaussian standard deviations), obtained by simulating unlensed event pairs in Gaussian noise matching the O3a sensitivity of the LIGO–Virgo network. We identify several high $\mathcal{B}^{\text{overlap}} > 50$ candidates, which we follow up on with a detailed joint-PE analysis. We have used abbreviated event names, quoting the last 4 digits of the date identifier (see Table 3 for full names).

distributions (Haris et al. 2018). We inject the simulated signals into Gaussian noise with O3a representative spectra for a LIGO–Virgo detector network. The underlying motivation in this step is to identify interesting candidates for joint-PE follow-up, which will give the final analysis verdict.

In Fig. 3 we show the scatter plot of $\log_{10} \mathcal{B}^{\text{overlap}}$ and $\log_{10} \mathcal{R}^{\text{gal}}$ for the O3a event pairs that have high combined ranking statistic. The dashed lines represent different significance levels as obtained from the simulations. The event pair GW190728_064510–GW190930_133541 gives the highest combined ranking statistic, $\log_{10}(\mathcal{B}^{\text{overlap}} \times \mathcal{R}^{\text{gal}}) = 3.6$; however, as can be seen from Fig. 3, its significance is above 1σ (68%) but much below the 2σ (95%) significance level.

To follow up on the most promising event pairs with the more detailed joint-PE analysis in the next section, we make a selection based on just the posterior overlap ranking statistic, $\mathcal{B}^{\text{overlap}}$, rather than the combined ranking statistic, $\mathcal{B}^{\text{overlap}} \times \mathcal{R}^{\text{gal}}$, because \mathcal{R}^{gal} depends strongly on the lens model. Our aim in the next section is to understand the high $\mathcal{B}^{\text{overlap}}$ event pairs in greater detail without resorting to any specific lens model. We thus select the most promising event pairs from Fig. 3, i.e., those with $\mathcal{B}^{\text{overlap}} > 50$, and carry out the joint-PE

analysis in the next section. The 19 selected pairs are listed in Table 3.

5.2. Joint parameter estimation analysis

Here we follow up on the most significant pairs of events from the posterior-overlap analysis with a more detailed but more computationally demanding joint-PE analysis. The benefit of this analysis is that it allows for more stringent constraints on the lensing hypothesis by investigating potential correlations in the full parameter space of BBH signals, instead of marginalizing over some parameters. Moreover, it also includes a test for the lensing image type by incorporating lensing phase information.

We perform our analysis using two independent pipelines, a LALINFERENCE-based pipeline (Liu et al. 2021) and a BILBY-based pipeline (HANABI; Lo & Magaña Hernandez 2021), giving us additional confidence in our results. Unlike the posterior-overlap analysis, the joint-PE analysis does not start from existing posterior samples. Instead, we start the inference directly using the detector strain data. In both pipelines, we follow the same data selection choices (calibration version, available detectors for each event, and noise subtraction procedures) as in the original GWTC-2 analysis (Abbott et al. 2020d), with special noise mitigation steps (glitch subtraction and frequency range limitations) taken for some events as listed in Table V of that paper. However, the two pipelines use different waveform models. In this section, we first describe how we quantify the evidence for the strong lensing hypothesis, then detail the two pipelines and finally present the results.

5.2.1. The coherence ratio and the Bayes factor

There will be three types of outputs for the joint-PE analysis. First, we compute a coherence ratio C_{U}^{L} , which is the ratio of the lensed and unlensed evidences, neglecting selection effects and using default priors in the joint-PE inference. We treat this as a ranking statistic, which quantifies how consistent two signals are with the lensed hypothesis. Large coherence ratios indicate that the parameters of the GWs agree with the expectations of multiple lensed events. This occurs, for example, when the masses and sky localization coincide. However, the coherence ratio does not properly account for the possibility that the parameters overlap by chance.

The likelihood that GW parameters overlap by chance sensitively depends on the underlying population of sources and lenses. For example, if there existed formation channels that produced GWs with similar frequency evolutions (as expected of lensing), the likelihood of an unlensed event mimicking lensing would increase substantially. Thus, we introduce a second output, the population-weighted coherence ratio $C_{\text{U}|\text{pop}}^{\text{L}}$, which incorporates prior information about the populations of BBHs and lenses. The value of $C_{\text{U}|\text{pop}}^{\text{L}}$ is subject to the choice of both the BBH and lens models.

Similarly, the probability that two signals agree with the multiple-image hypothesis is altered through selection effects, as some masses and sky orientations are preferentially detected. Thus, we also include the selection effects, which gives us our final output, the Bayes factor \mathcal{B}_U^L . The \mathcal{B}_U^L quantifies the evidence of the strong lensing hypothesis for a given detector network and population model. For the full derivations and detailed discussion on the difference between the coherence ratio and the Bayes factor, see [Lo & Magaña Hernandez \(2021\)](#).

5.2.2. LALINFERENCE-based pipeline

For the LALINFERENCE-based pipeline, we adopt the method presented by [Liu et al. \(2021\)](#), which was first used for analyzing pairs of events from GWTC-1 ([Abbott et al. 2019b](#)). LALINFERENCE-NEST ([Veitch et al. 2015](#)) implements nested sampling ([Skilling 2006](#)), which can compute evidences without explicitly carrying out the high-dimensional integral while sampling the posteriors. The LALINFERENCE-based pipeline uses the IMRPHENOMD waveform ([Husa et al. 2016](#); [Khan et al. 2016](#)), which is a phenomenological model that includes the inspiral, merger, and ringdown phases but assumes non-precessing binaries and only $\ell = |m| = 2$ multipole radiation. This is motivated by the fact that most events detected so far are well described by the dominant multipole moment ([Abbott et al. 2019b, 2020d](#)). Higher-order multipole moments, precession, or eccentricity could lead to non-trivial changes to the waveform for Type-II images, but such waveforms cannot currently be used with this pipeline. For a discussion of the events within GWTC-2 displaying measurable higher-order multipole moments or precession, see Appendix A of [Abbott et al. \(2020d\)](#).

As in the posterior-overlap analysis, we expect observed, lensed GWs to share the same parameters for the redshifted masses, spins, sky position, polarization angle and inclination, $\{(1+z)m_1, (1+z)m_2, \chi_1, \chi_2, \alpha, \delta, \psi, \theta_{JN}\}$. Hence, we force these parameters to be identical under the lensing hypothesis. For the unlensed hypothesis, we sample independent sets of parameters for each event. This is equivalent to performing two separate nested sampling runs and then combining their evidence. In total, LALINFERENCE samples in an 11-dimensional parameter space and provides C_U^L as the output.

We sample the apparent luminosity distance of the first event D_L^1 and the relative magnification μ_r ([Wang et al. 1996](#)) instead of the luminosity distance of the second event D_L^2 , using the relation $\sqrt{\mu_r} = D_L^1/D_L^2$. Since our waveform only includes the dominant $\ell = |m| = 2$ multipole moments, the lensing Morse phase is modeled by discrete shifts in the coalescence phase ϕ_c by an integer multiple of $\pi/4$ (with relation to the lensing phase shift $\Delta\phi = 2\Delta\phi_c$, [Dai & Venumadhav 2017](#); [Ezquiaga et al. 2021](#)). Thus, we consider all possi-

ble relative shifts $\Delta\phi_c \in \{0, \pi/4, \pi/2, 3\pi/4\}$ between two GW signals.

We set a uniform prior in $\log[(1+z)m_1]$ and $\log[(1+z)m_2]$ for both the lensed and unlensed hypothesis. The minimum and maximum component masses are respectively $3 M_\odot$ and $330 M_\odot$, with a minimum mass ratio of $q = m_2/m_1 = 0.05$. This choice reduces the prior volume by $10^2 - 10^3$ compared to the uniform prior used in GWTC-2. For the other parameters, the prior for the luminosity distance is $p(D_L) \propto D_L^2$ up to 20 Gpc, while the aligned spin magnitudes are uniform between 0 and 1.

5.2.3. The HANABI pipeline

The HANABI pipeline, on the other hand, adopts a hierarchical Bayesian framework that models the data generation process under the lensed and the unlensed hypothesis. This pipeline uses the IMRPHENOMXPHM waveform ([Pratten et al. 2020](#)), which models the full inspiral–merger–ringdown for generic precessing binaries including both the dominant and some sub-dominant multipole moments. Therefore, the parameter space of HANABI enlarges to 15 dimensions.

HANABI differs from the LALINFERENCE-based pipeline in the treatment of the Morse phase. Here the lensing phase is directly incorporated in the frequency-domain waveform, accounting for any possible distortion of Type-II images ([Ezquiaga et al. 2021](#)). Moreover, the lensed probability is computed against the different combinations of image types and, for this reason, HANABI only produces one evidence per pair and not one for each discrete phase difference as the LALINFERENCE-based pipeline. Unlike the LALINFERENCE-based pipeline, HANABI samples the observed masses in a uniform distribution. The mass ranges are different for each event pair, but an overall reweighting is applied later (see below). The rest of the prior choices for the intrinsic parameters are the same as for the LALINFERENCE-based pipeline with the addition of a discrete uniform prior on the Morse phase and isotropic spin priors.

In addition to computing the joint-PE coherence ratio, HANABI also incorporates prior information about the lens and BBH populations, as well as selection effects. In particular, the BBH population is chosen to follow a POWER LAW + PEAK MODEL in the primary mass following the best-fit parameters in [Abbott et al. \(2020f\)](#). Similarly, the secondary mass is fixed to a uniform distribution between the minimum and the primary mass. HANABI also uses an isotropic spin distribution and merger rate history following Model A in Sec. 3. The lens population is modeled by the optical depth described in [Hannuksela et al. \(2019\)](#) and a magnification distribution $p(\mu) \propto \mu^{-3}$ for $\mu \geq 2$. HANABI is thus able to output $C_U^L, C_U^L|_{\text{pop}}$ and \mathcal{B}_U^L . However, HANABI does not include any preference for a particular type of image, i.e., HANABI uses a discrete, uniform prior for the Morse phase shift $\Delta\phi_j$.

Table 3. Summary of joint-PE results for event pairs in O3a.

Event 1	Event 2	$\log_{10} \mathcal{R}^{\text{gal}}$	$\log_{10}(C_{\text{U}}^{\text{L}})$	$\log_{10}(C_{\text{U}}^{\text{L}} _{\text{pop}})$	$\log_{10}(\mathcal{B}_{\text{U}}^{\text{L}})$
			LALINFERENCE ($\Delta\phi$: 0, $\pi/2$, π , $3\pi/2$)	HANABI	HANABI
GW190412	GW190708_232457	-1.7	(+1.0, -9.7, -22.8, -4.4)	-6.6	-9.7
GW190421_213856	GW190910_112807	-	(+ 4.5 , +2.5, -1.5, -0.0)	-0.7	-3.8
GW190424_180648	GW190727_060333	-2.0	(+ 4.9 , +0.0, +1.1, + 4.0)	-0.8	-3.9
GW190424_180648	GW190910_112807	-	(+2.5, + 4.7 , + 4.3 , +1.6)	-0.8	-3.9
GW190513_205428	GW190630_185205	-0.7	(+0.8, + 4.3 , -1.9, -6.5)	-2.4	-5.5
GW190706_222641	GW190719_215514	+0.3	(+2.4, +2.4, -0.0, -0.5)	-0.3	-3.4
GW190707_093326	GW190930_133541	-1.7	(-4.6, -4.3, -3.5, -4.1)	-9.4	-12.5
GW190719_215514	GW190915_235702	-1.0	(+3.5, -2.1, -0.1, + 4.1)	-0.7	-3.8
GW190720_000836	GW190728_064510	+0.5	(-1.4, -0.9, -4.5, -5.4)	-6.7	-9.8
GW190720_000836	GW190930_133541	-1.3	(-3.5, -2.8, -3.9, -3.9)	-9.2	-12.3
GW190728_064510	GW190930_133541	-1.1	(-3.6, -2.5, -3.1, -2.9)	-8.5	-11.6
GW190413_052954	GW190424_180648	+0.4	(+0.6, -0.9, +0.4, -0.0)	-1.6	-4.7
GW190421_213856	GW190731_140936	-2.2	(+3.1, -1.9, +2.5, + 5.2)	-0.2	-3.3
GW190424_180648	GW190521_074359	-0.1	(+1.3, +3.8, +3.7, + 4.4)	-2.0	-5.1
GW190424_180648	GW190803_022701	-2.2	(+ 4.2 , +1.9, +2.6, +3.1)	-1.0	-4.1
GW190727_060333	GW190910_112807	-0.7	(+1.8, +3.3, +3.7, +3.4)	-1.4	-4.5
GW190731_140936	GW190803_022701	+0.9	(+ 4.1 , +3.2, +2.2, +3.4)	-0.9	-4.0
GW190731_140936	GW190910_112807	-0.6	(+0.1, + 4.5 , +0.8, -7.2)	-1.2	-4.3
GW190803_022701	GW190910_112807	-0.5	(+ 4.0 , + 5.5 , + 4.7 , +2.6)	-0.1	-3.2

NOTE— We select those events with posterior overlap ranking statistic larger than 50. For each pair of events presented in the first two columns, the third column lists the time-delay ranking statistic \mathcal{R}^{gal} as described in Section 5.1. The next column gives the coherence ratio of the lensed/unlensed hypothesis C_{U}^{L} obtained with the LALINFERENCE-based pipeline, including the results for the four possible lensing phase difference $\Delta\phi = 2\Delta\phi_c$. We highlight in boldface those pairs with $\log_{10}(C_{\text{U}}^{\text{L}}) > 4$ for at least one Morse phase shift. The fifth and sixth columns correspond to the HANABI results for the population-weighted coherence ratio $C_{\text{U}}^{\text{L}}|_{\text{pop}}$ and the Bayes factor $\mathcal{B}_{\text{U}}^{\text{L}}$. All quantities are given in \log_{10} . All high coherence ratio events display a small Bayes factor when including the population priors and selection effects. For the pairs GW190421_213856–GW190910_112807 and GW190424_180648–GW190910_112807, the time delays between events are larger than what we expect for galaxy lenses in our simulation, and thus $\mathcal{R}^{\text{gal}} = 0$.

5.2.4. Results

Within the O3a events, the LALINFERENCE-based pipeline finds 11 pairs with $\log_{10}(C_{\text{U}}^{\text{L}}) > 4$, indicating high parameter consistency. We have checked that the results of the LALINFERENCE-based pipeline are qualitatively consistent with those from HANABI. This reinforces our previous argument that the shift in the coalescence phase is a good approximate description of the lensing Morse phase given that in the present catalog most events are dominated by the $\ell = |m| = 2$ multipole moments. However, because of the pair-dependent prior choices of HANABI, we do not present its raw C_{U}^{L} results in Table 3.

We then include our prior expectation on the properties of the lensed images (derived from our BBH and lens population priors) and selection effects when computing the population-weighted HANABI coherence ratio and the Bayes factors $\mathcal{B}_{\text{U}}^{\text{L}}$. The results are summarized in Table 3. The event pair GW190728_064510–GW190930_133541, which seemed the most promising from the overlap analysis in Sec. 5.1, is disfavored by both joint-PE pipelines. After the inclusion of the population prior and selection effects, none of the event pairs display a preference for the lens hypothesis ($\log_{10} \mathcal{B}_{\text{U}}^{\text{L}} < 0$).

The population-weighted coherence ratio and the Bayes factor are subject to the BBH and lens model specifications. The population properties are not inferred taking into account the

possibility of lensing. This introduces an inevitable bias, but it can be justified *a posteriori* to be a good approximation given the expected low rate of strong lensing. Additionally, the population properties include significant uncertainties in the hyper-parameter estimates and presume a population model. In any case, to quantify this intrinsic uncertainty in the modeling, we consider different choices for the mass distribution and merger rate history. Varying the maximum BBH mass and the redshift evolution of the merger rate using the $R_{\min}(z)$ and $R_{\max}(z)$ of Model A in Sec. 3, we find that the strong lensing hypothesis is always disfavored. While these results are subject to assumptions on prior choices, our results are sufficient to reject the strong lensing hypothesis: Even if other prior choices favored the lensing hypothesis, the evidence would at best be inconclusive.

The impact of selection effects is considerable. Among other reasons, this is because present GW detectors preferentially observe higher mass events (Fishbach & Holz 2017), making coincidences in observed masses more probable. Along the same lines, given the specific antenna patterns of the current network of detectors, GW events are preferentially seen in specific sky regions with characteristic elongated localization areas (Chen et al. 2017), which favors the overlap between different events.

We also reanalyze the GW170104–GW170814 event pair in the O2 data previously studied by Dai et al. (2020); Liu et al. (2021). Using the LALINFERENCE-based pipeline, Liu et al. (2021) found that the coherence ratio, including selection effects associated with the Malmquist bias (Malmquist 1922), is $\log_{10}(C_{\text{U}}^{\text{L}}) \approx 4.3$ for a $\pi/2$ coalescence phase shift. However, when including together population and selection effects with HANABI, we find that the evidence drastically reduces to a Bayes factor of $\log_{10}(\mathcal{B}_{\text{U}}^{\text{L}}) \approx -2.0$.

In addition to the Bayes factor, it is vital to contrast the recovered number of candidate lensed pairs and their properties with astrophysical expectations. In Sec. 3.1 we found that the relative rate of GW events with at least two strongly lensed images above the detection threshold is below $\sim 1.3 \times 10^{-3}$ for all considered BBH population models. Thus, the lensing rate estimates significantly disfavor the lensing hypothesis *a priori*; even a moderate Bayes factor would not by itself yet make a compelling case for strong lensing. Additionally, the type of images, arrival times, and magnifications provide additional information on the lensing interpretation’s plausibility. For example, a quantification of the time-delay prior can be computed by multiplying the coherence ratio by \mathcal{R}^{gal} .

Although we do not find evidence of strong lensing, future electromagnetic follow-up of the candidates could allow for independent support for the hypothesis if we identified a lensed counterpart galaxy to these events (Sereno et al. 2011; Smith et al. 2018, 2017, 2019; Hannuksela et al. 2020; Robertson et al. 2020; Ryczanowski et al. 2020; Yu et al. 2020).

This identification could take place by matching GW and electromagnetic image properties when four GW images are available (Hannuksela et al. 2020). With two images, the number of hosts could also be constrained (Sereno et al. 2011; Yu et al. 2020), but to a lesser degree due to degeneracies with the lens and source alignment and uncertainties introduced by micro/millilensing – although strong lensing by galaxy clusters might allow us to identify a single cluster candidate (Smith et al. 2018, 2017, 2019; Robertson et al. 2020; Ryczanowski et al. 2020). Moreover, strong lensing could have produced additional images below the noise threshold. We perform a further investigation of such sub-threshold counterparts in the next section.

5.3. Search for sub-threshold lensed images

Here we search for sub-threshold counterpart images of the O3a events from GWTC-2 that would not have been identified as confident detections by the search pipelines used in Abbott et al. (2020d). As lensed images could in principle appear anywhere in the entire O3a data, we perform targeted template bank searches for these sub-threshold lensed counterparts over the whole O3a strain data set, following the data selection criteria described in Abbott et al. (2020d). We employ two matched-filter searches based on the GstLAL (Cannon et al. 2012; Messick et al. 2017; Hanna et al. 2020; Sachdev et al. 2019) and PyCBC (Usman et al. 2016; Nitz et al. 2018, 2019; Davies et al. 2020) pipelines, adapted to the lensing case in similar ways as in Li et al. (2019a) and McIsaac et al. (2020).

5.3.1. Search methods and setups

The lensed hypothesis states that the intrinsic masses and spins will remain consistent between multiple lensed images of the same event. Hence, we can perform searches that specifically target sub-threshold lensed counterparts of known events by creating reduced banks of template waveforms with masses and spins close to those inferred for the primary event. We use the public posterior mass and spin samples released with GWTC-2 (GWOSC 2020) to create these targeted template banks. This ensures that the known events will match well with the templates while simultaneously decreasing the FAR of the search for similar events, potentially returning new candidates that did not reach the search threshold in Abbott et al. (2020d). GstLAL’s reduced banks contain between 173 and 2698 templates per search, while for each PyCBC search we select a single aligned-spin template. The construction of these template banks closely follows Li et al. (2019a); McIsaac et al. (2020) and is further detailed in Appendix B. Template waveforms are generated using the aligned-spin SEOBNRv4_ROM waveform (Bohé et al. 2017; Pürrer 2014; Pürrer 2016) for both pipelines and all events, with the exception of GW190425 in the PyCBC search, where we use the TaylorF2 model (Blanchet et al. 1995; Sathyaprakash & Dhurandhar 1991; Poisson 1998; Damour et al. 2000; Mikoczi

et al. 2005; Blanchet et al. 2005; Arun et al. 2009; Buonanno et al. 2009; Faye et al. 2012; Bohé et al. 2013; Blanchet 2014; Bohé et al. 2015; Mishra et al. 2016).

Given these template banks, each search pipeline proceeds with configurations and procedures as outlined in Abbott et al. (2020d) to produce a priority list of potential lensed candidates matching each target event. To rank these, each pipeline uses a different method to estimate FARs.

GstLAL first identifies matched-filter triggers from one or more of the Hanford, Livingston, and Virgo data streams. Coincidences are identified with the same settings as in Abbott et al. (2020d). From each candidate’s recovered parameters, a likelihood-ratio ranking statistic is computed (Sachdev et al. 2019). Single-detector triggers are penalized using machine-learning based predictions (iDQ; Essick et al. 2020; Godwin et al. 2020) whereas for coincident triggers, no data quality products are used. We estimate the FAR of a trigger by comparing with the distribution of the ranking statistic from all non-coincident noise triggers, used to characterize the noise distribution, over the O3a data set.

PyCBC also first identifies single-detector matched-filter triggers, with a reduced clustering window compared to the GWTC-2 configuration (from 1 s to 0.01 s). These are tested for time coincidence between detectors and are required to have an SNR ≥ 4 in at least two detectors. While in the GWTC-2 analysis the PyCBC search was limited to the Hanford and Livingston detectors, here we also include Virgo data, using the methods described in Davies et al. (2020) to analyse the three detector network. FARs are estimated from a noise background measured using time-shifted data. All triggers within 0.1 s of the times of the events in GWTC-2 are removed from both the foreground (observed coincident events) and the background.

Candidates from both pipelines are further vetted by a sky localization consistency test against the targeted GWTC-2 event, as lensed images of the same event should come from consistent sky locations but the matched-filter searches do not check for this. For each new candidate, we generate a sky localization map $p(\Omega)$ using BAYESTAR (Singer & Price 2016), with Ω denoting parameters that specify the sky location. We compute the percentage overlap $O_{90\%CR}$ of the 90% credible regions between the sky localization $q(\Omega)$ of a GWTC-2 event and the sky localization $p(\Omega)$ of a sub-threshold event candidate as

$$O_{90\%CR} = \frac{\mathbf{1}_{90\%CR} [p(\Omega)q(\Omega)] d\Omega}{\min(\mathbf{1}_{90\%CR} [p(\Omega)] d\Omega, \mathbf{1}_{90\%CR} [q(\Omega)] d\Omega)}, \quad (8)$$

where $\mathbf{1}$ is the indicator function. To avoid false dismissal at this step, we only veto candidates with $O_{90\%CR} = 0$. All candidates with non-vanishing localization overlap are kept for further follow-up with data quality checks as discussed in Sec. 2 and with the joint-PE methods described in Sec. 5.2.

5.3.2. Results

In Table 4, we list the eight candidates with FAR < 1 in 16 years from the *individual* targeted searches for counterparts of the 39 detections reported in GWTC-2 found by at least one pipeline. Six of these are unique newly recovered candidates. This number, compared with ~ 2 expected noise events above this FAR from the number of searches performed, is consistent with additional astrophysical signals being present in the data set. However, in this work, we do not assess in detail the probability of astrophysical origin for each of these. The reported FARs also do not indicate how likely each trigger is to be a lensed counterpart of the targeted event, but only how likely it is to obtain a trigger with a similar ranking statistic from a pure noise background using these reduced template banks.

In contrast, Fig. 4 shows the *combined* search results from all 39 targets for each pipeline: GstLAL (top panel) and PyCBC (bottom panel), excluding triggers that correspond to other detections already reported in GWTC-2. Each panel shows the cumulative number of coincident triggers (observed) with inverse FARs greater than or equal to a given threshold value. For GstLAL, the combined results are obtained by a search over all O3a data using a combined template bank from the 39 targeted banks. For PyCBC, the FARs are obtained from the individual searches, but for triggers being found in several single-template searches, their inverse FARs are summed. In the same figure, we compare these results with estimated background distributions, accounting for the fact that we have re-analyzed the same data set of ~ 150 days multiple times, and find a slight excess in the rate of foreground triggers at high inverse FARs.

Instead, we perform follow-up analyses of the lensing hypothesis under the assumption of astrophysical origin, aiming to determine for each candidate pair in Table 4 whether it is more consistent with a pair of images of a single lensed event or with two independent astrophysical events. After taking into account the initial FAR thresholds, sky localization overlap, and data-quality checks, we have followed up six candidate pairs through LALINFERENCE joint Bayesian PE as described in Sec. 5.2.2. No special mitigation steps were required for data-quality reasons on any of the new candidates. The results are included in Table 4.

Compared with the results for GWTC-2 pairs in Table 3, the LALINFERENCE coherence ratios alone are insufficient to provide evidence of lensing while keeping in mind selection effects and prior odds. As another cross-check, we have also analysed the pair with the highest LALINFERENCE coherence ratio C_U^L (the candidate on 2019 September 16 found by the GW190620_030421 PyCBC search) with the HANABI pipeline described in Sec. 5.2.3. As with all pairs previously tested (see Table 3), after the inclusion of population priors and selection effects, there is no evidence favoring the lensing hypothesis

Table 4. Candidates from sub-threshold searches for strongly-lensed counterpart images of the 39 O3a events from GWTC-2.

UTC time	GWTC-2 targeted event	$ \Delta t $ [d]	$(1+z)\mathcal{M}$ [M_{\odot}]	FAR [yr^{-1}]		$O_{90\%CR}$ [%]	$\log_{10} C_{\text{U}}^{\text{L}}$ (LALINFERENCE) ($\Delta\phi$: 0, $\pi/2$, π , $3\pi/2$)
				PyCBC	GstLAL		
2019 Sep 25 23:28:45	GW190828.065509	28.69	17.3	0.003	98.681	0.0%	–
2019 Apr 26 19:06:42	GW190424.180648	2.04	65.5	–	0.017	63.8%	(–5.8, –5.8, –5.9, –5.6)
2019 Jul 11 03:07:56	GW190421.213856	80.23	47.7	0.032	0.341	1.2%	(+2.3, +1.1, +1.1, +2.6)
2019 Jul 25 17:47:28	GW190728.064510	2.54	9.0	–	0.038	0.0%	–
2019 Jul 11 03:07:56	GW190731.140936	20.46	47.4	0.045	0.944	2.9%	(+2.6, –1.2, –1.6, +0.9)
2019 Aug 05 21:11:37	GW190424.180648	103.13	68.8	–	0.051	26.9%	(–1.1, +0.6, –0.3, –0.7)
2019 Jul 11 03:07:56	GW190909.114149	60.36	49.0	0.053	1.196	12.6%	(+3.5, +2.2, +3.4, +2.9)
2019 Sep 16 20:06:58	GW190620.030421	88.71	53.3	0.055	1.389	49.5%	(+1.7, +3.6, +2.1, –3.2)

NOTE— The first column shows the UTC time of the newly found sub-threshold candidate. The second column lists the targeted O3a event from the catalog GWTC-2; see Table IV and Table VI of [Abbott et al. \(2020d\)](#) for details of these. The third column shows the absolute time difference between the candidate and the targeted event. The fourth column shows the redshifted chirp mass of the template that generated the trigger. The fifth and sixth columns show the corresponding FARs from the individual search for the target from the second column, from each of the two search pipelines (GstLAL and PyCBC), if the candidate has been recovered by it. The seventh column shows the percentage overlap of the 90% sky localization regions between the candidate and the targeted event, from the pipeline with the lower FAR. The eighth column shows the coherence ratio C_{U}^{L} for the pair from the LALINFERENCE joint-PE follow-up (only for candidate pairs with a localization overlap $> 0\%$). Candidates are only reported here if they pass a FAR threshold of < 1 in 16 years in at least one pipeline, and are sorted in ascending order by the lowest FAR from either pipeline.

for this pair either, with population-weighted coherence ratio $\log_{10}(C_{\text{U}}^{\text{L}}|_{\text{pop}}) = -0.1$ and Bayes factor $\log_{10}(\mathcal{B}_{\text{U}}^{\text{L}}) = -3.2$.

As lensing can produce more than two images of the same source, cases where several searches find the same trigger are of particular interest. We find that the same candidate on 2019 July 11 has been found with low FARs by three searches (targeting the GWTC-2 events GW190421.213856, GW190731.140936, and GW190909.114149). In addition, the trigger on 2019 August 05 is only found with sufficient FAR for inclusion in Table 4 by a single GstLAL search (for GW190424.180648), but was also recovered by those for GW190413.052954 and GW190803.022701 with FARs just below the cut. However, the GWTC-2 pairs involved in these possible quadruple sets have already been significantly disfavored by the HANABI analysis including population priors and selection effects. We also expect such multiple matches from an unlensed BBH population due to the clustering of the GWTC-2 events in parameter space ([Abbott et al. 2020d,f](#)).

Also, as discussed in detail in [McIsaac et al. \(2020\)](#), if any high-mass GW detections are interpreted as highly magnified images of lower-mass sources, then counterpart images for these would be more likely. However, we did not find any promising sub-threshold candidates for the five events discussed under the lensing magnification hypothesis in Sec. 4.

In summary, the sub-threshold searches can recover additional promising candidates that were not included in GWTC-2, which match other events closely and, in that sense, are consistent with the lensing hypothesis. However, we do not find sufficient evidence that they are indeed lensed images, as the set of results is also consistent with a population of physically independent and only coincidentally similar events.

6. SEARCH FOR MICROLENSING EFFECTS

Micro lensing by smaller lenses produces image separations on the order of microarcseconds. For GWs, it can also induce frequency-dependent wave-optics effects similar to femtolensing of light ([Nakamura 1998](#); [Takahashi & Nakamura 2003](#)). More specifically, when the characteristic wavelengths are comparable to the Schwarzschild radius of the lens, i.e., $\lambda_{\text{GW}} \sim R_{\text{Sch}}^{\text{lens}}$, it causes frequency-dependent magnification of the waveform. Moreover, the characteristic lensing time-delay due to microlensed images can be shorter than the GW signal duration, causing potentially observable beating patterns on the waveform ([Cao et al. 2014](#); [Jung & Shin 2019](#); [Lai et al. 2018](#); [Christian et al. 2018](#); [Diego et al. 2019](#); [Diego 2020](#); [Pagano et al. 2020](#)), due to waveform superposition. To observe GW microlensing, we search for these beating patterns instead of the time-dependent change in the flux traditionally observed for microlensing in electromagnetic signals.

Here we search for microlensing by isolated point masses. The microlensed waveform has the form

$$h^{\text{ML}}(f; \theta_{\text{ML}}) = h^{\text{U}}(f; \theta) F(f; M_{\text{L}}^z, y), \quad (9)$$

where h^{ML} and h^{U} are the microlensed and unlensed waveforms in the frequency domain, respectively. θ represents the set of parameters defining an unlensed GW signal, while $\theta_{\text{ML}} = \{\theta, M_{\text{L}}^z, y\}$. $F(f; M_{\text{L}}^z, y)$ is the frequency-dependent lensing magnification factor, which is a function of the redshifted lens mass $M_{\text{L}}^z = M_{\text{L}}(1+z_{\text{l}})$ and dimensionless impact parameter y , given in Eq. 2 of ([Lai et al. 2018](#)). The search involves re-estimating the parameters of previously identified events under the microlensed hypothesis as defined in Eq. (9), including those of the potential lens.




Figure 4. Combined results from the 39 sub-threshold searches with the GstLAL pipeline (top panel) and PyCBC pipeline (bottom panel). Each panel shows, as a solid line, the cumulative number of coincident triggers (observed) with inverse FARs greater than or equal to a given value. The dashed line is the expected distribution of background triggers, with the gray bands indicating uncertainties in multiples of the standard deviation σ of a Poisson distribution. For GstLAL, the results for this plot are obtained by a search over all O3a data using a combined bank from the 39 targeted banks. For PyCBC, the FARs are from the individual searches, but for triggers found by several of the single-template searches, their inverse FARs have been summed.

To measure the evidence of lensing signatures in a signal, we define a Bayes factor $\mathcal{B}_U^{\text{ML}}$, which is the evidence ratio between the microlensed and unlensed hypotheses. Higher positive values correspond to support for lensing. Hannuksela et al. (2019) searched for similar beating patterns due to point mass lenses in the O1 and O2 data, using an upper lens mass prior cutoff $M_L^z \lesssim 10^5 M_\odot$. They reported no evidence for such lensing patterns above $\log_{10} \mathcal{B}_U^{\text{ML}} > 0.2$.

For O3a, we analyze the 36 events from Abbott et al. (2020d) that confidently have both component masses above $3M_\odot$ and search for microlensing signatures following the same method as in Hannuksela et al. (2019). We perform PE using Bilby (Ashton et al. 2019; Romero-Shaw et al. 2020) and the nested sampling algorithm dynesty (Speagle 2020). For each event, we perform two PE runs using both unlensed and microlensed templates. For the unlensed case, which is

similar to the usual PE analysis, equivalent prior settings and data dictionaries such as strain data and power spectral densities (PSDs) are used as in Abbott et al. (2020d). The analysis uses the IMRPHEenomXPHM (Pratten et al. 2020) waveform for most events, except for GW190521, which is analyzed using the NRSUR7DQ4 waveform (Varma et al. 2019) and for the least massive event GW190924.021846 where the IMRPHEenomPv2 waveform is used. The prior on M_L^z is log uniform in the range $[1-10^5 M_\odot]$, above which the effect of microlensing is relatively small for the LIGO–Virgo sensitivity band. The impact parameter prior is $p(y) \propto y$ between $[0.1, 3]$, chosen due to geometry and isotropy (Lai et al. 2018).

In Fig. 5 we show violin plots of marginalized posterior distributions for the redshifted lens mass for each event, as well as the Bayes factors between the microlensed and unlensed hypotheses. The broad M_L^z posteriors correspond to broad posteriors on the impact parameter y , which is not well constrained for unlensed cases. In terms of Bayes factors, there is no substantial evidence of microlensing with a maximum $\log_{10} \mathcal{B}_U^{\text{ML}} = 0.5$ for the event GW190910.112807. Additionally, as can be seen in Appendix C, statistical fluctuations of the \log_{10} Bayes factors for injections without microlensing can be as high as 0.75. Thus, the observed Bayes factors are already by themselves consistent with random noise fluctuations and do not significantly favor the microlensing hypothesis for any of the events. The resulting posterior odds O_U^{ML} , which are the products of Bayes factors and the low prior odds of microlensing (Lai et al. 2018), would be even lower. Thus, we find no evidence of microlensing in this study.

We searched for microlensing due to isolated point masses. More complex models in which point mass lenses embedded in an external macromodel potential such as galaxies and galaxy clusters (Diego et al. 2019) can produce additional modulation on the magnified waveform, which could also prove important in the LIGO–Virgo frequency band. Future searches could be extended to cover a broader range of microlensing models.

7. CONCLUSIONS AND OUTLOOK

We have searched for gravitational lensing effects on the GW observations from O3a, the first half of the third LIGO–Virgo observing run, finding no strong evidence of lensing. First, we outlined estimates for the rate of strongly lensed GWs. Second, presuming a non-observation of lensing, we constrained the BBH merger-rate density at high redshift. Third, we used merger-rate density models obtained through the non-observation of a SGWB to estimate the GW lensing rate.

Next, we performed an analysis of apparent high-mass events under the hypothesis that they are lensed signals from lower-mass sources, finding that the highest-mass BBHs from O3a could be consistent with component masses below the




Figure 5. The marginalized posterior distribution of redshifted lens mass M_L^z and $\log_{10} \mathcal{B}_{UL}^{\text{ML}}$ between microlensed and unlensed hypotheses. The corresponding \log_{10} Bayes factors are noted to the right of the plot. We find no evidence of microlensing by point mass lenses.

PISN mass gap, while GW190425 and GW190426.152155 would require extreme magnifications to be compatible with the Galactic BNS population. This hypothesis is at the moment mainly disfavored by the expected lensing rates, but in the future, more quantitative constraints could also be set by connecting these magnification results with lens modeling to make predictions for the appearance of multiple images or the possibility of microlensing.

We then searched for signatures of multiple lensed images from a single source through several methods. We first investigated the parameter consistency among all pairs of O3a events from GWTC-2 using a posterior-overlap method, finding no significant event pairs but identifying several interesting candidates with high overlap.

We followed up on these candidate pairs using two detailed joint-PE analyses, finding high parameter consistency for 11 pairs. However, after the inclusion of a more appropriate population prior, selection effects, and the prior odds against the lensing hypothesis, these candidates do not provide sufficient evidence for a strong lensing claim.

Moreover, we used two targeted matched-filter approaches to search for additional lensed images of the known events that could be hidden beneath the thresholds of the corresponding broader analyses used to produce GWTC-2, identifying six new candidates. After follow-up by joint PE, we found no evidence to conclude that any of these sub-threshold triggers are lensed images.

Finally, we analyzed 36 events from GWTC-2 for microlensing effects by performing full PE with waveforms incorporating microlensing by point mass lenses. We found no evidence of microlensing.

In summary, our results on O3a data are consistent with the expected low rate of lensing at current detector sensitivities. However, improved analysis methods and lens modeling may allow digging deeper into potential lensing effects. Electromagnetic follow-up of lensing candidates, even if they are not significant enough based on the GW data alone, could also be promising (Serenio et al. 2011; Smith et al. 2018; Hannuksela et al. 2020; Yu et al. 2020). With the current generation of detectors further improving their sensitivity and the global network being extended (Abbott et al. 2020c), the chances of detecting clear lensing signatures will improve, and the field will offer many possibilities at the latest with third-generation (Punturo et al. 2010; Abbott et al. 2017; Reitze et al. 2019; Maggiore et al. 2020) and space-based detectors (Amaro-Seoane et al. 2017; Hu & Wu 2017) and their expected cosmological reach.

This material is based upon work supported by NSF’s LIGO Laboratory which is a major facility fully funded by the National Science Foundation. The authors also gratefully acknowledge the support of the Science and Technology Facili-

ties Council (STFC) of the United Kingdom, the Max-Planck-Society (MPS), and the State of Niedersachsen/Germany for support of the construction of Advanced LIGO and construction and operation of the GEO600 detector. Additional support for Advanced LIGO was provided by the Australian Research Council. The authors gratefully acknowledge the Italian Istituto Nazionale di Fisica Nucleare (INFN), the French Centre National de la Recherche Scientifique (CNRS) and the Netherlands Organization for Scientific Research, for the construction and operation of the Virgo detector and the creation and support of the EGO consortium. The authors also gratefully acknowledge research support from these agencies as well as by the Council of Scientific and Industrial Research of India, the Department of Science and Technology, India, the Science & Engineering Research Board (SERB), India, the Ministry of Human Resource Development, India, the Spanish Agencia Estatal de Investigación, the Vicepresidència i Conselleria d’Innovació, Recerca i Turisme and the Conselleria d’Educació i Universitat del Govern de les Illes Balears, the Conselleria d’Innovació, Universitats, Ciència i Societat Digital de la Generalitat Valenciana and the CERCA Programme Generalitat de Catalunya, Spain, the National Science Centre of Poland and the Foundation for Polish Science (FNP), the Swiss National Science Foundation (SNSF), the Russian Foundation for Basic Research, the Russian Science Foundation, the European Commission, the European Regional Development Funds (ERDF), the Royal Society, the Scottish Funding Council, the Scottish Universities Physics Alliance, the Hungarian Scientific Research Fund (OTKA), the French Lyon Institute of Origins (LIO), the Belgian Fonds de la Recherche Scientifique (FRS-FNRS), Actions de Recherche Concertées (ARC) and Fonds Wetenschappelijk Onderzoek – Vlaanderen (FWO), Belgium, the

Paris Île-de-France Region, the National Research, Development and Innovation Office Hungary (NKFIH), the National Research Foundation of Korea, the Natural Science and Engineering Research Council Canada, Canadian Foundation for Innovation (CFI), the Brazilian Ministry of Science, Technology, and Innovations, the International Center for Theoretical Physics South American Institute for Fundamental Research (ICTP-SAIFR), the Research Grants Council of Hong Kong, the National Natural Science Foundation of China (NSFC), the Leverhulme Trust, the Research Corporation, the Ministry of Science and Technology (MOST), Taiwan, the United States Department of Energy, the Kavli Foundation, and the Gordon and Betty Moore Foundation. The authors gratefully acknowledge the support of the NSF, STFC, INFN and CNRS for provision of computational resources.

We would like to thank all of the essential workers who put their health at risk during the COVID-19 pandemic, without whom we would not have been able to complete this work.

Software: Analyses in this paper made use of LAL-SUITE (LIGO Scientific Collaboration and Virgo Collaboration 2018), the GstLAL (Cannon et al. 2012; Messick et al. 2017; Hanna et al. 2020; Sachdev et al. 2019) and PyCBC (Usman et al. 2016; Nitz et al. 2018, 2019; Davies et al. 2020) pipelines; Bayesian inference with CPNest (Veitch et al. 2020), PyMULTINEST (Feroz et al. 2009, 2019), BILBY (Ashton et al. 2019; Smith et al. 2020; Romero-Shaw et al. 2020), and LALINFERENCE (Veitch et al. 2015); as well as the packages NUMPY (Harris et al. 2020), SciPY (Virtanen et al. 2020), ASTROPY (Robitaille et al. 2013; Price-Whelan et al. 2018), IPYTHON (Perez & Granger 2007), and LIGO.SKYPYMAP (Singer 2019). Plots were produced with MATPLOTLIB (Hunter 2007), and SEABORN (Waskom et al. 2020).

APPENDIX

A. LENSING STATISTICS SUPPLEMENTARY

Assuming a specific BBH formation channel, we can estimate the lensing rate for merger signals from that population. For example, suppose BBHs form as a consequence of isolated binary evolution. In that case, one can theoretically model BBH formation assuming that it traces the star-formation rate, modulated by the delay time distribution and by the stellar metallicity evolution (Belczynski et al. 2008, 2010; Dominik et al. 2013; Marchant et al. 2018; Eldridge et al. 2019; Neijssel et al. 2019; Boco et al. 2019; Santoliquido et al. 2021). However, note that if the BBHs form through other means or through multiple channels, the merger-rate density could be different (e.g., Miller & Lauburg 2009; Antonini & Rasio 2016; Rodriguez & Loeb 2018; Fragione & Silk 2020; De Luca et al. 2020; Antonini & Gieles 2020; Wong et al. 2020; Zevin et al. 2021; Bouffanais et al. 2021).

Here we assume two models for the merger-rate density. We base the first model on the assumption that the merger-rate density of the observed BBHs traces the star-formation rate density and the BBHs originate from Population I/II stars.

In this work, we did not consider the contribution of Population III stars. Population III stars have not been observed yet, and their physical properties, binary fraction, and initial mass function are still a matter of debate (Nakamura & Umemura 2001; Madau & Rees 2001; Bromm et al. 2002; Schaerer 2002; Norman 2008; Machida 2008; Ishigaki et al. 2018). As such, the contribution of Population III BBHs to gravitational-wave sources is also uncertain (e.g., Bond & Carr 1984; Kowalska et al. 2012; Belczynski et al. 2017; Liu & Bromm 2020). Should Population III stars dominated the BBH formation at high redshift, our results would need to be re-interpreted.

The first model, which we label Model A, uses the following fits that bracket the available population synthesis results from the literature (e.g., [Belczynski et al. 2008, 2010](#); [Dominik et al. 2013](#); [Marchant et al. 2018](#); [Eldridge et al. 2019](#); [Neijssel et al. 2019](#); [Boco et al. 2019](#); [Santoliquido et al. 2021](#)):

$$\begin{aligned}\mathcal{R}_m^{\min}(z_m) &= \frac{a_1 e^{a_2 z_m}}{a_3 + e^{a_4 z_m}} \text{Gpc}^{-3} \text{yr}^{-1}, \\ \mathcal{R}_m^{\max}(z_m) &= \frac{b_1 e^{b_2 z_m}}{b_3 + e^{b_4 z_m}} \text{Gpc}^{-3} \text{yr}^{-1},\end{aligned}\tag{A1}$$

where the fitting parameters $a_1 = 58.497$, $a_2 = 2.06424$, $a_3 = 2.82338$, $a_4 = 2.52898$, $b_1 = 105356$, $b_2 = 1.30278$, $b_3 = 2714.36$, and $b_4 = 2.22903$.

We base the second model, Model B, on the assumption that the merger-rate density follows the [Madau & Dickinson \(2014\)](#) ansatz:

$$\mathcal{R}_m(z_m; R_0, \alpha) = R_0 \frac{(1 + z_m)^\kappa}{1 + [(1 + z_m)/(1 + z_p)]^{(\gamma + \kappa)}}.\tag{A2}$$

To constrain the merger-rate density at high redshift, we assume that no strong lensing has occurred (Sec. 3.2). We further assume that events occur following a Poisson process.

Let us now assume Model B for the merger-rate density, Eq. A2. The distribution of merger-rate density parameters, given that no strong lensing has occurred,

$$p(R_0, \kappa, \gamma, z_p | N, \{d_i\}) \propto \mathcal{W} \times p(R_0, \kappa, \gamma, z_p | \{d_i\}),\tag{A3}$$

where $p(R_0, \kappa, \gamma, z_p | \{d_i\})$ follows the posterior distribution of parameters inferred from LIGO–Virgo population studies ([Abbott et al. 2020f](#)), and

$$\mathcal{W} = \frac{N_{\text{avg}}(R_0, \kappa, \gamma, z_p)^N \exp[-N_{\text{avg}}(R_0, \kappa, \gamma, z_p)]}{N!},\tag{A4}$$

with N being the number of observed, strongly lensed GW signals, and $N_{\text{avg}}(R_0, \kappa, \gamma, z_p)$ the expected number of events within a time Δt . Here, like in Sec. 3, we do not account for detector downtime, and instead as a proxy presume that the detectors are always online. The R_0 and κ value is measured at a low redshift ([Abbott et al. 2020f](#)). The γ and z_p values are unconstrained here and thus match an uninformative prior, with $p(\gamma) = \mathcal{SN}(5, 10, 3)$ being a split normal distribution and $p(z_p)$ being uniformly distributed between $[0, 4]$. The above equations give all the necessary ingredients to forecast the rate of strongly lensed events and place constraints on the merger-rate density based on the number of lensed signals observed by LIGO and Virgo.

B. CONSTRUCTION OF SUB-THRESHOLD COUNTERPART SEARCH TEMPLATE BANKS

For the GstLAL and PyCBC searches for sub-threshold lensed counterparts (Sec. 5.3) the targeted template banks for each event are constructed starting from a certain choice of posterior distributions released with GWTC-2 ([Abbott et al. 2020d](#); [GWOSC 2020](#)), aiming for a reduced-size template bank that is effective at recovering signals similar to the primary observed event.

For the GstLAL pipeline, we start, for all but three of the O3a events from GWTC-2, from non-spinning posteriors obtained with the IMRPHENOMD waveform ([Husa et al. 2016](#); [Khan et al. 2016](#)). In three cases, we instead start from posteriors obtained with the IMRPHENOMPV2 waveform ([Hannam et al. 2014](#); [Bohé et al. 2016](#)), which includes spin precession. These exceptions are GW190413_052954, GW190426_152155, and GW190909_114149. We then choose subsets of the original broad template bank from the GWTC-2 analysis by comparing against the posteriors of each event, using the following steps as introduced by [Li et al. \(2019a\)](#): We first draw $O(1000)$ of each event’s posterior samples with the highest likelihoods to account for the uncertainty in the event’s measured mass and spin parameters. For each sample we simulate, using the aligned-spin SEOBNRv4_ROM waveform model ([Bohé et al. 2017](#); [Pürrer 2014](#); [Pürrer 2016](#)), one signal with the event’s original optimal signal-to-noise ratio ρ_{opt} as given by Eq. (2) in ([Li et al. 2019a](#)) and nine extra signals with smaller ρ_{opt} , scaled by changing their effective distances D_{eff} ([Allen et al. 2012b](#)). The reduced template bank for an event is then constructed by searching the simulated data with the original GWTC-2 template bank (which also consists of SEOBNRv4_ROM waveforms) and keeping those templates that recover any of the simulated signals with a FAR < 1 in 30 days.

For PyCBC we select a single template for each search, choosing the maximum-posterior redshifted masses and aligned-spin components $\{(1 + z)m_1, (1 + z)m_2, \chi_1, \chi_2\}$ as estimated from a four-dimensional Gaussian KDE fit to the posterior samples from [GWOSC \(2020\)](#) for these parameters. Where available, we use aligned-spin posterior samples. In the case of GW190412 and GW190814, we use samples generated using the SEOBNRv4_ROM waveform; for GW190426_152155 we use a mixture of samples generated using the SEOBNRv4_ROM_NRTidalv2_NSBH and IMRPhenomNSBH waveforms; and for GW190425 we




Figure 6. Distribution of microlensing Bayes factors $\log_{10} \mathcal{B}_U^{\text{ML}}$ for unlensed simulated signals, recovered using a lensed template.

use samples generated using the IMRPhenomD_NRTidal, TEOBResumS, and SEOBNRv4T_surrogate waveforms. If aligned-spin posteriors are not available in the GWOSC (2020) data release, we use precessing posterior samples and marginalise over the transverse-spin components before applying the KDE. This produces an aligned-spin template with high matches at the peak of the posterior. In the case of GW190521, we use samples generated using the IMRPhenomPv3HM (Khan et al. 2020), NRSur7Dq4 (Varma et al. 2019) and SEOBNRv4PHM (Ossokine et al. 2020) waveforms. For all other events, we use samples generated using the SEOBNRv4P and IMRPhenomPv2 waveforms.

These choices of waveforms and posterior samples are not necessarily optimal, but they are valid for this analysis in the sense that the recovery of similar waveforms with parameters close to the best-fit ones for the targeted GWTC-2 events has been verified through injection studies. In addition, in the actual searches, the targeted banks constructed in this way successfully recovered the corresponding GWTC-2 events in all GstLAL searches, while for PyCBC triggers within 0.1 s of the target events were excluded from the final trigger list, but in all cases where the original events were observed with two or more detector, a coincident trigger was also recovered in the targeted search. In future work, revisiting the choice of posterior samples used to construct template banks may further improve sub-threshold searches’ effectiveness.

C. INJECTION STUDY FOR MICROLENSING ANALYSIS

A high Bayes factor $\mathcal{B}_U^{\text{ML}}$ itself is not conclusive evidence of microlensing in an observed event. We have performed an injection study to explore the impact of statistical fluctuations on the Bayes factor obtained from unlensed signals. We generate unlensed injections by randomly drawing from the parameter space of precessing BBH systems. Simulated Gaussian noise is used considering nominal O3 sensitivity (Abbott et al. 2020c), and we use the IMRPhenomPv2 waveform model (Hannam et al. 2014; Bohé et al. 2016) for all simulated injections. The statistical fluctuations of $\log_{10} \mathcal{B}_U^{\text{ML}}$ for 100 unlensed injections recovered using lensed templates can be seen in Fig. 6 which shows that the typical values found are $\log_{10} \mathcal{B}_U^{\text{ML}} < 0.75$.

REFERENCES

- Aasi, J., et al. 2015, *Classical Quantum Gravity*, 32, 074001, doi: [10.1088/0264-9381/32/7/074001](https://doi.org/10.1088/0264-9381/32/7/074001)
- Abbott, B. P., et al. 2016a, *Class. Quant. Grav.*, 33, 134001, doi: [10.1088/0264-9381/33/13/134001](https://doi.org/10.1088/0264-9381/33/13/134001)
- , 2016b, *Astrophys. J.*, 833, L1, doi: [10.3847/2041-8205/833/1/L1](https://doi.org/10.3847/2041-8205/833/1/L1)
- , 2017, *Class. Quant. Grav.*, 34, 044001, doi: [10.1088/1361-6382/aa51f4](https://doi.org/10.1088/1361-6382/aa51f4)
- , 2019a, *Astrophys. J. Lett.*, 882, L24, doi: [10.3847/2041-8213/ab3800](https://doi.org/10.3847/2041-8213/ab3800)
- , 2019b, *Phys. Rev.*, X9, 031040, doi: [10.1103/PhysRevX.9.031040](https://doi.org/10.1103/PhysRevX.9.031040)
- , 2020a, *Class. Quant. Grav.*, 37, 055002, doi: [10.1088/1361-6382/ab685e](https://doi.org/10.1088/1361-6382/ab685e)
- , 2020b, *Astrophys. J. Lett.*, 892, L3, doi: [10.3847/2041-8213/ab75f5](https://doi.org/10.3847/2041-8213/ab75f5)
- , 2020c, *Living Rev. Relativity*, 23, 3, doi: [10.1007/s41114-020-00026-9](https://doi.org/10.1007/s41114-020-00026-9)
- Abbott, R., et al. 2020d. <https://arxiv.org/abs/2010.14527>
- , 2020e, doi: [10.3847/2041-8213/ab960f](https://doi.org/10.3847/2041-8213/ab960f)
- , 2020f. <https://arxiv.org/abs/2010.14533>
- , 2020g, *Astrophys. J. Lett.*, 900, L13, doi: [10.3847/2041-8213/aba493](https://doi.org/10.3847/2041-8213/aba493)

- . 2020h, *Phys. Rev. Lett.*, 125, 101102, doi: [10.1103/PhysRevLett.125.101102](https://doi.org/10.1103/PhysRevLett.125.101102)
- . 2021a, *SoftwareX*, 100658, doi: [10.1016/j.softx.2021.100658](https://doi.org/10.1016/j.softx.2021.100658)
- . 2021b. <https://arxiv.org/abs/2101.12130>
- Acernese, F., et al. 2015, *Classical Quantum Gravity*, 32, 024001, doi: [10.1088/0264-9381/32/2/024001](https://doi.org/10.1088/0264-9381/32/2/024001)
- . 2018, *Class. Quant. Grav.*, 35, 205004, doi: [10.1088/1361-6382/aadf1a](https://doi.org/10.1088/1361-6382/aadf1a)
- Ade, P. A. R., et al. 2016, *A&A*, 594, A13, doi: [10.1051/0004-6361/201525830](https://doi.org/10.1051/0004-6361/201525830)
- Allen, B. 2005, *Phys. Rev. D*, 71, 062001, doi: [10.1103/PhysRevD.71.062001](https://doi.org/10.1103/PhysRevD.71.062001)
- Allen, B., Anderson, W. G., Brady, P. R., Brown, D. A., & Creighton, J. D. E. 2012a, *Phys. Rev. D*, 85, 122006, doi: [10.1103/PhysRevD.85.122006](https://doi.org/10.1103/PhysRevD.85.122006)
- . 2012b, *Phys. Rev. D*, 85, 122006, doi: [10.1103/PhysRevD.85.122006](https://doi.org/10.1103/PhysRevD.85.122006)
- Amaro-Seoane, P., et al. 2017. <https://arxiv.org/abs/1702.00786>
- Antonini, F., & Gieles, M. 2020, *Phys. Rev. D*, 102, 123016, doi: [10.1103/PhysRevD.102.123016](https://doi.org/10.1103/PhysRevD.102.123016)
- Antonini, F., & Rasio, F. A. 2016, *Astrophys. J.*, 831, 187, doi: [10.3847/0004-637X/831/2/187](https://doi.org/10.3847/0004-637X/831/2/187)
- Arun, K. G., Buonanno, A., Faye, G., & Ochsner, E. 2009, *PhRvD*, 79, 104023, doi: [10.1103/PhysRevD.79.104023](https://doi.org/10.1103/PhysRevD.79.104023)
- Ashton, G., et al. 2019, *Astrophys. J. Suppl. Ser.*, 241, 27, doi: [10.3847/1538-4365/ab06fc](https://doi.org/10.3847/1538-4365/ab06fc)
- Baker, T., & Trodden, M. 2017, *Phys. Rev. D*, 95, 063512, doi: [10.1103/PhysRevD.95.063512](https://doi.org/10.1103/PhysRevD.95.063512)
- Bartelmann, M. 2010, *Class. Quant. Grav.*, 27, 233001, doi: [10.1088/0264-9381/27/23/233001](https://doi.org/10.1088/0264-9381/27/23/233001)
- Belczynski, K., Dominik, M., Bulik, T., et al. 2010, *Astrophys. J. Lett.*, 715, L138, doi: [10.1088/2041-8205/715/2/L138](https://doi.org/10.1088/2041-8205/715/2/L138)
- Belczynski, K., Holz, D. E., Bulik, T., & O'Shaughnessy, R. 2016a, *Nature*, 534, 512, doi: [10.1038/nature18322](https://doi.org/10.1038/nature18322)
- Belczynski, K., Kalogera, V., Rasio, F. A., et al. 2008, *Astrophys. J. Suppl.*, 174, 223, doi: [10.1086/521026](https://doi.org/10.1086/521026)
- Belczynski, K., Ryu, T., Perna, R., et al. 2017, *Mon. Not. Roy. Astron. Soc.*, 471, 4702, doi: [10.1093/mnras/stx1759](https://doi.org/10.1093/mnras/stx1759)
- Belczynski, K., et al. 2016b, *Astron. Astrophys.*, 594, A97, doi: [10.1051/0004-6361/201628980](https://doi.org/10.1051/0004-6361/201628980)
- Blanchet, L. 2014, *Living Rev. Relativity*, 17, 2, doi: [10.12942/lrr-2014-2](https://doi.org/10.12942/lrr-2014-2)
- Blanchet, L., Damour, T., Esposito-Farèse, G., & Iyer, B. R. 2005, *Phys. Rev. D*, 71, 124004, doi: [10.1103/PhysRevD.71.124004](https://doi.org/10.1103/PhysRevD.71.124004)
- Blanchet, L., Damour, T., Iyer, B. R., Will, C. M., & Wiseman, A. G. 1995, *PhRvL*, 74, 3515, doi: [10.1103/PhysRevLett.74.3515](https://doi.org/10.1103/PhysRevLett.74.3515)
- Blandford, R., & Narayan, R. 1986, *Astrophys. J.*, 310, 568, doi: [10.1086/164709](https://doi.org/10.1086/164709)
- Boco, L., Lapi, A., Goswami, S., et al. 2019, *Astrophys. J.*, 881, 157, doi: [10.3847/1538-4357/ab328e](https://doi.org/10.3847/1538-4357/ab328e)
- Bohé, A., Hannam, M., Husa, S., et al. 2016, *PhenomPv2 - Technical Notes for LAL Implementation*, Tech. Rep. LIGO-T1500602, LIGO Project. <https://dcc.ligo.org/LIGO-T1500602>
- Bohé, A., Faye, G., Marsat, S., & Porter, E. K. 2015, *Class. Quant. Grav.*, 32, 195010, doi: [10.1088/0264-9381/32/19/195010](https://doi.org/10.1088/0264-9381/32/19/195010)
- Bohé, A., Marsat, S., & Blanchet, L. 2013, *Class. Quant. Grav.*, 30, 135009, doi: [10.1088/0264-9381/30/13/135009](https://doi.org/10.1088/0264-9381/30/13/135009)
- Bohé, A., et al. 2017, *Phys. Rev. D*, 95, 044028, doi: [10.1103/PhysRevD.95.044028](https://doi.org/10.1103/PhysRevD.95.044028)
- Bond, I. A., et al. 2004, *Astrophys. J. Lett.*, 606, L155, doi: [10.1086/420928](https://doi.org/10.1086/420928)
- Bond, J. R., & Carr, B. J. 1984, *Mon. Not. Roy. Astron. Soc.*, 207, 585, doi: [10.1093/mnras/207.3.585](https://doi.org/10.1093/mnras/207.3.585)
- Bouffanais, Y., Mapelli, M., Santoliquido, F., et al. 2021. <https://arxiv.org/abs/2102.12495>
- Broadhurst, T., Diego, J. M., & Smoot, G. 2018. <https://arxiv.org/abs/1802.05273>
- Broadhurst, T., Diego, J. M., & Smoot, G. F. 2019. <https://arxiv.org/abs/1901.03190>
- . 2020a. <https://arxiv.org/abs/2002.08821>
- . 2020b. <https://arxiv.org/abs/2006.13219>
- Bromm, V., Coppi, P. S., & Larson, R. B. 2002, *Astrophys. J.*, 564, 23, doi: [10.1086/323947](https://doi.org/10.1086/323947)
- Buikema, A., et al. 2020, *Phys. Rev. D*, 102, 062003, doi: [10.1103/PhysRevD.102.062003](https://doi.org/10.1103/PhysRevD.102.062003)
- Buonanno, A., Iyer, B. R., Ochsner, E., Pan, Y., & Sathyaprakash, B. S. 2009, *Phys.Rev.*, D80, 084043, doi: [10.1103/PhysRevD.80.084043](https://doi.org/10.1103/PhysRevD.80.084043)
- Buscicchio, R., Moore, C. J., Pratten, G., et al. 2020a, *Phys. Rev. Lett.*, 125, 141102, doi: [10.1103/PhysRevLett.125.141102](https://doi.org/10.1103/PhysRevLett.125.141102)
- Buscicchio, R., Moore, C. J., Pratten, G., Schmidt, P., & Vecchio, A. 2020b, *Phys. Rev. D*, 102, 081501, doi: [10.1103/PhysRevD.102.081501](https://doi.org/10.1103/PhysRevD.102.081501)
- Cahillane, C., et al. 2017, *Phys. Rev. D*, 96, 102001, doi: [10.1103/PhysRevD.96.102001](https://doi.org/10.1103/PhysRevD.96.102001)
- Callister, T., Fishbach, M., Holz, D., & Farr, W. 2020, *Astrophys. J. Lett.*, 896, L32, doi: [10.3847/2041-8213/ab9743](https://doi.org/10.3847/2041-8213/ab9743)
- Cannon, K., Cariou, R., Chapman, A., et al. 2012, *Astrophys.J.*, 748, 136, doi: [10.1088/0004-637X/748/2/136](https://doi.org/10.1088/0004-637X/748/2/136)
- Cao, S., Qi, J., Cao, Z., et al. 2019, *Sci. Rep.*, 9, 11608, doi: [10.1038/s41598-019-47616-4](https://doi.org/10.1038/s41598-019-47616-4)
- Cao, Z., Li, L.-F., & Wang, Y. 2014, *Phys. Rev. D*, 90, 062003, doi: [10.1103/PhysRevD.90.062003](https://doi.org/10.1103/PhysRevD.90.062003)
- Chen, H.-Y., Essick, R., Vitale, S., Holz, D. E., & Katsavounidis, E. 2017, *Astrophys. J.*, 835, 31, doi: [10.3847/1538-4357/835/1/31](https://doi.org/10.3847/1538-4357/835/1/31)
- Choi, Y.-Y., Park, C., & Vogeley, M. S. 2007, *Astrophys. J.*, 658, 884, doi: [10.1086/511060](https://doi.org/10.1086/511060)
- Christian, P., Vitale, S., & Loeb, A. 2018, *Phys. Rev. D*, 98, 103022, doi: [10.1103/PhysRevD.98.103022](https://doi.org/10.1103/PhysRevD.98.103022)

- Clowe, D., Gonzalez, A., & Markevitch, M. 2004, *Astrophys. J.*, 604, 596, doi: [10.1086/381970](https://doi.org/10.1086/381970)
- Coe, D., et al. 2013, *Astrophys. J.*, 762, 32, doi: [10.1088/0004-637X/762/1/32](https://doi.org/10.1088/0004-637X/762/1/32)
- Cornish, N. J., & Littenberg, T. B. 2015, *Class. Quant. Grav.*, 32, 135012, doi: [10.1088/0264-9381/32/13/135012](https://doi.org/10.1088/0264-9381/32/13/135012)
- Dai, L., & Venumadhav, T. 2017. <https://arxiv.org/abs/1702.04724>
- Dai, L., Venumadhav, T., & Sigurdson, K. 2017, *Phys. Rev. D*, 95, 044011, doi: [10.1103/PhysRevD.95.044011](https://doi.org/10.1103/PhysRevD.95.044011)
- Dai, L., Zackay, B., Venumadhav, T., Roulet, J., & Zaldarriaga, M. 2020. <https://arxiv.org/abs/2007.12709>
- Dal Canton, T., et al. 2014, *Phys. Rev. D*, 90, 082004, doi: [10.1103/PhysRevD.90.082004](https://doi.org/10.1103/PhysRevD.90.082004)
- Damour, T., Jaranowski, P., & Schaefer, G. 2000, *PhRvD*, 62, 084011, doi: [10.1103/PhysRevD.62.084011](https://doi.org/10.1103/PhysRevD.62.084011)
- Damour, T., & Nagar, A. 2016, *Lect. Notes Phys.*, 905, 273, doi: [10.1007/978-3-319-19416-5_7](https://doi.org/10.1007/978-3-319-19416-5_7)
- Davies, G. S., Dent, T., Tápai, M., et al. 2020, *Phys. Rev. D*, 102, 022004, doi: [10.1103/PhysRevD.102.022004](https://doi.org/10.1103/PhysRevD.102.022004)
- Davis, D., Massinger, T. J., Lundgren, A. P., et al. 2019, *Class. Quant. Grav.*, 36, 055011, doi: [10.1088/1361-6382/ab01c5](https://doi.org/10.1088/1361-6382/ab01c5)
- Davis, D., et al. 2021. <https://arxiv.org/abs/2101.11673>
- De Luca, V., Franciolini, G., Pani, P., & Riotto, A. 2020, *JCAP*, 04, 052, doi: [10.1088/1475-7516/2020/04/052](https://doi.org/10.1088/1475-7516/2020/04/052)
- Deguchi, S., & Watson, W. D. 1986, *Phys. Rev. D*, 34, 1708, doi: [10.1103/PhysRevD.34.1708](https://doi.org/10.1103/PhysRevD.34.1708)
- Diego, J., Hannuksela, O., Kelly, P., et al. 2019, *Astron. Astrophys.*, 627, A130, doi: [10.1051/0004-6361/201935490](https://doi.org/10.1051/0004-6361/201935490)
- Diego, J. M. 2020, *Phys. Rev. D*, 101, 123512, doi: [10.1103/PhysRevD.101.123512](https://doi.org/10.1103/PhysRevD.101.123512)
- Dietrich, T., et al. 2019, *Phys. Rev. D*, 99, 024029, doi: [10.1103/PhysRevD.99.024029](https://doi.org/10.1103/PhysRevD.99.024029)
- Dominik, M., Belczynski, K., Fryer, C., et al. 2013, *Astrophys. J.*, 779, 72, doi: [10.1088/0004-637X/779/1/72](https://doi.org/10.1088/0004-637X/779/1/72)
- Driggers, J. C., et al. 2019, *Phys. Rev. D*, 99, 042001, doi: [10.1103/PhysRevD.99.042001](https://doi.org/10.1103/PhysRevD.99.042001)
- Eldridge, J., Stanway, E., & Tang, P. N. 2019, *Mon. Not. Roy. Astron. Soc.*, 482, 870, doi: [10.1093/mnras/sty2714](https://doi.org/10.1093/mnras/sty2714)
- Essick, R., Godwin, P., Hanna, C., Blackburn, L., & Katsavounidis, E. 2020. <https://arxiv.org/abs/2005.12761>
- Ezquiaga, J. M., Holz, D. E., Hu, W., Lagos, M., & Wald, R. M. 2021, *Phys. Rev. D*, 103, 6, doi: [10.1103/PhysRevD.103.064047](https://doi.org/10.1103/PhysRevD.103.064047)
- Fan, X.-L., Liao, K., Biesiada, M., Piorkowska-Kurpas, A., & Zhu, Z.-H. 2017, *Phys. Rev. Lett.*, 118, 091102, doi: [10.1103/PhysRevLett.118.091102](https://doi.org/10.1103/PhysRevLett.118.091102)
- Farrow, N., Zhu, X.-J., & Thrane, E. 2019, *Astrophys. J.*, 876, 18, doi: [10.3847/1538-4357/ab12e3](https://doi.org/10.3847/1538-4357/ab12e3)
- Faye, G., Marsat, S., Blanchet, L., & Iyer, B. R. 2012, *Class. Quant. Grav.*, 29, 175004, doi: [10.1088/0264-9381/29/17/175004](https://doi.org/10.1088/0264-9381/29/17/175004)
- Feroz, F., Hobson, M., & Bridges, M. 2009, *Mon. Not. Roy. Astron. Soc.*, 398, 1601, doi: [10.1111/j.1365-2966.2009.14548.x](https://doi.org/10.1111/j.1365-2966.2009.14548.x)
- Feroz, F., Hobson, M., Cameron, E., & Pettitt, A. 2019, *Open J. Astrophys.*, 2, 10, doi: [10.21105/astro.1306.2144](https://doi.org/10.21105/astro.1306.2144)
- Fiori, I., et al. 2020, *Galaxies*, 8, 82, doi: [10.3390/galaxies8040082](https://doi.org/10.3390/galaxies8040082)
- Fishbach, M., & Holz, D. E. 2017, *Astrophys. J. Lett.*, 851, L25, doi: [10.3847/2041-8213/aa9bf6](https://doi.org/10.3847/2041-8213/aa9bf6)
- Fragione, G., & Silk, J. 2020, *Mon. Not. Roy. Astron. Soc.*, 498, 4591, doi: [10.1093/mnras/staa2629](https://doi.org/10.1093/mnras/staa2629)
- Fukugita, M., & Turner, E. L. 1991, *Mon. Not. Roy. Astron. Soc.*, 253, 99
- Godwin, P., et al. 2020. <https://arxiv.org/abs/2010.15282>
- Goyal, S., Haris, K., Mehta, A. K., & Ajith, P. 2021, *Phys. Rev. D*, 103, 024038, doi: [10.1103/PhysRevD.103.024038](https://doi.org/10.1103/PhysRevD.103.024038)
- GWOSC. 2020, GWTC-2 Data Release, doi: [10.7935/99gf-ax93](https://doi.org/10.7935/99gf-ax93)
- Hanna, C., et al. 2020, *Phys. Rev. D*, 101, 022003, doi: [10.1103/PhysRevD.101.022003](https://doi.org/10.1103/PhysRevD.101.022003)
- Hannam, M., Schmidt, P., Bohé, A., et al. 2014, *Phys. Rev. Lett.*, 113, 151101, doi: [10.1103/PhysRevLett.113.151101](https://doi.org/10.1103/PhysRevLett.113.151101)
- Hannuksela, O., Haris, K., Ng, K., et al. 2019, *Astrophys. J. Lett.*, 874, L2, doi: [10.3847/2041-8213/ab0c0f](https://doi.org/10.3847/2041-8213/ab0c0f)
- Hannuksela, O. A., Collett, T. E., Çalişkan, M., & Li, T. G. F. 2020, *Mon. Not. Roy. Astron. Soc.*, 498, 3395, doi: [10.1093/mnras/staa2577](https://doi.org/10.1093/mnras/staa2577)
- Haris, K., Mehta, A. K., Kumar, S., Venumadhav, T., & Ajith, P. 2018. <https://arxiv.org/abs/1807.07062>
- Harris, C. R., et al. 2020, *Nature*, 585, 357, doi: [10.1038/s41586-020-2649-2](https://doi.org/10.1038/s41586-020-2649-2)
- Hu, W.-R., & Wu, Y.-L. 2017, *Natl. Sci. Rev.*, 4, 685, doi: [10.1093/nsr/nwx116](https://doi.org/10.1093/nsr/nwx116)
- Hunter, J. D. 2007, *Computing in Science Engineering*, 9, 90, doi: [10.1109/MCSE.2007.55](https://doi.org/10.1109/MCSE.2007.55)
- Husa, S., Khan, S., Hannam, M., et al. 2016, *Phys. Rev. D*, 93, 044006, doi: [10.1103/PhysRevD.93.044006](https://doi.org/10.1103/PhysRevD.93.044006)
- Ishigaki, M. N., Tominaga, N., Kobayashi, C., & Nomoto, K. 2018, *Astrophys. J.*, 857, 46, doi: [10.3847/1538-4357/aab3de](https://doi.org/10.3847/1538-4357/aab3de)
- Jung, S., & Shin, C. S. 2019, *Phys. Rev. Lett.*, 122, 041103, doi: [10.1103/PhysRevLett.122.041103](https://doi.org/10.1103/PhysRevLett.122.041103)
- Karki, S., et al. 2016, *Rev. Sci. Instrum.*, 87, 114503, doi: [10.1063/1.4967303](https://doi.org/10.1063/1.4967303)
- Khan, S., Husa, S., Hannam, M., et al. 2016, *Phys. Rev. D*, 93, 044007, doi: [10.1103/PhysRevD.93.044007](https://doi.org/10.1103/PhysRevD.93.044007)
- Khan, S., Ohme, F., Chatziioannou, K., & Hannam, M. 2020, *Phys. Rev. D*, 101, 024056, doi: [10.1103/PhysRevD.101.024056](https://doi.org/10.1103/PhysRevD.101.024056)
- Kimball, C., et al. 2020. <https://arxiv.org/abs/2011.05332>
- Klimenko, S., Mohanty, S., Rakhmanov, M., & Mitselmakher, G. 2005, *Phys. Rev. D*, 72, 122002, doi: [10.1103/PhysRevD.72.122002](https://doi.org/10.1103/PhysRevD.72.122002)
- . 2006, *J. Phys. Conf. Ser.*, 32, 12, doi: [10.1088/1742-6596/32/1/003](https://doi.org/10.1088/1742-6596/32/1/003)

- Klimenko, S., Yakushin, I., Rakhmanov, M., & Mitselmakher, G. 2004, *Class. Quant. Grav.*, 21, S1685, doi: [10.1088/0264-9381/21/20/011](https://doi.org/10.1088/0264-9381/21/20/011)
- Klimenko, S., Vedovato, G., Drago, M., et al. 2011, *Phys. Rev. D*, 83, 102001, doi: [10.1103/PhysRevD.83.102001](https://doi.org/10.1103/PhysRevD.83.102001)
- Klimenko, S., et al. 2016, *Phys. Rev. D*, 93, 042004, doi: [10.1103/PhysRevD.93.042004](https://doi.org/10.1103/PhysRevD.93.042004)
- Kowalska, I., Bulik, T., & Belczynski, K. 2012, *A&A*, 541, A120, doi: [10.1051/0004-6361/201118604](https://doi.org/10.1051/0004-6361/201118604)
- Lai, K.-H., Hannuksela, O. A., Herrera-Martín, A., et al. 2018, *Phys. Rev. D*, 98, 083005, doi: [10.1103/PhysRevD.98.083005](https://doi.org/10.1103/PhysRevD.98.083005)
- Li, A. K., Lo, R. K., Sachdev, S., et al. 2019a, <https://arxiv.org/abs/1904.06020>
- Li, S.-S., Mao, S., Zhao, Y., & Lu, Y. 2018, *Mon. Not. Roy. Astron. Soc.*, 476, 2220, doi: [10.1093/mnras/sty411](https://doi.org/10.1093/mnras/sty411)
- Li, Y., Fan, X., & Gou, L. 2019b, *Astrophys. J.*, 873, 37, doi: [10.3847/1538-4357/ab037e](https://doi.org/10.3847/1538-4357/ab037e)
- Liao, K., Fan, X.-L., Ding, X.-H., Biesiada, M., & Zhu, Z.-H. 2017, *Nature Commun.*, 8, 1148, doi: [10.1038/s41467-017-01152-9](https://doi.org/10.1038/s41467-017-01152-9)
- LIGO Scientific Collaboration and Virgo Collaboration. 2018, LALSuite software, doi: [10.7935/GT1W-FZ16](https://doi.org/10.7935/GT1W-FZ16)
- . 2021, Data release for "Search for lensing signatures in the gravitational-wave observations from the first half of LIGO-Virgo's third observing run". <https://dcc.ligo.org/P2100173/public>
- Liu, B., & Bromm, V. 2020, *Mon. Not. Roy. Astron. Soc.*, 495, 2475, doi: [10.1093/mnras/staa1362](https://doi.org/10.1093/mnras/staa1362)
- Liu, X., Hernandez, I. M., & Creighton, J. 2021, *Astrophys. J.*, 908, 97, doi: [10.3847/1538-4357/abd7eb](https://doi.org/10.3847/1538-4357/abd7eb)
- Lo, R. K. L., & Magaña Hernandez, I. 2021, <https://arxiv.org/abs/2104.09339>
- Machida, M. N. 2008, *Astrophys. J. Lett.*, 682, L1, doi: [10.1086/590109](https://doi.org/10.1086/590109)
- Madau, P., & Dickinson, M. 2014, *Ann. Rev. Astron. Astrophys.*, 52, 415, doi: [10.1146/annurev-astro-081811-125615](https://doi.org/10.1146/annurev-astro-081811-125615)
- Madau, P., & Rees, M. J. 2001, *Astrophys. J. Lett.*, 551, L27, doi: [10.1086/319848](https://doi.org/10.1086/319848)
- Maggiore, M., et al. 2020, *JCAP*, 03, 050, doi: [10.1088/1475-7516/2020/03/050](https://doi.org/10.1088/1475-7516/2020/03/050)
- Malmquist, K. G. 1922, *Meddelanden fran Lunds Astronomiska Observatorium Serie I*, 100, 1
- Marchant, P., Renzo, M., Farmer, R., et al. 2018, doi: [10.3847/1538-4357/ab3426](https://doi.org/10.3847/1538-4357/ab3426)
- Markevitch, M., Gonzalez, A., Clowe, D., et al. 2004, *Astrophys. J.*, 606, 819, doi: [10.1086/383178](https://doi.org/10.1086/383178)
- McIsaac, C., Keitel, D., Collett, T., et al. 2020, *Phys. Rev. D*, 102, 084031, doi: [10.1103/PhysRevD.102.084031](https://doi.org/10.1103/PhysRevD.102.084031)
- Messick, C., et al. 2017, *Phys. Rev. D*, 95, 042001, doi: [10.1103/PhysRevD.95.042001](https://doi.org/10.1103/PhysRevD.95.042001)
- Mikoczi, B., Vasuth, M., & Gergely, L. A. 2005, *Phys. Rev.*, D71, 124043, doi: [10.1103/PhysRevD.71.124043](https://doi.org/10.1103/PhysRevD.71.124043)
- Miller, M. C., & Lauburg, V. M. 2009, *Astrophys. J.*, 692, 917, doi: [10.1088/0004-637X/692/1/917](https://doi.org/10.1088/0004-637X/692/1/917)
- Mishra, C. K., Kela, A., Arun, K. G., & Faye, G. 2016, *Phys. Rev.*, D93, 084054, doi: [10.1103/PhysRevD.93.084054](https://doi.org/10.1103/PhysRevD.93.084054)
- Mukherjee, S., Broadhurst, T., Diego, J. M., Silk, J., & Smoot, G. F. 2021, *Mon. Not. Roy. Astron. Soc.*, 501, 2451, doi: [10.1093/mnras/staa3813](https://doi.org/10.1093/mnras/staa3813)
- Nakamura, F., & Umemura, M. 2001, *Astrophys. J.*, 548, 19, doi: [10.1086/318663](https://doi.org/10.1086/318663)
- Nakamura, T. T. 1998, *Phys. Rev. Lett.*, 80, 1138, doi: [10.1103/PhysRevLett.80.1138](https://doi.org/10.1103/PhysRevLett.80.1138)
- Neijssel, C. J., Vigna-Gómez, A., Stevenson, S., et al. 2019, *Mon. Not. Roy. Astron. Soc.*, 490, 3740, doi: [10.1093/mnras/stz2840](https://doi.org/10.1093/mnras/stz2840)
- Ng, K. K., Wong, K. W., Broadhurst, T., & Li, T. G. 2018, *Phys. Rev. D*, 97, 023012, doi: [10.1103/PhysRevD.97.023012](https://doi.org/10.1103/PhysRevD.97.023012)
- Nguyen, P., et al. 2021. <https://arxiv.org/abs/2101.09935>
- Nitz, A. H., Dal Canton, T., Davis, D., & Reyes, S. 2018, *Phys. Rev. D*, 98, 024050, doi: [10.1103/PhysRevD.98.024050](https://doi.org/10.1103/PhysRevD.98.024050)
- Nitz, A. H., Dent, T., Dal Canton, T., Fairhurst, S., & Brown, D. A. 2017, *Astrophys. J.*, 849, 118, doi: [10.3847/1538-4357/aa8f50](https://doi.org/10.3847/1538-4357/aa8f50)
- Nitz, A. H., et al. 2019, *gwastro/pycbc: PyCBC Release v1.15.3*, doi: [10.5281/zenodo.3596447](https://doi.org/10.5281/zenodo.3596447)
- Norman, M. L. 2008, *AIP Conf. Proc.*, 990, 3, doi: [10.1063/1.2905629](https://doi.org/10.1063/1.2905629)
- Oguri, M. 2018, *Mon. Not. Roy. Astron. Soc.*, 480, 3842, doi: [10.1093/mnras/sty2145](https://doi.org/10.1093/mnras/sty2145)
- Oguri, M., & Marshall, P. J. 2010, *Mon. Not. Roy. Astron. Soc.*, 405, 2579, doi: [10.1111/j.1365-2966.2010.16639.x](https://doi.org/10.1111/j.1365-2966.2010.16639.x)
- Oguri, M., & Takahashi, R. 2020, *Astrophys. J.*, 901, 58, doi: [10.3847/1538-4357/abafab](https://doi.org/10.3847/1538-4357/abafab)
- Ohanian, H. 1974, *Int. J. Theor. Phys.*, 9, 425, doi: [10.1007/BF01810927](https://doi.org/10.1007/BF01810927)
- Ossokine, S., et al. 2020, *Phys. Rev. D*, 102, 044055, doi: [10.1103/PhysRevD.102.044055](https://doi.org/10.1103/PhysRevD.102.044055)
- Pagano, G., Hannuksela, O. A., & Li, T. G. F. 2020, *Astron. Astrophys.*, 643, A167, doi: [10.1051/0004-6361/202038730](https://doi.org/10.1051/0004-6361/202038730)
- Palenzuela, C. 2020, *Front. Astron. Space Sci.*, 7, 58, doi: [10.3389/fspas.2020.00058](https://doi.org/10.3389/fspas.2020.00058)
- Pang, P. T., Hannuksela, O. A., Dietrich, T., Pagano, G., & Harry, I. W. 2020, doi: [10.1093/mnras/staa1430](https://doi.org/10.1093/mnras/staa1430)
- Perez, F., & Granger, B. E. 2007, *Computing in Science Engineering*, 9, 21, doi: [10.1109/MCSE.2007.53](https://doi.org/10.1109/MCSE.2007.53)
- Poisson, E. 1998, *Phys. Rev. D*, 57, 5287, doi: [10.1103/PhysRevD.57.5287](https://doi.org/10.1103/PhysRevD.57.5287)
- Pratten, G., et al. 2020. <https://arxiv.org/abs/2004.06503>
- Price-Whelan, A., et al. 2018, *Astron. J.*, 156, 123, doi: [10.3847/1538-3881/aabc4f](https://doi.org/10.3847/1538-3881/aabc4f)

- Punturo, M., et al. 2010, *Class. Quant. Grav.*, 27, 084007, doi: [10.1088/0264-9381/27/8/084007](https://doi.org/10.1088/0264-9381/27/8/084007)
- Pürrer, M. 2014, *Class. Quant. Grav.*, 31, 195010, doi: [10.1088/0264-9381/31/19/195010](https://doi.org/10.1088/0264-9381/31/19/195010)
- Pürrer, M. 2016, *Phys. Rev. D*, 93, 064041, doi: [10.1103/PhysRevD.93.064041](https://doi.org/10.1103/PhysRevD.93.064041)
- Reitze, D., et al. 2019, *Bull. Am. Astron. Soc.*, 51, 035. <https://arxiv.org/abs/1907.04833>
- Robertson, A., Smith, G. P., Massey, R., et al. 2020, doi: [10.1093/mnras/staa1429](https://doi.org/10.1093/mnras/staa1429)
- Robitaille, T. P., et al. 2013, *Astron. Astrophys.*, 558, A33, doi: [10.1051/0004-6361/201322068](https://doi.org/10.1051/0004-6361/201322068)
- Rodriguez, C. L., & Loeb, A. 2018, *Astrophys. J. Lett.*, 866, L5, doi: [10.3847/2041-8213/aae377](https://doi.org/10.3847/2041-8213/aae377)
- Romero-Shaw, I. M., et al. 2020, *Mon. Not. Roy. Astron. Soc.*, 499, 3295, doi: [10.1093/mnras/staa2850](https://doi.org/10.1093/mnras/staa2850)
- Roulet, J., Venumadhav, T., Zackay, B., Dai, L., & Zaldarriaga, M. 2020, *Phys. Rev. D*, 102, 123022, doi: [10.1103/PhysRevD.102.123022](https://doi.org/10.1103/PhysRevD.102.123022)
- Rydzanowski, D., Smith, G. P., Bianconi, M., et al. 2020, *Mon. Not. Roy. Astron. Soc.*, 495, 1666, doi: [10.1093/mnras/staa1274](https://doi.org/10.1093/mnras/staa1274)
- Sachdev, S., et al. 2019. <https://arxiv.org/abs/1901.08580>
- Santoliquido, F., Mapelli, M., Giacobbo, N., Bouffanais, Y., & Artale, M. C. 2021, *Mon. Not. Roy. Astron. Soc.*, 502, 4877, doi: [10.1093/mnras/stab280](https://doi.org/10.1093/mnras/stab280)
- Sathyaprakash, B. S., & Dhurandhar, S. V. 1991, *Phys. Rev. D*, 44, 3819, doi: [10.1103/PhysRevD.44.3819](https://doi.org/10.1103/PhysRevD.44.3819)
- Schaerer, D. 2002, *Astron. Astrophys.*, 382, 28, doi: [10.1051/0004-6361:20011619](https://doi.org/10.1051/0004-6361:20011619)
- Schmidt, P. 2020, *Front. Astron. Space Sci.*, 7, 28, doi: [10.3389/fspas.2020.00028](https://doi.org/10.3389/fspas.2020.00028)
- Sereno, M., Jetzer, P., Sesana, A., & Volonteri, M. 2011, *Mon. Not. Roy. Astron. Soc.*, 415, 2773, doi: [10.1111/j.1365-2966.2011.18895.x](https://doi.org/10.1111/j.1365-2966.2011.18895.x)
- Singer, L. 2019, *ligo.skymap* <https://lscsoft.docs.ligo.org/ligo.skymap/>
- Singer, L. P., Goldstein, D. A., & Bloom, J. S. 2019. <https://arxiv.org/abs/1910.03601>
- Singer, L. P., & Price, L. R. 2016, *Phys. Rev.*, D93, 024013, doi: [10.1103/PhysRevD.93.024013](https://doi.org/10.1103/PhysRevD.93.024013)
- Skilling, J. 2006, *Bayesian Analysis*, 1, 833, doi: [10.1214/06-BA127](https://doi.org/10.1214/06-BA127)
- Smith, G., et al. 2017, *IAU Symp.*, 338, 98, doi: [10.1017/S1743921318003757](https://doi.org/10.1017/S1743921318003757)
- Smith, G. P., Bianconi, M., Massey, R., & Robertson, A. 2019, *GRB Coordinates Network*, 24250, 1
- Smith, G. P., Jauzac, M., Veitch, J., et al. 2018, *Mon. Not. Roy. Astron. Soc.*, 475, 3823, doi: [10.1093/mnras/sty031](https://doi.org/10.1093/mnras/sty031)
- Smith, G. P., Robertson, A., Bianconi, M., & Jauzac, M. 2019. <https://arxiv.org/abs/1902.05140>
- Smith, R. J. E., Ashton, G., Vajpeyi, A., & Talbot, C. 2020, *Mon. Not. Roy. Astron. Soc.*, 498, 4492, doi: [10.1093/mnras/staa2483](https://doi.org/10.1093/mnras/staa2483)
- Speagle, J. S. 2020, *Monthly Notices of the Royal Astronomical Society*, 493, 3132, doi: [10.1093/mnras/staa278](https://doi.org/10.1093/mnras/staa278)
- Sun, L., Goetz, E., Kissel, J. S., et al. 2020, *Classical and Quantum Gravity*, 37, 225008, doi: [10.1088/1361-6382/abb14e](https://doi.org/10.1088/1361-6382/abb14e)
- Takahashi, R., & Nakamura, T. 2003, *Astrophys. J.*, 595, 1039, doi: [10.1086/377430](https://doi.org/10.1086/377430)
- Thorne, K. 1982, in *Les Houches Summer School on Gravitational Radiation*, 1–57
- Tinker, J. L., Kravtsov, A. V., Klypin, A., et al. 2008, *Astrophys. J.*, 688, 709, doi: [10.1086/591439](https://doi.org/10.1086/591439)
- Usman, S. A., et al. 2016, *Class. Quant. Grav.*, 33, 215004, doi: [10.1088/0264-9381/33/21/215004](https://doi.org/10.1088/0264-9381/33/21/215004)
- Vajente, G., Huang, Y., Isi, M., et al. 2020, *Phys. Rev. D*, 101, 042003, doi: [10.1103/PhysRevD.101.042003](https://doi.org/10.1103/PhysRevD.101.042003)
- Varma, V., Field, S. E., Scheel, M. A., et al. 2019, *Phys. Rev. Research.*, 1, 033015, doi: [10.1103/PhysRevResearch.1.033015](https://doi.org/10.1103/PhysRevResearch.1.033015)
- Veitch, J., et al. 2015, *Phys. Rev. D*, 91, 042003, doi: [10.1103/PhysRevD.91.042003](https://doi.org/10.1103/PhysRevD.91.042003)
- Veitch, J., Pozzo, W. D., Williams, M., et al. 2020, *johnveitch/cpnest: Release 0.10.2, v0.10.2, Zenodo*, doi: [10.5281/zenodo.4309220](https://doi.org/10.5281/zenodo.4309220)
- Viets, A., et al. 2018, *Class. Quant. Grav.*, 35, 095015, doi: [10.1088/1361-6382/aab658](https://doi.org/10.1088/1361-6382/aab658)
- Virtanen, P., Gommers, R., Oliphant, T. E., et al. 2020, *Nature Methods*, 17, 261, doi: <https://doi.org/10.1038/s41592-019-0686-2>
- Wang, Y., Lo, R. K. L., Li, A. K. Y., & Chen, Y. 2021. <https://arxiv.org/abs/2101.08264>
- Wang, Y., Stebbins, A., & Turner, E. L. 1996, *Phys. Rev. Lett.*, 77, 2875, doi: [10.1103/PhysRevLett.77.2875](https://doi.org/10.1103/PhysRevLett.77.2875)
- Waskom, M., Botvinnik, O., Ostblom, J., et al. 2020, *mwaskom/seaborn: v0.10.1 (April 2020), v0.10.1, Zenodo*, doi: [10.5281/zenodo.3767070](https://doi.org/10.5281/zenodo.3767070)
- Wong, K. W. K., Breivik, K., Kremer, K., & Callister, T. 2020. <https://arxiv.org/abs/2011.03564>
- Yu, H., Zhang, P., & Wang, F.-Y. 2020, *Mon. Not. Roy. Astron. Soc.*, 497, 204, doi: [10.1093/mnras/staa1952](https://doi.org/10.1093/mnras/staa1952)
- Zevin, M., Bavera, S. S., Berry, C. P. L., et al. 2021, *Astrophys. J.*, 910, 152, doi: [10.3847/1538-4357/abe40e](https://doi.org/10.3847/1538-4357/abe40e)

THE LIGO SCIENTIFIC COLLABORATION AND THE VIRGO COLLABORATION, R. ABBOTT,¹ T. D. ABBOTT,² S. ABRAHAM,³ F. ACERNESE,^{4,5} K. ACKLEY,⁶ A. ADAMS,⁷ C. ADAMS,⁸ R. X. ADHIKARI,¹ V. B. ADYA,⁹ C. AFFELDT,^{10,11} D. AGARWAL,³ M. AGATHOS,^{12,13} K. AGATSUMA,¹⁴ N. AGGARWAL,¹⁵ O. D. AGUIAR,¹⁶ L. AIELLO,^{17,18,19} A. AIN,^{20,21} P. AJITH,²² K. M. ALEMAN,²³ G. ALLEN,²⁴ A. ALLOCCA,^{25,5} P. A. ALTIN,⁹ A. AMATO,²⁶ S. ANAND,¹ A. ANANYEVA,¹ S. B. ANDERSON,¹ W. G. ANDERSON,²⁷ S. V. ANGELOVA,²⁸ S. ANSOLDI,^{29,30} J. M. ANTELLIS,³¹ S. ANTIER,³² S. APPERT,¹ K. ARAI,¹ M. C. ARAYA,¹ J. S. AREEDA,²³ M. ARÈNE,³² N. ARNAUD,^{33,34} S. M. ARONSON,³⁵ K. G. ARUN,³⁶ Y. ASALI,³⁷ G. ASHTON,⁶ S. M. ASTON,⁸ P. ASTONE,³⁸ F. AUBIN,³⁹ P. AUFMUTH,^{10,11} K. AULTONEAL,³¹ C. AUSTIN,² S. BABAK,³² F. BADARACCO,^{18,19} M. K. M. BADER,⁴⁰ S. BAE,⁴¹ A. M. BAER,⁷ S. BAGNASCO,⁴² Y. BAI,¹ J. BAIRD,³² M. BALL,⁴³ G. BALLARDIN,³⁴ S. W. BALLMER,⁴⁴ M. BALS,³¹ A. BALSAMO,⁷ G. BALTUS,⁴⁵ S. BANAGIRI,⁴⁶ D. BANKAR,³ R. S. BANKAR,³ J. C. BARAYOGA,¹ C. BARBIERI,^{47,48,49} B. C. BARISH,¹ D. BARKER,⁵⁰ P. BARNEO,⁵¹ F. BARONE,^{52,5} B. BARR,⁵³ L. BARSOTTI,⁵⁴ M. BARSUGLIA,³² D. BARTA,⁵⁵ J. BARTLETT,⁵⁰ M. A. BARTON,⁵³ I. BARTOS,³⁵ R. BASSIRI,⁵⁶ A. BASTI,^{21,20} M. BAWAJ,^{57,58} J. C. BAYLEY,⁵³ A. C. BAYLOR,²⁷ M. BAZZAN,^{59,60} B. BÉCSY,⁶¹ V. M. BEDAKIHALE,⁶² M. BEJGER,⁶³ I. BELAHCENE,³³ V. BENEDETTO,⁶⁴ D. BENIWAL,⁶⁵ M. G. BENJAMIN,³¹ T. F. BENNETT,⁶⁶ J. D. BENTLEY,¹⁴ M. BENYAALA,²⁸ F. BERGAMIN,^{10,11} B. K. BERGER,⁵⁶ S. BERNUZZI,¹³ C. P. L. BERRY,^{53,15} D. BERSANETTI,⁶⁷ A. BERTOLINI,⁴⁰ J. BETZWIESER,⁸ R. BHANDARE,⁶⁸ A. V. BHANDARI,³ D. BHATTACHARJEE,⁶⁹ S. BHAUMIK,³⁵ J. BIDLER,²³ I. A. BILENKO,⁷⁰ G. BILLINGSLEY,¹ R. BIRNEY,⁷¹ O. BIRNHOLTZ,⁷² S. BISCANS,^{1,54} M. BISCHI,^{73,74} S. BISCOVEANU,⁵⁴ A. BISHT,^{10,11} B. BISWAS,³ M. BITOSSI,^{34,20} M.-A. BIZOUARD,⁷⁵ J. K. BLACKBURN,¹ J. BLACKMAN,⁷⁶ C. D. BLAIR,^{77,8} D. G. BLAIR,⁷⁷ R. M. BLAIR,⁵⁰ F. BOBBA,^{78,79} N. BODE,^{10,11} M. BOER,⁷⁵ G. BOGAERT,⁷⁵ M. BOLDRINI,^{80,38} F. BONDU,⁸¹ E. BONILLA,⁵⁶ R. BONNAND,³⁹ P. BOOKER,^{10,11} B. A. BOOM,⁴⁰ R. BORK,¹ V. BOSCHI,²⁰ N. BOSE,⁸² S. BOSE,³ V. BOSSILKOV,⁷⁷ V. BOUDART,⁴⁵ Y. BOUFFANAIS,⁴ A. BOZZI,³⁴ C. BRADASCIA,²⁰ P. R. BRADY,²⁷ A. BRAMLEY,¹ A. BRANCH,⁸ M. BRANCHESI,^{18,19} J. E. BRAU,⁴³ M. BRESCHI,¹³ T. BRIANT,⁸³ J. H. BRIGGS,⁵³ A. BRILLET,⁷⁵ M. BRINKMANN,^{10,11} P. BROCKILL,²⁷ A. F. BROOKS,¹ J. BROOKS,³⁴ D. D. BROWN,⁶⁵ S. BRUNETT,¹ G. BRUNO,⁸⁴ R. BRUNTZ,⁷ J. BRYANT,¹⁴ A. BUIKEMA,⁵⁴ T. BULIK,⁸⁵ H. J. BULTEN,^{40,86} A. BUONANNO,^{87,88} R. BUSCICCHIO,¹⁴ D. BUSKULIC,³⁹ R. L. BYER,⁵⁶ L. CADONATI,⁸⁹ M. CAESAR,⁹⁰ G. CAGNOLI,²⁶ C. CAHILLANE,¹ H. W. CAIN III,² J. CALDERÓN BUSTILLO,⁹¹ J. D. CALLAGHAN,⁵³ T. A. CALLISTER,^{92,93} E. CALLONI,^{25,5} J. B. CAMP,⁹⁴ M. CANEPA,^{95,67} M. CANNACCIUOLO,⁷⁸ K. C. CANNON,⁹⁶ H. CAO,⁶⁵ J. CAO,⁹⁷ E. CAPOTE,²³ G. CARAPPELLA,^{78,79} F. CARBOGNANI,³⁴ J. B. CARLIN,⁹⁸ M. F. CARNEY,¹⁵ M. CARPINELLI,^{99,100} G. CARULLO,^{21,20} T. L. CARVER,¹⁷ J. CASANUEVA DIAZ,³⁴ C. CASENTINI,^{101,102} G. CASTALDI,¹⁰³ S. CAUDILL,^{40,104} M. CAVAGLIA,⁶⁹ F. CAVALIERI,³³ R. CAVALIERI,³⁴ G. CELLA,²⁰ P. CERDÁ-DURÁN,¹⁰⁵ E. CESARINI,¹⁰² W. CHAIBI,⁷⁵ K. CHAKRAVARTI,³ B. CHAMPION,¹⁰⁶ C.-H. CHAN,¹⁰⁷ C. CHAN,⁹⁶ C. L. CHAN,⁹¹ K. CHANDRA,⁸² P. CHANIAL,³⁴ S. CHAO,¹⁰⁷ P. CHARLTON,¹⁰⁸ E. A. CHASE,¹⁵ E. CHASSANDE-MOTTIN,³² D. CHATTERJEE,²⁷ M. CHATURVEDI,⁶⁸ A. CHEN,⁹¹ H. Y. CHEN,¹⁰⁹ J. CHEN,¹⁰⁷ X. CHEN,⁷⁷ Y. CHEN,⁷⁶ Z. CHEN,¹⁷ H. CHENG,³⁵ C. K. CHEONG,⁹¹ H. Y. CHEUNG,⁹¹ H. Y. CHIA,³⁵ F. CHIADINI,^{110,79} R. CHIERICI,¹¹¹ A. CHINCARINI,⁶⁷ M. L. CHIOFALO,^{21,20} A. CHIUMMO,³⁴ G. CHO,¹¹² H. S. CHO,¹¹³ S. CHOATE,⁹⁰ R. K. CHOUDHARY,⁷⁷ S. CHOUDHARY,³ N. CHRISTENSEN,⁷⁵ Q. CHU,⁷⁷ S. CHUA,⁸³ K. W. CHUNG,¹¹⁴ G. CIANI,^{59,60} P. CIECIELAG,⁶³ M. CIEŚLAR,⁶³ M. CIFALDI,^{101,102} A. A. CIOBANU,⁶⁵ R. CIOLFI,^{115,60} F. CIPRIANO,⁷⁵ A. CIRONE,^{95,67} F. CLARA,⁵⁰ E. N. CLARK,¹¹⁶ J. A. CLARK,⁸⁹ L. CLARKE,¹¹⁷ P. CLEARWATER,⁹⁸ S. CLESSE,¹¹⁸ F. CLEVA,⁷⁵ E. COCCIA,^{18,19} P.-F. COHADON,⁸³ D. E. COHEN,³³ L. COHEN,² M. COLLEONI,¹¹⁹ C. G. COLLETTE,¹²⁰ M. COLPI,^{47,48} C. M. COMPTON,⁵⁰ M. CONSTANCIO JR.,¹⁶ L. CONTI,⁶⁰ S. J. COOPER,¹⁴ P. CORBAN,⁸ T. R. CORBITT,² I. CORDERO-CARRIÓN,¹²¹ S. COREZZI,^{58,57} K. R. CORLEY,³⁷ N. CORNISH,⁶¹ D. CORRE,³³ A. CORSI,¹²² S. CORTESE,³⁴ C. A. COSTA,¹⁶ R. COTESTA,⁸⁸ M. W. COUGHLIN,⁴⁶ S. B. COUGHLIN,^{15,17} J.-P. COULON,⁷⁵ S. T. COUNTRYMAN,³⁷ B. COUSINS,¹²³ P. COUVARES,¹ P. B. COVAS,¹¹⁹ D. M. COWARD,⁷⁷ M. J. COWART,⁸ D. C. COYNE,¹ R. COYNE,¹²⁴ J. D. E. CREIGHTON,²⁷ T. D. CREIGHTON,¹²⁵ A. W. CRISWELL,⁴⁶ M. CROQUETTE,⁸³ S. G. CROWDER,¹²⁶ J. R. CUDELL,⁴⁵ T. J. CULLEN,² A. CUMMING,⁵³ R. CUMMINGS,⁵³ E. CUOCO,^{34,127,20} M. CURYLO,⁸⁵ T. DAL CANTON,^{88,33} G. DÁLYA,¹²⁸ A. DANA,⁵⁶ L. M. DANESHGARANBAJASTANI,⁶⁶ B. D'ANGELO,^{95,67} S. L. DANILISHIN,¹²⁹ S. D'ANTONIO,¹⁰² K. DANZMANN,^{10,11} C. DARSOW-FROMM,¹³⁰ A. DASGUPTA,⁶² L. E. H. DATHIRI,³⁴ V. DATILLO,³⁴ I. DAVE,⁶⁸ M. DAVIER,³³ G. S. DAVIES,^{131,132} D. DAVIS,¹ E. J. DAW,¹³³ R. DEAN,⁹⁰ D. DEBRA,⁵⁶ M. DEENADAYALAN,³ J. DEGALLAIX,¹³⁴ M. DE LAURENTIS,^{25,5} S. DELÉGLISE,⁸³ V. DEL FAVERO,¹⁰⁶ F. DE LILLO,⁸⁴ N. DE LILLO,⁵³ W. DEL POZZO,^{21,20} L. M. DEMARCHI,¹⁵ F. DE MATTEIS,^{101,102} V. D'EMILIO,¹⁷ N. DEMOS,⁵⁴ T. DENT,¹³¹ A. DEPASSE,⁸⁴ R. DE PIETRI,^{135,136} R. DE ROSA,^{25,5} C. DE ROSSI,³⁴ R. DE SALVO,¹⁰³ R. DE SIMONE,¹¹⁰ S. DHURANDHAR,³ M. C. DÍAZ,¹²⁵ M. DIAZ-ORTIZ JR.,³⁵ N. A. DIDIO,⁴⁴ T. DIETRICH,⁸⁸ L. DI FIORE,⁵ C. DI FRONZO,¹⁴ C. DI GIORGIO,^{78,79} F. DI GIOVANNI,¹⁰⁵ T. DI GIROLAMO,^{25,5} A. DI LIETO,^{21,20} B. DING,¹²⁰ S. DI PACE,^{80,38} I. DI PALMA,^{80,38} F. DI RENZO,^{21,20} A. K. DIVAKARLA,³⁵ A. DMITRIEV,¹⁴ Z. DOCTOR,⁴³ L. D'ONOFRIO,^{25,5} F. DONOVAN,⁵⁴ K. L. DOOLEY,¹⁷ S. DORAVARI,³ I. DORRINGTON,¹⁷ M. DRAGO,^{18,19} J. C. DRIGGERS,⁵⁰ Y. DRORI,¹ Z. DU,⁹⁷ J.-G. DUCOIN,³³ P. DUPEJ,⁵³ O. DURANTE,^{78,79} D. D'URSO,^{99,100} P.-A. DUVERNE,³³ S. E. DWYER,⁵⁰ P. J. EASTER,⁶ M. EBERSOLD,¹³⁷ G. EDDOLLS,⁵³ B. EDELMAN,⁴³ T. B. EDO,^{1,133} O. EDY,¹³² A. EFFLER,⁸ J. EICHHOLZ,⁹ S. S. EIKENBERRY,³⁵ M. EISENMANN,³⁹ R. A. EISENSTEIN,⁵⁴ A. EJLLI,¹⁷ L. ERRICO,^{25,5} R. C. ESSICK,¹⁰⁹ H. ESTELLÉS,¹¹⁹ D. ESTEVEZ,¹³⁸ X. ÉTIENNE,¹³⁹ T. ETZEL,¹ M. EVANS,⁵⁴ T. M. EVANS,⁸ B. E. EWING,¹²³ J. M. EZQUIAGA,¹⁰⁹ V. FAFONE,^{101,102,18} H. FAIR,⁴⁴ S. FAIRHURST,¹⁷ X. FAN,⁹⁷ A. M. FARAH,¹⁰⁹ S. FARINON,⁶⁷ B. FARR,⁴³ W. M. FARR,^{92,93} N. W. FARRROW,⁶ E. J. FAUCHON-JONES,¹⁷ M. FAVATA,¹⁴⁰ M. FAYS,^{45,133} M. FAZIO,¹⁴¹ J. FEICHT,¹ M. M. FEJER,⁵⁶ F. FENG,³² E. FENYVESI,^{55,142} D. L. FERGUSON,⁸⁹ A. FERNANDEZ-GALIANO,⁵⁴ I. FERRANTE,^{21,20} T. A. FERREIRA,¹⁶ F. FIDECARO,^{21,20} P. FIGURA,⁸⁵ I. FIORI,³⁴ M. FISHBACH,^{15,109} R. P. FISHER,⁷ R. FITTIPALDI,^{143,79} V. FIUMARA,^{144,79} R. FLAMINIO,^{39,145} E. FLODEN,⁴⁶ E. FLYNN,²³ H. FONG,⁹⁶ J. A. FONT,^{105,146} B. FORMAL,¹⁴⁷ P. W. F. FORSYTH,⁹ A. FRANKE,¹³⁰ S. FRASCA,^{80,38} F. FRASCONI,²⁰ C. FREDERICK,¹⁴⁸ Z. FREI,¹²⁸ A. FREISE,¹⁴⁹ R. FREY,⁴³ P. FRITSCHEL,⁵⁴ V. V. FROLOV,⁸ G. G. FRONZÉ,⁴² P. FULDA,³⁵ M. FYFFE,⁸ H. A. GABBARD,⁵³ B. U. GADRE,⁸⁸ S. M. GAEBEL,¹⁴ J. R. GAIR,⁸⁸ J. GAIS,⁹¹ S. GALADAGE,⁶ R. GAMBA,¹³ D. GANAPATHY,⁵⁴ A. GANGULY,²² S. G. GAONKAR,³ B. GARAVENTA,^{67,95} C. GARCÍA-NÚÑEZ,⁷¹ C. GARCÍA-QUIRÓS,¹¹⁹ F. GARUFI,^{25,5} B. GATELEY,⁵⁰ S. AUDIO,³¹ V. GAYATHRI,³⁵ G. GEMME,⁶⁷ A. GENNAI,²⁰ J. GEORGE,⁶⁸ L. GERGELY,¹⁵⁰ P. GEWECKE,¹³⁰ S. GHONGE,⁸⁹ ABHIRUP GHOSH,⁸⁸ ARCHISMAN GHOSH,¹⁵¹ SHAON GHOSH,^{27,140} SHROBANA GHOSH,¹⁷ SOURATH GHOSH,³⁵ B. GIACOMAZZO,^{47,48,49} L. GIACOPPO,^{80,38} J. A. GIAIME,^{2,8} K. D. GIARDINA,⁸ D. R. GIBSON,⁷¹ C. GIER,²⁸ M. GIESLER,⁷⁶ P. GIRI,^{20,21} F. GISSI,⁶⁴ J. GLANZER,² A. E. GLECKL,²³ P. GODWIN,¹²³ E. GOETZ,¹⁵² R. GOETZ,³⁵ N. GOHLKE,^{10,11} B. GONCHAROV,⁶ G. GONZÁLEZ,² A. GOPAKUMAR,¹⁵³ M. GOSSELIN,³⁴ R. GOUATY,³⁹ S. GOYAL,²² B. GRACE,⁹ A. GRATO,^{154,5} M. GRANATA,¹³⁴ V. GRANATA,⁷⁸ A. GRANT,⁵³ S. GRAS,⁵⁴ P. GRASSIA,¹ C. GRAY,⁵⁰ R. GRAY,⁵³ G. GRECO,⁵⁷ A. C. GREEN,³⁵ R. GREEN,¹⁷ A. M. GRETARSSON,³¹ E. M. GRETARSSON,³¹ D. GRIFFITH,¹ W. GRIFFITHS,¹⁷ H. L. GRIGGS,⁸⁹ G. GRIGNANI,^{58,57} A. GRIMALDI,^{155,156} E. GRIMES,³¹ S. J. GRIMM,^{18,19} H. GROTE,¹⁷ S. GRUNEWALD,⁸⁸ P. GRUNING,³³ J. G. GUERRERO,²³ G. M. GUIDI,^{73,74} A. R. GUIMARAES,² G. GUIXÉ,⁵¹ H. K. GULATI,⁶² H.-K. GUO,¹⁴⁷ Y. GUO,⁴⁰ ANCHAL GUPTA,¹ ANURADHA GUPTA,¹⁵⁷ P. GUPTA,^{40,104} E. K. GUSTAFSON,¹ R. GUSTAFSON,¹⁵⁸ F. GUZMAN,¹¹⁶ L. HAEGEL,³² O. HALIM,^{159,30} E. D. HALL,⁵⁴ E. Z. HAMILTON,¹⁷ G. HAMMOND,⁵³ M. HANEY,¹³⁷ J. HANKS,⁵⁰ C. HANNA,¹²³ M. D. HANNAM,¹⁷ O. A. HANNUKSELA,^{104,40,91} H. HANSEN,⁵⁰ T. J. HANSEN,³¹ J. HANSON,⁸ T. HARDER,⁷⁵ T. HARDWICK,² K. HARRIS,^{40,104,22} J. HARMS,^{18,19} G. M. HARRY,¹⁶⁰ I. W. HARRY,¹³² D. HARTWIG,¹³⁰ B. HASKELL,⁶³ R. K. HASSKEW,⁸ C.-J. HASTER,⁵⁴ K. HAUGHIAN,⁵³ F. J. HAYES,⁵³ J. HEALY,¹⁰⁶ A. HEIDMANN,⁸³ M. C. HEINTZE,⁸ J. HEINZE,^{10,11} J. HEINZEL,¹⁶¹ H. HEITMANN,⁷⁵ F. HELLMAN,¹⁶² P. HELLO,³³ A. F. HELMLING-CORNELL,⁴³ G. HEMMING,³⁴ M. HENDRY,⁵³ I. S. HENG,⁵³ E. HENNES,⁴⁰ J. HENNIG,^{10,11} M. H. HENNIG,^{10,11} F. HERNANDEZ VIVANCO,⁶ M. HEURS,^{10,11} S. HILD,^{129,40} P. HILL,²⁸ A. S. HINES,¹¹⁶ S. HOCHHEIM,^{10,11} D. HOFMAN,¹³⁴ J. N. HOHMANN,¹³⁰ A. M. HOGGADO,²⁴ N. A. HOLLAND,⁹ I. J. HOLLOWAY,¹³³ Z. J. HOLMES,⁶⁵ K. HOLT,⁸ D. E. HOLZ,¹⁰⁹ P. HOPKINS,¹⁷ J. HOUGH,⁵³ E. J. HOWELL,⁷⁷ C. G. HOY,¹⁷ D. HOYLAND,¹⁴ A. HREIBI,^{10,11}

- Y. HSU,¹⁰⁷ Y. HUANG,⁵⁴ M. T. HÜBNER,⁶ A. D. HUDDART,¹¹⁷ E. A. HUERTA,²⁴ B. HUGHEY,³¹ V. HUI,³⁹ S. HUSA,¹¹⁹ S. H. HUTTNER,⁵³ R. HUXFORD,¹²³ T. HUYNH-DINH,⁸ B. IDZKOWSKI,⁸⁵ A. IESS,^{101,102} H. INCHAUSPE,³⁵ C. INGRAM,⁶⁵ G. INTINI,^{80,38} M. ISI,⁵⁴ K. ISLEIF,¹³⁰ B. R. IYER,²² V. JABERIANHAMEDAN,⁷⁷ T. JACQMIN,⁸³ S. J. JADHAV,¹⁶³ S. P. JADHAV,³ A. L. JAMES,¹⁷ A. Z. JAN,¹⁰⁶ K. JANI,⁸⁹ J. JANQUART,^{40,104} K. JANSSENS,¹⁶⁴ N. N. JANTHALUR,¹⁶³ P. JARANOWSKI,¹⁶⁵ D. JARIWALA,³⁵ R. JAUME,¹¹⁹ A. C. JENKINS,¹¹⁴ M. JEUNON,⁴⁶ W. JIA,⁵⁴ J. JIANG,³⁵ G. R. JOHNS,⁷ A. W. JONES,⁷⁷ D. I. JONES,¹⁶⁶ J. D. JONES,⁵⁰ P. JONES,¹⁴ R. JONES,⁵³ R. J. G. JONKER,⁴⁰ L. JU,⁷⁷ J. JUNKER,^{10,11} C. V. KALAGHATGI,¹⁷ V. KALOGERA,¹⁵ B. KAMAI,¹ S. KANDHASAMY,³ G. KANG,⁴¹ J. B. KANNER,¹ Y. KAO,¹⁰⁷ S. J. KAPADIA,²² D. P. KAPASI,⁹ S. KARAT,¹ C. KARATHANASIS,¹⁶⁷ S. KARKI,⁶⁹ R. KASHYAP,¹²³ M. KASPRZACK,¹ W. KASTAUN,^{10,11} S. KATSANEVAS,³⁴ E. KATSAVOUNIDIS,⁵⁴ W. KATZMAN,⁸ T. KAUR,⁷⁷ K. KAWABE,⁵⁰ F. KÉFÉLIAN,⁷⁵ D. KEITEL,¹¹⁹ J. S. KEY,¹⁶⁸ S. KHADKA,⁵⁶ F. Y. KHALILI,⁷⁰ I. KHAN,^{18,102} S. KHAN,¹⁷ E. A. KHAZANOV,¹⁶⁹ N. KHETAN,^{18,19} M. KHURSHEED,⁶⁸ N. KIBBUNCHOO,⁹ C. KIM,¹⁷⁰ J. C. KIM,¹⁷¹ K. KIM,¹⁷² W. S. KIM,¹⁷³ Y.-M. KIM,¹⁷⁴ C. KIMBALL,¹⁵ P. J. KING,⁵⁰ M. KINLEY-HANLON,⁵³ R. KIRCHHOFF,^{10,11} J. S. KISSEL,⁵⁰ L. KLEYBOLTE,¹³⁰ S. KLIMENKO,³⁵ A. M. KNEE,¹⁵² T. D. KNOWLES,¹³⁹ E. KNYAZEV,⁵⁴ P. KOCH,^{10,11} G. KOEKOEK,^{40,129} S. KOLEY,⁴⁰ P. KOLITSIDOU,¹⁷ M. KOLSTEIN,¹⁶⁷ K. KOMORI,⁵⁴ V. KONDRASHOV,¹ A. KONTOS,¹⁷⁵ N. KOPER,^{10,11} M. KOROBKO,¹³⁰ M. KOVALAM,⁷⁷ D. B. KOZAK,¹ V. KRINGEL,^{10,11} N. V. KRISHNENDU,^{10,11} A. KRÓLAK,^{176,177} G. KUEHN,^{10,11} F. KUEI,¹⁰⁷ A. KUMAR,¹⁶³ P. KUMAR,¹⁷⁸ RAHUL KUMAR,⁵⁰ RAKESH KUMAR,⁶² K. KUNS,⁵⁴ S. KWANG,²⁷ D. LAGHI,^{21,20} E. LALANDE,¹⁷⁹ T. L. LAM,⁹¹ A. LAMBERTS,^{75,180} M. LANDRY,⁵⁰ B. B. LANE,⁵⁴ R. N. LANG,⁵⁴ J. LANGE,^{181,106} B. LANTZ,⁵⁶ I. LA ROSA,³⁹ A. LARTAUD-VOLLARD,³³ P. D. LASKY,⁶ M. LAXEN,⁸ A. LAZZARINI,¹ C. LAZZARO,^{59,60} P. LEACI,^{80,38} S. LEAVEY,^{10,11} Y. K. LECOUCHE,⁵⁰ H. M. LEE,¹⁷² H. W. LEE,¹⁷¹ J. LEE,¹¹² K. LEE,⁵⁶ J. LEHMANN,^{10,11} A. LEMAITRE,¹⁸² E. LEON,²³ N. LEROY,³³ N. LETENDRE,³⁹ Y. LEVIN,⁶ J. N. LEVITON,¹⁵⁸ A. K. Y. LI,¹ B. LI,¹⁰⁷ J. LI,¹⁵ T. G. F. LI,⁹¹ X. LI,⁷⁶ F. LINDE,^{183,40} S. D. LINKER,⁶⁶ J. N. LINLEY,⁵³ T. B. LITTENBERG,¹⁸⁴ J. LIU,^{10,11} K. LIU,¹⁰⁷ X. LIU,²⁷ M. LLORENS-MONTEAGUDO,¹⁰⁵ R. K. L. LO,¹ A. LOCKWOOD,¹⁸⁵ M. L. LOLLIE,² L. T. LONDON,⁵⁴ A. LONGO,^{186,187} D. LOPEZ,¹³⁷ M. LORENZINI,^{101,102} V. LORIETTE,¹⁸⁸ M. LORMAND,⁸ G. LOSURDO,²⁰ J. D. LOUGH,^{10,11} C. O. LOUSTO,¹⁰⁶ G. LOVELACE,²³ H. LÜCK,^{10,11} D. LUMACA,^{101,102} A. P. LUNDGREN,¹³² R. MACAS,¹⁷ M. MACINNIS,⁵⁴ D. M. MACLEOD,¹⁷ I. A. O. MACMILLAN,¹ A. MACQUET,⁷⁵ I. MAGAÑA HERNANDEZ,²⁷ F. MAGAÑA-SANDOVAL,³⁵ C. MAGAZZÙ,²⁰ R. M. MAGEE,¹²³ R. MAGGIORE,¹⁴ E. MAJORANA,^{80,38} C. MAKAREM,¹ I. MAKSIMOVIC,¹⁸⁸ S. MALIAKAL,¹ A. MALIK,⁶⁸ N. MAN,⁷⁵ V. MANDIC,⁴⁶ V. MANGANO,^{80,38} J. L. MANGO,¹⁸⁹ G. L. MANSSELL,^{50,54} M. M. MANSKE,²⁷ M. MANTOVANI,³⁴ M. MAPELLI,^{59,60} F. MARCHESONI,^{190,57} F. MARION,³⁹ Z. MARK,⁷⁶ S. MÁRKA,³⁷ Z. MÁRKA,³⁷ C. MARKAKIS,¹² A. S. MARKOSYAN,⁵⁶ A. MARKOWITZ,¹ E. MAROS,¹ A. MARQUINA,¹²¹ S. MARSAT,³² F. MARTELLI,^{73,74} I. W. MARTIN,⁵³ R. M. MARTIN,¹⁴⁰ M. MARTINEZ,¹⁶⁷ V. MARTINEZ,²⁶ K. MARTINOVIC,¹¹⁴ D. V. MARTYNOV,¹⁴ E. J. MARX,⁵⁴ H. MASALEHDAN,¹³⁰ K. MASON,⁵⁴ E. MASSERA,¹³³ A. MASSEROT,³⁹ T. J. MASSINGER,⁵⁴ M. MASSO-REID,⁵³ S. MASTROGIOVANNI,³² A. MATAS,⁸⁸ M. MATEU-LUCENA,¹¹⁹ F. MATHICHARD,^{1,54} M. MATIUSHECHKINA,^{10,11} N. MAVALVALA,⁵⁴ J. J. McCANN,⁷⁷ R. MCCARTHY,⁵⁰ D. E. MCCLELLAND,⁹ P. McCLINCY,¹²³ S. McCORMICK,⁸ L. McCULLER,⁵⁴ G. I. McGHEE,⁵³ S. C. McGUIRE,¹⁹¹ C. McISAAC,¹³² J. McIVER,¹⁵² D. J. McMANUS,⁹ T. McRAE,⁹ S. T. McWILLIAMS,¹³⁹ D. MEACHER,²⁷ M. MEHMET,^{10,11} A. K. MEHTA,⁸⁸ A. MELATOS,⁹⁸ D. A. MELCHOR,²³ G. MENDELL,⁵⁰ A. MENENDEZ-VÁZQUEZ,¹⁶⁷ C. S. MENONI,¹⁴¹ R. A. MERCER,²⁷ L. MERENI,¹³⁴ K. MERFELD,⁴³ E. L. MERILH,⁵⁰ J. D. MERRITT,⁴³ M. MERZOUGUI,⁷⁵ S. MESHKOV,^{1,*} C. MESSENGER,⁵³ C. MESSICK,¹⁸¹ P. M. MEYERS,⁹⁸ F. MEYLAHN,^{10,11} A. MHASKE,³ A. MIANI,^{155,156} H. MIAO,¹⁴ I. MICHALOLIKOS,³⁵ C. MICHEL,¹³⁴ H. MIDDLETON,⁹⁸ L. MILANO,²⁵ A. L. MILLER,^{84,35} M. MILLHOUSE,⁹⁸ J. C. MILLS,¹⁷ E. MILOTTI,^{159,30} M. C. MILOVICH-GOFF,⁶⁶ O. MINAZZOLI,^{75,192} Y. MINENKOV,¹⁰² LL. M. MIR,¹⁶⁷ A. MISHKIN,³⁵ C. MISHRA,¹⁹³ T. MISHRA,³⁵ T. MISTRY,¹³³ S. MITRA,³ V. P. MITROFANOV,⁷⁰ G. MITSSELMACHER,³⁵ R. MITTLEMAN,⁵⁴ GEOFFREY MO,⁵⁴ K. MOGUSHI,⁶⁹ S. R. P. MOHAPATRA,⁵⁴ S. R. MOHITE,²⁷ I. MOLINA,²³ M. MOLINA-RUIZ,¹⁶² M. MONDIN,⁶⁶ M. MONTANI,^{73,74} C. J. MOORE,¹⁴ D. MORARU,⁵⁰ F. MORAWSKI,⁶³ A. MORE,³ C. MORENO,³¹ G. MORENO,⁵⁰ S. MORISAKI,⁹⁶ B. MOURS,¹³⁸ C. M. MOW-LOWRY,¹⁴ S. MOZZON,¹³² F. MUCIACCIA,^{80,38} ARUNAVA MUKHERJEE,^{194,53} D. MUKHERJEE,¹²³ SOMA MUKHERJEE,¹²⁵ SUBROTO MUKHERJEE,⁶² N. MUKUND,^{10,11} A. MULLAVEY,⁸ J. MUNCH,⁶⁵ E. A. MUÑOZ,⁴⁴ P. G. MURRAY,⁵³ R. MUSENICH,^{67,95} S. L. NADJI,^{10,11} A. NAGAR,^{42,195} I. NARDECCHIA,^{101,102} L. NATICCHIONI,³⁸ B. NAYAK,⁶⁶ R. K. NAYAK,¹⁹⁶ B. F. NEIL,⁷⁷ J. NEILSON,^{64,79} G. NELEMANS,¹⁹⁷ T. J. N. NELSON,⁸ M. NERY,^{10,11} A. NEUNZERT,¹⁶⁸ K. Y. NG,⁵⁴ S. W. S. NG,⁶⁵ C. NGUYEN,³² P. NGUYEN,⁴³ T. NGUYEN,⁵⁴ S. A. NICHOLS,² S. NISSANKE,^{198,40} F. NÓCERA,³⁴ M. NOH,¹⁵² M. NORMAN,¹⁷ C. NORTH,¹⁷ L. K. NUTTALL,¹³² J. OBERLING,⁵⁰ B. D. O'BRIEN,³⁵ J. O'DELL,¹¹⁷ G. OGANESYAN,^{18,19} J. J. OH,¹⁷³ S. H. OH,¹⁷³ F. OHME,^{10,11} H. OHTA,⁹⁶ M. A. OKADA,¹⁶ C. OLIVETTO,³⁴ R. ORAM,⁸ B. O'REILLY,⁸ R. G. ORMISTON,⁴⁶ N. D. ORMSBY,⁷ L. F. ORTEGA,³⁵ R. O'SHAUGHNESSY,¹⁰⁶ E. O'SHEA,¹⁷⁸ S. OSSOKINE,⁸⁸ C. OSTHELDER,¹ D. J. OTTAWAY,⁶⁵ H. OVERMIER,⁸ A. E. PACE,¹²³ G. PAGANO,^{21,20} M. A. PAGE,⁷⁷ G. PAGLIAROLI,^{18,19} A. PAI,⁸² S. A. PAI,⁶⁸ J. R. PALAMOS,⁴³ O. PALASHOV,¹⁶⁹ C. PALOMBA,³⁸ P. K. PANDA,¹⁶³ P. T. H. PANG,^{40,104} C. PANKOW,¹⁵ F. PANNARALE,^{80,38} B. C. PANT,⁶⁸ F. PAOLETTI,²⁰ A. PAOLI,³⁴ A. PAOLONE,^{38,199} W. PARKER,^{8,191} D. PASCUCCI,⁴⁰ A. PASQUALETTI,³⁴ R. PASSAQUETTI,^{21,20} D. PASSUELLO,²⁰ M. PATEL,⁷ B. PATRICELLI,^{34,20} E. PAYNE,⁶ T. C. PECHSIRI,³⁵ M. PEDRAZA,¹ M. PEGORARO,⁶⁰ A. PELE,⁸ S. PENN,²⁰⁰ A. PEREGO,^{155,156} A. PEREIRA,²⁶ T. PEREIRA,²⁰¹ C. J. PEREZ,⁵⁰ C. PÉRIGOIS,³⁹ A. PERRECA,^{155,156} S. PERRIÉS,¹¹¹ J. PETERMANN,¹³⁰ D. PETTERSON,¹ H. P. PFEIFFER,⁸⁸ K. A. PHAM,⁴⁶ K. S. PHUKON,^{40,183,3} O. J. PICCINI,³⁸ M. PICHOT,⁷⁵ M. PIENIBENE,^{21,20} F. PIERGIOVANNI,^{73,74} L. PIERINI,^{80,38} V. PIERRO,^{64,79} G. PILANT,³⁴ F. PILO,²⁰ L. PINARD,¹³⁴ I. M. PINTO,^{64,79,202} B. J. PIOTRKOWSKI,²⁷ K. PIOTRKOWSKI,⁸⁴ M. PIRELLO,⁵⁰ M. PITKIN,²⁰³ E. PLACIDI,^{80,38} W. PLASTINO,^{186,187} C. PLUCHAR,¹¹⁶ R. POGGIANI,^{21,20} E. POLINI,³⁹ D. Y. T. PONG,⁹¹ S. PONRATHNAM,³ P. POPOLIZIO,³⁴ E. K. PORTER,³² J. POWELL,²⁰⁴ M. PRACCHIA,³⁹ T. PRADIER,¹³⁸ A. K. PRAJAPATI,⁶² K. PRASAI,⁵⁶ R. PRASANNA,¹⁶³ G. PRATTEN,¹⁴ T. PRESTEGARD,²⁷ M. PRINCEPI,^{64,202,79} G. A. PRODI,^{205,156} L. PROKHOROV,¹⁴ P. PROPOSITO,^{101,102} L. PRUDENZI,⁸⁸ A. PUECHER,^{40,104} M. PUNTURO,⁵⁷ F. PUOSI,^{20,21} P. PUPPO,³⁸ M. PÜRNER,⁸⁸ H. QI,¹⁷ V. QUETSCHKE,¹²⁵ P. J. QUINONEZ,³¹ R. QUITZOW-JAMES,⁶⁹ F. J. RAAB,⁵⁰ G. RAAIJMAKERS,^{198,40} H. RADKINS,⁵⁰ N. RADULESCO,⁷⁵ P. RAFFAI,¹²⁸ S. X. RAIL,¹⁷⁹ S. RAJA,⁶⁸ C. RAJAN,⁶⁸ K. E. RAMIREZ,¹²⁵ T. D. RAMIREZ,²³ A. RAMOS-BUADES,⁸⁸ J. RANA,¹²³ P. RAPAGNANI,^{80,38} U. D. RAPOL,²⁰⁶ B. RATTO,³¹ V. RAYMOND,¹⁷ N. RAZA,¹⁵² M. RAZZANO,^{21,20} J. READ,²³ L. A. REES,¹⁶⁰ T. REGIMBAU,³⁹ L. REI,⁶⁷ S. REID,²⁸ D. H. REITZE,^{1,35} P. RELTON,¹⁷ P. RETTEGNO,^{207,42} F. RICCI,^{80,38} C. J. RICHARDSON,³¹ J. W. RICHARDSON,¹ L. RICHARDSON,¹¹⁶ P. M. RICKER,²⁴ G. RIEMENSCHNEIDER,^{207,42} K. RILES,¹⁵⁸ M. RIZZO,¹⁵ N. A. ROBERTSON,^{1,53} R. ROBBIE,¹ F. ROBINET,³³ A. ROCCHI,¹⁰² J. A. ROCHA,²³ S. RODRIGUEZ,²³ R. D. RODRIGUEZ-SOTO,³¹ L. ROLLAND,³⁹ J. G. ROLLINS,¹ V. J. ROMA,⁴³ M. ROMANELLI,⁸¹ R. ROMANO,^{4,5} C. L. ROMEL,⁵⁰ A. ROMERO,¹⁶⁷ I. M. ROMERO-SHAW,⁶ J. H. ROMIE,⁸ C. A. ROSE,²⁷ D. ROSIŃSKA,⁸⁵ S. G. ROSOFSKY,²⁴ M. P. ROSS,¹⁸⁵ S. ROWAN,⁵³ S. J. ROWLINSON,¹⁴ SANTOSH ROY,³ SOUMEN ROY,²⁰⁸ D. ROZZA,^{99,100} P. RUGGI,³⁴ K. RYAN,⁵⁰ S. SACHDEV,¹²³ T. SADECKI,⁵⁰ J. SADIQ,¹³¹ M. SAKELLARIADOU,¹¹⁴ O. S. SALAFIA,^{49,48,47} L. SALCONI,³⁴ M. SALEEM,³⁶ F. SALEMI,^{155,156} A. SAMAJDAR,^{40,104} E. J. SANCHEZ,¹ J. H. SANCHEZ,²³ L. E. SANCHEZ,¹ N. SANCHIS-GUAL,²⁰⁹ J. R. SANDERS,²¹⁰ A. SANUY,⁵¹ T. R. SARAVANAN,³ N. SARIN,⁶ B. SASSOLAS,¹³⁴ H. SATARI,⁷⁷ B. S. SATHYAPRAKASH,^{123,17} O. SAUTER,^{35,39} R. L. SAVAGE,⁵⁰ V. SAVANT,³ D. SAWANT,⁸² H. L. SAWANT,³ S. SAYAH,¹³⁴ D. SCHAETZL,¹ M. SCHEEL,⁷⁶ J. SCHEUER,¹⁵ A. SCHINDLER-TYKA,³⁵ P. SCHMIDT,¹⁴ R. SCHNABEL,¹³⁰ M. SCHNEEWIND,^{10,11} R. M. S. SCHOFIELD,⁴³ A. SCHÖNBECK,¹³⁰ B. W. SCHULTE,^{10,11} B. F. SCHUTZ,^{17,10} E. SCHWARTZ,¹⁷ J. SCOTT,⁵³ S. M. SCOTT,⁹ M. SEGLAR-ARROYO,³⁹ E. SEIDEL,²⁴ D. SELLERS,⁸ A. S. SENGUPTA,²⁰⁸ N. SENNETT,⁸⁸ D. SENTENAC,³⁴ E. G. SEO,⁹¹ V. SEQUINO,^{25,5} A. SERGEEV,¹⁶⁹ Y. SETYAWATI,^{10,11} T. SHAFFER,⁵⁰ M. S. SHAHRIAR,¹⁵ B. SHAMS,¹⁴⁷ S. SHARIFI,² A. SHARMA,^{181,19} P. SHARMA,⁶⁸ P. SHAWHAN,⁸⁷ N. S. SHCHEBLANOV,¹⁸² H. SHEN,²⁴ M. SHIKAUCHI,⁹⁶ R. SHINK,¹⁷⁹ D. H. SHOEMAKER,⁵⁴ D. M. SHOEMAKER,¹⁸ K. SHUKLA,¹⁶² S. SHYAMSUNDAR,⁶⁸ M. SIENIAWSKA,⁸⁵ D. SIGG,⁵⁰ L. P. SINGER,⁹⁴ D. SINGH,¹²³ N. SINGH,⁸⁵ A. SINGHA,^{129,40} A. M. SINTES,¹¹⁹ V. SIPALA,^{99,100} V. SKLRIS,¹⁷ B. J. J. SLAGMOLEN,⁹ T. J. SLAVEN-BLAIR,⁷⁷

J. SMETANA,¹⁴ J. R. SMITH,²³ R. J. E. SMITH,⁶ S. N. SOMALA,²¹¹ E. J. SON,¹⁷³ K. SONI,³ S. SONI,² B. SORAZU,⁵³ V. SORDINI,¹¹¹ F. SORRENTINO,⁶⁷ N. SORRENTINO,^{21,20} R. SOULARD,⁷⁵ T. SOURADEEP,^{206,3} E. SOWELL,¹²² V. SPAGNUOLO,^{129,40} A. P. SPENCER,⁵³ M. SPERA,^{59,60} A. K. SRIVASTAVA,⁶² V. SRIVASTAVA,⁴⁴ K. STAATS,¹⁵ C. STACHIE,⁷⁵ D. A. STEER,³² J. STEINLECHNER,^{129,40} S. STEINLECHNER,^{129,40} D. J. STOPS,¹⁴ M. STOVER,¹⁴⁸ K. A. STRAIN,⁵³ L. C. STRANG,⁹⁸ G. STRATTA,^{212,74} A. STRUNK,⁵⁰ R. STURANI,²⁰¹ A. L. STUVER,⁹⁰ J. SÜDBECK,¹³⁰ S. SUDHAGAR,³ V. SUDHIR,⁵⁴ H. G. SUH,²⁷ T. Z. SUMMERSCALES,²¹³ H. SUN,⁷⁷ L. SUN,^{9,1} S. SUNIL,⁶² A. SUR,⁶³ J. SURESH,⁹⁶ P. J. SUTTON,¹⁷ B. L. SWINKELS,⁴⁰ M. J. SZCZEPANČYK,³⁵ P. SZEWczyk,⁸⁵ M. TACCA,⁴⁰ S. C. TAIT,⁵³ C. TALBOT,¹ A. J. TANASIECZUK,⁸⁴ D. B. TANNER,³⁵ D. TAO,¹ A. TAPIA,²³ E. N. TAPIA SAN MARTIN,⁴⁰ J. D. TASSON,¹⁶¹ R. TENORIO,¹¹⁹ L. TERKOWSKI,¹³⁰ M. TEST,²⁷ M. P. THIRUGNANASAMBANDAM,³ M. THOMAS,⁸ P. THOMAS,⁵⁰ J. E. THOMPSON,¹⁷ S. R. THONDAPU,⁶⁸ K. A. THORNE,⁸ E. THRANE,⁶ SHUBHANSU TIWARI,¹³⁷ SRISHTI TIWARI,¹⁵³ V. TIWARI,¹⁷ K. TOLAND,⁵³ A. E. TOLLEY,¹³² M. TONELLI,^{21,20} A. TORRES-FORNÉ,¹⁰⁵ C. I. TORRIE,¹ I. TOSTA E MELO,^{99,100} D. TÖYRÄ,⁹ A. TRAPANANTI,^{190,57} F. TRAVASSO,^{57,190} G. TRAYLOR,⁸ M. C. TRINGALI,³⁴ A. TRIPATHEE,¹⁵⁸ L. TROIANO,^{214,79} A. TROVATO,³² R. J. TRUDEAU,¹ D. S. TSAI,¹⁰⁷ D. TSAI,¹⁰⁷ K. W. TSANG,^{40,215,104} M. TSE,⁵⁴ R. TSO,⁷⁶ L. TSUKADA,⁹⁶ D. TSUNA,⁹⁶ T. TSUTSUI,⁹⁶ M. TURCONI,⁷⁵ A. S. UBHI,¹⁴ R. P. UDALL,^{89,1} K. UENO,⁹⁶ D. UGOLINI,²¹⁶ C. S. UNNIKRISHNAN,¹⁵³ A. L. URBAN,² S. A. USMAN,¹⁰⁹ A. C. UTINA,^{129,40} H. VAHLBRUCH,^{10,11} G. VAJENTE,¹ A. VAJPEYI,⁶ G. VALDES,² M. VALENTINI,^{155,156} V. VALSAN,²⁷ N. VAN BAKEL,⁴⁰ M. VAN BEUZEKOM,⁴⁰ J. F. J. VAN DEN BRAND,^{129,86,40} C. VAN DEN BROECK,^{104,40} D. C. VANDER-HYDE,⁴⁴ L. VAN DER SCHAAF,⁴⁰ J. V. VAN HEIJNINGEN,^{77,84} J. VANOSKY,¹ M. VARDARO,^{183,40} A. F. VARGAS,⁹⁸ V. VARMA,⁷⁶ M. VASÚTH,⁵⁵ A. VECCHIO,¹⁴ G. VEDOVATO,⁶⁰ J. VEITCH,⁵³ P. J. VEITCH,⁶⁵ K. VENKATESWARA,¹⁸⁵ J. VENNEBERG,^{10,11} G. VENUGOPALAN,¹ D. VERKINDT,³⁹ Y. VERMA,⁶⁸ D. VESKE,³⁷ F. VETRANO,⁷³ A. VICERÉ,^{73,74} A. D. VIETS,¹⁸⁹ V. VILLA-ORTEGA,¹³¹ J.-Y. VINET,⁷⁵ S. VITALE,⁵⁴ T. VO,⁴⁴ H. VOCCA,^{58,57} E. R. G. VON REIS,⁵⁰ J. VON WRANGEL,^{10,11} C. VORVICK,⁵⁰ S. P. VYATCHANIN,⁷⁰ L. E. WADE,¹⁴⁸ M. WADE,¹⁴⁸ K. J. WAGNER,¹⁰⁶ R. C. WALET,⁴⁰ M. WALKER,⁷ G. S. WALLACE,²⁸ L. WALLACE,¹ S. WALSH,²⁷ J. Z. WANG,¹⁵⁸ W. H. WANG,¹²⁵ R. L. WARD,⁹ J. WARNER,⁵⁰ M. WAS,³⁹ N. Y. WASHINGTON,¹ J. WATCHI,¹²⁰ B. WEAVER,⁵⁰ L. WEI,^{10,11} M. WEINERT,^{10,11} A. J. WEINSTEIN,¹ R. WEISS,⁵⁴ C. M. WELLER,¹⁸⁵ F. WELLMANN,^{10,11} L. WEN,⁷⁷ P. WESSELS,^{10,11} J. W. WESTHOUSE,³¹ K. WETTE,⁹ J. T. WHELAN,¹⁰⁶ D. D. WHITE,²³ B. F. WHITING,³⁵ C. WHITTLE,⁵⁴ D. WILKEN,^{10,11} D. WILLIAMS,⁵³ M. J. WILLIAMS,⁵³ A. R. WILLIAMSON,¹³² J. L. WILLIS,¹ B. WILLKE,^{10,11} D. J. WILSON,¹¹⁶ W. WINKLER,^{10,11} C. C. WIPP,¹ T. WŁODARCZYK,⁸⁸ G. WOAN,⁵³ J. WOEHLENER,^{10,11} J. K. WOFFORD,¹⁰⁶ I. C. F. WONG,⁹¹ M. WRIGHT,⁵³ D. S. WU,^{10,11} D. M. WYSOCKI,^{27,106} L. XIAO,¹ H. YAMAMOTO,¹ F. W. YANG,¹⁴⁷ L. YANG,¹⁴¹ Y. YANG,³⁵ Z. YANG,⁴⁶ M. J. YAP,⁹ D. W. YEELES,¹⁷ A. B. YELIKAR,¹⁰⁶ M. YING,¹⁰⁷ A. YOON,⁷ HANG YU,⁷⁶ HAOCUN YU,⁵⁴ A. ZADROŻNY,¹⁷⁷ M. ZANOLIN,³¹ T. ZELENKOVA,³⁴ J.-P. ZENDRI,⁶⁰ M. ZEVIN,¹⁵ J. ZHANG,⁷⁷ L. ZHANG,¹ R. ZHANG,³⁵ T. ZHANG,¹⁴ C. ZHAO,⁷⁷ G. ZHAO,¹²⁰ Y. ZHAO,¹⁴⁷ Z. ZHOU,¹⁵ X. J. ZHU,⁶ A. B. ZIMMERMAN,¹⁸¹ M. E. ZUCKER,^{1,54} AND J. ZWEIZIG¹

THE LIGO SCIENTIFIC COLLABORATION AND THE VIRGO COLLABORATION

¹LIGO Laboratory, California Institute of Technology, Pasadena, CA 91125, USA

²Louisiana State University, Baton Rouge, LA 70803, USA

³Inter-University Centre for Astronomy and Astrophysics, Pune 411007, India

⁴Dipartimento di Farmacia, Università di Salerno, I-84084 Fisciano, Salerno, Italy

⁵INFN, Sezione di Napoli, Complesso Universitario di Monte S. Angelo, I-80126 Napoli, Italy

⁶OzGrav, School of Physics & Astronomy, Monash University, Clayton 3800, Victoria, Australia

⁷Christopher Newport University, Newport News, VA 23606, USA

⁸LIGO Livingston Observatory, Livingston, LA 70754, USA

⁹OzGrav, Australian National University, Canberra, Australian Capital Territory 0200, Australia

¹⁰Max Planck Institute for Gravitational Physics (Albert Einstein Institute), D-30167 Hannover, Germany

¹¹Leibniz Universität Hannover, D-30167 Hannover, Germany

¹²University of Cambridge, Cambridge CB2 1TN, United Kingdom

¹³Theoretisch-Physikalisches Institut, Friedrich-Schiller-Universität Jena, D-07743 Jena, Germany

¹⁴University of Birmingham, Birmingham B15 2TT, United Kingdom

¹⁵Center for Interdisciplinary Exploration & Research in Astrophysics (CIERA), Northwestern University, Evanston, IL 60208, USA

¹⁶Instituto Nacional de Pesquisas Espaciais, 12227-010 São José dos Campos, São Paulo, Brazil

¹⁷Gravity Exploration Institute, Cardiff University, Cardiff CF24 3AA, United Kingdom

¹⁸Gran Sasso Science Institute (GSSI), I-67100 L'Aquila, Italy

¹⁹INFN, Laboratori Nazionali del Gran Sasso, I-67100 Assergi, Italy

²⁰INFN, Sezione di Pisa, I-56127 Pisa, Italy

²¹Università di Pisa, I-56127 Pisa, Italy

²²International Centre for Theoretical Sciences, Tata Institute of Fundamental Research, Bengaluru 560089, India

²³California State University Fullerton, Fullerton, CA 92831, USA

²⁴NCSA, University of Illinois at Urbana-Champaign, Urbana, IL 61801, USA

²⁵Università di Napoli "Federico II", Complesso Universitario di Monte S. Angelo, I-80126 Napoli, Italy

²⁶Université de Lyon, Université Claude Bernard Lyon 1, CNRS, Institut Lumière Matière, F-69622 Villeurbanne, France

²⁷University of Wisconsin-Milwaukee, Milwaukee, WI 53201, USA

²⁸SUPA, University of Strathclyde, Glasgow G1 1XQ, United Kingdom

²⁹Dipartimento di Matematica e Informatica, Università di Udine, I-33100 Udine, Italy

³⁰INFN, Sezione di Trieste, I-34127 Trieste, Italy

³¹Embry-Riddle Aeronautical University, Prescott, AZ 86301, USA

³²Université de Paris, CNRS, Astroparticule et Cosmologie, F-75006 Paris, France

³³Université Paris-Saclay, CNRS/IN2P3, IJCLab, 91405 Orsay, France

- ³⁴European Gravitational Observatory (EGO), I-56021 Cascina, Pisa, Italy
- ³⁵University of Florida, Gainesville, FL 32611, USA
- ³⁶Chennai Mathematical Institute, Chennai 603103, India
- ³⁷Columbia University, New York, NY 10027, USA
- ³⁸INFN, Sezione di Roma, I-00185 Roma, Italy
- ³⁹Laboratoire d'Annecy de Physique des Particules (LAPP), Univ. Grenoble Alpes, Université Savoie Mont Blanc, CNRS/IN2P3, F-74941 Annecy, France
- ⁴⁰Nikhef, Science Park 105, 1098 XG Amsterdam, Netherlands
- ⁴¹Korea Institute of Science and Technology Information, Daejeon 34141, South Korea
- ⁴²INFN Sezione di Torino, I-10125 Torino, Italy
- ⁴³University of Oregon, Eugene, OR 97403, USA
- ⁴⁴Syracuse University, Syracuse, NY 13244, USA
- ⁴⁵Université de Liège, B-4000 Liège, Belgium
- ⁴⁶University of Minnesota, Minneapolis, MN 55455, USA
- ⁴⁷Università degli Studi di Milano-Bicocca, I-20126 Milano, Italy
- ⁴⁸INFN, Sezione di Milano-Bicocca, I-20126 Milano, Italy
- ⁴⁹INAF, Osservatorio Astronomico di Brera sede di Merate, I-23807 Merate, Lecco, Italy
- ⁵⁰LIGO Hanford Observatory, Richland, WA 99352, USA
- ⁵¹Institut de Ciències del Cosmos, Universitat de Barcelona, C/ Martí i Franquès 1, Barcelona, 08028, Spain
- ⁵²Dipartimento di Medicina, Chirurgia e Odontoiatria "Scuola Medica Salernitana", Università di Salerno, I-84081 Baronissi, Salerno, Italy
- ⁵³SUPA, University of Glasgow, Glasgow G12 8QQ, United Kingdom
- ⁵⁴LIGO Laboratory, Massachusetts Institute of Technology, Cambridge, MA 02139, USA
- ⁵⁵Wigner RCP, RMKI, H-1121 Budapest, Konkoly Thege Miklós út 29-33, Hungary
- ⁵⁶Stanford University, Stanford, CA 94305, USA
- ⁵⁷INFN, Sezione di Perugia, I-06123 Perugia, Italy
- ⁵⁸Università di Perugia, I-06123 Perugia, Italy
- ⁵⁹Università di Padova, Dipartimento di Fisica e Astronomia, I-35131 Padova, Italy
- ⁶⁰INFN, Sezione di Padova, I-35131 Padova, Italy
- ⁶¹Montana State University, Bozeman, MT 59717, USA
- ⁶²Institute for Plasma Research, Bhat, Gandhinagar 382428, India
- ⁶³Nicolaus Copernicus Astronomical Center, Polish Academy of Sciences, 00-716, Warsaw, Poland
- ⁶⁴Dipartimento di Ingegneria, Università del Sannio, I-82100 Benevento, Italy
- ⁶⁵OzGrav, University of Adelaide, Adelaide, South Australia 5005, Australia
- ⁶⁶California State University, Los Angeles, 5151 State University Dr, Los Angeles, CA 90032, USA
- ⁶⁷INFN, Sezione di Genova, I-16146 Genova, Italy
- ⁶⁸RRCAT, Indore, Madhya Pradesh 452013, India
- ⁶⁹Missouri University of Science and Technology, Rolla, MO 65409, USA
- ⁷⁰Faculty of Physics, Lomonosov Moscow State University, Moscow 119991, Russia
- ⁷¹SUPA, University of the West of Scotland, Paisley PA1 2BE, United Kingdom
- ⁷²Bar-Ilan University, Ramat Gan, 5290002, Israel
- ⁷³Università degli Studi di Urbino "Carlo Bo", I-61029 Urbino, Italy
- ⁷⁴INFN, Sezione di Firenze, I-50019 Sesto Fiorentino, Firenze, Italy
- ⁷⁵Artemis, Université Côte d'Azur, Observatoire de la Côte d'Azur, CNRS, F-06304 Nice, France
- ⁷⁶CaRT, California Institute of Technology, Pasadena, CA 91125, USA
- ⁷⁷OzGrav, University of Western Australia, Crawley, Western Australia 6009, Australia
- ⁷⁸Dipartimento di Fisica "E.R. Caianiello", Università di Salerno, I-84084 Fisciano, Salerno, Italy
- ⁷⁹INFN, Sezione di Napoli, Gruppo Collegato di Salerno, Complesso Universitario di Monte S. Angelo, I-80126 Napoli, Italy
- ⁸⁰Università di Roma "La Sapienza", I-00185 Roma, Italy
- ⁸¹Univ Rennes, CNRS, Institut FOTON - UMR6082, F-35000 Rennes, France
- ⁸²Indian Institute of Technology Bombay, Powai, Mumbai 400 076, India
- ⁸³Laboratoire Kastler Brossel, Sorbonne Université, CNRS, ENS-Université PSL, Collège de France, F-75005 Paris, France
- ⁸⁴Université catholique de Louvain, B-1348 Louvain-la-Neuve, Belgium
- ⁸⁵Astronomical Observatory Warsaw University, 00-478 Warsaw, Poland
- ⁸⁶VU University Amsterdam, 1081 HV Amsterdam, Netherlands
- ⁸⁷University of Maryland, College Park, MD 20742, USA
- ⁸⁸Max Planck Institute for Gravitational Physics (Albert Einstein Institute), D-14476 Potsdam, Germany
- ⁸⁹School of Physics, Georgia Institute of Technology, Atlanta, GA 30332, USA
- ⁹⁰Villanova University, 800 Lancaster Ave, Villanova, PA 19085, USA

- ⁹¹*The Chinese University of Hong Kong, Shatin, NT, Hong Kong*
- ⁹²*Stony Brook University, Stony Brook, NY 11794, USA*
- ⁹³*Center for Computational Astrophysics, Flatiron Institute, New York, NY 10010, USA*
- ⁹⁴*NASA Goddard Space Flight Center, Greenbelt, MD 20771, USA*
- ⁹⁵*Dipartimento di Fisica, Università degli Studi di Genova, I-16146 Genova, Italy*
- ⁹⁶*RESCEU, University of Tokyo, Tokyo, 113-0033, Japan.*
- ⁹⁷*Tsinghua University, Beijing 100084, China*
- ⁹⁸*OzGrav, University of Melbourne, Parkville, Victoria 3010, Australia*
- ⁹⁹*Università degli Studi di Sassari, I-07100 Sassari, Italy*
- ¹⁰⁰*INFN, Laboratori Nazionali del Sud, I-95125 Catania, Italy*
- ¹⁰¹*Università di Roma Tor Vergata, I-00133 Roma, Italy*
- ¹⁰²*INFN, Sezione di Roma Tor Vergata, I-00133 Roma, Italy*
- ¹⁰³*University of Sannio at Benevento, I-82100 Benevento, Italy and INFN, Sezione di Napoli, I-80100 Napoli, Italy*
- ¹⁰⁴*Institute for Gravitational and Subatomic Physics (GRASP), Utrecht University, Princetonplein 1, 3584 CC Utrecht, Netherlands*
- ¹⁰⁵*Departamento de Astronomía y Astrofísica, Universitat de València, E-46100 Burjassot, València, Spain*
- ¹⁰⁶*Rochester Institute of Technology, Rochester, NY 14623, USA*
- ¹⁰⁷*National Tsing Hua University, Hsinchu City, 30013 Taiwan, Republic of China*
- ¹⁰⁸*OzGrav, Charles Sturt University, Wagga Wagga, New South Wales 2678, Australia*
- ¹⁰⁹*University of Chicago, Chicago, IL 60637, USA*
- ¹¹⁰*Dipartimento di Ingegneria Industriale (DIIN), Università di Salerno, I-84084 Fisciano, Salerno, Italy*
- ¹¹¹*Institut de Physique des 2 Infinis de Lyon (IP2I), CNRS/IN2P3, Université de Lyon, Université Claude Bernard Lyon 1, F-69622 Villeurbanne, France*
- ¹¹²*Seoul National University, Seoul 08826, South Korea*
- ¹¹³*Pusan National University, Busan 46241, South Korea*
- ¹¹⁴*King's College London, University of London, London WC2R 2LS, United Kingdom*
- ¹¹⁵*INAF, Osservatorio Astronomico di Padova, I-35122 Padova, Italy*
- ¹¹⁶*University of Arizona, Tucson, AZ 85721, USA*
- ¹¹⁷*Rutherford Appleton Laboratory, Didcot OX11 0DE, United Kingdom*
- ¹¹⁸*Université libre de Bruxelles, Avenue Franklin Roosevelt 50 - 1050 Bruxelles, Belgium*
- ¹¹⁹*Universitat de les Illes Balears, IAC3—IEEC, E-07122 Palma de Mallorca, Spain*
- ¹²⁰*Université Libre de Bruxelles, Brussels 1050, Belgium*
- ¹²¹*Departamento de Matemáticas, Universitat de València, E-46100 Burjassot, València, Spain*
- ¹²²*Texas Tech University, Lubbock, TX 79409, USA*
- ¹²³*The Pennsylvania State University, University Park, PA 16802, USA*
- ¹²⁴*University of Rhode Island, Kingston, RI 02881, USA*
- ¹²⁵*The University of Texas Rio Grande Valley, Brownsville, TX 78520, USA*
- ¹²⁶*Bellevue College, Bellevue, WA 98007, USA*
- ¹²⁷*Scuola Normale Superiore, Piazza dei Cavalieri, 7 - 56126 Pisa, Italy*
- ¹²⁸*MTA-ELTE Astrophysics Research Group, Institute of Physics, Eötvös University, Budapest 1117, Hungary*
- ¹²⁹*Maastricht University, 6200 MD, Maastricht, Netherlands*
- ¹³⁰*Universität Hamburg, D-22761 Hamburg, Germany*
- ¹³¹*IGFAE, Campus Sur, Universidade de Santiago de Compostela, 15782 Spain*
- ¹³²*University of Portsmouth, Portsmouth, PO1 3FX, United Kingdom*
- ¹³³*The University of Sheffield, Sheffield S10 2TN, United Kingdom*
- ¹³⁴*Laboratoire des Matériaux Avancés (LMA), Institut de Physique des 2 Infinis (IP2I) de Lyon, CNRS/IN2P3, Université de Lyon, Université Claude Bernard Lyon 1, F-69622 Villeurbanne, France*
- ¹³⁵*Dipartimento di Scienze Matematiche, Fisiche e Informatiche, Università di Parma, I-43124 Parma, Italy*
- ¹³⁶*INFN, Sezione di Milano Bicocca, Gruppo Collegato di Parma, I-43124 Parma, Italy*
- ¹³⁷*Physik-Institut, University of Zurich, Winterthurerstrasse 190, 8057 Zurich, Switzerland*
- ¹³⁸*Université de Strasbourg, CNRS, IPHC UMR 7178, F-67000 Strasbourg, France*
- ¹³⁹*West Virginia University, Morgantown, WV 26506, USA*
- ¹⁴⁰*Montclair State University, Montclair, NJ 07043, USA*
- ¹⁴¹*Colorado State University, Fort Collins, CO 80523, USA*
- ¹⁴²*Institute for Nuclear Research, Hungarian Academy of Sciences, Bem t'er 18/c, H-4026 Debrecen, Hungary*
- ¹⁴³*CNR-SPIN, c/o Università di Salerno, I-84084 Fisciano, Salerno, Italy*
- ¹⁴⁴*Scuola di Ingegneria, Università della Basilicata, I-85100 Potenza, Italy*
- ¹⁴⁵*Gravitational Wave Science Project, National Astronomical Observatory of Japan (NAOJ), Mitaka City, Tokyo 181-8588, Japan*
- ¹⁴⁶*Osservatori Astronomic, Universitat de València, E-46980 Paterna, València, Spain*

- ¹⁴⁷*The University of Utah, Salt Lake City, UT 84112, USA*
- ¹⁴⁸*Kenyon College, Gambier, OH 43022, USA*
- ¹⁴⁹*Vrije Universiteit Amsterdam, 1081 HV, Amsterdam, Netherlands*
- ¹⁵⁰*University of Szeged, Dóm tér 9, Szeged 6720, Hungary*
- ¹⁵¹*Universiteit Gent, B-9000 Gent, Belgium*
- ¹⁵²*University of British Columbia, Vancouver, BC V6T 1Z4, Canada*
- ¹⁵³*Tata Institute of Fundamental Research, Mumbai 400005, India*
- ¹⁵⁴*INAF, Osservatorio Astronomico di Capodimonte, I-80131 Napoli, Italy*
- ¹⁵⁵*Università di Trento, Dipartimento di Fisica, I-38123 Povo, Trento, Italy*
- ¹⁵⁶*INFN, Trento Institute for Fundamental Physics and Applications, I-38123 Povo, Trento, Italy*
- ¹⁵⁷*The University of Mississippi, University, MS 38677, USA*
- ¹⁵⁸*University of Michigan, Ann Arbor, MI 48109, USA*
- ¹⁵⁹*Dipartimento di Fisica, Università di Trieste, I-34127 Trieste, Italy*
- ¹⁶⁰*American University, Washington, D.C. 20016, USA*
- ¹⁶¹*Carleton College, Northfield, MN 55057, USA*
- ¹⁶²*University of California, Berkeley, CA 94720, USA*
- ¹⁶³*Directorate of Construction, Services & Estate Management, Mumbai 400094 India*
- ¹⁶⁴*Universiteit Antwerpen, Prinsstraat 13, 2000 Antwerpen, Belgium*
- ¹⁶⁵*University of Białystok, 15-424 Białystok, Poland*
- ¹⁶⁶*University of Southampton, Southampton SO17 1BJ, United Kingdom*
- ¹⁶⁷*Institut de Física d'Altes Energies (IFAE), Barcelona Institute of Science and Technology, and ICREA, E-08193 Barcelona, Spain*
- ¹⁶⁸*University of Washington Bothell, Bothell, WA 98011, USA*
- ¹⁶⁹*Institute of Applied Physics, Nizhny Novgorod, 603950, Russia*
- ¹⁷⁰*Ewha Womans University, Seoul 03760, South Korea*
- ¹⁷¹*Inje University Gimhae, South Gyeongsang 50834, South Korea*
- ¹⁷²*Korea Astronomy and Space Science Institute, Daejeon 34055, South Korea*
- ¹⁷³*National Institute for Mathematical Sciences, Daejeon 34047, South Korea*
- ¹⁷⁴*Ulsan National Institute of Science and Technology, Ulsan 44919, South Korea*
- ¹⁷⁵*Bard College, 30 Campus Rd, Annandale-On-Hudson, NY 12504, USA*
- ¹⁷⁶*Institute of Mathematics, Polish Academy of Sciences, 00656 Warsaw, Poland*
- ¹⁷⁷*National Center for Nuclear Research, 05-400 Świerk-Otwock, Poland*
- ¹⁷⁸*Cornell University, Ithaca, NY 14850, USA*
- ¹⁷⁹*Université de Montréal/Polytechnique, Montreal, Quebec H3T 1J4, Canada*
- ¹⁸⁰*Laboratoire Lagrange, Université Côte d'Azur, Observatoire Côte d'Azur, CNRS, F-06304 Nice, France*
- ¹⁸¹*Department of Physics, University of Texas, Austin, TX 78712, USA*
- ¹⁸²*NAVIER, École des Ponts, Univ Gustave Eiffel, CNRS, Marne-la-Vallée, France*
- ¹⁸³*Institute for High-Energy Physics, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, Netherlands*
- ¹⁸⁴*NASA Marshall Space Flight Center, Huntsville, AL 35811, USA*
- ¹⁸⁵*University of Washington, Seattle, WA 98195, USA*
- ¹⁸⁶*Dipartimento di Matematica e Fisica, Università degli Studi Roma Tre, I-00146 Roma, Italy*
- ¹⁸⁷*INFN, Sezione di Roma Tre, I-00146 Roma, Italy*
- ¹⁸⁸*ESPCI, CNRS, F-75005 Paris, France*
- ¹⁸⁹*Concordia University Wisconsin, Mequon, WI 53097, USA*
- ¹⁹⁰*Università di Camerino, Dipartimento di Fisica, I-62032 Camerino, Italy*
- ¹⁹¹*Southern University and A&M College, Baton Rouge, LA 70813, USA*
- ¹⁹²*Centre Scientifique de Monaco, 8 quai Antoine 1er, MC-98000, Monaco*
- ¹⁹³*Indian Institute of Technology Madras, Chennai 600036, India*
- ¹⁹⁴*Saha Institute of Nuclear Physics, Bidhannagar, West Bengal 700064, India*
- ¹⁹⁵*Institut des Hautes Etudes Scientifiques, F-91440 Bures-sur-Yvette, France*
- ¹⁹⁶*Indian Institute of Science Education and Research, Kolkata, Mohanpur, West Bengal 741252, India*
- ¹⁹⁷*Department of Astrophysics/IMAPP, Radboud University Nijmegen, P.O. Box 9010, 6500 GL Nijmegen, Netherlands*
- ¹⁹⁸*GRAPPA, Anton Pannekoek Institute for Astronomy and Institute for High-Energy Physics, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, Netherlands*
- ¹⁹⁹*Consiglio Nazionale delle Ricerche - Istituto dei Sistemi Complessi, Piazzale Aldo Moro 5, I-00185 Roma, Italy*
- ²⁰⁰*Hobart and William Smith Colleges, Geneva, NY 14456, USA*
- ²⁰¹*International Institute of Physics, Universidade Federal do Rio Grande do Norte, Natal RN 59078-970, Brazil*
- ²⁰²*Museo Storico della Fisica e Centro Studi e Ricerche "Enrico Fermi", I-00184 Roma, Italy*

- ²⁰³*Lancaster University, Lancaster LA1 4YW, United Kingdom*
- ²⁰⁴*OzGrav, Swinburne University of Technology, Hawthorn VIC 3122, Australia*
- ²⁰⁵*Università di Trento, Dipartimento di Matematica, I-38123 Povo, Trento, Italy*
- ²⁰⁶*Indian Institute of Science Education and Research, Pune, Maharashtra 411008, India*
- ²⁰⁷*Dipartimento di Fisica, Università degli Studi di Torino, I-10125 Torino, Italy*
- ²⁰⁸*Indian Institute of Technology, Palaj, Gandhinagar, Gujarat 382355, India*
- ²⁰⁹*Centro de Astrofísica e Gravitação (CENTRA), Departamento de Física, Instituto Superior Técnico, Universidade de Lisboa, 1049-001 Lisboa, Portugal*
- ²¹⁰*Marquette University, 11420 W. Clybourn St., Milwaukee, WI 53233, USA*
- ²¹¹*Indian Institute of Technology Hyderabad, Sangareddy, Khandi, Telangana 502285, India*
- ²¹²*INAF, Osservatorio di Astrofisica e Scienza dello Spazio, I-40129 Bologna, Italy*
- ²¹³*Andrews University, Berrien Springs, MI 49104, USA*
- ²¹⁴*Dipartimento di Scienze Aziendali - Management and Innovation Systems (DISA-MIS), Università di Salerno, I-84084 Fisciano, Salerno, Italy*
- ²¹⁵*Van Swinderen Institute for Particle Physics and Gravity, University of Groningen, Nijenborgh 4, 9747 AG Groningen, Netherlands*
- ²¹⁶*Trinity University, San Antonio, TX 78212, USA*