

## GOPEN ACCESS

**Citation:** Janzen A, Richter KK, Mwebi O, Brown S, Onduso V, Gatwiri F, et al. (2021) Distinguishing African bovids using Zooarchaeology by Mass Spectrometry (ZooMS): New peptide markers and insights into Iron Age economies in Zambia. PLoS ONE 16(5): e0251061. https://doi.org/10.1371/ journal.pone.0251061

Editor: Justin W. Adams, Monash University, AUSTRALIA

Received: August 12, 2020

Accepted: April 20, 2021

Published: May 18, 2021

**Copyright:** © 2021 Janzen et al. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: We present archaeological data in the Supporting information files. Additionally, the raw data have been uploaded to the following locations for open access (this information and links are also in our Supporting information): MS/MS Data: ProteomeXchange through MassIVE Record Number: PXD020810 doi:10.25345/C5TJ3M password: reviewer Reference ZooMS Spectra: Zenodo doi:10.5281/

zenodo.3964709 Archaeological Zambia ZooMS Spectra: Zenodo doi:10.5281/zenodo.3971142. RESEARCH ARTICLE

# Distinguishing African bovids using Zooarchaeology by Mass Spectrometry (ZooMS): New peptide markers and insights into Iron Age economies in Zambia

Anneke Janzen <sup>1,2\*</sup>, Kristine Korzow Richter<sup>1,3</sup>, Ogeto Mwebi<sup>4</sup>, Samantha Brown <sup>1</sup>, Veronicah Onduso<sup>4</sup>, Filia Gatwiri<sup>5</sup>, Emmanuel Ndiema<sup>5</sup>, Maggie Katongo<sup>6</sup>, Steven T. Goldstein<sup>1</sup>, Katerina Douka<sup>1</sup>, Nicole Boivin<sup>1,7,8,9</sup>

Department of Archaeology, Max-Planck Institute for the Science of Human History, Jena, Germany,
Department of Anthropology, University of Tennessee, Knoxville, Tennessee, United States of America,
Department of Anthropology, Harvard University, Boston, Massachusetts, United States of America,
Department of Zoology, Osteology Section, National Museums of Kenya, Nairobi, Kenya, 5 Department of
Earth Sciences, Archaeology Section, National Museums of Kenya, Nairobi, Kenya, 6 Department of
Archaeology, Livingstone Museum, Livingstone, Zambia, 7 School of Social Science, The University of
Queensland, Brisbane, Australia, 8 Department of Anthropology and Archaeology, University of Calgary,
Calgary, Canada, 9 Department of Anthropology, National Museum of Natural History, Smithsonian
Institution, Washington, D.C., United States of America

\* ajanzen@utk.edu

## Abstract

Assessing past foodways, subsistence strategies, and environments depends on the accurate identification of animals in the archaeological record. The high rates of fragmentation and often poor preservation of animal bones at many archaeological sites across sub-Saharan Africa have rendered archaeofaunal specimens unidentifiable beyond broad categories, such as "large mammal" or "medium bovid". Identification of archaeofaunal specimens through Zooarchaeology by Mass Spectrometry (ZooMS), or peptide mass fingerprinting of bone collagen, offers an avenue for identification of morphologically ambiguous or unidentifiable bone fragments from such assemblages. However, application of ZooMS analysis has been hindered by a lack of complete reference peptide markers for African taxa, particularly bovids. Here we present the complete set of confirmed ZooMS peptide markers for members of all African bovid tribes. We also identify two novel peptide markers that can be used to further distinguish between bovid groups. We demonstrate that nearly all African bovid subfamilies are distinguishable using ZooMS methods, and some differences exist between tribes or sub-tribes, as is the case for Bovina (cattle) vs. Bubalina (African buffalo) within the subfamily Bovinae. We use ZooMS analysis to identify specimens from extremely fragmented faunal assemblages from six Late Holocene archaeological sites in Zambia. ZooMS-based identifications reveal greater taxonomic richness than analyses based solely on morphology, and these new identifications illuminate Iron Age subsistence economies c. 2200–500 cal BP. While the Iron Age in Zambia is associated with the transition from hunting and foraging to the development of farming and herding, our results demonstrate the continued reliance on wild bovids among Iron Age communities in central and southwestern

**Funding:** This work was funded by the Max Planck Society, with additional contribution from the European Research Council (FINDER Project, grant agreement No 715069).

**Competing interests:** The authors have declared that no competing interests exist. The funder had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Zambia Iron Age and herding focused primarily on cattle. We also outline further potential applications of ZooMS in African archaeology.

## Introduction

Secure identification of animal taxa in the archaeological record is critical for assessing past subsistence strategies, environments, foodways, and tracking species dispersals and translocations. A major challenge in exploring these questions, and in analysis of African archaeofaunal assemblages in general, is the sheer number of bovid species in most regions. Globally, the Bovidae family composes over 50% of ungulates with over 140 extant species. Bovids exhibit significant diversity in body size, morphology, geographical range, diet, and behavior (Fig 1), and inhabit a range of environments across Africa, which vary greatly in terms of topography,



Fig 1. Distribution maps of African species belonging to each of the wild bovid subfamilies, tribes, or sub-tribes analyzed in this study. Distribution data from IUCN [16]. Basemap from [17].

https://doi.org/10.1371/journal.pone.0251061.g001







climate, and vegetation (Fig 2). Sub-Saharan Africa is home to the greatest diversity of bovids on Earth, including species from eight of the nine subfamilies of Bovidae (Figs 1 and 3). Three domestic bovid species are also widespread across the continent: cattle (*Bos taurus* and *Bos indicus*), sheep (*Ovis aries*), and goat (*Capra hircus*). Given the considerable richness of bovid





species across much of the African continent, these morphologically similar taxa can be difficult to distinguish when fragmentary. In the absence of robust, diagnostic elements or teeth, highly fragmented assemblages are often characterized by mostly coarse identifications to the family level and body size class (e.g. Bovid 2) [1]. A range of processes contribute to fragmentation as well as deletion of identifiable bone ends, including culinary processing [2, 3], carnivore damage [4–6], and density-mediated attrition, in which less-dense elements or element portions fail to survive taphonomic processes [7, 8]. Manufacture of bone tools or objects also typically involves removal of epiphyses and obliterates identifiable features [9] Where reported in faunal analyses, the less- or non-identifiable component comprises the bulk of faunal assemblages, and bovids comprise the majority of minimally-identifiable mammal remains [10–15]. Given this large proportion of remains that are not attributable to taxon in so many archaeological sites, zooarchaeologists have been limited in their interpretations of African subsistence strategies, foodways, and the economic and symbolic importance of animals in the past.

These taphonomic issues make it particularly difficult to evaluate taxonomic frequencies of bovid species in sites with high bone fragmentation and poor preservation, hindering assessments of hunting strategies, foodways, past environments, and subsistence economies. In particular, tracking the expansion and nature of food production in Africa is challenging because initial evidence for livestock in a region is often patchy [22–25], and identifications may be hotly debated [26–29]. For example, early evidence for domesticates in Kenya south of the Lake Turkana Basin before 3300 BP are sparse, and limited to a few identifiable elements at a handful of sites, such as in Kansyore levels (c. 4400 to 4000 cal. BP) at Wadh Lang'o [25] and at c. 4100 BP at Enkapune Ya Muto [23]. In southern Africa, early zooarchaeological evidence for livestock is also limited to a few morphologically identifiable specimens. Sites including Toteng 1 [30] and Spoegrivier [31–33] have early evidence for sheep at c. 2070 and 2100 BP, respectively. Efforts in identifying early domesticates through molecular methods such as ancient DNA (aDNA) have been met with controversy [26–29]. Many early herders and farmers relied

on domesticates and terrestrial wild species [13, 34–36] and reliance on wild bovids, particularly in forested areas, has continued into recent centuries in many parts of the continent [37]. Despite their centrality in archaeological discussions of African economies and diets, the considerable richness of native bovid species and their gross morphological similarity, as well as their osteological resemblance to domesticates, complicates assessments of their exploitation in the past.

Zooarchaeology by Mass Spectrometry (ZooMS), or peptide mass fingerprinting of bone collagen to taxonomically identify morphological ambiguous mammal, fish, and reptile bone, has been successfully applied in many parts of the world [38-42]. Unlike ancient aDNA analysis, ZooMS derives identifications from the study of bone collagen, which is well-preserved in many archaeological contexts. Given the often poor preservation of aDNA, particularly outside of temperate regions and when petrous bone is unavailable, ZooMS-based studies accordingly have higher success rates than aDNA studies [43]. In East African contexts, ZooMS has proven to be effective in examining translocations of Asian species and identifying domesticates, but in limited contexts [44, 45]. Coutu and colleagues [46] also used ZooMS to investigate the early ivory trade in southern Africa in the 7<sup>th</sup> century CE. Despite the clear value of ZooMS, however, very few studies have applied the method to study indigenous African taxa, and those that have done so have been focused on worked bone artifacts in limited geographical areas, and have drawn upon a small reference library of African bovid species [47-49]. Two recent studies have explored early evidence for caprines. Le Meillour and colleagues used paleoproteomics on potential caprines to clarify the absence of domestic livestock at Leopard Cave in central Namibia until c. 960 BP [49]. Most recently, Coutu et al [33] identified a new ZooMS marker that separates sheep from wild bovids to explore the introduction of domestic sheep into southern Africa. However, the lack of a comprehensive peptide marker reference library for wild African bovids has greatly limited the applicability of ZooMS in African archaeology. The present study remedies this situation by providing these reference markers.

Here, we present the first extensive study identifying ZooMS peptide markers for all indigenous African bovid tribes, and character two new peptide markers for bovids. This database allows, for the first time, large-scale systematic identifications of otherwise unidentifiable fragments in African archaeofaunal assemblages, as well as rapid confirmation of identifications of domesticate and wild species that lack confidence due to morphological similarity. We apply the newly generated reference African bovid peptide sequences produced in this study to the investigation of extremely fragmented archaeofaunal assemblages from several recently excavated Iron Age (c. 2200-500 cal BP) archaeological sites in central and southwestern Zambia. Today, Zambia is home to members of most of the bovid groups described in this study (Fig 1), and may have included more taxa in the past. Our results reveal that subsistence economies were grounded in cattle-based pastoralism, and included hunting of wild bovids, particularly cephalophines (duikers). We also demonstrate the validity and reliability of ZooMS methods in African contexts and provide a baseline for collagen recovery rates for typical highly fragmented African assemblages. The newly developed African bovid ZooMS reference database offers a novel avenue for addressing key challenges to traditional zooarchaeological analyses, and thus more effectively exploring African hunting and subsistence strategies, foodways, and past environments.

#### Faunal identifications and key research questions

While osteologically similar, bovids are incredibly diverse in their diets and the environments they inhabit. For example, extant members of the subfamily Alcelaphinae—wildebeest (*Conno-chaetes taurinus and Connochaetes gnou*), hartebeest (*Alcelaphus buselaphus*), tsessebe/topi

(*Damaliscus lunatus*), and blesbok/bontebok (*Damaliscus pygargus*)—are dedicated grazers that inhabit open grasslands, shrublands, and floodplains [50]. The sole member of the sub-family Aepycerotinae, the impala (*Aepyceros melampus*), is an obligate drinker and mixed feeder, preferring woodland and savannah ecotones [51]. In contrast, members of the subfamily Cephalophinae (e.g. *Sylvicapra grimmia, Cephalophus sp.*)—the duikers—are primarily browsers. These smaller-bodied bovids prefer more wooded habitats, though substantial variation in habitat preference exists among species within this group [52]. While variation in diet and habitat preference exists among species within all bovid groups, these broad differences in diet among bovid tribes are apparent in carbon stable isotope data, reflecting differences in the photosynthetic pathways of grasses versus trees and shrubs at low elevations in the tropics [53, 54].

Given their habitat and dietary differences [55, 56], taxonomic identifications of bovids beyond the family level are crucial for inferring past environments [57–63]. Species-level identifications are most effective, allowing for detailed morphological studies that can further clarify environmental context [64, 65], but broad identifications to tribe or subfamily can still provide useful information, particularly in poorly preserved or small assemblages in which few specimens are morphologically identifiable to genus or species. Coarse identifications of bovids to tribe also allow for improved interpretation of stable isotope data. For example, carbon isotope analysis of mixed feeder species (e.g. impala or goat) can clarify the degree of brushy vegetation or grass in an environment. Similarly, identifications of broad taxonomic groups are useful for interpreting  $\delta^{18}$ O values of evaporation-sensitive and -insensitive animals [66–69].

Assessment of prey choice and subsistence economies is not possible without secure taxonomic identifications. Bovid size classes have typically been used to assess broad hunting, transport, and carcass processing strategies [12, 15], but as bovids inhabit varying environments and exhibit behavioral differences, narrower identifications reveal more detail about forager decisions and tactics [23, 58, 62, 70–73] (Fig 4). Finally, bovid and mammal size classes used by zooarchaeologists across Africa vary [1, 58, 74], and therefore narrower identifications can aid in inter-regional comparisons of zooarchaeological data.

A major focus in Holocene African archaeology involves the study of the development of pastoral lifeways. Pastoralism in Africa is unique in that it developed much earlier than farming, first appearing in northern Africa around 8,000 years ago, and gradually spreading throughout the continent over the next several millennia. Tracking the spread of pastoralism and subsequent mixed agro-pastoralists across Africa, as well as exploring the diversity of pastoral management strategies across the continent, hinges on secure identifications of domesticates. Cattle, sheep, and goat expanded beyond the range of their wild progenitors: while there is some debate whether cattle were domesticated in Africa [75–78], wild African cattle (*Bos primigenius*) were restricted to northern Africa. Sheep and goat were domesticated in southwest Asia [79–82], and are the only members of subfamily Caprinae in sub-Saharan Africa. Still, identifying domesticates in archaeofaunal assemblages comes with some challenges. Cattle are osteologically similar to the widespread African buffalo (*Syncerus caffer*) [83], and while many osteological guides exist for identifying caprines [84–87], fragmentary post-cranial remains are difficult to distinguish from like-sized wild bovids [49].

Detailed analyses of herd management strategies have been carried out across eastern [3, 11, 88], and southern Africa [89–92], but somewhat patchily—large parts of the continent have not seen such studies (e.g. [93] in western Africa). Domestic caprines are also infrequently identified to species [45], precluding deeper analyses of herd management strategies and explorations of biological connections between communities (eg. [94]). A great deal of work remains to be done in establishing a firm chronology for the spread of domesticates across



**Fig 4. Examples of research questions in African archaeology that can be aided by ZooMS of bovids.** Relevant time periods for each question are indicated. Elevation data from [19].

much of Africa, and major questions still exist regarding the mechanisms behind these dispersals, as well as the nature of pastoral economies over the last few millennia (e.g., [95]). Additionally, the exact trajectories of the expansion of herding throughout Africa are still poorly understood, as are the mechanisms by which herding took hold in different parts of the continent, i.e., whether through demographic migration or local adoption [25, 96]. Furthermore, herding was not adopted wholesale as it spread across the continent: clear archaeological evidence for the long-term persistence of foragers exists [58], and many food-producing populations also continued to rely on terrestrial wild taxa, as seen in Pastoral Neolithic [11, 97], as well as Iron Age sites in eastern and southern Africa [98–101]. In West Africa, recent studies highlight the range of wild bovid species exploited by humans until relatively recently, reflecting wild species' roles in elite foodways, as seen in 17<sup>th</sup> and 18<sup>th</sup> century Dahomean palace sites [37] and revealed through investigation of the trans-Saharan trade in 11<sup>th</sup> to 14<sup>th</sup> century Senegal [102]. Animal diversity in ritual contexts, such as at Hierakopolis in predynastic Egypt [103] and 2<sup>nd</sup> millennium CE rain control sites such as Ratho Kroonkop in South Africa [104], also highlights the need for a comprehensive set of reference markers for African fauna.

## ZooMS in African archaeology

To date, ZooMS analyses of faunal remains in Africa have been limited to a few pioneering studies. The first applications of ZooMS in African archaeological contexts were of ivory artifacts. Coutu et al. [46] used ZooMS to identify the species used in artifacts from Early Iron Age sites (c. 1400 to 1100 BP) in KwaZulu-Natal, South Africa, finding that elephants were the source of all ivory analyzed. Two recent studies have focused on bone artifacts: Desmond and colleagues [48] analyzed bone tools from the Later Stone Age cave site of Taforalt in eastern Morocco, and Bradfield and colleagues [47] used ZooMS to identify bone arrowheads from Iron Age (c. 1700 to 400 BP) sites in southern Africa. Both studies provided partial sets of reference markers for 6 and 13 bovids respectively, which allowed for coarse identifications of species used to manufacture bone artifacts.

Pastoral strategies have been explored by Prendergast et al. [45], who utilized ZooMS-based identifications to support and improve morphological identifications of sheep and goat at the Pastoral Neolithic site of Luxmanda in northern Tanzania. Le Meillour and colleagues [105] employed a paleoproteomic approach to identify bone fragments from the Later Stone Age sites of Toteng 1 and 3 in southern Africa. The authors demonstrate the usefulness of the method to identify suid, equid, and bovid (including domestic sheep) specimens. Prendergast et al. [44] used ZooMS to track species translocations. Circumventing the challenged posed by the osteological similarity between rat-sized native African murid species and introduced Asian black rats, ZooMS enabled significant improvement of species identifications. This study thus chronicled the late appearance of black rat in island and coastal East African sites in the second millennium BP, supporting the hypothesis that Asian species were introduced through Indian Ocean trade, but challenging ideas of early maritime-mediated biological exchange between Asia and Africa.

Most recently Coutu et al. [33] used ZooMS to separate domestic sheep from several similarly sized, wild bovids to confirm identifications of earliest domestic sheep in southern Africa. While only a small subset of bovids were analyzed, the development of a new marker that allows for separation between domestic and some wild bovids is critical in the use of ZooMS in Africa.

These initial studies highlight the potential of ZooMS in Africa, where, as in other parts of the world [42, 106–110], it has proven valuable in addressing a variety of questions. By providing identifications of morphologically ambiguous and fragmentary specimens, ZooMS has the

potential to revive research interest in, and add value to, assemblages previously considered too poorly preserved for analysis. Through its role in clarifying taxonomic identifications, ZooMS opens new opportunities for addressing questions of paleoenvironment, subsistence strategies, foodways, the spread of and development of herding economies, and the symbolic importance of animals in the African past. However, application of ZooMS analysis across the continent has been hindered by a lack of complete reference peptide markers for African taxa, particularly bovids. Here we present the first comprehensive set of peptide markers for wild African bovids, based upon genetic and proteomic data from 34 different species across all subfamilies present in Africa. We also identify two additional markers that allow for further discrimination within subfamilies. Our results show that ZooMS can be used to distinguish among most bovid tribes, and in some cases provide narrower taxonomic identifications, which also allows for unambiguous identification of domestic livestock. We use these reference sets to identify wild and domestic taxa from six archaeological sites with extremely poor faunal preservation. These sites are located in southern and central Zambia, and span the Iron Age period of Zambia (c. 2000-500 years BP), an archaeological phase associated with the spread of farming, longer-term village settlements, and the formation of complex polities in southern Africa.

## Materials & methods

For proteomic analysis, modern reference samples were taken from twenty taxa across all six of the eight subfamilies present in Africa. In addition, to ensure that results were comparable with published markers, samples from modern cattle (Bos taurus) were also obtained to ensure similarity with previously published reference data for Bos (Table 1). Three individuals from each species were sampled from the Osteology Section at the National Museums of Kenya in Nairobi (all specimen information can be found in S1 Table). In the few cases where three individuals were not available for a species, at least three individuals were taken from the genus. 50 mg of bone from each individual was sampled using a handheld Dremel outfitted with a diamond drill bit at the Osteology Laboratory at the National Museums of Kenya, Nairobi. Between samples, the drill bits were sonicated and rinsed with distilled water. Published reference LC-MS/MS data for impala (Aepyceros melampus) and springbok (Antidorcas marsupialis) [33, 49] were used to confirm the presence of the correct marker sequences in these species, although MALDI-TOF-MS analysis was not conducted for these taxa in this study. Genetic data was mined from published Bovidae genomes (Chen et al, 2019, S1 File) for 28 African species. Combining both genomic and proteomic reference material, a total of 34 African species were analyzed. Genetic data from eight additional Bovid species and three species from two outgroups (Equidae and Cervidae) were also used along with previously published ZooMS markers (Table 1).

Faunal specimens from the six archaeological sites included in this study are: Kapiri Mposhi A (n = 69), Fibobe II (n = 1), and Muteteshi (n = 4) in the Irumide Belt of Central Zambia, the Kalomo culture site of Kalundu Mound on the Batoka Plateau of southern Zambia (n = 102), and Salumano (n = 9) and Jakobo West (n = 7) on the margin of the Kalahari Depression of southwestern Zambia (Fig 5). All necessary permits were obtained for the described study for both modern and archaeological specimens, which complied with all relevant regulations. All sampled reference material is housed in the Osteology Section at the National Museums of Kenya in Nairobi (all specimen information can be found in S1 Table). Research and export permits in Kenya were granted by the National Commission for Science, Technology and Innovation (ref. NACOSTI/P/17/66411/14936), and the National Museums of Kenya (ref. NMK/ESC/ACL/j/VOL.1). Research and export permits for the samples were granted by

Family	Subfamily	Tribe	Subtribe	Genus	Species	Common Name	DNA	ZooMS
Bovidae	Aepycerotinae			Aepyceros	melampus	impala	X <sup>a</sup>	* d, g
	Alcelaphinae			Connochaetes	taurinus	blue wildebeest	X <sup>a</sup>	X** <sup>e</sup>
				Damaliscus (Beatragus)	hunteri	hirola		X
				Damaliscus	lunatus	topi, tsessebe	X <sup>a</sup>	X
				Alcelaphus	buselaphus	hartebeest	X <sup>a</sup>	X**e,f
	Antilopinae	Antilopini		Nanger	granti	Grant's gazelle	X <sup>a</sup>	X
				Eudorcas	thomsonii	Thomson's gazelle	X <sup>a</sup>	X
				Gazella	rufifrons	red-fronted gazelle		* * d,f
				Antidorcas	marsupialis	springbok	X <sup>a</sup>	X <sup>g,h</sup>
				Litocranius	walleri	gerenuk	Xa	
		Neotragini		Procapra	przewalskii	Przewalskyi's gazelle	Xa	
				Ourebia	ourebi	oribi	Xa	X** <sup>e</sup>
				Oreotragus	oreotragus	klipspringer	Xa	X** <sup>e</sup>
				Madoqua	kirkii	Kirk's dik-dik	Xa	X
				Neotragus	moschatus	suni	Xa	X
				Raphicerus	campestris	steenbok	Xa	
	Pantholopinae			Pantholops	hodgsonii	Tibetan antelope	Xa	*d
	Bovinae	Bovini	Bovina	Bos	taurus	cattle	X	X*b
				Bos	grunniens	yak	Xa	
				Bison	bison	bison	X	* d
			Bubalina	Bubalus	bubalis	water buffalo	X	*b
				Syncerus	caffer	African buffalo	Xa	X
		Tragelaphini		Taurotragus	oryx	common eland	Xa	X**e
				Taurotragus	derbianus	giant eland		X
				Tragelaphus	eurycerus	bongo	Xa	X
				Tragelaphus	scriptus	bushbuck	Xa	
				Tragelaphus	imberbis	lesser kudu	Xa	
				Tragelaphus	strepsiceros	greater kudu	Xa	**e
				Tragelaphus	buxtoni	mountain nyala	Xa	
				Tragelaphus	spekii	sitatunga	Xa	
				Tragelaphus	angasii	nalya		**e
	Caprinae			Ovis	aries	sheep	X	*b,c,d,h
				Capra	hirca	goat	X	*b,c,d
				Capra	ibex	ibex	Xa	*b,c
				Ammotragus	lervia	Barbary sheep		*f
	Cephalophinae			Cephalophus	harveyi	Harvey's duiker	Xa	
				Philantomba	maxwellii	Maxwell's duiker	Xa	
				Sylvicapra	grimmia	bush duiker	Xa	X <sup>h</sup>
				Cephalophus	adersi	Aders's duiker		x
	Hippotraginae			Hippotragus	niger	sable antelope		X**e
				Hippotragus	equinus	roan antelope		X**e,f
				Oryx	gazella	gemsbok	X <sup>a</sup>	-
	Reduncinae		-	Kobus	kob	kob		X
				Kobus	ellipsiprvmnus	waterbuck	Xa	X
			-	Redunca	redunca	bohor reedbuck	X <sup>a</sup>	

#### Table 1. List of the modern reference species used in this study.

(Continued)

#### Table 1. (Continued)

Family	Subfamily	Tribe	Subtribe	Genus	Species	Common Name	DNA	ZooMS
Cervidae	Capreolinae			Odocoileus	virginianus	white tailed deer	Х	*d
				Rangifer	tarandus	reindeer		* d
Equidae				Equus	caballus	horse	X	*b,d

ZooMS data was generated for three individuals from each of the species indicated, with LC-MS/MS conducted on one individual from the species indicated in bold. Members from Families Cervidae and Equidae with existing markers and sequence data were used to generate comparative reference spectra for validation. An 'X' in DNA column indicates species where genetic sequence data for COL1A1 and COL1A2 was available. An 'X' in the ZooMS column indicates species where data was generated for this publication. Bold indicates species analyzed by LC-MS/MS.

\*Indicates where all 9 common ZooMS markers are published.

\*\*Indicates where some, but not all of the common ZooMS markers are published.

<sup>a</sup>Genetic data from Chen et al (2019). Other genetic data was mined from NCBI.

<sup>b</sup>[38].

°[111].

<sup>d</sup>[112].

<sup>e</sup>[<u>47</u>].

<sup>f</sup>[48].

<sup>g</sup>LC-MS/MS data was analyzed from LeMeillour et al [49].

<sup>h</sup> LC-MS/MS data was analyzed from Coutu et al. [33].

https://doi.org/10.1371/journal.pone.0251061.t001





https://doi.org/10.1371/journal.pone.0251061.g005

NHCC, Lusaka, and fauna from Zambia sites were exported with permission of The Livingstone Museum. Faunal remains from these sites are highly fragmented and often poorly preserved due to culinary processing and unfavorable taphonomic conditions, particularly in the Kalahari sands at Salumano and Jakobo West [113].

Salumano is one of the type sites of the "Situmpa Group" of southwestern Zambia (also including Lusu Rapids, Situmpa, and Machili Forest Station), defined by Clark and Fagan [114] and Katanekwa [115] as reflecting a pre-Iron Age appearance of ceramics and livestock in the region. The early occurrences of Situmpa pottery and cattle at Salumano date to c. 2000 BP, overlaid by Early and Later Iron Age horizons extending into the last few hundred years [101, 116]. Salumano was revisited and re-tested by researchers at the Max Planck Institute for the Science of Human History and the Livingstone Museum in 2017, confirming this sequence. During the course of this research, a nearby locality of Jakobo West was discovered, which had only a single archaeological horizon with stone tools, bone fragments, and pot sherds that match typical "Situmpa Group" styles [115] encountered at about 1.3 m below surface.

Muteteshi and Fibobe II in the Mulungushi River Basin were first excavated by the Zambian National Monuments Commission between 1977 and 1979 under the direction of John Robertson [117]. Muteteshi is a single component Early Iron Age site with radiocarbon dates ranging from roughly 2000 to 1600 years BP, with early evidence of domesticated sorghum (*Sorghum bicolor*) [117]. Fibobe II is a nearby village site with substantial evidence for hut structures, iron working areas, and cultivation of domesticated crops dating to the Later Iron Age (approximately 1175–1410 years BP) [117]. Both sites were revisited in 2018 and the limited fauna analyzed here were recovered. Both sites had very poor faunal preservation.

Kapiri Mposhi A was identified during the course of new fieldwork in 2018. The site is not yet dated, but a transition from Later Stone Age to Iron Age material culture within a 4 meter sequence is apparent. Based on stone tool and ceramics styles, the site may have accumulated from c. 4000 to 600 years ago, likely by stone tool-using foragers who were in contact with Iron Age farmers [118].

Kalundu Mound is a roughly 140 x 106m low-sloped mound consisting of 3 meters of stratified archaeological material on the Batoka Plateau, southern Zambia. Excavations by Inskeep [119] and Fagan [98] identified Kalundu Mound as being diagnostic of the Later Iron Age 'Kalomo Culture' of the region based on ceramic styles, iron and ornamental artifacts, and burial traditions. Kalomo economies appear to have relied on sorghum cultivation and cattlekeeping, as well hunting of wild fauna [98]. However, current models for animal subsistence are based only on identification of teeth alone, and do not necessarily provide an accurate reflection of the breadth of animal species exploited during the Later Iron Age. Early radiocarbon dates on bulk charcoal samples from Kalundu Mound suggested use and formation of the mound occurred from ~1650–960 years ago, overlaying pit features that yielded dates in the mid 3<sup>rd</sup> millennium BP. Re-excavation of a 1x3m trench and subsequent direct dating of macrobotanical and bone remains recovered in 2017 indicates a more precise date range of ~740–550 cal. BP based on Bayesian modeling of AMS radiocarbon dates on short-lived macrobotanical remains and bone fragments [120]. These excavations also yielded the largest faunal assemblage of the sites included in this study.

Before ZooMS analysis, each assemblage underwent a complete zooarchaeological analysis by AJ. This included all specimens that were either identifiable to element or measuring at least 20 mm. Each specimen was identified to element and portion, and to the narrowest taxonomic category (e.g. "Mammal", "Bovid", "*Bos* sp.") based upon morphology. When possible, size was recorded, using Brain's [1] bovid size classes. In most cases this was not possible, and thus the specimen was assigned to a minimum size class based upon thickness of cortical bone





and identifiability relative to specimen size (e.g. mammal  $\geq 2$ , indicating the specimen came from size 2 or larger mammal).

The sites analyzed in this study have small faunal assemblages, ranging from 1 to 522 in number of identifiable specimens (NISP). Fragmentation rates are considerable at all sites (Fig 6)—the modal fragment size for all sites except for Jakobo West (ZAM, NISP = 7) is between 20 and 30mm. Such high rates of bone comminution rendered most fragments identifiable only to broader taxonomic groupings (S2 Table). ZooMS analysis of bone fragments was carried out on 100% of specimens from Jakobo West, 4 of 6 NISP from Muteteshi, 9 of 24 NISP from Salumano and 69 of 84 NISP from Kapiri Mposhi, excluding fragments that were clearly not mammalian, exhibited signs of burning, or were in such poor condition that collagen extraction was highly unlikely. Finally, 102 specimens, ~20% of the large assemblage from Kalundu (NISP = 522), were analyzed.

#### Collagen extraction and digestion

For both archeological and modern bones, 10–20 mg of bone was subsampled and collagen was extracted based upon published ZooMS extraction protocols [38, 42, 121]. Bone powder or chips were demineralized in 500  $\mu$ l of 0.5M hydrochloric acid at 4°C for one to two days.

The supernatant was removed and the samples were rinsed twice with 50mM ammonium bicarbonate (NH<sub>4</sub>HCO<sub>3</sub>), pH 8.0 (AmBic). Then the samples were incubated in 100  $\mu$ l of AmBic at 65 °C for 1 hour. 50  $\mu$ L of the resulting supernatant was incubated with 0.2  $\mu$ g of trypsin (ThermoFisher Pierce<sup>TT</sup> Trypsin Protease) at 37 °C overnight. Following digestion, the samples were acidified to 0.1% trifuoroacitic acid (TFA) and purified using a 100  $\mu$ l C18 resin ZipTip<sup>®</sup> pipette tip (EMD Millipore) with conditioning and eluting solutions composed of 50% acetonitrile and 0.1% TFA and a lower hydrophobicity wash buffer of 0.1% TFA. Peptides were eluted in 50  $\mu$ l.

## Peptide mass fingerprinting

Tryptic collagen peptides from both modern and archaeological samples were diluted 1:10 with the eluting solution, and  $0.5\mu$ l of the resulting tryptic peptides were mixed with a 0.5  $\mu$ l of  $\alpha$ -cyano-4-hydroxycinnamic acid matrix solution and spotted in triplicate along with calibration standards onto a ground steel 385 spot MALDI target plate. Samples were run on a Bruker Autoflex Speed MALDI-TOF mass spectrometer (Bruker Daltonics). Spectra were visually inspected using the mMass software [122] and low-quality spectra were rerun on the MALDI-TOF-MS without dilution or at 1:100 dilution depending upon the quality of the sample. MALDI-TOF-MS spectra for the modern references and protein sequence data can be found at Zenodo, doi:10.5281/zenodo.3964709 and for the archaeological samples at Zenodo, doi:10. 5281/zenodo.3971142.

## LC-MS/MS

In order to confirm the sequences of peptide markers, one sample from each of the modern reference species was characterized by LC-MS/MS. The individuals chosen had the highest quality MALDI-TOF-MS spectra. 25µl of the tryptic peptide extract was sent for LC-MS/MS at the Functional Genomics Center Zurich using a Q-Exactive HF mass spectrometer (Thermo Scientific, Bremen, Germany) coupled with an ACQUITY UPLC M-Class system (Waters AG, Baden-Dättwil, Switzerland). For liquid chromatography, 0.1% formic acid in H2O was used for channel A and 0.1% formic acid in acetonitrile for channel B. Column temperature was  $50^{\circ}$ C. For each peptide sample, 4  $\mu$ L was loaded onto the column with a flow rate of 15  $\mu$ l/min of 99% solvent A for 30 seconds at room temperature. Eluting peptides were separated on a C18 column (HSS T3 C18, 100Å, 1.8 µm, 75 µm x 250 mm, Waters AG, Baden-Dättwil, Switzerland). After gradient stabilization for 1.5 min at 99% solvent A, peptides were separated over 120 minutes with a gradient of 1% - 5% solvent B in 30 seconds, 5% - 40% solvent B in 120 min. The column was cleaned after the run with 98% solvent B for 5 min and holding 98% B for 8 min prior to re-establishing loading condition. Full scan MS spectra (350-1500 m/z) were acquired in the Orbitrap with a resolution of 120'000 at 200 m/z, an automatic gain control target of 3e6, and maximum injection time of 50 ms. Precursors for MS2 scans were isolated with an isolation window of 1.2 Da, automated gain control of 1e5, a maximum fill time of 50 ms and HCD fragmented with a normalized collision energy of 28. From each MS1 scan, the 12 most intense precursor ions were fragmented and scanned with a resolution of 30'000 at 200 m/z and a fixed first mass of 130 m/z. For MS2 selection, the filter intensity threshold was 9e4. Unassigned singly charged ions and ions were excluded. Precursor masses already selected for MS/MS were excluded for further selection for 30 seconds. LC-MS/MS data is available at PXD020810 through MassIVE (doi:10.25345/C5TJ3M).

LC-MS/MS data was then quality checked using Byonic v.3.2.0 (Protein Metrics Inc. [123]) against SwissProt (downloaded 30 April 2019) with the following parameters: cleavage sites fully specific C-term R and K; 2 missed cleavages allowed; mass changes– 3 common, 1 rare;

common: oxidation on M, P, and K, deamidation on N and Q; rare: pyro-Glu on N-term Q and E; no sequence variations allowed, wildcard search disabled, with decoys to confirm that the peptides were primarily Bovine collagen and to identify other significant protein hits. Proteins were considered present if they had 4 or more peptides with PEP 2D scores < 0.001 and a log probability > 5 (S3 and S4 Tables).

## Biomarker identification and confirmation

*Bos taurus* collagen protein sequences (P02453, P02465) were used to locate the collagen genes in the genomic data from bovids [124]. Collagen sequences were obtained from genomic data as nucleotide sequences, the exons were identified where possible and translated. Collagen sequences for all of the species were aligned in Geneious Prime 2019.0.4 (https://www. geneious.com). The locations of the nine well-characterized mammal markers [38, 42] were mapped onto the aligned sequences to predict the masses of these peptides in each species for which genetic data was available (S1 File). MALDI data for the reference samples, was peakpicked in mMass [122], with a signal to noise ratio of 6 and screened for peaks at the locations of the predicted masses. Markers were considered present for a species if they were found in 2 or more individuals at the predicted masses.

For confirmation of biomarkers, the MS/MS spectral data was then analyzed using Byonic with the following parameters: cleavage sites fully specific C-term R and K; 2 missed cleavages allowed; mass changes– 5 common, 1 rare; common: oxidation on M, P, and K, deamidation on N and Q; rare: pyro-Glu on N-term Q and E; no sequence variations allowed, wildcard search disabled, with decoys. The database was formed from the published sequences from the bovids (*Bos taurus* (P02453, P02465), *Bison bison* (XP\_010841089.1, XP\_010838069.1), *Bubalus bubalus* (XP\_006041214.2, XP\_006054012.1), *Ovis aries* (XP\_011983013.1, XP\_004007775.1), *Capra hircus* (XP\_017920382.1, XP\_005678993.1)), all of the mined bovid sequences from genetic data, the collagen sequences for outgroup species (*Odocolieus virginia-nus* (XP\_020769668.1, XP\_020769440.1), *Equus caballus* (XP\_023508478.1, XP\_001492989.1), the top proteins for the SwissProt search (168 proteins–S4 Table), and the common contaminants from Byonic. The peptide spectral matches (psms) to COL1 were filtered for an FDR < 1% and a PEP 2D score < 0.001. Markers were confirmed if there were at least three psms to the tryptic peptide corresponding to the genetic sequence of the marker with a precursor mass within 0.5 Da of the m/z identified in the MALDI spectra.

To identify new markers, the aligned bovine sequences were searched for sequence variants in Geneious. The masses of the tryptic peptides were predicted using PeptideMass from ExPASy [125, 126]. Peptides with masses between 800 and 3500 Daltons were then confirmed as authentic and the only peptide present at that mass in the LC-MS/MS results as above. MALDI-MS spectra were then analyzed for the consistent presence of any of these new candidate markers and only those which are present in over 75% of the reference spectra were presented as new diagnostic markers (S5 Table).

#### Identification of archaeological samples

The archaeological spectra were first screened for good signal to noise and collagen signal by eye. Using mMass [122], high quality spectra were identified by presence of shared collagen markers. Spectra were processed using baseline correction (precision 52, relative offset 39), smoothing (Savitzky-Golay method with a window size of 0.3 m/z), deisotoping (maximum charge 1, isotope mass tolerance 0.1 m/z, isotope intensity tolerance 50%, isotope mass shift 0) and then peakpicked (signal to noise threshold of 3, relative intensity threshold 1%). The

resulting peak lists were used to identify the samples to the lowest taxonomic level possible using the bovid markers.

## **Results and discussion**

#### Data quality control

**MALDI data.** All of the modern reference samples yielded high quality MALDI data with the presence of shared collagen peaks and the 9 published markers. For the 192 archaeological samples analyzed, 62 were able to be identified, 44 yielded partial identifications, one yielded collagen was not identifiable to a known taxon, and 85 did not yield sufficient collagen for identification (S2 Table).

LC-MS/MS data. LC-MS/MS data from all of the species tested had over 85% coverage of both COL1 proteins <u>S3 Table</u>). 168 proteins were found during the analysis against SwissProt between all of the samples (<u>S4 Table</u>). These proteins were added to the col1 proteins for subsequent biomarker determination and confirmation. There was 100% agreement between the genetic data, the LC-MS/MS data, and the MALDI-TOF data for all of the markers presented here this includes the LC-MS/MS data from impala and springbok which aligns with the published genetic sequences for both species.

**Common Bovidae ZooMS markers.** The results of the reference specimens presented here broadly align with genetic studies of African bovids [124, 127–130]. All bovid species studied share four of the existing ZooMS markers:  $COL1\alpha 2$  793–816 (D) at m/z 2131,  $COL1\alpha 2$  454–483 (E) at m/z 2792,  $COL1\alpha 1$  508–519 (P1) at m/z 1105, and  $COL1\alpha 2$  292–309 (P2) at m/z 1648 (Table 2; S1 and S2 Figs). The other five existing ZooMS markers vary among bovid groups, and therefore combinations of these markers can be used to uniquely identify members of subtribe Bubalina, Bovina, tribe Tragelaphini, and subfamilies Hippotraginae, Alcelaphinae, Aepycerotinae, Reduncinae, and Cephalophinae. Several genera from the polyphyletic subfamily Antiliopinae can also be identified uniquely (Table 2; S1 and S2 Figs).

**Novel ZooMS markers for bovids.** Our results provide two new markers,  $COL1\alpha2 375$  and  $COL1\alpha2 889-906$  (see S1 File for discussion on marker nomenclature), that can be used to differentiate among several bovid groups.  $COL1\alpha2 889-906$  has six versions in the genetic data which was supported by LC-MS/MS analysis (Table 3). Two of these versions have identical masses and therefore cannot be distinguished using peptide mass fingerprinting techniques (S5 Table). The version found in most bovids has a peak at m/z 1532 [33]. Caprinae, Bovinae, some members of Alcelaphinae and Cephalophinae have peaks at m/z 1560. However, a peak at m/z 1560 cannot be used reliably for identification because another peptide at m/z 1560 is found in all bovids.  $COL1\alpha2 889-906$  is most useful for helping to distinguish Hippotraginae (m/z 1588). It also can be used to identify the genus *Cephalophus* (m/z 1574) within Cephalophinae, and to identify *Alcelaphus* (m/z 1590) within Alcelaphinae.

Marker COL1 $\alpha$ 2 375 is more complicated to use as tryptic digestion is sometimes incomplete when the digestion site is followed by a proline (when indicating both markers only the first position is used in the name). This is illustrated by the presence of peaks corresponding to peptides with no missed cleavages (COL1 $\alpha$ 2 375–386) and one missed cleavage (COL1 $\alpha$ 2 375–396). Bovinae, Reduncinae, and some members of Antilopinae and Cephalophinae have peaks corresponding to COL1 $\alpha$ 2 375–396 (m/z 2056/2072) that cannot be used reliably for identifications because they are composed of several peptides. COL1 $\alpha$ 2 375–386 (m/z 1182) can also overlap with the COL1 $\alpha$ 2 978–990 (A') marker present at the same m/z value in some species of Neotragini (e.g. *Madoqua*). Therefore, caution is needed when using m/z 1182. However, m/z 1154 (COL1 $\alpha$ 2 375–386) and m/z 2028/2044 (COL1 $\alpha$ 2 375–396) can be used to uniquely identify Caprinae, Alcelaphinae, and Hippotraginae. The presence of peaks at m/z 1154/2028/

or Bovidae.
markers f
ZooMS
Table 2.

			COLIGI		102		1A237		L1a2	2011a2	COL1a2	COL1a2	COL1a2	COL1a2		[]a]	COI	1a2
			508-519	978-	990			484	4-498	389-906	502-519	292-309	793-816	454-483	586	-618	757-	789
Subfamily	Tribe	Subtribe or Genus	P1	A	A'				в		υ	P2	D	щ	ц	ŕ	IJ	ۍ ا
Bovinae	Bovini	Bovina**	1105	1192	1208	1182	2056 20	072 1	427	1532	1580	1648	2131	2792	2853	2869	3017	3033
Bovinae	Bovini	Bubalina	1105	1192	1208	1182	2056 20	072 1	455	1532	1580	1648	2131	2792	2853	2869	3059	3075
Bovinae	Tragelaphini	Group 1	1105	1192	1208	1182	2056 20	072 1	427	1560	1580	1648	2131	2792	2883	2899	3043	3059
Bovinae	Tragelaphini	Group 2*	1105	1192	1208	1182	2056 20	072 1	427	1532	1580	1648	2131	2792	2883	2899	3043	3059
A epycerotinae**			1105	1180	1196	1182	2056 20	072 1	427	1532	1550	1648	2131	2792	2883	2899	3017	3033
Antilopinae	Antilopini	Nanger, Eudorcas, Antidorcas**	1105	1180	1196	1182	2056 20	072 1	427	1532	1550	1648	2131	2792	2883	2899	3017	3033
Antilopinae	Neotragini	Raphicerus*, Ourebia	1105	1180	1196	1182	2056 20	072 1	427	1532	1550	1648	2131	2792	2883	2899	3017	3033
Antilopinae	Neotragini	Neotragus	1105	1180	1196	1182	2056 20	072 1	427	1532	1580	1648	2131	2792	2883	2899	3017	3033
Antilopinae	Neotragini	Oreotragus	1105	1192	1208	1154	2028 20	044 1	427	1532	1580	1648	2131	2792	2883	2899	3017	3033
Antilopinae	Neotragini	Madoqua	1105	1166	1182	1182	2056 20	072 1	427	1532	1550	1648	2131	2792	2883	2899	3017	3033
Cephalophinae		Sylvicapra	1105	1192	1208	1182	2056 20	072 1	427	1532	1580	1648	2131	2792	2853	2869	3043	3059
Cephalophinae		Cephalophus	1105	1192	1208	1154	2028 20	044 1	427	1574	1580	1648	2131	2792	2853	2869	3043	3059
Cephalophinae		$Philantomba^*$	1105	1192	1208	1154	2028 20	044 1	427	1560	1580	1648	2131	2792	2883	2899	3043	3059
Reduncinae			1105	1150	1166	1182	2056 20	072 1	427	1532	1550	1648	2131	2792	2883	2899	3017	3033
Hippotraginae			1105	1180	1196	1154	2028 20	044 1	427	1588	1580	1648	2131	2792	2883	2899	3043	3059
Alcelaphinae		Alcelaphus	1105	1180	1196	1154	2028 20	044 1	427	1590	1550	1648	2131	2792	2883	2899	3017	3033
Alcelaphinae		Connochaetes, Damaliscus	1105	1180	1196	1154	2028 20	044 1	427	1560	1550	1648	2131	2792	2883	2899	3017	3033
Pantholopinae*			1105	1180	1196	1182	2056 20	072 1	427	1560	1580	1648	2131	2792	2857	2873	3017	3033
Caprinae		Capra**	1105	1180	1196	1154	2028 20	044 1	427	1560	1580	1648	2131	2792	2883	2899	3077	3093
Caprinae		$Ovis^{**}$	1105	1180	1196	1154	2028 20	044 1	427	1560	1580	1648	2131	2792	2883	2899	3017	3033
Masses in dark sh	ided cells need	to be used carefully for in	nterpretatior	ı. Mass	es in li	ght shae	led cells	are prese	ent in th	e LC-MS/N	4S but are	not suitable	e for specie	s identifica	tion u	sing M	ALDI. I	3oth

COL102 375-386 and COL102 375-396 are present in the LC-MS/MS and MALDI data, so both are presented. Group 1: Taurotragus, Tragelaphus buxtoni, Tragelaphus euryceros; \*Predicted from the genetic sequence data only, not confirmed with MALDI or LC-MS/MS data.

\*\* Previously published species. Neotragus pygmaeus has markers different from the other Neotragus species predicted from the genetic data. See S1 File for more detail.

https://doi.org/10.1371/journal.pone.0251061.t002

Marker	Letter Name	Tryptic Name	Sequence	Mass	Mass'
COL1a2 978-	А	A2T85	AGQPGAVGPAGIR	1150	1166
990			TGQPGAVGPAGIR	1180	1196
			IGQPGAVGPAGIR	1192	1208
			SGQPGAVGPAGIR	1166	1182
COL1a2 375		A2T34	EGPVGLPGIDGR*	1182	
			EGPVGLPGIDGRPGPIGPAGAR*	2056	2072
			EGPAGLPGIDGR*	1154	
			EGPAGLPGIDGRPGPIGPAGAR*	2028	2044
COL1a2 484-	В	A2T43	GIPGEFGLPGPAGAR	1427	
498			GIPGEFGLPGPAGVR	1455	
COL1a2 889-		A2T75	GEPGPAGAVGPAGAVGPR	1532	
906			GEPGPVGAVGPAGAVGPR	1560	
			GEPGPAGVVGPAGAVGPR	1560	
			GEPGPVGAVGPTGAVGPR	1590	
			GEPGPVGAVGPVGAVGPR	1588	
			GEPGPVGAIGPAGAVGPR	1574	
COL1a2 502-	С	A2T45	GPPGESGAAGPAGPIGSR	1550	
519			GPPGESGAAGPTGPIGSR	1580	
COL1a1 586-	F	A1T55/56	GLTGPIGPPGPAGAPGDKGEAGPSGPAGPTGAR	2853	2869
618			GLTGPIGPPGPAGAPGDKGETGPSGPAGPTGAR	2883	2899
			GLTGPIGPPGPAGAAGDKGETGPSGPAGPTGAR	2857	2873
COL1a2 757-	G	A2T67	GPSGEPGTAGPPGTPGPQGLLGAPGFLGLPGSR	3017	3033
789			GPSGEPGTAGPPGTPGPQGFLGPGFLGLPGSR	3077	3093
			GPSGEPGTAGPPGTPGPQGLLGPPGFLGLPGSR	3043	3059
			GPSGEPGTAGPPGTPGPQGLLGLPGFLGLPGSR	3059	3075

Table 3. Sequences of markers used to differentiate members of Bovidae.

The sequences correspond to the masses for the diagnostic markers in Table 2. The names of the markers in different naming conventions used for ZooMS analysis in previous publications are also presented where relevant. Bold letters in the sequence data indicate locations of amino acid differences between the different versions of the markers. Masses in dark shaded cells need to be used carefully for interpretation. Masses in light shaded cells are present in the LC MS/MS, but are not suitable for species identification using MALDI.

\*Both the fully cleaved peptide and the peptide with one missed cleavage are present in the LC-MS/MS and the MALDI.

https://doi.org/10.1371/journal.pone.0251061.t003

2044 can also be used to identify the genera *Cephalophus* and *Philantomba* within Cephalophinae and *Oreotragus* within Antilopinae.

**G peptide (COL1a2 757–789).** The high mass COL1a2 757–789 (G and G') peptides are commonly used for identification of sheep and goat [111]. They also show diversity among other bovid groups. The version seen in sheep and cattle has peaks at m/z 3017 and 3033. Frequently however, a peak at m/z 3017 (but not m/z 3033) has been observed in *Capra* spectra along with their unique markers (m/z 3077/3093) (Fig 7; S3 Fig). In investigating the cause for this, our LC-MS/MS data showed that another peptide, conserved across all of the bovid species as shown from the sequence data, is present at m/z 3017 (GPPGASGAPGPQGFQGPPGE PGEPGQTGPAGAR–COL1a2 89–130) (S3 Fig). This not the same peptide used in the identification of sheep/cattle and is not a reliable marker for ZooMS identification. Therefore, we suggest that m/z 3017 should no longer be used in the identification of bovids as there is a high



Fig 7. Examples of differences between bovid groups for some published marker sets. Panel A shows the difference between the COL1 $\alpha$ 2 484–498 (B) marker at m/z 1455 for *Bubalina* (shown: *Syncerus caffer*) and m/z 1427 for all other bovids (shown: *Bos taurus*). Panel B shows the difference between the COL1 $\alpha$ 2 502–519 (C) marker at m/z 1580 (shown: *Hippotragus niger*) and at m/z 1550 (shown: *Eudorcas thomsonii*). Panel C shows the difference between the COL1 $\alpha$ 2 757–789 (G/G') marker at 3033 (shown: *Alcelaphus buselaphus*, G' only), m/z 3043/3059 (shown: *Tragelaphus eurycerus*), and m/z 3059/3075 (shown: *Syncerus caffer*). The peak at m/z 3017 is composed of multiple peptides, one of which is shared between all Bovids (shown in *Syncerus caffer*), and therefore should not be used for identification.

likelihood for misidentification. The peak at m/z 3033, should still be considered a reliable marker for COL1 $\alpha$ 2 757–789 (G') and can continue to be used for identification without its counterpart.

#### Distinguishing among bovid groups

**Subfamily Bovinae.** All members of the subfamily Bovinae share the COL1 $\alpha$ 2 978–990 (A/A') marker at m/z 1192/1208 and the COL1 $\alpha$ 2 502–519 (C) marker at m/z 1580. Within Bovinae are two tribes, Tragelaphini and Bovini, which can be distinguished from each other using the COL1 $\alpha$ 1 586–618 (F/F') marker with m/z of 2853/2869 for Bovinae and m/z 2883/2869 for Tragelaphini, and the COL1 $\alpha$ 2 757–789 (G') marker with an m/z of 3033 for Bovinae and 3043/3059 for Tragelaphini (Table 2). Both Tragelaphini and Bovini share the new marker COL1 $\alpha$ 2 375 (m/z 1182/2056/2072), but within Tragelaphini there is variation in COL1 $\alpha$ 2 889–906. *Tragelaphus imberis* (lesser kudu), *Tragelaphus scriptus* (harnessed bushbuck), *Tragelaphus spekii* (sitatunga), and *Tragelaphus strepsiceros* (greater kudu) share the variation at m/z 1530. *Tragelaphus buxtoni* (mountain nyala), *Tragelaphus euryceros* (bongo) and both species of *Taurotragus* (elands) have a peak at m/z 1560; so care should be taken when using this marker to distinguish between tragelaphines, as the latter is not a reliable indicator. In addition, because there is variation with the genus *Tragelaphus*, only the specific species reported here can be identified using COL1 $\alpha$ 2 889–906.

Bovini includes two subtribes of which members have known peptide markers: Bovina, containing *Bos* and *Bison* and Bubalina, containing *Bubalus* (water buffalo and their relatives) and *Syncerus caffer* (African buffalo). We confirm that *Syncerus* shares the same COL1 $\alpha$ 2 484–498 (B) marker at m/z 1455 and COL1 $\alpha$ 2 757–789 (G/G') markers at m/z 3059/3075 with *Bubalus* [38, 112], allowing for African buffalo to be distinguished from cattle.

**Subfamilies Reduncinae and Hippotraginae.** Members of Reduncinae can be distinguished from all other bovid groups by the COL1 $\alpha$ 2 978–990 (A/A') markers at m/z 1150/1166. In Reduncinae the COL1 $\alpha$ 2 502–519 (C), COL1 $\alpha$ 1 586–618 (F/F'), and COL1 $\alpha$ 2 757–789 (G/G') markers are shared with several other groups, with a peak at m/z 1550, m/z 2883/2899, and m/z 3043/3059, respectively (Table 2). Members of Hippotraginae share COL1 $\alpha$ 2 978–990 (A), COL1 $\alpha$ 2 484–498 (B), COL1 $\alpha$ 2 502–519 (C), COL1 $\alpha$ 1 586–618 (F), and COL1 $\alpha$ 2 757–789 (G) markers with other groups, but can be uniquely identified with the new marker COL1 $\alpha$ 2 889 at m/z 1572 and 1588 (Table 2). Caution should be used with m/z 1588 as it overlaps with a shared collagen peptide at m/z 1586 (Fig 8). In addition, the combination of the COL1 $\alpha$ 2 978–990 (A/A') marker at m/z 1180/1196 and the COL1 $\alpha$ 2 757–789 (G/G') marker at m/z 3043/3059 can also be used to uniquely identify Hippotraginae.

**Subfamily Alcelaphinae.** All Alcelaphinae analyzed in this study share COL1 $\alpha$ 2 978–990 (A/A') marker at m/z 1180/1196, COL1 $\alpha$ 2 502–519 (C) marker at m/z 1550, COL1 $\alpha$ 1 586–618 (F/F') at m/z 2883/2899, and COL1 $\alpha$ 2 757–789 (G/G') at m/z 3017/3033, and thus they are indistinguishable from *Aepyceros melampus* (impala), *Nanger granti* (Grant's gazelle), *Eudorcas thomsonii* (Thomson's gazelle), and *Ourebia ourebi* (oribi) using only the existing markers. However, all alcelaphine bovids have COL1 $\alpha$ 2 375 at m/z 1154/2056/2072, which can be used to distinguish them uniquely. Furthermore, the new COL1 $\alpha$ 2 889–906 marker at m/z 1590 can be used to uniquely identify the genus *Alcelaphus* (hartebeest). The two other genera we analyzed, *Connochaetes* (wildebeest) and members of the genus *Damaliscus* (topi and tsessebe) have m/z 1560, which is not useful for identification.

**Subfamily Cephalophinae.** We only analyzed members of two genera, *Cephalophus* and *Sylvicapra*, which share all of the existing ZooMS markers with members of the subtribe Bovina except for the COL1 $\alpha$ 2 757–789 (G/G') marker, which in Cephalophinae has an m/z of



Fig 8. Novel markers shown for four different families. Most bovids share the  $COL1\alpha 2$  889–906 marker at m/z 1532 (shown: *Syncerus caffer*). Hippotraginae can be distinguished using m/z 1588. However, this must be used carefully as the marker overlaps with the peak shared by all species at m/z 1586 (shown: *Hippotragus equinus*). Members of the genus *Cephalophus* can be identified based upon the  $COL1\alpha 2$  889–906 marker at m/z 1574 (shown: *Cephalophus adersi*) and the genus *Alcelaphus* can be identified from other Alcelaphines at m/z 1590 (shown: *Alcelaphus buselaphus*). The  $COL1\alpha 2$  375–396 marker is shown (m/z 2056 and 2072, shown in *Hippotragus equinus*). Subfamilies Hippotraginae, Caprinae, and Alcelaphinae, along with *Oreotragus* and some genera of Cephalophinae can be distinguished with markers at m/z 2028 and m/z 2044. *Syncerus caffer* has peaks at m/z 2056/2072, which are shared with another peptide present in all bovids and therefore not suitable for diagnostic purposes.

3043/3059 (Table 2). Cephalophus adersi (Aders's duiker) and Sylvicapra grimmia (bush duiker) share the same existing marker profile but can be distinguished based upon the two novel markers. Cephalophus adersi can be identified by COL1 $\alpha$ 2 375 at m/z 1182/2056/2072 and COL1 $\alpha$ 2 889–906 at m/z 1574 and Sylvicapra grimmia by COL1 $\alpha$ 2 889–906 at m/z 1532 (Table 2). Genetic data for a third species, Philantomba maxwellii (Maxwell's duiker), indicates that this genus could be uniquely identified using the COL1 $\alpha$ 1 586–618 (F/F') and COL1 $\alpha$ 2 375 markers. While we consider all three genera within the subfamily Cephalophinae here, many species have not been assessed, particularly within Cephalophus which includes 15 extant species. These taxa are also widespread across the continent, particularly in forest zones (Fig 1). Given the differences we have presented here and the diversity within Cephalophinae [131] it is possible that other members of Cephalophinae might have different marker profiles (e.g. different masses for various markers), so care should be taken when identifying new species of Cephalophinae, verifying ZooMS markers with LC-MS/MS and/or genetic data.

**Subfamily Antilopinae.** There is considerable diversity within Antilopinae. Traditionally members of this subfamily have been grouped into two tribes: Antilopini (gazelles and their relatives)—for which the specimens analyzed in this study include *Eudrocas thomsonii* (Thomson's gazelle) and *Nanger granti* (Grant's gazelle)—and Neotragini (African dwarf antelopes), for which the specimens analyzed in this study historically has included *Ourebia ourebi, Neotragus* spp., *Madoqua* (dik-diks), and *Oreotragus* (klipspringer). However, genetic studies have shown that Antilopinae is actually polyphyletic [124, 127, 132]. In particular, the tribe Neotragini has long been considered a catch-all group for small antelopes, which is reflected in the

lack of consistent markers for the entire tribe with four different marker profiles for the five genera studied.

All Antilopinae share the same COL1 $\alpha$ 1 586–618 (F/F') markers at m/z 2883/2899, COL1a2 757-789 (G') marker at m/z 3033, and COL1a2 889-906 marker at m/z 1532. All studied Antiopini and some of the Neotraini (genera Ourebia and Raphicerus) all share the existing marker profiles with Alcelaphinae and Aepyceros (Table 2). While  $COL1\alpha 2$  375 allows Alcelaphinae to be uniquely identified, Antilopini, Ourebia, and Raphicerus have the marker at m/z 1182/2056/2072 and cannot be distinguished. Several of the neotragine bovids have unique profiles. Oreotragus oreotragus (klipspringer) is highly divergent from the other Neotragini with the COL1α2 978–990 (A/A') marker at m/z 1192/1208, COL1α2 502–519 (C) marker at m/z 1580, and the COL1α2 375 marker at m/z 1154/2028/2044. Madoqua kirkii (Kirk's dikdik) has a unique COL1a2 978–990 (A/A') marker at m/z 1166/1182, distinguishing it from all other bovid groups (Table 2). Neotragus moschatus (suni) can be distinguished from the other Antilopinae species with a combination of  $COL1\alpha 2\ 978-990$  (A/A') markers at m/z 1180/1196 and COL1a2 502-519 (C) marker at m/z 1580. Although Neotragus moschatus and Ovis share identical existing marker sets, they can be uniquely distinguished by the presence of the COL1 $\alpha$ 2 375 marker in Ovis at m/z 1154/2028/2044 and the COL1 $\alpha$ 2 889–906 marker in Neotragus moschatus at m/z 1532. Given the diversity within the Antilopinae, the markers are indicated here only for the species tested. Further sampling for ZooMS and LC-MS/MS analysis will need to be done to confirm markers of other taxa within this subfamily, such as gerenuk (Litocranius walleri) and steenbok and grysbok (genus Raphicerus).

The results of this study show that it is possible to distinguish among major bovid groups and securely identify domestic livestock from wild bovids in sub-Saharan Africa using ZooMS [38]. In some cases, body size can also be used to refine ZooMS identifications when only partial ZooMS spectra are recovered (see also [47]). For example, *Bos, Cephalophus*, and *Sylvicapra* share all markers but the COL1 $\alpha$ 2 757–789 (G') peptide. The substantial size difference between *Bos* and members of Cephalophinae can be used to differentiate these taxa in cases when the COL1 $\alpha$ 2 757–789 (G') peptide is not preserved, which is a common occurrence (S2 Table).

In some cases identification is more difficult. *Aepyceros melampus, Nanger granti, Eudorcas thomsonii, Ourebia ourebi*, and *Raphicerus* all share the same suite of peptide markers. Impala and Grant's gazelle are similar in size, and while both are larger than *Eudorcas, Ourebia*, and especially *Raphicerus* [52], it is likely that fragmentary bone—not identifiable to element— would not be attributable to a size class, and therefore will not be useful in refining ZooMS-based identifications. The presence of juvenile specimens presents another confounding factor, particularly in cases where bone fragments are not identifiable to element.

### Comparison with published markers

Our study provides a comprehensive set of markers for the bovid groups, and our results compare favorably with published markers in other studies. Our results agree with the incomplete published marker profiles for *Alcelaphus buselaphus* [48], members of Antilopini from [48, 49, 112], and *Antidorcas marsupialis* and *Sylvicapra grimmia* from [33]. Bradfield and colleagues [47] present incomplete marker profiles for a number of bovid species that were unconfirmed with LC-MS/MS. In three cases our markers differ from those presented in [47]. We confirm using both sequence and LC-MS/MS data that *Connochaetes taurinus* (blue wildebeest) has the same existing marker profiles as the other alcelaphines, contradicting the differences in the COL1 $\alpha$ 2 978–990 (A) and COL1 $\alpha$ 2 757–789 (G) markers reported by Bradfield et al. [47]. In our work all Hippotragine species tested have the same COL1 $\alpha$ 2 757–789 (G/G') markers at m/z 3043/3059. In addition, we do not find the differences in the existing markers between Tragelaphini species as reported by Bradfield et al. [47]. We analyzed predicted protein sequences and/or LC-MS/MS data for two species from the genus *Taurotragus* and six from *Tragelaphus*, including two of the three species that were analyzed in [47] and find that the COL1 $\alpha$ 2 757–789 (G) makers are identical across all eight Tragelaphini species tested. The difference in the COL1 $\alpha$ 2 757–789 (G) maker profiles may be related to the difficulty of using the peak at m/z 3017 which is why we recommend that it no longer be used to distinguish between bovid species. Our LC-MS/MS analysis of the data from Le Meillour et al. [49] agrees with that of Coutu et al. [33] in that we have identified the same inconsistencies in the LC-MS/MS data and their published full sequence. We do not challenge species identifications proposed by Le Meillour et al. (2020), only additional parts of the sequence data which were not used for their identifications, but our used in this study and in Coutu et al. [33]. These differences highlight the importance of confirming markers using sequence and/or LC-MS/MS data to ensure accuracy.

#### ZooMS analysis and archaeofaunal identifications

Collagen preservation among the sampled archaeological sites is extremely uneven. Eight of nine specimens analyzed from Salumano yielded good spectra and taxonomic identifications, but none of the four analyzed specimens from Muteteshi yielded any collagen, and only eight of 69 specimens analyzed from Kapiri Mposhi yielded complete spectra for identification beyond family, while another three were identifiable to other families (e.g. Leporidae) (Table 4). Forty-one of 102 specimens from Kalundu yielded good spectra sufficient for identifications (Table 4). Only two of the Kalundu specimens were not attributable to any known taxa on the basis of ZooMS. One of the four good spectra obtained from the seven analyzed Jakobo West specimens was not identifiable to a known taxon. Finally, the single specimen analyzed from Fibobe yielded identifiable results. The detailed results for all sites are presented

Site	Bone recove site by wei	ered fro ght in §	m each grams	R	esults of	f ZooMS analysis (number	of specimens)	
	ID/ analyzed	NID	Total	Analyzed	Full ID	Full spectra (unknown animal)	Partial spectra and ID	Fail
Kapiri Mposhi	119.1	7.4	126.5	69	11	1	12	45
Muteteshi	5.1	0	5.1	4	-	-	-	4
Fibobe	1.1	0	1.1	1	-	_	1	-
Salumano	58.5	11.8	70.6	9	7	_	1	1
Kalundu	1280.2	185.7	1465.9	102	41	-	29	32
Jakobo West	6.9	1.0	7.9	7	3	1	-	3

Table 4. Identifiable/analyzed versus unidentifiable bone recovered from each site by weight in grams, and number of specimens analyzed and number of successful, partial, and failed spectra.

For bone recovered from the site, "ID/analyzed" includes the fraction of the assemblage that measured at least 20 mm in maximum dimension or was either identifiable to element or taxon. "NID" includes all other faunal material that was unidentifiable and measuring under 20 mm. "Full ID" includes bovids identified to subfamily or narrower taxonomic grouping, as well as wild species identified to family (e.g. Mustelidae, Leporidae). "Full spectra (unknown animal)" includes specimens that yielded good spectra but could not be attributed to any taxon because many African taxa do not have published peptide markers. "Partial spectra and ID" includes specimens that yielded some peptide markers, but not enough for definitive identifications. "Fail" includes specimens that did not produce any peptide markers.

https://doi.org/10.1371/journal.pone.0251061.t004

in S2 Table (Table 4). In sum, c. 44% of analyzed specimens failed to yield collagen. A range of factors likely contributed to the lack of collagen preservation at some sites. Kalahari sands potentially impacted preservation at Jakobo West, consistent with other studies of collagen preservation in sandy sediments [113], although nearby Salumano yielded better results. More generally, collagen preservation in tropical areas can be poor due to high temperatures, fluctuations in temperature, and a dynamic hydrologic environment [133, 134]. Culinary processing, including boiling, may also be responsible for loss of collagen [135]. Future ZooMS analyses would benefit from additional screening techniques to assess collagen preservation prior to collagen extraction and analyses [105, 109, 136].

Specimens yielding full and partial spectra spectra revealed considerably more taxonomic diversity at the sites than did morphological identifications. In all sites except for Muteteshi in which all four specimens failed to yield sufficient or well-preserved collagen for ZooMSbased taxonomic identification—taxonomic richness (here NTAXA includes bovid group or other family such as Canidae or Leporidae) increased with ZooMS analysis (Fig 9; Table 5). Analysis of the archaeofaunal remains show that, in some cases, ZooMS analysis of even the smallest assemblages can reveal greater taxonomic richness than morphological identifications. Analysis of only 9 specimens from Salumano resulted in identification of three taxa. The two largest assemblages, Kalundu and Kapiri Mposhi, also saw increases in NTAXA. Better preservation at Kalundu resulted in the greatest NTAXA increase, from 3 to 10. Poor preservation at Kapiri Mposhi resulted in many failed samples and several partial identifications (e.g., Aepycerotinae/Alcelaphinae/Nanger/Eudorcas/Raphicerus/Oribi). However, because these specimens were originally identified as either indeterminate mammals or bovids, these partial identifications are still an improvement over identifications based solely upon morphology, and demonstrate the presence of wild bovids at the site (S2 Table).

The most common identification at the archaeological sites was to Bos, with 50 identifications, followed by Cephalophinae with 9 identifications. These include ZooMS-based identifications informed by specimen size (see below). Not all markers are needed for certain identifications. For example, the identification of Reduncinae does not require the COL1 $\alpha$ 2 757-789 (G/G') markers if the COL1 $\alpha$ 2 978–990 (A/A') markers are present. In some cases where necessary markers were not visible, specimen size could be used to narrow the identification: for example, when the COL1 $\alpha$ 2 757–789 (G/G') marker was missing, size could be used to distinguish Bos from Sylvicapra or Cephalophus (Table 4; S2 Table). Some specimens had peptide markers that matched closely related species for which peptides are known. For example, one specimen each from Kalundu and Salumano had markers matching those published for suids [38]. Those markers also agree with the partial set identified for warthog (Phacochoerus africanus) by [47], and therefore this identification of suid is likely correct. Similarly, several fragments from small mammals had peptide markers matching those of leporids [112], and although not all peptide markers for members of the Leporidae are known, these identifications are likely correct, and morphology is consistent with Leporidae. Finally, one specimen also yielded good spectra, but was not attributable to any known taxon (Table 3; S2 Table) A wide range of wild fauna, including viverrids, rodents, and mustelids were recovered from earlier excavations at Kalundu [98]. Peptide markers for many of these African taxa have not been established and therefore further work on a comprehensive reference library is needed before they can be identified using ZooMS.

#### Herding economies and the persistence of hunting in Iron Age Zambia

Despite uneven preservation and the often coarse nature of ZooMS-based identifications, the results of our analysis provide insights into the Iron Age economies of southwestern and



Fig 9. Percent of sampled specimens identified to broad taxonomic categories. Upper chart shows identifications based on morphology alone. Lower chart shows identifications using ZooMS and morphology.

central Zambia. Cattle were identified at every site except for Muteteshi, which had a small sample size (n = 4) and exceptionally poor collagen preservation. Caprines were only identified at Kalundu—evidenced by a single specimen. Kalundu, Kapiri Mposhi, and Salumano also yielded evidence for wild bovids, but the three smallest assemblages did not. The wild bovid assemblage at Kalundu is largely comprised of duikers, but these were not identified at Kapiri Mposhi, although based on partial spectra, one reduncine and several less identifiable wild bovids were recognized in the assemblage. The Kalundu assemblage also yielded the largest diversity of wild fauna, including Hippotragine, Tragelaphine, and Antilopine bovids, as well as other large mammals, such as suid, elephant, and rhinoceros. The ZooMS-based

Taxon	KAV Morph	KAV ZooMS + Morph	KMP Morph	KMP ZooMS + Morph	SAL morph	SAL ZooMS + Morph	ZAM Morph	ZAM ZooMS + Morph	FIB Morph	FIB ZooMS + Morph	MTS Morph	MTS ZooMS + Morph
Subfamily/tribe	7	56	0	9	0	6	0	2	0	1	0	0
Bos	-	35	_	7	-	5	-	2	-	1	-	-
Cf. Bos	5	5	_	_	_	_	_	-	_	-	_	-
Tragelaphini	_	2	_	_	_	1	_	-	_	-	_	-
Cephalophinae cf. Sylvicapra	-	4	-	-	-	-	-	-	_	-	-	-
Cephalophinae	-	5	_	-	-	-	-	-	_	-	-	_
Hippotraginae	-	2	-	-	-	-	-	-	-	-	-	-
Reduncinae	-	-	-	2	-	-	-	-	-	-	-	-
Antilopini: Eudorcas/ Ourebia	-	-	-	-	-	-	-	-	_	-	-	-
Caprinae	1	1	-	-	-	-	-	-	-	-	-	-
Elephant	-	1	-	-	-	-	-	-	-	-	-	-
Rhinoceros	-	1	-	-	-	-	-	-	-	-	-	-
Cf. Human	1	-	-	-	-	-	-	-	-	-	-	-
Family	26	23	4	17	1	2	0	2	0	0	0	0
Mustelidae	-	-	-	1	-	-	-	-	-	-	-	-
Canidae	-	2	-	-	-	-	-	-	-	-	-	-
cf. Suid	-	1	-	-	-	1	-	-	-	-	-	-
Bos/Tragelaphini size $\geq 3$	-	2	-	-	-	-	-	-	-	-	-	-
Bos/Cephalophinae	-	1	-	1	-	-	-	-	-	-	-	-
Bos/Cephalophinae cf. Sylvicapra	-	4	-	3	-	-	-	2	-	-	-	-
Bos/Tragelaphini/ Cephalophinae	-	-	-	1	-	-	-	-	-	-	-	-
Bos/Tragelaphini/ Cephalophinae cf. <i>Sylvicapra</i>	-	-	_	-	-	-	-	_	-	-	-	_
Hippotraginae/ Tragelaphini	-	1	-	-	-	-	-	-	-	-	-	-
Eudorcas/Ourebia	-	1	-	-	-	-	-	-	-	-	-	-
Bos/Tragelaphinae/ Oreotragus/ Cephalophinae	-	1	_	-	-	-	-	-	-	-	-	-
Aepycerotinae/ Alcelaphinae/ Antilopinae/Caprinae/ Hippotraginae	_	_	_	1	_	-	-	_	_	-	-	_
Aepycerotinae/ Antilopinae/Reduncinae/ Caprinae—Ovis	-	1	_	-	-	-	-	-	_	-	-	_
Alcelaphinae/ Aepycereotinae/Nanger/ Eudorcas/Ourebia	-	-	-	1	-	1	-	-	-	-	-	-
Aepycerotinae/Nanger/ Eudorcas/Raphicerus/ Ourebia	-	-	_	1	-	-	-	-	_	-	-	-
Aepycerotinae/Nanger/ Eudorcas/Raphicerus/ Ourebia/Madoqua/ Reduncinae	_	-	_	2	-	-	_	-	_	-	-	-

#### Table 5. Identifications of analyzed archaeological specimens.

(Continued)

Taxon	KAV Morph	KAV ZooMS + Morph	KMP Morph	KMP ZooMS + Morph	SAL morph	SAL ZooMS + Morph	ZAM Morph	ZAM ZooMS + Morph	FIB Morph	FIB ZooMS + Morph	MTS Morph	MTS ZooMS + Morph
Bovid 2	6	-	1	1	-	-	-	-	-	-	-	-
Bovid 3	4	-	1	1	-	-	-	-	-	-	-	-
Cf. Bovid 3	1	-	-	-	-	-	-	-	-	-	-	-
Leporidae	1	3	-	3	-	-	-	-	-	-	-	-
Bovid 2/3	-	-	1	-	-	-	-	-	-	-	-	-
Bovid $\geq 3$	14	6	-	-	1	-	-	-	-	-	-	-
Rodentia	-		1	1	-	-	-	-	-	-	-	-
Mammal size/indet	69	23	65	43	8	1	7	3	1	0	4	4
Felidae/Canidae/ Loxodonta/Mustelidae	-	-	-	1	-	-	-	-	-	-	-	-
Felidae/Mustelidae/ Hyaenidae	-	1	-	-	-	-	-	-	-	-	-	-
Hypsodont $\geq 2$	1	1	-	-	-	-	-	-	-	-	-	-
Hypsodont $\geq 3$	8	1	1	1	-	-	-	-	1	-	-	-
Mammal 0.5	1	-	1	1	-	-	-	-	-	-	-	-
Mammal 1	3	-	5	3	-	-	-	-	-	-	-	-
Mammal 1 or 2	1	-	2	2	-	-	-	-	-	-	-	-
Mammal 2	7	2	5	3	1	-	-	-	-	-	1	1
Mammal 3	2	1	3	2	-	-	-	-	-	-	-	-
Mammal ≥3	22	10	-	-	3	-	-	-	-	-	-	-
Mammal ≥2	18	5	42	25	4	1	4	1	-	-	2	2
Mammal $\geq 1$	5	2	3	2	-	-	2	1	-	-	1	1
Vertebrate indet.	1	-	3	3	-	-	1	1	-	-	-	-
Total analyzed	102	102	69	69	9	9	7	7	1	1	4	4
NTAXA	3	10	1	5	1	3	0	1	0	1	0	0

#### Table 5. (Continued)

"Morph" indicates identifications using morphology only. "Morph + ZooMS" indicates identifications considering ZooMS results and morphology. Site codes: KAV, Kalundu Mound; KMP, SAL, Salumano; ZAM; Jakobo West Kapiri Mposhi; FIB, Fibobe II; MTS, Muteteshi.

https://doi.org/10.1371/journal.pone.0251061.t005

identifications from the sites analyzed here align well with reported faunal remains from previous excavations at these and other Iron Age sites in Zambia.

Analysis of fauna from previous excavations at Salumano (c. 2300 BP) reveal cattle in early horizons of the site, with caprines appearing later [116]. Similarly, cattle are the only domesticate present in the earlier component at Nanga (c. 1240 to 1070 BP) [101, 137], with cattle and caprines occurring together only in the later component of the site (c. 980 BP) [137]. These patterns point to a broadening of pastoral economies over the course of the Iron Age. Earlier excavations at Kalundu, as well as Isamu Pati, another Kalomo culture site, revealed that diverse wild fauna, particularly duiker, comprised a major portion of the assemblages [98]. Both Kalundu and Isamu Pati assemblages yielded cattle remains, as well as cattle figurines. Caprine remains are also present at both sites—cattle and caprine mandibles are in equal number at Isamu Pati, though no quantification data are given for the Kalundu caprines. Identifiable fauna from Iron Age levels at Makwe in eastern Zambia also show a predominance of cattle, in addition to wild fauna, primarily suids [100]. Identifiable fauna from the Early Iron Age site of Namakala (c.1400 BP) revealed only wild species [101, 138]. The continued reliance on wild fauna has been attributed to the constant threat of trypanosomiasis, communicated by the tsetse fly (*Glossina* spp.), which is abundant across southern and southeastern Zambia [139]. Caprines, particularly goats, are able to avoid tsetse fly blood meals much more effectively than are cattle [140]. Goats are mixed feeders that prefer to browse, and may be used to suppress brush, particularly in tsetse-ridden environments [141, 142]. Although sample sizes are small, the abundance of cattle and relative paucity of goat at Kalundu and Kapiri Mposhi hint at a relatively tsetse-free environment. Our analyses thus point to a unique pattern of cattle-based pastoralism supplemented with hunting, particularly duikers, as evidenced at Kalundu. These wild bovids would have provided an abundant source of meat—even intensive hunting as part of tsetse control efforts in the 1960s did not reduce duiker populations in east-ern Zambia [143].

## Conclusion

With the development of a reference set of peptide markers for African bovids, we demonstrate that identification to subfamily, and sometimes narrower taxonomic categories, is possible through ZooMS. By incorporating ZooMS into zooarchaeological analyses, the contours of foodways in Iron Age Zambia become more clear. Results show that in central and southwestern Zambia, Iron Age subsistence economies and foodways were based largely on cattle, but, as seen at Kalundu, wild bovids were also important. Further analyses of larger assemblages are necessary to reveal more detailed information on herding economies, culinary processing techniques, and foodways.

More broadly, identification of fragmentary material to subfamily and tribe is incredibly useful, given the dietary and habitat requirements of these different bovid groups. Members of Alcelaphinae are grassland-dwelling animals, whereas many species of Cephalophinae are typically found in more forested, closed-canopy environments. Such identifications can provide some insight into past environments, and provide identifications that can inform further analyses, including stable isotope analysis. In some cases, genus-level identifications are possible for wild bovids such as the klipspringer, dikdik, and suni. Such specific identifications provide avenues for assessing deeper questions regarding shifts in climate, environment, as well as hunting and land-use pressures that may have contributed to changes in faunal distributions over time in a variety of African contexts.

Different bovid groups share many peptide markers with identical amino acid sequences, and thus m/z ratios, secure taxonomic identifications can typically only be made on specimens with well-preserved collagen and full spectra. Because size class can sometimes help narrow taxonomic identifications, we recommend that, if available, specimens identifiable to element and size be analyzed for ZooMS. Furthermore, while this study includes species from all bovid tribes, out results point to considerable diversity in peptide markers within the Antilopinae, and some differences in the new markers exist within subfamilies or tribes. Further analyses of bovids within some subfamilies are necessary to clarify these differences.

Our results demonstrate the applicability of ZooMS for the study of even very highly fragmented and poorly preserved assemblages in African archaeological contexts. Using ZooMS, domesticates, including cattle, sheep, and goat, can usually be distinguished from most wild African bovid species, and therefore this method will be extremely useful in investigations of the dispersal of these domestic livestock across the African continent. And though ZooMSbased identifications are restricted to broader taxonomic categories in most cases, the method will still be of great benefit in the analysis of very fragmentary and poorly preserved assemblages, and can provide greater consistency in identifications among analysts. ZooMS certainly does not render zooarchaeology obsolete of course; identification of African bovids through this method instead opens up more possibilities for integrating this method with standard zooarchaeological analyses, allowing for deeper analyses of taxon-specific differences in transport, carcass handling, and culinary processing [2, 144], as well as taphonomic processes, which contribute to major identification biases [145–147]. Recent work elsewhere has shown the usefulness of ZooMS in assessing differential processing strategies [110]. Application of ZooMS on a large scale has great potential for clarifying taxon-specific patterns of preservation.

## Supporting information

**S1 Fig. MALDI-TOF spectra from the bovids.** The following pages contain one example spectra from each of the species used in the study. Each subfamily has a page with grouped spectra and then each individual has a peak picked spectra individually. The first spectra in the grouped spectra for each subfamily is always *Bos taurus* (Bovinae) for easy comparison to a previously published species.

(PDF)

**S2 Fig. MS/MS sequence identification of biomarkers.** The following pages show the MS/MS spectra of the peptides at each of the diagnostic biomarkers presented in the manuscript. All images from Byonic.

(PDF)

S3 Fig. MS/MS sequence identification of the peptides at m/z 3017. In Bovidae the peak at m/z 3017 is composed of two peptides. One peptide is diagnostic between different species (COL1 $\alpha$ 2 757–789 or G). The other peptide is shared between all bovids and therefore non-diagnostic (COL1 $\alpha$ 2 89–130). This is why m/z 3017 should not be used for taxonomic identification. The diagnostic peptide is the only peptide which composes the MALDI peak with a different number of oxidations (m/z 3033) and therefore that is diagnostic. Images from Byonic. (PDF)

**S1 Table. Modern species which were collected for reference data used in this study along with their numbers.** OM numbers are the specimen numbers for the samples we collected, which are housed in the Osteology Section at the National Museums of Kenya in Nairobi. Samples in bold were analyzed by LC-MS/MS. (XLSX)

**S2 Table.** Archaeological data. ZooMS & Morphology ID takes into account morphological indicators, such as body size, to refine the ZooMS-based identification. Site codes: KAV, Kalundu Mound; KMP, SAL, Salumano; ZAM; Jakobo West Kapiri Mposhi; FIB, Fibobe II; MTS, Muteteshi. (XLSX)

**S3 Table. MS/MS information.** Table shows the numbers for the LC-MS/MS samples as well as summary statistics from the runs. (XLSX)

S4 Table. List of all top proteins from the search against SwissProt. A total of 168 proteins

were found in at least one sample and were then added to the collagen sequence database for the marker authentication.

(XLSX)

**S5** Table. Marker list for all species including sequences, masses, and numbers of oxidations. Masses in light italics are present in the LC-MS/MS, but are not suitable for species identification using MALDI because they are present in more than one marker or those masses are also the masses of common contaminants. Boviane—Tragaelaphini—Group1: *Taurotragus*, *Tragelaphus buxtoni*; *Tragelaphus eurycerus*; Boviane—Tragaelaphini—Group2: *Tragelaphus imberis*, *Tragelaphus strepsiceros*, *Tragelaphus scriptus*, *Tragelaphus spekii*. \*Predicted from the genetic sequence data only, not confirmed with MALDI data. \*\*The both the fully cleaved peptide and the peptide with one missed cleavage are present in the LC-MS/MS and the MALDI so are presented here as a joint marker. \*\*\*Previously published species. \*\*\*\*New markers presented in this paper. 1588 for Neotraginae is in the trailing peaks of another peptide at 1586. (XLSX)

S1 File. Information on data accessibility, nomenclature of ZooMS markers, and a note on *Neotragus pygmaeus*.

(PDF)

## Acknowledgments

We would like to thank C. Bochaton for reptile identifications. Archaeological sites were excavated by a joint team from the Max Planck Institute of the Science of Human History, University of Queensland, Livingstone Museum, National Heritage Conservation Commission (NHCC), and Universities of Zambia. Genetic data was generated by the Ruminant Genome Project and assistance was provided by R. Heller, M. Collins, and J.A. Samaniego. Finally, we would like to thank two anonymous reviewers for their thoughtful comments which greatly improved the manuscript.

## **Author Contributions**

Conceptualization: Anneke Janzen, Ogeto Mwebi, Emmanuel Ndiema, Nicole Boivin.

Data curation: Kristine Korzow Richter.

**Formal analysis:** Anneke Janzen, Kristine Korzow Richter, Samantha Brown, Veronicah Onduso, Filia Gatwiri.

Funding acquisition: Anneke Janzen, Katerina Douka.

Investigation: Anneke Janzen, Samantha Brown.

Methodology: Kristine Korzow Richter, Samantha Brown.

Project administration: Katerina Douka, Nicole Boivin.

Resources: Ogeto Mwebi, Emmanuel Ndiema, Maggie Katongo, Steven T. Goldstein.

Supervision: Ogeto Mwebi, Katerina Douka, Nicole Boivin.

Visualization: Anneke Janzen, Kristine Korzow Richter.

Writing – original draft: Anneke Janzen, Kristine Korzow Richter, Ogeto Mwebi, Samantha Brown, Steven T. Goldstein, Nicole Boivin.

#### References

- 1. Brain CK. The Hunters or the Hunted? An Introduction to African Cave Taphonomy. Chicago: The University of Chicago Press; 1981.
- Lupo KD. What explains the carcass field processing and transport decisions of contemporary huntergatherers? Measures of economic anatomy and zooarchaeological skeletal part representation. J Archaeol Method Theory. 2006; 13(1):19–66.
- 3. Marshall FB. Cattle herds and caprine flocks. In: Robertshaw P, editor. Early Pastoralists of Southwestern Kenya. Memoir 11. Nairobi: British Institute in Eastern Africa; 1990. p. 205–60.

- Brain CK. The contribution of Namib Desert Hottentots to an understanding of australopithecine bone accumulations. Scientific Papers of the Namib Desert Research Station. 1969; 1969(39):13–22.
- 5. Marean CW, Spencer LM. Impact of Carnivore Ravaging on Zooarchaeological Measures of Element Abundance. AmerAnt. 1991; 56(4):645–58.
- Marean CW, Spencer LM, Blumenschine RJ, Capaldo SD. Captive hyaena bone choice and destruction, the schlepp effect and Olduvai archaeofaunas. JAS. 1992; 19:101–21.
- 7. Lam YM, Chen X, Marean CW, Frey CJ. Bone density and long bone representation in archaeological faunas: comparing results from CT and photon densitometry. JAS. 1998; 25(6):559–70.
- 8. Lyman RL. Density-mediated attrition of bone assemblages: new insights. From bones to behavior. 1993:324–41.
- 9. Smith AB, Poggenpoel C. The technology of bone tool fabrication in the south-western Cape, South Africa. WorldArch. 1988; 20(1):103–15.
- Badenhorst S, Plug I. Unidentified specimens in zooarcheaology. Palaeontologica Africana. 2011; 46:89–92.
- 11. Gifford DP, Isaac GL, Nelson CM. Prolonged Drift. Azania. 1980; 15:57–108.
- Marean CW, Abe Y, Frey CJ, Randall RC. Zooarchaeological and taphonomic analysis of the Die Kelders Cave 1 Layers 10 and 11 Middle Stone Age larger mammal fauna. J Hum Evol. 2000; 38 (1):197–233. https://doi.org/10.1006/jhev.1999.0356 PMID: 10627403
- 13. Marshall FB, Barthelme JW, Stewart K. Early domestic stock at Dongodien. Azania. 1984; 19:120–7.
- Sadr K, Plug I. Faunal remains in the transition from hunting to herding in southeastern Botswana. The South African Archaeological Bulletin. 2001:76–82.
- Thompson JC, Henshilwood CS. Taphonomic analysis of the Middle Stone Age larger mammal faunal assemblage from Blombos Cave, southern Cape, South Africa. J Hum Evol. 2011; 60(6):746–67. https://doi.org/10.1016/i.jhevol.2011.01.013 PMID: 21470662
- IUCN. The IUCN Red List of Threatened Species. Version 2020–3. <u>https://www.iucnredlist.org</u>. Downloaded on 15 May 2020 2020 [
- ESRI. World Hillshade [basemap]. Scale Not Given. "World Hillshade". Aug 13, 2020. https://www. arcgis.com/home/item.html?id=1b243539f4514b6ba35e7d995890db1d. Accessed 25 November 2020.
- ESRI. World Imagery [basemap]. Scale not given. "World Imagery". 16 December 2020. https://www. arcgis.com/home/item.html?id=10df2279f9684e4a9f6a7f08febac2a9 (23 December 2020).
- NASA. Digital elevation model [basemap]. "NASA Shuttle Radar Topography Mission Global 30 arc second V002". https://cmr.earthdata.nasa.gov/search/concepts/C204582036-LPDAAC\_ECS.html (25 November 2020).
- FAO. 2000 Decadal Rainfall. http://www.fao.org/geonetwork/srv/en/metadata.show?id= 30900&currTab=simple (accessed 25 November 2000)
- Olson DM, Dinerstein E, Wikramanayake ED, Burgess ND, Powell GVN, Underwood EC, et al. Terrestrial ecoregions of the world: a new map of life on Earth. Bioscience. 2001; 51(11):933–8.
- **22.** Ambrose SH. Chronology of the Later Stone Age and food production in East Africa. JAS. 1998; 25:377–92.
- Marean CW. Hunter to herder: large mammal remains from the hunter-gatherer occupation at Enkapune Ya Muto rock shelter, Central Rift, Kenya. The African Archaeological Review. 1992; 10:65–127.
- 24. Pleurdeau D, Imalwa E, Détroit F, Lesur J, Veldman A, Bahain J-J, et al. "Of sheep and men": earliest direct evidence of caprine domestication in southern Africa at Leopard Cave (Erongo, Namibia). PLoS One. 2012; 7(7):e40340. https://doi.org/10.1371/journal.pone.0040340 PMID: 22808138
- 25. Prendergast ME. Kansyore fisher-foragers and transitions to food production in East Africa: the view from Wadh Lang'o, Nyanza Province, Western Kenya. Azania. 2010; 45(1):83–111.
- 26. Horsburgh KA, Moreno-Mayar JV, Gosling AL. Revisiting the Kalahari debate in the highlands: ancient DNA provides new faunal identifications at Sehonghong, Lesotho. Azania: Archaeological Research in Africa. 2016; 51(3):295–306.
- Horsburgh KA, Orton J, Klein RG. Beware the Springbok in Sheep's Clothing: How Secure Are the Faunal Identifications upon Which We Build Our Models? Afr Archaeol Rev. 2016:1–9.
- **28.** Horsburgh KA. A reply to Plug 2017: science requires self-correction. Azania: Archaeological Research in Africa. 2018; 53(1):114–8.
- **29.** Plug I. Reply to Horsburgh et al. 2016: 'Revisiting the Kalahari debate in the highlands.'. Azania: Archaeological Research in Africa. 2018; 53(1):98–113.

- Robbins LH, Campbell AC, Murphy ML, Brook GA, Liang F, Skaggs SA, et al. Recent archaeological research at Toteng, Botswana: early domesticated livestock in the Kalahari. Journal of African Archaeology. 2008; 6(1):131–49.
- Sealy JC, Yates R. Direct radiocarbon dating of early sheep bones: two further results. The South African Archaeological Bulletin. 1996; 51(164):109–10.
- Vogel J, Plug I, Webbley L. New dates for the introduction of sheep into South Africa: the evidence from Spoegrivier Cave in Namaqualand. 1997.
- Coutu AN, Taurozzi AJ, Mackie M, Jensen TZT, Collins MJ, Sealy J. Palaeoproteomics confirm earliest domesticated sheep in southern Africa ca. 2000 BP. Sci Rep. 2021; 11(1):1–11. https://doi.org/10.1038/s41598-020-79139-8 PMID: 33414495
- Carter PL, Flight C. A report on the fauna from the sites of Ntereso and Kintampo Rock Shelter Six in Ghana: with evidence for the practice of animal husbandry during the second millennium b.c. Man. 1972; 7(2):277–82.
- **35.** Holl A. Subsistence patterns of the Dhar Tichitt Neolithic, Mauritania. Afr Archaeol Rev. 1985; 3 (1):151–62.
- Plug I. Hunters and herders: An archaeozoological study of some prehistoric communities in the Kruger National Park. 1990.
- Monroe JC, Janzen A. The Dahomean Feast: Royal Women, Private Politics, and Culinary Practices in Atlantic West Africa. Afr Archaeol Rev. 2014; 31(2):299–337.
- Buckley M, Collins M, Thomas-Oates J, Wilson JC. Species identification by analysis of bone collagen using matrix-assisted laser desorption/ionisation time-of-flight mass spectrometry. Rapid Commun Mass Spectrom. 2009; 23(23):3843–54. https://doi.org/10.1002/rcm.4316 PMID: 19899187
- Charlton S, Alexander M, Collins M, Milner N, Mellars P, O'Connell TC, et al. Finding Britain's last hunter-gatherers: A new biomolecular approach to 'unidentifiable'bone fragments utilising bone collagen. JAS. 2016; 73:55–61.
- Harvey VL, Daugnora L, Buckley M. Species identification of ancient Lithuanian fish remains using collagen fingerprinting. JAS. 2018; 98:102–11.
- Richter KK, McGrath K, Masson-MacLean E, Hickinbotham S, Tedder A, Britton K, et al. What's the catch? Archaeological application of rapid collagen-based species identification for Pacific Salmon. JAS. 2020; 116:105116.
- 42. Welker F, Soressi M, Rendu W, Hublin J-J, Collins M. Using ZooMS to identify fragmentary bone from the late Middle/Early Upper Palaeolithic sequence of Les Cottés, France. JAS. 2015; 54:279–86.
- Hendy J, Welker F, Demarchi B, Speller C, Warinner C, Collins MJ. A guide to ancient protein studies. Nature Ecology & Evolution. 2018; 2(5):791–9.
- Prendergast ME, Buckley M, Crowther A, Frantz L, Eager H, Lebrasseur O, et al. Reconstructing Asian faunal introductions to eastern Africa from multi-proxy biomolecular and archaeological datasets. PLoS One. 2017; 12(8):e0182565. <u>https://doi.org/10.1371/journal.pone.0182565</u> PMID: 28817590
- **45.** Prendergast ME, Janzen A, Buckley M, Grillo KM. Sorting the sheep from the goats in the Pastoral Neolithic: morphological and biomolecular approaches at Luxmanda, Tanzania. Archaeological and Anthropological Sciences. 2019; 11(6):3047–62.
- 46. Coutu AN, Whitelaw G, le Roux P, Sealy JC. Earliest evidence for the ivory trade in southern Africa: Isotopic and ZooMS analysis of seventh–tenth century AD ivory from KwaZulu-Natal. Afr Archaeol Rev. 2016; 33(4):411–35.
- Bradfield J, Forssman T, Spindler L, Antonites AR. Identifying the animal species used to manufacture bone arrowheads in South Africa. Archaeological and Anthropological Sciences. 2019; 11(6):2419– 34.
- Desmond A, Barton N, Bouzouggar A, Douka K, Fernandez P, Humphrey L, et al. ZooMS identification of bone tools from the North African Later Stone Age. JAS. 2018; 98:149–57.
- Le Meillour L, Zirah S, Zazzo A, Cersoy S, Détroit F, Imalwa E, et al. Palaeoproteomics gives new insight into early southern African pastoralism. Sci Rep. 2020; 10(1):1–11. https://doi.org/10.1038/ s41598-019-56847-4 PMID: 31913322
- Stanley-Price MR. The nutritional ecology of Coke's hartebeest (Alcelaphus buselaphus cokei) in Kenya. J Appl Ecol. 1978:33–49.
- 51. Stewart D. Food preferences of an impala herd. The Journal of wildlife management. 1971:86–93.
- 52. Kingdon J. East African Mammals. Chicago: University of Chicago Press; 1984.
- 53. Cerling TE, Harris JM, Passey BH. Diets of East African Bovidae based on stable isotope analysis. Journal of Mammology. 2003; 84(2):456–70.

- Sponheimer M, Lee-Thorp JA, DeRuiter DJ, Smith J, Van Der Merwe NJ, Reed K, et al. Diets of southern African Bovidae: stable isotope evidence. J Mammal. 2003; 84(2):471–9.
- Gagnon M, Chew AE. Dietary preferences in extant African Bovidae. J Mammal. 2000; 81(2):490– 511.
- **56.** Vrba ES. 14. The significance of bovid remains as indicators of environment and predation patterns: University of Chicago Press, Chicago; 1980.
- 57. Faith JT. Ungulate community richness, grazer extinctions, and human subsistence behavior in southern Africa's Cape Floral Region. Palaeogeogr, Palaeoclimatol, Palaeoecol. 2011; 306(3–4):219–27.
- Gifford-Gonzalez D. The fauna from Ele Bor: evidence for the persistence of foragers into the later Holocene of arid north Kenya. Afr Archaeol Rev. 2003; 20(2):81–119.
- Peters J. Late Quaternary Mammalian remains from Central and Eastern Sudan and their palaeoenvironment significance. 1992.
- Jousse H, Escarguel G. The use of Holocene bovid fossils to infer palaeoenvironment in Africa. QSRv. 2006; 25(7–8):763–83.
- Roberts P, Prendergast ME, Janzen A, Shipton C, Blinkhorn J, Zech J, et al. Late Pleistocene to Holocene human palaeoecology in the tropical environments of coastal eastern Africa. Palaeogeogr, Palaeoclimatol, Palaeoecol. 2020; 537:109438.
- Marean CW. Implications of late Quaternary mammalian fauna from Lukenya Hill (south-central Kenya) for paleoenvironmental change and faunal extinctions. QuRes. 1992; 37(2):239–55.
- 63. Tryon CA, Faith JT, Peppe DJ, Beverly EJ, Blegen N, Blumenthal SA, et al. The Pleistocene prehistory of the Lake Victoria basin. Quat Int. 2016; 404:100–14.
- DeGusta D, Vrba E. A method for inferring paleohabitats from the functional morphology of bovid astragali. JAS. 2003; 30(8):1009–22.
- Plummer TW, Bishop LC, Hertel F. Habitat preference of extant African bovids based on astragalus morphology: operationalizing ecomorphology for palaeoenvironmental reconstruction. JAS. 2008; 35 (11):3016–27.
- Blumenthal SA, Levin NE, Brown FH, Brugal J-P, Chritz KL, Harris JM, et al. Aridity and hominin environments. Proceedings of the National Academy of Sciences. 2017; 114(28):7331–6. https://doi.org/ 10.1073/pnas.1700597114 PMID: 28652366
- 67. Chritz KL, Cerling TE, Freeman KH, Hildebrand EA, Janzen A, Prendergast ME. Climate, ecology, and the spread of herding in eastern Africa. QSRv. 2019; 204:119–32.
- Lehmann SB, Braun DR, Dennis KJ, Patterson DB, Stynder DD, Bishop LC, et al. Stable isotopic composition of fossil mammal teeth and environmental change in southwestern South Africa during the Pliocene and Pleistocene. Palaeogeogr, Palaeoclimatol, Palaeoecol. 2016; 457:396–408.
- Robinson JR, Rowan J. Holocene paleoenvironmental change in southeastern Africa (Makwe Rockshelter, Zambia): Implications for the spread of pastoralism. QSRv. 2017; 156:57–68.
- Dewar G, Halkett D, Hart T, Orton J, Sealy JC. Implications of a mass kill site of springbok (Antidorcas marsupialis) in South Africa: hunting practices, gender relations, and sharing in the Later Stone Age. JAS. 2006; 33(9):1266–75.
- Jenkins KE, Nightingale S, Faith JT, Peppe DJ, Michel LA, Driese SG, et al. Evaluating the potential for tactical hunting in the Middle Stone Age: Insights from a bonebed of the extinct bovid, Rusingoryx atopocranion. J Hum Evol. 2017; 108:72–91. <u>https://doi.org/10.1016/j.jhevol.2016.11.004</u> PMID: 28622933
- Jones MB, Brandt SA, Marshall F. Hunter-gatherer reliance on inselbergs, big game, and dwarf antelope at the Rifle Range Site, Buur Hakaba, southern Somalia ~20,000–5,000 BP. Quat Int. 2017; 30:1e11.
- Lupo KD, Schmitt DN. Small prey hunting technology and zooarchaeological measures of taxonomic diversity and abundance: ethnoarchaeological evidence from Central African forest foragers. JAnthArch. 2005; 24(4):335–53.
- 74. Plug I, Mitchel P. Sehonghong: hunter-gatherer utilization of animal resources in the highlands of Lesotho. Ann Transvaal Mus. 2008; 45(1):31–53.
- 75. Hanotte O, Bradley DG, Ochieng JW, Verjee Y, Hill EW, Rege JEO. African pastoralism: Genetic imprints of origins and migrations. Sci. 2002; 296(5566):336–9. https://doi.org/10.1126/science. 1069878 PMID: 11951043
- 76. Achilli A, Bonfiglio S, Olivieri A, Malusa A, Pala M, Kashani BH, et al. The multifaceted origin of taurine cattle reflected by the mitochondrial genome. PLoS One. 2009; 4(6):e5753. <u>https://doi.org/10.1371/journal.pone.0005753</u> PMID: 19484124

- Bradley DG, MacHugh DE, Cunningham P, Loftus R. Mitochondrial diversity and the origins of African and European cattle. Proceedings National Academy of Science. 1996; 93:5131–5.
- Stock F, Gifford-Gonzalez D. Genetics and African cattle domestication. Afr Archaeol Rev. 2013; 30 (1):51–72.
- 79. Naderi S, Rezaei H-R, Pompanon F, Blum MG, Negrini R, Naghash H-R, et al. The goat domestication process inferred from large-scale mitochondrial DNA analysis of wild and domestic individuals. Proceedings of the National Academy of Sciences. 2008; 105(46):17659–64. <u>https://doi.org/10.1073/pnas.0804782105</u> PMID: 19004765
- Zeder MA, Hesse B. The initial domestication of goats (*Capra hircus*) in the Zagros mountains 10,000 years ago. Sci. 2000; 287:2254–7. https://doi.org/10.1126/science.287.5461.2254 PMID: 10731145
- Daly KG, Delser PM, Mullin VE, Scheu A, Mattiangeli V, Teasdale MD, et al. Ancient goat genomes reveal mosaic domestication in the Fertile Crescent. Sci. 2018; 361(6397):85–8. https://doi.org/10. 1126/science.aas9411 PMID: 29976826
- Bruford MW, Townsend SJ. Mitochondrial DNA diversity in modern sheep: Implications for domestication. In: Zeder MA, Bradley DG, Emshwiller E, Smith BD, editors. Documenting Domestication: New Genetic and Archaeological Paradigms. Berkeley: University of California Press; 2006. p. 306–16.
- Peters J. Osteomorphological features of the appendicular skeleton of African buffalo, Syncerus caffer (Sparrman, 1779) and of domestic cattle, Bos primigenius f. taurus Bojanus, 1827. Zeitschrift für Säugetierkunde. 1988:108–23.
- Boessneck J. Osteological differences between sheep (Ovis aries Linné) and goat (Capra hircus Linné): Thames and Hudson; 1971.
- Halstead P, Collins P, Isaakidou V. Sorting the sheep from the goats: Morphological distinctions between the mandibles and mandibular teeth of adult *Ovis* and *Capra*. JAS. 2002; 29:545–53.
- Zeder MA, Lapham HA. Assessing the reliability of criteria used to identify postcranial bones in sheep, Ovis, and goats, Capra. JAS. 2010; 37(11):2887–905.
- Zeder MA, Pilaar SE. Assessing the reliability of criteria used to identify mandibles and mandibular teeth in sheep, *Ovis*, and goats, *Capra*. JAS. 2010; 37 (2):225–42.
- Janzen A, Balasse M, Ambrose SH. Early Pastoral Mobility in Kenya Assessed through Stable Isotope Analysis. JAS. 2020; 117.
- Arnold ER, Greenfield HJ, Creaser RA. Domestic cattle mobility in early farming villages in southern Africa: harvest profiles and strontium (87Sr/86Sr) isotope analyses from Early Iron Age sites in the lower Thukela River Valley of South Africa. Archaeological and Anthropological Sciences. 2013; 5 (2):129–44.
- **90.** Balasse M, Ambrose SH, Smith AB, Price TD. The seasonal mobility model for prehistoric herders in the south-western Cape of South Africa assessed by isotopic analysis of sheep tooth enamel. JAS. 2002; 29:917–32.
- Balasse M, Smith AB, Ambrose SH, Leigh SR. Determining sheep birth seasonality by analysis of tooth enamel oxygen isotope ratios: The Late Stone Age site of Kasteelberg (South Africa). JAS. 2003; 30(2):205–15.
- Plug I, Badenhorst S. Notes on the fauna from three Late Iron Age mega-sites, Boitsemagano, Molokwane and Mabjanamatshwana, North West Province, South Africa. The South African Archaeological Bulletin. 2006:57–67.
- **93.** Stone AC. Finding the ephemeral: Herding strategies and socio-economic organization in an urban West African context. Quat Int. 2018; 471:160–74.
- 94. Haruda A, Varfolomeev V, Goriachev A, Yermolayeva A, Outram A. A new zooarchaeological application for geometric morphometric methods: distinguishing Ovis aries morphotypes to address connectivity and mobility of prehistoric Central Asian pastoralists. JAS. 2019; 107:50–7.
- 95. Fraser L-M, Badenhorst S. Livestock use in the Limpopo Valley of southern Africa during the Iron Age. The South African Archaeological Bulletin. 2014:192–8.
- 96. Sadr K. Livestock first reached southern Africa in two separate events. PLoS One. 2015; 10(8): e0134215. https://doi.org/10.1371/journal.pone.0134215 PMID: 26295347
- 97. Marshall FB, Stewart K. Hunting, fishing and herding pastoralists of western Kenya: The fauna from Gogo Falls. Archaeozoologia. 1995; 7:7–27.
- Fagan BM, Phillipson D, Daniels S. Iron Age Cultures in Zambia: Kalomo and Kangila: Chatto & Windus; 1967.
- 99. Grody E. Of buffalo and butchers: Coupling traditional procurement studies with taphonomic analyses to explore intensive wild animal processing patterns at two Early Iron Age sites in the Kruger National Park. Afr Archaeol Rev. 2016; 33(4):385–409.

- 100. Phillipson DW. The prehistory of eastern Zambia: British Institute in Eastern Africa; 1976.
- 101. Katanekwa N. Namakala and Nanga Sites and the Chronology of the Early Iron Age in Southern & South-West Zambia. The South African Archaeological Bulletin. 1979:120–2.
- 102. Dueppen SA, Gokee C. Hunting on the margins of medieval West African states: a preliminary study of the zooarchaeological record at Diouboye, Senegal. Azania: Archaeological Research in Africa. 2014; 49(3):354–85.
- **103.** Linseele V, Van Neer W, Friedman R. Special animals from a special place? The fauna from HK29A at Predynastic Hierakonpolis. JARCE. 2009:105–36.
- 104. Brunton S, Badenhorst S, Schoeman MH. Ritual fauna from Ratho Kroonkop: a second millennium AD rain control site in the Shashe-Limpopo Confluence area of South Africa. Azania: Archaeological Research in Africa. 2013; 48(1):111–32.
- 105. Le Meillour L, Zazzo A, Lesur J, Cersoy S, Marie A, Lebon M, et al. Identification of degraded bone and tooth splinters from arid environments using palaeoproteomics. Palaeogeogr, Palaeoclimatol, Palaeoecol. 2018; 511:472–82.
- Brandt LØ, Haase K, Collins MJ. Species identification using ZooMS, with reference to the exploitation of animal resources in the medieval town of Odense. Danish Journal of Archaeology. 2018; 7(2):139–53.
- 107. McGrath K, Rowsell K, St-Pierre CG, Tedder A, Foody G, Roberts C, et al. identifying Archaeological Bone via non-Destructive ZooMS and the Materiality of Symbolic expression: examples from iroquoian Bone points. Sci Rep. 2019; 9(1):1–10. https://doi.org/10.1038/s41598-018-37186-2 PMID: 30626917
- **108.** Birch SEP, Scheu A, Buckley M, Çakırlar C. Combined osteomorphological, isotopic, aDNA, and ZooMS analyses of sheep and goat remains from Neolithic Ulucak, Turkey. Archaeological and Anthropological Sciences. 2019; 11(5):1669–81.
- **109.** Pothier Bouchard G, Riel-Salvatore J, Negrino F, Buckley M. Archaeozoological, taphonomic and ZooMS insights into The Protoaurignacian faunal record from Riparo Bombrini. Quat Int. 2020.
- Sinet-Mathiot V, Smith GM, Romandini M, Wilcke A, Peresani M, Hublin J-J, et al. Combining ZooMS and zooarchaeology to study Late pleistocene hominin behaviour at Fumane (Italy). Sci Rep. 2019; 9 (1):1–13. https://doi.org/10.1038/s41598-018-37186-2 PMID: 30626917
- 111. Buckley M, Kansa SW, Howard S, Campbell S, Thomas-Oates J, Collins M. Distinguishing between archaeological sheep and goat bones using a single collagen peptide. JAS. 2010; 37(1):13–20.
- 112. Welker F, Hajdinjak M, Talamo S, Jaouen K, Dannemann M, David F, et al. Palaeoproteomic evidence identifies archaic hominins associated with the Châtelperronian at the Grotte du Renne. PNAS. 2016; 113(40):11162–7. https://doi.org/10.1073/pnas.1605834113 PMID: 27638212
- 113. Pate FD. Bone collagen diagenesis at Roonka Flat, South Australia: Implications for isotopic analysis. Archaeol Oceania. 1997; 32(2):170–5.
- Clark JD, Fagan BM. Charcoals, sands, and channel decorated pottery from Northern Rhodesia. Amer Anthrop. 1965; 67(2):354–71.
- 115. Katanekwa N. Some Early Iron Age sites from the Machili valley of south western Zambia. AZANIA: Journal of the British Institute in Eastern Africa. 1978; 13(1):135–66.
- **116.** Plug I. The fauna from Salumano, an Early Iron Age site in Zambia. Archaeologia Zambiana. 1981; 20:20–2.
- 117. Robertson JH. Early iron age archaeology in central Zambia. Azania. 2000; 35(1):147-82.
- **118.** Musonda FB. The significance of pottery in Zambian Later Stone Age contexts. Afr Archaeol Rev. 1987; 5(1):147–58.
- **119.** Inskeep RR. Some Iron Age Sites in Northern Rhodesia. The South African Archaeological Bulletin. 1962; 17(67):136–80.
- **120.** Goldstein ST, Crowther A, Henry ER, Janzen A, Katongo M, Brown S, et al. Revisiting Kalundu Mound, Zambia: Implications for the timing of social and subsistence transitions in Iron Age southern Africa. Afr Archaeol Rev. in review:Forthcoming.
- 121. Brown S, Hebestreit S, Wang N, Boivin N, Douka K, Richter KK. Zooarchaeology by Mass Spectrometry (ZooMS)- Pretreatment protocols for bone material v1. protocolsio. 2020.
- 122. Strohalm M, Hassman M, Košata B, Kodíček M. mMass data miner: an open source alternative for mass spectrometric data analysis. Rapid Commun Mass Spectrom. 2008; 22(6):905–8. <u>https://doi.org/10.1002/rcm.3444</u> PMID: 18293430
- Bern M, Kil YJ, Becker C. Byonic: advanced peptide and protein identification software. Current protocols in bioinformatics. 2012; 40(1):13–20. https://doi.org/10.1002/0471250953.bi1320s40 PMID: 23255153
- 124. Chen L, Qiu Q, Jiang Y, Wang K, Lin Z, Li Z, et al. Large-scale ruminant genome sequencing provides insights into their evolution and distinct traits. Sci. 2019; 364(6446):eaav6202. https://doi.org/10.1126/ science.aav6202 PMID: 31221828

- 125. Gasteiger E, Hoogland C, Gattiker A, Wilkins MR, Appel RD, Bairoch A. Protein identification and analysis tools on the ExPASy server. The proteomics protocols handbook: Springer; 2005. p. 571–607.
- 126. Wilkins MR, Lindskog I, Gasteiger E, Bairoch A, Sanchez JC, Hochstrasser DF, et al. Detailed peptide characterization using PEPTIDEMASS–a World-Wide-Web-accessible tool. Electrophoresis. 1997; 18 (3-4):403–8. https://doi.org/10.1002/elps.1150180314 PMID: 9150918
- 127. Bärmann EV, Rössner GE, Wörheide G. A revised phylogeny of Antilopini (Bovidae, Artiodactyla) using combined mitochondrial and nuclear genes. Mol Phylogen Evol. 2013; 67(2):484–93. <u>https://doi.org/10.1016/j.ympev.2013.02.015</u> PMID: 23485920
- 128. Fernández MH, Vrba ES. A complete estimate of the phylogenetic relationships in Ruminantia: a dated species-level supertree of the extant ruminants. Biological reviews. 2005; 80(2):269–302. https://doi.org/10.1017/s1464793104006670 PMID: 15921052
- 129. Gatesy J, Amato G, Vrba E, Schaller G, DeSalle R. A Cladistic Analysis of Mitochondrial Ribosomal DNA from the Bovidae. Mol Phylogen Evol. 1997; 7(3):303–19. https://doi.org/10.1006/mpev.1997. 0402 PMID: 9187090
- Hassanin A, Douzery EJP. The Tribal Radiation of the Family Bovidae (Artiodactyla) and the Evolution of the Mitochondrial Cytochrome b Gene. Mol Phylogen Evol. 1999; 13(2):227–43.
- 131. Johnston AR, Anthony NM. A multi-locus species phylogeny of African forest duikers in the subfamily Cephalophinae: evidence for a recent radiation in the Pleistocene. BMC Evol Biol. 2012; 12(1):120. https://doi.org/10.1186/1471-2148-12-120 PMID: 22823504
- Matthee CA, Robinson TJ. Cytochrome b phylogeny of the family Bovidae: Resolution within the Alcelaphini, Antilopini, Neotragini, and Tragelaphini. Mol Phylogen Evol. 1999; 12(1):31–46.
- **133.** Klinken GJv. Dating aand dietary reconstruction by isotopic analysis of amino acids in fossil bone collagen-with special reference to the caribbean: University of Groningen; 1991.
- 134. Trueman CN, Behrensmeyer AK, Tuross N, Weiner S. Mineralogical and compositional changes in bones exposed on soil surfaces in Amboseli National Park, Kenya: diagenetic mechanisms and the role of sediment pore fluids. JAS. 2004; 31(6):721–39.
- Roberts SJ, Smith CI, Millard A, Collins MJ. The taphonomy of cooked bone: characterizing boiling and its physico-chemical effects. Archa. 2002; 44(3):485–94.
- Naihui W, Samantha B, Peter D, Sandra H, Maxim K, Sindy L, et al. Testing the efficacy and comparability of ZooMS protocols on archaeological bone. J Proteomics. 2020:104078. <u>https://doi.org/10.1016/j.jprot.2020.104078 PMID: 33338688</u>
- Plug I. Nanga: an Early Iron Age cattle herding community in the Machili Valley, Zambia. Archaeologia Zambiana. 1981; 20:17–9.
- **138.** Plug I. Namakala and Nanga: faunal report on two Early Iron Age sites, Zambia. The South African Archaeological Bulletin. 1979:123–6.
- **139.** Ford J, Katondo K. Maps of tsetse fly (Glossina) distribution in Africa, 1973 according to sub-generic groups on scale of 1: 5,000,000. Bull Anim Hlth Prod Afr. 1977; 25(2):188–94.
- 140. Connor RJ. African animal trypanosomiases. In: Coertzer JAW, Thomson GR, Tustun RC, Kriek NPJ, editors. Infectious Diseases of Livestock with Special Reference to Southern Africa. Vol. 1. New York: Oxford University Press; 1994. p. 167–205.
- 141. Lamprey R, Waller R. The Loita-Mara region in historical times: patterns of subsistence, settlement and ecological change In: Robertshaw P, editor. Early Pastoralists of South-western Kenya. Nairobi: British Institute in Eastern Africa; 1990. p. 16–35.
- 142. Pratt DJ, Gwynne MD. Rangeland Management and Ecology in East Africa. London: Hodder & Stoughton; 1977.
- Wilson V, Roth H. The effects of tsetse control operations on common duiker in Eastern Zambia. Afr J Ecol. 1967; 5(1):53–64.
- O'Connell JF, Hawkes K, Jones NB. Hadza hunting, butchering, and bone transport and their archaeological implications. J Anthrop Res. 1988; 44(2):113–61.
- Faith JT, Behrensmeyer AK. Changing patterns of carnivore modification in a landscape bone assemblage, Amboseli Park, Kenya. JAS. 2006; 33(12):1718–33.
- **146.** Marean CW, Domínguez Rodrigo M, Pickering TR. Skeletal element equifinality in zooarchaeology begins with method: the evolution and status of the "shaft critique". J Taphon. 2004; 2(1):69–98.
- 147. Yeshurun R, Marom N, Bar-Oz G. Differential Fragmentation of Different Ungulate Body-Size: A Comparison of Gazelle and Fallow Deer Bone Fragmentation in Levantine Prehistoric Assemblages. J Taphon. 2007; 5(3):137–48.