



Strange Seal Sounds: Claps, Slaps, and Multimodal Pinniped Rhythms

Laura Verga^{1,2} and Andrea Ravnani^{1*}

¹ Comparative Bioacoustics Group, Max Planck Institute for Psycholinguistics, Nijmegen, Netherlands, ² Department NP&PP, Faculty of Psychology and Neuroscience, Maastricht University, Maastricht, Netherlands

Keywords: multimodal communication, pinniped, seal (*Pinnipedia*), bioacoustics, exaptation, honest signalling

BEYOND VOCAL COMMUNICATION

Within mammalian bioacoustics, vocal communication has received much attention. Efforts to understand sound production often focus on sounds generated *via* apparatuses that specifically evolved to phonate, such as the larynx. However, while mammals mostly perceive sounds *via* one organ, the ear, they can produce sounds *via* limbs, tails, flippers, tools, and several other mechanisms which, at first, may not seem to have primarily evolved for sonation (Tyack and Miller, 2002; Frankel, 2009; Clark, 2016). For example, kangaroo rats drum their foot to communicate (Randall, 1984), while non-human primates drum using artificial tools (Remedios et al., 2009), resonant surfaces (Ravnani et al., 2013), and their hands (e.g., Dufour et al., 2015). Also aquatic mammals can produce a variety of non-vocal sounds (such as whistles, snorts, and others; Tyack and Miller, 2002). These sound production modes may enable communication even when laryngeal phonation is ineffective or impaired (Munoz and Blumstein, 2012; Partan, 2017). Research on sound production *beyond phonation* is key to properly characterise the richness of animal communication.

Recent exploratory work (Hocking et al., 2020) provides an example of non-vocal sound production in a pinniped: grey seals (*Halichoerus grypus*) clapping their fore flippers underwater, a signalling behaviour previously attributed to vocalising. While Hocking et al.'s observation is limited to few events, it is reminiscent of previous, seemingly unrelated work reporting water-slapping behaviour in other species, including a close relative, the harbour seal (*Phoca vitulina*) (Venables and Venables, 1957; Newby, 1973; Hanlan, 1998; Hayes et al., 2004; see also humpback whales: Dunlop et al., 2010). The preliminary data reported by Hocking et al. naturally invite a host of questions, whose answers rely on a characterisation of when, how often, and under what circumstances these claps occur. Are they a frequent or seasonal phenomenon? Are they modulated by social context? Furthermore, the mechanism of knock production should be considered: while clapping the fore flippers can generate loud knock-like sounds, other mechanisms have also been proposed (e.g., in walrus: teeth clacking, tongue movement, or suction; Sjare and Stirling, 1981; Sjare et al., 2003; Reichmuth et al., 2009; Larsen and Reichmuth, 2012). This is a good starting place for future research, though, clearly, more observations are required to answer these questions. In particular, the field should perform more empirical, foundational work. This should (1) provide robust observations and descriptions in addition to anecdotes, (2) evaluate context, timing and seasonality in the production of percussive sounds, (3) determine which sex produces these signals and potential sexually dimorphic characteristics, and (4) design rigorous experiments that test potential function of percussive sounds. Once these absolutely necessary foundations are established, we suggest exploring more complex topics related to bioenergetics, signal evolution, multimodality, and rhythm production/perception. In this Opinion piece, we discuss these more hypothetical research directions, which however can only be performed after more thorough biological descriptions of the basic phenomena.

OPEN ACCESS

Edited by:

Carl Soulsbury,
University of Lincoln, United Kingdom

Reviewed by:

John Terhune,
University of New Brunswick Saint
John, Canada
Caroline Casey,
University of California Santa Cruz,
United States

*Correspondence:

Andrea Ravnani
andrea.ravnani@mpi.nl

Specialty section:

This article was submitted to
Behavioral and Evolutionary Ecology,
a section of the journal
Frontiers in Ecology and Evolution

Received: 21 December 2020

Accepted: 02 July 2021

Published: 02 August 2021

Citation:

Verga L and Ravnani A (2021)
Strange Seal Sounds: Claps, Slaps,
and Multimodal Pinniped Rhythms.
Front. Ecol. Evol. 9:644497.
doi: 10.3389/fevo.2021.644497

MECHANISMS FOR (DIS)HONEST SIGNALLING?

If replicated, this research may encourage to re-evaluate previous evidence. Could previously recorded “vocalisations [which] sounded like a loud piercing clap” (pg. 61, McCulloch, 2000) and “knocks” (pg. 2213, Asselin et al., 1993) in grey seals have been actual underwater claps? A hypothetical reassessment of the production mechanism underlying a sound would entail several implications. Awareness of the sound source could be helpful to test potential sound-body allometric links.

One testable hypothesis is that claps may be a partly dishonest signal as they give away limited information about body size while their source level is surprisingly high, especially compared to captive individuals (Wahlberg et al., 2002). This hypothesis dovetails with some empirical evidence of water claps functioning as aggressive and territorial behaviours in harbour seals (Hayes et al., 2004), since claps have been so far observed mostly in males and in presence of other seals (for example in Weddell seals: Russell et al., 2016). Furthermore, limited underwater visibility, as reported by Hocking et al. (2020), would promote a signalling strategy concealing body size. To test whether clapping is a dishonest signal, field studies should investigate the conditions under which underwater claps take place (e.g., water visibility, social context).

Conversely, claps could be honest signals, since the strength of the animal or the size of its flipper may determine the intensity of the clap and the perceived loudness. This second hypothesis would comply with allometric scaling (e.g. Garcia et al., 2017): if sound-producing structures scale with body size, honest signalling ensues (Garcia and Ravignani, 2020). Larger individuals should also be able to produce stronger or (visually) larger claps (Partan, 2013). Anatomical observations in, and tests of allometry across, individuals producing underwater claps might contribute to disentangling these contrasting hypotheses.

EXAPTATION AND REPURPOSING OF BIOMECHANICAL PROCESSES

This research underlines the process of evolutionary exaptations for communicative purposes (Gould and Lewontin, 1979; Buss et al., 1998). Seal clapping “may be a ritualised version of a swimming stroke” (pg. 1, Hocking et al., 2020). In other words, a movement evolved for essential in-water displacement (Fish, 2000; Kuhn and Frey, 2012) may have been repurposed for acoustic communication (see also Clark, 2016). As a parallel, the tree drumming of woodpeckers is now a purely communicative signal, probably repurposed from what originally was a simpler foraging behaviour (Dodenhoff et al., 2001; Miles et al., 2018, 2020; Garcia et al., 2020). Both seals and woodpeckers may provide fascinating examples of the evolution and repurposing of biomechanical processes. At what point in pinniped phylogeny may swimming strokes have been recruited for communication, and did this happen in

multiple pinniped species? Communicative non-vocal sounds have been observed in several pinnipeds (Schusterman and Van Parijs, 2003; Russell et al., 2016¹); more specifically, claps have been reported, to our knowledge, in grey seals (e.g., Hocking et al., 2020), harbour seals (Venables and Venables, 1957; Newby, 1973; Hanlan, 1998; Hayes et al., 2004), and walruses (e.g., Reichmuth et al., 2009). Underwater recordings of similar non-vocal sounds in other pinnipeds might contribute to answering this question. With limited evidence it is difficult and unwise to generalise to all pinnipeds. However, one may hypothesise that the swimming style of phocids, propelled by their hindflippers, may free up their foreflippers for communicative purposes, while the swimming style of otariids, using their foreflippers to “fly” underwater, may have hindered their exaptation for communication (Kuhn and Frey, 2012).

MULTIMODALITY AND ENERGETICS

Most work mentioned above advocates multimodal approaches to communication. Multimodality is sometimes neglected, with some research programs only focusing on one production and one perception channel (Slocombe et al., 2011). Grey seals exhibit a communicative behaviour that is motorically produced (bypassing specialised laryngeal neurons) and might be perceived acoustically at long ranges and visually at short ranges (Wahlberg et al., 2002; Ravignani et al., 2016; Hocking et al., 2020). As the larynx is hidden from sight, the act of mammalian vocalisation is invisible to the receiver (cf. Fitch and Reby, 2001; Higham and Hebets, 2013; Nowak, 2020). A clap, instead, could potentially reach the receiver visually, acoustically, or haptically, also allowing for multisensory integration.

Multimodality also entails energetic considerations. While laryngeal phonation is relatively cheap, other modes of controlled sound production may be more energy-expensive. Indeed, research on the energetic costs of communication generally assumes higher expenditure for multi-modal as compared to uni-modal interactions (Partan, 2013). For example, in sympatric wolf spiders, multi-modal displays (as in *Schizocosa ocreata*) require higher energy levels than unimodal displays (as in *S. rovneri*) (Cady et al., 2011). Rather than being disadvantageous, such a costly display might serve as an honest signal indicating a male's good condition (Zahavi, 1975; Byers et al., 2010; Mitoyen et al., 2019). Yet, questions concerning the relative costs and benefits of pinniped sound production *via* clapping and slapping (Beier and Wartzok, 1979; Wahlberg et al., 2002; Gillooly and Ophir, 2010) remain open, as well as their function (e.g., territorial, reproductive, etc.; Russell et al., 2016). If they indeed relate to mating, one may expect a variation in clapping/slapping abilities due to ontogeny (Rado et al., 1991), an increase at puberty onset, a cyclical variation entrained with mating seasonality or a decline due to senescence (Soulsbury and

¹We are here referring to jaw claps, which the authors originally grouped as vocalisations for methodological reasons.

Halsey, 2018). The reliable evidence of such seasonal and developmental trends in the vocal displays of seals (e.g., Van Parijs et al., 1999; Galimberti et al., 2008; Reichmuth and Schusterman, 2009) may invite comparisons with non-vocal displays and facilitate the understanding of their function. By combining quantitative techniques (e.g., Gillooly and Ophir, 2010), allometric considerations (Garcia and Ravignani, 2020), and field observations (Hocking et al., 2020), some of these questions may be addressed.

COMMUNICATIVE RHYTHMS IN THE MILLISECOND-SECOND RANGE

More observations on the clapping behaviour of grey seals, their characteristics, and context of use are needed: Are these sounds produced occasionally or routinely? Do they contain rhythmic components? The presence of rhythmic features in claps would allow to link Hocking et al.'s (2020) finding to research on communicative rhythms and could spur a subfield of ecologically-relevant percussive rhythms in mammals. Recently, cross-species evidence has shown rhythmic capacities, sometimes employed for communication, in pinnipeds (Cook et al., 2013; Rouse et al., 2016; Mathevon et al., 2017; Ravignani, 2019). "Rhythm" is not meant here in its circadian sense, studied for instance in ecology, but instead as "temporal structure" at short timescales (de Reus et al., 2020). When little information is encoded in the frequency domain, as in seals' claps and slaps (Wahlberg et al., 2002; Hocking et al., 2020), this temporal structure could emerge in sound signals and serve to encode information. Within animal cognition and behaviour, evidence for rhythm in pinnipeds is particularly interesting (Ravignani et al., 2016; Wilson and Cook, 2016). In fact, pinnipeds constitute a key taxon to test a cross-species hypothesis which links rhythm and vocal learning capacities (Patel, 2006). Still, work on rhythm in mammals is relatively limited, especially when compared to the richness of rhythm production research, for instance, in insects and frogs (Greenfield, 1994; Hartbauer and Römer, 2016). While these species can produce extremely fast rhythms, the rate and complexity of non-vocal rhythms in mammals may be hampered by the physical limitations occurring when moving a limb (but see, for example, Randall, 1984 on the foot drumming behaviour of kangaroo rats). Mammalogists and comparative psychologists may still be inspired and benefit from decades of work on rhythm and percussive behaviour in arthropods and anurans (Ravignani et al., 2014). In particular, since the 1960's, entomologists and then herpetologists have been measuring the communicative rhythms of their species with almost millisecond accuracy (e.g., Buck and Buck, 1968). Avian researchers followed, while mammalogists and primatologists are slightly lagging behind (de Reus et al., 2020). Applying concepts such as phase resetting and period correction, for instance, to communicative rhythms in apes could also inform the evolution of rhythmic capacities in our own species (cf. Bittman, 2020).

CONCLUSIONS AND FUTURE WORK

Hocking et al.'s (2020) finding, albeit preliminary, can inspire at least six hypothetical strands of future work. First, to establish a base from psychophysics, *propagation* experiments could test how far the sound of grey seals clapping carries underwater (Wahlberg et al., 2002); this strand of research would help disentangle the role of claps as either honest or dishonest signals, as the latter may be more relevant at short distances (e.g., Tyack and Miller, 2002). Second, *biomechanics* and metabolic work could pinpoint the energetic costs and evolutionary benefits of clapping (Fish, 2000; Kuhn and Frey, 2012); this research should consider the context in which a signalling behaviour occurs (e.g., occasional vs. prolonged use; environmental and social conditions). Third, a larger dataset (McCulloch, 2000) would allow onset-to-onset temporal measurements to investigate whether claps may feature putative *rhythmic structures*, linking either claps within a series (e.g., based on their inter-onset intervals) or repeated series performed in succession. After that, to test for homologies and analogies, comparative analyses could be attempted with water slaps in harbour seals and other pinniped percussive behaviour (Wahlberg et al., 2002). Fourth, it would be important to test how nearby conspecifics perceive claps. Therefore, connecting to recent work on pinniped timing (Heinrich et al., 2016, 2020), one could explore how grey seals *perceive temporal information* in sequences of claps. Fifth, one could target the proximate and ultimate *function of clapping*, and its potential role in sexual or natural selection. Sixth, methodological advances in neuroimaging techniques (e.g., Cook et al., 2021) may be employed to inform on the neural underpinnings of cross-modality and their interface with the physiological and physical constraints imposed on flippers by their original function (i.e., swimming). All this work, we stress, can only come once more fundamental research is performed to tackle basic biological questions. For the time being, we will keep looking for more percussive performances by these fascinating mammals.

AUTHOR CONTRIBUTIONS

AR and LV developed the arguments and co-wrote the paper. All authors contributed to the article and approved the submitted version.

FUNDING

The work of LV and AR was supported by a Max Planck Research Group (MPRG) Awarded to AR.

ACKNOWLEDGMENTS

We are grateful to the Sealcentre Pieterburen, which let us observe pinniped behaviour and communication, which in turn spurred some of the hypotheses presented here.

REFERENCES

- Asselin, S., Hammill, M. O., and Barrette, C. (1993). Underwater vocalizations of ice breeding grey seals. *Can. J. Zool.* 71, 2211–2219. doi: 10.1139/z93-310
- Beier, J. C., and Wartzok, D. (1979). Mating behaviour of captive spotted seals (*Phoca largha*). *Anim. Behav.* 27, 772–781. doi: 10.1016/0003-3472(79)90013-7
- Bittman, E. L. (2020). Entrainment is NOT synchronization: an important distinction and its implications. *J. Biol. Rhythms.* 36:196–199. doi: 10.1177/0748730420972817
- Buck, J., and Buck, E. (1968). Mechanism of rhythmic synchronous flashing of fireflies: fireflies of Southeast Asia may use anticipatory time-measuring in synchronizing their flashing. *Science* 159, 1319–1327. doi: 10.1126/science.159.3821.1319
- Buss, D. M., Haselton, M. G., Shackelford, T. K., Bleske, A. L., and Wakefield, J. C. (1998). Adaptations, exaptations, and spandrels. *Am. Psychol.* 53:533. doi: 10.1037/0003-066X.53.5.533
- Byers, J., Hebets, E., and Podos, J. (2010). Female mate choice based upon male motor performance. *Anim. Behav.* 79, 771–8. doi: 10.1016/j.anbehav.2010.01.009
- Cady, A. B., Delaney, K. J., and Uetz, G. W. (2011). Contrasting energetic costs of courtship signaling in two wolf spiders having divergent courtship behaviors. *J. Arachnol.* 39, 161–165. doi: 10.1636/Hi09-70.1
- Clark, C. J. (2016). “Locomotion-induced sounds and sonations: mechanisms, communication function, and relationship with behavior,” In *Vertebrate Sound Production and Acoustic Communication*, eds. R. Suthers and E. Al (Cham: Springer I), 83–117.
- Cook, P., Rouse, A., Wilson, M., and Reichmuth, C. (2013). A California sea lion (*Zalophus californianus*) can keep the beat: motor entrainment to rhythmic auditory stimuli in a non vocal mimic. *J. Comp. Psychol.* 127, 412–427. doi: 10.1037/a0032345
- Cook, P. F., Hoard, V. A., Dolui, S., deB Frederick, B., Redfern, R., Dennison, S. E., et al. (2021). An MRI protocol for anatomical and functional evaluation of the California sea lion brain. *J. Neurosci. Methods* 353:109097. doi: 10.1016/j.jneumeth.2021.109097
- de Reus, K., Soma, M., Anichini, M., Gamba, M., de Heer Kloots, M., Lense, M. D., et al. (2020). Rhythm in dyadic interactions. *Philos. Trans. B.* doi: 10.31234/osf.io/9yrkv
- Dodenhoff, D. J., Stark, R. D., and Johnson, E. V. (2001). Do woodpecker drums encode information for species recognition? *Condor* 103, 143–150. doi: 10.1093/condor/103.1.143
- Dufour, V., Poulin, N., Charlotte Cur, é, and Sterck, E. H. M. (2015). Chimpanzee drumming: a spontaneous performance with characteristics of human musical drumming. *Sci. Rep.* 5:11320. doi: 10.1038/srep11320
- Dunlop, R. A., Cato, D. H., and Noad, M. J. (2010). Your attention please: increasing ambient noise levels elicits a change in communication behaviour in humpback whales (*Megaptera novaeangliae*). *Proc. R. Soc. B Biol. Sci.* 277, 2521–2529. doi: 10.1098/rspb.2009.2319
- Fish, F. E. (2000). Biomechanics and energetics in aquatic and semiaquatic mammals: platypus to whale. *Physiol. Biochem. Zool.* 73, 683–698. doi: 10.1086/318108
- Fitch, W. T., and Reby, D. (2001). The descended larynx is not uniquely human. *Proc. R. Soc. London B Biol. Sci.* 268, 1669–1675. doi: 10.1098/rspb.2001.1704
- Frankel, A. S. (2009). “Sound production,” in *Encyclopedia of Marine Mammals*, eds W. F. Perrin, B. Würsig, and J. G. M. Thewissen (Academic Press), 1056–1071.
- Galimberti, F., Sanvito, S., and Miller, E. (2008). Development of aggressive vocalizations in male southern elephant seals (*Mirounga leonina*): maturation or learning? *Behaviour* 145, 137–170. doi: 10.1163/156853907783244729
- Garcia, M., Herbst, C. T., Bowling, D. L., Dunn, J. C., and Fitch, W. T. (2017). Acoustic allometry revisited: morphological determinants of fundamental frequency in primate vocal production. *Sci. Rep.* 7, 1–11. doi: 10.1038/s41598-017-11000-x
- Garcia, M., and Ravignani, A. (2020). Acoustic allometry and vocal learning in mammals. *Biol. Lett.* 16:20200081. doi: 10.1098/rsbl.2020.0081
- Garcia, M., Theunissen, F., Sèbe, F., Clavel, J., Ravignani, A., Marin-Cudraz, T., et al. (2020). Evolution of communication signals and information during species radiation. *Nat. Commun.* 11, 1–5. doi: 10.1038/s41467-020-18772-3
- Gillooly, J. F., and Ophir, A. G. (2010). The energetic basis of acoustic communication. *Proc. R. Soc. B Biol. Sci.* 277, 1325–1331. doi: 10.1098/rspb.2009.2134
- Gould, S. J., and Lewontin, R. C. (1979). The spandrels of San Marco and the Panglossian paradigm: a critique of the adaptationist programme. *Proc. R. Soc. London B* 205, 581–598. doi: 10.1098/rspb.1979.0086
- Greenfield, M. D. (1994). Synchronous and alternating choruses in insects and anurans: common mechanisms and diverse functions. *Am. Zool.* 34, 605–615. doi: 10.1093/icb/34.6.605
- Hanlan, S. K. (1998). *Nosing Behaviour in Captive Harbour Seals (Phoca vitulina Concolor): Implications for Olfaction and Affiliation*. Doctoral Dissertation, Memorial University of Newfoundland.
- Hartbauer, M., and Römer, H. (2016). Rhythm generation and rhythm perception in insects: the evolution of synchronous choruses. *Front. Neurosci.* 10:223. doi: 10.3389/fnins.2016.00223
- Hayes, S. A., Kumar, A., Costa, D. P., Mellinger, D. K., Harvey, J. T., Southall, B. L., et al. (2004). Evaluating the function of the male harbour seal, *Phoca vitulina*, roar through playback experiments. *Anim. Behav.* 67, 1133–1139. doi: 10.1016/j.anbehav.2003.06.019
- Heinrich, T., Dehnhardt, G., and Hanke, F. D. (2016). Harbour seals (*Phoca vitulina*) are able to time precisely. *Anim. Cogn.* 19, 1133–1142. doi: 10.1007/s10071-016-1020-3
- Heinrich, T., Ravignani, A., and Hanke, F. H. (2020). Visual timing abilities of a harbour seal (*Phoca vitulina*) and a South African fur seal (*Arctocephalus pusillus pusillus*) for sub- and supra-second time intervals. *Anim. Cogn.* 23, 851–859. doi: 10.1007/s10071-020-01390-3
- Higham, J. P., and Hebets, E. A. (2013). An introduction to multimodal communication. *Behav. Ecol. Sociobiol.* 67, 1381–1388. doi: 10.1007/s00265-013-1590-x
- Hocking, D. P., Burville, B., Parker, W. M. G., Evans, A. R., Park, T., and Marx, F. G. (2020). Percussive underwater signaling in wild gray seals. *Mar. Mammal Sci.* 36, 728–732. doi: 10.1111/mms.12666
- Kuhn, C., and Frey, E. (2012). Walking like caterpillars, flying like bats-pinniped locomotion. *Palaeobiodivers. Palaeoenvir.* 92, 197–210. doi: 10.1007/s12549-012-0077-5
- Larsen, O. N., and Reichmuth, C. (2012). Proximal mechanisms for sound production in male Pacific walrus. *Can. Acoust.* 40:139.
- Mathevon, N., Casey, C., Reichmuth, C., and Charrier, I. (2017). Northern elephant seals memorize the rhythm and timbre of their rivals’ voices. *Curr. Biol.* 27, 2352–2356. doi: 10.1016/j.cub.2017.06.035
- McCulloch, S. (2000). *The Vocal Behaviour of the Grey Seal (Halichoerus grypus)*. Ph.D. thesis, University of St. Andrews, St. Andrews.
- Miles, M. C., Schuppe, E. R., and Fuxjager, M. J. (2020). Selection for rhythm as a trigger for recursive evolution in the elaborate display system of woodpeckers. *Am. Nat.* 195, 772–787. doi: 10.1086/707748
- Miles, M. C., Schuppe, E. R., Ligon, I. V., R. M., and Fuxjager, M. J. (2018). Macroevolutionary patterning of woodpecker drums reveals how sexual selection elaborates signals under constraint. *Proc. R. Soc. B Biol. Sci.* 285:20172628. doi: 10.1098/rspb.2017.2628
- Mitoyen, C., Quigley, C., and Fusani, L. (2019). Evolution and function of multimodal courtship displays. *Ethology* 125, 503–515. doi: 10.1111/eth.12882
- Munoz, N. E., and Blumstein, D. T. (2012). Multisensory perception in uncertain environments. *Behav. Ecol.* 23, 457–462. doi: 10.1093/beheco/arr220
- Newby, T. C. (1973). Observations on the breeding behavior of the harbor seal in the state of Washington. *J. Mammal.* 54, 540–543. doi: 10.2307/1379151
- Nowak, L. J. (2020). Observations on mechanisms and phenomena underlying underwater and surface vocalisations of grey seals. *Bioacoustics.* 1–20. doi: 10.1080/09524622.2020.1851298
- Partan, S. R. (2013). Ten unanswered questions in multimodal communication. *Behav. Ecol. Sociobiol.* 67, 1523–1539. doi: 10.1007/s00265-013-1565-y
- Partan, S. R. (2017). Multimodal shifts in noise: switching channels to communicate through rapid environmental change. *Anim. Behav.* 124, 325–337. doi: 10.1016/j.anbehav.2016.08.003
- Patel, A. D. (2006). Musical rhythm, linguistic rhythm, and human evolution. *Music Percept.* 24, 99–104. doi: 10.1525/mp.2006.24.1.99
- Rado, R., Wollberg, Z., and Terkel, J. (1991). The ontogeny of seismic communication during dispersal in the blind mole rat. *Anim. Behav.* 42, 15–21. doi: 10.1016/S0003-3472(05)80601-3

- Randall, J. A. (1984). Territorial defense and advertisement by footdrumming in bannertail kangaroo rats (*Dipodomys spectabilis*) at high and low population densities. *Behav. Ecol. Sociobiol.* 16, 11–20. doi: 10.1007/BF00293099
- Ravignani, A. (2019). Timing of antisynchronous calling: a case study in a harbor seal pup (*Phoca vitulina*). *J. Comp. Psychol.* 133:272. doi: 10.1037/com0000160
- Ravignani, A., Bowling, D. L., and Fitch, W. (2014). Chorusing, synchrony, and the evolutionary functions of rhythm. *Front. Psychol.* 5:1118. doi: 10.3389/fpsyg.2014.01118
- Ravignani, A., Fitch, W. T., Hanke, F. D., Heinrich, T., Hurgitsch, B., Kotz, S. A., et al. (2016). What pinnipeds have to say about human speech, music, and the evolution of rhythm. *Front. Neurosci.* 10:274. doi: 10.3389/fnins.2016.00274
- Ravignani, A., Olivera, V. M., Gingras, B., Hofer, R., Hernández, C. R., Sonnweber, R. S., et al. (2013). Primate drum kit: a system for studying acoustic pattern production by non-human primates using acceleration and strain sensors. *Sensors* 13, 9790–9820. doi: 10.3390/s130809790
- Reichmuth, C., Mulsow, J., and Schusterman, R. J. (2009). “Underwater acoustic displays of a Pacific walrus (*Odobenus rosmarus divergens*): Source level estimates and temporal patterning,” in *18th Biennial Conference on the Biology of Marine Mammals*. (Quebec City), 210.
- Reichmuth, C., and Schusterman, R. J. (2009). Annual temporal patterning in the vocalizations of captive seals: two long-term case studies. *J. Acoust. Soc. Am.* 125, 2676–2676. doi: 10.1121/1.4784215
- Remedios, R., Logothetis, N. K., and Kayser, C. (2009). Monkey drumming reveals common networks for perceiving vocal and nonvocal communication sounds. *Proc. Natl. Acad. Sci. U. S. A.* 106, 18010–18015. doi: 10.1073/pnas.0909756106
- Rouse, A. A., Cook, P. F., Large, E. W., and Reichmuth, C. (2016). Beat keeping in a sea lion as coupled oscillation: implications for comparative understanding of human rhythm. *Front. Neurosci.* 10:257. doi: 10.3389/fnins.2016.00257
- Russell, L., Purdy, J., and Davis, R. (2016). Social context predicts vocalization use in the courtship behaviors of weddell seals (*Leptonychotes weddellii*): a case study. *Anim. Behav. Cogn.* 3, 95–119. doi: 10.12966/abc.04.05.2016
- Schusterman, R. J., and Van Parijs, S. M. (2003). Pinniped vocal communication: an introduction. *Aquat. Mamm.* 29, 177–180.
- Sjare, B., and Stirling, I. (1981). I hear you knocking. *Nat. History* 3, 60–63.
- Sjare, B., Stirling, I., and Spencer, C. (2003). Structural variation in the songs of Atlantic walrus breeding in the Canadian High Arctic. *Aquat. Mammals* 29, 297–318. doi: 10.1578/016754203101024121
- Slocombe, K. E., Waller, B. M., and Liebal, K. (2011). The language void: the need for multimodality in primate communication research. *Anim. Behav.* 81, 919–924. doi: 10.1016/j.anbehav.2011.02.002
- Soulsbury, C. D., and Halsey, L. G. (2018). Does physical activity age wild animals? *Front. Ecol. Evol.* 6:222. doi: 10.3389/fevo.2018.00222
- Tyack, P. L., and Miller, E. H. (2002). “Vocal anatomy, acoustic communication and echolocation,” in *Marine Mammal Biology: An Evolutionary Approach*, ed. A. R. Hoelzel (Blackwell), 142–184.
- Van Parijs, S. M., Hastie, G. D., and Thompson, P. M. (1999). Geographical variation in temporal and spatial vocalization patterns of male harbour seals in the mating season. *Anim. Behav.* 58, 1231–1239. doi: 10.1006/anbe.1999.1258
- Venables, U. M., and Venables, L. S. V. (1957). Mating behaviour of the seal *Phoca vitulina* in Shetland. *Proc. Zool. Soc. London* 128, 387–396. doi: 10.1111/j.1096-3642.1957.tb00332.x
- Wahlberg, M., Lunneryd, S.-G., and Westerberg, H. (2002). The source level of harbour seal flipper slaps. *Aquat. Mammals* 28, 90–92.
- Wilson, M., and Cook, P. F. (2016). Rhythmic entrainment: why humans want to, fireflies can't help it, pet birds try, and sea lions have to be bribed. *Psychon. Bull. Rev.* 23, 1647–1659. doi: 10.3758/s13423-016-1013-x
- Zahavi, A. (1975). Mate selection-A selection for a handicap. *J. Theor. Biol.* 53, 205–214. doi: 10.1016/0022-5193(75)90111-3

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2021 Verga and Ravignani. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.