



# Brains in space: the importance of understanding the impact of long-duration spaceflight on spatial cognition and its neural circuitry

Alexander C. Stahn<sup>1,2</sup> · Simone Kühn<sup>3,4</sup>

Received: 16 July 2021 / Accepted: 21 July 2021 / Published online: 18 August 2021  
© The Author(s) 2021

## Abstract

Fifty years after the first humans stepped on the Moon, space faring nations have entered a new era of space exploration. NASA's reference mission to Mars is expected to comprise 1100 days. Deep space exploratory class missions could even span decades. They will be the most challenging and dangerous expeditions in the history of human spaceflight and will expose crew members to unprecedented health and performance risks. The development of adverse cognitive or behavioral conditions and psychiatric disorders during those missions is considered a critical and unmitigated risk factor. Here, we argue that spatial cognition, i.e., the ability to encode representations about self-to-object relations and integrate this information into a spatial map of the environment, and their neural bases will be highly vulnerable during those expeditions. Empirical evidence from animal studies shows that social isolation, immobilization, and altered gravity can have profound effects on brain plasticity associated with spatial navigation. We provide examples from historic spaceflight missions, spaceflight analogs, and extreme environments suggesting that spatial cognition and its neural circuitry could be impaired during long-duration spaceflight, and identify recommendations and future steps to mitigate these risks.

**Keywords** Spatial cognition · Hippocampus · Spaceflight · Extreme environments

## Moon, mars and beyond: pushing the limits of human performance

On July 20, 1969, the world held its breath when Neil Armstrong and Buzz Aldrin landed on the Moon and became the first humans to explore the lunar surface. Now about fifty years later, space faring civilizations such as China, Japan, Europe, and India have joined the US and Russia in a new era of space exploration that goes well beyond the Moon. Spaceflight will play an increasingly important role in accelerating technological developments and transfer, establishing gateways to foster deep space exploration, seeking extraterrestrial life, establishing lunar colonies, venturing into deep space, and sending humans to Mars. Private partnerships and entities are fueling this race. *SpaceX* has been the first private company to transport astronauts to orbit and the International Space Station (ISS). Reusable rockets and super heavy launch systems such as *Starship* are expected to reduce costs of interplanetary flight and accelerate the developments needed for the colonization of other planets. Similar approaches and technologies have been recently announced by the China Academy of Launch

---

This article is a contribution to the proceedings of the “8th International Conference on Spatial Cognition: Cognition and Action in a Plurality of Spaces” (ICSC 2021).

---

✉ Alexander C. Stahn  
astahn@pennmedicine.upenn.edu

<sup>1</sup> Department of Psychiatry, Unit of Experimental Psychiatry, Perelman School of Medicine, University of Pennsylvania, 4233 Guardian Dr, 1016 Blockley Hall, Philadelphia, PA 19104, USA

<sup>2</sup> Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin and Humboldt-Universität zu Berlin, Institute of Physiology, Charitéplatz 1, 10117 Berlin, Germany

<sup>3</sup> Lise Meitner Group for Environmental Neuroscience, Max Planck Institute for Human Development, 14195 Berlin, Germany

<sup>4</sup> Department of Psychiatry and Psychotherapy, University Medical Center Hamburg-Eppendorf, 20246 Hamburg, Germany

Vehicle Technology, coinciding with China's announcement to send its first human crew to Mars in 2033 and establish a large-scale settlement on the *Red Planet*. No matter which nation will win the next phase of this new *Space Race*, it will push the limits of human health and performance.

Space is a naturally hostile environment characterized by reduced gravity levels and various environmental, operational, and psychological stressors (e.g., radiation, hypercapnia, little separation between rest and works schedules, social isolation and confinement). Future exploratory missions will be considerably longer than current standard ISS missions. NASA's reference mission to Mars is expected to comprise 1100 days. Deep space exploratory class missions could even span decades. These expeditions are considered the most dangerous and difficult explorations in the history of human spaceflight, and will expose crew members to unprecedented health and performance risks. The development of adverse cognitive or behavioral conditions and psychiatric disorders during those missions is considered a critical and unmitigated risk factor (Slack et al. 2016). To ensure successful future human space exploration, the risks must be precisely identified. Tools need to be provided that foster the efficient their efficient monitoring and prediction of adverse effects of spaceflight on brain and cognitive performance. In addition, target-specific countermeasures have to be established that help mitigating neurobehavioral impairments that may put astronaut health and mission success at risk.


### Risk of adverse brain and cognitive changes in response to long-duration spaceflight

Neuroimaging findings have revealed structural brain changes in response to spaceflight, including an upward shift of the brain, redistribution of cerebrospinal fluid, ventricular volume decreases, and widespread decreases in gray matter volume (Jillings et al. 2020; Roberts et al. 2017; Van Ombergen et al. 2018, 2019). These findings seem to stand in contrast to data on cognitive performance, revealing only minor impairments in response to spaceflight (Strangman et al. 2014). Whether and to what extent these changes lead to operational impairments and adverse behavioral conditions is currently not well understood. Retrospective analyses investigating the relationships between changes in cognitive performance using NASA's cognitive test battery *WinSCAT* (Kane et al. 2005) and whole-brain structural analyses are inconclusive. Higher total ventricular volume was associated with reduced accuracy on a symbol substitution task ("Code Substitution" test), and faster response speed in an n-back paradigm ("Running Memory Continuous Performance" test) (Roberts et al. 2019). Other findings from standard 6-month ISS missions have shown significant decrements

in manual dexterity, dual-tasking, motion perception, and a considerable degradation of a virtual navigation, i.e., car driving task immediately after return from space (Moore et al. 2019). Whereas these effects are expected to be relatively short-lived, i.e., minutes up to several days, they are considered a substantial risk for exploratory space missions, which involve transitions between gravitational levels (Harm et al. 2015). Data from a 1-year mission using NASA's *Cognition* battery (Basner et al. 2015), assessing the cognitive performance of ten neuropsychological tests, suggests that adverse cognitive effects can persist up to 6 months postflight (Garrett-Bakelman et al. 2019). According to a review of studies that were performed during space missions, current data do not support cognitive deficits in low Earth orbit (Strangman et al. 2014). However, because of different approaches, methodologies, study durations, and small sizes, the effects of spaceflight on cognitive functions remain to be determined. However, because of different approaches, methodologies, study durations, and small sizes, the effects of spaceflight on cognitive functions remain to be determined (Mammarella 2020). Together, these data raise several key questions: (1) Could more complex cognitive and operational tasks be more sensitive to changes of inflight performance? (2) Is the neural circuitry underlying these tasks affected by spaceflight? (3) What are the long-term consequences of spaceflight on these tasks?

### Need to monitor visuospatial abilities before, during and after spaceflight

Operational performance can be assessed by simulating spaceflight-related tasks such as using robotic arms to capture a transiting spacecraft or to control a spacecraft to maneuver it and dock it to another vehicle (Ivkovic et al. 2019; Johannes et al. 2017). Successful completion of the maneuvers relates to various cognitive domains, including but not limited to situational awareness, planning, decision-making, object orientation, mental rotation, visual processing, fine motor control, and visual motor integration (Wong et al. 2020). It can be hypothesized that complex tasks assessing the encoding, processing, storage, and retrieval of visuospatial information could be particularly vulnerable during spaceflight. An analysis of shuttle missions revealed that touchdown speed in 20% of orbiter landings was outside acceptable limits, some of which were associated with pilot-induced oscillations, i.e., increasing flight corrections in opposite directions (Moore et al. 2008). In 1997, piloting errors during control of the TORU (Teleoperated Mode of (spacecraft) Control) system resulted in the collision of the supply spacecraft Progress M-34 with the MIR station, damaging the *Spektr* module and a solar panel (Morgan 2013). Likewise, several telerobotic incidents occurred on ISS,



**Fig. 1** Traverse of Astronauts Edgar D. Mitchell and Alan Shepard during extravehicular activity (EVA-2) of the Apollo 14 mission. (a) Edgar D. Mitchell moves across the lunar surface as he is studying a map, trying to figure out where they are in vain; both astronauts thought they were much closer to Cone Crater than they actually were, and they did not recognize any landmarks in their view (picture was taken at location B1). (b) Outline of the traverse from Lunar

Module to Cone Crater via B1 and back. Station C1 indicates “Saddle Rock,” where the last sample was retrieved before returning to the Lunar Module. Neither astronaut noticed that “Saddle Rock” was depicted as a landmark on their map. It was only after the completion of the mission that they had realized that were only 30 m from the rim of Cone Crater. Picture Credit: NASA/USGS and Google


including a collision between the *Canadarm2*, the shuttle payload door, an external antenna, and Canadarm in which the two robotic arms crossed within 1.5 m of each other (Moore et al. 2019).

Spatial updating, path integration, route learning, way-finding, and cognitive mapping are also key to successfully navigating in small- and large-scale environments. The criticality of encoding representations about self-to-object relations and integrating this information into a spatial map of the environment for spaceflight operations was highlighted during the Apollo 14 mission. Astronauts Ed Mitchell and Alan Shepard had to walk to a crater located within a mile from their landing module. Having nearly reached the target destination, they had to abort the assignment because of spatial disorientation (Fig. 1). The difficulties associated with spatial navigation on the lunar surface were confirmed during the post-mission debriefs when Alan Shepard explained, “I felt that we had a navigation problem on EVA-2. I don’t know why we didn’t worry a little bit more about that pre-flight (...) Second, there’s no questions that it is easy to misjudge distances, not only above the surface [that is during the landing or from the Lunar Module windows] (...) but also distances along the surface.” (Heiken and Jones 2007).

Finding your way in a new territory has always been a significant challenge for explorers. From an evolutionary perspective, the “(...) ability to estimate one’s own position

and track and plan one’s own path in physical space is key to survival” (*Focus on Spatial Cognition* 2017). To identify the time course of visuospatial abilities in response to spaceflight, we have developed a specific battery of tasks that was recently flight-certified for use on the ISS. This battery assesses visuospatial memory formation, topographic mapping, path integration, and spatial updating. The battery has been tested in various spaceflight analogs on Earth. Experiments performed during parabolic flight maneuvers have shown that spatial updating is sensitive to gravitational changes, including micro- and hyper-gravity (Stahn et al. 2020). Recently, *Spatial Cognition* was selected as part of NASA’s CIPHER<sup>1</sup> project. *Spatial Cognition* will investigate visuospatial changes and their neural basis in a total of 30 astronauts, equally assigned to 2-month, 6-month, and 1-year missions. Likewise, it will be essential to identify cortical and subcortical brain areas associated with the spatial encoding of landmark identities, retrieving spatial information, and processing visual features important for landmark recognition such as the hippocampal formation, striatum, precuneus, retrosplenial complex, parahippocampal place area,

<sup>1</sup> CIPHER stands for Complement of Integrated Protocols for Human Exploration Research, and is a project comprising 17 investigators that integrate various disciplines to investigate multiple physiological, biological, and psychological aspects to short-duration (2-months), standard-duration (6-months), and long-duration (12-months) human spaceflight missions.



**Fig. 2** Environmental, operational, and psychological stressors associated with spaceflight. Ionizing radiation, hypercapnia (increased CO<sub>2</sub> levels), altered vestibular stimulation and reduced physical activity in response to weightlessness, circadian disruptions and poor sleep due to altered day and night cycles, isolation and confinement, and sensory deprivation can have adverse effects on hippocampal plasticity. The hippocampus is critical for declarative memory formation, emotion processing, and spatial cognition. Together with the entorhinal cortex, the hippocampus supports the encoding, consolidation,

and retrieval of spatial information by processing information about location (place cells of the hippocampus) relative to a grid map characterized by a hexagonal pattern (grid cells in the entorhinal cortex). Picture Credit: Schematic brain and hippocampus were created with [BioRender.com](https://www.biorender.com). Spacecraft (middle) and right icon in bottom row (sensory deprivation): NASA; astronaut silhouette (third icon, bottom row) by Natasha Sinagina/CC BY); icon depicting Non-24h Day/Night cycles by [icon-library.com](https://www.icon-library.com)



and the occipital place area (Epstein et al. 2017; Geerts et al. 2020; Hartley et al. 2014; Wolbers et al. 2008). The hippocampus is considered the human “inner” GPS by providing information about location (place cells of the hippocampus) relative to a grid map characterized by a hexagonal pattern generated by grid cell firing activity in the entorhinal cortex (Moser et al. 2015). Together, the hippocampus and entorhinal cortex play a critical role in exploring unfamiliar terrains, navigating on new planets, and performing complex operational visuospatial tasks (Fig. 2). To this aim, *Spatial Cognition* will combine behavioral data with multi-modal neuroimaging that are expected to provide new knowledge on the dose–response relationships between the length of spaceflight missions, brain changes, and their implications for spatial orientation and navigation. In addition, data will be collected up to a year after return from space to identify the time course of recovery.

### Environmental stressors, hippocampal plasticity and spatial cognition

The hippocampus, known as a highly plastic brain region that is key to complex spatial navigation, is vulnerable to various stressors associated with spaceflight, including but not limited to radiation, hypercapnia, altered vestibular stimulation, reduced physical activity levels, circadian disorders and poor sleep, sensory deprivation, and social isolation and confinement (Fig. 2). It is also possible that these stressors interact with each other. The strength and direction of such effects is currently not well understood, and also deserves further research.

#### Radiation

Cosmic radiation is expected to be a critical risk for adverse neurobehavioral effects during spaceflight. Recent reviews reported considerable structural and functional damage of the brain that is particularly prominent in the prefrontal cortex and hippocampus, and associated with a broad range of adverse behavioral conditions, including taste aversion, reversal learning deficits, disrupted reinforcement behavior, contextual fear conditioning, and spatial learning and memory formation (Kiffer et al. 2019).

#### Carbon dioxide

The limited capacity of air recycling systems on spacecraft can increase carbon dioxide (CO<sub>2</sub>) levels and lead to hypoxia/hypercapnia. Typical concentrations of CO<sub>2</sub> on ISS range between 2 and 4 mmHg and can be up to ten-fold higher than on Earth (outdoors about 0.3 mmHg and

in well-ventilated rooms about 0.5 mmHg). Animal studies have shown that chronic exposure to 0.3% CO<sub>2</sub> concentrations can impair brain plasticity and behavior during early development (Kiray et al. 2014). In contrast to earlier reports (Satish et al. 2012), recent data found no or little change of varying levels of CO<sub>2</sub> acutely on cognitive performance (Basner et al. 2021; Lee et al. 2019; Scully et al. 2019). The chronic effects of heightened CO<sub>2</sub> levels on complex visuospatial abilities remain to be determined.

#### Weightlessness

Weightlessness also affects the vestibular system, which goes beyond maintaining gaze and postural stabilization. Acute exposure to weightlessness and transitions between gravity levels significantly challenge the integration of neuro-vestibular signaling, associated with motion sickness and alterations in spatial abilities and sensorimotor functioning (Reschke and Clément 2018). Otoliths, graviceptors in the inner ear, rely on information from linear acceleration, and therefore cannot respond to tilt when gravity is lacking. Furthermore, projections of the vestibular pathways to the limbic system and neocortex play a critical role for brain plasticity, including spatial learning and memory formation (Smith 2017). Peripheral lesions of the vestibular pathways have been linked to atrophy in the hippocampus and spatial memory impairments that are long-lasting and may even be permanent (Smith et al. 2010).

#### Body unloading

The lack of gravity also diminishes physical activity, known as a critical driver for brain plasticity and cognition (Voss et al. 2013). We recently showed that long-duration bed rest (> 1 month) has detrimental effects on various cognitive processes, including episodic memory formation in the hippocampus and parahippocampus (Brauns et al. 2019, 2021; Friedl-Werner et al. 2020). We also found that prolonged physical inactivity associated with bed rest induces circadian disruptions (Mendt et al. 2021a, b).

#### Non-24 h light–dark cycles

Spaceflight is associated with circadian disruptions and adverse sleep (Barger et al. 2014; Flynn-Evans et al. 2016). A recent analysis of astronauts on standard ISS missions (average stay of 155 days) reported circadian misalignment in 20% of flight days and corresponding sleep losses of one hour per night (Flynn-Evans et al. 2016). Poor sleep is associated with neurodegenerative and neuropsychiatric conditions and hippocampal atrophy (Fjell et al. 2020).

Chronic sleep deprivation has also been shown to lead to hippocampal atrophy across the adult lifespan (Fjell et al. 2020). The detrimental effects of sleep on the hippocampus are independent of stress hormones (Mueller et al. 2008), and structural hippocampal alterations have been observed after brief periods of sleep deprivation (Raven et al. 2019).

### Isolation, confinement, and sensory deprivation


Reduced sensory stimulation and sensory monotony experienced in isolated, confined, and extreme (ICE) environments are expected to be major contributors to adverse neurobehavioral conditions. Monotonous sensory stimulation, boredom, and isolation and confinement are severe stressors can lead to interpersonal tension and conflict, negative affect, work place errors, and increased mortality (Eastwood et al. 2012). For more than 30 years, space agencies have been investigating the effects of isolation and confinement using facilities designed to simulate spaceflight missions. These “laboratory” studies are characterized by highly controlled settings and have been referred to as isolated and controlled confinement (ICC) (Choukér and Stahn 2020). With few exceptions such as the Russian Mars500 study or the more recent SIRIUS projects using the NEK facility at the Institute of Biomedical Problems (IBMP) in Moscow, these studies are typically limited to short durations (<60 days). Laboratory experiments also lack the complexity, unpredictability and risks associated with actual expeditions in extreme environments. Exploration expeditions bear the potential to study the effects of prolonged isolation and confinement in natural extreme environments and have been become known as isolated confined and extreme environments (ICE). The first reported data on the behavioral effects of isolation and confinement date back to Polar explorers, providing anecdotal evidence of the psychological and physiological challenges associated with long-duration Antarctic expeditions (Palinkas and Suedfeld 2008). Previous research in ICCs and ICEs focused on mood disorders, asthenia, psychosomatic reactions, psychosocial adaptations, and psychiatric emergencies (McPhee and Charles 2009). To what extent social isolation directly causes brain changes, and cognitive performance impairments is less clear. Animal studies have shown that stress and social isolation disrupt hippocampal neurogenesis (e.g., Cinini et al. 2014; Gould et al. 1997; Schloesser et al. 2010), prevents exercise-induced hippocampal neurogenesis (Leasure and Decker 2009; Pereda-Pérez et al. 2013; Stranahan et al. 2006), selectively reduces hippocampal brain-derived neurotrophic factor (Scaccianoce et al. 2006), and impairs hippocampal long-term potentiation (Kamal et al. 2014).

To investigate whether similar effects can be detected in humans, we recently investigated the neurobehavioral responses to prolonged isolation associated with Antarctic overwintering (Stahn et al. 2019). T<sub>1</sub>- and T<sub>2</sub>-weighted magnetic resonance imaging (MRI) data were collected before

and 1.5 months after a 14-month expedition to Antarctica to assess structural brain changes and compare these data to a control group matched for sex, age, and educational background. Hippocampal subfield volumes decreased after the expedition, namely bilateral dentate gyrus volume was significantly smaller in the expeditioners than in the control group (mean group decrease in volume  $\pm$  SE:  $32 \pm 13$  mm<sup>3</sup>, equivalent to a  $7.2 \pm 3\%$  volume reduction). Whole-brain analyses using voxel-based morphometry (VBM) revealed further decreases of gray matter probability in the left parahippocampus (mean group decrease  $\pm$  SE:  $3.84 \pm 0.72\%$ ), and in the right lateral and left medial and right lateral prefrontal cortex (PFC) (mean group decrease  $\pm$  SE:  $3.33 \pm 0.48\%$ ; left medial PFC: mean group decrease  $\pm$  SE:  $2.99 \pm 0.25\%$ ). Brain-derived trophic factor (BDNF), a protein key to brain plasticity and learning and memory formation (Egan et al. 2003; Harward et al. 2016), was determined in serum blood samples collected before, ten times during, and once after the expedition. After the first quarter of the expedition, serum BDNF concentration was reduced compared to the baseline measurement before the expedition and did not recover at 1.5 months after the end of the expedition (mean reduction  $\pm$  SE:  $11 \pm 1.5$  ng/mL,  $45 \pm 4.9\%$ ). Reductions in BDNF from pre- to post-mission were associated with decreases in dentate gyrus volume ( $R^2=0.47$ ). The reductions in dentate gyrus volume were also associated with lower cognitive performance in tests of spatial processing ( $R^2=0.87$ ) and the resolution of response conflict ( $R^2=0.82$ ), but there was no reduction in performance in other cognitive tests (i.e., Digit Symbol Substitution, Stroop Congruent task).

### Need for target-specific countermeasures

To mitigate adverse neurobehavioral effects of prolonged spaceflight on spatial cognition and its neural basis, target-specific countermeasure will be needed that go beyond current practices such as exercise, lower body negative pressure, and nutritional supplementation. For instance, specific types of video gaming have the potential to enhance brain plasticity (Kühn et al. 2014a, b; 2017). Further, specific training programs aimed at improving operational performance skills (e.g., Johannes et al. 2017) can be expected to improve visuospatial abilities and affect their neural circuitry. Moreover, combining physical activity with virtual environments could be promising (Vessel and Russo 2015). Preliminary data analyses of the NASA sponsored project *Hybrid Training* showed that voluntary exercise on bicycle ergometer combined with a visual sensory stimulation could mitigate some of the neurobehavioral effects in response to 14 months of isolation and confinement associated with overwintering at Neumayer III station in Antarctica. These effects were manifested as increased BDNF concentrations and reductions in



**Fig. 3** Individualized countermeasures (ICount) to mitigate adverse neurobehavioral effects. To mitigate the neurobehavioral risks associated with long-duration spaceflight, a comprehensive “toolbox” of countermeasures is needed. The figure lists some examples that will play a critical role in reducing the risks of adverse cognitive and behavioral effects and psychiatric disorders during long-duration expeditions. Some of the countermeasures listed such as exercise, videogaming, diet and nutritional supplementation, sleep hygiene, and self-adapted visuo-spatial learning tasks will also help to main-

tain hippocampal plasticity and spatial cognition. The relationships between individual countermeasures will vary between and within individuals. The relative importance of specific strategies will vary during the course of mission, requiring a constant reevaluation of the crewmembers’ individual needs. Note that the figure is a schematic illustration, and the weights of the interventions are used to reflect their dynamic nature through an expedition, but do not suggest any importance of one countermeasure over the other

hippocampal subfield volume and whole-brain gray matter and support the role of physical activity as a key driver of brain plasticity (Vivar and van Praag 2017).

Given the range of environmental, operational and psychological conditions and stressors, it is expected that there is no single countermeasure that will serve as a universal remedy. In addition, it is possible that the responses to the countermeasures will vary between individuals. Countermeasure must therefore be understood as a dynamic construct that is optimized relative to the individual needs as a function of mission duration. The concept of individualized countermeasure that we like to term as “ICount” is summarized in Fig. 3. A multiplicity of methodologies and approaches could be combined in a toolbox that is flexibly adapted to address the crews’ individual needs. The countermeasures range from habitat design (e.g., lighting, personal and social space), to exercise, workload considerations including variations in meaningful work, sleep/rest schedules, relaxation techniques, videogaming, entertainment, virtual reality, plants, food, and other sensory augmentation measures, to strategies for maintaining and enhancing crew cohesion, psychological counseling and family support. The variety of approaches will be critical to maximize the stimulation and their synergies, and consider phenotypic differences, and the dynamic nature of individual preferences for specific needs during long-duration expeditions.

## Summary and Conclusions

Outer space is considered the most extreme environment for human mankind. Without support systems and protective suits, life in space or on other planets in our solar system is

not possible. Even prolonged stays in the habitat of a spacecraft pose significant physiological and psychological challenges. Spaceflight affects every organ system. In addition to microgravity the spacecraft setting is characterized by multiple environmental toxicants and operational stressors such as radiation, noise, hypercapnia, hypoxia, decompression, dietary restrictions, fluid shifts, increased intracranial pressure, non-24 h light–dark cycles, acute operational shifts in sleep timing, psychological factors related to high workload under pressure, operational and interpersonal distress, and isolation and confinement. The effects of space travel on brain and behavior are currently not well understood but are considered a high and unmitigated risk for future long-duration space missions.

Studies in animals and ground-based spaceflight analogs suggest that the spatial cognition and its neural basis could be particularly vulnerable to future long-duration space missions. Future studies are therefore critically needed to (1) understand the effects of extreme environments and spaceflight on spatial cognition and its neural circuitry, (2) demonstrate and verify the techniques needed to monitor, diagnose, and prevent such effects, and (3) develop target-specific countermeasures to mitigate adverse effects on visuospatial abilities. In addition, imaging and cognitive data should be complemented by biochemical assessments, and advances in multi-omics technologies such as genomics, transcriptomics, proteomics, and metabolomics. They will be critical to close knowledge gaps of the underlying molecular mechanisms and genetic drivers of neurobehavioral adaptations in extreme environments. Combining brain imaging, cognitive and biochemical methodologies, and outcomes could provide the basis to better understand and characterize the type,

extent, cause, and mechanisms of adverse neurobehavioral effects and their phenotypic signatures.

The integration of imaging, physiological, biochemical, and behavioral data will contribute to the space agencies' goal to provide knowledge, technologies, and tools to enable safe, reliable, and productive human space exploration. At the same time, they can also benefit research and applications on Earth. Spaceflight analogs such as isolation experiments, Antarctic expeditions, and bed rest studies, can provide unique standardized settings that induce neurophysiological and psychological conditions that typically evolve over long time spans, and cannot be replicated in typical laboratory settings. The opportunity to study prospectively the time course of brain and behavioral changes in time lapse in healthy adults, characterize them before any clinical manifestations occur, and follow them up through recovery can help understand the effects and the biological basis of aging-related cognitive decline, social isolation, clinical manifestations associated with impaired physical mobility, or lifestyle changes in response to pandemics.

**Funding** Open Access funding enabled and organized by Projekt DEAL. The work indicated here are supported by the German Aerospace Agency (DLR) through Grants 50WB1525, 50WB1915, and 50WB2115.

## Declarations

**Conflict of interest** The authors declare that there is no conflict of interest.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## References

- Barger LK, Flynn-Evans EE, Kubey A, Walsh L, Ronda JM, Wang W, Wright KP, Czeisler CA (2014) Prevalence of sleep deficiency and use of hypnotic drugs in astronauts before, during, and after spaceflight: an observational study. *Lancet Neurol* 13(9):904–912. [https://doi.org/10.1016/S1474-4422\(14\)70122-X](https://doi.org/10.1016/S1474-4422(14)70122-X)
- Basner M, Savitt A, Moore TM, Port AM, McGuire S, Ecker AJ, Nasrini J, Mollicone DJ, Mott CM, McCann T, Dinges DF, Gur RC (2015) Development and validation of the cognition test battery for spaceflight. *Aerosp Med Hum Perform* 86(11):942–952. <https://doi.org/10.3357/AMHP.4343.2015>
- Basner M, Stahn AC, Nasrini J, Dinges DF, Moore TM, Gur RC, Mühl C, Macias BR, Laurie SS (2021) Effects of head-down tilt bed rest plus elevated CO<sub>2</sub> on cognitive performance. *J Appl Physiol* 130(4):1235–1246. <https://doi.org/10.1152/jappphysiol.00865.2020>
- Brauns K, Werner A, Gunga H-C, Maggioni MA, Dinges DF, Stahn A (2019) Electrocortical evidence for impaired affective picture processing after long-term immobilization. *Sci Rep* 9(1):16610. <https://doi.org/10.1038/s41598-019-52555-1>
- Brauns K, Friedl-Werner A, Gunga H-C, Stahn AC (2021) Effects of two months of bed rest and antioxidant supplementation on attentional processing. *Cortex* 141:81–93. <https://doi.org/10.1016/j.cortex.2021.03.026>
- Choukér A, Stahn AC (2020) COVID-19—the largest isolation study in history: the value of shared learnings from spaceflight analogs. *Npj Microgravity* 6(1):32. <https://doi.org/10.1038/s41526-020-00122-8>
- Cinini SM, Barnabe GF, Galvão-Coelho N, de Medeiros MA, Perez-Mendes P, Sousa MBC, Covolan L, Mello LE, Galvao-Coelho N, de Medeiros MA, Perez-Mendes P, Sousa MBC, Covolan L, Mello LE (2014) Social isolation disrupts hippocampal neurogenesis in young non-human primates. *Front Neurosci* 8:45. <https://doi.org/10.3389/fnins.2014.00045>
- Eastwood JD, Frischen A, Fenske MJ, Smilek D (2012) The unengaged mind: defining boredom in terms of attention. *Perspect Psychol Sci* 7(5):482–495. <https://doi.org/10.1177/1745691612456044>
- Egan MF, Kojima M, Callicott JH, Goldberg TE, Kolachana BS, Bertolino A, Zaitsev E, Gold B, Goldman D, Dean M, Lu B, Weinberger DR (2003) The BDNF val66met polymorphism affects activity-dependent secretion of BDNF and human memory and hippocampal function. *Cell* 112(2):257–269
- Epstein RA, Patai EZ, Julian JB, Spiers HJ (2017) The cognitive map in humans: spatial navigation and beyond. *Nat Neurosci* 20(11):1504–1513. <https://doi.org/10.1038/nn.4656>
- Fjell AM, Sørensen Ø, Amlien IK, Bartrés-Faz D, Bros DM, Buchmann N, Demuth I, Drevon CA, Düzel S, Ebmeier KP, Idland A-V, Kietzmann TC, Kievit R, Kühn S, Lindenberger U, Mowinckel AM, Nyberg L, Price D, Sexton CE, Walhovd KB (2020) Self-reported sleep relates to hippocampal atrophy across the adult lifespan: results from the Lifebrian consortium. *Sleep* 43(5):zsz280. <https://doi.org/10.1093/sleep/zsz280>
- Flynn-Evans EE, Barger LK, Kubey AA, Sullivan JP, Czeisler CA (2016) Circadian misalignment affects sleep and medication use before and during spaceflight. *Npj Microgravity* 2(1):1–6. <https://doi.org/10.1038/npjmgrav.2015.19>
- Focus on Spatial Cognition (2017) *Nat Neurosci* 20(11):1431. <https://doi.org/10.1038/nn.4666>
- Friedl-Werner A, Brauns K, Gunga H-C, Kühn S, Stahn A (2020) Exercise-induced changes in brain activity during memory encoding and retrieval after long-term bed rest. *Neuroimage* 223:117359. <https://doi.org/10.1016/j.neuroimage.2020.117359>
- Garrett-Bakelman FE, Darshi M, Green SJ, Gur RC, Lin L, Macias BR, McKenna MJ, Meydan C, Mishra T, Nasrini J, Piening BD, Rizzardi LF, Sharma K, Siamwala JH, Taylor L, Vitaterna MH, Afkarian M, Afshinnekoo E, Ahadi S, Turek FW (2019) The NASA Twins Study: a multidimensional analysis of a year-long human spaceflight. *Science (new York, n.y.)* 364(6436):127–128. <https://doi.org/10.1126/science.aau8650>
- Geerts JP, Chersi F, Stachenfeld KL, Burgess N (2020) A general model of hippocampal and dorsal striatal learning and decision making. *Proc Natl Acad Sci USA* 117(49):31427–31437. <https://doi.org/10.1073/pnas.2007981117>
- Gould E, McEwen BS, Tanapat P, Galea LA, Fuchs E (1997) Neurogenesis in the dentate gyrus of the adult tree shrew is regulated



- by psychosocial stress and NMDA receptor activation. *J Neurosci* 17(7):2492–2498
- Harm DL, Reschke MF, Wood SJ (2015) Spatial orientation and motion perception in microgravity. In: Hoffman R, Hancock P, Scerbo M, Parasuraman R, Szalma J (eds) *The Cambridge handbook of applied perception research*. Cambridge University Press, pp 912–929. <https://doi.org/10.1017/CBO9780511973017.054>
- Hartley T, Lever C, Burgess N, O’Keefe J (2014) Space in the brain: how the hippocampal formation supports spatial cognition. *Philos Trans R Soc Lond B Biol Sci* 369(1635):20120510. <https://doi.org/10.1098/rstb.2012.0510>
- Harward SC, Hedrick NG, Hall CE, Parra-Bueno P, Milner TA, Pan E, Laviv T, Hempstead BL, Yasuda R, McNamara JO (2016) Autocrine BDNF-TrkB signalling within a single dendritic spine. *Nature* 538(7623):99–103. <https://doi.org/10.1038/nature19766>
- Heiken G, Jones EM (2007) *On the moon: the Apollo journals*. Springer
- Ivkovic V, Sommers B, Cefaratti DA, Newman G, Thomas DW, Alexander DG, Strangman GE (2019) Operationally relevant behavior assessment using the robotic on-board trainer for research (ROBoT-r). *Aerosp Med Hum Perform* 90(9):819–825. <https://doi.org/10.3357/AMHP.5324.2019>
- Jillings S, Van Ombergen A, Tomilovskaya E, Rumshiskaya A, Litvinova L, Nosikova I, Pechenkova E, Rukavishnikov I, Kozlovskaya IB, Manko O, Danilichev S, Sunaert S, Parizel PM, Sinityn V, Petrovichev V, Laureys S, Eulenburg P, Sijbers J, Wuyts FL, Jeurissen B (2020) Macro- and microstructural changes in cosmonauts’ brains after long-duration spaceflight. *Sci Adv* 6(36):eaaz9488. <https://doi.org/10.1126/sciadv.aaz9488>
- Johannes B, Bronnikov S, Bubeev Y, Dudukin A, Hoermann H, Frett T, Rittweger J, Gaillard A (2017) A tool to facilitate learning in a complex manual control task. *Int J Appl Psychol* 7(4):79–85. <https://doi.org/10.5923/j.jiap.20170704.01>
- Kamal A, Ramakers GMJ, Altinbilek B, Kas MJH (2014) Social isolation stress reduces hippocampal long-term potentiation: effect of animal strain and involvement of glucocorticoid receptors. *Neuroscience* 256:262–270. <https://doi.org/10.1016/j.neuroscience.2013.10.016>
- Kane RL, Short P, Sipes W, Flynn CF (2005) Development and validation of the Spaceflight Cognitive Assessment Tool for Windows (WinSCAT). *Aviat Space Environ Med* 76(6 SUPPL.):B183–B191
- Kiffer F, Boerma M, Allen A (2019) Behavioral effects of space radiation: a comprehensive review of animal studies. *Life Sci Space Res* 21:1–21. <https://doi.org/10.1016/j.lssr.2019.02.004>
- Kiray M, Sisman AR, Camsari UM, Evren M, Dayi A, Baykara B, Aksu I, Ates M, Uysal N (2014) Effects of carbon dioxide exposure on early brain development in rats. *Biotech Histochem* 89(5):371–383. <https://doi.org/10.3109/10520295.2013.872298>
- Kühn S, Gleich T, Lorenz RC, Lindenberger U, Gallinat J (2014a) Playing Super Mario induces structural brain plasticity: gray matter changes resulting from training with a commercial video game. *Mol Psychiatry* 19(2):265–271. <https://doi.org/10.1038/mp.2013.120>
- Kühn S, Lorenz R, Banaschewski T, Barker GJ, Büchel C, Conrod PJ, Flor H, Garavan H, Ittermann B, Loth E, Mann K, Nees F, Artiges E, Paus T, Rietschel M, Smolka MN, Ströhle A, Walaszek B, Schumann G, Gallinat J (2014b) Positive association of video game playing with left frontal cortical thickness in adolescents. *PLoS ONE* 9(3):91506. <https://doi.org/10.1371/journal.pone.0091506>
- Kühn S, Lorenz RC, Weichenberger M, Becker M, Haesner M, O’Sullivan J, Steinert A, Steinhagen-Thiessen E, Brandhorst S, Bremer T, Gallinat J (2017) Taking control! Structural and behavioural plasticity in response to game-based inhibition training in older adults. *Neuroimage* 156:199–206. <https://doi.org/10.1016/j.neuroimage.2017.05.026>
- Leasure JL, Decker L (2009) Social isolation prevents exercise-induced proliferation of hippocampal progenitor cells in female rats. *Hippocampus* 19(10):907–912. <https://doi.org/10.1002/hipo.20563>
- Lee JK, De Dios Y, Kofman I, Mulavara AP, Bloomberg JJ, Seidler RD (2019) Head down tilt bed rest plus elevated CO<sub>2</sub> as a spaceflight analog: effects on cognitive and sensorimotor performance. *Front Hum Neurosci* 13:355. <https://doi.org/10.3389/fnhum.2019.00355>
- Mammarella N (2020) The Effect of Microgravity-Like Conditions on High-Level Cognition: A Review. *Astron. Space Sci.* <https://doi.org/10.3389/fspas.2020.00006>
- McPhee JC, Charles JB (2009) Human health and performance risks of space exploration missions (SP-2009-3405). Human Research Program, National Aeronautics and Space Administration, Lyndon B. Johnson Space Center. <https://humanresearchroadmap.nasa.gov/evidence/reports/EvidenceBook.pdf>
- Mendt S, Brauns K, Friedl-Werner A, Belavy DL, Steinach M, Schlabs T, Werner A, Gunga H-C, Stahn AC (2021a) Long-term bed rest delays the circadian phase of core body temperature. *Front Physiol* 12:658707. <https://doi.org/10.3389/fphys.2021.658707>
- Mendt S, Gunga H-C, Felsenberg D, Belavy DL, Steinach M, Stahn AC (2021b) Regular exercise counteracts circadian shifts in core body temperature during long-duration bed rest. *Npj Microgravity* 7(1):1. <https://doi.org/10.1038/s41526-020-00129-1>
- Moore ST, MacDougall HG, Lesceu X, Speyer JJ, Wuyts F, Clark JB (2008) Head-eye coordination during simulated orbiter landing. *Aviat Space Environ Med.* <https://doi.org/10.3357/ASEM.2209.2008>
- Moore ST, Dilda V, Morris TR, Yungheer DA, MacDougall HG, Wood SJ (2019) Long-duration spaceflight adversely affects post-landing operator proficiency. *Sci Rep* 9(1):2677. <https://doi.org/10.1038/s41598-019-39058-9>
- Morgan C (2013) Shuttle-MIR: the United States and Russia share history’s highest stage. National Aeronautics and Space Administration
- Moser M-B, Rowland DC, Moser EI (2015) Place Cells, Grid Cells, and Memory. *Cold Spring Harb Perspect Biol* 7(2):a021808. <https://doi.org/10.1101/cshperspect.a021808>
- Mueller AD, Pollock MS, Lieblich SE, Epp JR, Galea LAM, Mistlberger RE (2008) Sleep deprivation can inhibit adult hippocampal neurogenesis independent of adrenal stress hormones. *Am J Physiol Integr Comp Physiol* 294(5):R1693–R1703. <https://doi.org/10.1152/ajpregu.00858.2007>
- Palinkas LA, Suedfeld P (2008) Psychological effects of polar expeditions. *Lancet* 371(9607):153–163. [https://doi.org/10.1016/S0140-6736\(07\)61056-3](https://doi.org/10.1016/S0140-6736(07)61056-3)
- Pereda-Pérez I, Popović N, Otalora BB, Popović M, Madrid JA, Rol MA, Venero C (2013) Long-term social isolation in the adulthood results in CA1 shrinkage and cognitive impairment. *Neurobiol Learn Mem* 106:31–39. <https://doi.org/10.1016/j.nlm.2013.07.004>
- Raven F, Meerlo P, Van der Zee EA, Abel T, Havekes R (2019) A brief period of sleep deprivation causes spine loss in the dentate gyrus of mice. *Neurobiol Learn Mem* 160:83–90. <https://doi.org/10.1016/j.nlm.2018.03.018>
- Reschke MF, Clément G (2018) Vestibular and sensorimotor dysfunction during space flight. *Curr Pathobiol Rep* 6(3):177–183. <https://doi.org/10.1007/s40139-018-0173-y>
- Roberts DR, Albrecht MH, Collins HR, Asemanni D, Chatterjee AR, Spampinato MV, Zhu X, Chimowitz MI, Antonucci MU (2017) Effects of spaceflight on astronaut brain structure as indicated on MRI. *N Engl J Med* 377(18):1746–1753. <https://doi.org/10.1056/NEJMoa1705129>
- Roberts DR, Asemanni D, Nietert PJ, Eckert MA, Inglesby DC, Bloomberg JJ, George MS, Brown TR (2019) Prolonged microgravity affects human brain structure and function. *Am J Neuroradiol* 40(11):1878–1885. <https://doi.org/10.3174/ajnr.A6249>

- Satish U, Mendell MJ, Shekhar K, Hotchi T, Sullivan D, Streufert S, Fisk WJ (2012) Is CO<sub>2</sub> an indoor pollutant? Direct effects of low-to-moderate CO<sub>2</sub> concentrations on human decision-making performance. *Environ Health Perspect* 120(12):1671–1677. <https://doi.org/10.1289/ehp.1104789>
- Scaccianoce S, Del Bianco P, Paolone G, Caprioli D, Modafferi AM, Nencini P, Badiani A (2006) Social isolation selectively reduces hippocampal brain-derived neurotrophic factor without altering plasma corticosterone. *Behav Brain Res* 168(2):323–325. <https://doi.org/10.1016/j.bbr.2005.04.024>
- Schloesser RJ, Lehmann M, Martinowich K, Manji HK, Herkenham M (2010) Environmental enrichment requires adult neurogenesis to facilitate the recovery from psychosocial stress. *Mol Psychiatry* 15(12):1152–1163. <https://doi.org/10.1038/mp.2010.34>
- Scully RR, Basner M, Nasrini J, Lam C, Hermosillo E, Gur RC, Moore T, Alexander DJ, Satish U, Ryder VE (2019) Effects of acute exposures to carbon dioxide on decision making and cognition in astronaut-like subjects. *Npj Microgravity* 5(1):17. <https://doi.org/10.1038/s41526-019-0071-6>
- Slack K, Williams T, Schneiderman J, Whitmire A, Picano J (2016) Evidence report: risk of adverse cognitive or behavioral conditions and psychiatric disorders. <https://humanresearchroadmap.nasa.gov/evidence/reports/bmed.pdf>
- Smith PF (2017) Is hippocampal neurogenesis modulated by the sensation of self-motion encoded by the vestibular system? *Neurosci Biobehav Rev* 83:489–495. <https://doi.org/10.1016/j.neubiorev.2017.09.013>
- Smith PF, Darlington CL, Zheng Y (2010) Move it or lose it—stimulation of the vestibular system necessary for normal spatial memory. *Hippocampus* 20(1):36–43. <https://doi.org/10.1002/hipo.20588>
- Stahn AC, Gunga HC, Kohlberg E, Gallinat J, Dinges DF, Kühn S (2019) Brain changes in response to long Antarctic expeditions. *N Engl J Med* 381(23):2273–2275. <https://doi.org/10.1056/NEJMc1904905>
- Stahn AC, Riemer M, Wolbers T, Werner A, Brauns K, Besnard S, Denise P, Kühn S, Gunga H-C (2020) Spatial updating depends on gravity. *Front Neural Circuits* 14:20. <https://doi.org/10.3389/fncir.2020.00020>
- Stranahan AM, Khalil D, Gould E (2006) Social isolation delays the positive effects of running on adult neurogenesis. *Nat Neurosci* 9(4):526–533. <https://doi.org/10.1038/nn1668>
- Strangman GE, Sipes W, Beven G (2014) Human cognitive performance in spaceflight and analogue environments. *Aviation space and environmental medicine* 85(10):1033–1048. <https://doi.org/10.3357/ASEM.3961.2014>
- Van Ombergen A, Jillings S, Jeurissen B, Tomilovskaya E, Ruhl RM, Rumshiskaya A, Nosikova I, Litvinova L, Annen J, Pechenkova EV, Kozlovskaya IB, Sunaert S, Parizel PM, Sinitsyn V, Laureys S, Sijbers J, Zu Eulenburg P, Wuyts FL (2018) Brain tissue-volume changes in cosmonauts. *N Engl J Med* 379(17):1678–1680. <https://doi.org/10.1056/NEJMc1809011>
- Van Ombergen A, Jillings S, Jeurissen B, Tomilovskaya E, Rumshiskaya A, Litvinova L, Nosikova I, Pechenkova E, Rukavishnikov I, Manko O, Danylichev S, Rühl RM, Kozlovskaya IB, Sunaert S, Parizel PM, Sinitsyn V, Laureys S, Sijbers J, Zu Eulenburg P, Wuyts FL (2019) Brain ventricular volume changes induced by long-duration spaceflight. *Proc Natl Acad Sci USA* 116(21):10531–10536. <https://doi.org/10.1073/pnas.1820354116>
- van Praag H, Kempermann G, Gage FH (2000) Neural consequences of environmental enrichment. *Nat Rev Neurosci* 1(3):191–198. <https://doi.org/10.1038/35044558>
- Vessel EA, Russo S (2015) Effects of reduced sensory stimulation and assessment of countermeasures for sensory stimulation augmentation. A report for NASA behavioral health and performance research: sensory stimulation augmentation tools for long duration spaceflight (NASA/TM-2015). NASA Center for AeroSpace Information
- Vivar C, van Praag H (2017) Running changes the brain: the long and the short of it. *Physiology* 32(6):410–424. <https://doi.org/10.1152/physiol.00017.2017>
- Voss MW, Vivar C, Kramer AF, van Praag H (2013) Bridging animal and human models of exercise-induced brain plasticity. *Trends Cogn Sci* 17(10):525–544. <https://doi.org/10.1016/j.tics.2013.08.001>
- Wolbers T, Hegarty M, Büchel C, Loomis JM, Büchel C, Loomis JM (2008) Spatial updating: how the brain keeps track of changing object locations during observer motion. *Nat Neurosci* 11(10):1223–1230. <https://doi.org/10.1038/nn.2189>
- Wong L, Pradhan S, Karasinski J, Hu C, Strangman G, Ivkovic V, Arsintescu L, Flynn-Evans E (2020) Performance on the robotics on-board trainer (ROBoT-r) spaceflight simulation during acute sleep deprivation. *Front Neurosci*. <https://doi.org/10.3389/fnins.2020.00697>

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.