



This postprint was originally published by Elsevier as:  
Coricelli, C., Rumiati, R. I., & Rioux, C. (2022). **Implicit and explicit safety evaluation of foods: The importance of food processing.** *Appetite*, 175, Article 106062.  
<https://doi.org/10.1016/j.appet.2022.106062s>

Supplementary material to this article is available. For more information see  
<https://hdl.handle.net/21.11116/0000-0009-56B1-E>

### **Nutzungsbedingungen:**

Dieser Text wird unter einer Deposit-Lizenz (Keine Weiterverbreitung - keine Bearbeitung) zur Verfügung gestellt. Gewährt wird ein nicht exklusives, nicht übertragbares, persönliches und beschränktes Recht auf Nutzung dieses Dokuments. Dieses Dokument ist ausschließlich für den persönlichen, nicht-kommerziellen Gebrauch bestimmt. Auf sämtlichen Kopien dieses Dokuments müssen alle Urheberrechtshinweise und sonstigen Hinweise auf gesetzlichen Schutz beibehalten werden. Sie dürfen dieses Dokument nicht in irgendeiner Weise abändern, noch dürfen Sie dieses Dokument für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, aufführen, vertreiben oder anderweitig nutzen. Mit der Verwendung dieses Dokuments erkennen Sie die Nutzungsbedingungen an.

### **Terms of use:**

This document is made available under Deposit Licence (No Redistribution - no modifications). We grant a non-exclusive, nontransferable, individual and limited right to using this document. This document is solely intended for your personal, non-commercial use. All of the copies of this documents must retain all copyright information and other information regarding legal protection. You are not allowed to alter this document in any way, to copy it for public or commercial purposes, to exhibit the document in public, to perform, distribute or otherwise use the document in public. By using this particular document, you accept the above-stated conditions of use.

### **Provided by:**

Max Planck Institute for Human Development  
Library and Research Information  
[library@mpib-berlin.mpg.de](mailto:library@mpib-berlin.mpg.de)

This postprint was originally published as:

Coricelli, C., Rumiati, R. I., & Rioux, C. (2022). Implicit and explicit safety evaluation of foods: The importance of food processing. *Appetite*, 175, Article 106062. <https://doi.org/10.1016/j.appet.2022.106062>.

## Implicit and explicit safety evaluation of foods: The importance of food processing

C. Coricelli<sup>a,b,\*</sup>, R.I. Rumiati<sup>a</sup>, C. Rioux<sup>c</sup>

<sup>a</sup> *International School for Advanced Studies (SISSA), via Bonomea, 265 - 34136, Trieste, Italy*

<sup>b</sup> *Western University, 1151 Richmond St, London, ON, N6A 3K7, Canada*

<sup>c</sup> *Max Planck Institute for Human Development, Lentzeallee, 94, 14195, Berlin, Germany*

\* Corresponding author. Department of Psychology, Western University, Western Interdisciplinary Research Building, 1151 Richmond St, London, ON, N6A 3K7, Canada.

E-mail addresses: [ccoricel@uwo.ca](mailto:ccoricel@uwo.ca) (C. Coricelli), [rumiati@sissa.it](mailto:rumiati@sissa.it) (R.I. Rumiati), [rioux@mpib-berlin.mpg.de](mailto:rioux@mpib-berlin.mpg.de) (C. Rioux).

---

Keywords: Food cognition, Go/No-Go association task, Eating behaviors, Inter-individual differences, Control

---

Supplementary material to this article is available. For more information see <https://hdl.handle.net/21.11116/0000-0009-56B1-E>

---

### Abstract

Identifying beneficial foods in the environment, while avoiding ingesting something toxic, is a crucial task humans face on a daily basis. Here we directly examined adults' implicit and explicit safety evaluations of the same foods presented with different degrees of processing, ranging from unprocessed (raw) to processed (cut or cooked). Moreover, we investigated whether individual characteristics (e.g., Body Mass Index, food neophobia and hunger) modulated their evaluations. We hypothesized that adults would associate the processed form of a food with safety more than its unprocessed form since processing techniques, which are ubiquitously applied in different cultures, often reduce the toxicity of foods, and signal previous human intervention and intended consumption. Adults (N = 109, 43 females) performed an implicit Go/No-Go association task (GNAT) online, assessing the association between safety attributes and food images differing on their degree of processing, both unfamiliar and familiar foods were used. Then each food was explicitly evaluated. Results revealed that individual self-reported characteristics affected both implicit and explicit evaluations. Individuals with excess weight and obesity had a strong and positive implicit association between processed foods and safety attributes, but explicitly rated cooked foods as the least safe overall, this latter result was found in highly neophobic individuals as well. Yet, at the explicit level, when looking at unfamiliar foods only, processed foods were rated safer than unprocessed foods by all participants. Our results are the first evidence that directly highlights the relevance of the degree of processing in food safety evaluation and suggest that thinking of the important tasks humans face regarding food selection enriches our understanding of food behaviors.

---

### 1. Introduction

Food is certainly one of the most salient and rewarding stimuli in our environment. As omnivores, we need to balance the necessity of gathering a great variety of food to ensure nutritional health against the risks posed by ingesting something toxic and so we engage in hundreds of food choices daily (Rozin & Todd, 2015; Wirt & Collins, 2009). Food is a complex and multi-attribute stimulus and several factors influencing food choices have been well-documented by previous literature, such as energy density, palatability and healthiness of the food (Hare et al., 2011; Papies et al., 2007; Toepel et al., 2009), studied along with differences related to the perceiver (*dietary habits*, Houben et al., 2010; *hunger level*, Hoefling & Strack, 2008; *body mass index* (BMI), Craeynest et al., 2005; *food neophobia*, Reilly, 2019; *personality* Nederkooft et al., 2004). However, little is known regarding evaluation of the safety of food, namely whether it is safe to consume or can lead to negative consequences such as poisoning.

#### 1.1. Food processing as a signal of food safety

How we evaluate the safety of food deserves further investigation. The environment in which our ancestors lived and our brains evolved strongly differ from modern Western circumstances (Barkow et al., 1992). Nowadays, we engage in food selection in an environment where the presence of ready-to-eat food and food cues is omnipresent and overwhelming (e.g., Cunningham & Egeth, 2018; Sanger, 2019). It rarely occurs to us to question whether a food item is safe to eat during a trip to the grocery store where foods are already packaged. Yet, such

evaluation had to be made by our ancestors each time a new type of food was encountered (Rozin & Todd, 2015; Wertz, 2019), especially new plant-based foods - an essential component of human diets across evolutionary times (Hardy & Kubiak-Martens, 2016; Ungar & Sponheimer, 2011) - as many plants contain poisonous parts (e.g., cassava root, Cashdan, 1998; see Włodarczyk et al., 2018, for a short review). The importance of identifying edible and non-toxic items appears clear when such capacity is lost as in the case of *pica*, the psychological disorder in which non-edible materials such as metal, soil, pebbles, or clay are frequently consumed, endangering patients' lives, (Sekiya et al., 2018). If *pica* disorder represents an extreme example, the adaptive problem of finding beneficial food likely shaped our brains in ways that constrain the cognitive processes underlying our food behaviors. One possibility could be to render them sensitive to a range of cues indicating safety and edibility (Wertz, 2019). For instance, infants are selective when inferring edibility from conspecifics' actions, ingesting a particular entity after observing an adult eating it, but not when the adult handles it in other ways without eating it (Wertz & Wynn, 2014a).

Diverse cues can signal which foods are safe to eat in a given environment, one of these is food processing. Based on previous work, we define food processing as actions altering the naturally occurring state of the food (e.g., a fruit growing on a tree), such as cutting, cooking or aggregation of different ingredients (Froni et al., 2013; Froni & Rumiati, 2017; Rumiati & Froni, 2016). These are the main, simple processing techniques still being used in most of human societies (e.g., Mombo, Dumat, Shahid, & Schreck, 2017). Food processing is a prior, and often necessary, step for human food consumption the world over and has been an important part of human life for millennia (Wrangham, 2009). In the *cooking hypothesis*, Wrangham argues that by providing a significant increase in the net energy gain and softening foods (Boback et al., 2007; Carmody et al., 2011), food processing activities had a key role in human evolution, leading to physical transformations in hominids (e.g., reduction of tooth and gut size and increase of brain size) and freeing our time to engage in other activities rather than chewing and hunting (Wrangham, 2009; Wrangham et al., 1999; Zink & Lieberman, 2016). In contrast, primates spend on average 8 h per day chewing raw foods (Wrangham & Conklin-Brittain, 2003). Importantly, many processing techniques reduce the toxicity of raw food items, lessening the risk of infection and poisoning (Carmody & Wrangham, 2009). For example, cassava root is a staple food in many societies, however, it cannot be eaten raw and requires complex processing techniques, such as soaking in water for several days, prior to consumption (Mombo et al., 2017).

Only a handful of studies investigating food evaluation have considered the degree of processing. Yet they reveal promising evidence that both individuals who are healthy and patients (e.g., Alzheimer dementia) perceive processed foods differently to unprocessed foods (Aiello et al., 2018; Vignando et al., 2018; see Froni & Rumiati, 2017 for a review). For instance, Coricelli et al. (2019a) found significant differences in reaction times when participants performed a food vs. non-food categorization task, with participants being significantly faster in categorizing processed foods as foods compared to the unprocessed foods. This behavioral advantage in recognition was supported by differences in the brain responses recorded using electroencephalography. Around 130 ms (ms) post-stimulus presentation, greater activation in visual areas in response to processed foods was shown, supporting the perceptual relevance of highly rewarding stimuli when compared to less rewarding stimuli (here unprocessed foods; Coricelli et al., 2019a; see also Toepel et al., 2009). In this early time window, the human brain also responds differently to edible (foods) and non-edible (objects and rotten foods) items (around 100 ms post-stimulus onset) (Tsourides et al., 2016). This time is strikingly fast if compared to other effects related to food found in windows between 400 and 800 ms post-stimulus onset such as appetitive conditioning (Blechert et al., 2016). Behavioral data revealed that, adults and even children see processed foods as human-made objects that bear markers of previous intervention and require less work prior consumption, and see unprocessed foods as more naturally occurring (Froni et al., 2013; Girgis & Nguyen, 2020). Moreover, developmental data revealed that infants and children might view cues of processing as a signal of food safety. Infants display attenuated wariness behaviors towards novel processed plant foods (e.g., unfamiliar fruits and vegetables cut into pieces) compared to novel unprocessed whole plants with fruits (Rioux & Wertz, 2021). Children assign negative properties (e.g., "This food makes you throw up") less often to processed foods compared to unprocessed foods (Foinant et al., 2021a). Such evidence of a distinction between processed and unprocessed foods present at this early stage in life seems to reveal that the level of processing plays an important role in the development of core building blocks of humans' food cognition. The safety signal conveyed by the level of processing likely arises both from the fact that processing techniques often reduce the toxicity of raw foods and because cues of processing reveal that another person has already interacted with a candidate food and has deemed it to be edible.

However, in this line of work, participants were not asked directly about the *safety* of the foods and were always presented with processed foods that differ from unprocessed foods on at least one of the following variables, leading to potential confounds: *food types* (e.g., meat vs. vegetables, Coricelli et al., 2019a; pear vs. star fruit, Foinant et al., 2021a), *caloric density* (e.g., cookie vs. fruit, Girgis & Nguyen, 2020) or *overall shape and color* (cut papaya vs. whole plant with fruits; Rioux & Wertz, 2021). Whether individuals perceive the *same* food differently depending on their degree of food processing, remains to be investigated.

## 1.2. Implicit and explicit evaluation of food

Food evaluation is known to be influenced by both explicit and implicit factors (Marty et al., 2017; Monnery-Patris & Chamberon, 2020). Explicit evaluations are assumed to influence responses described as conscious or controlled, while implicit evaluations are assumed to influence responses described as non-conscious and uncontrolled (Marty et al., 2017; Perugini, Costantini, Richetin, & Zogmaister, 2015).

Investigations of explicit evaluations of food usually consist of direct self-reports asking participants to rate different dimensions on a scale (i.e., liking, wanting, willingness to pay, frequency of consumption; Roefs & Jansen, 2002; Finlayson et al., 2007; Romero et al., 2018). Individuals' explicit evaluation of food regarding its caloric content (high vs. low) received most of the attention, with participants often reporting negative evaluations of high-calorie (or high-fat) palatable foods, when compared to the low-calorie (low-fat) unpalatable counterparts (Czyzewska & Graham, 2008; Houben et al., 2010; Papies et al., 2009; Roefs & Jansen, 2002; Rothmund et al., 2007). One of the few studies to investigate explicit evaluation of food regarding its level of processing found that participants view processed foods as more ready-to-eat and requiring less work prior to consumption (Froni et al., 2013), yet they did not assess *perceived safety* of the different foods. Despite their widespread use, such measures are vulnerable to biases such as the social desirability bias in which participants seek to present a positive image of themselves (e.g., underestimating the liking of junk foods, Cerri et al., 2019; Czyzewska et al., 2011). Moreover, such measures do not capture well responses influenced by unconscious factors (Monnery-Patris & Chamberon, 2020).

To capture such responses and reduce the social desirability bias, implicit evaluations of food have been studied using indirect behavioral measures, such as the Implicit Association Test (IAT; Greenwald et al., 1998), the Affective Priming Task (Fazio, 1995) or more recently, the Go/No-Go Association Task (GNAT; Nosek & Banaji, 2001). These computerized tasks require fast responses to sequences of stimuli (words, images, etc.) presented for a few hundred milliseconds. The main assumption is that participant responses are facilitated (better accuracy and faster reaction times) if consecutive stimuli, or stimuli which require the same response pattern (i.e., share the same key for

response) are closely associated. As with explicit measures, mainly high vs. low calorie/fat evaluations of the foods have been investigated. In sharp contrast with the explicit food evaluation literature, most of the work reports implicit positive attitude towards high-fat palatable foods (Lamote et al., 2004; Papies et al., 2009; Roefs et al., 2005). Indeed, explicit and implicit measures often do not converge, and both measures should be assessed when investigating food evaluation and choices (see Hofmann et al., 2005 for a meta-analysis; Hoefling & Strack, 2008). Regarding implicit food evaluation depending on its degree of processing, using the IAT and controlling for caloric content, Coricelli and colleagues found that participants held a positive implicit association between both processed and unprocessed food and positivity (i.e., words associated with positivity such as joy, peace and holidays; Coricelli et al., 2019b). To our knowledge, this is the only study investigating implicit evaluation of food regarding its degree of food processing, yet the association between food processing and the *safety* attribute was not directly assessed.

### 1.3. The present experiment

In line with the existing findings, the aim of the present research was to investigate for the first time (i) implicit and explicit safety evaluations of food depending on its degree of processing and (ii) whether individual characteristics modulated participants' responses. First, we predicted that individuals would hold a positive implicit association between processed foods and safety, evaluating the processed forms safer than the unprocessed forms. Our primary focus was on implicit association because it rarely occurs to us today to question whether a food is safe to eat or not, but it was a recurrent task over evolutionary time, and therefore, natural selection likely favored cognitive systems sensitive to cues of food safety. Second, we predicted that explicit evaluations would diverge partially from implicit evaluations, because in our modern food environment processed foods are often high in calories/fat and are viewed as "junk foods".

For the implicit evaluation, we used the Go/No-Go association task (GNAT) where participants had to press the spacebar in the presence of a target concept (Go trials) and refrain from pressing it if the presented items belong to other concepts (No-Go trials). The GNAT has considerable methodological advances over the more common IAT (Williams & Kaufmann, 2012). The primary one is its ability to assess associations between a single concept (e.g., food) and attributes (e.g., safety and toxicity) without having to measure the relative associations between two concepts (e.g., food and non-food) and attributes. The GNAT has been successfully used in several experiments (e.g., Ashford et al., 2018; Buhlmann et al., 2011), notably with food stimuli (Gerdan & Kurt, 2020; Mas et al., 2020; Spence & Townsend, 2007) and shows good psychometric qualities, such as internal consistency and reliability (Bar-Anan & Nosek, 2014; Williams & Kaufmann, 2012). Individual characteristics such as hunger level, food neophobia, dietary habits and BMI were measured because previous work has shown its influence on food evaluation (Coricelli et al., 2019a; Foinant et al., 2021a; Houben et al., 2010; Mas et al., 2020).

Adults completed the GNAT and explicit ratings task on the same- colored images depicting foods differing only on their degree of processing: (i) unprocessed fruits and vegetables, (ii) the *same* foods cut into pieces and (iii) cooked into a puree. Cut foods were included since previous work has shown that infants' neophobic behaviors are attenuated when they are confronted by cut foods compared to whole foods (Rioux & Wertz, 2021). Moreover, children are less prone to assign negative properties to cut foods compared to whole foods (Foinant et al., 2021a; Lafraire et al., 2020). Food cutting is also a common component of more complex food processing techniques and a clear cue of human intervention. In addition, cooked foods were included because cooking food is a more advanced technique and often efficiently reduces the toxicity of the raw foods (e.g., Mombo et al., 2017; see also Carmody & Wrangham, 2009). We chose fruit and vegetable stimuli because it is an important class of foods with the potential to be poisonous and even deadly to humans (Hardy & Kubiak-Martens, 2016; Henry et al., 2014; Mithöfer & Boland, 2012; Włodarczyk et al., 2018). Importantly, we also chose fruits and vegetables because consumption of these foods is notoriously low and below recommended intakes (Hall et al., 2009). Therefore, it is of crucial importance to shed light on the mechanisms underpinning the evaluation of fruits and vegetables to pave the way towards effective interventions for promoting the adoption of healthy eating behaviors.

## 2. Materials and methods

### 2.1. Participants

Participants were 109 Italian adults (43 females) with normal or corrected-to-normal vision. Age of the participants was between 18 and 34 years ( $M = 24.4$ ,  $SD = 4.0$ ) and their Body Mass Index (BMI,  $\text{kg}/\text{m}^2$ ) ranged from 17 to 37 ( $M = 23.7$ ,  $SD = 3.9$ ). This sample size was chosen based on a power analysis with pilot participants (with the *powerSim* function from the *simr* package in R; Green & MacLeod, 2016), assuming a small effect size in the implicit Go/No-Go Association task (difference in reaction time of 10 ms between experimental conditions, as in studies using similar design, e.g., Mas et al., 2020) and a power of 80%. Data of additional 14 subjects were collected but excluded based on participants' performance on the implicit task (see *data preparation and statistical analysis* section below).


### 2.2. Procedure

Due to the COVID-19 pandemic, the study was conducted online and participants were recruited through the platform Prolific (Prolific, Oxford, UK; <https://www.prolific.co/>). Each participant provided informed consent prior to beginning the experiment. The study conformed to the Declaration of Helsinki and was approved by SISSA's Ethic Committee. The study lasted approximately 40 min and comprised of three separate phases with the following constant order: (a) assessment of the implicit evaluations using the GNAT; (b) explicit ratings of the food stimuli used in the GNAT; and (c) questionnaires on participants' characteristics. Stimulus presentation and registration of responses for the GNAT was controlled by PsychoPy 3.0 (Peirce et al., 2019; retrieved from <http://www.psychopy.org/>) and ran through the online repository and launch platform Pavlovia (<http://www.pavlovia.org/>) (the GNAT has already been successfully implemented in online settings, e.g., Ashford et al., 2018). Stimulus presentation and registration of responses for the explicit ratings and questionnaires was controlled by Qualtrics (<http://www.qualtrics.com/>).

Participants received a compensation of 5 GBP after completion of the study. As this is usual practice in online settings, to ensure participants' attention throughout the study, they were informed they could receive a 1.5 GBP extra payment if they did not answer randomly during the GNAT (criterion: more than 60% of correct answers), and if they passed two attentional checks added to the questionnaires (i.e., "respond never to this question").

#### 2.2.1. Implicit evaluations: Go/No-Go association task (GNAT)

The GNAT (Nosek & Banaji, 2001) was chosen in order to investigate participants' implicit evaluations of different foods in terms of safety. In this task, participants must (a) respond to target stimuli (Go trials) by pressing the spacebar on a computer keyboard while (b) withholding their response to distractor stimuli (No-Go trials). The response deadline was set to 600 ms with an inter-trial interval of 100 ms consisting of a white screen (see Fig. 1a). Participants were asked to commit as few errors as possible in order to avoid a speed-accuracy trade-off (Zimmerman, 2011, p. 2344), in which participants commit too many errors while trying to respond quickly. These times were selected, based on piloting the task, to balance the need for time pressure while keeping an error rate that could vary between participants. Feedback was given



b)

	Familiar		Unfamiliar	
<b>Raw whole foods (Condition P0)</b>				
<b>Raw cut foods (Condition P1)</b>				
<b>Cooked pureed foods (Condition P2)</b>				

**Fig. 1.** Example of trials in the condition P0 and Block *Safety* (panel a) and example of food stimuli used in the Go/No-Go association task (GNAT) (panel b). In the Go-trials a whole food [carrot] and a safety-related word [Harmless]; in the No-Go trials a kitchen utensil [ice-cream spoon] and a toxicity-related word [Infected]. **Note.** In panel a) exemplar trials of the GNAT in the condition P0 and Block *Safety* (raw whole foods presented). Participants had to press the spacebar in the Go trials and refrain from pressing the spacebar in the No-Go trials. Images were presented on the screen for 600 ms and a blank screen was presented for 100 ms as Inter-Trial Interval (ITI). In panel b) exemplar food stimuli used in the GNAT with the three different degrees of processing.

after error trials, with a red “X” appearing below the stimulus for 150 ms.

The GNAT began with four training blocks (see Table 1 and SM Video S1 Training + Condition P0 for a demonstration of the GNAT). In the training blocks participants had to respond (i.e., press the space bar) to only one stimuli category (either foods, kitchen utensils, words related to safety, or words related to toxicity respectively in each of the four training blocks). There were 6 trials in each training block. Following the training blocks, participants completed three conditions consisting of two experimental blocks each: Block *Safety* and Block *Toxicity* (see Table 1 and SM Video S1). In Block *Safety*, participants had to respond to food images and words related to safety and refrain from responding when viewing kitchen utensils and words related to toxicity (see Fig. 1a). In Block *Toxicity*, participants had to respond to food images and words related to toxicity and refrain from responding when viewing kitchen utensils and words related to safety. There were 96 trials in each of the experimental blocks (24 foods, 24 kitchen utensils, 24 words related to safety, 24 words related to toxicity, see Table 1) presented in a pseudo-randomized order, with the constraint that an image trial (food or kitchen utensil) was followed by a word trial (see Fig. 1a). One separate Block *Safety* and Block *Toxicity* per condition were created based on our experimental manipulation of the food stimuli namely: raw whole foods (condition P0), raw cut foods (condition P1) and cooked pureed foods (condition P2), see Table 1 for an overview of the GNAT

Condition	Press the spacebar	Don't press the spacebar	Number of trials (1:1 ratio)
Training 1	Food	Utensil	6
Training 2	Utensil	Food	6
Training 3	Safety word	Toxicity word	6
Training 4	Toxicity word	Safety word	6
Condition P0 – Block <i>Safety</i> (Raw whole foods)	Food or Safety word	Utensil or Toxicity word	96 (8 practice trials)
Condition P0 – Block <i>Toxicity</i> (Raw whole foods)	Food or Toxicity word	Utensil or Safety word	96 (8 practice trials)
Condition P1- Block <i>Safety</i> (Raw cut foods)	Food or Safety word	Utensil or Toxicity word	96 (8 practice trials)
Condition P1- Block <i>Toxicity</i> (Raw cut foods)	Food or Toxicity word	Utensil or Safety word	96 (8 practice trials)
Condition P2 -Block <i>Safety</i> (Cooked pureed foods)	Food or Safety word	Utensil or Toxicity word	96 (8 practice trials)
Condition P2 -Block <i>Toxicity</i> (Cooked pureed foods)	Food or Toxicity word	Utensil or Safety word	96 (8 practice trials)

**Table 1** Overview of the implicit Go/No-Go association task. **Note.** 1:1 ratio indicates an equal number of Go and No-Go trials.

structure. Participants' accuracy and latency to press the spacebar (Reaction Time/RT) reflect the ease to associate the target concept (different food) to the attribute (safety vs. toxicity) in the two different blocks. Order of presentation of Block *Safety* and Block *Toxicity* within each condition, as well as condition order, were counterbalanced across participants.

Supplementary data related to this article can be found at <https://hdl.handle.net/21.11116/0000-0009-56B1-E>.

The experimental stimuli for the three different conditions consisted of pictures depicting eight fruits and vegetables, further subdivided in familiar (*carrot, tomato, peach* and *apple*) and unfamiliar items (*Buddha hand citron, pink guava, jackfruit* and *starfruit*, see Fig. 1b). In condition P0 the fruits and vegetables were presented raw and whole (henceforth *whole*). In condition P1, they were presented raw and cut into familiar shapes (e.g., slices for tomato, quarters for peach) (henceforth *cut*). In condition P2, they were presented cooked and pureed, without any container (henceforth *cooked*). Before each experimental condition began, a short description of the foods depicted in the condition was presented to participants (e.g., 'you will now be presented with pictures of food raw and whole', see SM Video S1). Pictures of 24 different kitchen utensils (8 per condition) matched for overall shape, size and color of the food pictures (e.g., ice-cream spoon, lemon squeezer, pan) were used for distractor picture stimuli. All pictures were color photographs in jpeg-format (1920 × 1080 pixels), selected from online free from copyright image search, then modified (i.e., cropped) using GNU Image Manipulation Program (GIMP; <http://www.gimp.org/>) and placed on a white background (see SM Section 1.1 and Fig. S1 for a complete presentation of picture stimuli). Note that we used a distractor concept (i.e., kitchen utensils that are not obviously dangerous like knives), to keep some difficulty to the task, but the GNAT performs robustly without a distractor concept (Nosek & Banaji, 2001). In addition, the GNAT is not affected significantly by the relatedness of the target and distractor concepts, namely whether they are from related concepts (e.g., fruits and vegetables) or from distant concepts (e.g., foods and clothes) (Nosek & Banaji, 2001).

Finally, we used eight Italian words, four associated to safety (*safe [sicuro], immaculate [immacolato], pure [puro]* and *harmless [innocuo]*) and four associated to toxicity (*infected [infetto], poisoned [dannoso], damaging [avvelenato]* and *dangerous [pericoloso]*). The complete set of images and a demonstration video of the GNAT, showing the training blocks and condition P0, are available in the Supplementary Materials (SM) and on OSF: <https://osf.io/snrgk/>.

Stimuli selection was based on the results of a pilot study. A separate set of 29 Italian healthy participants (20 females) aged between 20 and 33 years ( $M = 26.8$ ,  $SD = 3.9$ ) were asked to rate 12 pictures of foods depicted in their whole, cut and cooked forms, on the following dimensions: *familiarity* (whole foods), *degree of processing* and *degree of cooking* (cut and cooked foods), *similarity* (between whole and cut/cooked form of the same food), followed, in brackets, by labels at the extremes of the scale:

- Familiarity: 'How familiar are you with the depicted food?' ('Very unfamiliar' [0] – 'Very familiar' [100]);
- Degree of processing: 'How prepared (transformed by human intervention for eating purposes) is the depicted food?' ('Not at all prepared' [0] – 'Very prepared' [100]);
- Degree of cooking: 'How cooked is the depicted food?' ('Not at all cooked' [0] – 'Very cooked' [100]);
- Similarity: 'How similar are the depicted foods?' ('Very dissimilar' [0] – 'Very similar' [100]).

For each rating a Visual Analog Scale (VAS) was positioned below the picture which measured 1920 × 1080 pixels. Picture presentation order was randomized across participants. Moreover, participants had to rate 11 words *a priori* related to safety and 12 words *a priori* related to danger on the following dimensions, followed, in brackets, by labels at the extremes of the scale:

- Association to safety/danger: 'How much is the present word associated with the concept of safety/danger?' ('Not related to safety/danger at all' [0] – 'Very related to safety/danger' [100]);
- Familiarity: 'How familiar are you with the presented word?' ('Not familiar at all' [0] – 'Very familiar' [100]);
- Valence: 'How negative/positive is the present word?' ('Very negative' [0] – 'Very positive' [100]).

The number of syllables and word length were also calculated for each word. For the analysis, VAS distances were converted to a scale ranging from 0 to 100, although this was not explicitly displayed to the participants.

We selected food pictures for inclusion in our GNAT that significantly differed in familiarity, with our selected familiar stimuli being significantly more familiar than the unfamiliar foods ( $t(3) = 39.02$ ,  $p < .001$ ). Our selected cooked foods were significantly viewed as more processed than our cut foods ( $t(3) = 14.12$ ,  $p < .001$ ). Our cooked foods were also significantly viewed as less similar to whole foods than our cut foods ( $t(3) = 7.65$ ,  $p < .001$ ). Selected positive words were highly associated with safety and negative words were highly associated with toxicity, with no significant difference in the strength of association ( $t(3) = 1.11$ ,  $p = .35$ ). Moreover, our words differed on valence with the words associated to safety being significantly more positive than the words associated with toxicity ( $t(3) = 16.94$ ,  $p < .001$ ), our words did not differ on familiarity ( $t(3) = 1.17$ ,  $p = .33$ ), nor on word length ( $t(3) = 0.87$ ,  $p = .50$ ) and number of syllables ( $t(3) = 0.68$ ,  $p = .55$ ).

### 2.2.2. Explicit evaluations: food pictures ratings

After having completed the implicit task, participants rated all the 24 food images presented in the task on different dimensions by selecting their response along a VAS. The VAS was positioned below the images which measured 1920 × 1080 pixels, and picture presentation order was randomized across participants. For the analysis, VAS distances were converted to a scale ranging from 0 to 100, although this was not explicitly displayed to the participants (see Fig. S2 in the SM section 1.2). Participants rated the images of whole foods, cut foods and cooked foods on the following five dimensions: *safety*, *valence*, *wanting*, *healthiness* and *frequency of consumption*, followed, in brackets, by labels at the extremes of the scale:

- Safety: 'How safe is ingesting the food represented in the picture?' ('Not safe at all' [0] – 'Very safe' [100]);
- Valence: 'How negative/positive is the content of the picture for you?' ('Very negative' [0] – 'Very positive' [100]);
- Wanting: 'How much do you want to eat the food represented in the picture at this moment?' ('Don't want to eat it now' [0] – 'Want to eat it now' [100]);
- Healthiness value: 'How healthy is the food represented in the picture?' ('Not healthy' [0] – 'Very healthy' [100]);
- Frequency of consumption: 'How often do you eat the food represented in the picture?' ('I never eat this food' [0] – 'I eat this food very often' [100]).

### 2.2.3. Questionnaires on participants' characteristics

After having completed the implicit and explicit tasks, participants reported their characteristics (i.e., age, gender, height and weight) and hunger level using a VAS. Due to the COVID-19 pandemic these characteristics were self-reported by participants, instead of measured in person in the lab, but several studies indicate that in-lab measured and self-reported anthropometric data (e.g., height and weight) are strongly positively correlated and those self-reports can be a valid method of collecting anthropometric data (Bonn et al., 2013; Lassale et al., 2013; Huang et al., 2020; Pursey et al., 2014; van der Laan et al., 2022). For the analysis, VAS distances were converted to a scale ranging from 0 to 100,



although this was not explicitly displayed to the participants. Two separate questions regarding hunger level were presented investigating *pre-task hunger level* and *post-task hunger level*, followed, in brackets, by labels at the extremes of the scale:

- (a) How hungry were you before beginning the study? (“Not at all [0] – A lot [100]”);
- (b) How hungry are you at the moment? (“Not at all [0] – A lot [100]”).

Moreover, participants had to report “How many hours ago did you have your last meal?” by inserting the number of hours.

A questionnaire regarding participants’ dietary habits was then completed; participants had to report whether they had food allergies (i.e., gluten, lactose, nuts, other), food intolerances (i.e., gluten, lactose, sulfites, fructose, other), how would they define their diet (i.e., omnivore, vegetarian, vegan, other) and whether they had other dietary restrictions other than caused by allergies or food intolerances - for example, ones based on personal, ethical or religious reasons. Finally, participants had to fill in the standardized and validated questionnaire investigating novel food avoidance: the Italian translation of the Food Neophobia Scale (FNS) (for the complete set of Italian questions see Proserpio et al., 2016), which consists of 10 statements regarding individual tendencies to approach or avoid unfamiliar foods (exemplar statements: “I like foods from different countries” or “I am very picky”). Participants had to report their agreement on each statement on a 7-point Likert-like scale (“Strongly disagree” – “Strongly agree”). For the analysis, each answer was then numerically coded with high scores indicating high food neophobia (possible range 10–70).

### 2.3. Data preparation and statistical analysis

Data and scripts used for statistical analysis are available on OSF: <https://osf.io/snrqk/>. All analyses were performed in the R environment (version 3.6.3; <http://www.r-project.org/>). To investigate whether participants hold implicit and explicit associations between processed foods and safety, participants’ answers have been analyzed using a Linear Mixed-effects Model approach (LMM, Bates et al., 2015) using the *lmer* function (*lme4* package; [cran.r-project.org/web/packages/lme4/index.html](https://cran.r-project.org/web/packages/lme4/index.html)). This method exploits the inter-trial variability by analyzing each data point per participant to investigate the modulation of different factors. Such models are called mixed since they include *fixed* effects, which represent population-level effects which should persist across experiments and *random* effects, which vary across level of grouping factors (i. e., participants) (Brown, 2021; Meteyard & Davies, 2020).

To investigate implicit associations, we tested separate LMM models with average Reaction Times (RTs) and Error rates during the GNAT as dependent variables, because meaningful information about task performance can be found in both average reaction times and errors due to a potential speed-accuracy trade-off (Nosek & Banaji’s, 2001). The main assumption in GNATs is that participant responses are facilitated (better accuracy and faster reaction times) if stimuli that require the same response pattern (i.e., share the same key for response – here space bar) are closely associated. In the present research, we expected faster reaction times and better accuracy in Blocks *Safety*, and especially in the condition P2, when participants had to respond simultaneously to food images and safety words. To investigate explicit associations, we tested separate LMM models with each Explicit ratings during the food picture rating task as dependent variables (Safety, Valence, Wanting, Healthiness and Frequency of consumption). In total nine LMM models were tested (see details below). In all LMM models, participants served as a random effect to account for shared variances within subjects (see also Aiello et al., 2018; Coricelli et al., 2019b; for a similar approach). The fit of each LMM model was tested by comparing it to the fit of its null model (containing no predictors) through the Aikake Information Criterion (AIC) values. Furthermore, for the full models, analysis of deviance was first inspected using the *Wald chi square test* and then post hoc comparisons were performed using the *emmeans* function (R packages *car* and *emmeans*). Multiple comparisons were controlled for using the Tukey’s method.

#### 2.3.1. Go/No-Go association task: Reaction times (RTs)

Before running the LMM models we checked whether some RTs data should be excluded from analysis. Following Nosek & Baniji’s (2001) recommendations, RTs were examined to determine if any participants had more than 10% trials with responses under 300 ms, as they reflect stimulus anticipation and random responding (Buhmann et al., 2011), or overall accuracy below 60%. Fourteen participants were excluded based on these criteria, leaving a sample of 109 participants. Next, we removed from analysis the first 8 trials (out of 96) in each of the 6 experimental blocks (Block *Safety* and Block *Toxicity* in each condition, see Table 1) as they could be regarded as practice trials. Further, as recommended, erroneous RTs to distractor trials (i.e., trials with kitchen utensils) were not included in the RTs analysis, so that only target trials (i.e., trials with foods) were kept. This deletion occurred because the distractors are considered noise. Finally, RTs inferior to 300 ms were excluded from analysis based on previous literature (Buhmann et al., 2011) and the actual distribution of our data ( $M = 428.3$  ms,  $SD = 77.7$ ). The task did not register RTs greater than 600 ms, therefore there were no extreme slow RTs to discard.

After the data cleaning step, we computed a first LMM model with the remaining cleaned RTs data as our dependent variable and the following fixed effects: Block (Block *Safety* and Block *Toxicity*), Condition (P0, P1 and P2), the five covariates Food familiarity (familiar and unfamiliar), BMI (continuous variable), FNS (continuous variable), Pre- task Hunger levels (continuous variable – henceforth Hunger levels), and Explicit Safety ratings (continuous variable), as well as the interaction between Block, Condition and covariates (see SM Section 2.1.1. and Table S1 for the complete model). A second identical LMM model with Explicit Valence ratings, instead of Explicit Safety ratings, was also tested (see SM Section 2.1.2. and Table S3 for the complete model).

#### 2.3.2. Go/No-Go association task: Errors

In order to analyze participants’ error distributions both target trials (i.e., trials with foods - Go trials) and distractor trials (i.e., trials with kitchen utensils - No-Go trials) were included in this analysis (on the sample of 109 participants and with practice trials excluded). Following Mas et al. (2020) and Gerdan & Kurt (2020), from these trials the number of *misses* (incorrect responses in Go trials), and *false alarms* (incorrect responses in No-Go trials), were extracted from the data. Misses are omission errors, where participants do not detect the signal (for instance in Block *Safety*, participants miss to press the space bar while a food or a safety word is displayed on the screen). False alarms are commission errors, where participants believe the signal is present while it is actually not (for instance in Block *Safety*, participants press the space bar while a kitchen utensil or a toxicity word is displayed on the screen). Both types of errors are important to take into account. Overall, a large number of misses indicate low capacity to detect the signal during the task and leads to low accuracy, while a large number of false alarms indicates a liberal decision bias (e.g., participant tending to press the spacebar no matter whether the signal is present or not). None of the 109 participants included based on the previous RTs analysis had an overall error rate >40% (indicating low accuracy, Nosek & Baniji’s 2001). Therefore, no further participants were excluded for this analysis.

After this data preparation, a third LMM model with the miss rates as a dependent variable was tested, and the following fixed effects: Block (Block *Safety* and Block *Toxicity*), Condition (P0, P1 and P2), the three participant covariates BMI (continuous variable), FNS (continuous variable) and Hunger levels (continuous variable), as well as the interaction between Block, Condition and covariates. A fourth LMM identical to the previous one with false alarm rates as a dependent variable was also tested (see SM Section 2.2. and Table S5 for the complete models).

### 2.3.3. Food pictures ratings

Explicit ratings (Safety, Valence, Wanting, Healthiness and Frequency of consumption) from the 109 participants have been analyzed by converting the VAS scale ratings to a scale ranging from 0 to 100. One separate LMM model with each of the Explicit ratings as dependent variables was tested, with the following fixed effects: Degree of processing (whole food, cut food, cooked food), the four covariates: Food familiarity (familiar and unfamiliar), BMI (continuous variable), FNS (continuous variable), and Hunger levels (continuous variable), as well as the interaction between Degree of processing and the covariates. This analysis resulted in five separate LMM models (see SM Section 2.3. for the complete models).

## 3. Results

### 3.1. Implicit evaluations: Go/No-Go association task

#### 3.1.1. Reaction times (RTs)

Mean participants' RT to the task was 438 ms (SD = 54.3), which is similar to previous average reaction times to food stimuli in GNATs (e.g., Gerdan & Kurt, 2020; Mas et al., 2020). The results of our LMM models with RTs in response to target food stimuli as a dependent variable are now described. In the full model, Block, Condition, the five covariates (Food familiarity, BMI, FNS, Hunger levels, and Explicit Safety ratings), as well as the interaction between Block, Condition and covariates were modeled as fixed effects. The full model had a better fit than the null model (containing no predictors) as shown with a significant drop in AIC ( $\chi^2(35) = 327.20$ ,  $p < .001$ , marginal  $R^2 = 0.026$ , conditional  $R^2 = 0.24$ ). Significant main and interaction effects are presented in Table 2 (see SM Section 2.1.1. and Table S1 for the description of all the main and interaction effects tested in the full model).

In the full model the following main effects were significant: Block Condition and Explicit Safety ratings (see Table 2). The main effect of Explicit Safety ratings indicated that, as participants rated the pictures as safer, they were faster in pressing the space bar in Go trials (i.e., trials with foods) during the GNAT.

The only 2-way interaction that was significant was the Condition\*Food familiarity interaction (see Table 2). Post hoc comparisons revealed that participants were significantly faster in responding to familiar foods compared to unfamiliar foods but only in condition P0 where the foods were whole. Overall, participants were slower in responding to cut foods compared to the other foods (see SM Section 2.1.1. and Table S2 for a full description of the contrasts revealed from the interaction effect between Condition and Food familiarity).

Finally, only the 3-way interaction of Block\*Condition\*BMI was significant (see Fig. 2 and Table 2). Post hoc comparisons revealed that participants were slower in Block *Toxicity* compared to Block *Safety* only in response to cooked foods (condition P2) and as a function of their BMI ( $b = -1.19$ ,  $SE = 0.39$ ,  $z = -3.06$ ,  $p = .027$ ). None of the other comparisons were significant.

We found a similar pattern of results for our second LMM model with Explicit Valence ratings, instead of Explicit Safety ratings (for the description of all the main and interaction effects tested in the full model see Table S3 in the SM section 2.1.2.).

#### 3.1.2. Errors

Mean participants' miss rate to the task was 17.74% (SD = 21.07). This indicated that the task was overall easy for the participants. Mean false alarm rate was 9.36% (SD = 13.03), indicating that overall participants showed a conservative criterion (e.g., tendency to refrain from pressing the spacebar for Go trials). Results with the miss and false alarm rates as dependent variables of the LMM models are now described (see SM section 2.2. for a complete description of the models).

In the full model with the miss rates as the dependent variable, Block, Condition, the three participant covariates BMI, FNS and Hunger levels, as well as the interaction between Block, Condition and covariates were modeled as fixed effects. The full model had a better fit than the null model (containing no predictors) as shown with a significant drop in AIC ( $\chi^2(23) = 39.47$ ,  $p = .018$ , marginal  $R^2 = 0.046$ , conditional  $R^2 = 0.63$ ). Significant main effects are presented in Table 3 (see SM Section 2.2. and Table S5 for the description of all the main and interaction effects tested in the full model).

The main effect of Condition was significant (see Table 3), with participants being more accurate in responding to cooked foods (P2) compared to cut foods ( $b = 3.00$ ,  $SE = 1.01$ ,  $z = 2.97$ ,  $p = .0089$ ). The main effect of Block was also significant (see Table 3), with participants being more accurate in Block *Safety* ( $b = 2.95$ ,  $SE = 0.83$ ,  $z = 3.57$ ,  $p < .001$ ). The main effect of BMI approached significance (see Table 3), indicating that as their BMI increased, participants tended to have fewer misses ( $r = -0.14$ ,  $p < .001$ ).

In the full model with the false alarm rates as the dependent variable, Block, Condition, the three participant covariates BMI, FNS and Hunger levels, as well as the interaction between Block, Condition and covariates were modeled as fixed effects. The full model had a better fit than the null model (containing no predictors) as shown with a significant drop in AIC ( $\chi^2(23) = 35.21$ ,  $p = .049$ , marginal  $R^2 = 0.046$ , conditional  $R^2 = 0.23$ ). Significant main effects are presented in Table 3 (see SM Section 2.2. and Table S5 for the description of all the main and interaction effects tested in the full model).

The main effect of Condition was significant (see Table 3), with participants being more conservative (i.e., making fewer false alarms) in responding to cooked foods (P2) compared to cut foods ( $b = 2.90$ ,  $SE = 0.91$ ,  $z = 3.20$ ,  $p = .0042$ ). The 2-way interaction Block\*FNS was also significant (see Table 3). In Block *Toxicity*, as their neophobia increased, participants were more liberal (i.e., making more false alarms  $r = 0.013$ ,  $p = .018$ ). In Block *Safety*, participants' numbers of false alarms did not change depending of their neophobia levels ( $r = 0.021$ ,  $p = .70$ ).

### 3.2. Explicit evaluations: food pictures ratings

Mean participants' Explicit Safety rating was 76 (SD = 24), indicating that overall participants judged all foods quite safe to eat. Results with the Explicit Safety ratings in response to food stimuli as a dependent variable of the LMM model is now described. In the full model, Degree of processing, the four covariates (Food familiarity, BMI, FNS, and Hunger levels), as well as the interaction between Degree of processing and the covariates were modeled as fixed effects. The full model had a better fit than the null model (containing no predictors) as shown with a significant drop in AIC ( $\chi^2(14) = 4919.50$ ,  $p < .001$ , marginal  $R^2 = 0.29$ , conditional  $R^2 = 0.50$ ). Significant main and interaction effects are presented in Table 4 (see SM Section 2.3.1. and Table S6 for the description of all the main and interaction effects tested in the full model).

The following main effects were significant: Degree of processing, Food familiarity and FNS (see Table 4). All the 2-way interactions were significant: Degree of processing\*Food familiarity, Degree of processing\*BMI, Degree of processing\*FNS and Degree of processing\*Hunger levels (see Table 4).


Post hoc comparisons revealed that in the 2-way Degree of processing\*Food familiarity interaction (see Fig. 3), participants reported familiar foods as significantly safer compared to unfamiliar foods for

Effect	$\chi^2(df)$	p
Block	63.09(1)	<.001
Condition	175.96(2)	<.001
Safety ratings	10.79(1)	.0010
Condition * Food familiarity	14.25(2)	.0010
Block * Condition * BMI	9.94(2)	.0070

**Table 2** ANOVA significant results for linear mixed-effect model for participants' Reaction Times in the Go/No-Go association task with Explicit Safety ratings.

**Note.**  $\chi^2$ -values for effects using Type II Wald chi-square tests.





**Fig. 2.** Participants' Reaction Times in the Go/No-Go association task depending on Condition, Block and BMI. **Note.** Linear regression lines with 95% confidence intervals. Condition P0: whole foods; Condition P1: cut foods; Condition P2: cooked foods. Block *Safety*: food is associated with safety words; Block *Toxicity*: food is associated with toxicity words. In condition P2 only, as their BMI increased, participants associated cooked foods with safety more than toxicity, as they were significantly faster in Block *Safety* compared to Block *Toxicity*.

Effect	$\chi^2(df)$	p
<b>LMM with Misses</b>		
Condition	9.21(2)	.010
Block	12.71(1)	<.001
BMI	3.53(1)	.060
<b>LMM with False alarms</b>		
Condition	10.54(2)	.0051
Block*FNS	4.81	.028

**Table 3**

ANOVA significant results for linear mixed-effect models for participants' Errors rates in the Go/No-Go association task.


**Note.**  $\chi^2$ -values for effects using Type II Wald chi-square tests. FNS = Participants' food neophobia scores.

Effect	$\chi^2(df)$	p
Degree of processing	456.28(2)	<.001
Food familiarity	3519.50(1)	<.001
FNS	18.41(1)	<.001
Degree of processing * Food familiarity	1694.44(2)	<.001
Degree of processing * BMI	83.92(2)	<.001
Degree of processing * FNS	187.31(2)	<.001
Degree of processing * Hunger levels	56.19(2)	<.001

**Table 4**

ANOVA significant results for linear mixed-effect model for Participants' Explicit Safety ratings.

**Note.**  $\chi^2$ -values for effects using Type II Wald chi-square tests. FNS = Participants' food neophobia scores.



**Fig. 3.** Participants' Explicit Safety ratings depending on Degree of processing and Food familiarity. **Note.** Raw means and standard errors of participants' explicit ratings of safety. For familiar foods, whole and cut foods were rated the safest. For unfamiliar foods, cooked foods were rated the safest. For cooked foods, safety ratings did not differ as a function of food familiarity.

whole foods ( $b = 30.06$ ,  $SE = 0.54$ ,  $z = 55.68$ ,  $p < .001$ ), and cut foods ( $b = 24.97$ ,  $SE = 0.54$ ,  $z = 45.94$ ,  $p < .001$ ) but not for the cooked foods. For familiar foods, cooked foods were rated significantly less safe than whole foods ( $b = -20.99$ ,  $SE = 0.53$ ,  $z = -39.10$ ,  $p < .001$ ) and cut foods ( $b = -19.72$ ,  $SE = 0.54$ ,  $z = -36.63$ ,  $p < .001$ ), that did not differ. On the contrary, for unfamiliar foods, cooked foods were rated the safest (compared to whole foods:  $b = 8.32$ ,  $SE = 0.54$ ,  $z = 15.39$ ,  $p < .001$ ; compared to cut foods:  $b = 4.44$ ,  $SE = 0.54$ ,  $z = 8.20$ ,  $p < .001$ ). Cut foods were also rated safer than whole foods ( $b = 3.88$ ,  $SE = 0.55$ ,  $z =$

7.11,  $p < .001$ ).

Post hoc comparisons for the other interaction effects revealed that as their BMI increased, participants significantly rated cooked foods as less safe compared to both whole foods ( $b = -0.78$ ,  $SE = 0.09$ ,  $z = -7.87$ ,  $p < .001$ ) and cut foods ( $b = -0.78$ ,  $SE = 0.09$ ,  $z = 7.93$ ,  $p < .001$ , see Fig. 4 panel A).

Similar results were found as participants' food neophobia increased (whole vs. cooked food:  $b = -0.36$ ,  $SE = 0.03$ ,  $z = -11.90$ ,  $p < .001$ , cut vs. cooked food:  $b = -0.36$ ,  $SE = 0.03$ ,  $z = -11.8$ ,  $p < .001$ , see Fig. 4 panel B) and their hunger levels decreased (whole vs. cooked food:  $b = -0.78$ ,  $SE = 0.01$ ,  $z = -5.96$ ,  $p < .001$ , cut vs. cooked food:  $b = -0.09$ ,  $SE = 0.01$ ,  $z = -6.88$ ,  $p < .001$ , see Fig. 4 panel C).

As the main focus of the present manuscript was to investigate safety evaluations, in SM sections 2.3.2. - 2.3.5 we listed and summarized the results of the other LMM models with, respectively, Explicit ratings of Valence, Wanting, Healthiness and Frequency of consumption as dependent variables of the models with Degree of processing, the four covariates (Food familiarity, BMI, FNS, and Hunger levels), as well as the interaction between Degree of processing and the covariates modeled as fixed effects. Overall similar patterns were found across models, with the notable exception that for Healthiness ratings, cooked foods, regardless of their familiarity, were rated less healthy than whole and cut foods. In addition, participants rated more positively and reported to eat more frequently familiar whole foods compared to familiar cut foods, while safety ratings for these two foods did not differ.


#### 4. Discussion

Identifying beneficial foods in the environment is a task we all face daily. A handful of studies revealed that, from an early age, individuals view processed foods more positively than unprocessed foods (Aiello et al., 2018; Coricelli et al., 2019a, Coricelli et al., 2019b; Foinant et al., 2021a; Feroni & Rumiati, 2017; Girgis & Nguyen, 2020; Rioux & Wertz, 2021), showing the importance of taking into account the degree of processing when investigating human food behaviors. Therefore, the aim of the present study was to expand on this limited line of work and directly investigate, for the first time, whether individuals differently evaluate the safety of a food depending on its degree of processing, both at an implicit and explicit level. Overall, we found evidence that individuals evaluate the cooked form of a food safer than its less processed forms, albeit with some important modulations depending on participants' and foods' characteristics.

A Go/No-Go association task was employed to investigate the implicit evaluations, given the advantage of this measure compared to other implicit measures (e.g., IAT) in being able to assess a single target concept (Nosek & Banaji, 2001). First, participants' miss error rates showed that, when food was associated with words related to toxicity (Block *Toxicity*), participants were less accurate in their responses, meaning that they were less willing to say the signal (food) was present. The result shows that, overall fruits and vegetables were implicitly associated with a positive attribute, and this is also revealed by the overall high explicit ratings for all the foods. Because we only used low-calorie foods, without comparing them with high-calorie foods, our results can't be directly compared to previous work on implicit attitudes (e.g., Roefs et al., 2005). Instead, our results add to the existing literature, which has shown that in healthy adults, when compared to non-food (e.g., kitchen utensils), in healthy adults, food overall is associated with positive attributes, even at the implicit level (using the affective priming task, Czyzewska & Graham, 2008; using the IAT, Coricelli et al., 2019b).

In line with our first prediction, all participants were faster to respond to cooked foods compared to the other foods, made fewer misses and false alarms in respond to cooked foods, and certain individuals evaluated the cooked form of a food safer than its less processed forms. Indeed, as their self-reported BMI increased (overall range 17–37) individuals were faster in associating cooked foods with safety (Block *Safety*) compared to toxicity (Block *Toxicity*) (see Fig. 2). In sum, individuals with higher BMIs especially associated the cooked form of a food with safety, and none of the individuals associated the less processed forms of a food with higher levels of safety. These results are in line with previous findings showing a greater and faster activation in adult brains in response to processed foods compared to unprocessed foods (Coricelli et al., 2019a), but go further by suggesting that certain individuals represent differently the *same* food depending on its degree of processing, evaluating the processed forms *safer* than the unprocessed forms.

As in Coricelli and colleagues (2019a), in the present research all participants responded faster to cooked foods, but the finding that only individuals with higher self-reported BMIs associated more the cooked form of a food with safety compared to toxicity was less expected.



**Fig. 4.** Participants' Explicit Safety ratings depending on Degree of processing and BMI (panel A), Food Neophobia (panel B) or Hunger level (panel C). **Note.** Linear regression lines with 95% confidence intervals. FNS = Participants' food neophobia scores. High Food neophobia scores indicate high food neophobia (range 10–64,  $M = 29.51$ ,  $SD = 12.83$ ). High Hunger levels indicate high hunger before the task (range 10–100,  $M = 29.13$ ,  $SD = 28.46$ ).

Individuals with excess weight and obesity tended to be better at the task resulting in a smaller number of miss errors in our task (i.e., incorrect responses for Go trials regardless of blocks) and previous GNAT studies reported shorter RTs to food images (e.g., Gerdan & Kurt, 2020; Mas et al., 2020; but see Osimo et al., 2019 for an opposite finding). However, we believe that our findings speak against a mere easiness account because individuals with excess weight and obesity were actually slower to respond to cooked food when it was associated with toxicity (Condition P2, Block *Toxicity*). It is possible that a stronger positive implicit association between safety and cooked foods in certain individuals leads to a higher consumption of these types of foods, resulting in weight gain in modern circumstances because highly processed foods are often also high in fat and calories. For instance, Marty and colleagues (2017) directly assessed whether implicit attitudes towards foods can predict actual eating behaviors in children and found that they consumed more of a food they previously implicitly rated high on a hedonic level. However, in our study, neither participants' wanting of the foods nor frequency of consumption changed depending on their BMI, but we did not have highly processed foods in our stimuli set. An important outstanding question for future work is then whether an implicit association between safety and processed foods leads to a higher consumption of these particular foods.

After they completed the implicit task, participants completed an explicit rating task on the same images used in the GNAT, given that food evaluation is known to be influenced by both implicit and explicit factors (Marty et al., 2017; Monnery-Patris & Chamberon, 2020). Overall food pictures were rated positively. Participants might have inferred from the familiar foods (e.g., tomato and apple) that, in this experiment, they were presented with edible items, hence the relatively high explicit ratings in all of the presented foods. It is also possible that the explicit ratings were relatively high because fruits and vegetables are considered healthy foods and eating these products leads to positive long-term consequences on health. Nonetheless we found interesting differences, notably depending on the level of processing. In line with our second prediction that implicit and explicit evaluations would partially diverge, participants rated cooked foods less safe than the other foods, especially people with high food neophobia and people with excess weight and obesity, who had a strong positive association between these processed foods and safety attributes at the implicit level (GNAT RTs results). The explicit results converge with previous literature showing that individuals often report more negative evaluations of highly processed foods, which are also high in calories, compared to the low-calorie counterparts (Czyzewska & Graham, 2008; Houben et al., 2010; Papies et al., 2009; Roefs & Jansen, 2002; Rothmund et al., 2007).

It is common to find implicit and explicit results, which go in opposite directions (Hoefling & Strack, 2008; Hofmann et al., 2005). Previous literature has explained this phenomenon in light of two models. On one hand, the *Dual Attitudes Model* (Wilson et al., 2000) states that for a given object different evaluations can coexist (i.e., holding an implicit positive evaluation and an explicit negative evaluation for ice cream). On the other, the *Reflective-Impulsive Model* of behavior regulation by Strack and Deutsch (2004) proposes the existence of two separate systems, an impulsive and a reflective one, which produce different behavioral outcomes depending on whether the decision is based on motivational orientations (e.g., food palatability) or based on knowledge (e.g., long-term health consequences). A prediction derived from Strack and Deutsch's model is that, when control resources are reduced (e.g., by time pressure or hunger), the functioning of the reflective system is limited, and impulsive behaviors are increased (Czyzewska et al., 2011; Friese et al., 2008), as would be expected in implicit tasks. In line with the prediction of the *Reflective-Impulsive Model*, in the present study, individuals explicitly rated cooked foods safer as their hunger increased (i.e., when control resources were reduced), converging with their implicit evaluations.

In the present study, it is not surprising that people with high neophobia rated foods overall less positively than their counterparts. Neophobia is thought to be a protective strategy against the risk of ingesting potentially poisonous items (Dovey et al., 2008; Lafraire et al., 2016; Reilly, 2019; Rioux, 2019; Rozin & Todd, 2015) and neophobic individuals assign more negative properties to foods compared to individuals with less neophobic disposition (e.g., Foinant et al., 2021a; 2021b). Accordingly, during our GNAT, neophobic participants were more willing to associate food with toxicity. In general, neophobic individuals tend to show wariness when presented with unknown foods, as these foods may be harmful once ingested. In the present study, neophobic participants might have rated cooked foods even less positively because these foods were overall less familiar to participants (as indicated by the Frequency of consumption ratings). Regarding participants with excess weight and obesity, it is possible that these individuals, who might be concerned about weight gain, explicitly rated cooked foods less positively because nowadays processed foods are often high in calories and industrialized pureed foods often contain additives (e.g., sugar, salt, conservatives). Thus, we need to consume them in small quantities in order to avoid negative long-term health consequences. Accordingly, our results revealed that all foods were rated less healthy when they were cooked into a puree (as indicated by the Explicit Healthiness ratings), and participants were less willing to eat these foods overall (as indicated by the Explicit Wanting ratings). It remains an open question to what degree our current findings would generalize to other cultures, including non-WEIRD populations (Henrich et al., 2010) where industrialized processed foods are less common and foraging for wild plant-food resources is still practiced.

Nevertheless, the further examination of the explicit ratings, revealed that all individuals rated cooked foods safer than its less processed forms, in a particular situation: when they were confronted with unfamiliar foods. Indeed, for unfamiliar foods, which participants could not recognize easily, participants reported lower values of safety for whole foods compared to cut foods and then lower values of safety for cut foods compared to cooked foods (see Fig. 3). The familiarity status of each food (familiar vs. unfamiliar) was defined based on the pilot study and validated in the main study, as participants ate more often the familiar foods overall compared to the unfamiliar ones, indicated by the Frequency of consumption ratings. The result that cooked unfamiliar foods were explicitly rated safer than its less processed forms converges with the pattern found with the GNAT, showing that under a state of uncertainty, the degree of processing is used as a cue for better safety in food evaluation. Further, the degree of processing seems to influence food choices as well because participants were more willing to eat the foods they rated safer, as indicated by the Wanting ratings that parallel the Safety ratings.

It is important to note that, in the case of familiar foods, cooked foods were actually rated less safe than raw foods. Indeed, whole and cut familiar foods (e.g., whole and cut tomato) were rated the safest by the participants and they rated cooked familiar foods (e.g., cooked tomato puree) less safe. In our modern food environment, it is clear that familiar foods like tomatoes, carrots, peaches and apples are safe to eat. During a trip to the grocery store, it would never occur to us to question the edibility of such familiar foods. On the other hand, because industrialized processed foods might contain unhealthy additives, it is more likely that individuals would question the safety attribute of familiar processed foods. In addition, in the present study participants might have been familiar with the packaging of processed foods, while we presented them with images of plain purees without any containers. Accordingly, participants reported that they more often ate the familiar unprocessed foods compared to the cooked counterparts. In sum, it appears that participants rated the foods they knew the best the safest, namely familiar raw foods, but when they were confronted with unfamiliar foods, they used the degree of processing to make finer-grained safety evaluation. The association we found in adults of familiar foods with positive properties has also been recently found in children by Foinant et al. (2021a). Children would generalize positive properties such as

“gives strength” to familiar foods and negative properties such as “gives nausea” to unfamiliar foods when presented with various types of fruits. Importantly, in this study the degree of processing was also taken into account (though including only cut foods) and the results showed that children significantly generalized more positive properties to cut foods compared to whole foods in the case of unfamiliar foods (Foinant et al., 2021a), converging with our results with unfamiliar foods. Taken together, Foinant and colleagues’, along with our findings, suggest that when both adults and children have no prior knowledge on foods, cues of food processing afford more positive properties, even at the explicit level.

#### 4.1 Future directions and limitations

In summary, the findings from the present study show that adults evaluate the cooked form of a food, both at the implicit and explicit level, safer than its less processed forms, albeit with some important modulations depending on participants’ (i.e., BMI, food neophobia and hunger) and foods’ characteristics (i.e., familiarity). The results add to the growing literature highlighting the role of cues indicating processing in the evaluation of food safety (Foinant et al., 2021a; Rioux & Wertz, 2021) and converge with research showing that the degree of processing has a key role in food cognition (Aiello et al., 2018; Coricelli et al., 2019a; Foroni & Rumiati, 2017; Girgis & Nguyen, 2020). The relative difference in safety evaluation (i.e., rating two fruits rather safe, but the processed form safer than the unprocessed form), has the potential to reveal what drives the choices subjects will make when they are confronted with several food items, which can have important health implications in a context of an environment where food cues are omnipresent and overwhelming.

However, there is much that remains to be investigated. Such as what kind of processing action is needed to trigger a safety signal. In the present study, cooked foods seemed to be the foods most associated with safety (compared to cut foods). It suggests that more advanced processing techniques might be needed to trigger a safety signal. Indeed, these complex techniques (e.g., cooking, soaking in hot water) are often needed to reduce the toxicity of raw foods (Carmody & Wrangham, 2009; Mombo et al., 2017) while cutting a food does not alter its chemical properties. It is important to note that, as a manipulation check, our pilot study confirmed that individuals considered the cooked foods as both more processed and cooked compared to the whole and cut versions of the same food. Remarkably, the procedure of cutting a food is a clear cue of human intervention (Foroni et al., 2013), which could signal intended consumption and modulate *edibility* evaluations (i.e., food vs. non-food). However, the cut food might require further processing actions before being safe to consume, thereby affecting *safety* evaluations. In line with this idea, Foinant et al. (2021b) found that when children performed a food vs. non-food categorization task, they more often miscategorized cut non-food items as foods compared to whole non-food items. Future studies can further examine the association between cut foods and edibility, by using in a similar GNAT with words associated with edibility rather than safety. Another important and related future line of research is to include other types of processing actions and processed foods to further investigate the relative importance of cues of previous human interaction (e.g., cutting, grinding) and chemical alteration (e.g., cooking, pickling, frying) in evaluations of the safety of a food. Finally, it is also an open question whether our results would hold in younger populations. Childhood is a critical period to examine food evaluation as many foods are initially unfamiliar to children. Yet very few studies have investigated implicit food evaluation in this population, probably due notably to the high demands of implicit tasks. Recently a version of the IAT has been adapted for children as young as four years of age (DeJesus et al., 2020). Therefore, future studies should examine whether children also hold implicit or explicit associations between safety and food processing.

While our current results are consistent with the proposal that cues of processing can act as a signal of food safety, we acknowledge that the present study suffers from several limitations. First, due to the COVID-19 pandemic this research was conducted online and therefore relied on self-reports to collect sensitive data such as weight and height. Despite some evidence that self-reports can be a valid method of collecting anthropometric data (Bonn et al., 2013; Lassale et al., 2013; Huang et al., 2020; Pursey et al., 2014), height and weight data might be especially prone to reporting bias, resulting in a possible underestimation of BMI in our sample (e.g., Pursey et al., 2014). And further testing in the lab is required to assess the robustness of our findings. Due to the online set-up we were also not able to measure participants’ actual eating behaviors toward the foods they saw in the task. A behavioral food choice measure must be included in future studies to investigate, in detail, how individuals’ evaluation of food safety predicts actual eating behaviors and to what extent food processing alone can explain cravings for industrialized processed foods that are often both highly processed and high in calories/fat. In this future line of work, it would also be important to assess participants’ expectation of the taste of the foods (of both familiar and unfamiliar foods), something that we did not measure here and that is known to influence wanting and choices of foods (Rangel, 2013). Second, while the Explicit Frequency of consumption ratings give an indication of participants’ familiarity with the foods, we did not directly ask participants to name the different foods. A categorization task performed after the implicit and explicit tasks could have provided a better indication of the role of recognition in safety evaluation. In the present study, we chose cooked pureed foods as the most processed foods to match the visual complexity of the whole, cut, and cooked foods (i.e., having all images composed by one single color and element). However, one aspect which might have affected both familiarity and reported frequency of consumption is that rarely pureed foods would be presented without packaging. This is a potential limitation of the study which future studies, including a vaster continuum of processed foods, could address.

## 5. Conclusion

As a first-of-its-kind study, here we present results showing how humans use cues of processing to assign different safety attributes to unprocessed and processed foods. It is of crucial importance to shed light on the mechanisms underpinning the evaluation of foods to identify important mechanisms that can increase acceptance of healthy foods such as fruits and vegetables and pave the way towards effective interventions for promoting the consumption of such healthy food products. This is especially important in populations with high neophobia (mainly children), who have less varied diets, eat less fruits and vegetables, and assign more negative properties to these healthy foods (Proserpio et al., 2018; Foinant et al., 2021a; 2021b). Our findings provide a critical first step toward future work that could develop such interventions.

### Ethical statement

The research was conducted via an online experimental protocol approved by the Ethics Committee of the International School of Advanced Studies of Trieste (SISSA). The work was carried out in accordance with the international guidelines for research involving humans established in the Declaration of Helsinki. All subjects provided informed consent prior to starting the experiment.

### Declaration of interest

The authors declare no conflict of interest.

### Declaration of funding source

The present study was funded by the International School for

## Authors contributions

CC and CR conceived the original idea of the study, CR designed the experimental task, CC and CR collected the data. CC and CR have conducted data analysis and interpreted the results. CC, CR wrote the manuscript, RIR, CC and CR reviewed and finalized the manuscript.

## Declaration of competing interest

The authors declare no conflicts of interest with this publication.

## Acknowledgments

We are grateful to SISSA for funding the present study with funds (UA.00.NEUR.CI53) awarded to Raffaella Ida Rumiati. We thank J'érémie Lafraire and members of the Max Planck Research Group 'Naturalistic Social Cognition', for helpful discussions and comments on previous versions of this manuscript.

## Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://hdl.handle.net/21.11116/0000-0009-56B1-E>.

## References

- Aiello, M., Vignando, M., Foroni, F., Pergola, G., Rossi, P., Silveri, M. C., & Rumiati, R. I. (2018). Episodic memory for natural and transformed food. *Cortex*, *107*, 13–20.
- Ashford, R. D., Brown, A. M., & Curtis, B. (2018). Substance use, recovery, and linguistics: The impact of word choice on explicit and implicit bias. *Drug and Alcohol Dependence*, *189*, 131–138.
- Bar-Anan, Y., & Nosek, B. A. (2014). A comparative investigation of seven indirect attitude measures. *Behavior Research Methods*, *46*(3), 668–688.
- Barkow, J. H., Cosmides, L., & Tooby, J. (Eds.). (1992). *The adapted mind: Evolutionary psychology and the generation of culture*. USA: Oxford University Press.
- Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*, *67*(1), 1–48.
- Blechert, J., Testa, G., Georgii, C., Klimesch, W., & Wilhelm, F. H. (2016). The Pavlovian craver: Neural and experiential correlates of single trial naturalistic food conditioning in humans. *Physiology & Behavior*, *158*, 18–25.
- Boback, S. M., Cox, C. L., Ott, B. D., Carmody, R., Wrangham, R. W., & Secor, S. M. (2007). Cooking and grinding reduces the cost of meat digestion. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, *148*(3), 651–656.
- Bonn, S. E., Troll Lagerros, Y., & Baller, K. (2013). How valid are Web-based self-reports of weight? *Journal of Medical Internet Research*, *15*(4), e52.
- Brown, V. A. (2021). An introduction to linear mixed-effects modeling in R. *Advances in Methods and Practices in Psychological Science*, *4*(1), Article 2515245920960351.
- Buhlmann, U., Teachman, B. A., & Kathmann, N. (2011). Evaluating implicit attractiveness beliefs in body dysmorphic disorder using the Go/No-go Association Task. *Journal of Behavior Therapy and Experimental Psychiatry*, *42*(2), 192–197.
- Carmody, R. N., Weintraub, G. S., & Wrangham, R. W. (2011). Energetic consequences of thermal and nonthermal food processing. *Proceedings of the National Academy of Sciences*, *108*(48), 19199–19203.
- Carmody, R. N., & Wrangham, R. W. (2009). The energetic significance of cooking. *Journal of Human Evolution*, *57*(4), 379–391.
- Cashdan, E. (1998). Adaptiveness of food learning and food aversions in children. *Social Science Information*, *37*(4), 613–632.
- Cerri, J., Thøgersen, J., & Testa, F. (2019). Social desirability and sustainable food research: A systematic literature review. *Food Quality and Preference*, *71*, 136–140.
- Coricelli, C., Foroni, F., Osimo, S. A., & Rumiati, R. I. (2019b). Implicit and explicit evaluations of foods: The natural and transformed dimension. *Food Quality and Preference*, *73*, 143–153.
- Coricelli, C., Toepel, U., Notter, M. L., Murray, M. M., & Rumiati, R. I. (2019a). Distinct brain representations of processed and unprocessed foods. *European Journal of Neuroscience*, *50*(8), 3389–3401.
- Craeynest, M., Crombez, G., De Houwer, J., Deforche, B., Tanghe, A., & De Bourdeaudhuij, I. (2005). Explicit and implicit attitudes towards food and physical activity in childhood obesity. *Behaviour Research and Therapy*, *43*(9), 1111–1120.
- Cunningham, C. A., & Egeth, H. E. (2018). The capture of attention by entirely irrelevant pictures of calorie-dense foods. *Psychonomic Bulletin & Review*, *25*(2), 586–595.
- Czyzewska, M., & Graham, R. (2008). Implicit and explicit attitudes to high-and low- calorie food in females with different BMI status. *Eating Behaviors*, *9*(3), 303–312.
- Czyzewska, M., Graham, R., & Ceballos, N. A. (2011). Explicit and implicit attitudes to food. *Handbook of Behavior, Food and Nutrition*, 673–692.
- DeJesus, J. M., Gelman, S. A., & Lumeng, J. C. (2020). Children's implicit food cognition: Developing a food Implicit Association Test. *Cognitive Development*, *54*, Article 100889.
- Dovey, T. M., Staples, P. A., Gibson, E. L., & Halford, J. C. G. (2008). Food neophobia and "picky/fussy" eating in children: A review. *Appetite*, *50*(2–3), 181–193.
- Fazio, R. H. (1995). Attitudes as object-evaluation associations: Determinants, consequences, and correlates of attitude accessibility. *Attitude Strength: Antecedents and Consequences*, *4*, 247–282.
- Finlayson, G., King, N., & Blundell, J. E. (2007). Liking vs. wanting food: Importance for human appetite control and weight regulation. *Neuroscience & Biobehavioral Reviews*, *31*(7), 987–1002.
- Foinant, D., Lafraire, J., & Thibaut, J.-P. (2021a). Strength or nausea? Children's reasoning about the health consequences of food consumption. *Frontiers in Psychology*, *12*, 1119.
- Foinant, D., Lafraire, J., & Thibaut, J.-P. (2021b). Fatal errors in the food domain: children's categorization performance and strategy depend on both food processing and neophobic dispositions. *Preprint*. <https://doi.org/10.13140/RG.2.2.2204.97929>
- Foroni, F., Pergola, G., Arqiris, G., & Rumiati, R. I. (2013). The FoodCast research image database (FRIDa). *Frontiers in Human Neuroscience*, *7*.
- Foroni, F., & Rumiati, R. I. (2017). Food perception and categorization: From food/no- food to different types of food. In H. D. Cohen, & C. Lefebvre (Eds.), *Handbook of categorization in cognitive science* (2nd ed., pp. 271–287). Elsevier.
- Friese, M., Hofmann, W., & Wanke, M. (2008). When impulses take over: Moderated predictive validity of explicit and implicit attitude measures in predicting food choice and consumption behaviour. *British Journal of Social Psychology*, *47*, 397–419.
- Gerdan, G., & Kurt, M. (2020). Response inhibition according to the stimulus and food type in exogenous obesity. *Appetite*, *150*, 104651.
- Girgis, H., & Nguyen, S. P. (2020). Grown or made? Children's determination of the origins of natural versus processed foods. *Cognitive Development*, *56*, Article 100887.
- Green, P., & MacLeod, C. J. (2016). SIMR: an R package for power analysis of generalized linear mixed models by simulation. *Methods in Ecology and Evolution*, *7*, 493–498.
- Greenwald, A. G., McGhee, D. E., & Schwartz, J. L. (1998). Measuring individual differences in implicit cognition: The implicit association test. *Journal of Personality and Social Psychology*, *74*(6), 1464.
- Hall, J. N., Moore, S., Harper, S. B., & Lynch, J. W. (2009). Global variability in fruit and vegetable consumption. *American Journal Of Preventive Medicine*, *36*(5), 402–409. e5.
- Hardy, K., & Kubiak-Martens, L. K. (2016). *Wild harvest: Plants in the hominin and pre-agrarian human worlds*. Oxbow books.
- Hare, T. A., Malmaud, J., & Rangel, A. (2011). Focusing attention on the health aspects of foods changes value signals in vmPFC and improves dietary choice. *Journal of Neuroscience*, *31*(30), 11077–11087.
- Henrich, J., Heine, S. J., & Norenzayan, A. (2010). The weirdest people in the world? *Behavioral and Brain Sciences*, *33*(2–3), 61–135.
- Henry, A. G., Brooks, A. S., & Piperno, D. R. (2014). Plant foods and the dietary ecology of Neanderthals and early modern humans. *Journal of Human Evolution*, *69*, 44–54.
- Hoefling, A., & Strack, F. (2008). The tempting effect of forbidden foods. High calorie content evokes conflicting implicit and explicit evaluations in restrained eaters. *Appetite*, *51*(3), 681–689.
- Hofmann, W., Gawronski, B., Gschwendner, T., Le, H., & Schmitt, M. (2005). A meta-analysis on the correlation between the Implicit Association Test and explicit self-report measures. *Personality and Social Psychology Bulletin*, *31*(10), 1369–1385.
- Houben, K., Roefs, A., & Jansen, A. (2010). Guilty pleasures. Implicit preferences for high calorie food in restrained eating. *Appetite*, *55*(1), 18–24.
- Huang, L., Chen, W., Renzaho, A. M. N., & Hall, B. J. (2020). Validation of obesity status based on self-reported data among Filipina and Indonesian female migrant domestic workers in Macao (SAR), China. *International Journal of Environmental Research and Public Health*, *17*(16), 5927.
- van der Laan, L. N., Papias, E. K., Ly, A., & Smeets, P. A. M. (2022). Examining the neural correlates of goal priming with the NeuroShop, a novel virtual reality fMRI paradigm. *Appetite*, *170*, Article 105901.
- Lafraire, J., Rioux, C., Giboreau, A., & Picard, D. (2016). Food rejections in children: Cognitive and social/environmental factors involved in food neophobia and picky/ fussy eating behavior. *Appetite*, *96*, 347–357.
- Lafraire, J., Rioux, C., Hamaoui, J., Girgis, H., Nguyen, S., & Thibaut, J.-P. (2020). Food as a borderline domain of knowledge: The development of domain-specific inductive reasoning strategies in young children. *Cognitive Development*, *56*, Article 100946.
- Lamote, S., Hermans, D., Baeyens, F., & Felen, P. (2004). An exploration of affective priming as an indirect measure of food attitudes. *Appetite*, *42*(3), 279–286.
- Lassale, C., Péneau, S., Touvier, M., Julia, C., Galan, P., Hercberg, S., & Kesse-Guyot, E. (2013). Validity of web-based self-reported weight and height: results of the Nutrinet-Santé study. *Journal of medical internet research*, *15*(8), Article e2575.
- Marty, L., Chamberon, S., Bournez, M., Nicklaus, S., & Monnery-Patris, S. (2017). Comparison of implicit and explicit attitudes towards food between normal- and overweight French children. *Food Quality and Preference*, *60*, 145–213.
- Mas, M., Brindisi, M.-C., Chabanet, C., & Chamberon, S. (2020). Implicit food odour priming effects on reactivity and inhibitory control towards foods. *PLoS One*, *15*(6), Article e0228830.
- Meteyard, L., & Davies, R. A. (2020). Best practice guidance for linear mixed-effects models in psychological science. *Journal of Memory and Language*, *112*, Article 104092.
- Mithöfer, A., & Boland, W. (2012). Plant defense against herbivores: Chemical aspects. *Annual Review of Plant Biology*, *63*, 431–450.
- Mombo, S., Dumat, C., Shahid, M., & Schreck, E. (2017). A socio-scientific analysis of the environmental and health benefits as well as potential risks of cassava production and consumption. *Environmental Science and Pollution Research*, *24*(6), 5207–5221.



- Mombo, S., Dumat, C., Shahid, M., & Schreck, E. (2017). A socio-scientific analysis of the environmental and health benefits as well as potential risks of cassava production and consumption. *Environmental Science and Pollution Research*, 24(6), 5207–5221.
- Monnery-Patris, S., & Chambaron, S. (2020). Added-value of indirect methods to assess the relationship between implicit memory and food choices in adult consumers as well as in children. *Current Opinion in Food Science*, 33, 14–20.
- Nederkorn, C., Van Eijs, Y., & Jansen, A. (2004). Restrained eaters act on impulse. *Personality and Individual Differences*, 37(6), 1651–1658.
- Nosek, B. A., & Banaji, M. R. (2001). The go/no-go association task. *Social Cognition*, 19(6), 625–666.
- Osimo, S. A., Korb, S., & Aiello, M. (2019). Obesity, subliminal perception and inhibition: Neuromodulation of the prefrontal cortex. *Behaviour Research and Therapy*, 119, Article 103408.
- Papies, E. K., Stroebe, W., & Aarts, H. (2007). Pleasure in the mind: Restrained eating and spontaneous hedonic thoughts about food. *Journal of Experimental Social Psychology*, 43(5), 810–817.
- Papies, E. K., Stroebe, W., & Aarts, H. (2009). Who likes it more? Restrained eaters' implicit attitudes towards food. *Appetite*, 53(3), 279–287.
- Peirce, J. W., Gray, J. R., Simpson, S., MacAskill, M. R., Hochenberger, R., Sogo, H., ... Lindeløv, J. K. (2019). PsychoPy2: experiments in behavior made easy. *Behavior Research Methods*. <https://doi.org/10.3758/s13428-018-01193-y>
- Perugini, M., Costantini, G., Richetin, J., & Zanna, M. P. (2015). *Implicit association tests, then and now. Behavior-based assessment: Going beyond self-report in the personality, affective, motivation, and social domains*. Göttingen.
- Proserpio, C., Laureati, M., Bertoli, S., Battezzati, A., & Pagliarini, E. (2016). Determinants of obesity in Italian adults: The role of taste sensitivity, food liking, and food neophobia. *Chemical Senses*, 41(2), 169–176.
- Proserpio, C., Laureati, M., Invitti, C., & Pagliarini, E. (2018). Reduced taste responsiveness and increased food neophobia characterize obese adults. *Food Quality and Preference*, 63, 73–79.
- Pursey, K., Burrow, S. T. L., Stanwell, P., & Collins, C. E. (2014). How accurate is web-based self-reported height, weight, and body mass index in young adults? *Journal of Medical Internet Research*, 16(1), e4.
- Rangel, A. (2013). Regulation of dietary choice by the decision-making circuitry. *Nature Neuroscience*, 16(12), 1717–1724.
- Reilly, S. (2019). *Food neophobia* (1st ed.). Elsevier.
- Rioux, C. (2019). Food neophobia in childhood. In H. L. Meiselman (Ed.), *Handbook of eating and drinking: Interdisciplinary perspectives* (pp. 413–432). Springer Press.
- Rioux, C., & Wertz, A. E. (2021). Avoidance of plant foods in infancy. *Developmental Psychology*, 57(5), 609.
- Roefs, A., Herman, C. P., MacLeod, C. M., Smulders, F. T. Y., & Jansen, A. (2005). At first sight: How do restrained eaters evaluate high-fat palatable foods? *Appetite*, 44(1), 103–114.
- Roefs, A., & Jansen, A. (2002). Implicit and explicit attitudes toward high-fat foods in obesity. *Journal of Abnormal Psychology*, 111(3), 517.
- Romero, C. A., Compton, M. T., Yang, Y., & Snow, J. C. (2018). The real deal: Willingness-to-pay and satiety expectations are greater for real foods versus their images. *Cortex*, 107, 78–91.
- Rothmund, Y., Preuschhof, C., Bohner, G., Bauknecht, H. C., Klingebiel, R., Flor, H., & Klapp, B. F. (2007). Differential activation of the dorsal striatum by high-calorie visual food stimuli in obese individuals. *NeuroImage*, 37(2), 410–421.
- Rozin, P., & Todd, P. M. (2015). The evolutionary psychology of food intake and food choice. In D. M. Buss (Ed.), *The handbook of evolutionary psychology* (2nd ed., 1 pp. 183–205). Hoboken, NJ: Wiley.
- Rumiati, R. I., & Foroni, F. (2016). We are what we eat: How food is represented in our mind/brain. *Psychonomic Bulletin & Review*, 23(4), 1043–1054.
- Sänger, J. (2019). Can't take my eyes off you—How task irrelevant pictures of food influence attentional selection. *Appetite*, 133, 313–323.
- Sekiya, K., Mori, S., & Otomo, Y. (2018). Coin pica-induced gastric perforation resulting from ingestion of 1,894 coins. 8 kg in total: Case report and review of published works. *Acute Medicine and Surgery*, 5(2), 177.
- Spence, A., & Townsend, E. (2007). Predicting behaviour towards genetically modified food using implicit and explicit attitudes. *British Journal of Social Psychology*, 46(2), 437–457.
- Strack, F., & Deutsch, R. (2004). Reflective and impulsive determinants of social behavior. *Personality and Social Psychology Review: An Official Journal of the Society for Personality and Social Psychology, Inc.* 8(3), 220–247.
- Toepel, U., Knebel, J. F., Hudry, J., le Coutre, J., & Murray, M. M. (2009). The brain tracks the energetic value in food images. *NeuroImage*, 44(3), 967–974.
- Tsourides, K., Shariat, S., Nejadi, H., Gandhi, T. K., Cardinaux, A., Simons, C. T., ... Sinha, P. (2016). Neural correlates of the food/non-food visual distinction. *Biological Psychology*, 115, 35–42.
- Ungar, P. S., & Sponheimer, M. (2011). The diets of early hominins. *Science*, 334(6053), 190–193.
- Vignando, M., Aiello, M., Foroni, F., Marcon, G., Tettamanti, M., & Rumiati, R. I. (2018). How experience modulates semantic memory for food: Evidence from elderly adults and centenarians. *Scientific Reports*, 8(1), 1–9.
- Wertz, A. E. (2019). How plants shape the mind. *Trends in Cognitive Sciences*, 23(7), 528–531.
- Wertz, A. E., & Wynn, K. (2014a). Selective social learning of plant edibility in 6- and 18-month-old infants. *Psychological Science*, 25(4), 874–882.
- Williams, B. J., & Kaufmann, L. M. (2012). Reliability of the go/no go association task. *Journal of Experimental Social Psychology*, 48(4), 879–891.
- Wilson, T. D., Lindsey, S., & Schooler, T. Y. (2000). A model of dual attitudes. *Psychological Review*, 107(1), 101–126.
- Wirt, A., & Collins, C. E. (2009). Diet quality—what is it and does it matter? *Public Health Nutrition*, 12(12), 2473–2492.
- Włodarczyk, A., Elsner, C., Schmitterer, A., & Wertz, A. E. (2018). Every rose has its thorn: Infants' responses to pointed shapes in naturalistic contexts. *Evolution and Human Behavior*, 39(6), 583–593.
- Wrangham, R. W. (2009). *Catching fire: How cooking made us human*. New York, New York, NY: Basic Books.
- Wrangham, R. W., & Conklin-Brittain, N. (2003). Cooking as a biological trait. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, 136(1), 35–46.
- Wrangham, R. W., Jones, J. H., Laden, G., Pilbeam, D., & Conklin-Brittain, N. (1999). The raw and the stolen. *Current Anthropology*, 40(5), 567–594.
- Zimmerman, M. E. (2011). *Speed-accuracy tradeoff*. Encyclopedia of clinical neuropsychology.
- Zink, K. D., & Lieberman, D. E. (2016). Impact of meat and Lower Palaeolithic food processing techniques on chewing in humans. *Nature*, 531(7595), 500–503.