

CASE REPORT

Open Access



Hyperkinetic stereotyped movements in a boy with biallelic CNTNAP2 variants

Marcello Scala^{1,2†} , Midas Anijs^{3†}, Roberta Battini^{4,5}, Francesca Madia⁶, Valeria Capra⁶, Paolo Scudieri^{2,6}, Alberto Verrotti⁷, Federico Zara⁶, Carlo Minetti^{1,2}, Sonja C. Vernes^{3,8,9*} and Pasquale Striano^{1,2*}

Abstract

Background: Heterozygous variants in *CNTNAP2* have been implicated in a wide range of neurological phenotypes, including intellectual disability (ID), epilepsy, autistic spectrum disorder (ASD), and impaired language. However, heterozygous variants can also be found in unaffected individuals. Biallelic *CNTNAP2* variants are rarer and cause a well-defined genetic syndrome known as CASPR2 deficiency disorder, a condition characterised by ID, early-onset refractory epilepsy, language impairment, and autistic features.

Case-report: A 7-year-old boy presented with hyperkinetic stereotyped movements that started during early infancy and persisted over childhood. Abnormal movements consisted of rhythmic and repetitive shaking of the four limbs, with evident stereotypic features. Additional clinical features included ID, attention deficit-hyperactivity disorder (ADHD), ASD, and speech impairment, consistent with CASPR2 deficiency disorder. Whole-genome array comparative genomic hybridization detected a maternally inherited 0.402 Mb duplication, which involved intron 1, exon 2, and intron 2 of *CNTNAP2* (c.97 +?_209-?dup). The affected region in intron 1 contains a binding site for the transcription factor FOXP2, potentially leading to abnormal *CNTNAP2* expression regulation. Sanger sequencing of the coding region of *CNTNAP2* also identified a paternally-inherited missense variant c.2752C > T, p.(Leu918Phe).

Conclusion: This case expands the molecular and phenotypic spectrum of CASPR2 deficiency disorder, suggesting that Hyperkinetic stereotyped movements may be a rare, yet significant, clinical feature of this complex neurological disorder. Furthermore, the identification of an in-frame, largely non-coding duplication in *CNTNAP2* points to a sophisticated underlying molecular mechanism, likely involving impaired FOXP2 binding.

Keywords: CASPR2, Intragenic duplication, Hyperkinetic movement disorder, Hyperkinesia, Speech impairment, Intellectual disability, Autism

Background

Biallelic *CNTNAP2* variants cause CASPR2 deficiency disorder (CDD), a syndromic neurodevelopmental disorder involving refractory epilepsy, ID, language impairment, and autistic features [1–4]. Heterozygous variants

in *CNTNAP2* have also been identified in patients with a wide range of complex neurological phenotypes, but such variants can also be found in unaffected individuals [5, 6].

CASPR2, the protein product of *CNTNAP2*, is a trans-membrane cell adhesion molecule from the neuroligin family that is widely expressed throughout the brain [5, 7]. CASPR2 localises to juxtaparanodes of myelinated axons, where it is involved in neuron-glia interactions, and mediates the clustering of potassium channels via interaction with contactin 2 (also known as TAG-1) [8, 9]. CASPR2 is also localised to the synapse where it is

* Correspondence: Sonja.vernes@mpi.nl; strianop@gmail.com; pasqualestriano@gaslini.org

[†]Marcello Scala and Midas Anijs contributed equally to this work.

³Neurogenetics of Vocal Communication Group, Max Planck Institute for Psycholinguistics, Nijmegen, The Netherlands

¹Pediatric Neurology and Muscular Diseases Unit, IRCCS Istituto Giannina Gaslini, Genoa, Italy

Full list of author information is available at the end of the article



© The Author(s). 2021 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

involved in several additional processes, such as neuronal migration, neurite development and synapse maturation, stability, and function [10–14].

We describe a patient carrying compound heterozygous variants in *CNTNAP2* including a missense variant and an intragenic duplication that were inherited from the father and mother, respectively. In addition to the common features found in *CASPR2* deficiency disorder (ID, ADHD, ASD, and speech impairment), the boy presented with peculiar hyperkinetic stereotyped movements, expanding the molecular and phenotypic spectrum of CDD.

Case presentation

This case was a 7-year-old boy without a family history of neurodevelopmental disability, born at term to non-consanguineous healthy parents (Fig. 1A) following a twin pregnancy complicated by intrauterine growth restriction and preeclampsia. His dizygotic twin brother was healthy at birth but was diagnosed with absence epilepsy during infancy. The neonatal course was characterised by feeding difficulties leading to failure to thrive. At 3 months of age, recurrent episodes of crying associated with semi-continuous, repetitive jerky movements of upper and lower limbs were observed, which were diagnosed as hyperkinetic stereotyped movements (Supplementary Video 1).

These paroxysmal hyperkinetic-dyskinetic episodes recurred periodically without any trigger, lasted 10–12 h, and mimicked infantile colics. Their frequency was temporarily reduced when anti-reflux formulas and ranitidine were administered for concomitant recurrent vomiting. Electroencephalogram and brain magnetic resonance imaging were unremarkable.

During his first years of life, the patient displayed ADHD and autistic traits. The boy also featured a phonological processing deficiency and clumsiness in gross and fine movements was also observed. His neuropsychological evaluation (WPPSI-III at age 6 years) revealed a global IQ of 71 (verbal, 86; performance, 68; processing, 55). The movement disorder persisted during the following years through non-triggered fast, high-amplitude, rhythmic, continuous and repetitive shaking involving the four limbs with stereotypic features (Supplementary Videos 2–3). His stereotypic movement disorder had been misdiagnosed as tics before, which indeed tend to appear from age 6–7 years [15]. However, these repetitive movements already started before the age of 3 and consisted of intense patterns of movement that ran longer than tics and were more bilateral than tics. Several medications, including carbamazepine, valproate, gabapentin, levodopa, flunarizine, benzodiazepines, did not improve his symptoms. The boy is currently receiving clonazepam, 0.2 mg/Kg in two daily doses.

Genetic findings


Written informed consent was obtained from the patient's parents. Whole-genome array-CGH on peripheral blood genomic DNA of the family quartet revealed a maternally inherited *CNTNAP2* variant c.97 +?_209-?dup in the proband. Real-time quantitative PCR confirmed that this 0.402 Mbp duplication involved part of intron 1 and exon 2, with a breakpoint within intron 2 (Fig. 1C). This duplication is not predicted to introduce a frameshift in the *CASPR2* protein. The duplication was not inherited by the sibling.

PCR amplification and Sanger sequencing was performed to screen *CNTNAP2* exons in the family, revealing a paternally inherited missense variant c.2752C > T, p.(Leu918Phe) in the proband (Fig. 1B). This variant is absent from the gnomAD dataset (<https://gnomad.broadinstitute.org/variant/>) and affects a conserved residue within the Laminin G-like (LG) 3 domain (GERP = 5.49). It is reported in ClinVar (allele ID 924741, <https://www.ncbi.nlm.nih.gov/clinvar/variation/949154/>) and predicted damaging by in silico tools (e.g., DANN, 0.9977; Mutation Taster, 1; CADD, 24.1). Both variants were absent in the proband's sibling. Further details are available in [Supplementary Methods](#).

Discussion

The range of clinical symptoms found in patients harboring biallelic *CNTNAP2* variants are collectively described as *CASPR2* deficiency disorder. Dyskinetic features are not included amongst the classic neurological manifestations of this condition [1–4]. In this patient, we observed hyperkinetic stereotyped movements consisting of continuous, repetitive and rhythmic shaking of the four limbs, which became evident during the first months of life and persisted over the years. These abnormal movements did not resemble generalised dystonia, paroxysmal non-kinesigenic dyskinesia, or paroxysmal kinesigenic dyskinesia, but rather represent a novel neurological manifestation of *CASPR2* deficiency disorder. This dyskinetic phenotype may be related to the role of *CASPR2* in facilitating nerve conduction and synaptic connectivity, particularly given its widespread expression in the central and peripheral nervous system [5, 9, 14]. In patients harboring biallelic variants in *CNTNAP2*, brain MRI may be normal, such as in our case, or show cortical dysplasia (cortical dysplasia-focal epilepsy syndrome, CDFES).^{1–3} Additional findings include cerebellar abnormalities (vermian hypoplasia or atrophy) and nonspecific white matter abnormalities.^{1,2}

The c.97 +?_209-?dup and p.(Leu918Phe) variants were inherited from unaffected parents, supporting the incomplete penetrance of heterozygous *CNTNAP2* variants [5, 6]. The maternally inherited variant (c.97 +?_209-?dup) involved a complete duplication of exon 2



(99 bp) (Fig. 1D), which may negatively impact protein folding and/or protein-protein interactions. Indeed, exon 2 encodes part of the discoidin homology domain, which is found in the extracellular portion CASPR2 and is known to mediate protein-protein interactions, pointing to a potential perturbation of this mechanism in the patient (Fig. 1E). The paternally inherited missense variant p.(Leu918Phe) is also in the extracellular portion of CASPR2, but is found in the third LG domain (Fig. 1E). Missense variants in the extracellular portion of CASPR2 have been shown to impair interactions with contactin 2 and affect axon growth in cortical neurons [12]. This

effect is likely related to the trans-synaptic bridge formed in neurons by the interaction of CASPR2 and contactin 2 that contributes to synaptic organization and synaptic transmission [16]. It would therefore be of value to determine if the variants identified in this patient have functional consequences for protein-protein interactions, or for synaptic or juxtapanodal organisation and signalling. Such functional studies may shed light on why heterozygous *CNTNAP2* variants show incomplete penetrance and why biallelic mutations together produce a range of phenotypes, including the novel motor phenotype described herein.

The maternal duplication also has the potential to alter the regulation of *CNTNAP2*. The transcription factor FOXP2 targets a binding site in intron 1 of *CNTNAP2* and regulates its expression. Mutations in *FOXP2* are a monogenic cause of childhood apraxia of speech (SPCH1, OMIM #602081) [17] and *CNTNAP2* is functionally implicated in the aetiology of this condition as part of the downstream network of FOXP2 target genes [18]. This is supported by a genetic association of *CNTNAP2* variants with phonological memory performance in children with specific language impairment [18]. Duplication of the FOXP2 target site in intron 1 of the maternally inherited allele could alter FOXP2-mediated regulation of *CNTNAP2* and lead to aberrant CASPR2 levels, a hypothesis that needs functional testing in cell or animal models. In this way, it is possible that the intronic duplication could contribute to neurodevelopmental or speech phenotypes in patients that are related to both FOXP2 and *CNTNAP2*, such as phonological processing.

This study has two main limitations. First, a proper functional characterization of the two detected *CNTNAP2* variants to evaluate their impact on protein structure and function could not be performed. Second, it was not possible to further investigate the presence of additional variants with a potential modifier effect on the clinical phenotype through Next Generation Sequencing (NGS) techniques, such as whole exome sequencing (WES) or whole genome sequencing (WGS).

Conclusion

This case of a boy with hyperkinetic stereotyped movements and biallelic *CNTNAP2* mutations expands our knowledge about *CNTNAP2*-related disorders. It presents a new, rare neurological manifestation for CDD and posits a remarkable molecular mechanism in which coding and non-coding *CNTNAP2* mutations could contribute to the observed phenotypes. Accordingly, we suggest screening *CNTNAP2* regulatory regions in patients with a CDD-suggestive phenotype even if a single heterozygous *CNTNAP2* variant has been identified or if atypical neurological phenotypes are also present. This will lead to better diagnoses that can improve the management of patients with these disorders. This, together with functional studies of the consequences of the identified mutations, will advance our scientific understanding of disease genes like *CNTNAP2*.

Abbreviations

ADHD: Attention deficit-hyperactivity disorder; ASD: Autism spectrum disorder; CDD: CASPR2 deficiency disorder; ID: Intellectual disability; NGS: Next Generation Sequencing; WES: Whole exome sequencing; WGS: Whole genome sequencing

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s13052-021-01162-w>.

Additional file 1.

Additional file 2.

Additional file 3.

Additional file 4.

Acknowledgements

We thank the patient and his parents for their kind cooperation and the consent to the publication of this study.

Authors' contributions

MS, MA, SCV, and PS conceived and designed the study. MS, RB, FM, VC, and PS clinically reviewed the case and contributed clinical data. MA and SCV designed and carried out molecular studies. MS, MA, SCV, PS, AV, FZ, and CM analysed and interpreted the data. MS and MA drafted the manuscript. SCV and PS supervised the drafting of and edited the manuscript. All authors read, critically revised, and approved the final manuscript.

Funding

SCV was supported by a Max Planck Research Group awarded by the Max Planck Gesellschaft, a Human Frontiers Science Program Grant (RGP0058/2016), and a UKRI Future Leaders Fellowship (MR/T021985/1). MA was supported by an International Max Planck Research School (IMPRS) PhD Fellowship from the Max Planck Institute for Psycholinguistics.

Availability of data and materials

All data generated or analysed during this study are included in this published article.

Declarations

Ethics approval and consent to participate

The patient's parents provided written informed consent to exome sequencing and inclusion into the research study approved by the Giannina Gaslini Children's Institute Ethics Committee.

Consent for publication

A signed consent for this publication has been obtained from the patient and his parents.

Competing interests

All authors declare no disclosures relevant to the manuscript.

Author details

¹Pediatric Neurology and Muscular Diseases Unit, IRCCS Istituto Giannina Gaslini, Genoa, Italy. ²Department of Neurosciences, Rehabilitation, Ophthalmology, Genetics, Maternal and Child Health University of Genoa, Genoa, Italy. ³Neurogenetics of Vocal Communication Group, Max Planck Institute for Psycholinguistics, Nijmegen, The Netherlands. ⁴Department of Clinical and Experimental Medicine, University of Pisa, Pisa, Italy. ⁵IRCCS Fondazione Stella Maris, Pisa, Italy. ⁶Medical Genetic Unit, IRCCS Istituto G. Gaslini, Genoa, Italy. ⁷Department of Pediatrics, University of Perugia, Perugia, Italy. ⁸School of Biology, University of St Andrews, Fife, UK. ⁹Donders Institute for Brain, Cognition and Behaviour, Nijmegen, The Netherlands.

Received: 7 July 2021 Accepted: 25 September 2021

Published online: 12 October 2021

References

- Rodenas-Cuadrado P, Pietrafusa N, Francavilla T, La Neve A, Striano P, Vernes SC. Characterisation of CASPR2 deficiency disorder—a syndrome involving autism, epilepsy and language impairment. *BMC Med Genet*. 2016;17(1):8. <https://doi.org/10.1186/s12881-016-0272-8>.
- Smogavec M, Cleall A, Hoyer J, Lederer D, Nassogne MC, Palmer EE, et al. Eight further individuals with intellectual disability and epilepsy carrying biallelic *CNTNAP2* aberrations allow delineation of the mutational and

- phenotypic spectrum. *J Med Genet.* 2016;53(12):820–7. <https://doi.org/10.1136/jmedgenet-2016-103880>.
3. Zweier C, de Jong EK, Zweier M, Orrico A, Ousager LB, Collins AL, et al. CNTNAP2 and NRXN1 are mutated in autosomal-recessive Pitt-Hopkins-like mental retardation and determine the level of a common synaptic protein in *Drosophila*. *Am J Hum Genet.* 2009;85(5):655–66. <https://doi.org/10.1016/j.ajhg.2009.10.004>.
 4. Watson CM, Crinnion LA, Tzika A, Mills A, Coates A, Pendlebury M, et al. Diagnostic whole genome sequencing and split-read mapping for nucleotide resolution breakpoint identification in CNTNAP2 deficiency syndrome. *Am J Med Genet A.* 2014;164a(10):2649–55.
 5. Rodenas-Cuadrado P, Ho J, Vernes SC. Shining a light on CNTNAP2: complex functions to complex disorders. *Eur J Hum Genet.* 2014;22(2):171–8. <https://doi.org/10.1038/ejhg.2013.100>.
 6. Toma C, Pierce KD, Shaw AD, Heath A, Mitchell PB, Schofield PR, et al. Comprehensive cross-disorder analyses of CNTNAP2 suggest it is unlikely to be a primary risk gene for psychiatric disorders. *PLoS Genet.* 2018;14(12):e1007535. <https://doi.org/10.1371/journal.pgen.1007535>.
 7. Gordon A, Salomon D, Barak N, Pen Y, Tsoory M, Kimchi T, et al. Expression of Cntnap2 (Caspr2) in multiple levels of sensory systems. *Mol Cell Neurosci.* 2016;70:42–53. <https://doi.org/10.1016/j.mcn.2015.11.012>.
 8. Poliak S, Gollan L, Salomon D, Berglund EO, Ohara R, Ranscht B, et al. Localization of Caspr2 in myelinated nerves depends on axon-glia interactions and the generation of barriers along the axon. *J Neurosci.* 2001; 21(19):7568–75. <https://doi.org/10.1523/JNEUROSCI.21-19-07568.2001>.
 9. Poliak S, Salomon D, Elhanany H, Sabanay H, Kiernan B, Pevny L, et al. Juxtaparanodal clustering of shaker-like K⁺ channels in myelinated axons depends on Caspr2 and TAG-1. *J Cell Biol.* 2003;162(6):1149–60. <https://doi.org/10.1083/jcb.200305018>.
 10. Peñagarikano O, Abrahams BS, Herman EI, Winden KD, Gdalyahu A, Dong H, et al. Absence of CNTNAP2 leads to epilepsy, neuronal migration abnormalities, and core autism-related deficits. *Cell.* 2011;147(1):235–46. <https://doi.org/10.1016/j.cell.2011.08.040>.
 11. Anderson GR, Galfin T, Xu W, Aoto J, Malenka RC, Südhof TC. Candidate autism gene screen identifies critical role for cell-adhesion molecule CASPR2 in dendritic arborization and spine development. *Proc Natl Acad Sci U S A.* 2012;109(44):18120–5. <https://doi.org/10.1073/pnas.1216398109>.
 12. Goullancourt A, Canali G, Girault J-A, Oguievetskaia K, Garcia M, Goutebroze L, et al. Genetic variants in autism-related CNTNAP2 impair axonal growth of cortical neurons. *Hum Mol Genet.* 2018;27(11):1941–54. <https://doi.org/10.1093/hmg/ddy102>.
 13. Varea O, Martin-de-Saavedra MD, Kopeikina KJ, Schürmann B, Fleming HJ, Fawcett-Patel JM, et al. Synaptic abnormalities and cytoplasmic glutamate receptor aggregates in contactin associated protein-like 2/Caspr2 knockout neurons. *Proc Natl Acad Sci U S A.* 2015;112(19):6176–81. <https://doi.org/10.1073/pnas.1423205112>.
 14. Lazaro MT, Taxidis J, Shuman T, Bachmutsky I, Ikrar T, Santos R, et al. Reduced Prefrontal Synaptic Connectivity and Disturbed Oscillatory Population Dynamics in the CNTNAP2 Model of Autism. *Cell Rep.* 2019; 27(9):2567–78.e6.
 15. Zinner SH, Mink JW. Movement disorders I: tics and stereotypies. *Pediatr Rev.* 2010;31(6):223–33. <https://doi.org/10.1542/pir.31-6-223>.
 16. Lu Z, Reddy MV, Liu J, Kalichava A, Liu J, Zhang L, et al. Molecular architecture of Contactin-associated protein-like 2 (CNTNAP2) and its interaction with Contactin 2 (CNTN2). *J Biol Chem.* 2016;291(46):24133–47. <https://doi.org/10.1074/jbc.M116.748236>.
 17. Lai CS, Fisher SE, Hurst JA, Vargha-Khadem F, Monaco AP. A forkhead-domain gene is mutated in a severe speech and language disorder. *Nature.* 2001;413(6855):519–23. <https://doi.org/10.1038/35097076>.
 18. Vernes SC, Newbury DF, Abrahams BS, Winchester L, Nicod J, Groszer M, et al. A functional genetic link between distinct developmental language disorders. *N Engl J Med.* 2008;359(22):2337–45. <https://doi.org/10.1056/NEJMoa0802828>.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

