

All-sky search for gravitational wave emission from scalar boson clouds around spinning black holes in LIGO O3 data

The LIGO Scientific Collaboration, The Virgo Collaboration, and The KAGRA Collaboration*
(Dated: December 1, 2021)

This paper describes the first all-sky search for long-duration, quasi-monochromatic gravitational-wave signals emitted by ultralight scalar boson clouds around spinning black holes using data from the third observing run of Advanced LIGO. We analyze the frequency range from 20 Hz to 610 Hz, over a small frequency derivative range around zero, and use multiple frequency resolutions to be robust towards possible signal frequency wanderings. Outliers from this search are followed up using two different methods, one more suitable for nearly monochromatic signals, and the other more robust towards frequency fluctuations. We do not find any evidence for such signals and set upper limits on the signal strain amplitude, the most stringent being $\approx 10^{-25}$ at around 130 Hz. We interpret these upper limits as both an “exclusion region” in the boson mass/black hole mass plane and the maximum detectable distance for a given boson mass, based on an assumption of the age of the black hole/boson cloud system.

I. INTRODUCTION

Theories of beyond standard model physics allow for the production of ultralight bosons that could constitute a portion or all of dark matter [1]. If such ultralight bosons exist, they could appear around rotating black holes due to quantum fluctuations [2]. They would then scatter off and extract angular momentum from these black holes, and form macroscopic clouds, i.e. boson condensates, through a superradiance process [2]. The energy structure of the clouds resembles that of a hydrogen atom, earning boson clouds the name “gravitational atoms” in the sky. After the black hole spin decreases below a threshold, the superradiance process stops and the cloud depletes over time mainly through annihilation of bosons into gravitons, which generates quasi-monochromatic, long-duration gravitational waves when the self-interaction for bosons is weak.

The signal would also have a small, positive frequency variation, known as a “spin-up”, that would arise from the classical contraction of the cloud over time as it loses mass [2]. The values of the spin-up depend on whether we consider scalar [3], vector [4, 5], or tensor [6] boson clouds, but in this search, we consider scalar boson cloud signals with very small spin-ups, of maximum $\mathcal{O}(10^{-12})$ Hz/s. Recently, the effect on gravitational-wave emission from boson self-interactions has been studied for the scalar case [7], showing that the signal can be significantly affected, including the magnitude of the spin-up, as the self-interaction increases.

The gravitational-wave signal frequency depends primarily on the mass of the boson, and weakly on the mass and spin of the black hole. Recently, exclusion regions of the black hole / scalar boson mass parameter space were calculated using upper limits from an all-sky search for spinning neutron stars in LIGO/Virgo data from the second observing run (O2) [8]. Furthermore, a search

for boson clouds from Cygnus X-1 was performed with a hidden Markov model (HMM) method [9] on the same data set, and disfavored particular boson masses for this object [10]. It has been suggested [11] that current methods for all-sky searches, such as the one used in [8], could suffer a significant sensitivity penalty in the extreme case of a population of 10^8 black holes in the Milky Way, due to the superposition of many signals in a small frequency range (as a consequence of the signal’s frequency dependence at leading order on the boson mass). A detailed study [12] has quantified this effect, showing that the actual average loss of sensitivity is at most of about 15% for a signal “density” of 1-3 signals per frequency bin, while it rapidly reduces, and becomes negligible, for both lower and higher densities.

In addition to quasi-monochromatic gravitational waves emitted by boson clouds around isolated black holes, it is also possible to probe the existence of boson clouds in binary black hole coalescences. One avenue requires measurements of spin: in principle, boson clouds will extract mass and spin from black holes, resulting in low spin values for black holes compared to those in a universe without boson clouds; in practice, individual black hole spins are hard to measure [13, 14], and the current spin distribution of black holes depends on both the mass of boson clouds and the distribution of spins when the black holes were born [15, 16]. It is therefore interesting to hierarchically combine spin measurements from various black hole mergers to obtain constraints on boson cloud/spin interactions [17, 18]. Additionally, the presence of a boson cloud will induce a change in the waveforms used in compact binary searches, e.g. new multipole moments and tidal effects (parameterized as Love numbers) [19, 20], and such differences in binary black hole mergers may actually be detectable depending on the boson mass and field strength [21]. It has also been proposed to combine multi-band observations of black hole mergers and boson cloud signals when future detectors are online [22], and to search for a stochastic gravitational-wave background from scalar and vector boson cloud signals [23–25]. Complementary meth-

* Full author list given at the end of the article.

ods [26, 27] and searches [28, 29] for ultralight scalar and vector bosons have also been developed that use the gravitational-wave interferometers as particle detectors, which provide additional constraints on the boson mass. Ultralight bosons can therefore have different effects on different types of gravitational-wave signals.

This paper presents results from the first all-sky search tailored to gravitational waves from depleting scalar boson clouds around black holes using the Advanced LIGO [30] data in the third observing run (O3). Although the expected gravitational-wave signal is almost monochromatic, it could come from anywhere in the sky; thus, the Doppler modulations from the relative motion of the Earth and the source, of $\mathcal{O}(10^{-4}f)$, where f is the frequency of the signal, are a priori unknown. A fully coherent search, in which we take a single Fourier Transform of the whole data set and look for peaks in the power spectrum after demodulating the data, is therefore impossible to perform in each sky direction because of limited computational power. Indeed, the number of sky positions to search over scales with the square of both the Fourier Transform length and the frequency, and would amount to $\mathcal{O}(10^{20})$ at high frequencies [31]. This means that we must employ semi-coherent methods, in which we break the data into smaller chunks in time that are analyzed coherently, and then combined incoherently, to keep the computational cost under control while retaining the desired sensitivity [32, 33]. Moreover, we adopt a multi-resolution approach, equivalent to considering several different Fourier Transform lengths [34], in order to be sensitive towards possible unpredicted frequency fluctuations of the gravitational-wave signal.

This paper is organized as follows: in section II, we describe our model for a gravitational-wave signal resulting from a depleting scalar boson cloud around a rotating black hole; in section III, we explain our method to search for such signals; in section IV, we provide information on the data sets we use in our analysis; in section V, we present the results of the search and our upper limits, including their astrophysical interpretations in terms of constraints in the boson/black hole mass plane and on the maximum detectable distance; finally, in section VI, we discuss prospects for future work. Appendices contain details on the follow-up of selected outliers.

II. THE SIGNAL

An isolated black hole is born with a defining mass, spin and charge [35]. Ultralight boson particles around a black hole will scatter off it and extract energy and momentum from it if the so-called *superradiant* condition is satisfied, i.e. $\omega < m\Omega$ [2, 36] where ω is the angular frequency of the boson (related to the boson mass linearly), m is the azimuthal quantum number in the hydrogen-like gravitational atom, and Ω is the angular frequency of the rotating black hole's outer horizon. Because the bosons have a finite mass, they become gravitationally

bound to the black hole, which allows for successive scatterings and a build-up of a macroscopic cloud, extracting up to about 10% of the black hole mass [2]. This process is maximized when the particle's reduced Compton wavelength is comparable to the black hole radius:

$$\frac{\hbar c}{m_b} \simeq \frac{GM_{\text{BH}}}{c^2}, \quad (1)$$

where m_b is the boson mass-energy, M_{BH} is the black hole mass, \hbar is the reduced Planck's constant, and c is the speed of light. The typical time scale of the superradiance phase is [3]

$$\tau_{\text{inst}} \approx 27 \left(\frac{M_{\text{BH}}}{10M_{\odot}} \right) \left(\frac{\alpha}{0.1} \right)^{-9} \left(\frac{1}{\chi_i} \right) \text{ days}, \quad (2)$$

where χ_i is the black hole initial dimensionless spin, and

$$\alpha = \frac{GM_{\text{BH}}}{c^3} \frac{m_b}{\hbar}, \quad (3)$$

is the fine-structure constant in the gravitational atom.

Once the superradiant condition is no longer satisfied, the growth stops, and the cloud begins to give off energy primarily via annihilation of particles in the form of gravitational waves [9], over a typically much longer time scale which, for $m = 1$ and in the limit $\alpha \ll 0.1$, is given by

$$\tau_{\text{gw}} \approx 6.5 \times 10^4 \left(\frac{M_{\text{BH}}}{10M_{\odot}} \right) \left(\frac{\alpha}{0.1} \right)^{-15} \left(\frac{1}{\chi_i} \right) \text{ years}. \quad (4)$$

For scalar bosons, these clouds have a much shorter growth time scale compared to the time scale for them to deplete [3]; thus, if they exist now, they are more likely to be depleting rather than growing. The gravitational-wave emission is at a frequency f_{gw} [8]:

$$f_{\text{gw}} \simeq 483 \text{ Hz} \left(\frac{m_b}{10^{-12} \text{ eV}} \right) \times \left[1 - 7 \times 10^{-4} \left(\frac{M_{\text{BH}}}{10M_{\odot}} \frac{m_b}{10^{-12} \text{ eV}} \right)^2 \right]. \quad (5)$$

In fact, this frequency slowly increases in time by an amount that depends on the boson self-interaction constant F_b [7]. As in [7], we consider a scalar boson that, in addition to a non-zero mass, has a quartic interaction with an extremely small coupling λ , and characterize this coupling in terms of the parameter $F_b := m_b/\sqrt{\lambda}$. In the context of an axion-like particle, F_b would roughly correspond to the symmetry breaking energy scale.

Specifically, if $F_b \gtrsim 2 \times 10^{18}$ GeV, the spin-up is dominated by the depletion of the cloud through particle annihilation and is given by

$$\dot{f}_{\text{gw}} \approx 7 \times 10^{-15} \left(\frac{m_b}{10^{-12} \text{ eV}} \right)^2 \left(\frac{\alpha}{0.1} \right)^{17} \text{ Hz/s}. \quad (6)$$

With this condition on F_b , our search probes energies at the Planck scale, and is sensitive to the QCD axion with a mass of $\sim 10^{-13}\text{--}10^{-12}$ eV [7]. However, if

the bosons have a non-negligible self-interaction (i.e. for smaller F_b), two other spin-up terms, one due to energy level transitions and another due to the variation of the self-interaction energy, appear. The former is given by

$$\dot{f}_{\text{tr}} \approx 10^{-10} \left(\frac{m_b}{10^{-12} \text{ eV}} \right)^2 \left(\frac{\alpha}{0.1} \right)^{17} \left(\frac{10^{17} \text{ GeV}}{F_b} \right)^4 \text{ Hz/s}, \quad (7)$$

which is dominant in the intermediate self-interaction regime, for $8.5 \times 10^{16}(\alpha/0.1) \text{ GeV} \lesssim F_b \lesssim 2 \times 10^{18} \text{ GeV}$. In this regime, the signal duration significantly shortens, thus reducing the chance of detection. In the strong self-interaction regime, when $F_b \lesssim 8.5 \times 10^{16}(\alpha/0.1) \text{ GeV}$, the spin-up can be parametrized as

$$\dot{f}_{\text{se}} \approx 10^{-10} \left(\frac{m_b}{10^{-12} \text{ eV}} \right)^2 \left(\frac{\alpha}{0.1} \right)^{17} \left(\frac{10^{17} \text{ GeV}}{F_b} \right)^6 \text{ Hz/s}, \quad (8)$$

and the signal strength rapidly decreases with increasing interaction, making the detection of annihilation signals basically impossible for current detectors. In Sec. V C, we will briefly discuss the implication of our search results in relation to the value of F_b .

These signals have lower amplitudes than those from compact binary coalescences [11], with initial values (for $\alpha \ll 0.1$)

$$h_0 \approx 6 \times 10^{-24} \left(\frac{M_{\text{BH}}}{10 M_{\odot}} \right) \left(\frac{\alpha}{0.1} \right)^7 \left(\frac{1 \text{ kpc}}{D} \right) (\chi_i - \chi_c), \quad (9)$$

where

$$\chi_c \approx \frac{4\alpha}{4\alpha^2 + 1}, \quad (10)$$

is the black hole spin when the superradiance is saturated, and D is the source distance. The emitted signal amplitude decreases in time as the clouds are depleted, according to

$$h(t) = \frac{h_0}{1 + \frac{t}{\tau_{\text{gw}}}}. \quad (11)$$

The equations above are analytic approximations to the true behavior of a gravitational wave originating from boson clouds. When we consider $\alpha \sim 0.1$, we must account for the difference between the numerical and analytic solutions, which is ~ 3 in energy at the highest α considered here, $\alpha \sim 0.15$. We obtain this factor of ~ 3 comparing the black and red curves in Fig. 2 of [3] at $\alpha \sim 0.15$.

III. SEARCH METHOD

The semi-coherent method we employ is based on the Band Sampled Data (BSD) framework [37], which is being used in an increasing number of continuous-wave searches, due to its flexibility and computational efficiency [38–40]. BSD files are data structures that store

a reduced analytic strain-calibrated time series in 10-Hz/one-month chunks. To construct BSD files, we take a Fourier transform of a chunk of time-series strain data, extract a 10-Hz band, keep only the positive frequency components, and inverse Fourier Transform to obtain the reduced analytic signal.

The main steps of the analysis pipeline are schematically shown in Fig. 1 and are summarized below.




FIG. 1. Scheme of the search pipeline.

A. Peakmap construction

Our analysis starts with a set of BSD files covering the whole third observing run and frequencies between 20 Hz and 610 Hz. The maximum frequency is chosen to ensure the computational cost of the analysis is reasonable, and is consistent with the fact that the most relevant part of the accessible parameter space of the black hole-boson system corresponds to the intermediate frequencies. Each of these time series is divided into chunks of duration

$$T_{\text{FFT}} = \frac{1}{\Omega_{\text{rot}}} \sqrt{\frac{c}{2f_{10}R_{\text{rot}}}}, \quad (12)$$

where f_{10} is the maximum frequency of each corresponding 10-Hz band, $\Omega_{\text{rot}} = 2\pi/86164.09 \text{ rad/s}$ is the Earth sidereal angular frequency, and R_{rot} is the Earth rotational radius, which we conservatively take as that corresponding to the detector at the lowest latitude ($R_{\text{rot}} = 5.5 \times 10^6 \text{ km}$ for the Livingston detector). This T_{FFT} length guarantees that, if we take the Fourier Transform of the chunk, the power spread of any possible continuous-wave signal, caused by the motion of the Earth with respect to the source, is fully contained in one frequency bin with a width of $\delta f = 1/T_{\text{FFT}}$ in each chunk. In Fig. 2, the chunk duration is shown as a function of the frequency. For each of these chunks, the square modulus of the Fourier Transform is computed (by means of




FIG. 2. Chunk duration T_{FFT} as a function of frequency. The T_{FFT} is fixed within each 10-Hz band.

the FFT algorithm) and divided by an estimation of the average spectrum over the same time window. The resulting quantity has a mean value of one independently of the frequency - for this reason it is called the *equalized* spectrum - and takes values significantly different from one when narrow frequency features are present in the data. We select prominent peaks in the equalized spectra, defined as time-frequency pairs that correspond to local maxima and have a magnitude above a threshold of $\theta = 2.5$ [31]. The collection of the these time-frequency peaks forms the so-called *peakmap*, see e.g. [31] for more details.

B. Doppler effect correction

We build a suitable grid in the sky such that, after the Doppler effect correction for a given sky direction, any residual Doppler effect always produces an error in frequency within half a frequency bin [31]. For each sky direction in the grid, the peaks in the peakmap are shifted in order to compensate for the time-dependent Doppler modulation, according to the relation

$$f_{0,t_k} = \frac{f_{\text{obs},t_k}}{1 + \frac{\vec{v}_{t_k} \cdot \hat{n}}{c}}, \quad (13)$$

where f_{obs,t_k} is an observed frequency peak at the time t_k , \vec{v}_{t_k} is the detector's velocity, and \hat{n} is the unit vector identifying the sky direction.

As we have described in Sec. II, the signal has an intrinsic spin-up, associated with the cloud depletion. Although we do not apply an explicit correction for the spin-up, the analysis retains most of the signal power - with a maximum sensitivity loss less than a few percent¹

- as long as the corresponding frequency variation during the full observation time T_{obs} is confined within $\pm 1/2$ bin of the frequency evolution rate $\delta \dot{f}$, given by

$$\delta \dot{f} = \frac{1}{2T_{\text{obs}}T_{\text{FFT}}}. \quad (14)$$

The $\delta \dot{f}$ value is plotted as a function of the frequency in Fig. 3.




FIG. 3. Bin size of the frequency time derivative, $\delta \dot{f}$, a function of the frequency, given the full observing time of O3 and T_{FFT} chosen for each 10-Hz band. The search by default covers a frequency time derivative range of $\dot{f} \in [-\delta \dot{f}/2, \delta \dot{f}/2]$. Such a range contributes to set constraints on the portion of the boson mass/black hole mass explored in the search.

C. Peakmap projection

Once we have applied the Doppler correction, we project the peakmap onto the frequency axis and select the most significant frequencies for further analysis. Indeed, if a monochromatic signal comes from a given direction, we expect its associated peaks to be aligned in frequency and appear as a prominent peak in the projected peakmap, when the signal is strong enough. The statistics of the peakmap are described in detail in [31], so here we only provide a brief review. In the absence of signals, the probability p_0 of selecting a peak depends on the noise distribution. In the case of Gaussian noise, it is given by

$$p_0 = e^{-\theta} - e^{-2\theta} + \frac{1}{3}e^{-3\theta}, \quad (15)$$

where $\theta = 2.5$ is the threshold we apply to construct the peakmap. On the other hand, if a signal with spectral amplitude (in units of equalized noise)

$$\lambda = \frac{4|\tilde{h}(f)|^2}{T_{\text{FFT}}S_n(f)}, \quad (16)$$

¹ In principle, the maximum sensitivity loss would be of about 36%, when the corrected signal frequency falls in the middle among two consecutive frequency bins. In practice, however, the application of the moving average on the peakmaps, see section

III D, allows us to recover part of the lost signal power, reducing the maximum loss to about 19% (when applying a moving average of two bins, i.e. a window with $W=2$).

where $\tilde{h}(f)$ is the signal Fourier Transform, and $S_n(f)$ is the detector noise power spectral density, is present, the corresponding probability p_λ of selecting a signal peak, for weak signals with respect to the noise level, is given by

$$p_\lambda = p_0 + \frac{\lambda}{2} \theta (e^{-\theta} - 2e^{-2\theta} + e^{-3\theta}). \quad (17)$$

The statistic of the peakmap projection is, for well-behaved noise, a binomial with expectation value $\mu = Np_0$, where N is the number of FFTs, and standard deviation $\sigma = \sqrt{Np_0(1-p_0)}$. In the presence of a signal, the expected number of peaks at the signal frequency (after the Doppler correction) is Np_λ . It is then useful to introduce the Critical Ratio CR:

$$\text{CR} = \frac{n - \mu}{\sigma}, \quad (18)$$

where n is the actual number of peaks in a given frequency bin, which, in the limit of large N (in practice, greater than a ~ 30), closely follows a Gaussian distribution with zero mean and unity standard deviation.

D. Peakmap moving averages

The signal emitted by a boson cloud is not exactly monochromatic. In order to deal with the uncertainty in the signal morphology, we apply a sequence of moving averages (MA) over a window width W varying from two to ten frequency bins, starting from the Doppler corrected peakmap projection. This procedure can be shown to be equivalent – and computationally much more efficient – to build peakmap projections with an effective FFT duration $\approx T_{\text{FFT}}/W$. While these moving averages reduce the sensitivity to signals with a characteristic coherence time larger than T_{FFT} , e.g. purely monochromatic signals, they provide better sensitivity to signals with a typical coherence time comparable to the effective FFT duration [34].

E. Candidate selection and coincidences

We use the CR as a detection statistic to identify potentially interesting outliers. Outliers are uniformly selected in the parameter space by taking the two points with the highest CR in each peakmap projection, for each sky position in every 0.05-Hz frequency band, independent of the actual CR values, as long as it is above a minimum value of 3.8.

In the next step, we identify coincident pairs among outliers found in the Hanford and Livingston data, corresponding to the same W , by using a dimensionless coincidence window of 3. In other words, two outliers, one from Hanford and the other from Livingston, are consid-

ered coincident if the dimensionless distance

$$d = \sqrt{\left(\frac{\Delta f}{\delta f}\right)^2 + \left(\frac{\Delta \lambda}{\delta \lambda}\right)^2 + \left(\frac{\Delta \beta}{\delta \beta}\right)^2} \leq 3 \quad (19)$$

where Δf , $\Delta \lambda$, $\Delta \beta$ are the differences between parameter values of the outliers found in Hanford (H1) and Livingston (L1), and δf , $\delta \lambda$, $\delta \beta$ are the corresponding bin sizes². The coincidence window of 3 is chosen according to the studies carried out using simulated signals and is consistent with the choice in standard all-sky searches for neutron stars [41, 42].

F. Post-processing

Coincident outliers are subject to a sequence of post-processing steps in order to discard those incompatible with an astrophysical signal.

The first veto is to check if some of the outliers are compatible with known narrow-band instrumental disturbances, also known as *noise lines*. An outlier is considered as caused by a noise line, and thus discarded, if its Doppler band, defined by $\Delta f_{\text{dopp}} = \pm 10^{-4}f$, where f is the outlier frequency, intersects the noise line.

The next is a consistency veto, in which a pair of coincident outliers are discarded if their CR is not compatible with the detectors' noise level at the corresponding frequency. Specifically, we veto those coincident outliers for which the CR in one detector is more than a factor of 5 different than that in the other detector, after being normalized by the detector's power spectral density [8, 42]. This choice, also used in standard all-sky searches for continuous waves from neutron stars, is motivated by the desire to safely eliminate an outlier only if the discrepancy is really significant and the CR is large in at least one of the two detectors (i.e., a real astrophysical signal is not expected to behave that way).

The previous steps are all applied to coincident outliers separately for each value of the MA window width W . In case of outliers coincident across different W values, only the one with the highest CR is kept³.

G. Follow-up

Outliers with $W = 1$, the case in which we do not apply moving averages, could be due to monochromatic signals or signals with very small fluctuations in frequency, while outliers with $W > 1$ are more compatible with signals characterized by a larger degree of frequency fluctuation.

² The sky position bin sizes in general are different for two outliers identified at different sky positions, so the average among the corresponding bin sizes is actually used.

³ This restriction reduces the computational cost with only a small loss in sensitivity.

These two sets of candidates are then followed up with different methods, one based on the Frequency-Hough algorithm, and used e.g. in [41], and the other using the Viterbi method, see e.g. [10]. Both methods are briefly explained in Appendix B.

IV. DATA

We use data collected by the Advanced LIGO gravitational-wave detectors over the O3 run, which lasted from April 1st, 2019 to March 27th, 2020, with a one-month break in data collection in October 2019. The duty cycles of the two detectors, H1 and L1, are $\sim 76\%$ and $\sim 77\%$, respectively, during O3.

In the case of a detection, the calibration uncertainties in the detector strain data stream could impact the estimates of the boson properties. Without a detection, these uncertainties could affect the estimated instrument sensitivity and inferred upper limits on astrophysical source properties. The analysis uses the “C01” version of calibrated data, which has estimated maximum uncertainties (68% confidence interval) of $\sim 7\%/\sim 11\%$ in magnitude and $\sim 4\text{ deg}/\sim 9\text{ deg}$ in phase, over the first/second halves of O3 [43, 44]. The frequency-dependent uncertainties vary over the course of a run but do not change by large values. The time-dependent variations could lead to errors in opposite directions at a given frequency during different time periods. In addition, the calibration errors and uncertainties are different at the two LIGO sites. By integrating the data over the whole run, we expect that the impact from the time-dependent, frequency-dependent calibration uncertainties cancels out to some extent, and the overall impact on the inferred upper limits is within a level of $\sim 2\%$. Thus we do not explicitly consider the calibration uncertainties in our analysis.

Due to the presence of a large number of transient noise artifacts, a *gating* procedure [45, 46] has also been applied to the LIGO data. This procedure applies an inverse Tukey window to the LIGO data at times when the root-mean-square value of the whitened strain channel in the band of 25–50 Hz or 70–110 Hz exceeds a certain threshold. Only 0.4% and $\sim 1\%$ of data is removed for Hanford and Livingston, respectively, and the improvement in data quality from applying such gating is significant, as seen in the stochastic and continuous gravitational-wave analyses in O3 [47].

V. RESULTS

In this section, we present the results of the analysis described in Sec. III. With no candidates surviving the follow-up, we present the upper limits on the signal strain amplitude in Sec. V B, and the astrophysical implications in Sec. V C.

Coincident outliers produced by the main search are first pruned in order to reduce their number to a manageable level. A further reduction is based on a close case-by-case inspection of the strongest outliers, as described in the following.

A. Outlier vetos

First, we remove outliers due to *hardware injections*, which simulate the gravitational wave signals expected from spinning neutron stars, added for testing our analysis methods by physically moving the mirrors as if a signal had arrived. See Appendix A for more details.

Moreover, we find several bunches of outliers which are not compatible with astrophysical signals, and thus are also discarded and not followed up. In these cases, the time-frequency peakmaps and spectra show the presence of broadband disturbances or lines. Examples of such disturbances are shown in Fig. 4, where we plot on the left the time-frequency peakmap in the frequency range of 28.2–28.35 Hz over the whole run and on the right the corresponding projection onto the frequency axis. A wide and strong transient disturbance responsible for many outliers with frequency in the range of 28.279–28.283 Hz is clearly visible.

In addition, we find an excess of outliers at several multiples of 10 Hz in both H1 and L1 data, starting from 230 Hz. These are artifacts produced by the procedure to build BSDs⁴. As it is extremely unlikely that the real astrophysical signals are coincident with multiples of 10 Hz, we veto all candidates within a range of $\pm 10^{-4}$ Hz around each multiple of 10 Hz.

Finally, we apply a final threshold on the CR in order to further reduce the number of outliers to a small enough number to permit follow up. In particular, we use different thresholds $\text{CR}_{\text{thr}} = 4.1$ and 4.4 for outliers with $W = 1$ and $W > 1$, respectively. This procedure leaves us with 9449 outliers with $W = 1$ and 7816 outliers with $W > 1$. The choice of threshold does not affect the upper limits presented later, since those are based on the outliers with the highest CR.

Such outliers are followed-up with two different methods: a method based on the Frequency-Hough algorithm if the outliers are characterized by $W = 1$ (Appendix B 1), and a method based on a HMM tracking scheme (the Viterbi algorithm) if the outliers are characterized by $W > 1$ (Appendix B 2). No potential CW candidate remains after the follow-up. Details of the follow-up procedure and results are presented in Appendix C.

⁴ The artifacts are a well-known consequence of the 10 Hz baseline used to build BSD files. They could be avoided by constructing the BSDs with interlaced frequency bands.




FIG. 4. Time-frequency peakmap (left) and corresponding projection (right) in the frequency band of 28.2–28.35 Hz over the full run. Several instrumental artifacts are visible, e.g., strong lines with varying amplitudes (~ 28.22 Hz), weaker lines with sudden frequency variation (~ 28.33 Hz), and broad transient disturbances (vertical bright lines). In particular, the features around 28.28 Hz in the second half of O3 are responsible for a big bunch of outliers with frequency in the range of 28.279–28.283 Hz.

B. Upper limits

Having concluded that none of the outliers is compatible with an astrophysical signal, we compute 95% confidence level (C.L.) upper limits placed on the signal strain amplitude.

The limits are computed, for every 1-Hz sub-band in the range of 20–610 Hz, with the analytic formula of the Frequency-Hough sensitivity, given by Eqn. (67) in [31], where the relevant quantities, i.e. an estimate of the noise power spectral density S_n and the CR, are obtained from the O3 data used in this search. Specifically, for each 1-Hz subband, we take the maximum CR of the outliers, separately for Hanford and Livingston detectors, and the detectors' averaged noise spectra in the same subband. We then compute the two 95% confidence-level upper limits using the equation, and take the worse (i.e. the higher) among the two.

This procedure has been validated by injecting simulated signals into LIGO and Virgo O2 and O3 data [48]. In particular, this procedure has been shown to produce conservative upper limits with respect to those obtained with a much more computationally demanding injection campaign. It has been also verified that the upper limits obtained with a classical procedure based on injections are always above those based on the same analytical formula but using the minimum CR in each 1-Hz subband. The two curves, based, respectively, on the highest and the smallest CR, define a belt containing both a more stringent upper limit estimation and the sensitivity estimation of our search. When we will discuss the astrophysical implications of the search, we will always refer to the upper limits obtained using Eqn. (67) in [31], which are conservative with respect to limits derived from injections. This same procedure is being used for the O3 standard all-sky search for continuous waves from spinning neutron stars [49].

The resulting upper limits, marginalized over the sky and the source polarization parameters, are shown in Fig. 5, as circles connected by a dashed line. The mini-

mum value is 1.04×10^{-25} at 130.5 Hz. In the same figure, the dots connected by the dashed line correspond to the lower bound computed using the minimum CR. A comparison with previous all-sky searches, see e.g. Fig. 4 of [50], shows that our conservative results are better than most of the past searches, including the recent early O3 analysis [50]. In particular, our minimum upper limit value improves upon that in [50] by $\sim 30\%$. The main factors that contribute to the improvement are: 1) the use of the full O3 C01 gated data, 2) the use of longer FFTs at least in a portion of the frequency band, and 3) the restricted spin-up/down range that could impact the maximum CR and thus the upper limit in each subband. On the other hand, the all-sky search described in [50], as well as most of the other past searches, cover a significantly larger parameter space in both the spin-down/up and frequency ranges, and thus are sensitive to signals that are not considered in this analysis. An O2 all-sky search reported in [51] produces upper limits slightly better than ours, by about 5 – 10% on average (due to the use of a much longer coherent integration time of days), but obtained over a smaller parameter space and more limited in scope, being focused on low-ellipticity sources. See Sec. VI for a more detailed discussion.

We want to stress two important points here. First, our conservative procedure to compute upper limits, based on the maximum CR in each subband, produces strain values that are typically larger than the minimum detectable amplitudes at the same frequency. That is, the search sensitivity is expected to be better than the upper limits. Second, this is the first all-sky search that is optimized for frequency wandering signals, both at the level of outlier selection and follow-up. This allows us to achieve a better sensitivity to such signals compared to that achievable using the maximum possible FFT duration (which is the best choice for nearly monochromatic signals).




FIG. 5. 95% C.L. upper limits on the signal strain amplitude obtained from this search (circles, connected by the dashed line). Values are given in each 1-Hz subband. No value is given for the subbands where no outlier survives. The dots connected by a dashed line define the lower bound obtained using the minimum CR in each subband.

C. Astrophysical implications

The upper limits presented above can be translated into physical constraints on the source properties. We interpret the results in two different ways. First, similar to what has been done in [8], we compute the exclusion regions in the boson mass and black hole mass plane, assuming fixed values of the other relevant parameters, namely the distance to the system D , the initial dimensionless spin of the black hole χ_i , and a time equal to the age of the cloud $t = t_{\text{age}}$. Indeed, from Eqs. (9) and (11), once D , χ_i , and t_{age} are fixed, the signal strain amplitude depends only on the boson mass m_b and black hole mass M_{BH} . Thus for each pair of (m_b, M_{BH}) , we can exclude the presence of a source with those assumed parameters, if it would have produced a signal whose strain is larger than the upper limit at a given frequency in figure 5. In our search, the α values probed roughly fall in the range of $[0.02, 0.13]$. Eqs. (4) and (9) do not hold in the range of $\alpha \gtrsim 0.1$, as shown in Fig. 2 of [3]. Specifically, we look at the Brito+ curve derived analytically compared to the black curve obtained from numerical simulations, and see that they differ in energy by a factor of 3 at the largest α considered in this search, $\alpha = 0.15$. This implies that we underestimate τ_{gw} up to a factor of 3 and overestimate h_0 up to a factor of $\sqrt{3}$ (at $\alpha \simeq 0.13$). Thus we correct Eqs. (4) and (9) by these factors such that the resulting exclusion regions are slightly conservative in the

whole parameter space. In Fig. 6, the exclusion regions are plotted for $D = 1$ kpc (left) and $D = 15$ kpc (right), assuming a high initial spin of $\chi_i = 0.9$. For each distance assumption, three different values of the black hole age are considered. In Fig. 7, the exclusion regions are shown for the same parameters as before, except that a lower initial spin value, $\chi_i = 0.5$ is assumed. In both cases, as expected, the constrained region is smaller when the source is assumed to be at a farther distance as, in this case, the signal amplitude at the detector is smaller. For any given black hole mass, these results improve upon the constraints obtained in Advanced LIGO O2 data [8] for lower boson masses, while they are slightly less constraining for higher boson masses. This is because for a given black hole mass, higher boson masses correspond to higher signal spin-up (see Eq. (6)), and we simply do not cover a large spin-up range. On the other hand, the constraints described in [8] were obtained from the results of a search not specifically designed for boson clouds. In particular, restricting the exploration to the parameter space relevant for the expected boson cloud signals significantly reduces the trial factor with a consequent implicit gain in the sensitivity with respect to standard wider parameter space searches.

A second type of interpretation, as shown in Fig. 8, is represented by the maximum distance D_{max} at which we can exclude, as a function of the boson mass, the presence of an emitting system of a given age t_{age} , assuming a




FIG. 6. Exclusion regions in the boson mass (m_b) and black hole mass (M_{BH}) plane for an assumed distance of $D = 1$ kpc (left) and $D = 15$ kpc (right), and an initial black hole dimensionless spin $\chi_i = 0.9$. For $D = 1$ kpc, three possible values of the black hole age, $t_{\text{age}} = 10^3, 10^6, 10^8$ years, are considered; for $D = 15$ kpc, $t_{\text{age}} = 10^3, 10^{4.5}, 10^6$ years are considered.




FIG. 7. Same as Fig. 6 but for black hole initial spin $\chi_i = 0.5$. The assumed distance is $D = 1$ kpc (left), and $D = 15$ kpc (right).

population of black holes. The black hole population has been simulated using a Kroupa mass distribution, with probability density described by $f(m) \propto m^{-2.3}$ [52], with two different ranges, of $[5, 50] M_\odot$ and $[5, 100] M_\odot$, and a uniform initial spin distribution in the range of $[0.2, 0.9]$. The criterion to compute D_{max} at a given boson mass is that at least 5% of the simulated signals would produce a strain amplitude larger than the upper limit at the corresponding frequency in the detectors, and thus would have been detected by our search. This choice of 5% should be conservative, given the large black hole population in the galaxy (10^7 - 10^8) from which we seek a single signal. As before, we correct Eqs. (4) and (9) by a factor of 3 and $\sqrt{3}$, respectively, to obtain more accurate τ_{gw} and h_0 estimates at $\alpha \gtrsim 0.1$ and thus slightly conservative constraints in the full range. As expected, on average when

the maximum black hole mass is smaller, the maximum distance is also smaller, as the signal strain increases with a high power of the black hole mass. Instead of focusing on the specific properties of the emitting system, as for the exclusion regions previously discussed, this constraint on distance depends on the chosen Kroupa mass and uniform spin distributions of the black holes, and thus it reflects the ensemble properties of the assumed black hole population.

Overall, the distance constraints allow us to draw semi-quantitative conclusions on the possible presence of emitting boson clouds in our Galaxy. For instance, young systems, with t_{age} smaller than about 10³ years are disfavored in the whole Galaxy for boson masses above about 2.5×10^{-13} eV for a maximum black hole mass of $50 M_\odot$ and above about 1.2×10^{-13} eV for a maximum black




FIG. 8. Maximum distance at which at least 5% of a simulated population of black holes with a boson cloud would produce a gravitational-wave signal with strain amplitude larger than the upper limit in the detectors. The left plot refers to a maximum black hole mass of $50M_{\odot}$, while the right plot to a maximum mass of $100M_{\odot}$. The different colored markers correspond to different system ages, ranging from 10^3 years to 10^7 years, as indicated in the legend. The alignment of points for different ages at the smallest boson masses (and distances) is the result of a discretization effect due to the finite size grid used in distance.

hole mass of $100M_{\odot}$. As expected, older systems, expected to be more abundant, are more likely to be ruled out only at smaller distances, as they produce on average weaker gravitational-wave signals. As an example, bosons with masses between $\sim 2 \times 10^{-13}$ – 8×10^{-13} eV ($\sim 10^{-13}$ – 8×10^{-13} eV) would be excluded within a distance of 1 kpc from Earth for a maximum black hole mass of $50M_{\odot}$ ($100M_{\odot}$).

The general shape of the maximum distance curves is a result of the combination of different factors pushing in different directions: higher boson masses tend to produce signals with stronger initial amplitudes, but also faster decay rates as a function of t_{age} . Towards the lower m_b end in the plot, D_{max} increases with m_b because the effect of the increasing initial signal amplitudes dominates; towards the higher m_b end, D_{max} tends to decrease again since the faster signal decay rate dominates. At the same time, for higher boson masses, lower black hole masses are required to form a cloud producing signals that last long enough to be detectable. Hence the black hole mass distribution (larger population at lower M_{BH}) comes into play as well. In addition, the results are weighted by the frequency-dependent detector sensitivity (reflected in the upper limit curve in Fig. 5), which also affects the final shape of the D_{max} curves.

For both of the constraints presented above, we do not take into account factors that are largely uncertain but could also be relevant to the interpretation, e.g., the black hole spatial distribution or the formation rate. We defer the integration of these factors to future work when they are better constrained.

Concerning the interpretation of our results with respect to the assumed boson self-interaction, the search covers a small range of spin-up values (see Sec III and, in

particular, Fig. 3), mainly corresponding to a regime of small boson self-interaction, i.e. one in which the spin-up is dominated by Eq (6). For a subset of smaller boson and black hole masses, the spin-up dominated by Eq (7) in the intermediate regime is also covered in the search. However, as we have clarified in Sec II, the signal duration is expected to be shorter in this intermediate regime, and the signal amplitude is smaller due to the smaller boson and black hole masses [see Eq (9)]. Thus the signals in the intermediate regime are less likely to be detected compared to those in the small self-interaction regime.

VI. CONCLUSIONS

This paper describes the first all-sky search tailored to the predicted continuous-wave emission from scalar boson clouds around spinning black holes. We cover a frequency range between 20 Hz and 610 Hz, and a small frequency-dependent spin-up range corresponding to the small self-interaction regime.

We use a multiple frequency-resolution approach, in order to optimize the sensitivity for signals characterized by a slightly wandering frequency, for which a search based on the fixed maximum possible Fourier transform duration would cause a sensitivity loss. No candidate survives the multi-step follow-up procedures we have implemented.

Following up outliers in this analysis was significantly more challenging due, in part, to not having a spin-down or spin-up range to consider, typical of standard all-sky continuous-wave analyses, for which instrumental artifacts lead to clusters over extended ranges in source spin-down. Establishing an instrumental source of such clus-

ters is more straightforward in standard analyses than in this analysis. Manual checks by visual inspection of many spectra had to be performed, along with additional methods necessary to reject an outlier due to a transient spectral artifact in H1 (see section C 1). Therefore, this work not only motivates further searches for boson cloud systems, but also refined methods to mitigate or conclusively veto persistent or transient spectral disturbances at the level of detector commissioning and characterization.

The resulting upper limits are significantly better than those obtained in previous all-sky searches for continuous waves, including a recent search in the early O3 data [50], although our search typically covers a much smaller spin-down/up range with respect to those. On the other hand, our upper limits are slightly worse than those obtained in the O2 search carried out with the Falcon pipeline [51], which however was mainly focused on low-ellipticity sources, and thus covered a much smaller spin-down/up range than we do (from a factor of ~ 4 at lower frequencies up to a factor of ~ 18 at higher frequencies), and moreover did not account for the possibility of signal frequency wandering. The same pipeline was also run from 500–1700 Hz [53] with the same caveats, and although the limits presented in both of these searches are quite stringent, they rely on low-ellipticity sources emitting almost monochromatic gravitational waves.

Code improvements are possible to make the search more sensitive and computationally more efficient. In this way, future boson cloud searches using the basic methodology adopted in this paper will expand the parameter space, covering a wider frequency band and a larger spin-up range, as well as search for vector boson cloud signals that could have much higher spin-ups than those considered here. This, together with the expected detector improvements in upcoming runs of Advanced LIGO, Advanced Virgo and KAGRA detectors, will significantly increase the chance of detecting gravitational radiation from these interesting sources, or at least to better constrain the parameter space of these systems.

Appendix A: Hardware injections

Both H1 and L1 data contain 14 simulated continuous-wave signals, called *hardware injections*, added via hardware for testing purposes. Only one of them, named P5, is within the searched frequency and f range and has an amplitude high enough to be detectable in this search. Indeed we find it with a maximum CR $\simeq 33$ and 36, respectively in the two detectors (and with an averaged dimensionless distance $d \approx 0.7$). As this injected signal is very strong, it produces tens of thousands of outliers at slightly different points in the parameter space. Thus we remove these outliers caused by the hardware injection P5. Fig. 9 shows the outliers found in H1 data caused by this injected signal.




FIG. 9. Critical ratio of the candidates produced by *hardware injection* P5 in H1 data and found in the search with $W = 1$. The distance of the highest CR candidate from the injected signal is about 0.7.

Appendix B: Follow-up methodology

Here we provide details about the two methods used for outlier follow-up. The first one, applied to outliers with $W = 1$, is more suitable for monochromatic or nearly monochromatic signals, while the second works well for signals characterized by some amount of frequency fluctuations. In principle, both methods can be applied to all outliers. The sensitivities of the two methods cannot be directly compared since the two methods assume different signal models. To keep the number of outliers at a practical level for both methods, we only use one method for each category. We assume that for outliers with $W = 1$, the impact from any potential random frequency fluctuation is negligible.

1. Follow-up for outliers with $W = 1$

The follow-up for monochromatic or nearly monochromatic signals consists of several steps, applied to each outlier, and is described in detail in [41]. First, given the frequency and sky coordinates of an outlier, the Doppler modulation of a possible gravitational-wave signal is removed from the data by heterodyne⁵. This allows us to build a new set of FFTs, with a longer duration with respect to the original value, as well as the corresponding peakmap. The factor by which the FFTs duration is increased is a function of the frequency and is shown in Fig. 10. It is chosen as a compromise between the need to improve sensitivity and to keep the computational cost of the follow-up under control. Using different lengthening factors at different frequencies aims to produce similar durations for the new set of FFTs, which

⁵ This is a standard technique that involves multiplying the data by a phase factor $\exp[-j\phi_d(t)]$, where $\phi_d(t)$ is the time-dependent phase evolution induced by the Doppler effect.




FIG. 10. FFT length increase factor, with respect to the main search FFT length, as a function of frequency for the first follow-up stage (FU1). This factor is reduced at the highest frequencies in order to keep the computer memory used by the analysis jobs under control.

would correspond to similar follow-up sensitivity, taking into account that the initial FFTs are shorter at higher frequencies (see Fig. 2). In practice, due to computer memory limitations, we have to limit the increase at the highest frequencies between 500 Hz and 600 Hz.

A follow-up volume, defined by ± 3 frequency bins and ± 3 bins for each sky coordinates (these bins, called *coarse* bins, are those of the initial search setup) around the outlier, is considered. For each sky position and frequency in the follow-up volume, by linear extrapolation of the detector velocity vector, we evaluate the residual Doppler modulation (with respect to the center of the grid), which is corrected in the peakmap by properly shifting each peak. This extrapolation procedure is orders of magnitude faster than the exact Doppler correction and is accurate as long as the follow-up volume is sufficiently small (see [41]).

The Frequency-Hough algorithm [31] is then applied to the resulting ensemble of corrected peakmaps (covering ± 1 coarse bin in \dot{f}). It maps the detector time-frequency plane of the peakmap to the source frequency- \dot{f} plane. The absolute maximum identified among all frequency- \dot{f} Frequency-Hough histograms provides the refined parameters of the original outlier. This procedure is applied to each element of coincident pairs of outliers.

Next, we apply a series of vetoes to each pair of refined outliers, detailed as follows. First, for each detector, a new peakmap is computed using the data coherently corrected with the refined parameters of the outlier and then projected onto the frequency axis. We take the maximum of this projection in a range of ± 2 coarse bins around the outlier frequency⁶.

The rest of a 0.2-Hz band around the outlier, which we consider as the “off-source” region, is divided into a frequency-dependent number of intervals of the same width (in practice, this number varies from about 600 at the lowest frequencies, to about 100 at the highest frequencies). We take the maximum of the peakmap projection in each of these intervals and sort all these maxima in a decreasing order. We keep the outlier if it ranks in the 1% top list in both detectors.

Surviving outliers are then subject to three further tests. The first is a test based on the consistency of the outlier CR in the two detectors, similar to that already used in the main search. The second is a test in which an outlier is discarded, if its CR after the follow-up (in both detectors) drops compared to the original CR from the main search. Finally, we discard coincident outliers with $d > 6$ after the follow-up, indicating that they can no longer be considered as coincident. Outliers surviving this first follow-up round (FU1) are subject to a second round (FU2) in which the same procedure is used but the FFT duration is further increased. Fig. 11 shows the lengthening factor in FU2, as before with respect to the initial duration. The follow-up volume around each out-




FIG. 11. FFT length increase factor, with respect to the main search length, as a function of frequency for the second follow-up stage (FU2).

lier consists, as before, of ± 3 coarse frequency bins, ± 3 bin for each sky coordinate and $\pm 1 \dot{f}$ bin around the outlier (note that these are *coarse* bins of the initial main search). If needed, a third follow-up stage (FU3) along the same line, is applied. The follow-up volume is the same as in previous steps, while the FFT length increase is chosen specifically depending on the outlier parameters. Any outliers surviving these follow-up steps will be eventually analyzed with ad-hoc methods.

⁶ The ± 2 bins used at this level do not have any particular relation

with the ± 3 bins used to define the follow-up volume. The latter choice is due to the uncertainty on the parameters of the outlier, while the former is chosen to estimate the outlier significance.

2. Follow-up for outliers with $W > 1$

Outliers found with $W > 1$ are expected to have some level of frequency evolution that leads to power leakage into adjacent frequency bins. We use the Viterbi method, based on a HMM tracking scheme, which has been used in various continuous-wave searches [9, 10, 54–56]. The Viterbi method can efficiently track secular evolution as well as random fluctuation in the signal frequency, and thus is an ideal follow-up tool to verify candidates with $W > 1$.

For each outlier identified with $W > 1$, we first estimate the maximum \dot{f} value assuming that all the signal power is distributed in W frequency bins,

$$\dot{f}_{\max} = \frac{W}{2T_{\text{obs}}T_{\text{FFT}}}, \quad (\text{B1})$$

and select a coherent time T_{coh} such that the \dot{f}_{\max} can be covered in the Viterbi follow-up search. See details about how to select T_{coh} in Section IV A of [9]. In this follow-up study, three different values of $T_{\text{coh}} = 1d$, $2d$, and $4d$, are used, with shorter T_{coh} applied to larger \dot{f}_{\max} . For $T_{\text{coh}} = 1d$, $2d$, and $4d$, a narrow frequency band of 0.5 Hz, 0.125 Hz, and 0.03125 Hz, centered at the outlier frequency, is searched, respectively. The best estimated sky position in the main search is used for the follow-up analysis. This follow-up procedure consists of three steps.

First, we search in the combined data from LIGO Hanford and Livingston using the Viterbi method with the T_{coh} and frequency band configurations listed above. The detection score S is defined, such that the log likelihood of the optimal Viterbi path equals the mean log likelihood of all paths ending in different frequency bins plus S standard deviations in each band searched [55]. We discard outliers which are not found in the Viterbi follow-up, either S is below the threshold (corresponding to 1% false alarm probability) or the optimal Viterbi path is completely irrelevant to the original outlier (returned at a nonoverlapping frequency). Second, we eliminate the outliers whose significance is higher when analyzing one detector only than combining the two, while searching the other detector alone yields S below the threshold, i.e., the outlier is dominantly louder in a single detector. Third, we carry out manual inspections for any remaining outliers that survive the first two steps.

Appendix C: Candidates and follow-up results

The outliers produced by the main search, with average window $W = 1$ and $W > 1$, are shown, respectively, in Figs. 12 and 13 in terms of their CR as a function of the frequency. It is clear that the majority of the strongest outliers is due to H1 data, and is concentrated in the lower frequency range below ~ 150 Hz. In the following, we present details of the follow-up results for the outliers.

1. Follow-up results for outliers with $W = 1$

For outliers with $W = 1$, we apply the follow-up procedure described in Appendix B 1.

We start with the first follow-up stage (FU1), which produces outliers with refined parameters, on which various vetoes are applied. The application of the significance veto removes most of the outliers, with only 160 surviving. Among these, 14 are discarded as their CR decreases, and an additional 10 are eliminated because they yield a post-follow-up distance of $d > 6$ (i.e., they are no longer coincident).

The CRs of the 136 outliers passing FU1 are shown, as a function of frequency, in Fig. 14. By visual inspection of the refined time-frequency peakmaps produced in FU1, we are able to conclude that many of the strongest outliers are not compatible with an astrophysical signal. In particular, this happens for one outlier at 24.20 Hz, six at ~ 31.14 Hz, two at ~ 37.31 Hz, one at ~ 41.72 Hz, one at ~ 57.59 Hz, three at ~ 69.39 Hz and 22 at ~ 69.55 Hz. As an example, we show in Fig. 15 the refined peakmap built from H1 data, indicating a clear transient disturbance in the second half of the run and responsible for the outliers at ~ 69.55 Hz.

The remaining 97 outliers are analyzed in the second follow-up stage (FU2), which reduces the number of surviving candidates to three (91 are removed by the significance veto and 3 by the distance veto).

The final three remaining outliers have average parameters (averaged values of the parameters obtained from the two detectors) shown in Table I. The first one, at ~ 61.44 Hz, has been discarded by looking at the corresponding refined peakmap, where a transient disturbance in H1 data, starting ~ 20 days before the end of the run and at the exact frequency of the candidate, can be identified, see Fig. 16. The other two have been subject to the third follow-up step (FU3), with an increase of the FFT duration of, respectively, a factor of 10 and 12 with respect to the initial value used in the main search. Neither of the outliers pass the significance veto and, moreover, the CR reduces for both outliers (in both detectors). Thus we finally discard them as not compatible with astrophysical signals.

2. Follow-up results for outliers with $W > 1$

For the original outliers with $W > 1$, we apply the follow-up procedure described in Sec. B 2. In total, 7816 outliers are identified with $W > 1$ in the main search. After the first step of searching data from both LIGO detectors using the Viterbi method, only a small number of the original outliers (267 out of 7816) are identified above the Viterbi threshold (1% false alarm probability in each sub-band searched). After the second step, in which the Viterbi searches are carried out in data from Hanford and Livingston detectors separately, we eliminate an additional 73 outliers, and the remaining 194 are




FIG. 12. Critical ratio of the main search H1 (left) and L1 (right) outliers with $W = 1$.




FIG. 13. Critical ratio of the main search H1 (left) and L1 (right) outliers with $W > 1$.




FIG. 14. Critical ratios of outliers passing FU1 follow-up stage as a function of their frequencies (circles: Livingston, stars: Hanford).

subject to further manual scrutiny.

Finally, we carry out manual inspections for the remaining 194 outliers, clustered at six different frequencies. We find 178 of them clustered near 26.33 Hz are caused by a pulsar hardware injection in O3, and thus

are not of astrophysical origin. Another 15 are associated with various instrumental disturbances. For the last outlier found at ~ 190.82 Hz, further investigations in every few days' of data find that it results from disturbances around that frequency at the Hanford detector in the last ~ 20 days.

ACKNOWLEDGMENTS

This material is based upon work supported by NSF's LIGO Laboratory which is a major facility fully funded by the National Science Foundation. The authors also gratefully acknowledge the support of the Science and Technology Facilities Council (STFC) of the United Kingdom, the Max-Planck-Society (MPS), and the State of Niedersachsen/Germany for support of the construction of Advanced LIGO and construction and operation of the GEO600 detector. Additional support for Advanced LIGO was provided by the Australian Research Council. The authors gratefully acknowledge the Italian Istituto Nazionale di Fisica Nucleare (INFN), the French Centre National de la Recherche Scientifique (CNRS) and the Netherlands Organization for Scientific Research (NWO), for the construction and operation of the Virgo detector and the creation and support of the EGO consortium. The authors also gratefully acknowledge research

frequency [Hz]	λ (deg)	β (deg)	\dot{f} [Hz/s]	CR
61.4416	76.6500	60.5462	$-4.6745 \cdot 10^{-13}$	6.1300
104.1381	85.5924	-59.5761	-5.8598×10^{-13}	4.5150
305.1118	23.2246	50.5265	-5.3121×10^{-12}	4.3600

TABLE I. Average parameters of outliers passing the second follow-up stage (FU2).




FIG. 15. Time-frequency peakmap of H1 data after FU1 follow-up stage showing a distinct broad disturbance appearing in the second half of the run and responsible for the 22 outliers identified with frequency ~ 69.55 Hz. Time is measured in days since the starting of the run. The colorbar represents a power spectrum in units of 10^{-40} [1/Hz].




FIG. 16. Time-frequency peakmap of H1 data after the second follow-up stage (FU2) around the outlier with frequency ~ 61.44 Hz. Time is measured in days since the starting of the run. A transient disturbance, starting from ~ 20 days before the end of the run, is clearly visible. The color bar represents power spectrum in units of 10^{-40} [1/Hz].

support from these agencies as well as by the Council of Scientific and Industrial Research of India, the Department of Science and Technology, India, the Science & Engineering Research Board (SERB), India, the Ministry of Human Resource Development, India, the Spanish Agencia Estatal de Investigación (AEI), the Spanish Ministerio de Ciencia e Innovación and Ministerio de Universidades, the Conselleria de Fons Europeus, Universitat i Cultura and the Direcció General de Política Universitaria i Recerca del Govern de les Illes Balears, the Conselleria d'Innovació, Universitats, Ciència i Societat Digital de la Generalitat Valenciana and the CERCA Programme Generalitat de Catalunya, Spain, the National Science Centre of Poland and the European Union – European Regional Development Fund; Foundation for Polish Science (FNP), the Swiss National Science Foundation (SNSF), the Russian Foundation for Basic Research, the Russian Science Foundation, the European Commission, the European Social Funds (ESF), the European Regional Development Funds (ERDF), the Royal Society, the Scottish Funding Council, the Scottish Universities Physics Alliance, the Hungarian Scientific Research Fund (OTKA), the French Lyon Institute of Origins (LIO), the Belgian Fonds de la Recherche Scientifique (FRS-FNRS), Actions de Recherche Concertées

(ARC) and Fonds Wetenschappelijk Onderzoek – Vlaanderen (FWO), Belgium, the Paris Île-de-France Region, the National Research, Development and Innovation Office Hungary (NKFIH), the National Research Foundation of Korea, the Natural Science and Engineering Research Council Canada, Canadian Foundation for Innovation (CFI), the Brazilian Ministry of Science, Technology, and Innovations, the International Center for Theoretical Physics South American Institute for Fundamental Research (ICTP-SAIFR), the Research Grants Council of Hong Kong, the National Natural Science Foundation of China (NSFC), the Leverhulme Trust, the Research Corporation, the Ministry of Science and Technology (MOST), Taiwan, the United States Department of Energy, and the Kavli Foundation. The authors gratefully acknowledge the support of the NSF, STFC, INFN and CNRS for provision of computational resources.

This work was supported by MEXT, JSPS Leading-edge Research Infrastructure Program, JSPS Grant-in-Aid for Specially Promoted Research 26000005, JSPS Grant-in-Aid for Scientific Research on Innovative Areas 2905: JP17H06358, JP17H06361 and JP17H06364, JSPS Core-to-Core Program A. Advanced Research Networks, JSPS Grant-in-Aid for Scientific Research (S) 17H06133 and 20H05639 , JSPS Grant-in-Aid for Trans-

formative Research Areas (A) 20A203: JP20H05854, the joint research program of the Institute for Cosmic Ray Research, University of Tokyo, National Research Foundation (NRF) and Computing Infrastructure Project of KISTI-GSDC in Korea, Academia Sinica (AS), AS Grid Center (ASGC) and the Ministry of Science and Technology (MoST) in Taiwan under grants including AS-CDA-

105-M06, Advanced Technology Center (ATC) of NAOJ, Mechanical Engineering Center of KEK.

We would like to thank all of the essential workers who put their health at risk during the COVID-19 pandemic, without whom we would not have been able to complete this work.

-
- [1] A. Arvanitaki, M. Baryakhtar, and X. Huang, “Discovering the qcd axion with black holes and gravitational waves,” *Physical Review D*, vol. 91, no. 8, p. 084011, 2015.
- [2] R. Brito, V. Cardoso, and P. Pani, “Superradiance: New Frontiers in Black Hole Physics,” *Lect. Notes Phys.*, vol. 906, pp. pp.1–237, 2015.
- [3] R. Brito, S. Ghosh, E. Barausse, E. Berti, V. Cardoso, I. Dvorkin, A. Klein, and P. Pani, “Gravitational wave searches for ultralight bosons with LIGO and LISA,” *Phys. Rev. D*, vol. 96, no. 6, p. 064050, 2017.
- [4] M. Baryakhtar, R. Lasenby, and M. Teo, “Black hole superradiance signatures of ultralight vectors,” *Physical Review D*, vol. 96, no. 3, p. 035019, 2017.
- [5] N. Siemonsen and W. E. East, “Gravitational wave signatures of ultralight vector bosons from black hole superradiance,” *Physical Review D*, vol. 101, no. 2, p. 024019, 2020.
- [6] R. Brito, S. Grillo, and P. Pani, “Black Hole Superradiant Instability from Ultralight Spin-2 Fields,” *Phys. Rev. Lett.*, vol. 124, no. 21, p. 211101, 2020.
- [7] M. Baryakhtar, M. Galanis, R. Lasenby, and O. Simon, “Black hole superradiance of self-interacting scalar fields,” *Phys. Rev. D*, vol. 103, p. 095019, May 2021.
- [8] C. Palomba, S. D’Antonio, P. Astone, S. Frasca, G. Intini, I. La Rosa, P. Leaci, S. Mastrogiovanni, A. L. Miller, F. Muciaccia, et al., “Direct constraints on the ultralight boson mass from searches of continuous gravitational waves,” *Physical Review Letters*, vol. 123, no. 17, p. 171101, 2019.
- [9] M. Isi, L. Sun, R. Brito, and A. Melatos, “Directed searches for gravitational waves from ultralight bosons,” *Physical Review D*, vol. 99, no. 8, p. 084042, 2019.
- [10] L. Sun, R. Brito, and M. Isi, “Search for ultralight bosons in Cygnus X-1 with Advanced LIGO,” *Phys. Rev. D*, vol. 101, no. 6, p. 063020, 2020. [Erratum: *Phys. Rev. D* 102, 089902 (2020)].
- [11] S. J. Zhu, M. Baryakhtar, M. A. Papa, D. Tsuna, N. Kawanaka, and H.-B. Eggenstein, “Characterizing the continuous gravitational-wave signal from boson clouds around galactic isolated black holes,” *Phys. Rev. D*, vol. 102, p. 063020, Sep 2020.
- [12] L. Pierini and et al. *in preparation*.
- [13] S. Vitale, R. Lynch, J. Veitch, V. Raymond, and R. Sturani, “Measuring the spin of black holes in binary systems using gravitational waves,” *Phys. Rev. Lett.*, vol. 112, no. 25, p. 251101, 2014.
- [14] M. Pürer, M. Hannam, and F. Ohme, “Can we measure individual black-hole spins from gravitational-wave observations?,” in *14th Marcel Grossmann Meeting on Recent Developments in Theoretical and Experimental General Relativity, Astrophysics, and Relativistic Field Theories*, 2017.
- [15] K. Belczynski et al., “Evolutionary roads leading to low effective spins, high black hole masses, and O1/O2 rates for LIGO/Virgo binary black holes,” *Astron. Astrophys.*, vol. 636, p. A104, 2020.
- [16] D. Gerosa, E. Berti, R. O’Shaughnessy, K. Belczynski, M. Kesden, D. Wysocki, and W. Gladysz, “Spin orientations of merging black holes formed from the evolution of stellar binaries,” *Phys. Rev. D*, vol. 98, no. 8, p. 084036, 2018.
- [17] K. K. Y. Ng, O. A. Hannuksela, S. Vitale, and T. G. F. Li, “Searching for ultralight bosons within spin measurements of a population of binary black hole mergers,” *Phys. Rev. D*, vol. 103, no. 6, p. 063010, 2021.
- [18] K. K. Y. Ng, S. Vitale, O. A. Hannuksela, and T. G. F. Li, “Constraints on Ultralight Scalar Bosons within Black Hole Spin Measurements from the LIGO-Virgo GWTC-2,” *Phys. Rev. Lett.*, vol. 126, no. 15, p. 151102, 2021.
- [19] D. Baumann, H. S. Chia, and R. A. Porto, “Probing ultralight bosons with binary black holes,” *Physical Review D*, vol. 99, no. 4, p. 044001, 2019.
- [20] Q. Yang, L.-W. Ji, B. Hu, Z.-J. Cao, and R.-G. Cai, “An axion-like scalar field environment effect on binary black hole merger,” *Res. Astron. Astrophys.*, vol. 18, no. 6, p. 065, 2018.
- [21] S. Choudhary, N. Sanchis-Gual, A. Gupta, J. C. De-gollado, S. Bose, and J. A. Font, “Gravitational waves from binary black hole mergers surrounded by scalar field clouds: Numerical simulations and observational implications,” *Phys. Rev. D*, vol. 103, no. 4, p. 044032, 2021.
- [22] K. K. Y. Ng, M. Isi, C.-J. Haster, and S. Vitale, “Multi-band gravitational-wave searches for ultralight bosons,” *Phys. Rev. D*, vol. 102, no. 8, p. 083020, 2020.
- [23] R. Brito, S. Ghosh, E. Barausse, E. Berti, V. Cardoso, I. Dvorkin, A. Klein, and P. Pani, “Stochastic and resolvable gravitational waves from ultralight bosons,” *Phys. Rev. Lett.*, vol. 119, no. 13, p. 131101, 2017.
- [24] L. Tsukada, T. Callister, A. Matas, and P. Meyers, “First search for a stochastic gravitational-wave background from ultralight bosons,” *Phys. Rev. D*, vol. 99, no. 10, p. 103015, 2019.
- [25] L. Tsukada, R. Brito, W. E. East, and N. Siemonsen, “Modeling and searching for a stochastic gravitational-wave background from ultralight vector bosons,” 11 2020.
- [26] A. Pierce, Z. Zhang, Y. Zhao, et al., “Dark photon dark matter produced by axion oscillations,” *Physical Review D*, vol. 99, no. 7, p. 075002, 2019.
- [27] A. L. Miller et al., “Probing new light gauge bosons with gravitational-wave interferometers using an adapted semicoherent method,” *Phys. Rev. D*, vol. 103, no. 10, p. 103002, 2021.
- [28] H.-K. Guo, K. Riles, F.-W. Yang, and Y. Zhao, “Search-

- ing for dark photon dark matter in ligo o1 data,” *Communications Physics*, vol. 2, no. 1, pp. 1–7, 2019.
- [29] R. Abbott *et al.*, “Constraints on dark photon dark matter using data from LIGO’s and Virgo’s third observing run,” 5 2021.
- [30] J. Aasi, B. P. Abbott, R. Abbott, T. Abbott, M. R. Abernathy, K. Ackley, C. Adams, T. Adams, P. Addesso, and et al., “Advanced LIGO,” *CQGra*, vol. 32, p. 074001, Apr. 2015.
- [31] P. Astone, A. Colla, S. D’Antonio, S. Frasca, and C. Palomba, “Method for all-sky searches of continuous gravitational wave signals using the Frequency-Hough transform,” *Physical Review D*, vol. 90, no. 4, p. 042002, 2014.
- [32] K. Riles, “Recent searches for continuous gravitational waves,” *Modern Physics Letters A*, vol. 32, no. 39, p. 1730035, 2017.
- [33] M. Sieniawska *et al.*, “Continuous waves from neutron stars: current status and prospects,” *Universe*, vol. 5, no. 11, p. 217, 2019.
- [34] S. D’Antonio, C. Palomba, P. Astone, S. Frasca, G. Intini, I. La Rosa, P. Leaci, S. Mastrogiovanni, A. Miller, F. Muciaccia, *et al.*, “Semicohherent analysis method to search for continuous gravitational waves emitted by ultralight boson clouds around spinning black holes,” *Physical Review D*, vol. 98, no. 10, p. 103017, 2018.
- [35] K. S. Thorne, C. W. Misner, and J. A. Wheeler, *Gravitation*. Freeman, 2000.
- [36] J. D. Bekenstein, “Extraction of energy and charge from a black hole,” *Phys. Rev. D*, vol. 7, pp. 949–953, 1973.
- [37] O. Piccinni, P. Astone, S. D’Antonio, S. Frasca, G. Intini, P. Leaci, S. Mastrogiovanni, A. Miller, C. Palomba, and A. Singhal, “A new data analysis framework for the search of continuous gravitational wave signals,” *Classical and Quantum Gravity*, vol. 36, no. 1, p. 015008, 2018.
- [38] O. J. Piccinni, P. Astone, S. D’Antonio, S. Frasca, G. Intini, I. La Rosa, P. Leaci, S. Mastrogiovanni, A. Miller, and C. Palomba, “Directed search for continuous gravitational-wave signals from the galactic center in the advanced ligo second observing run,” *Physical Review D*, vol. 101, no. 8, p. 082004, 2020.
- [39] R. Abbott, T. Abbott, *et al.*, “Searches for continuous gravitational waves from young supernova remnants in the early third observing run of advanced ligo and virgo,” *arXiv:2105.11641*, 2021.
- [40] R. Abbott, T. Abbott, S. Abraham, F. Acernese, K. Ackley, A. Adams, C. Adams, R. Adhikari, V. Adya, C. Affeldt, *et al.*, “Gravitational-wave constraints on the equatorial ellipticity of millisecond pulsars,” *arXiv preprint arXiv:2007.14251*, 2020.
- [41] B. Abbott *et al.*, “All-sky search for continuous gravitational waves from isolated neutron stars using Advanced LIGO O2 data,” *Physical Review D*, vol. 100, no. 2, p. 024004, 2019.
- [42] B. P. Abbott *et al.*, “All-sky search for periodic gravitational waves in the o1 ligo data,” *Physical Review D*, vol. 96, no. 6, p. 062002, 2017.
- [43] L. Sun *et al.*, “Characterization of systematic error in Advanced LIGO calibration,” *Class. Quant. Grav.*, vol. 37, no. 22, p. 225008, 2020.
- [44] L. Sun *et al.*, “Characterization of systematic error in Advanced LIGO calibration in the second half of O3,” 2021.
- [45] <https://dcc.ligo.org/T2000384/>.
- [46] <https://dcc.ligo.org/T2000512/>.
- [47] R. Abbott *et al.*, “Upper Limits on the Isotropic Gravitational-Wave Background from Advanced LIGO’s and Advanced Virgo’s Third Observing Run,” 1 2021.
- [48] M. Di Cesare, *Master Thesis*. 2021.
- [49] LVK *in preparation*.
- [50] R. Abbott *et al.*, “All-sky search for continuous gravitational waves from isolated neutron stars in the early o3 ligo data,” 2021.
- [51] V. Dergachev and M. A. Papa, “The search for continuous gravitational waves from small-ellipticity sources at low frequencies,” *Physical Review D*, vol. 104, p. 043003, 2021.
- [52] O. D. Ebert, B. J. S., and M. Kaplinghat, “Counting black holes: The cosmic stellar remnant population and implications for ligo,” *MNRAS*, vol. 120, no. 473, p. 1186, 2018.
- [53] V. Dergachev and M. A. Papa, “Results from the First All-Sky Search for Continuous Gravitational Waves from Small-Ellipticity Sources,” *Phys. Rev. Lett.*, vol. 125, no. 17, p. 171101, 2020.
- [54] S. Suvorova, L. Sun, A. Melatos, W. Moran, and R. J. Evans, “Hidden Markov model tracking of continuous gravitational waves from a neutron star with wandering spin,” *Physical Review D*, vol. D93, no. 12, p. 123009, 2016.
- [55] L. Sun, A. Melatos, S. Suvorova, W. Moran, and R. Evans, “Hidden Markov model tracking of continuous gravitational waves from young supernova remnants,” *Physical Review D*, vol. D97, no. 4, p. 043013, 2018.
- [56] L. Sun and A. Melatos, “Application of hidden markov model tracking to the search for long-duration transient gravitational waves from the remnant of the binary neutron star merger gw170817,” *Physical Review D*, vol. 99, no. 12, p. 123003, 2019.

The LIGO Scientific Collaboration, Virgo Collaboration, and KAGRA Collaboration

R. Abbott,¹ H. Abe,² F. Acernese,^{3,4} K. Ackley,⁵ N. Adhikari,⁶ R. X. Adhikari,¹ V. K. Adkins,⁷ V. B. Adya,⁸ C. Affeldt,^{9,10} D. Agarwal,¹¹ M. Agathos,^{12,13} K. Agatsuma,¹⁴ N. Aggarwal,¹⁵ O. D. Aguilar,¹⁶ L. Aiello,¹⁷ A. Ain,¹⁸ P. Ajith,¹⁹ T. Akutsu,^{20,21} S. Albanesi,^{22,23} R. A. Alfaidi,²⁴ A. Allocata,^{25,4} P. A. Altin,⁸ A. Amato,²⁶ C. Anand,⁵ S. Anand,¹ A. Ananyeva,¹ S. B. Anderson,¹ W. G. Anderson,⁶ M. Ando,^{27,28} T. Andrade,²⁹ N. Andres,³⁰ M. Andrés-Carcasona,³¹ T. Andrić,³² S. V. Angelova,³³ S. Ansoldi,^{34,35} J. M. Antelis,³⁶ S. Antier,^{37,38} T. Apostolatos,³⁹ E. Z. Appavuravther,^{40,41} S. Appert,¹ S. K. Apple,⁴² K. Arai,¹ A. Araya,⁴³ M. C. Araya,¹ J. S. Areeda,⁴⁴ M. Arène,⁴⁵ N. Aritomi,²⁰ N. Arnaud,^{46,47} M. Arogeti,⁴⁸ S. M. Aronson,⁷ K. G. Arun,⁴⁹ H. Asada,⁵⁰ Y. Asali,⁵¹ G. Ashton,⁵² Y. Aso,^{1,53,54} M. Assiduo,^{55,56} S. Assis de Souza Melo,⁴⁷ S. M. Aston,⁵⁷ P. Astone,⁵⁸ F. Aubin,⁵⁶ K. AultONeal,^{1,36} C. Austin,⁷ S. Babak,⁴⁵ F. Badaracco,⁵⁹ M. K. M. Bader,⁶⁰ C. Badger,⁶¹ S. Bae,⁶² Y. Bae,⁶³ A. M. Baer,⁶⁴ S. Bagnasco,²³ Y. Bai,¹ J. Baird,⁴⁵ R. Bajpai,⁶⁵ T. Baka,⁶⁶ M. Ball,⁶⁷ G. Ballardin,⁴⁷ S. W. Ballmer,⁶⁸ A. Balsamo,⁶⁴ G. Baltus,⁶⁹ S. Banagiri,¹⁵ B. Banerjee,³² D. Bankar,¹¹ J. C. Barayoga,¹ C. Barbieri,^{70,71,72} B. C. Barish,¹ D. Barker,⁷³ P. Barneo,²⁹ F. Barone,^{74,4} B. Barr,²⁴ L. Barsotti,⁷⁵ M. Barsuglia,⁴⁵ D. Barta,⁷⁶ J. Bartlett,⁷³ M. A. Barton,²⁴ I. Bartos,⁷⁷ S. Basak,¹⁹ R. Bassiri,⁷⁸ A. Basti,^{79,18} M. Bawaj,^{1,40,80} J. C. Bayley,²⁴ M. Bazzan,^{81,82} B. R. Becher,⁸³ B. Bécsy,⁸⁴ V. M. Bedakihale,⁸⁵ F. Beirnaert,⁸⁶ M. Bejger,^{1,87} I. Belahcene,⁴⁶ V. Benedetto,⁸⁸ D. Beniwal,⁸⁹ M. G. Benjamin,⁹⁰ T. F. Bennett,⁹¹ J. D. Bentley,^{1,14} M. BenYaala,³³ S. Bera,¹¹ M. Berbel,⁹² F. Bergamin,^{9,10} B. K. Berger,^{1,78} S. Bernuzzi,^{1,13} D. Bersanetti,^{1,93} A. Bertolini,⁶⁰ J. Betzwieser,^{1,57} D. Beveridge,^{1,94} R. Bhandare,⁹⁵ A. V. Bhandari,¹¹ U. Bhardwaj,^{1,38,60} R. Bhatt,¹ D. Bhattacharjee,^{1,96} S. Bhaumik,^{1,77} A. Bianchi,^{60,97} I. A. Bilenko,⁹⁸ G. Billingsley,^{1,1} S. Bini,^{99,100} R. Birney,¹⁰¹ O. Birnholtz,^{1,102} S. Biscans,^{1,75} M. Bischi,^{55,56} S. Biscoveanu,^{1,75} A. Bisht,^{9,10} B. Biswas,¹¹ M. Bitossi,^{47,18} M.-A. Bizouard,^{1,37} J. K. Blackburn,¹ C. D. Blair,⁹⁴ D. G. Blair,⁹⁴ R. M. Blair,⁷³ F. Bobba,^{103,104} N. Bode,^{9,10} M. Boér,³⁷ G. Bogaert,³⁷ M. Boldrini,^{105,58} G. N. Bolingbrooke,^{1,89} L. D. Bonavena,⁸¹ F. Bondu,¹⁰⁶ E. Bonilla,^{1,78} R. Bonnand,^{1,30} P. Booker,^{9,10} B. A. Boom,⁶⁰ R. Bork,¹ V. Boschi,^{1,18} N. Bose,¹⁰⁷ S. Bose,¹¹ V. Bossilkov,⁹⁴ V. Boudart,^{1,69} Y. Bouffanais,^{81,82} A. Bozzi,⁴⁷ C. Bradaschia,¹⁸ P. R. Brady,^{1,6} A. Bramley,⁵⁷ A. Branch,⁵⁷ M. Branchesi,^{1,32,108} J. E. Brau,^{1,67} M. Breschi,^{1,13} T. Briant,^{1,109} J. H. Briggs,²⁴ A. Brillet,³⁷ M. Brinkmann,^{9,10} P. Brockill,⁶ A. F. Brooks,^{1,1} J. Brooks,⁴⁷ D. D. Brown,⁸⁹ S. Brunett,¹ G. Bruno,⁵⁹ R. Bruntz,^{1,64} J. Bryant,¹⁴ F. Bucci,⁵⁶ T. Bulik,¹¹⁰ H. J. Bulten,⁶⁰ A. Buonanno,^{1,111,112} K. Burtnyk,⁷³ R. Buscicchio,^{1,14} D. Buskulic,³⁰ C. Buy,^{1,113} R. L. Byer,⁷⁸ G. S. Cabourn Davies,^{1,52} G. Cabras,^{1,34,35} R. Cabrita,^{1,59} L. Cadonati,^{1,48} M. Caesar,^{1,114} G. Cagnoli,^{1,26} C. Cahillane,^{1,73} J. Calderón Bustillo,^{1,115} J. D. Callaghan,²⁴ T. A. Callister,^{1,116,117} E. Calloni,^{1,25,4} J. Cameron,⁹⁴ J. B. Camp,^{1,118} M. Canepa,^{1,119,93} S. Canevarolo,⁶⁶ M. Cannavacciuolo,^{1,103} K. C. Cannon,^{1,28} H. Cao,⁸⁹ Z. Cao,^{1,120} E. Capocasa,^{1,45,20} E. Capote,⁶⁸ G. Carapella,^{1,103,104} F. Carbognani,^{1,47} M. Carlassara,^{1,9,10} J. B. Carlin,^{1,121} M. F. Carney,^{1,15} M. Carpinelli,^{1,122,123,47} G. Carrillo,^{1,67} G. Carullo,^{1,79,18} T. L. Carver,^{1,17} J. Casanueva Diaz,^{1,47} C. Casentini,^{1,124,125} G. Castaldi,^{1,126} S. Caudill,^{1,60,66} M. Cavaglià,^{1,96} F. Cavalier,^{1,46} R. Cavalieri,^{1,47} G. Cella,^{1,18} P. Cerdá-Durán,^{1,127} E. Cesarini,^{1,125} W. Chaibi,^{1,37} S. Chalathadka Subrahmanyam,^{1,128} E. Champion,^{1,129} C.-H. Chan,^{1,130} C. Chan,^{1,28} C. L. Chan,^{1,131} K. Chan,^{1,131} M. Chan,^{1,132} K. Chandra,^{1,107} I. P. Chang,^{1,130} P. Chanial,^{1,47} S. Chao,^{1,130} C. Chapman-Bird,^{1,24} P. Charlton,^{1,133} E. A. Chase,^{1,15} E. Chassande-Mottin,^{1,45} C. Chatterjee,^{1,1,94} Debarati Chatterjee,^{1,11} Deep Chatterjee,^{1,6} M. Chaturvedi,^{1,95} S. Chaty,^{1,45} C. Chen,^{1,134,135} D. Chen,^{1,53} H. Y. Chen,^{1,75} J. Chen,^{1,130} K. Chen,^{1,136} X. Chen,^{1,94} Y.-B. Chen,^{1,137} Y.-R. Chen,^{1,138} Z. Chen,^{1,17} H. Cheng,^{1,77} C. K. Cheong,^{1,131} H. Y. Cheung,^{1,131} H. Y. Chia,^{1,77} F. Chiadini,^{1,139,104} C.-Y. Chiang,^{1,140} G. Chiarini,^{1,82} R. Chierici,^{1,141} A. Chincarini,^{1,93} M. L. Chiofalo,^{1,79,18} A. Chiummo,^{1,47} R. K. Choudhary,^{1,94} S. Choudhary,^{1,11} N. Christensen,^{1,37} Q. Chu,^{1,94} Y.-K. Chu,^{1,140} S. S. Y. Chua,^{1,8} K. W. Chung,^{1,61} G. Ciani,^{1,81,82} P. Ciecielag,^{1,87} M. Cieślar,^{1,87} M. Cifaldi,^{1,124,125} A. A. Ciobanu,^{1,89} R. Ciolfi,^{1,142,82} F. Cipriano,^{1,37} F. Clara,^{1,73} J. A. Clark,^{1,1,48} P. Clearwater,^{1,143} S. Clesse,^{1,144} F. Cleva,^{1,37} E. Coccia,^{1,32,108} E. Codazzo,^{1,32} P.-F. Cohadon,^{1,109} D. E. Cohen,^{1,46} M. Colleoni,^{1,145} C. G. Collette,^{1,146} A. Colombo,^{1,70,71} M. Colpi,^{1,70,71} C. M. Compton,^{1,73} M. Constancio Jr.,^{1,16} L. Conti,^{1,82} S. J. Cooper,^{1,14} P. Corban,^{1,57} T. R. Corbitt,^{1,7} I. Cordero-Carrión,^{1,147} S. Corezzi,^{1,80,40} K. R. Corley,^{1,51} N. J. Cornish,^{1,84} D. Corre,^{1,46} A. Corsi,^{1,148} S. Cortese,^{1,47} C. A. Costa,^{1,16} R. Cotesta,^{1,112} R. Cottingham,^{1,57} M. W. Coughlin,^{1,149} J.-P. Coulon,^{1,37} S. T. Countryman,^{1,51} B. Cousins,^{1,150} P. Couvares,^{1,1} D. M. Coward,^{1,94} M. J. Cowart,^{1,57} D. C. Coyne,^{1,1} R. Coyne,^{1,151} J. D. E. Creighton,^{1,6} T. D. Creighton,^{1,90} A. W. Criswell,^{1,149} M. Croquette,^{1,109} S. G. Crowder,^{1,152} J. R. Cudell,^{1,69} T. J. Cullen,^{1,7} A. Cumming,^{1,24} R. Cummings,^{1,24} L. Cunningham,^{1,24} E. Cuoco,^{1,47,153,18} M. Curylo,^{1,110} P. Dabadie,^{1,26} T. Dal Canton,^{1,46} S. Dall'Osso,^{1,32} G. Dálya,^{1,86,154} A. Dana,^{1,78} B. D'Angelo,^{1,119,93} S. Danilishin,^{1,155,60}

- S. D'Antonio,¹²⁵ K. Danzmann,^{9, 10} C. Darsow-Fromm ,¹²⁸ A. Dasgupta,⁸⁵ L. E. H. Datrier,²⁴ Sayak Datta,¹¹ Sayantani Datta ,⁴⁹ V. Dattilo,⁴⁷ I. Dave,⁹⁵ M. Davier,⁴⁶ D. Davis ,¹ M. C. Davis ,¹¹⁴ E. J. Daw ,¹⁵⁶ R. Dean,¹¹⁴ D. DeBra,⁷⁸ M. Deenadayalan,¹¹ J. Degallaix ,¹⁵⁷ M. De Laurentis,^{25, 4} S. Deléglise ,¹⁰⁹ V. Del Favero,¹²⁹ F. De Lillo ,⁵⁹ N. De Lillo,²⁴ D. Dell'Aquila ,¹²² W. Del Pozzo,^{79, 18} L. M. DeMarchi,¹⁵ F. De Matteis,^{124, 125} V. D'Emilio,¹⁷ N. Demos,⁷⁵ T. Dent ,¹¹⁵ A. Depasse ,⁵⁹ R. De Pietri ,^{158, 159} R. De Rosa ,^{25, 4} C. De Rossi,⁴⁷ R. DeSalvo ,^{126, 160} R. De Simone,¹³⁹ S. Dhurandhar,¹¹ M. C. Díaz ,⁹⁰ N. A. Didio,⁶⁸ T. Dietrich ,¹¹² L. Di Fiore,⁴ C. Di Fronzo,¹⁴ C. Di Giorgio ,^{103, 104} F. Di Giovanni ,¹²⁷ M. Di Giovanni,³² T. Di Girolamo ,^{25, 4} A. Di Lieto ,^{79, 18} A. Di Michele ,⁸⁰ B. Ding,¹⁴⁶ S. Di Pace ,^{105, 58} I. Di Palma ,^{105, 58} F. Di Renzo ,^{79, 18} A. K. Divakarla,⁷⁷ A. Dmitriev ,¹⁴ Z. Doctor,¹⁵ L. Donahue,¹⁶¹ L. D'Onofrio ,^{25, 4} F. Donovan,⁷⁵ K. L. Dooley,¹⁷ S. Doravari ,¹¹ M. Drago ,^{105, 58} J. C. Driggers ,⁷³ Y. Drori,¹ J.-G. Ducoin,⁴⁶ P. Dupej,²⁴ U. Dupletsza,³² O. Durante,^{103, 104} D. D'Urso ,^{122, 123} P.-A. Duverne,⁴⁶ S. E. Dwyer,⁷³ C. Eassa,⁷³ P. J. Easter,⁵ M. Ebersold,¹⁶² T. Eckhardt ,¹²⁸ G. Eddolls ,²⁴ B. Edelman ,⁶⁷ T. B. Edo,¹ O. Edy ,⁵² A. Effler ,⁵⁷ S. Eguchi ,¹³² J. Eichholz ,⁸ S. S. Eikenberry,⁷⁷ M. Eisenmann,^{30, 20} R. A. Eisenstein,⁷⁵ A. Ejlli ,¹⁷ E. Engelby,⁴⁴ Y. Enomoto ,²⁷ L. Errico,^{25, 4} R. C. Essick ,¹⁶³ H. Estellés,¹⁴⁵ D. Estevez ,¹⁶⁴ Z. Etienne,¹⁶⁵ T. Etzel,¹ M. Evans ,⁷⁵ T. M. Evans,⁵⁷ T. Evstafyeva,¹² B. E. Ewing,¹⁵⁰ F. Fabrizi ,^{55, 56} F. Faedi,⁵⁶ V. Fafone ,^{124, 125, 32} H. Fair,⁶⁸ S. Fairhurst,¹⁷ P. C. Fan ,¹⁶¹ A. M. Farah ,¹⁶⁶ S. Farinon,⁹³ B. Farr ,⁶⁷ W. M. Farr ,^{116, 117} E. J. Fauchon-Jones,¹⁷ G. Favaro ,⁸¹ M. Favata ,¹⁶⁷ M. Fays ,⁶⁹ M. Fazio,¹⁶⁸ J. Feicht,¹ M. M. Fejer,⁷⁸ E. Fenyvesi ,^{76, 169} D. L. Ferguson ,¹⁷⁰ A. Fernandez-Galiana ,⁷⁵ I. Ferrante ,^{79, 18} T. A. Ferreira,¹⁶ F. Fidecaro ,^{79, 18} P. Figura ,¹¹⁰ A. Fiori ,^{18, 79} I. Fiori ,⁴⁷ M. Fishbach ,¹⁵ R. P. Fisher,⁶⁴ R. Fittipaldi,^{171, 104} V. Fiumara,^{172, 104} R. Flaminio,^{30, 20} E. Floden,¹⁴⁹ H. K. Fong,²⁸ J. A. Font ,^{127, 173} B. Fornal ,¹⁶⁰ P. W. F. Forsyth,⁸ A. Franke,¹²⁸ S. Frasca,^{105, 58} F. Frasconi ,¹⁸ J. P. Freed,³⁶ Z. Frei ,¹⁵⁴ A. Freise ,^{60, 97} O. Freitas,¹⁷⁴ R. Frey ,⁶⁷ P. Fritschel,⁷⁵ V. V. Frolov,⁵⁷ G. G. Fronzé ,²³ Y. Fujii,¹⁷⁵ Y. Fujikawa,¹⁷⁶ Y. Fujimoto,¹⁷⁷ P. Fulda,⁷⁷ M. Fyffe,⁵⁷ H. A. Gabbard,²⁴ B. U. Gadre ,¹¹² J. R. Gair ,¹¹² J. Gais,¹³¹ S. Galaudage,⁵ R. Gamba,¹³ D. Ganapathy ,⁷⁵ A. Ganguly ,¹¹ D. Gao ,¹⁷⁸ S. G. Gaonkar,¹¹ B. Garaventa ,^{93, 119} C. García Núñez,¹⁰¹ C. García-Quirós,¹⁴⁵ F. Garufi ,^{25, 4} B. Gateley,⁷³ V. Gayathri,⁷⁷ G.-G. Ge ,¹⁷⁸ G. Gemme ,⁹³ A. Gennai ,¹⁸ J. George,⁹⁵ O. Gerberding ,¹²⁸ L. Gergely ,¹⁷⁹ P. Gewecke,¹²⁸ S. Ghonge ,⁴⁸ Abhirup Ghosh ,¹¹² Archisman Ghosh ,⁸⁶ Shaon Ghosh ,¹⁶⁷ Shrobona Ghosh,¹⁷ Tathagata Ghosh ,¹¹ B. Giacomazzo ,^{70, 71, 72} L. Giacoppo,^{105, 58} J. A. Giaime ,^{7, 57} K. D. Giardina,⁵⁷ D. R. Gibson,¹⁰¹ C. Gier,³³ M. Giesler ,¹⁸⁰ P. Giri ,^{18, 79} F. Gissi,⁸⁸ S. Gkaitatzis ,^{18, 79} J. Glanzer,⁷ A. E. Gleckl,⁴⁴ P. Godwin,¹⁵⁰ E. Goetz ,¹⁸¹ R. Goetz ,⁷⁷ N. Gohlke,^{9, 10} J. Golomb,¹ B. Goncharov ,³² G. González ,⁷ M. Gosselin,⁴⁷ R. Gouaty,³⁰ D. W. Gould,⁸ S. Goyal,¹⁹ B. Grace,⁸ A. Grado ,^{182, 4} V. Graham,²⁴ M. Granata ,¹⁵⁷ V. Granata,¹⁰³ A. Grant,²⁴ S. Gras,⁷⁵ P. Grassia,¹ C. Gray,⁷³ R. Gray ,²⁴ G. Greco,⁴⁰ A. C. Green ,⁷⁷ R. Green,¹⁷ A. M. Gretarsson,³⁶ E. M. Gretarsson,³⁶ D. Griffith,¹ W. L. Griffiths ,¹⁷ H. L. Griggs ,⁴⁸ G. Grignani,^{80, 40} A. Grimaldi ,^{99, 100} E. Grimes,³⁶ S. J. Grimm,^{32, 108} H. Grote ,¹⁷ S. Grunewald,¹¹² P. Gruning,⁴⁶ A. S. Gruson,⁴⁴ D. Guerra ,¹²⁷ G. M. Guidi ,^{55, 56} A. R. Guimaraes,⁷ G. Guixé,²⁹ H. K. Gulati,⁸⁵ A. M. Gunny,⁷⁵ H.-K. Guo ,¹⁶⁰ Y. Guo,⁶⁰ Anchal Gupta,¹ Anuradha Gupta ,¹⁸³ I. M. Gupta,¹⁵⁰ P. Gupta,^{60, 66} S. K. Gupta,¹⁰⁷ R. Gustafson,¹⁸⁴ F. Guzman ,¹⁸⁵ S. Ha,¹⁸⁶ I. P. W. Hadiputrawan,¹³⁶ L. Haegel ,⁴⁵ S. Haino,¹⁴⁰ O. Halim ,³⁵ E. D. Hall ,⁷⁵ E. Z. Hamilton,¹⁶² G. Hammond,²⁴ W.-B. Han ,¹⁸⁷ M. Haney ,¹⁶² J. Hanks,⁷³ C. Hanna,¹⁵⁰ M. D. Hannam,¹⁷ O. Hannuksela,^{66, 60} H. Hansen,⁷³ T. J. Hansen,³⁶ J. Hanson,⁵⁷ T. Harder,³⁷ K. Haris,^{60, 66} J. Harms ,^{32, 108} G. M. Harry ,⁴² I. W. Harry ,⁵² D. Hartwig ,¹²⁸ K. Hasegawa,¹⁸⁸ B. Haskell,⁸⁷ C.-J. Haster ,⁷⁵ J. S. Hathaway,¹²⁹ K. Hattori,¹⁸⁹ K. Haughian,²⁴ H. Hayakawa,¹⁹⁰ K. Hayama,¹³² F. J. Hayes,²⁴ J. Healy ,¹²⁹ A. Heidmann ,¹⁰⁹ A. Heidt,^{9, 10} M. C. Heintze,⁵⁷ J. Heinze ,^{9, 10} J. Heinzel,⁷⁵ H. Heitmann ,³⁷ F. Hellman ,¹⁹¹ P. Hello,⁴⁶ A. F. Helmling-Cornell ,⁶⁷ G. Hemming ,⁴⁷ M. Hendry ,²⁴ I. S. Heng,²⁴ E. Hennes ,⁶⁰ J. Hennig,¹⁹² M. H. Hennig ,¹⁹² C. Henshaw,⁴⁸ A. G. Hernandez,⁹¹ F. Hernandez Vivanco,⁵ M. Heurs ,^{9, 10} A. L. Hewitt ,¹⁹³ S. Higginbotham,¹⁷ S. Hild,^{155, 60} P. Hill,³³ Y. Himemoto,¹⁹⁴ A. S. Hines,¹⁸⁵ N. Hirata,²⁰ C. Hirose,¹⁷⁶ T.-C. Ho,¹³⁶ S. Hochheim,^{9, 10} D. Hofman,¹⁵⁷ J. N. Hohmann,¹²⁸ D. G. Holcomb ,¹¹⁴ N. A. Holland,⁸ I. J. Hollows ,¹⁵⁶ Z. J. Holmes ,⁸⁹ K. Holt,⁵⁷ D. E. Holz ,¹⁶⁶ Q. Hong,¹³⁰ J. Hough,²⁴ S. Hourihane,¹ E. J. Howell ,⁹⁴ C. G. Hoy ,¹⁷ D. Hoyland,¹⁴ A. Hreibi,^{9, 10} B.-H. Hsieh,¹⁸⁸ H.-F. Hsieh ,¹⁹⁵ C. Hsiung,¹³⁴ Y. Hsu,¹³⁰ H.-Y. Huang ,¹⁴⁰ P. Huang ,¹⁷⁸ Y.-C. Huang ,¹³⁸ Y.-J. Huang ,¹⁴⁰ Yiting Huang,¹⁵² Yiwen Huang,⁷⁵ M. T. Hübner ,⁵ A. D. Huddart,¹⁹⁶ B. Hughey,³⁶ D. C. Y. Hui ,¹⁹⁷ V. Hui ,³⁰ S. Husa,¹⁴⁵ S. H. Huttner,²⁴ R. Huxford,¹⁵⁰ T. Huynh-Dinh,⁵⁷ S. Ide,¹⁹⁸ B. Idzkowski ,¹¹⁰ A. Iess,^{124, 125} K. Inayoshi ,¹⁹⁹ Y. Inoue,¹³⁶ P. Iosif ,²⁰⁰ M. Isi ,⁷⁵ K. Isleif,¹²⁸ K. Ito,²⁰¹ Y. Itoh ,^{177, 202} B. R. Iyer ,¹⁹ V. JaberianHamedan ,⁹⁴ T. Jacqmin ,¹⁰⁹ P.-E. Jacquet ,¹⁰⁹ S. J. Jadhav,²⁰³ S. P. Jadhav ,¹¹ T. Jain,¹²

- A. L. James,¹⁷ A. Z. Jan,¹⁷⁰ K. Jani,²⁰⁴ J. Janquart,^{66, 60} K. Janssens,^{205, 37} N. N. Janthalur,²⁰³
 P. Jaranowski,²⁰⁶ D. Jariwala,⁷⁷ R. Jaume,¹⁴⁵ A. C. Jenkins,⁶¹ K. Jenner,⁸⁹ C. Jeon,²⁰⁷ W. Jia,⁷⁵
 J. Jiang,⁷⁷ H.-B. Jin,^{208, 209} G. R. Johns,⁶⁴ R. Johnston,²⁴ A. W. Jones,⁹⁴ D. I. Jones,²¹⁰ P. Jones,¹⁴
 R. Jones,²⁴ P. Joshi,¹⁵⁰ L. Ju,⁹⁴ A. Jue,¹⁶⁰ P. Jung,⁶³ K. Jung,¹⁸⁶ J. Junker,^{9, 10} V. Juste,¹⁶⁴ K. Kaihotsu,²⁰¹
 T. Kajita,²¹¹ M. Kakizaki,¹⁸⁹ C. V. Kalaghatgi,^{17, 66, 60, 212} V. Kalogera,¹⁵ B. Kamai,¹ M. Kamiizumi,¹⁹⁰
 N. Kanda,^{177, 202} S. Kandhasamy,¹¹ G. Kang,²¹³ J. B. Kanner,¹ Y. Kao,¹³⁰ S. J. Kapadia,¹⁹ D. P. Kapasi,⁸
 C. Karathanasis,³¹ S. Karki,⁹⁶ R. Kashyap,¹⁵⁰ M. Kasprzack,¹ W. Kastaun,^{9, 10} T. Kato,¹⁸⁸ S. Katsanevas,⁴⁷
 E. Katsavounidis,⁷⁵ W. Katzman,⁵⁷ T. Kaur,⁹⁴ K. Kawabe,⁷³ K. Kawaguchi,¹⁸⁸ F. Kéfélian,³⁷ D. Keitel,¹⁴⁵
 J. S. Key,²¹⁴ S. Khadka,⁷⁸ F. Y. Khalili,⁹⁸ S. Khan,¹⁷ T. Khanam,¹⁴⁸ E. A. Khazanov,²¹⁵ N. Khetan,^{32, 108}
 M. Khursheed,⁹⁵ N. Kijbunchoo,⁸ A. Kim,¹⁵ C. Kim,²⁰⁷ J. C. Kim,²¹⁶ J. Kim,²¹⁷ K. Kim,²⁰⁷ W. S. Kim,⁶³
 Y.-M. Kim,¹⁸⁶ C. Kimball,¹⁵ N. Kimura,¹⁹⁰ M. Kinley-Hanlon,²⁴ R. Kirchhoff,^{9, 10} J. S. Kissel,⁷³
 S. Klimentko,⁷⁷ T. Klinger,¹² A. M. Knee,¹⁸¹ T. D. Knowles,¹⁶⁵ N. Knust,^{9, 10} E. Knyazev,⁷⁵ Y. Kobayashi,¹⁷⁷
 P. Koch,^{9, 10} G. Koekoek,^{60, 155} K. Kohri,²¹⁸ K. Kokeyama,²¹⁹ S. Koley,³² P. Kolitsidou,¹⁷ M. Kolstein,³¹
 K. Komori,⁷⁵ V. Kondrashov,¹ A. K. H. Kong,¹⁹⁵ A. Kontos,⁸³ N. Koper,^{9, 10} M. Korobko,¹²⁸ M. Kovalam,⁹⁴
 N. Koyama,¹⁷⁶ D. B. Kozak,¹ C. Kozakai,⁵³ V. Kringel,^{9, 10} N. V. Krishnendu,¹⁹ A. Królak,^{220, 221}
 G. Kuehn,^{9, 10} F. Kuei,¹³⁰ P. Kuijper,⁶⁰ S. Kulkarni,¹⁸³ A. Kumar,²⁰³ Prayush Kumar,¹⁹ Rahul Kumar,⁷³
 Rakesh Kumar,⁸⁵ J. Kume,²⁸ K. Kuns,⁷⁵ Y. Kuromiya,²⁰¹ S. Kuroyanagi,^{222, 223} K. Kwak,¹⁸⁶
 G. Lacaille,²⁴ P. Lagabbe,³⁰ D. Laghi,¹¹³ E. Lalande,²²⁴ M. Lalleman,²⁰⁵ T. L. Lam,¹³¹ A. Lamberts,^{37, 225}
 M. Landry,⁷³ B. B. Lane,⁷⁵ R. N. Lang,⁷⁵ J. Lange,¹⁷⁰ B. Lantz,⁷⁸ I. La Rosa,³⁰ A. Lartaux-Vollard,⁴⁶
 P. D. Lasky,⁵ M. Laxen,⁵⁷ A. Lazzarini,¹ C. Lazzaro,^{81, 82} P. Leaci,^{105, 58} S. Leavey,^{9, 10} S. LeBohec,¹⁶⁰
 Y. K. Lecoeuche,¹⁸¹ E. Lee,¹⁸⁸ H. M. Lee,²²⁶ H. W. Lee,²¹⁶ K. Lee,²²⁷ R. Lee,¹³⁸ I. N. Legred,¹
 J. Lehmann,^{9, 10} A. Lemaître,²²⁸ M. Lenti,^{56, 229} M. Leonardi,²⁰ E. Leonova,³⁸ N. Leroy,⁴⁶ N. Letendre,³⁰
 C. Levesque,²²⁴ Y. Levin,⁵ J. N. Leviton,¹⁸⁴ K. Leyde,⁴⁵ A. K. Y. Li,¹ B. Li,¹³⁰ J. Li,¹⁵ K. L. Li,²³⁰ P. Li,²³¹
 T. G. F. Li,¹³¹ X. Li,¹³⁷ C.-Y. Lin,²³² E. T. Lin,¹⁹⁵ F.-K. Lin,¹⁴⁰ F.-L. Lin,²³³ H. L. Lin,¹³⁶
 L. C.-C. Lin,²³⁰ F. Linde,^{212, 60} S. D. Linker,^{126, 91} J. N. Linley,²⁴ T. B. Littenberg,²³⁴ G. C. Liu,¹³⁴ J. Liu,⁹⁴
 K. Liu,¹³⁰ X. Liu,⁶ F. Llamas,⁹⁰ R. K. L. Lo,¹ T. Lo,¹³⁰ L. T. London,^{38, 75} A. Longo,²³⁵ D. Lopez,¹⁶²
 M. Lopez Portilla,⁶⁶ M. Lorenzini,^{124, 125} V. Loriette,²³⁶ M. Lormand,⁵⁷ G. Losurdo,¹⁸ T. P. Lott,⁴⁸
 J. D. Lough,^{9, 10} C. O. Lousto,¹²⁹ G. Lovelace,⁴⁴ J. F. Lucaccioni,²³⁷ H. Lück,^{9, 10} D. Lumaca,^{124, 125}
 A. P. Lundgren,⁵² L.-W. Luo,¹⁴⁰ J. E. Lynam,⁶⁴ M. Ma'arif,¹³⁶ R. Macas,⁵² J. B. Machtlinger,¹⁵ M. MacInnis,⁷⁵
 D. M. Macleod,¹⁷ I. A. O. MacMillan,¹ A. Macquet,³⁷ I. Magaña Hernandez,⁶ C. Magazzù,¹⁸
 R. M. Magee,¹ R. Maggiore,¹⁴ M. Magnozzi,^{93, 119} S. Mahesh,¹⁶⁵ E. Majorana,^{105, 58} I. Maksimovic,²³⁶
 S. Maliakal,¹ A. Malik,⁹⁵ N. Man,³⁷ V. Mandic,¹⁴⁹ V. Mangano,^{105, 58} G. L. Mansell,^{73, 75} M. Manske,⁶
 M. Mantovani,⁴⁷ M. Mapelli,^{81, 82} F. Marchesoni,^{41, 40, 238} D. Marín Pina,²⁹ F. Marion,³⁰ Z. Mark,¹³⁷
 S. Márka,⁵¹ Z. Márka,⁵¹ C. Markakis,¹² A. S. Markosyan,⁷⁸ A. Markowitz,¹ E. Maros,¹ A. Marquina,¹⁴⁷
 S. Marsat,⁴⁵ F. Martelli,^{55, 56} I. W. Martin,²⁴ R. M. Martin,¹⁶⁷ M. Martinez,³¹ V. A. Martinez,⁷⁷ V. Martinez,²⁶
 K. Martinovic,⁶¹ D. V. Martynov,¹⁴ E. J. Marx,⁷⁵ H. Masalehdan,¹²⁸ K. Mason,⁷⁵ E. Massera,¹⁵⁶ A. Masserot,³⁰
 M. Masso-Reid,²⁴ S. Mastrogiovanni,⁴⁵ A. Matas,¹¹² M. Mateu-Lucena,¹⁴⁵ F. Matichard,^{1, 75}
 M. Matiushechkina,^{9, 10} N. Mavalvala,⁷⁵ J. J. McCann,⁹⁴ R. McCarthy,⁷³ D. E. McClelland,¹⁸
 P. K. McClintey,¹⁵⁰ S. McCormick,⁵⁷ L. McCuller,⁷⁵ G. I. McGhee,²⁴ S. C. McGuire,⁵⁷ C. McIsaac,⁵²
 J. McIver,¹⁸¹ T. McRae,⁸ S. T. McWilliams,¹⁶⁵ D. Meacher,⁶ M. Mehmet,^{9, 10} A. K. Mehta,¹¹²
 Q. Meijer,⁶⁶ A. Melatos,¹²¹ D. A. Melchor,⁴⁴ G. Mendell,⁷³ A. Menendez-Vazquez,³¹ C. S. Menoni,¹⁶⁸
 R. A. Mercer,⁶ L. Mereni,¹⁵⁷ K. Merfeld,⁶⁷ E. L. Merilh,⁵⁷ J. D. Merritt,⁶⁷ M. Merzougui,³⁷ S. Meshkov,^{1, *}
 C. Messenger,²⁴ C. Messick,⁷⁵ P. M. Meyers,¹²¹ F. Meylahn,^{9, 10} A. Mhaske,¹¹ A. Miami,^{99, 100} H. Miao,¹⁴
 I. Michaloliakos,⁷⁷ C. Michel,¹⁵⁷ Y. Michimura,²⁷ H. Middleton,¹²¹ D. P. Mihaylov,¹¹² L. Milano,^{25, †}
 A. L. Miller,⁵⁹ A. Miller,⁹¹ B. Miller,^{38, 60} M. Millhouse,¹²¹ J. C. Mills,¹⁷ E. Milotti,^{239, 35} Y. Minenkov,¹²⁵
 N. Mio,²⁴⁰ Ll. M. Mir,³¹ M. Miravet-Tenés,¹²⁷ A. Mishkin,⁷⁷ C. Mishra,²⁴¹ T. Mishra,⁷⁷ T. Mistry,¹⁵⁶
 S. Mitra,¹¹ V. P. Mitrofanov,⁹⁸ G. Mitselmakher,⁷⁷ R. Mittleman,⁷⁵ O. Miyakawa,¹⁹⁰ K. Miyo,¹⁹⁰
 S. Miyoki,¹⁹⁰ Geoffrey Mo,⁷⁵ L. M. Modaffer,¹⁴⁵ E. Moguel,²³⁷ K. Mogushi,⁹⁶ S. R. P. Mohapatra,⁷⁵
 S. R. Mohite,⁶ I. Molina,⁴⁴ M. Molina-Ruiz,¹⁹¹ M. Mondin,⁹¹ M. Montani,^{55, 56} C. J. Moore,¹⁴ J. Moragues,¹⁴⁵
 D. Moraru,⁷³ F. Morawski,⁸⁷ A. More,¹¹ C. Moreno,³⁶ G. Moreno,⁷³ Y. Mori,²⁰¹ S. Morisaki,⁶ N. Morisue,¹⁷⁷
 Y. Moriwaki,¹⁸⁹ B. Mours,¹⁶⁴ C. M. Mow-Lowry,^{60, 97} S. Mozzon,⁵² F. Muciaccia,^{105, 58} Arunava Mukherjee,²⁴²
 D. Mukherjee,¹⁵⁰ Soma Mukherjee,⁹⁰ Subroto Mukherjee,⁸⁵ Suvodip Mukherjee,^{163, 38} N. Mukund,^{9, 10}
 A. Mullavey,⁵⁷ J. Munch,⁸⁹ E. A. Muñiz,⁶⁸ P. G. Murray,²⁴ R. Musenich,^{93, 119} S. Muusse,⁸⁹ S. L. Nadji,^{9, 10}

- K. Nagano ^{1D},²⁴³ A. Nagar,^{23, 244} K. Nakamura ^{1D},²⁰ H. Nakano,²⁴⁵ M. Nakano,¹⁸⁸ Y. Nakayama,²⁰¹ V. Napolano,⁴⁷
 I. Nardecchia ^{1D},^{124, 125} T. Narikawa,¹⁸⁸ H. Narola,⁶⁶ L. Naticchioni ^{1D},⁵⁸ B. Nayak,⁹¹ R. K. Nayak ^{1D},²⁴⁶ B. F. Neil,⁹⁴
 J. Neilson,^{88, 104} A. Nelson,¹⁸⁵ T. J. N. Nelson,⁵⁷ M. Nery,^{9, 10} P. Neubauer,²³⁷ A. Neunzert,²¹⁴ K. Y. Ng,⁷⁵
 S. W. S. Ng ^{1D},⁸⁹ C. Nguyen ^{1D},⁴⁵ P. Nguyen,⁶⁷ T. Nguyen,⁷⁵ L. Nguyen Quynh ^{1D},²⁴⁷ J. Ni,¹⁴⁹ W.-T. Ni ^{1D},^{208, 178, 138}
 S. A. Nichols,⁷ T. Nishimoto,¹⁸⁸ A. Nishizawa ^{1D},²⁸ S. Nissanke,^{38, 60} E. Nitoglia ^{1D},¹⁴¹ F. Nocera,⁴⁷
 M. Norman,¹⁷ C. North,¹⁷ S. Nozaki,¹⁸⁹ G. Nurbek,⁹⁰ L. K. Nuttall ^{1D},⁵² Y. Obayashi ^{1D},¹⁸⁸ J. Oberling,⁷³
 B. D. O'Brien,⁷⁷ J. O'Dell,¹⁹⁶ E. Oelker ^{1D},²⁴ W. Ogaki,¹⁸⁸ G. Oganesyan,^{32, 108} J. J. Oh ^{1D},⁶³ K. Oh ^{1D},¹⁹⁷
 S. H. Oh ^{1D},⁶³ M. Ohashi ^{1D},¹⁹⁰ T. Ohashi,¹⁷⁷ M. Ohkawa ^{1D},¹⁷⁶ F. Ohme ^{1D},^{9, 10} H. Ohta,²⁸ M. A. Okada,¹⁶
 Y. Okutani,¹⁹⁸ C. Olivetto,⁴⁷ K. Oohara ^{1D},^{188, 248} R. Oram,⁵⁷ B. O'Reilly ^{1D},⁵⁷ R. G.Ormiston,¹⁴⁹
 N. D. Ormsby,⁶⁴ R. O'Shaughnessy ^{1D},¹²⁹ E. O'Shea ^{1D},¹⁸⁰ S. Oshino ^{1D},¹⁹⁰ S. Ossokine ^{1D},¹¹² C. Osthelder,¹
 S. Otabe,² D. J. Ottaway ^{1D},⁸⁹ H. Overmier,⁵⁷ A. E. Pace,¹⁵⁰ G. Pagano,^{79, 18} R. Pagano,⁷ M. A. Page,⁹⁴
 G. Pagliaroli,^{32, 108} A. Pai,¹⁰⁷ S. A. Pai,⁹⁵ S. Pal,²⁴⁶ J. R. Palamos,⁶⁷ O. Palashov,²¹⁵ C. Palomba ^{1D},⁵⁸ H. Pan,¹³⁰
 K.-C. Pan ^{1D},^{138, 195} P. K. Panda,²⁰³ P. T. H. Pang,^{60, 66} C. Pankow,¹⁵ F. Pannarale ^{1D},^{105, 58} B. C. Pant,⁹⁵
 F. H. Panther,⁹⁴ F. Paoletti ^{1D},¹⁸ A. Paoli,⁴⁷ A. Paolone,^{58, 249} G. Pappas,²⁰⁰ A. Parisi ^{1D},¹³⁴ H. Park,⁶ J. Park ^{1D},²⁵⁰
 W. Parker ^{1D},⁵⁷ D. Pascucci ^{1D},^{60, 86} A. Pasqualetti,⁴⁷ R. Passaquieti ^{1D},^{79, 18} D. Passuello,¹⁸ M. Patel,⁶⁴ M. Pathak,⁸⁹
 B. Patricelli ^{1D},^{47, 18} A. S. Patron,⁷ S. Paul ^{1D},⁶⁷ E. Payne,⁵ M. Pedraza,¹ R. Pedurand,¹⁰⁴ M. Pegoraro,⁸²
 A. Pele,⁵⁷ F. E. Peña Arellano ^{1D},¹⁹⁰ S. Penano,⁷⁸ S. Penn ^{1D},²⁵¹ A. Perego,^{99, 100} A. Pereira,²⁶ T. Pereira ^{1D},²⁵²
 C. J. Perez,⁷³ C. Périsoigis,³⁰ C. C. Perkins,⁷⁷ A. Perreca ^{1D},^{99, 100} S. Perriès,¹⁴¹ D. Pesios,²⁰⁰ J. Petermann ^{1D},¹²⁸
 D. Petterson,¹ H. P. Pfeiffer ^{1D},¹¹² H. Pham,⁵⁷ K. A. Pham ^{1D},¹⁴⁹ K. S. Phukon ^{1D},^{60, 212} H. Phurailatpam,¹³¹
 O. J. Piccinni ^{1D},⁵⁸ M. Pichot ^{1D},³⁷ M. Piendibene,^{79, 18} F. Piergiovanni,^{55, 56} L. Pierini ^{1D},^{105, 58} V. Pierro ^{1D},^{88, 104}
 G. Pillant,⁴⁷ M. Pillas,⁴⁶ F. Pilo,¹⁸ L. Pinard,¹⁵⁷ C. Pineda-Bosque,⁹¹ I. M. Pinto,^{88, 104, 253} M. Pinto,⁴⁷
 B. J. Piotrkowski,⁶ K. Piotrkowski,⁵⁹ M. Pirello,⁷³ M. D. Pitkin ^{1D},¹⁹³ A. Placidi ^{1D},^{40, 80} E. Placidi,^{105, 58}
 M. L. Planas ^{1D},¹⁴⁵ W. Plastino ^{1D},^{254, 235} C. Pluchar,²⁵⁵ R. Poggiani ^{1D},^{79, 18} E. Polini ^{1D},³⁰ D. Y. T. Pong,¹³¹
 S. Ponrathnam,¹¹ E. K. Porter,⁴⁵ R. Poulton ^{1D},⁴⁷ A. Poverman,⁸³ J. Powell,¹⁴³ M. Pracchia,³⁰ T. Pradier,¹⁶⁴
 A. K. Prajapati,⁸⁵ K. Prasai,⁷⁸ R. Prasanna,²⁰³ G. Pratten ^{1D},¹⁴ M. Principe,^{88, 253, 104} G. A. Prodi ^{1D},^{256, 100}
 L. Prokhorov,¹⁴ P. Prosposito,^{124, 125} L. Prudenzi,¹¹² A. Puecher,^{60, 66} M. Punturo ^{1D},⁴⁰ F. Puosi,^{18, 79} P. Puppo,⁵⁸
 M. Pürer ^{1D},¹¹² H. Qi ^{1D},¹⁷ N. Quartey,⁶⁴ V. Quetschke,⁹⁰ P. J. Quinonez,³⁶ R. Quitzow-James,⁹⁶ F. J. Raab,⁷³
 G. Raaijmakers,^{38, 60} H. Radkins,⁷³ N. Radulesco,³⁷ P. Raffai ^{1D},¹⁵⁴ S. X. Rail,²²⁴ S. Raja,⁹⁵ C. Rajan,⁹⁵
 K. E. Ramirez ^{1D},⁵⁷ T. D. Ramirez,⁴⁴ A. Ramos-Buades ^{1D},¹¹² J. Rana,¹⁵⁰ P. Rapagnani,^{105, 58} A. Ray,⁶
 V. Raymond ^{1D},¹⁷ N. Raza ^{1D},¹⁸¹ M. Razzano ^{1D},^{79, 18} J. Read,⁴⁴ L. A. Rees,⁴² T. Regimbau,³⁰ L. Rei ^{1D},⁹³ S. Reid,³³
 S. W. Reid,⁶⁴ D. H. Reitze,^{1, 77} P. Relton ^{1D},¹⁷ A. Renzini,¹ P. Rettewno ^{1D},^{22, 23} B. Revenu ^{1D},⁴⁵ A. Reza,⁶⁰
 M. Rezac,⁴⁴ F. Ricci,^{105, 58} D. Richards,¹⁹⁶ J. W. Richardson ^{1D},²⁵⁷ L. Richardson,¹⁸⁵ G. Riemenschneider,^{22, 23}
 K. Riles ^{1D},¹⁸⁴ S. Rinaldi ^{1D},^{79, 18} K. Rink ^{1D},¹⁸¹ N. A. Robertson,¹ R. Robie,¹ F. Robinet,⁴⁶ A. Rocchi ^{1D},¹²⁵
 S. Rodriguez,⁴⁴ L. Rolland ^{1D},³⁰ J. G. Rollins ^{1D},¹ M. Romanelli,¹⁰⁶ R. Romano,^{3, 4} C. L. Romel,⁷³ A. Romero ^{1D},³¹
 I. M. Romero-Shaw,⁵ J. H. Romie,⁵⁷ S. Ronchini ^{1D},^{32, 108} L. Rosa,^{4, 25} C. A. Rose,⁶ D. Rosińska,¹¹⁰ M. P. Ross ^{1D},²⁵⁸
 S. Rowan,²⁴ S. J. Rowlinson,¹⁴ S. Roy,⁶⁶ Santosh Roy,¹¹ Soumen Roy,²⁵⁹ D. Rozza ^{1D},^{122, 123} P. Ruggi,⁴⁷
 K. Ruiz-Rocha,²⁰⁴ K. Ryan,⁷³ S. Sachdev,¹⁵⁰ T. Sadecki,⁷³ J. Sadiq ^{1D},¹¹⁵ S. Saha ^{1D},¹⁹⁵ Y. Saito,¹⁹⁰ K. Sakai,²⁶⁰
 M. Sakellariadou ^{1D},⁶¹ S. Sakon,¹⁵⁰ O. S. Salafia ^{1D},^{72, 71, 70} F. Salces-Carcoba ^{1D},¹ L. Salconi,⁴⁷ M. Saleem ^{1D},¹⁴⁹
 F. Salemi ^{1D},^{99, 100} A. Samajdar ^{1D},⁷¹ E. J. Sanchez,¹ J. H. Sanchez,⁴⁴ L. E. Sanchez,¹ N. Sanchis-Gual ^{1D},²⁶¹
 J. R. Sanders,²⁶² A. Sanuy ^{1D},²⁹ T. R. Saravanan,¹¹ N. Sarin,⁵ B. Sassolas,¹⁵⁷ H. Satari,⁹⁴ O. Sauter ^{1D},⁷⁷
 R. L. Savage ^{1D},⁷³ V. Savant,¹¹ T. Sawada ^{1D},¹⁷⁷ H. L. Sawant,¹¹ S. Sayah,¹⁵⁷ D. Schaetzl,¹ M. Scheel,¹³⁷
 J. Scheuer,¹⁵ M. G. Schiworski ^{1D},⁸⁹ P. Schmidt ^{1D},¹⁴ S. Schmidt,⁶⁶ R. Schnabel ^{1D},¹²⁸ M. Schneewind,^{9, 10}
 R. M. S. Schofield,⁶⁷ A. Schönbeck,¹²⁸ B. W. Schulte,^{9, 10} B. F. Schutz,^{17, 9, 10} E. Schwartz ^{1D},¹⁷ J. Scott ^{1D},²⁴
 S. M. Scott ^{1D},⁸ M. Seglar-Arroyo ^{1D},³⁰ Y. Sekiguchi ^{1D},²⁶³ D. Sellers,⁵⁷ A. S. Sengupta,²⁵⁹ D. Sentenac,⁴⁷ E. G. Seo,¹³¹
 V. Sequino,^{25, 4} A. Sergeev,²¹⁵ Y. Setyawati ^{1D},^{9, 10, 66} T. Shaffer,⁷³ M. S. Shahriar ^{1D},¹⁵ M. A. Shaikh ^{1D},¹⁹
 B. Shams,¹⁶⁰ L. Shao ^{1D},¹⁹⁹ A. Sharma,^{32, 108} P. Sharma,⁹⁵ P. Shawhan ^{1D},¹¹¹ N. S. Shcheblanov ^{1D},²²⁸ A. Sheela,²⁴¹
 Y. Shikano ^{1D},^{264, 265} M. Shikauchi,²⁸ H. Shimizu ^{1D},²⁶⁶ K. Shimode ^{1D},¹⁹⁰ H. Shinkai ^{1D},²⁶⁷ T. Shishido,⁵⁴ A. Shoda ^{1D},²⁰
 D. H. Shoemaker ^{1D},⁷⁵ D. M. Shoemaker ^{1D},¹⁷⁰ S. ShyamSundar,⁹⁵ M. Sieniawska,⁵⁹ D. Sigg ^{1D},⁷³ L. Silenzi ^{1D},^{40, 41}
 L. P. Singer ^{1D},¹¹⁸ D. Singh ^{1D},¹⁵⁰ M. K. Singh ^{1D},¹⁹ N. Singh ^{1D},¹¹⁰ A. Singha ^{1D},^{155, 60} A. M. Sintes ^{1D},¹⁴⁵
 V. Sipala,^{122, 123} V. Skliris,¹⁷ B. J. J. Slagmolen ^{1D},⁸ T. J. Slaven-Blair,⁹⁴ J. Smetana,¹⁴ J. R. Smith ^{1D},⁴⁴ L. Smith,²⁴
 R. J. E. Smith ^{1D},⁵ J. Soldateschi ^{1D},^{229, 268, 56} S. N. Somala ^{1D},²⁶⁹ K. Somiya ^{1D},² I. Song ^{1D},¹⁹⁵ K. Soni ^{1D},¹¹
 V. Sordini,¹⁴¹ F. Sorrentino,⁹³ N. Sorrentino ^{1D},^{79, 18} R. Soulard,³⁷ T. Souradeep,^{270, 11} E. Sowell,¹⁴⁸
 V. Spagnuolo,^{155, 60} A. P. Spencer ^{1D},²⁴ M. Spera ^{1D},^{81, 82} P. Spinicelli,⁴⁷ A. K. Srivastava,⁸⁵ V. Srivastava,⁶⁸

- K. Staats,¹⁵ C. Stachie,³⁷ F. Stachurski,²⁴ D. A. Steer ,⁴⁵ J. Steinlechner,^{155, 60} S. Steinlechner ,^{155, 60}
 N. Stergioulas,²⁰⁰ D. J. Stops,¹⁴ M. Stover,²³⁷ K. A. Strain ,²⁴ L. C. Strang,¹²¹ G. Stratta ,^{271, 58}
 M. D. Strong,⁷ A. Strunk,⁷³ R. Sturani,²⁵² A. L. Stuver ,¹¹⁴ M. Suchenek,⁸⁷ S. Sudhagar ,¹¹ V. Sudhir ,⁷⁵
 R. Sugimoto ,^{272, 243} H. G. Suh ,⁶ A. G. Sullivan ,⁵¹ T. Z. Summerscales ,²⁷³ L. Sun ,⁸ S. Sunil,⁸⁵
 A. Sur ,⁸⁷ J. Suresh ,²⁸ P. J. Sutton ,¹⁷ Takamasa Suzuki ,¹⁷⁶ Takanori Suzuki,² Toshikazu Suzuki,¹⁸⁸
 B. L. Swinkels ,⁶⁰ M. J. Szczepańczyk ,⁷⁷ P. Szewczyk,¹¹⁰ M. Tacca,⁶⁰ H. Tagoshi,¹⁸⁸ S. C. Tait ,²⁴
 H. Takahashi ,²⁷⁴ R. Takahashi ,²⁰ S. Takano,²⁷ H. Takeda ,²⁷ M. Takeda,¹⁷⁷ C. J. Talbot,³³ C. Talbot,¹
 K. Tanaka,²⁷⁵ Taiki Tanaka,¹⁸⁸ Takahiro Tanaka ,²⁷⁶ A. J. Tanasiyczuk,⁵⁹ S. Tanioka ,¹⁹⁰ D. B. Tanner,⁷⁷
 D. Tao,¹ L. Tao ,⁷⁷ R. D. Tapia,¹⁵⁰ E. N. Tapia San Martín ,⁶⁰ C. Taranto,¹²⁴ A. Taruya ,²⁷⁷ J. D. Tasson ,¹⁶¹
 R. Tenorio ,¹⁴⁵ J. E. S. Terhune ,¹¹⁴ L. Terkowski ,¹²⁸ M. P. Thirugnanasambandam,¹¹ M. Thomas,⁵⁷
 P. Thomas,⁷³ E. E. Thompson,⁴⁸ J. E. Thompson ,¹⁷ S. R. Thondapu,⁹⁵ K. A. Thorne,⁵⁷ E. Thrane,⁵
 Shubhanshu Tiwari ,¹⁶² Srishti Tiwari,¹¹ V. Tiwari ,¹⁷ A. M. Toivonen,¹⁴⁹ A. E. Tolley ,⁵² T. Tomaru ,²⁰
 T. Tomura ,¹⁹⁰ M. Tonelli,^{79, 18} Z. Tornasi,²⁴ A. Torres-Forné ,¹²⁷ C. I. Torrie,¹ I. Tosta e Melo ,¹²³ D. Töyrä,⁸
 A. Trapananti ,^{41, 40} F. Travasso ,^{40, 41} G. Traylor,⁵⁷ M. Trevor,¹¹¹ M. C. Tringali ,⁴⁷ A. Tripathee ,¹⁸⁴
 L. Troiano,^{278, 104} A. Trovato ,⁴⁵ L. Trozzo ,^{4, 190} R. J. Trudeau,¹ D. Tsai,¹³⁰ K. W. Tsang,^{60, 279, 66}
 T. Tsang ,²⁸⁰ J.-S. Tsao,²³³ M. Tse,⁷⁵ R. Tso,¹³⁷ S. Tsuchida,¹⁷⁷ L. Tsukada,¹⁵⁰ D. Tsuna,²⁸ T. Tsutsui ,²⁸
 K. Turbang ,^{281, 205} M. Turconi,³⁷ D. Tuyenbayev ,¹⁷⁷ A. S. Ubhi ,¹⁴ N. Uchikata ,¹⁸⁸ T. Uchiyama ,¹⁹⁰
 R. P. Udall,¹ A. Ueda,²⁸² T. Uehara ,^{283, 284} K. Ueno ,²⁸ G. Ueshima,²⁸⁵ C. S. Unnikrishnan,²⁸⁶ A. L. Urban,⁷
 T. Ushiba ,¹⁹⁰ A. Utina ,^{155, 60} G. Vajente ,¹ A. Vajpeyi,⁵ G. Valdés ,¹⁸⁵ M. Valentini ,^{183, 99, 100}
 V. Valsan,⁶ N. van Bakel,⁶⁰ M. van Beuzekom ,⁶⁰ M. van Dael,^{60, 287} J. F. J. van den Brand ,^{155, 97, 60}
 C. Van Den Broeck,^{66, 60} D. C. Vander-Hyde,⁶⁸ H. van Haevermaet ,²⁰⁵ J. V. van Heijningen ,⁵⁹ M. H. P. M. van Putten,²⁸⁸ N. van Remortel ,²⁰⁵ M. Vardaro,^{212, 60} A. F. Vargas,¹²¹ V. Varma ,¹¹² M. Vasíth ,⁷⁶ A. Vecchio ,¹⁴
 G. Vedovato,⁸² J. Veitch ,²⁴ P. J. Veitch ,⁸⁹ J. Venneberg ,^{9, 10} G. Venugopalan ,¹ D. Verkindt ,³⁰ P. Verma,²²¹
 Y. Verma ,⁹⁵ S. M. Vermeulen ,¹⁷ D. Veske ,⁵¹ F. Vetrano,⁵⁵ A. Viceré ,^{55, 56} S. Vidyant,⁶⁸ A. D. Viets ,²⁸⁹
 A. Vijaykumar ,¹⁹ V. Villa-Ortega ,¹¹⁵ J.-Y. Vinet,³⁷ A. Virtuoso,^{239, 35} S. Vitale ,⁷⁵ H. Vocca,^{80, 40}
 E. R. G. von Reis,⁷³ J. S. A. von Wrangel,^{9, 10} C. Vorwick ,⁷³ S. P. Vyatchanin ,⁹⁸ L. E. Wade,²³⁷ M. Wade ,²³⁷
 K. J. Wagner ,¹²⁹ R. C. Walet,⁶⁰ M. Walker,⁶⁴ G. S. Wallace,³³ L. Wallace,¹ J. Wang ,¹⁷⁸ J. Z. Wang,¹⁸⁴
 W. H. Wang,⁹⁰ R. L. Ward,⁸ J. Warner,⁷³ M. Was ,³⁰ T. Washimi ,²⁰ N. Y. Washington,¹ J. Watchi ,¹⁴⁶
 B. Weaver,⁷³ C. R. Weaving,⁵² S. A. Webster,²⁴ M. Weinert,^{9, 10} A. J. Weinstein ,¹ R. Weiss,⁷⁵ C. M. Weller,²⁵⁸
 R. A. Weller ,²⁰⁴ F. Wellmann,^{9, 10} L. Wen,⁹⁴ P. Weßels,^{9, 10} K. Wette ,⁸ J. T. Whelan ,¹²⁹ D. D. White,⁴⁴
 B. F. Whiting ,⁷⁷ C. Whittle ,⁷⁵ D. Wilken,^{9, 10} D. Williams ,²⁴ M. J. Williams ,²⁴ A. R. Williamson ,⁵²
 J. L. Willis ,¹ B. Willke ,^{9, 10} D. J. Wilson,²⁵⁵ C. C. Wipf,¹ T. Włodarczyk,¹¹² G. Woan ,²⁴ J. Woehler,^{9, 10}
 J. K. Wofford ,¹²⁹ D. Wong,¹⁸¹ I. C. F. Wong ,¹³¹ M. Wright,²⁴ C. Wu ,¹³⁸ D. S. Wu ,^{9, 10} H. Wu,¹³⁸
 D. M. Wysocki,⁶ L. Xiao ,¹ T. Yamada,²⁶⁶ H. Yamamoto ,¹ K. Yamamoto ,¹⁸⁹ T. Yamamoto ,¹⁹⁰
 K. Yamashita,²⁰¹ R. Yamazaki,¹⁹⁸ F. W. Yang ,¹⁶⁰ K. Z. Yang ,¹⁴⁹ L. Yang ,¹⁶⁸ Y.-C. Yang,¹³⁰ Y. Yang ,²⁹⁰
 Yang Yang,⁷⁷ M. J. Yap,⁸ D. W. Yeeles,¹⁷ S.-W. Yeh,¹³⁸ A. B. Yelikar ,¹²⁹ M. Ying,¹³⁰ J. Yokoyama ,^{28, 27}
 T. Yokozawa,¹⁹⁰ J. Yoo,¹⁸⁰ T. Yoshioka,²⁰¹ Hang Yu ,¹³⁷ Haocun Yu ,⁷⁵ H. Yuzurihara,¹⁸⁸ A. Zadrożny,²²¹
 M. Zanolini,³⁶ S. Zeidler ,²⁹¹ T. Zelenova,⁴⁷ J.-P. Zendri,⁸² M. Zevin ,¹⁶⁶ M. Zhan,¹⁷⁸ H. Zhang,²³³ J. Zhang ,⁹⁴
 L. Zhang,¹ R. Zhang ,⁷⁷ T. Zhang,¹⁴ Y. Zhang,¹⁸⁵ C. Zhao ,⁹⁴ G. Zhao,¹⁴⁶ Y. Zhao ,^{188, 20} Yue Zhao,¹⁶⁰
 R. Zhou,¹⁹¹ Z. Zhou,¹⁵ X. J. Zhu ,⁵ Z.-H. Zhu ,^{120, 231} A. B. Zimmerman ,¹⁷⁰ M. E. Zucker,^{1, 75} and J. Zweizig ,¹
- (The LIGO Scientific Collaboration, the Virgo Collaboration, and the KAGRA Collaboration)

¹ LIGO Laboratory, California Institute of Technology, Pasadena, CA 91125, USA

² Graduate School of Science, Tokyo Institute of Technology, Meguro-ku, Tokyo 152-8551, Japan

³ Dipartimento di Farmacia, Università di Salerno, I-84084 Fisciano, Salerno, Italy

⁴ INFN, Sezione di Napoli, Complesso Universitario di Monte S. Angelo, I-80126 Napoli, Italy

⁵ OzGrav, School of Physics & Astronomy, Monash University, Clayton 3800, Victoria, Australia

⁶ University of Wisconsin-Milwaukee, Milwaukee, WI 53201, USA

⁷ Louisiana State University, Baton Rouge, LA 70803, USA

⁸ OzGrav, Australian National University, Canberra, Australian Capital Territory 0200, Australia

⁹ Max Planck Institute for Gravitational Physics (Albert Einstein Institute), D-30167 Hannover, Germany

¹⁰ Leibniz Universität Hannover, D-30167 Hannover, Germany

¹¹ Inter-University Centre for Astronomy and Astrophysics, Pune 411007, India

¹² University of Cambridge, Cambridge CB2 1TN, United Kingdom

¹³ Theoretisch-Physikalisches Institut, Friedrich-Schiller-Universität Jena, D-07743 Jena, Germany

¹⁴ University of Birmingham, Birmingham B15 2TT, United Kingdom

- ¹⁵Northwestern University, Evanston, IL 60208, USA
¹⁶Instituto Nacional de Pesquisas Espaciais, 12227-010 São José dos Campos, São Paulo, Brazil
¹⁷Cardiff University, Cardiff CF24 3AA, United Kingdom
¹⁸INFN, Sezione di Pisa, I-56127 Pisa, Italy
¹⁹International Centre for Theoretical Sciences, Tata Institute of Fundamental Research, Bengaluru 560089, India
²⁰Gravitational Wave Science Project, National Astronomical Observatory of Japan (NAOJ), Mitaka City, Tokyo 181-8588, Japan
²¹Advanced Technology Center, National Astronomical Observatory of Japan (NAOJ), Mitaka City, Tokyo 181-8588, Japan
²²Dipartimento di Fisica, Università degli Studi di Torino, I-10125 Torino, Italy
²³INFN Sezione di Torino, I-10125 Torino, Italy
²⁴SUPA, University of Glasgow, Glasgow G12 8QQ, United Kingdom
²⁵Università di Napoli "Federico II", Complesso Universitario di Monte S. Angelo, I-80126 Napoli, Italy
²⁶Université de Lyon, Université Claude Bernard Lyon 1, CNRS, Institut Lumière Matière, F-69622 Villeurbanne, France
²⁷Department of Physics, The University of Tokyo, Bunkyo-ku, Tokyo 113-0033, Japan
²⁸Research Center for the Early Universe (RESCEU), The University of Tokyo, Bunkyo-ku, Tokyo 113-0033, Japan
²⁹Institut de Ciències del Cosmos (ICCUB), Universitat de Barcelona, C/ Martí i Franquès 1, Barcelona, 08028, Spain
³⁰Univ. Savoie Mont Blanc, CNRS, Laboratoire d'Annecy de Physique des Particules - IN2P3, F-74000 Annecy, France
³¹Institut de Física d'Altes Energies (IFAE), Barcelona Institute of Science and Technology, and ICREA, E-08193 Barcelona, Spain
³²Gran Sasso Science Institute (GSSI), I-67100 L'Aquila, Italy
³³SUPA, University of Strathclyde, Glasgow G1 1XQ, United Kingdom
³⁴Dipartimento di Scienze Matematiche, Informatiche e Fisiche, Università di Udine, I-33100 Udine, Italy
³⁵INFN, Sezione di Trieste, I-34127 Trieste, Italy
³⁶Embry-Riddle Aeronautical University, Prescott, AZ 86301, USA
³⁷Artemis, Université Côte d'Azur, Observatoire de la Côte d'Azur, CNRS, F-06304 Nice, France
³⁸GRAPPA, Anton Pannekoek Institute for Astronomy and Institute for High-Energy Physics, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, Netherlands
³⁹National and Kapodistrian University of Athens, School of Science Building, 2nd floor, Panepistimiopolis, 15771 Ilissia, Greece
⁴⁰INFN, Sezione di Perugia, I-06123 Perugia, Italy
⁴¹Università di Camerino, Dipartimento di Fisica, I-62032 Camerino, Italy
⁴²American University, Washington, D.C. 20016, USA
⁴³Earthquake Research Institute, The University of Tokyo, Bunkyo-ku, Tokyo 113-0032, Japan
⁴⁴California State University Fullerton, Fullerton, CA 92831, USA
⁴⁵Université de Paris, CNRS, Astroparticule et Cosmologie, F-75006 Paris, France
⁴⁶Université Paris-Saclay, CNRS/IN2P3, IJCLab, 91405 Orsay, France
⁴⁷European Gravitational Observatory (EGO), I-56021 Cascina, Pisa, Italy
⁴⁸Georgia Institute of Technology, Atlanta, GA 30332, USA
⁴⁹Chennai Mathematical Institute, Chennai 603103, India
⁵⁰Department of Mathematics and Physics, Columbia University, New York, NY 10027, USA
⁵²University of Portsmouth, Portsmouth, PO1 3FX, United Kingdom
⁵³Kamioka Branch, National Astronomical Observatory of Japan (NAOJ), Kamioka-cho, Hida City, Gifu 506-1205, Japan
⁵⁴The Graduate University for Advanced Studies (SOKENDAI), Mitaka City, Tokyo 181-8588, Japan
⁵⁵Università degli Studi di Urbino "Carlo Bo", I-61029 Urbino, Italy
⁵⁶INFN, Sezione di Firenze, I-50019 Sesto Fiorentino, Firenze, Italy
⁵⁷LIGO Livingston Observatory, Livingston, LA 70754, USA
⁵⁸INFN, Sezione di Roma, I-00185 Roma, Italy
⁵⁹Université catholique de Louvain, B-1348 Louvain-la-Neuve, Belgium
⁶⁰Nikhef, Science Park 105, 1098 XG Amsterdam, Netherlands
⁶¹King's College London, University of London, London WC2R 2LS, United Kingdom
⁶²Korea Institute of Science and Technology Information, Daejeon 34141, Republic of Korea
⁶³National Institute for Mathematical Sciences, Daejeon 34047, Republic of Korea
⁶⁴Christopher Newport University, Newport News, VA 23606, USA
⁶⁵School of High Energy Accelerator Science, The Graduate University for Advanced Studies (SOKENDAI), Tsukuba City, Ibaraki 305-0801, Japan
⁶⁶Institute for Gravitational and Subatomic Physics (GRASP), Utrecht University, Princetonplein 1, 3584 CC Utrecht, Netherlands
⁶⁷University of Oregon, Eugene, OR 97403, USA
⁶⁸Syracuse University, Syracuse, NY 13244, USA
⁶⁹Université de Liège, B-4000 Liège, Belgium

- ⁷⁰Università degli Studi di Milano-Bicocca, I-20126 Milano, Italy
⁷¹INFN, Sezione di Milano-Bicocca, I-20126 Milano, Italy
⁷²INAF, Osservatorio Astronomico di Brera sede di Merate, I-23807 Merate, Lecco, Italy
⁷³LIGO Hanford Observatory, Richland, WA 99352, USA
⁷⁴Dipartimento di Medicina, Chirurgia e Odontoiatria "Scuola Medica Salernitana", Università di Salerno, I-84081 Baronissi, Salerno, Italy
⁷⁵LIGO Laboratory, Massachusetts Institute of Technology, Cambridge, MA 02139, USA
⁷⁶Wigner RCP, RMKI, H-1121 Budapest, Konkoly Thege Miklós út 29-33, Hungary
⁷⁷University of Florida, Gainesville, FL 32611, USA
⁷⁸Stanford University, Stanford, CA 94305, USA
⁷⁹Università di Pisa, I-56127 Pisa, Italy
⁸⁰Università di Perugia, I-06123 Perugia, Italy
⁸¹Università di Padova, Dipartimento di Fisica e Astronomia, I-35131 Padova, Italy
⁸²INFN, Sezione di Padova, I-35131 Padova, Italy
⁸³Bard College, Annandale-On-Hudson, NY 12504, USA
⁸⁴Montana State University, Bozeman, MT 59717, USA
⁸⁵Institute for Plasma Research, Bhat, Gandhinagar 382428, India
⁸⁶Universiteit Gent, B-9000 Gent, Belgium
⁸⁷Nicolaus Copernicus Astronomical Center, Polish Academy of Sciences, 00-716, Warsaw, Poland
⁸⁸Dipartimento di Ingegneria, Università del Sannio, I-82100 Benevento, Italy
⁸⁹OzGrav, University of Adelaide, Adelaide, South Australia 5005, Australia
⁹⁰The University of Texas Rio Grande Valley, Brownsville, TX 78520, USA
⁹¹California State University, Los Angeles, Los Angeles, CA 90032, USA
⁹²Departamento de Matemáticas, Universitat Autònoma de Barcelona, Edifici C Facultad de Ciencias 08193 Bellaterra (Barcelona), Spain
⁹³INFN, Sezione di Genova, I-16146 Genova, Italy
⁹⁴OzGrav, University of Western Australia, Crawley, Western Australia 6009, Australia
⁹⁵RRCAT, Indore, Madhya Pradesh 452013, India
⁹⁶Missouri University of Science and Technology, Rolla, MO 65409, USA
⁹⁷Vrije Universiteit Amsterdam, 1081 HV Amsterdam, Netherlands
⁹⁸Lomonosov Moscow State University, Moscow 119991, Russia
⁹⁹Università di Trento, Dipartimento di Fisica, I-38123 Povo, Trento, Italy
¹⁰⁰INFN, Trento Institute for Fundamental Physics and Applications, I-38123 Povo, Trento, Italy
¹⁰¹SUPA, University of the West of Scotland, Paisley PA1 2BE, United Kingdom
¹⁰²Bar-Ilan University, Ramat Gan, 5290002, Israel
¹⁰³Dipartimento di Fisica "E.R. Caianiello", Università di Salerno, I-84084 Fisciano, Salerno, Italy
¹⁰⁴INFN, Sezione di Napoli, Gruppo Collegato di Salerno, Complesso Universitario di Monte S. Angelo, I-80126 Napoli, Italy
¹⁰⁵Università di Roma "La Sapienza", I-00185 Roma, Italy
¹⁰⁶Univ Rennes, CNRS, Institut FOTON - UMR6082, F-3500 Rennes, France
¹⁰⁷Indian Institute of Technology Bombay, Powai, Mumbai 400 076, India
¹⁰⁸INFN, Laboratori Nazionali del Gran Sasso, I-67100 Assergi, Italy
¹⁰⁹Laboratoire Kastler Brossel, Sorbonne Université, CNRS, ENS-Université PSL, Collège de France, F-75005 Paris, France
¹¹⁰Astronomical Observatory Warsaw University, 00-478 Warsaw, Poland
¹¹¹University of Maryland, College Park, MD 20742, USA
¹¹²Max Planck Institute for Gravitational Physics (Albert Einstein Institute), D-14476 Potsdam, Germany
¹¹³L2IT, Laboratoire des 2 Infinis - Toulouse, Université de Toulouse, CNRS/IN2P3, UPS, F-31062 Toulouse Cedex 9, France
¹¹⁴Villanova University, Villanova, PA 19085, USA
¹¹⁵IGFAE, Universidade de Santiago de Compostela, 15782 Spain
¹¹⁶Stony Brook University, Stony Brook, NY 11794, USA
¹¹⁷Center for Computational Astrophysics, Flatiron Institute, New York, NY 10010, USA
¹¹⁸NASA Goddard Space Flight Center, Greenbelt, MD 20771, USA
¹¹⁹Dipartimento di Fisica, Università degli Studi di Genova, I-16146 Genova, Italy
¹²⁰Department of Astronomy, Beijing Normal University, Beijing 100875, China
¹²¹OzGrav, University of Melbourne, Parkville, Victoria 3010, Australia
¹²²Università degli Studi di Sassari, I-07100 Sassari, Italy
¹²³INFN, Laboratori Nazionali del Sud, I-95125 Catania, Italy
¹²⁴Università di Roma Tor Vergata, I-00133 Roma, Italy
¹²⁵INFN, Sezione di Roma Tor Vergata, I-00133 Roma, Italy
¹²⁶University of Sannio at Benevento, I-82100 Benevento, Italy and INFN, Sezione di Napoli, I-80100 Napoli, Italy
¹²⁷Departamento de Astronomía y Astrofísica, Universitat de València, E-46100 Burjassot, València, Spain

- ¹²⁸Universität Hamburg, D-22761 Hamburg, Germany
¹²⁹Rochester Institute of Technology, Rochester, NY 14623, USA
¹³⁰National Tsing Hua University, Hsinchu City, 30013 Taiwan, Republic of China
¹³¹The Chinese University of Hong Kong, Shatin, NT, Hong Kong
¹³²Department of Applied Physics, Fukuoka University, Jonan, Fukuoka City, Fukuoka 814-0180, Japan
¹³³OzGrav, Charles Sturt University, Wagga Wagga, New South Wales 2678, Australia
¹³⁴Department of Physics, Tamkang University, Danshui Dist., New Taipei City 25137, Taiwan
¹³⁵Department of Physics and Institute of Astronomy, National Tsing Hua University, Hsinchu 30013, Taiwan
¹³⁶Department of Physics, Center for High Energy and High Field Physics, National Central University, Zhongli District, Taoyuan City 32001, Taiwan
¹³⁷CaRT, California Institute of Technology, Pasadena, CA 91125, USA
¹³⁸Department of Physics, National Tsing Hua University, Hsinchu 30013, Taiwan
¹³⁹Dipartimento di Ingegneria Industriale (DIIN), Università di Salerno, I-84084 Fisciano, Salerno, Italy
¹⁴⁰Institute of Physics, Academia Sinica, Nankang, Taipei 11529, Taiwan
¹⁴¹Université Lyon, Université Claude Bernard Lyon 1, CNRS, IP2I Lyon / IN2P3, UMR 5822, F-69622 Villeurbanne, France
¹⁴²INAF, Osservatorio Astronomico di Padova, I-35122 Padova, Italy
¹⁴³OzGrav, Swinburne University of Technology, Hawthorn VIC 3122, Australia
¹⁴⁴Université libre de Bruxelles, Avenue Franklin Roosevelt 50 - 1050 Bruxelles, Belgium
¹⁴⁵IAC3–IEEC, Universitat de les Illes Balears, E-07122 Palma de Mallorca, Spain
¹⁴⁶Université Libre de Bruxelles, Brussels 1050, Belgium
¹⁴⁷Departamento de Matemáticas, Universitat de València, E-46100 Burjassot, València, Spain
¹⁴⁸Texas Tech University, Lubbock, TX 79409, USA
¹⁴⁹University of Minnesota, Minneapolis, MN 55455, USA
¹⁵⁰The Pennsylvania State University, University Park, PA 16802, USA
¹⁵¹University of Rhode Island, Kingston, RI 02881, USA
¹⁵²Bellevue College, Bellevue, WA 98007, USA
¹⁵³Scuola Normale Superiore, Piazza dei Cavalieri, 7 - 56126 Pisa, Italy
¹⁵⁴Eötvös University, Budapest 1117, Hungary
¹⁵⁵Maastricht University, P.O. Box 616, 6200 MD Maastricht, Netherlands
¹⁵⁶The University of Sheffield, Sheffield S10 2TN, United Kingdom
¹⁵⁷Université Lyon, Université Claude Bernard Lyon 1, CNRS, Laboratoire des Matériaux Avancés (LMA), IP2I Lyon / IN2P3, UMR 5822, F-69622 Villeurbanne, France
¹⁵⁸Dipartimento di Scienze Matematiche, Fisiche e Informatiche, Università di Parma, I-43124 Parma, Italy
¹⁵⁹INFN, Sezione di Milano Bicocca, Gruppo Collegato di Parma, I-43124 Parma, Italy
¹⁶⁰The University of Utah, Salt Lake City, UT 84112, USA
¹⁶¹Carleton College, Northfield, MN 55057, USA
¹⁶²University of Zurich, Winterthurerstrasse 190, 8057 Zurich, Switzerland
¹⁶³Perimeter Institute, Waterloo, ON N2L 2Y5, Canada
¹⁶⁴Université de Strasbourg, CNRS, IPHC UMR 7178, F-67000 Strasbourg, France
¹⁶⁵West Virginia University, Morgantown, WV 26506, USA
¹⁶⁶University of Chicago, Chicago, IL 60637, USA
¹⁶⁷Montclair State University, Montclair, NJ 07043, USA
¹⁶⁸Colorado State University, Fort Collins, CO 80523, USA
¹⁶⁹Institute for Nuclear Research, Bem t'er 18/c, H-4026 Debrecen, Hungary
¹⁷⁰University of Texas, Austin, TX 78712, USA
¹⁷¹CNR-SPIN, c/o Università di Salerno, I-84084 Fisciano, Salerno, Italy
¹⁷²Scuola di Ingegneria, Università della Basilicata, I-85100 Potenza, Italy
¹⁷³Observatori Astronòmic, Universitat de València, E-46980 Paterna, València, Spain
¹⁷⁴Centro de Física das Universidades do Minho e do Porto, Universidade do Minho, Campus de Gualtar, PT-4710 - 057 Braga, Portugal
¹⁷⁵Department of Astronomy, The University of Tokyo, Mitaka City, Tokyo 181-8588, Japan
¹⁷⁶Faculty of Engineering, Niigata University, Nishi-ku, Niigata City, Niigata 950-2181, Japan
¹⁷⁷Department of Physics, Graduate School of Science, Osaka City University, Sumiyoshi-ku, Osaka City, Osaka 558-8585, Japan
¹⁷⁸State Key Laboratory of Magnetic Resonance and Atomic and Molecular Physics, Innovation Academy for Precision Measurement Science and Technology (APM), Chinese Academy of Sciences, Xiao Hong Shan, Wuhan 430071, China
¹⁷⁹University of Szeged, Dóm tér 9, Szeged 6720, Hungary
¹⁸⁰Cornell University, Ithaca, NY 14850, USA
¹⁸¹University of British Columbia, Vancouver, BC V6T 1Z4, Canada

- ¹⁸²INAF, Osservatorio Astronomico di Capodimonte, I-80131 Napoli, Italy
¹⁸³The University of Mississippi, University, MS 38677, USA
¹⁸⁴University of Michigan, Ann Arbor, MI 48109, USA
¹⁸⁵Texas A&M University, College Station, TX 77843, USA
¹⁸⁶Ulsan National Institute of Science and Technology, Ulsan 44919, Republic of Korea
¹⁸⁷Shanghai Astronomical Observatory, Chinese Academy of Sciences, Shanghai 200030, China
¹⁸⁸Institute for Cosmic Ray Research (ICRR), KAGRA Observatory,
The University of Tokyo, Kashiwa City, Chiba 277-8582, Japan
¹⁸⁹Faculty of Science, University of Toyama, Toyama City, Toyama 930-8555, Japan
¹⁹⁰Institute for Cosmic Ray Research (ICRR), KAGRA Observatory,
The University of Tokyo, Kamioka-cho, Hida City, Gifu 506-1205, Japan
¹⁹¹University of California, Berkeley, CA 94720, USA
¹⁹²Maastricht University, 6200 MD, Maastricht, Netherlands
¹⁹³Lancaster University, Lancaster LA1 4YW, United Kingdom
¹⁹⁴College of Industrial Technology, Nihon University, Narashino City, Chiba 275-8575, Japan
¹⁹⁵Institute of Astronomy, National Tsing Hua University, Hsinchu 30013, Taiwan
¹⁹⁶Rutherford Appleton Laboratory, Didcot OX11 0DE, United Kingdom
¹⁹⁷Department of Astronomy & Space Science, Chungnam National University, Yuseong-gu, Daejeon 34134, Republic of Korea
¹⁹⁸Department of Physical Sciences, Aoyama Gakuin University, Sagamihara City, Kanagawa 252-5258, Japan
¹⁹⁹Kavli Institute for Astronomy and Astrophysics,
Peking University, Haidian District, Beijing 100871, China
²⁰⁰Aristotle University of Thessaloniki, University Campus, 54124 Thessaloniki, Greece
²⁰¹Graduate School of Science and Engineering, University of Toyama, Toyama City, Toyama 930-8555, Japan
²⁰²Nambu Yoichiro Institute of Theoretical and Experimental Physics (NITEP),
Osaka City University, Sumiyoshi-ku, Osaka City, Osaka 558-8585, Japan
²⁰³Direktorate of Construction, Services & Estate Management, Mumbai 400094, India
²⁰⁴Vanderbilt University, Nashville, TN 37235, USA
²⁰⁵Universiteit Antwerpen, Prinsstraat 13, 2000 Antwerpen, Belgium
²⁰⁶University of Bialystok, 15-424 Bialystok, Poland
²⁰⁷Ewha Womans University, Seoul 03760, Republic of Korea
²⁰⁸National Astronomical Observatories, Chinese Academic of Sciences, Chaoyang District, Beijing, China
²⁰⁹School of Astronomy and Space Science, University of Chinese Academy of Sciences, Chaoyang District, Beijing, China
²¹⁰University of Southampton, Southampton SO17 1BJ, United Kingdom
²¹¹Institute for Cosmic Ray Research (ICRR), The University of Tokyo, Kashiwa City, Chiba 277-8582, Japan
²¹²Institute for High-Energy Physics, University of Amsterdam,
Science Park 904, 1098 XH Amsterdam, Netherlands
²¹³Chung-Ang University, Seoul 06974, Republic of Korea
²¹⁴University of Washington Bothell, Bothell, WA 98011, USA
²¹⁵Institute of Applied Physics, Nizhny Novgorod, 603950, Russia
²¹⁶Inje University Gimhae, South Gyeongsang 50834, Republic of Korea
²¹⁷Department of Physics, Myongji University, Yongin 17058, Republic of Korea
²¹⁸Institute of Particle and Nuclear Studies (IPNS),
High Energy Accelerator Research Organization (KEK), Tsukuba City, Ibaraki 305-0801, Japan
²¹⁹School of Physics and Astronomy, Cardiff University, Cardiff, CF24 3AA, UK
²²⁰Institute of Mathematics, Polish Academy of Sciences, 00656 Warsaw, Poland
²²¹National Center for Nuclear Research, 05-400 Świerk-Otwock, Poland
²²²Instituto de Fisica Teorica, 28049 Madrid, Spain
²²³Department of Physics, Nagoya University, Chikusa-ku, Nagoya, Aichi 464-8602, Japan
²²⁴Université de Montréal/Polytechnique, Montreal, Quebec H3T 1J4, Canada
²²⁵Laboratoire Lagrange, Université Côte d'Azur,
Observatoire Côte d'Azur, CNRS, F-06304 Nice, France
²²⁶Seoul National University, Seoul 08826, Republic of Korea
²²⁷Sungkyunkwan University, Seoul 03063, Republic of Korea
²²⁸NAVIER, École des Ponts, Univ Gustave Eiffel, CNRS, Marne-la-Vallée, France
²²⁹Università di Firenze, Sesto Fiorentino I-50019, Italy
²³⁰Department of Physics, National Cheng Kung University, Tainan City 701, Taiwan
²³¹School of Physics and Technology, Wuhan University, Wuhan, Hubei, 430072, China
²³²National Center for High-performance computing, National Applied Research Laboratories,
Hsinchu Science Park, Hsinchu City 30076, Taiwan
²³³Department of Physics, National Taiwan Normal University, sec. 4, Taipei 116, Taiwan
²³⁴NASA Marshall Space Flight Center, Huntsville, AL 35811, USA
²³⁵INFN, Sezione di Roma Tre, I-00146 Roma, Italy
²³⁶ESPCI, CNRS, F-75005 Paris, France
²³⁷Kenyon College, Gambier, OH 43022, USA

- ²³⁸School of Physics Science and Engineering, Tongji University, Shanghai 200092, China
²³⁹Dipartimento di Fisica, Università di Trieste, I-34127 Trieste, Italy
- ²⁴⁰Institute for Photon Science and Technology, The University of Tokyo, Bunkyo-ku, Tokyo 113-8656, Japan
²⁴¹Indian Institute of Technology Madras, Chennai 600036, India
- ²⁴²Saha Institute of Nuclear Physics, Bidhannagar, West Bengal 700064, India
²⁴³Institute of Space and Astronautical Science (JAXA), Chuo-ku, Sagamihara City, Kanagawa 252-0222, Japan
- ²⁴⁴Institut des Hautes Etudes Scientifiques, F-91440 Bures-sur-Yvette, France
- ²⁴⁵Faculty of Law, Ryukoku University, Fushimi-ku, Kyoto City, Kyoto 612-8577, Japan
- ²⁴⁶Indian Institute of Science Education and Research, Kolkata, Mohanpur, West Bengal 741252, India
²⁴⁷Department of Physics, University of Notre Dame, Notre Dame, IN 46556, USA
- ²⁴⁸Graduate School of Science and Technology, Niigata University, Nishi-ku, Niigata City, Niigata 950-2181, Japan
- ²⁴⁹Consiglio Nazionale delle Ricerche - Istituto dei Sistemi Complessi, Piazzale Aldo Moro 5, I-00185 Roma, Italy
²⁵⁰Korea Astronomy and Space Science Institute (KASI), Yuseong-gu, Daejeon 34055, Republic of Korea
- ²⁵¹Hobart and William Smith Colleges, Geneva, NY 14456, USA
- ²⁵²International Institute of Physics, Universidade Federal do Rio Grande do Norte, Natal RN 59078-970, Brazil
²⁵³Museo Storico della Fisica e Centro Studi e Ricerche "Enrico Fermi", I-00184 Roma, Italy
- ²⁵⁴Dipartimento di Matematica e Fisica, Università degli Studi Roma Tre, I-00146 Roma, Italy
²⁵⁵University of Arizona, Tucson, AZ 85721, USA
- ²⁵⁶Università di Trento, Dipartimento di Matematica, I-38123 Povo, Trento, Italy
²⁵⁷University of California, Riverside, Riverside, CA 92521, USA
²⁵⁸University of Washington, Seattle, WA 98195, USA
- ²⁵⁹Indian Institute of Technology, Palaj, Gandhinagar, Gujarat 382355, India
- ²⁶⁰Department of Electronic Control Engineering, National Institute of Technology, Nagaoka College, Nagaoka City, Niigata 940-8532, Japan
- ²⁶¹Departamento de Matemática da Universidade de Aveiro and Centre for Research and Development in Mathematics and Applications, Campus de Santiago, 3810-183 Aveiro, Portugal
²⁶²Marquette University, Milwaukee, WI 53233, USA
- ²⁶³Faculty of Science, Toho University, Funabashi City, Chiba 274-8510, Japan
- ²⁶⁴Graduate School of Science and Technology, Gunma University, Maebashi, Gunma 371-8510, Japan
- ²⁶⁵Institute for Quantum Studies, Chapman University, Orange, CA 92866, USA
- ²⁶⁶Accelerator Laboratory, High Energy Accelerator Research Organization (KEK), Tsukuba City, Ibaraki 305-0801, Japan
²⁶⁷Faculty of Information Science and Technology, Osaka Institute of Technology, Hirakata City, Osaka 573-0196, Japan
- ²⁶⁸INAF, Osservatorio Astrofisico di Arcetri, Largo E. Fermi 5, I-50125 Firenze, Italy
- ²⁶⁹Indian Institute of Technology Hyderabad, Sangareddy, Khammam, Telangana 502285, India
- ²⁷⁰Indian Institute of Science Education and Research, Pune, Maharashtra 411008, India
²⁷¹Istituto di Astrofisica e Planetologia Spaziali di Roma, Via del Fosso del Cavaliere, 100, 00133 Roma RM, Italy
- ²⁷²Department of Space and Astronautical Science, The Graduate University for Advanced Studies (SOKENDAI), Sagamihara City, Kanagawa 252-5210, Japan
²⁷³Andrews University, Berrien Springs, MI 49104, USA
- ²⁷⁴Research Center for Space Science, Advanced Research Laboratories, Tokyo City University, Setagaya, Tokyo 158-0082, Japan
- ²⁷⁵Institute for Cosmic Ray Research (ICRR), Research Center for Cosmic Neutrinos (RCCN), The University of Tokyo, Kashiwa City, Chiba 277-8582, Japan
- ²⁷⁶Department of Physics, Kyoto University, Sakyou-ku, Kyoto City, Kyoto 606-8502, Japan
²⁷⁷Yukawa Institute for Theoretical Physics (YITP), Kyoto University, Sakyou-ku, Kyoto City, Kyoto 606-8502, Japan
- ²⁷⁸Dipartimento di Scienze Aziendali - Management and Innovation Systems (DISA-MIS), Università di Salerno, I-84084 Fisciano, Salerno, Italy
²⁷⁹Van Swinderen Institute for Particle Physics and Gravity, University of Groningen, Nijenborgh 4, 9747 AG Groningen, Netherlands
- ²⁸⁰Faculty of Science, Department of Physics, The Chinese University of Hong Kong, Shatin, N.T., Hong Kong
²⁸¹Vrije Universiteit Brussel, Pleinlaan 2, 1050 Brussel, Belgium
- ²⁸²Applied Research Laboratory, High Energy Accelerator Research Organization (KEK), Tsukuba City, Ibaraki 305-0801, Japan
²⁸³Department of Communications Engineering, National Defense Academy of Japan, Yokosuka City, Kanagawa 239-8686, Japan
- ²⁸⁴Department of Physics, University of Florida, Gainesville, FL 32611, USA
²⁸⁵Department of Information and Management Systems Engineering, Nagaoka University of Technology, Nagaoka City, Niigata 940-2188, Japan
- ²⁸⁶Tata Institute of Fundamental Research, Mumbai 400005, India
- ²⁸⁷Eindhoven University of Technology, Postbus 513, 5600 MB Eindhoven, Netherlands
- ²⁸⁸Department of Physics and Astronomy, Sejong University, Gwangjin-gu, Seoul 143-747, Republic of Korea

²⁸⁹*Concordia University Wisconsin, Mequon, WI 53097, USA*

²⁹⁰*Department of Electrophysics, National Yang Ming Chiao Tung University, Hsinchu, Taiwan*

²⁹¹*Department of Physics, Rikkyo University, Toshima-ku, Tokyo 171-8501, Japan*

(Dated: December 1, 2021)

* Deceased, August 2020.

† Deceased, April 2021.