

Bottom-up reconstitution of focal adhesion complexes

Stephanie Schumacher¹, Roberto Vazquez Nunez², Christian Biertümpfel² and Naoko Mizuno^{2,3} 

¹ Department of Structural Cell Biology, Max Planck Institute of Biochemistry, Martinsried, Germany

² Laboratory of Structural Cell Biology, National Heart, Lung, and Blood Institute, National Institutes of Health, Bethesda, MD, USA

³ National Institute of Arthritis and Musculoskeletal and Skin Diseases, National Institutes of Health, Bethesda, MD, USA

Keywords

actin; integrin; PIP2; talin; vinculin

Correspondence

N. Mizuno, Laboratory of Structural Cell Biology, National Heart, Lung, and Blood Institute, National Institutes of Health, 50 South Dr., Bethesda, MD, 20892, USA
 Tel: +1-301-443-3118
 E-mail: naoko.mizuno@nih.gov

(Received 25 January 2021, revised 13 April 2021, accepted 14 May 2021)

doi:10.1111/febs.16023

Focal adhesions (FA) are large macromolecular assemblies relevant for various cellular and pathological events such as migration, polarization, and metastatic cancer formation. At FA sites at the migrating periphery of a cell, hundreds of players gather and form a network to respond to extra cellular stimuli transmitted by the integrin receptor, the most upstream component within a cell, initiating the FA signaling pathway. Numerous cellular experiments have been performed to understand the FA architecture and functions; however, their intricate network formation hampers unraveling the precise molecular actions of individual players. Here, *in vitro* bottom-up reconstitution presents an advantageous approach for elucidating the FA machinery and the hierarchical crosstalk of involved cellular players.

Cellular adhesions are crucial for the development of multicellular organisms and tissue morphogenesis, as they enable cells to connect to each other and to their environment. Cells attach to their surroundings via large intracellular molecular assemblies called focal adhesions (FA), which are localized at the plasma membrane. They are responsible for two major cellular processes: First, they provide direct mechanical links between the extracellular matrix (ECM) and the cell through connections to the cytoskeleton. Second, they sense the environment and bidirectionally transmit signals across the plasma membrane [1–3]. Besides providing structural connections, FAs also play a role in transmitting signals that can affect cell survival or differentiation by altering gene expression (reviewed in [4]). Aberrant FA functions have detrimental effects and are linked to several pathologies. Therefore, it is critical to understand how cells manage to properly function during adhesion and how they dynamically connect to their surrounding tissue.

Abbreviations

ECM, extracellular matrix; FAs, Focal adhesions; NMR, Nuclear magnetic resonance; PIP2, phosphatidylinositol-4,5-bisphosphate; TM, transmembrane.

A central component of FAs is the transmembrane integrin receptor, recognizing extracellular cues [5]. The activation of integrin can be triggered either by the binding of an extracellular ligand like fibronectin (outside-in signaling) or by intracellular proteins talin and kindlin (inside-out signaling) [5]; therefore, it transmits bi-directional signals. In both signaling directions, activated integrin further facilitates the recruitment of FA players and the FAs mature into a vast complex containing ~200 different proteins that extensively connect to the actin cytoskeleton [2,6,7]. Using super-resolution microscopy techniques [8–10], organizational properties of adhesions, both in vertical (perpendicular to plasma membrane) and lateral (parallel to membrane) directions, have been studied. 3D imaging revealed that a ~40-nm-wide region lies between integrins and the actin cytoskeleton, which consists of three layers: the integrin signaling, the force transduction, and the actin regulatory layers [9] (Fig. 1). Each layer contains distinct sets of proteins, with an

exception of talin that connects between integrin and actin, thus, spanning over all three layers [11]. Laterally, FAs extend along the actin retrograde flow as maturation proceeds, and proteins such as talin and vinculin align in this direction [12–14].

The mechanism of interactions of the key proteins with integrin, actin fibers, and other FA proteins has been extensively studied, employing numerous techniques such as biophysical techniques, mass spectrometry, super-resolution, and structural biology methods like electron microscopy, X-ray crystallography, or NMR [9,15–18]. However, the complicated FA network prevents us from understanding the precise molecular functions of the FA components. This is because individual components are part of various submodules within FAs and their functions cannot be easily dissected within the synergistic network or with only truncated domains lacking full regulatory

functions. Therefore, building up submachineries of FAs from individual, fully regulatable components, gives a great advantage to elucidate the hierarchical interactions of FA players. This ‘bottom-up’ reconstitution allows to systematically connect different functional subcomplexes toward a comprehensive understanding of FAs.

In this review, we will focus on the efforts of such bottom-up reconstitutions, particularly from the FA initiation process to the attachment of actin bundles onto the plasma membrane (Fig. 1). We will discuss the recent findings on the molecular crosstalk toward FA network formation. The elucidation of the fundamental molecular mechanisms of these players will lead to a general understanding of how cells attach and react to their surroundings, which is the first step in comprehending how aberrations in these processes contribute to diseases such as cancer [2].




Fig. 1. Simplified schematic of the FA machinery focusing on the activation of integrin. Components of the ECM (gray) bind to integrin receptors (α subunit in light green and β subunit in light blue), which reach through the plasma membrane (dark gray) into the cytosol. Intracellular proteins kindlin (pink) and talin (blue) bind to the cytoplasmic tail of β -integrin together with additional signaling factors like FAK, Pax and Skelemin (gray). Activated talin extends through all FA layers from the integrin receptor to the actin cytoskeleton (red/pink) and vinculin (dark yellow) enforces the talin-actin interaction. The FA machinery is tightly regulated and allows bidirectional signal transduction from outside-in and from inside-out.

FA core initiation machinery

Integrins

The main conductor of FAs, integrins, are type I transmembrane receptors that link the extracellular environment to the cytoskeleton of a cell (Figs 1 and 2). Integrins are heterodimers, consisting of a noncovalently linked α and β subunit (Fig. 2A). In vertebrates, 18 α and 8 β isoforms assemble into 24 different integrins, recognizing various ligands and exhibiting diverse expression profiles [5]. Integrins can be divided into four subclasses, depending on their extracellular ligand recognition profile, namely by binding to RGD, collagen, laminin, or leucocyte-specific receptors [5]. Upon ligand binding, integrins undergo conformational changes that enable intracellular proteins to bind (Fig. 2B). Intracellular linker proteins such as talin and kindlin connect further to the actin cytoskeleton (Fig. 1) and transduce chemical and mechanical signals into the cell (outside-in signaling). On the other hand, their ligand binding affinity can also be regulated by binding of intracellular proteins (inside-out signaling). By changing their structural conformation from a bent-closed to an extended-open state [5], integrins can integrate and transduce stimuli across the plasma membrane, which affect downstream signaling and cell fate [4].

Integrin–direct communicator between the outside and inside of a cell

Integrin α and β subunits are comprised of three main parts: the N-terminal ligand-binding ectodomain (~ 800 a.a.), a transmembrane (TM) helix (~ 20 a.a.), and the C-terminal cytoplasmic tail (CT; ~ 13–79 a.a.) [19] (Fig. 2A). The ectodomains themselves are divided into head, upper leg, and lower leg domains [20]. The ligands are typically recognized by a cleft between the β -propeller of the α subunit head and the β I domain of the β subunit head. The well-studied RGD motif, a three amino acid sequence on many integrin ligands, such as fibronectin or fibrinogen, is recognized by both subunits [21–23]. Studies employing techniques such as X-ray crystallography, electron microscopy, NMR, and light microscopy have revealed that integrins exist in three main conformations: bent, extended-closed, and extended-open (Fig. 2A) [5,20,22,24–27]. The head is in close proximity to the closed legs in the bent state with an exception of the recently reported integrin α 5 β 1 showing an incomplete bent conformation [28]. In the bent conformation, integrin displays a low ligand binding affinity [20]. When extended, the

integrin ectodomain is pointed away from the membrane [20]; the legs however are still closed (extended-closed). Further opening of the integrin headpiece leads to the separation of the legs (extended-open), and this conformation shows high ligand binding affinity. It has been suggested that integrins are constantly shifting between these different conformational states, which can be described as a conformational equilibrium or molecular breathing [5,20]. The downstream TM domains of α - and β -integrin form a coiled-coil helix pair, securing the inactive form of integrin when closed [29–32]. Upon opening of integrin, the separation of the coiled-coil is thought to occur, accommodating a large conformational change of integrin [25,29,32]. The CT domains are relatively short, ranging from 13 to 79 amino acids in length, except for the β 4 tail containing ~ 1000 amino acids. Despite their small size, integrin CT domains are considered as ‘interaction hubs’ for proteins of the intracellular signaling network [30,33]. Particularly, the β CT domain is of importance, as it contains several recognition motifs for interaction partners, such as the NPxY and NxxY motifs (Fig. 2B) [19,33]. The membrane-proximal NPxY motif is recognized by phosphotyrosine binding (PTB)-containing proteins, such as talin and DOK1 [34,35] and, the membrane-distal motif NxxY is recognized by proteins such as integrin cytoplasmic domain-associated protein 1 (ICAP1), CD98, and kindlin [19,30,33,36–41].

Since multiple proteins can bind to partially overlapping regions on the β integrin CT domains and possibly compete with each other, the process of the protein interactions must be regulated [33]. Phosphorylation could be employed to switch between adaptor protein binding, as in the case of talin and DOK1 [42]. Similarly, the role of lipids in the recruitment of adaptor proteins has been suggested [43–46]. Several adaptor proteins contain domains that are known for lipid binding (e.g., PH domain in kindlin) [33]. It has also been observed that increased lipid phosphorylation can take place within FAs by phosphatidylinositol 3-kinase (PI3Ks) and phosphatidylinositol(4) phosphate 5 kinase type I gamma (PIPKI gamma) [47–49], which affects the affinity and recruitment of FA proteins like talin and kindlin to the plasma membrane.

Intracellular focal adhesion adaptor proteins

Out of the over 200 proteins involved in FA signaling, only a few of intracellular proteins are involved in the initial integrin activation and adhesion assembly steps. Talin and kindlin are two major FA components, which are exclusive integrin activators and key factors




Fig. 2. Domain architecture of integrin receptors. (A) α -integrins consist of a β -propeller head, thigh, Genu, Calf-1, calf-2, TM helix, and cytoplasmic tail domains. β -integrins comprise a β I head, hybrid, PSI, I-EGF1-4, β -tail, TM helix, and cytoplasmic tail domains. Integrin receptors can adopt a bent (left), an extended-closed (center), and an extended-open (right) conformation. They bind ligands typically at the cleft between the α - and β -subunit heads (depicted in orange). (B) The cytoplasmic tails of β -integrins can contain two different linear binding motifs (NPxY or NxxY) for various interaction partners (in boxes on the right) depending on β -integrin isotype.

to trigger FA assembly. Both talin and kindlin contain 4.1-ezrin-radixin-moesin (FERM) domains (Fig. 3) and bind to the CT domain of the β subunit of integrin (Fig. 2). Both proteins cooperate with each other and are needed to fully activate integrins [19,50].

Talin

Talin, as a main integrin activator, acts as a direct link between integrin and the actin cytoskeleton [50,51]. As it is present in most integrin-based adhesions and crucial for inside-out activation of integrin, talin has been denoted as the core [52] or master of FAs [53]. Talin is also important for mechanotransduction, as it senses mechanical force and transduces it into biochemical signals [53]. In mammals, two isoforms of talin are present; talin-1 is ubiquitously expressed and talin-2 is mostly found in heart, brain, and kidney [52]. The significance of talin for proper cell adherence and function has been demonstrated by a plethora of studies. Knockout of talin leads to embryonic lethality in mice at E 8.5 [54] or severe defects when targeted to specific tissues [50]. On a cellular level, depleted talin can lead to decreased integrin activation and impaired cell spreading [55–57].

Talin is a 270-kDa large protein consisting of an N-terminal head followed by a rod domain connected by a long unstructured linker region (Fig. 3A,B). The talin head is an atypical FERM domain having an additional F0 domain. The rod domain is comprised of 62 α -helices arranged into 13 helical bundles (R1–

R13), consisting of 4–5 helices per bundle followed by a dimerization domain (DD) at the C terminus [50,58].

Regulation of talin

As talin is important for many cellular processes, several control and recruitment mechanisms tightly regulate its activation. *In vitro* studies showed that talin can exist either in a \sim 15-nm globular, autoinhibited form or in a \sim 80-nm extended conformation [53,59,60]. A recent cryo-EM structure of autoinhibited talin displays how the long rod domains are folded in autoinhibited talin [60] (Fig. 3A), which occludes most major interaction sites to FA proteins such as actin, vinculin, and integrin. Two intramolecular interactions between the FERM domain and the rod domain, namely F3–R9 and F2–R12, were shown to be critical for autoinhibition. The F3–R9 connection shields the main integrin binding site 1 (IBS1) on F3 [45], while the F2–R12 connection shields the phosphatidylinositol-4,5-bisphosphate (PIP₂) binding pocket on F2 [60] (Fig. 3A). The FERM domain F0–F1 is connected to F2 by a flexible linker and appears to be accessible when talin employs an autoinhibited conformation. This notion is also supported by the two crystal structures showing the arrangement of F0–F1 either as a linear extension from F2–F3 [61] or folding into a canonical compact form [62], reflecting its conformational dynamics. The recruitment of talin to membrane is suggested to be facilitated by the GTPase Rap1 [63]. Rap1 binds to the tip of F0, which




Fig. 3. Domain architecture of talin, kindlin, and vinculin. (A) Talin consists of a globular FERM head (F0-3) and a tail of 13 helical rod domains (R1-R13, in different colors) and a dimerization helix (DD, behind R13). In the autoinhibited form, all rod domains are entangled and the structure is secured by key interactions between F2-R12 and F3-R13 (insert). (B) Model of activated talin in an extended conformation with individual domains in different colors. Actin-binding sites (ABS2 and ABS3) become accessible. Potential vinculin-binding sites are highlighted with violet balls. (C) Kindlin consists of a globular FERM domain (F0-F3) and an additional PH domain. (D) Vinculin consists of a head domain comprising helical bundles D1-D4 and a helical tail that folds back in the autoinhibited state. The tail domain is released upon activation and opens binding sites for talin as well as for actin.

is still accessible in the inhibited form of talin, which may suggest that talin may be recruited to the membrane surface in the autoinhibited form. Upon

activation of talin through the engagement to FAs, the FERM domain may align linearly on the membrane surface in a tight interaction when PIP_2 is present.

Interestingly, full-length talin was shown to act as a monomer in solution, while talin contains a short ~ 20 a.a. C-terminal dimerization domain [58,64] (Fig. 3B), which has been suggested to contribute to the cluster formation of FA components, necessary for FA maturation. Whether talin acts as a monomer or dimer when engaged in FAs is an important question to explore in future. For talin to act as a dimer, it may be necessary to have an interaction partner to align the extended talin for dimerization.

Kindlin

The other known integrin activator is kindlin. It contains ~ 75 kDa F0–F3 FERM domain resembling the talin head, except for an additional inserted pleckstrin homology (PH) domain with affinity for PIP₂ [19,65] (Fig. 3b). In mammals, three isoforms exist: kindlin 1, 2, and 3, which exhibit varying expression patterns. Kindlin 1 is mainly expressed in epithelial cells and kindlin 3 in hematopoietic cells, while kindlin 2 is expressed ubiquitously, except for hematopoietic cells [19,66]. Consistent with these expression profiles, knockout experiments in mice have shown that kindlin 1 deletion leads to severe defects in epithelial tissues, such as skin blistering [40]; kindlin 2 knockout leads to embryonic lethality [38] and deletion of kindlin 3 leads to severe bleeding disorders [41]. On a cellular level, kindlin depletion leads to impaired inside-out integrin activation, FA formation and cell spreading, even when talin is present [38,40,41,67]. Kindlins are reported to bind to the membrane-distal NxxY motif of integrin β CT domains [19] (Fig. 2B). A crystal structure of kindlin 2 in complex with integrin β 1 CT revealed a dimeric form of kindlin with an additional binding motif on the β tail, TTV, which is necessary for the recruitment of kindlin to FAs in cells [68]. Furthermore, a recent study showed a possibility that the autoinhibition of kindlin may be achieved through homotrimerization [69,70], which blocks kindlin's integrin binding site in the trimer structure [69]. While these studies implicated scenarios how kindlin is regulated, the exact role of kindlin, particularly whether oligomerization aids or inhibits integrin binding, is still an open question.

Regulatory proteins of talin and kindlin

Talin, when elongated, can expose binding sites to a number of proteins [53,60,71]. Among the interactions provided by talin, particular importance lies in the connection between talin and actin as well as between talin, vinculin, and actin, as these complete the structural scaffolding of FAs. Besides the direct binding of

talin to actin, vinculin plays a critical role for strengthening the connection by crosslinking talin and actin [72–76]. Vinculin is a 120 kDa globular protein containing 4 α -helical bundles (D1–D4) in the talin-binding head domain and another α -helical bundle in its actin-binding tail domain [77,78] (Fig. 3D). In the autoinhibited conformation, they fold onto each other occluding binding sites to actin and talin (Fig. 3D). 11 potential vinculin binding sites are predicted along the talin sequence [79,80] (Fig. 3B), and multiple vinculin binding on the talin surface may facilitate actin bundle formation in FAs. Truncated vinculin head, tail, and deregulated mutants have been used to circumvent the lack of understanding of the regulation of vinculin activation [81–83].

Other talin-binding factors include RIAM (Rap1-GTP-interacting adaptor molecule) [64,84], talin-activator Kank [85], filamin [86], integrin cytoplasmic domain-associated protein 1 (ICAP1) [37], DOK1 [34,35], PI3PK γ [48,49], and α -actinin [87]. Each of these components interact not only with talin but also often with each other, facilitating to form an intricate network. However, how all these components are interplaying still has to be elucidated.

Kindlin also plays a role in transducing signals by interacting with several signaling proteins. Particularly, kindlin interacts with integrin-linked kinase (ILK) [88], which then forms a complex with particularly interesting new cysteine-histidine-rich protein (PINCH) and parvin, forming the ILK-PINCH-parvin (IPP) complex. The IPP complex also plays a role in connecting integrin to the actin cytoskeleton (reviewed in [89]). Other kindlin interacting proteins include migfilin [90], paxillin [91], and actin-related proteins 2/3 (Arp2/3) [92]. Paxillin binds to FAK and triggers Rho and Src signaling pathways that ultimately affect cell fate [93,94].

Bottom-up analysis of the interplay within the FA initiation machinery

FAs undergo a complex network formation that contains multiple layers of regulation; therefore, it is challenging to elucidate the functions of individual factors at a molecular level. Many molecular studies have used truncated or deregulated proteins to focus on how distinct interactions of proteins of interest occur. However, using full-length proteins and building up a machinery by a bottom-up *in vitro* reconstitution approach gives valuable hints on their regulations. A particular interest lies in the initiation of the FA machinery. How exactly talin, kindlin, and other factors dock onto the integrin-embedded plasma membrane surface and how they build up a machinery

connecting to the actin cytoskeleton provides insight into the molecular basis of FAs.

Integrin-PIP₂ containing membrane plus integrin activator talin and kindlin

As most of the FA proteins are tightly autoregulated, finding key to activate them is the critical first step for building up functional FA submachineries. Recent studies revealed that signaling lipid PIP₂ plays an important role in the regulation and recruitment of the integrin activators talin and kindlin [61,95–98]. Our recent study additionally showed that autoinhibited full-length talin binds to PIP₂-rich membrane surface, indicating that its autoinhibition is released upon binding to PIP₂-rich membrane [99].

By combining those with structural studies of the full-length autoinhibited talin 1 [60] and the truncated talin 2 FERM domain in complex with the integrin β 1D tail peptide [97], we can extrapolate our understanding on how activated talin then activates integrin on the membrane surface (Fig. 4A). When talin's FERM domain is docked to the PIP₂-containing membrane surface, the tangled rod domain is likely detached from the FERM domain, which would expose the integrin binding surface on the F3 domain. Subsequently, the rod domain could flip away from the membrane so that it orients toward the cytosol. Membrane-attached talin may readily recognize integrin because of its proximity to the plasma membrane as well as its exposure of the integrin binding site on the F3 domain. The exposed talin F3 domain binds to the NPxY motif in the integrin β CT. This interaction inhibits the salt bridge interaction between the integrin CT domains of the α and β subunits [34,56,97,100–103] inducing the separation of leg domains and the activation of integrin.

In contrast, it is still unclear how the regulation of kindlin and the involvement of PIP₂ takes place. Kindlin might be necessary to cluster integrins [104] rather than opening the integrin α - and β -CT domains like talin [105]. Future studies describing the dynamic process of kindlin and its interaction with talin, the PIP₂ membrane as well as integrin are awaited to understand what the precise regulatory role of kindlin is. Nevertheless, neither talin nor kindlin alone are thought to be sufficient to activate integrins by themselves *in vivo*, and are thus considered as co-dependent activators [19]. The binding mode remains elusive [19,50] as it is not clear whether talin and kindlin interact directly on the same β -tail (Fig. 4B). Other possibilities include that they bind sequentially to the same tail or simultaneously to different tails within the

same integrin cluster (Fig. 4B) [19,50]. It should however be noted that kindlin has been shown to be dispensable for integrin activation in an *in vitro* environment [18], and therefore, it may have rather an assisting role. Nevertheless, these are important questions that can be analyzed by *in vitro* reconstitution approaches.

Recruitment of actin to the membrane surface

The docking of actin bundles to the FA initiation machinery located at the plasma membrane surface is a critical step for the development of FAs. That is mainly mediated by talin as it can directly bind to integrin, the plasma membrane through PIP₂ and actin. Furthermore, vinculin strengthens the binding of talin and actin. These assemblies are the structural foundation of FAs. How actin can be recruited to the FA initiation site has been suggested through recent *in vitro* reconstitution studies [60,99]. Those studies showed that autoinhibited talin is still accessible for actin through actin-binding site 3 (ABS3) located at R13, while the other actin-binding site ABS2 at R4–R8 is occluded and its actin-binding function inhibited (Fig. 3A). ABS2 is ready to engage actin once the autoinhibition of talin is released by binding to PIP₂-enriched membranes (Figs. 3B and 4A). The opening of both ABS2 and ABS3 together facilitates binding of more actin filaments to talin. Upon binding of talin to integrin and actin, talin can extend to a 60- to 100-nm-long fibrous strand [11,59,106] (Fig. 4a). This elongation is thought to enable talin to span the distance between integrins and actin fibers [9] and to act as a cytoskeletal linker and mechanosensor [53]. At the same time, extended talin provides a platform for the binding of vinculin (Fig. 3B), reinforcing the engagement of actin to FAs. Interestingly, neither activated talin alone nor PIP₂-enriched membranes were shown to be sufficient to allow vinculin to bind to these components (Fig. 3D). It was however shown that activated talin attached to PIP₂-enriched membranes is able to activate vinculin and recruit it to the membrane surface where talin is localized [99] (Fig. 4A). While this is advantageous for the hierarchical regulation toward the engagement of actin bundles to the FA site, the precise molecular mechanism of action of vinculin regulation is still elusive.

Mechanosensitive behavior of the FA components

As FAs provide a link between the ECM and cells, the molecules within FAs are constantly exposed to




Fig. 4. Schematic of integrin activation and FA initiation. (A) Autoinhibited talin can approach to the PIP2-enriched membranes, resulting in the release of talin head and rod domains autoinhibition. Binding of talin to the cytoplasmic tails of β -integrin primes the integrin receptor, which then binds to ligands in the extended-open conformation. Opened talin can bind to actin, and this interaction is strengthened by crosslinking of talin and actin by activated vinculin. Further force-dependent extension of talin (indicated by lightning bolts) uncovers additional vinculin binding sites. (B) Models of the cooperative activation of integrin by kindlin and talin. Integrin receptors can be activated by sequential binding of kindlin and talin (violet path), by simultaneous binding of both proteins to the same integrin receptor (cis cooperation, green path), or by synchronous binding of both proteins to different, clustered integrin receptors (trans cooperation, blue path).

mechanical forces generated by both constant actin polymerization toward the membrane as well as actomyosin contraction of the cytoskeleton. Several of the components, such as integrin, talin, and vinculin, are found to be mechanosensitive [107–110]. To characterize the force induced behavior of FA molecules, *in vitro* biophysical analyses have been instrumental. Increase of the strength of the integrin-ECM bond with applied force was observed, both on isolated integrin as well as on cells [111,112], which marks it as a so-called catch-bond [15]. The force that integrin-ECM bonds can withstand until the bond breaks has been measured to be 50–100 pN [113]. In contrast, the conformation of talin is altered upon force, as it unfolds, which marks it as a slip-bond [107]. The unfolding of talin leads to an exposure of cryptic binding sites for proteins like vinculin [79]. This suggested to expose some of the 11 cryptic vinculin binding sites, while only one vinculin binds to talin without applied force [60]. The recruitment of more vinculin to the stretched talin surface is suggested to promote FA growth [79,80], but the precise mode of action of talin under force is still not well understood. Furthermore, the stretching of actin has been suggested to change the affinity of actin binding proteins such as vinculin [114]. Interestingly, the affinity of vinculin to actin depends on the direction of the applied force, implicating another layer of regulation to control vinculin binding at the FA site. As talin's binding modules resemble the structural folding of vinculin, it is possible that talin may also be able to sense the stretching of actin, as for the case of catenin [115–117]. These examples hint at the regulation of several protein functions and the assembly of the FA initiation machinery by force. Taking the mechanosensitivity into consideration will therefore give more comprehensive insights into FA assembly.

Maturation of FA and conclusion

The assembly process of FA complexes progresses in a highly coordinated and dynamic manner [51,118]. Upon generation of a positive feedback loop, which is directly linked to mechanical force, that is, ECM stiffness and

myosin II contractility [17,119], the maturation of FAs is thought to occur where hundreds of FA components start forming an interconnected network. Building up the FA machinery from minimal components gives a great advantage for elucidating key interactions within FAs and provides a hierarchical understanding of the molecular actions of individual players. That provides a basis for the understanding of the layers of the vast FA network formation.

Acknowledgements

We thank members of the Mizuno laboratory for helpful discussions. NM acknowledges the Boehringer Ingelheim Foundation Plus 3 Program, and the European Research Council (ERC-CoG, 724209) and German Research Foundation, DFG SFB863 for facilitating the related research. NM is a recipient of the EMBO Young Investigator award. NM is supported by the Intramural Research Program of the National Heart Lung and Blood Institute, and the National Institute of Arthritis and Musculoskeletal and Skin Diseases of National Institutes of Health, USA.

Conflict of interest

The authors declare no conflicts of interest.

Author contributions

SS, CB., and NM collected the relevant literature and wrote the review manuscript. RVN prepared the figures. All the authors read and approved the submitted version.

References

- 1 Geiger B, Bershadsky A, Pankov R & Yamada KM (2001) Transmembrane crosstalk between the extracellular matrix–cytoskeleton crosstalk. *Nat Rev Mol Cell Biol* **2**, 793–805.
- 2 Winograd-Katz SE, Fassler R, Geiger B & Legate KR (2014) The integrin adhesome: from genes and

- proteins to human disease. *Nat Rev Mol Cell Biol* **15**, 273–288.
- 3 Wozniak MA, Modzelewska K, Kwong L & Keely PJ (2004) Focal adhesion regulation of cell behavior. *Biochim Biophys Acta* **1692**, 103–119.
 - 4 Legate KR, Wickstrom SA & Fassler R (2009) Genetic and cell biological analysis of integrin outside-in signaling. *Genes Dev* **23**, 397–418.
 - 5 Hynes RO (2002) Integrins: bidirectional, allosteric signaling machines. *Cell* **110**, 673–687.
 - 6 Zaidel-Bar R, Itzkovitz S, Ma'ayan A, Iyengar R & Geiger B (2007) Functional atlas of the integrin adhesome. *Nat Cell Biol* **9**, 858–867.
 - 7 Zaidel-Bar R & Geiger B (2010) The switchable integrin adhesome. *J Cell Sci* **123**, 1385–1388.
 - 8 Case LB, Baird MA, Shtengel G, Campbell SL, Hess HF, Davidson MW & Waterman CM (2015) Molecular mechanism of vinculin activation and nanoscale spatial organization in focal adhesions. *Nat Cell Biol* **17**, 880–892.
 - 9 Kanchanawong P, Shtengel G, Pasapera AM, Ramko EB, Davidson MW, Hess HF & Waterman CM (2010) Nanoscale architecture of integrin-based cell adhesions. *Nature* **468**, 580–584.
 - 10 Paszek MJ, DuFort CC, Rubashkin MG, Davidson MW, Thorn KS, Liphardt JT & Weaver VM (2012) Scanning angle interference microscopy reveals cell dynamics at the nanoscale. *Nat Methods* **9**, 825–827.
 - 11 Liu J, Wang Y, Goh WI, Goh H, Baird MA, Ruchland S, Teo S, Bate N, Critchley DR, Davidson MW *et al.* (2015) Talin determines the nanoscale architecture of focal adhesions. *Proc Natl Acad Sci USA* **112**, E4864–E4873.
 - 12 Choi CK, Vicente-Manzanares M, Zareno J, Whitmore LA, Mogilner A & Horwitz AR (2008) Actin and alpha-actinin orchestrate the assembly and maturation of nascent adhesions in a myosin II motor-independent manner. *Nat Cell Biol* **10**, 1039–1050.
 - 13 Margadant F, Chew LL, Hu X, Yu H, Bate N, Zhang X & Sheetz M (2011) Mechanotransduction in vivo by repeated talin stretch-relaxation events depends upon vinculin. *PLoS Biol* **9**, e1001223.
 - 14 Zaidel-Bar R, Ballestrem C, Kam Z & Geiger B (2003) Early molecular events in the assembly of matrix adhesions at the leading edge of migrating cells. *J Cell Sci* **116**, 4605–4613.
 - 15 Kong F, Garcia AJ, Mould AP, Humphries MJ & Zhu C (2009) Demonstration of catch bonds between an integrin and its ligand. *J Cell Biol* **185**, 1275–1284.
 - 16 Patla I, Volberg T, Elad N, Hirschfeld-Warneken V, Grashoff C, Fassler R, Spatz JP, Geiger B & Medalia O (2010) Dissecting the molecular architecture of integrin adhesion sites by cryo-electron tomography. *Nat Cell Biol* **12**, 909–915.
 - 17 Schiller HB & Fassler R (2013) Mechanosensitivity and compositional dynamics of cell-matrix adhesions. *EMBO Rep* **14**, 509–519.
 - 18 Ye F, Hu G, Taylor D, Ratnikov B, Bobkov AA, McLean MA, Sliagar SG, Taylor KA & Ginsberg MH (2010) Recreation of the terminal events in physiological integrin activation. *J Cell Biol* **188**, 157–173.
 - 19 Moser M, Legate KR, Zent R & Fassler R (2009) The tail of integrins, talin, and kindlins. *Science* **324**, 895–899.
 - 20 Luo BH, Carman CV & Springer TA (2007) Structural basis of integrin regulation and signaling. *Annu Rev Immunol* **25**, 619–647.
 - 21 Nagae M, Re S, Mihara E, Nogi T, Sugita Y & Takagi J (2012) Crystal structure of alpha5beta1 integrin ectodomain: atomic details of the fibronectin receptor. *J Cell Biol* **197**, 131–140.
 - 22 Xiao T, Takagi J, Collier BS, Wang JH & Springer TA (2004) Structural basis for allostery in integrins and binding to fibrinogen-mimetic therapeutics. *Nature* **432**, 59–67.
 - 23 Xiong JP, Stehle T, Zhang R, Joachimiak A, Frech M, Goodman SL & Arnaout MA (2002) Crystal structure of the extracellular segment of integrin alpha Vbeta3 in complex with an Arg-Gly-Asp ligand. *Science* **296**, 151–155.
 - 24 Campbell ID & Humphries MJ (2011) Integrin structure, activation, and interactions. *Cold Spring Harb Perspect Biol* **3**, a004994.
 - 25 Kim M, Carman CV & Springer TA (2003) Bidirectional transmembrane signaling by cytoplasmic domain separation in integrins. *Science* **301**, 1720–1725.
 - 26 Takagi J, Petre BM, Walz T & Springer TA (2002) Global conformational rearrangements in integrin extracellular domains in outside-in and inside-out signaling. *Cell* **110**, 599–611.
 - 27 Takagi J, Strokovich K, Springer TA & Walz T (2003) Structure of integrin alpha5beta1 in complex with fibronectin. *EMBO J* **22**, 4607–4615.
 - 28 Schumacher S, Dedden D, Vazquez Nunez R, Matoba K, Takagi J, Biertümpfel C & Mizuno N (2021) Structural insights into integrin $\alpha 5 \beta 1$ opening by fibronectin ligand. *Science. Advances* **7**, eabe9716.
 - 29 Lau TL, Kim C, Ginsberg MH & Ulmer TS (2009) The structure of the integrin alphaIIb beta3 transmembrane complex explains integrin transmembrane signalling. *EMBO J* **28**, 1351–1361.
 - 30 Wegener KL & Campbell ID (2008) Transmembrane and cytoplasmic domains in integrin activation and protein-protein interactions (review). *Mol Membr Biol* **25**, 376–387.
 - 31 Partridge AW, Liu S, Kim S, Bowie JU & Ginsberg MH (2005) Transmembrane domain helix packing

- stabilizes integrin alphaIIb beta3 in the low affinity state. *J Biol Chem* **280**, 7294–7300.
- 32 Luo BH, Springer TA & Takagi J (2004) A specific interface between integrin transmembrane helices and affinity for ligand. *PLoS Biol* **2**, e153.
 - 33 Legate KR & Fassler R (2009) Mechanisms that regulate adaptor binding to beta-integrin cytoplasmic tails. *J Cell Sci* **122**, 187–198.
 - 34 Calderwood DA, Yan B, de Pereda JM, Alvarez BG, Fujioka Y, Liddington RC & Ginsberg MH (2002) The phosphotyrosine binding-like domain of talin activates integrins. *J Biol Chem* **277**, 21749–21758.
 - 35 Yamanashi Y & Baltimore D (1997) Identification of the Abl- and rasGAP-associated 62 kDa protein as a docking protein, Dok. *Cell* **88**, 205–211.
 - 36 Zent R, Fenczik CA, Calderwood DA, Liu S, Dello M & Ginsberg MH (2000) Class- and splice variant-specific association of CD98 with integrin beta cytoplasmic domains. *J Biol Chem* **275**, 5059–5064.
 - 37 Chang DD, Hoang BQ, Liu J & Springer TA (2002) Molecular basis for interaction between Icap1 alpha PTB domain and beta 1 integrin. *J Biol Chem* **277**, 8140–8145.
 - 38 Montanez E, Ussar S, Schifferer M, Bosl M, Zent R, Moser M & Fassler R (2008) Kindlin-2 controls bidirectional signaling of integrins. *Genes Dev* **22**, 1325–1330.
 - 39 Moser M, Bauer M, Schmid S, Ruppert R, Schmidt S, Sixt M, Wang HV, Sperandio M & Fassler R (2009) Kindlin-3 is required for beta2 integrin-mediated leukocyte adhesion to endothelial cells. *Nat Med* **15**, 300–305.
 - 40 Ussar S, Moser M, Widmaier M, Rognoni E, Harrer C, Genzel-Boroviczeny O & Fassler R (2008) Loss of Kindlin-1 causes skin atrophy and lethal neonatal intestinal epithelial dysfunction. *PLoS Genet* **4**, e1000289.
 - 41 Moser M, Nieswandt B, Ussar S, Pozgajova M & Fassler R (2008) Kindlin-3 is essential for integrin activation and platelet aggregation. *Nat Med* **14**, 325–330.
 - 42 Oxley CL, Anthis NJ, Lowe ED, Vakonakis I, Campbell ID & Wegener KL (2008) An integrin phosphorylation switch: the effect of beta3 integrin tail phosphorylation on Dok1 and talin binding. *J Biol Chem* **283**, 5420–5426.
 - 43 Chinthalapudi K, Rangarajan ES, Patil DN, George EM, Brown DT & Izard T (2014) Lipid binding promotes oligomerization and focal adhesion activity of vinculin. *J Cell Biol* **207**, 643–656.
 - 44 Chinthalapudi K, Rangarajan ES & Izard T (2018) The interaction of talin with the cell membrane is essential for integrin activation and focal adhesion formation. *Proc Natl Acad Sci USA* **115**, 10339–10344.
 - 45 Song X, Yang J, Hirbawi J, Ye S, Perera HD, Goksoy E, Dwivedi P, Plow EF, Zhang R & Qin J (2012) A novel membrane-dependent on/off switch mechanism of talin FERM domain at sites of cell adhesion. *Cell Res* **22**, 1533–1545.
 - 46 Janmey PA, Bucki R & Radhakrishnan R (2018) Regulation of actin assembly by PI(4,5)P2 and other inositol phospholipids: an update on possible mechanisms. *Biochem Biophys Res Commun* **506**, 307–314.
 - 47 Chen HC & Guan JL (1994) Association of focal adhesion kinase with its potential substrate phosphatidylinositol 3-kinase. *Proc Natl Acad Sci USA* **91**, 10148–10152.
 - 48 Di Paolo G, Pellegrini L, Letinic K, Cestra G, Zoncu R, Voronov S, Chang S, Guo J, Wenk MR & De Camilli P (2002) Recruitment and regulation of phosphatidylinositol phosphate kinase type 1 gamma by the FERM domain of talin. *Nature* **420**, 85–89.
 - 49 Ling K, Doughman RL, Firestone AJ, Bunce MW & Anderson RA (2002) Type I gamma phosphatidylinositol phosphate kinase targets and regulates focal adhesions. *Nature* **420**, 89–93.
 - 50 Calderwood DA, Campbell ID & Critchley DR (2013) Talins and kindlins: partners in integrin-mediated adhesion. *Nat Rev Mol Cell Biol* **14**, 503–517.
 - 51 Bachir AI, Zareno J, Moissoglu K, Plow EF, Gratton E & Horwitz AR (2014) Integrin-associated complexes form hierarchically with variable stoichiometry in nascent adhesions. *Curr Biol* **24**, 1845–1853.
 - 52 Gough RE & Goult BT (2018) The tale of two talins - two isoforms to fine-tune integrin signalling. *FEBS Lett* **592**, 2108–2125.
 - 53 Klapholz B & Brown NH (2017) Talin - the master of integrin adhesions. *J Cell Sci* **130**, 2435–2446.
 - 54 Monkley SJ, Zhou XH, Kinston SJ, Giblett SM, Hemmings L, Priddle H, Brown JE, Pritchard CA, Critchley DR & Fassler R (2000) Disruption of the talin gene arrests mouse development at the gastrulation stage. *Dev Dyn* **219**, 560–574.
 - 55 Nieswandt B, Moser M, Pleines I, Varga-Szabo D, Monkley S, Critchley D & Fassler R (2007) Loss of talin1 in platelets abrogates integrin activation, platelet aggregation, and thrombus formation in vitro and in vivo. *J Exp Med* **204**, 3113–3118.
 - 56 Tadokoro S, Shattil SJ, Eto K, Tai V, Liddington RC, de Pereda JM, Ginsberg MH & Calderwood DA (2003) Talin binding to integrin beta tails: a final common step in integrin activation. *Science* **302**, 103–106.
 - 57 Zhang X, Jiang G, Cai Y, Monkley SJ, Critchley DR & Sheetz MP (2008) Talin depletion reveals independence of initial cell spreading from integrin activation and traction. *Nat Cell Biol* **10**, 1062–1068.
 - 58 Gingras AR, Bate N, Goult BT, Hazelwood L, Canestrelli I, Grossmann JG, Liu H, Putz NS, Roberts GC, Volkmann N *et al.* (2008) The structure of the C-

- terminal actin-binding domain of talin. *EMBO J* **27**, 458–469.
- 59 Winkler J, Lunsdorf H & Jockusch BM (1997) Energy-filtered electron microscopy reveals that talin is a highly flexible protein composed of a series of globular domains. *Eur J Biochem* **243**, 430–436.
- 60 Dedden D, Schumacher S, Kelley CF, Zacharias M, Biertumpfel C, Fassler R & Mizuno N (2019) The architecture of talin1 reveals an autoinhibition mechanism. *Cell* **179**, 120–131.e13.
- 61 Elliott PR, Goult BT, Kopp PM, Bate N, Grossmann JG, Roberts GC, Critchley DR & Barsukov IL (2010) The Structure of the talin head reveals a novel extended conformation of the FERM domain. *Structure* **18**, 1289–1299.
- 62 Zhang P, Azizi L, Kukkurainen S, Gao T, Baikoghli M, Jacquier M-C, Sun Y, Määttä JAE, Cheng RH, Wehrle-Haller B *et al.* (2020) Crystal structure of the FERM-folded talin head reveals the determinants for integrin binding. *Proc Natl Acad Sci USA* **117**, 32402–32412.
- 63 Zhu L, Yang J, Bromberger T, Holly A, Lu F, Liu H, Sun K, Klapproth S, Hirbawi J, Byzova TV *et al.* (2017) Structure of Rap1b bound to talin reveals a pathway for triggering integrin activation. *Nat Commun* **8**, 1744–1812.
- 64 Goult BT, Xu XP, Gingras AR, Swift M, Patel B, Bate N, Kopp PM, Barsukov IL, Critchley DR, Volkmann N *et al.* (2013) Structural studies on full-length talin1 reveal a compact auto-inhibited dimer: implications for talin activation. *J Struct Biol* **184**, 21–32.
- 65 Qu H, Tu Y, Shi X, Larjava H, Saleem MA, Shattil SJ, Fukuda K, Qin J, Kretzler M & Wu C (2011) Kindlin-2 regulates podocyte adhesion and fibronectin matrix deposition through interactions with phosphoinositides and integrins. *J Cell Sci* **124**, 879–891.
- 66 Rognoni E, Ruppert R & Fassler R (2016) The kindlin family: functions, signaling properties and implications for human disease. *J Cell Sci* **129**, 17–27.
- 67 Ma YQ, Qin J, Wu C & Plow EF (2008) Kindlin-2 (Mig-2): a co-activator of beta3 integrins. *J Cell Biol* **181**, 439–446.
- 68 Li H, Deng Y, Sun K, Yang H, Liu J, Wang M, Zhang Z, Lin J, Wu C, Wei Z *et al.* (2017) Structural basis of kindlin-mediated integrin recognition and activation. *Proc Natl Acad Sci USA* **114**, 9349–9354.
- 69 Bu W, Levitskaya Z, Loh ZY, Jin S, Basu S, Ero R, Yan X, Wang M, Ngan SFC, Sze SK *et al.* (2020) Structural basis of human full-length kindlin-3 homotrimer in an auto-inhibited state. *PLoS Biol* **18**, e3000755.
- 70 Kadry YA, Maisuria EM, Huet-Calderwood C & Calderwood DA (2020) Differences in self-association between kindlin-2 and kindlin-3 are associated with differential integrin binding. *J Biol Chem* **295**, 11161–11173.
- 71 Goult BT, Yan J & Schwartz MA (2018) Talin as a mechanosensitive signaling hub. *J Cell Biol* **217**, 3776–3784.
- 72 Thievensen I, Thompson PM, Berlemont S, Plevock KM, Plotnikov SV, Zemljic-Harpe A, Ross RS, Davidson MW, Danuser G, Campbell SL *et al.* (2013) Vinculin-actin interaction couples actin retrograde flow to focal adhesions, but is dispensable for focal adhesion growth. *J Cell Biol* **202**, 163–177.
- 73 Thompson PM, Tolbert CE, Shen K, Kota P, Palmer SM, Plevock KM, Orlova A, Galkin VE, Burridge K, Egelman EH *et al.* (2014) Identification of an actin binding surface on vinculin that mediates mechanical cell and focal adhesion properties. *Structure* **22**, 697–706.
- 74 Bays JL & DeMali KA (2017) Vinculin in cell-cell and cell-matrix adhesions. *Cell Mol Life Sci* **74**, 2999–3009.
- 75 Johnson RP & Craig SW (1994) An intramolecular association between the head and tail domains of vinculin modulates talin binding. *J Biol Chem* **269**, 12611–12619.
- 76 Carisey A, Tsang R, Greiner AM, Nijenhuis N, Heath N, Nazgiewicz A, Kemkemer R, Derby B, Spatz J & Ballestrem C (2013) Vinculin regulates the recruitment and release of core focal adhesion proteins in a force-dependent manner. *Curr Biol* **23**, 271–281.
- 77 Bakolitsa C, Cohen DM, Bankston LA, Bobkov AA, Cadwell GW, Jennings L, Critchley DR, Craig SW & Liddington RC (2004) Structural basis for vinculin activation at sites of cell adhesion. *Nature* **430**, 583–586.
- 78 Borgon RA, Vonrhein C, Bricogne G, Bois PRJ & Izard T (2004) Crystal structure of human vinculin. *Structure* **12**, 1189–1197.
- 79 del Rio A, Perez-Jimenez R, Liu R, Roca-Cusachs P, Fernandez JM & Sheetz MP (2009) Stretching single talin rod molecules activates vinculin binding. *Science* **323**, 638–641.
- 80 Gingras AR, Ziegler WH, Frank R, Barsukov IL, Roberts GC, Critchley DR & Emsley J (2005) Mapping and consensus sequence identification for multiple vinculin binding sites within the talin rod. *J Biol Chem* **280**, 37217–37224.
- 81 Cohen DM, Kutscher B, Chen H, Murphy DB & Craig SW (2006) A conformational switch in vinculin drives formation and dynamics of a talin-vinculin complex at focal adhesions. *J Biol Chem* **281**, 16006–16015.
- 82 Humphries JD, Wang P, Streuli C, Geiger B, Humphries MJ & Ballestrem C (2007) Vinculin controls focal adhesion formation by direct interactions with talin and actin. *J Cell Biol* **179**, 1043–1057.

- 83 Weekes J, Barry ST & Critchley DR (1996) Acidic phospholipids inhibit the intramolecular association between the N- and C-terminal regions of vinculin, exposing actin-binding and protein kinase C phosphorylation sites. *Biochem J* **314** (Pt 3), 827–832.
- 84 Lee HS, Anekal P, Lim CJ, Liu CC & Ginsberg MH (2013) Two modes of integrin activation form a binary molecular switch in adhesion maturation. *Mol Biol Cell* **24**, 1354–1362.
- 85 Sun Z, Tseng HY, Tan S, Senger F, Kurzawa L, Dedden D, Mizuno N, Wasik AA, They M, Dunn AR *et al.* (2016) Kank2 activates talin, reduces force transduction across integrins and induces central adhesion formation. *Nat Cell Biol* **18**, 941–953.
- 86 Calderwood DA, Huttenlocher A, Kiosses WB, Rose DM, Woodside DG, Schwartz MA & Ginsberg MH (2001) Increased filamin binding to beta-integrin cytoplasmic domains inhibits cell migration. *Nat Cell Biol* **3**, 1060–1068.
- 87 Roca-Cusachs P, del Rio A, Puklin-Faucher E, Gauthier NC, Biaias N & Sheetz MP (2013) Integrin-dependent force transmission to the extracellular matrix by alpha-actinin triggers adhesion maturation. *Proc Natl Acad Sci USA* **110**, E1361–E1370.
- 88 Fukuda K, Bledzka K, Yang J, Perera HD, Plow EF & Qin J (2014) Molecular basis of kindlin-2 binding to integrin-linked kinase pseudokinase for regulating cell adhesion. *J Biol Chem* **289**, 28363–28375.
- 89 Wickström SA, Lange A, Montanez E & Fässler R (2010) The ILK/PINCH/parvin complex: the kinase is dead, long live the pseudokinase!. *EMBO J* **29**, 281–291.
- 90 Tu Y, Wu S, Shi X, Chen K & Wu C (2003) Migfilin and Mig-2 link focal adhesions to filamin and the actin cytoskeleton and function in cell shape modulation. *Cell* **113**, 37–47.
- 91 Zhu L, Liu H, Lu F, Yang J, Byzova TV & Qin J (2019) Structural basis of paxillin recruitment by kindlin-2 in regulating cell adhesion. *Structure* **27**, 1686–1697.e5.
- 92 Bottcher RT, Veelders M, Rombaut P, Faix J, Theodosiou M, Stradal TE, Rottner K, Zent R, Herzog F & Fassler R (2017) Kindlin-2 recruits paxillin and Arp2/3 to promote membrane protrusions during initial cell spreading. *J Cell Biol* **216**, 3785–3798.
- 93 Huvenceers S & Danen EH (2009) Adhesion signaling - crosstalk between integrins, Src and Rho. *J Cell Sci* **122**, 1059–1069.
- 94 Theodosiou M, Widmaier M, Böttcher RT, Rognoni E, Veelders M, Bharadwaj M, Lambacher A, Austen K, Müller DJ, Zent R & *et al.* (2016) Kindlin-2 cooperates with talin to activate integrins and induces cell spreading by directly binding paxillin. *Elife* **5**, e10130.
- 95 Liu J, Fukuda K, Xu Z, Ma Y-Q, Hirbawi J, Mao X, Wu C, Plow EF & Qin J (2011) Structural basis of phosphoinositide binding to kindlin-2 protein pleckstrin homology domain in regulating integrin activation. *J Biol Chem* **286**, 43334–43342.
- 96 Perera HD, Ma Y-Q, Yang J, Hirbawi J, Plow EF & Qin J (2011) Membrane binding of the N-terminal ubiquitin-like domain of kindlin-2 is crucial for its regulation of integrin activation. *Structure* **19**, 1664–1671.
- 97 Anthis NJ, Wegener KL, Ye F, Kim C, Goult BT, Lowe ED, Vakonakis I, Bate N, Critchley DR, Ginsberg MH *et al.* (2009) The structure of an integrin/talin complex reveals the basis of inside-out signal transduction. *EMBO J* **28**, 3623–3632.
- 98 Goksoy E, Ma YQ, Wang X, Kong X, Perera D, Plow EF & Qin J (2008) Structural basis for the autoinhibition of talin in regulating integrin activation. *Mol Cell* **31**, 124–133.
- 99 Kelley CF, Litschel T, Schumacher S, Dedden D, Schwille P & Mizuno N (2020) Phosphoinositides regulate force-independent interactions between talin, vinculin, and actin. *Elife* **9**, e56110.
- 100 Calderwood DA, Zent R, Grant R, Rees DJ, Hynes RO & Ginsberg MH (1999) The Talin head domain binds to integrin beta subunit cytoplasmic tails and regulates integrin activation. *J Biol Chem* **274**, 28071–28074.
- 101 Garcia-Alvarez B, de Pereda JM, Calderwood DA, Ulmer TS, Critchley D, Campbell ID, Ginsberg MH & Liddington RC (2003) Structural determinants of integrin recognition by talin. *Mol Cell* **11**, 49–58.
- 102 Vinogradova O, Velyvis A, Velyviene A, Hu B, Haas T, Plow E & Qin J (2002) A structural mechanism of integrin alpha(IIB)beta(3) “inside-out” activation as regulated by its cytoplasmic face. *Cell* **110**, 587–597.
- 103 Wegener KL, Partridge AW, Han J, Pickford AR, Liddington RC, Ginsberg MH & Campbell ID (2007) Structural basis of integrin activation by talin. *Cell* **128**, 171–182.
- 104 Ye F, Petrich BG, Anekal P, Lefort CT, Kasirer-Friede A, Shattil SJ, Ruppert R, Moser M, Fassler R & Ginsberg MH (2013) The mechanism of kindlin-mediated activation of integrin alphaIIb beta3. *Curr Biol* **23**, 2288–2295.
- 105 Bledzka K, Liu J, Xu Z, Perera HD, Yadav SP, Bialkowska K, Qin J, Ma YQ & Plow EF (2012) Spatial coordination of kindlin-2 with talin head domain in interaction with integrin beta cytoplasmic tails. *J Biol Chem* **287**, 24585–24594.
- 106 Molony L, McCaslin D, Abernethy J, Paschal B & Burridge K (1987) Properties of talin from chicken gizzard smooth muscle. *J Biol Chem* **262**, 7790–7795.

- 107 Elosegui-Artola A, Trepats X & Roca-Cusachs P (2018) Control of mechanotransduction by molecular clutch dynamics. *Trends Cell Biol* **28**, 356–367.
- 108 Eyckmans J, Boudou T, Yu X & Chen CS (2011) A hitchhiker's guide to mechanobiology. *Dev Cell* **21**, 35–47.
- 109 Roca-Cusachs P, Conte V & Trepats X (2017) Quantifying forces in cell biology. *Nat Cell Biol* **19**, 742–751.
- 110 Yusko EC & Asbury CL (2014) Force is a signal that cells cannot ignore. *Mol Biol Cell* **25**, 3717–3725.
- 111 Chen W, Lou J, Evans EA & Zhu C (2012) Observing force-regulated conformational changes and ligand dissociation from a single integrin on cells. *J Cell Biol* **199**, 497–512.
- 112 Katsumi A, Naoe T, Matsushita T, Kaibuchi K & Schwartz MA (2005) Integrin activation and matrix binding mediate cellular responses to mechanical stretch. *J Biol Chem* **280**, 16546–16549.
- 113 Roca-Cusachs P, Iskratsch T & Sheetz MP (2012) Finding the weakest link: exploring integrin-mediated mechanical molecular pathways. *J Cell Sci* **125**, 3025–3038.
- 114 Huang DL, Bax NA, Buckley CD, Weis WI & Dunn AR (2017) Vinculin forms a directionally asymmetric catch bond with F-actin. *Science* **357**, 703–706.
- 115 Mei L, de Los E, Reyes S, Reynolds MJ, Leicher R, Liu S & Alushin GM (2020) Molecular mechanism for direct actin force-sensing by α -catenin. *Elife* **9**, 213.
- 116 Xu X-P, Pokutta S, Torres M, Swift MF, Hanein D, Volkman N & Weis WI (2020) Structural basis of α E-catenin-F-actin catch bond behavior. *Elife* **9**, 531.
- 117 Buckley CD, Tan J, Anderson KL, Hanein D, Volkman N, Weis WI, Nelson WJ & Dunn AR (2014) Cell adhesion. The minimal cadherin-catenin complex binds to actin filaments under force. *Science* **346**, 1254211.
- 118 Geiger B & Yamada KM (2011) Molecular architecture and function of matrix adhesions. *Cold Spring Harb Perspect Biol* **3**, a005033.
- 119 Wehrle-Haller B (2012) Structure and function of focal adhesions. *Curr Opin Cell Biol* **24**, 116–124.