

Online Search for a Hyperplane in High-Dimensional Euclidean Space

Antonios Antoniadis*, Ruben Hoeksma*,
Sándor Kisfaludi-Bak[†] and Kevin Schewior[‡]

September 2021

Abstract

We consider the online search problem in which a server starting at the origin of a d -dimensional Euclidean space has to find an arbitrary hyperplane. The best-possible competitive ratio and the length of the shortest curve from which each point on the d -dimensional unit sphere can be seen are within a constant factor of each other. We show that this length is in $\Omega(d) \cap O(d^{3/2})$.

1 Introduction

In d -dimensional Euclidean space, given a set of objects $\mathcal{T} \subseteq \mathcal{P}(\mathbb{R}^d)$, the *online search problem* (for an object in \mathcal{T}) asks for a curve (or *search strategy*) $\zeta : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}^d$ with $\zeta(0) = 0$ that minimizes the competitive ratio [14]. We say that ζ is *c-competitive* if there exists an $\alpha \in \mathbb{R}_{\geq 0}$ such that

$$\text{length}(\zeta|_{[0,t^*]}) \leq c \cdot \text{dist}(0, T) + \alpha$$

for all $T \in \mathcal{T}$, where $t^* = \inf\{t : \zeta(t) \in T\}$. The infimum over all such c is the *competitive ratio* of ζ . Problems of this type date back to Beck [4] and Bellman [6].

The arguably most basic and well-known such online search problem from this class is the version in which $d = 1$ and $\mathcal{T} = \mathbb{R}$, also known as the cow-path problem, where the best-possible competitive ratio is 9 [5], achieved by visiting the points $(-2)^0, (-2)^1, \dots$ sequentially. In fact, the class of all 9-competitive strategies has been investigated more closely [1]. Moreover, the competitive ratio can be improved to about 4.591 using randomization [5]. The problem has been generalized to more than two paths starting at the origin [13], searching various types of objects in the plane or lattice [3], and many more scenarios. For a (slightly outdated) survey, see the book by Gal [11].

*Department of Applied Mathematics, University of Twente, Enschede, The Netherlands.
{a.antoniadis,r.p.hoeksma}@utwente.nl

[†]Institute for Theoretical Studies, ETH Zürich, Zurich, Switzerland.
sander.kisfaludi-bak@mpi-inf.mpg.de

[‡]Department of Mathematics and Computer Science, University of Cologne, Cologne, Germany.
schewior@cs.uni-koeln.de

Another very natural generalization is the case of general d and \mathcal{T} being equal to \mathcal{H}^{d-1} , the set of all hyperplanes in \mathbb{R}^d . We call this version the *d-Dimensional Hyperplane Search Problem*. When \mathcal{T} is a set of affine subspaces, this is arguably the most interesting case. Note that, if \mathcal{T} contains all affine subspaces of dimension $d' \leq d - 2$, any constant competitive ratio is ruled out, even when d is fixed. Surprisingly, in addition to the case of $d = 1$, results only seem to be known for $d = 2$ and for offline versions [9, 2], leaving a gap in the online-algorithms literature. For $d = 2$, it is conjectured that a logarithmic spiral that achieves a competitive ratio of about 13.811 is optimal [3, 10]. The present paper addresses the aforementioned gap by providing the first asymptotic results for $d \rightarrow \infty$.

For the remainder of the paper, we will consider the essentially equivalent but arguably cleaner *Sphere Inspection Problem*: The goal is to find a d -dimensional minimum-length closed curve, γ , that *inspects the unit sphere* in \mathbb{R}^d , i.e., γ sees every point p on the unit sphere S^{d-1} . Here, we say that an object $O \subseteq \mathbb{R}^d$ *sees* a point p' on the surface of the unit sphere if there is a point $p \in O$ such that the line segment pp' intersects the unit ball exclusively at p' . Note that the curve is not required to start in the origin any more. Such a minimum-length curve exists for any dimension [12]. While this problem is trivial for $d \in \{1, 2\}$, it has only been shown recently that the best-possible length for $d = 3$ is 4π [12]. No results for higher dimensions are known. Such visibility problems have also been considered from an algorithmic point of view, e.g., [7, 8].

While the connection between hyperplane search and sphere inspection seems to be folklore (e.g., [12]), we state it formally and provide a short proof for completeness. Let I be an interval, and for $\beta \in \mathbb{R}$, a set $T \subseteq \mathbb{R}^d$, and curve $\gamma : I \rightarrow \mathbb{R}^d$, we denote by $\beta \cdot T$ the set $\{\beta \cdot x : x \in T\}$ and by $\beta \cdot \gamma$ the curve $I \rightarrow \mathbb{R}^d, t \mapsto \beta \cdot \gamma(t)$.

Proposition 1. *Let $f : \mathbb{N} \rightarrow \mathbb{R}_{\geq 0}$. The following statements are equivalent:*

- (i) *There exists a length- $O(f(d))$ closed curve in \mathbb{R}^d that inspects S^{d-1} .*
- (ii) *There exists an $O(f(d))$ -competitive strategy for the d -Dimensional Hyperplane Search Problem.*

Proof. We first show that (i) implies (ii). Fix $d \in \mathbb{N}$, let γ be the length- ℓ curve for the Sphere Inspection Problem in \mathbb{R}^d , and let γ_0 be a point on γ . We show that there exists a (12ℓ) -competitive strategy for the d -Dimensional Hyperplane Search Problem where the additive constant α is equal to 3ℓ . Our strategy works in an infinite number of phases, starting with Phase 0. In each Phase i , the strategy ζ consists of a straight line from the origin to $2^i\gamma_0$, the curve $2^i \cdot \gamma$, and a straight line back from $2^i\gamma_0$ to the origin.

In the following, we show that, in Phase i , all hyperplanes of distance at most 2^i from the origin are visited, and the total distance traversed in Phase i is at most $3 \cdot 2^i \cdot \ell$, which implies the claim. We show two parts separately:

- We show that all claimed hyperplanes are visited by ζ . Consider Phase i and a hyperplane $H \in \mathcal{H}^{d-1}$ at distance at most 2^i from the origin. Let n be a unit normal vector of H . Note that $2^i \cdot \gamma$ sees both the point $2^i n$ and the point $-2^i n$. Let p_1 be a point from which $2^i \cdot \gamma$ sees $2^i n$, and let p_2 be a point from which it sees $-2^i n$. Note that, if $\text{dist}(0, H) = 2^i$, then p_1 or p_2 may be on H . In that case we are done. Otherwise p_1 and p_2 are on different sides of H , and, by the intermediate value theorem, $2^i \cdot \gamma$ intersects H .

- We next bound the distance traversed by ζ in Phase i . Note that it suffices to show that the start point of $2^i \cdot \gamma$, which is equal to its end point, has distance at most $2^i \ell$ from the origin. By scaling, we can restrict to showing $\|\gamma_0\|_2 \leq \ell$. Let H be the hyperplane $\langle \gamma_0, x \rangle = 0$. Note that γ must intersect H since otherwise γ does not see $-\gamma_0/\|\gamma_0\|_2$, thus $\text{dist}(\gamma_0, H) \leq \ell$. Since this distance (of γ_0 to H) is precisely $\|\gamma_0\|_2$, it follows that $\|\gamma_0\|_2 \leq \ell$.

We now show that (ii) implies (i). Again fix $d \in \mathbb{N}$. Assume that there exists a curve γ starting in the origin and $\alpha \in \mathbb{R}_{\geq 0}$ such that, for all hyperplanes $H \in \mathcal{H}^{d-1}$, γ visits H after traversing at most a length of $c \cdot \text{dist}(0, H) + \alpha$. We show that, for any $\varepsilon > 0$, there exists a, not necessarily closed, length- $(c + \varepsilon)$ curve that inspects the sphere. This then implies the existence of a length- $(2c + \varepsilon)$ closed curve that inspects the sphere. We define

$$\zeta := \frac{\varepsilon}{\alpha} \cdot \gamma|_{[0, t^*]}, \text{ where } t^* := \sup_{H \in \mathcal{H}^{d-1}: \text{dist}(0, H) \leq \alpha/\varepsilon} \min\{t : \gamma(t) \in H\}.$$

Note that the minimum exists because H is closed. Since for each $H \in \mathcal{H}^{d-1}$ with $\text{dist}(0, H) \leq \alpha/\varepsilon$, γ visits H after traversing at most a length of $c \cdot \alpha/\varepsilon + \alpha$, the length of $\gamma|_{[0, t^*]}$ is at most $c \cdot \alpha/\varepsilon + \alpha$, so the length of ζ is at most $c + \varepsilon$.

It remains to be shown that ζ indeed sees every point $p \in S^{d-1}$. Let H be the hyperplane tangent to the unit sphere in p . Further let $t := \min\{t : \gamma(t) \in \alpha/\varepsilon \cdot H\}$. By definition of t^* , we have $t \leq t^*$. Further, by definition of ζ , we have $\zeta(t) \in H$, implying that $\zeta(t)$ sees p . \square

In Section 2, we give an auxiliary lemma. In Sections 3 and 4 we will use that lemma and show the following two theorems.

Theorem 1. *Any curve in \mathbb{R}^d that inspects S^{d-1} has length at least $2d$.*

Theorem 2. *There exists a closed curve γ in \mathbb{R}^d of length $(2d)^{3/2}$ that inspects S^{d-1} .*

2 An Auxiliary Lemma

The following lemma simplifies thinking about the Sphere Inspection Problem.

Lemma 3. *Let $P \subset \mathbb{R}^d$ be the convex hull of some point set V . Then, the following two statements are equivalent.*

(i) *We have $S^{d-1} \subset P$.*

(ii) *The set V sees every point $p \in S^{d-1}$.*

Proof. We start by showing that (i) implies (ii). Let H be the hyperplane that is tangent to the unit sphere in p . If H contains a point $v \in V$, then the lemma directly follows since the line segment vp lies completely within H and therefore does not intersect the unit sphere other than at p . So assume that there is no point $v \in V$ that is contained in H . Then, and since P contains one point of the hyperplane ($p \in S^{d-1}$, which is contained in P by (i)), there must exist two points $v_1, v_2 \in V$ which are separated by H . Without loss of generality, let v_1 be the point in the halfspace defined by H whose interior is disjoint




Figure 1: The apex of the cone C sees all points on the unit sphere that B does.

from S^{d-1} . Then the line segment v_1p is completely contained in that halfspace, and since the interior of the halfspace is disjoint from S^{d-1} , v_1 can see p .

Now we show that (ii) implies (i). Towards a contradiction, assume that there exists $p^* \in S^{d-1}$ with $p^* \notin P$. Then consider a hyperplane H that separates p^* from P , and let n be a unit normal vector of H pointing away from P . Consider the hyperplane H' that is tangent to S^{d-1} in n . Clearly, both P and $S^{d-1} \setminus \{n\}$ are contained in one open halfspace defined by H' . Note that no point in this halfspace can see n . Therefore, no $v \in V$ can see n ; a contradiction. \square

3 Lower Bound

The goal of this section is to prove Theorem 1. Towards this, let γ be a curve in \mathbb{R}^d that inspects the unit sphere. We cut γ into a minimum number of contiguous portions of length at most δ for some fixed $\delta < 2$. Let ξ_1, \dots, ξ_n be the resulting tour portions, where $n = \lceil |\gamma|/\delta \rceil$. Choose a portion ξ , and let x be its midpoint. Clearly ξ is contained in the ball B that has center x and radius $\delta/2$. Further define C to be the cone that is the intersection of all halfspaces that contain both B and S^{d-1} and whose defining hyperplanes are tangent to both B and S^{d-1} . Note that the set of points on the sphere that can be seen by the curve ξ can also be seen from the apex of C , as visualized by Figure 1. This holds since the radius of B is $\delta/2 < 1$. Note that a single point $p \in \mathbb{R}^d$ can see some subset of an open hemisphere H of the unit sphere. Let H_1, \dots, H_n denote a set of open hemispheres such that H_i covers the portion of the sphere seen by ξ_i . Since γ inspects the sphere, we have that $(H_i)_{i=1}^n$ covers the sphere.

We need the following lemma.

Lemma 4. *The minimum number of open hemispheres that cover S^{d-1} is $d + 1$.*

Proof. To show that $d + 1$ points are sufficient, choose a simplex containing the sphere. Now consider the set of open hemispheres whose poles are colinear with the origin and a vertex of that simplex. Indeed, since the simplex contains the sphere, by Lemma 3, the set of vertices of the simplex sees every point $p \in S^{d-1}$. Since each point sees only a subset of the corresponding open hemisphere, the upper bound follows.

For the lower bound, we can use induction on d . Clearly the circle S^1 needs at least three open half-circles to be covered. For S^{d-1} , we have that the boundary of the first hemisphere H_1 is S^{d-2} , and each hemisphere H_i can cover at most an open hemisphere of ∂H_1 . So by induction at least $(d - 2) + 2 = d$ hemispheres are needed to cover ∂H_1 , and thus we have that at least $d + 1$ hemispheres are needed to cover S^{d-1} . \square

By Lemma 4 we have that $n \geq d + 1$, implying $\lceil \text{length}(\gamma)/\delta \rceil \geq d + 1$ and therefore $\text{length}(\gamma)/\delta \geq d$. With $\delta = 2 - \varepsilon$, we have that $\text{length}(\gamma) \geq (2 - \varepsilon) \cdot d$ for all $\varepsilon > 0$, and thus $\text{length}(\gamma) \geq 2d$.

4 Upper Bound

In this section we prove Theorem 2. Let C^d be the d -dimensional cross-polytope, i.e., the polytope $\{x \in \mathbb{R}^d : \|x\|_1 \leq 1\}$. Define $\bar{C}^d := \sqrt{d} \cdot C^d = \{x \in \mathbb{R}^d : \|x\|_1 \leq \sqrt{d}\}$, as shown in Figure 2. We claim that the vertices of \bar{C}^d inspect the unit sphere.

Lemma 5. *For any point $p \in S^{d-1}$, there exists a vertex v of \bar{C}^d that sees p .*

Proof. We prove that \bar{C}^d contains the unit sphere; then Lemma 3 shows the claim. In other words we prove that for any point $x \in \mathbb{R}^d$ with $\|x\|_2 \leq 1$ (so, $x \in S^{d-1}$), it also holds that $\|x\|_1 \leq \sqrt{d}$. Indeed, by the Cauchy-Schwartz inequality we have that $\|x\|_1 \leq \sqrt{d} \cdot \|x\|_2$ for any x which in turn can be upper bounded by \sqrt{d} for any $x \in \mathbb{R}^d$ with $\|x\|_2 \leq 1$. \square

Let $G(\bar{C}^d)$ be the graph¹ of polytope \bar{C}^d . Note that $G(\bar{C}^d)$ is the so-called *cocktail party graph* which can be obtained by removing a perfect matching from a complete graph on $2d$ vertices (indeed, in \bar{C}^d any vertex v has an edge to any other vertex except $-v$). We next prove that $G(\bar{C}^d)$ is Hamiltonian.

Lemma 6. *The graph $G(\bar{C}^d)$ is Hamiltonian.*

Proof. The proof is by induction on d . The statement clearly holds for \mathbb{R}^2 where the cross polytope is a square, and $G(\bar{C}^2)$ itself is a Hamiltonian cycle that we denote by c^2 . Consider a Hamiltonian cycle c^{d-1} for $G(\bar{C}^{d-1})$. Note that $G(\bar{C}^d)$ can be constructed by $G(\bar{C}^{d-1})$ and adding two nodes, v^* and $-v^*$ that are connected to each of the nodes of $G(\bar{C}^{d-1})$. To construct a cycle c^d of $G(\bar{C}^d)$, take any two distinct edges $\{v_1, v_2\}$ and $\{v_3, v_4\}$ contained in c^{d-1} and replace them with the edges $\{v_1, v^*\}, \{v^*, v_2\}$ and $\{v_3, -v^*\}, \{-v^*, v_4\}$, respectively. The resulting tour is connected, visits all nodes of

¹The graph $G(P)$ of a polytope P is a graph with a node for each vertex of P and an edge connecting those nodes if P has an edge between the corresponding vertices. The graph of P is also referred to as its 1-skeleton.




Figure 2: The cross polytope that has vertices at distance \sqrt{d} from the origin contains the unit sphere, and thus, a tour of its vertices (thick) inspects the unit sphere.

$G(\bar{C}^d)$ exactly once and the used edges are contained in the edge set of $G(\bar{C}^d)$, thus ensuring the feasibility of c^d . \square

By Lemma 5 it directly follows that any closed curve that visits all the vertices of \bar{C}^d inspects the unit sphere, and therefore so does the closed curve γ corresponding to the Hamiltonian cycle c^d in the skeleton of \bar{C}_d . To complete the proof of Theorem 2, note that each edge of \bar{C}_d has a length of $\sqrt{2d}$ and that γ traverses $2d$ such edges.

5 Conclusion

In this paper, we narrowed down the optimal competitive ratio for the d -Dimensional Hyperplane Search Problem to $\Omega(d) \cap O(d^{3/2})$. The obvious open problem is closing this gap.

Acknowledgements. We thank Paula Roth for helpful discussions.

References

- [1] Spyros Angelopoulos, Christoph Dürr, and Shendan Jin. Best-of-two-worlds analysis of online search. In *International Symposium on Theoretical Aspects of Computer Science (STACS)*, volume 126, pages 7:1–7:17, 2019.

- [2] Antonios Antoniadis, Krzysztof Fleszar, Ruben Hoeksma, and Kevin Schewior. A PTAS for euclidean TSP with hyperplane neighborhoods. *ACM Trans. Algorithms*, 16(3):38:1–38:16, 2020.
- [3] Ricardo A. Baeza-Yates, Joseph C. Culberson, and Gregory J. E. Rawlins. Searching in the plane. *Information and Computation*, 106(2):234–252, 1993.
- [4] Anatole Beck. On the linear search problem. *Naval Research Logistics*, 2:221–228, 1964.
- [5] Anatole Beck and D. J. Newman. Yet more on the linear search problem. *Israel Journal of Mathematics*, 8(4):419–429, 1970.
- [6] Richard Bellmann. An optimal search problem. *SIAM Review*, 5:274, 1963.
- [7] Svante Carlsson, Håkan Jonsson, and Bengt J. Nilsson. Finding the shortest watchman route in a simple polygon. *Discret. Comput. Geom.*, 22(3):377–402, 1999.
- [8] Moshe Dror, Alon Efrat, Anna Lubiw, and Joseph S. B. Mitchell. Touring a sequence of polygons. In *ACM Symposium on Theory of Computing (STOC)*, pages 473–482. ACM, 2003.
- [9] Adrian Dumitrescu and Csaba D. Tóth. The traveling salesman problem for lines, balls, and planes. *ACM Trans. Algorithms*, 12(3):43:1–43:29, 2016.
- [10] Steven R. Finch and Li-Yan Zhu. Searching for a shoreline. *arXiv preprint math/0501123*, 2005.
- [11] Shmuel Gal. *Search Games*. Academic Press, 1980.
- [12] Mohammad Ghomi and James Wenk. Shortest closed curve to inspect a sphere. *Journal für die reine und angewandte Mathematik (Crelle’s Journal)*, 2021. Forthcoming, preprint on arXiv under abs/2010.15204.
- [13] Ming-Yang Kao, John H. Reif, and Stephen R. Tate. Searching in an unknown environment: An optimal randomized algorithm for the cow-path problem. *Information and Computation*, 131(1):63–79, 1996.
- [14] Anna R. Karlin, Mark S. Manasse, Larry Rudolph, and Daniel Dominic Sleator. Competitive snoopy caching. *Algorithmica*, 3:77–119, 1988.