

RIF1 and KAP1 differentially regulate the choice of inactive versus active X chromosomes

Elin Enverald^{1,2,†}, Lynn Marie Powell¹ , Lora Boteva¹, Rossana Foti^{2,‡}, Nerea Blanes Ruiz³, Gözde Kibar⁴, Agnieszka Piszczek^{2,§}, Fatima Cavaleri^{2,¶}, Martin Vingron⁴, Andrea Cerase³  & Sara B C Buonomo^{1,2,*} 

Abstract

The onset of random X chromosome inactivation in mouse requires the switch from a symmetric to an asymmetric state, where the identities of the future inactive and active X chromosomes are assigned. This process is known as X chromosome choice. Here, we show that RIF1 and KAP1 are two fundamental factors for the definition of this transcriptional asymmetry. We found that at the onset of differentiation of mouse embryonic stem cells (mESCs), biallelic up-regulation of the long non-coding RNA *Tsix* weakens the symmetric association of RIF1 with the *Xist* promoter. The *Xist* allele maintaining the association with RIF1 goes on to up-regulate *Xist* RNA expression in a RIF1-dependent manner. Conversely, the promoter that loses RIF1 gains binding of KAP1, and KAP1 is required for the increase in *Tsix* levels preceding the choice. We propose that the mutual exclusion of *Tsix* and RIF1, and of RIF1 and KAP1, at the *Xist* promoters establish a self-sustaining loop that transforms an initially stochastic event into a stably inherited asymmetric X-chromosome state.

Keywords KAP1; RIF1; *Tsix*; X chromosome inactivation; *Xist*

Subject Categories Chromatin, Transcription & Genomics; Development

DOI 10.15252/emboj.2020105862 | Received 7 June 2020 | Revised 5 October 2021 | Accepted 19 October 2021 | Published online 17 November 2021

The EMBO Journal (2021) 40: e105862

Introduction

X chromosome inactivation (XCI) is the process leading to the stable transcriptional silencing of one of the two X chromosomes in female placental mammals, with the aim of equalising the expression of X-linked genes between males and females (Lyon, 1961). This process represents one of the best-studied examples of how different nuclear processes, such as epigenetic control, 3D organisation of chromatin contacts, sub-nuclear positioning and, potentially, replication-timing

regulation, are integrated to achieve transcriptional control. Random XCI (rXCI) is initiated when *Xist*, an X-encoded long non-coding RNA (lncRNA) is up-regulated from one of the two X chromosomes, the future inactive X chromosome (Xi) (Brockdorff *et al*, 1991; Brown *et al*, 1991; Penny *et al*, 1996; Marahrens *et al*, 1997). *In vivo*, this happens around the time of embryo implantation (Monk & Harper, 1978; Rastan, 1982), while in cultured female mouse embryonic stem cells (mESCs), XCI takes place during a narrow time-window at the onset of differentiation (Wutz & Jaenisch, 2000). Monoallelic up-regulation of *Xist* is coupled to loss of pluripotency and several activating and repressing factors of this process have been described (Lee & Lu, 1999; Navarro *et al*, 2008; Jonkers *et al*, 2009; Tian *et al*, 2010; Chureau *et al*, 2011; Gontan *et al*, 2012; Makhlof *et al*, 2014; Furlan *et al*, 2018). Guided by the three-dimensional (3D) conformation of the X chromosome (Engreitz *et al*, 2013; Simon *et al*, 2013), *Xist* spreads *in cis* and recruits SPEN to enhancers and promoters of the X-linked genes to trigger their silencing (Chu *et al*, 2015; McHugh *et al*, 2015; Moindrot *et al*, 2015; Monfort *et al*, 2015; Dossin *et al*, 2020), and the exclusion of RNA polymerase II (Chaumeil *et al*, 2006; Kucera *et al*, 2011). This in turn promotes the recruitment of the Polycomb Repressive Complexes (PRC1/2) and the accumulation of tri-methylated H3K27 (H3K27me3) (Sun *et al*, 2006; Zhao *et al*, 2008) and monoubiquitinated H2AK119 (H2AK119ub) (de Napoles *et al*, 2004) on the future inactive X chromosome (Xi). Contextually, Lamin B receptor (LBR) tethers the future Xi to the nuclear periphery to facilitate *Xist* spreading into active gene regions and the maintenance of the silent state (Chen *et al*, 2016). Finally, the entire Xi switches the replication timing to mid-late S-phase (Takagi *et al*, 1982). The combination of all these events facilitates the attainment of an exceptionally stable transcriptionally silent status, so robustly controlled that it is maintained throughout the entire life of the organism. One of the least understood of all these steps is the mechanism that, in the initiating phase of XCI, directs the random choice of which one of the two *Xist* alleles to up-regulate, marking the future Xi, and which to silence (marking the future active X chromosome, Xa). We will refer to this

¹ Institute of Cell Biology, School of Biological Sciences, University of Edinburgh, Edinburgh, UK

² Epigenetics & Neurobiology Unit, European Molecular Biology Laboratory (EMBL Rome), Monterotondo, Italy

³ Blizard Institute, Centre for Genomics and Child Health, Barts and The London School of Medicine and Dentistry, Queen Mary University of London, London, UK

⁴ Max-Planck-Institut fuer molekulare Genetik, Berlin, Germany

*Corresponding author (Lead contact). E-mail: sara.buonomo@ed.ac.uk

[†]Present address: Department of Molecular Biosciences, The Wenner-Gren Institute, Stockholm University, Stockholm, Sweden

[‡]Present address: Center for Basic and Translational Neuromedicine, Faculty of Health and Medical Sciences, University of Copenhagen, København N, Denmark

[§]Present address: Histology Facility Vienna BioCenter Core Facilities (VBCF), Vienna, Austria

[¶]Present address: Evotec SE, Hamburg, Germany

process as the “choice” (Avner & Heard, 2001). This is a key stage, as failure to establish monoallelic *Xist* expression can result in either both X chromosomes being silenced or both remaining active, consequently leading to embryonic lethality (Takagi & Abe, 1990; Marahrens et al, 1997; Borensztein et al, 2017). *Tsix* is a lncRNA encoded by a gene that overlaps, in the antisense orientation, with *Xist*, and plays a well-established role as an *in cis* repressor of *Xist* (Lee & Lu, 1999). In female mESCs, *Tsix* is bi-allelically expressed, to become down-regulated on one of the two X chromosomes, the future Xi, at the onset of differentiation, hence allowing for *in cis* *Xist* up-regulation (Lee et al, 1999; Stavropoulos et al, 2001). The switch to mono-allelic expression of *Tsix* is important in determining the destinies of the future Xi (*Tsix* silenced) and Xa (*Tsix* transiently maintained). In fact, interfering with the expression of one of the two *Tsix* alleles in female mESCs results in a non-random choice, with the *Tsix*-defective chromosome pre-determined as the future Xi (reviewed in (Starmer & Magnuson, 2009)). Although *Tsix* down-regulation is essential for *in cis* up-regulation of *Xist*, the molecular mechanism of *Tsix*-driven silencing is still unclear. The *Tsix* terminator region overlaps with the *Xist* promoter, and *Tsix* transcription through this region and/or *Tsix* RNA have been proposed to be essential for *Xist* repression (Shibata & Lee, 2004) by promoting a transient silenced chromatin state (Navarro et al, 2005, 2006; Sado et al, 2005; Ohhata et al, 2008). The establishment of the opposite *Xist/Tsix* expression patterns on the two genetically identical X chromosomes must rely on the coordinated asymmetric distribution of activators and/or repressors of transcription.

Mathematical modelling can recapitulate the experimental features of XCI by postulating the existence of an *in cis*-negative regulator of *Xist* (cXR) and an *in trans*, X-linked, *Xist* activator (tXA) (Mutzel et al, 2019). While a cXR is sufficient to explain the maintenance of mono-allelic *Xist* expression, a tXA is needed to explain: 1. the establishment of the *Xist* mono-allelic expression; 2. the female specificity of XCI; 3. the resolution of bi-allelic *Xist* expression detected, to various extents, in different organisms (Mutzel et al, 2019). In mouse, *Tsix* is the most likely cXR, while RNF12, an X-linked ubiquitin ligase that functions as a dose-dependent initiator of XCI (Jonkers et al, 2009; Gontan et al, 2012), has been proposed as one of the potential tXA. However, while overexpression of *Rlim* (*Rnf12*) in male cells can induce XCI (Jonkers et al, 2009), its deletion in females is not sufficient to prevent *Xist* up-regulation (Shin et al, 2014; Wang et al, 2017). Thus, RNF12 could account for the X-linked aspects of the tXA function, such as female specificity and resolution of bi-allelic expression, but one or multiple other transactivators must be contributing to the asymmetric control of *Xist* expression. Moreover, conceptually, the expression level of a single, X-linked gene, does not constitute a switch robust or sensitive enough to be the only element to control a clear-cut bi-stable state for *Xist* (active on one and silent on the other allele) (Mutzel & Schulz, 2020). The establishment of *in cis*, self-reinforcing and mutually exclusive circuits on the two *Xist* alleles could create the ultrasensitivity required to generate a binary switch-type of control (Mutzel & Schulz, 2020). Key to this model, is the idea that the initial stochastic events will trigger a chain of local, mutually exclusive and self-sustaining events to bookmark both Xi and Xa.

RIF1 is a multifaceted protein, required for the regulation of several of the nuclear processes that take place during XCI. RIF1 is the only known genome-wide regulator of replication timing

(Cornacchia et al, 2012; Hayano et al, 2012; Yamazaki et al, 2012; Hiraga et al, 2014; Peace et al, 2014; Foti et al, 2016; Seller & O’Farrell, 2018). It confines long-range chromatin contacts within the respective boundaries of the nuclear A/B compartments (Gnan et al, 2021) and plays an as yet unclear function in the control of gene expression (Daxinger et al, 2013; Foti et al, 2016; Tanaka et al, 2016; Zofall et al, 2016; Li et al, 2017; Toteva et al, 2017). RIF1 is an adaptor for Protein Phosphatase 1 (PP1), one of the main Ser/Thr phosphatases in eukaryotic cells (Trinkle-Mulcahy et al, 2006; Dave et al, 2014; Hiraga et al, 2014, 2017; Mattarocci et al, 2014; Sreesankar et al, 2015; Alver et al, 2017). In *Drosophila melanogaster*, the RIF1-PP1 interaction was shown to be essential during embryonic development (Seller & O’Farrell, 2018). In addition, the RIF1-PP1 interaction is essential for RIF1-dependent organisation of chromatin contacts (Gnan et al, 2021). Others (Chapman et al, 2013; Daxinger et al, 2013) and we (this work) have observed that mouse RIF1 deficiency is associated with a sex-linked differential lethality, with the female embryos dying around the time of implantation. These data have suggested that RIF1 could be important during XCI. Here we find that RIF1, present biallelically on the *Xist* P2-promoter in female mESCs, becomes asymmetrically enriched at P2 on the future Xi, concomitant with the choice, at the time when *Tsix* expression switches from bi- to mono-allelic. RIF1 then plays an essential role in *Xist* up-regulation, thus determining the future Xi. The removal of RIF1 from the future Xa arises from the KAP1-dependent increase of *Tsix* bi-allelic expression that leads to the choice. Our data identify the KAP1-dependent regulation of *Tsix* levels and the consequent transition of RIF1 association with *Xist* promoter from symmetric to asymmetric, as key steps in the molecular cascade that leads to the identification of the future Xi and Xa.

Results

RIF1 is required for X inactivation during embryonic development and for *Xist* up-regulation

The analysis of the progeny derived from inter-crosses of mice heterozygous for a *Rif1* null allele (*Rif1*^{+/-}, Appendix Fig S1A and B) in a pure C57BL/6J genetic background has revealed that *Rif1* is essential for embryonic development (Fig 1A). In contrast, in a mixed genetic C57BL/6J-129/SvJ background, *Rif1* deletion results in a differential lethality between the sexes (Fig 1B). Indeed, in this case, only a small proportion of the expected *Rif1*^{-/-} mice, exclusively males, are recovered at weaning. In order to pinpoint more precisely the time of the onset of lethality, we have analysed the frequency of recovery of *Rif1*^{-/-} embryos at different stages of development in a C57BL/6J pure background. We found that, up to the blastocyst stage (E3.5), there are no obvious differences in the number of male and female *Rif1* null and wild-type embryos recovered (our unpublished observation). However, by E7.5, although still recoverable, *Rif1* null female embryos are already dead/abnormal (Fig 1C and D). In contrast, males appear to die only around mid-gestation (Fig 1C). This early-onset lethality observed specifically in females suggests that the lack of RIF1 could interfere with the process of XCI, as the timing coincides with the onset of random XCI.

Given the diversity of its roles, RIF1 could act at one or several of the multiple steps during XCI. To dissect at what stage(s) of the




Figure 1. *Rif1* deficiency leads to female embryonic lethality at peri-implantation.

A, B Tables summarising the number and the sex of the pups recovered at weaning from *Rif1*^{+/-} x *Rif1*^{+/-} mice inter-crosses, either in a C57BL/6j (A) or in a mixed C57BL/6j-129/Svj genetic background (B). The observed number of mice is compared to the expected number, based on the Mendelian ratio. *P* calculated by χ^2 .
C The table summarises the number and the sex of the embryos of the indicated genotypes, recovered from timed matings of *Rif1*^{+/-} x *Rif1*^{+/-} mice, in a C57BL/6j genetic background. The day of gestation (E) is indicated. (D). Representative images of *Rif1*^{-/-} E7.5 embryos, female top and male bottom.

process RIF1 is required, we generated female mESCs carrying homozygous conditional *Rif1* allele (*Rif1*^{Flox/Flox}) and a tamoxifen-inducible CRE recombinase (*Rosa26*^{Cre-ERT/+}, Buonomo *et al*, 2009). To trigger XCI in the absence of *Rif1*, we set up a protocol in which we combined differentiation by embryoid body (EB) formation (Doetschman *et al*, 1985) and tamoxifen treatment (Fig 2A and Materials and Methods). By RT-qPCR as well as by RNA sequencing, we found that *Rif1* deletion (Fig 2B) severely impairs *Xist* up-regulation (Figs 2C, EV1A and B, and EV2A) and, consequently, the enrichment of H3K27me3 on the future Xi (Fig 2D). Failure of *Xist* up-regulation in the absence of *Rif1* is not due to a general defect in exit from pluripotency (Figs EV1C and EV2B, D and E) or to failed commitment to differentiation (Figs EV1C and EV2C–E). Moreover, during the early stages of differentiation the levels of the main negative regulator of *Xist*, Tsix, appear to be reduced faster in *Rif1*

knockout cells compared to the control (Appendix Fig S2A). Finally, the overall dynamics of RNF12 appear comparable between control and *Rif1* knockout cells (Fig EV1B and Appendix Fig S2B). Overall, these results indicate that failure of *Xist* up-regulation is the likely cause of defective XCI in *Rif1* null female embryos and that RIF1 could directly and positively regulate *Xist* expression.

RIF1 is a positive regulator of *Xist* and its binding specifically bookmarks the future Xi

Xist is controlled from two promoters, P1 and P2 (Johnston *et al*, 1998), separated by a repetitive region essential for the silencing properties of *Xist* (Wutz *et al*, 2002). While the epigenetic control of the upstream P1 promoter was shown to be important for *Xist* regulation (Navarro *et al*, 2005), P2 appears to serve as an internal




Figure 2. *Rif1* null female mESCs fail to up-regulate *Xist* upon differentiation.

- A** Overview of the experimental design. *Rif1*^{+/+} and *Rif1*^{F/F} mESCs were grown for 2 days in medium supplemented with 4-Hydroxytamoxifen (OHT) to induce the translocation of the Cre recombinase into the nucleus, leading to *Rif1* deletion in the *Rif1*^{F/F} cells (*Rif1*^{-/-}). The embryoid body (EB) differentiation protocol was then started to trigger XCI. OHT was kept in the medium during the first 24 h of EB differentiation. Cells were differentiated up to 4 (RNA analysis) or 7 days (H3K27me3 IF).
- B** Representative western blot to monitor RIF1 levels after Cre-mediated *Rif1* deletion and EB differentiation. SMC1: loading control.
- C** Time course analysis of *Xist* RNA expression by RT-qPCR during EB differentiation of *Rif1*^{+/+} (*Rif1*^{+/+} +OHT) and *Rif1*^{-/-} (*Rif1*^{F/F} +OHT) cells at the indicated timepoints. *Rif1*^{+/+} (solid line) and *Rif1*^{-/-} (dashed line), female (black) and male (grey). Data are presented as mean ± standard deviation from three (female lines) or two (male lines) independent experiments. Statistical significance was determined using two-way ANOVA comparing female *Rif1*^{+/+} to female *Rif1*^{-/-} cell lines (*****P* ≤ 0.0001). *Xist* RT-primers *Xist* ex7 F and R were used. Values are normalised to a geometric mean consisting of the expression of *Gapdh*, *Ubiquitin* and *β-Actin*.
- D** Bar plot summarising the number of cells showing H3K27me3-marked Xi as a percentage of total cells counted, in *Rif1*^{+/+} (*Rif1*^{+/+} +OHT) and *Rif1*^{-/-} (*Rif1*^{F/F} +OHT) female mESCs at the indicated days of EB differentiation. Averages ± standard deviation from three (day 4 and 7) and two (day 2) independent experiments (*n* > 200).

regulatory unit, possibly modulating the expression from P1 (Makhoul *et al*, 2014). We found that RIF1 is enriched specifically at *Xist* P2 promoter, both in mESCs (Fig 3A, Appendix Fig S2C and D) and in early EBs (Fig 3B), supporting the hypothesis that RIF1 could be a direct regulator of *Xist* expression. In agreement with this, we found that P2 harbours two potential RIF1-binding sites, defined by the presence of a consensus sequence derived from the analysis of RIF1 genome-wide distribution by ChIP-seq in female mESCs (Foti *et al*, 2016) (Appendix Fig S3A). To confirm that RIF1 association with *Xist* promoter has a positive effect on *Xist* expression, we used a

reporter assay system, where *Xist* promoter has been cloned upstream of a firefly Luciferase gene (Gontan *et al*, 2012). We found that, upon differentiation, in the absence of RIF1, the induction of Luciferase from the *Xist* promoter is significantly reduced (Fig 3C), supporting the hypothesis that RIF1 association with P2 exerts a positive, direct effect on *Xist* transcription.

Upon differentiation, *Xist* is mono-allelically transcribed, up-regulated only from the future Xi. If RIF1 acts as a positive regulator of *Xist*, we would expect it to be associated mono-allelically, specifically with P2 on the future Xi. In order to test this hypothesis, we




Figure 3.

Figure 3. RIF1 associates with *Xist* promoter on the future Xi.

- A, B RIF1 association with the *Xist* promoter assessed by ChIP-qPCR in two independent *Rif1*^{+/+} (*Rif1*^{+/+} +OHT, black) and two *Rif1*^{-/-} (*Rif1*^{F/F} +OHT, grey) female cell lines, in ESCs (A) and at 2 days of EB differentiation (B). P1 and P2 indicate the two *Xist* promoters, 5' indicates a region 2 kb upstream of *Xist* TSS. Inter1 and 2 are two intergenic regions that serve as negative controls. Peak and cRAD represent two previously identified regions of RIF1 association (positive control). See Appendix Fig S2C for primer positions within *Xist*. Mean \pm standard deviation from three independent experiments (A) and two independent experiments (B). *P* calculated by Student's two-tailed, paired *t* test comparing RIF1 association in *Rif1*^{+/+} cells on *Xist* P2 and P1 versus 5'. **P* \leq 0.05, *****P* \leq 0.0001 and ns = not significant.
- C *Rif1* deletion decreases the efficiency of up-regulation of a *Luciferase* reporter under the control of *Xist* promoter (*Xist*-2p-luc), at 2 days of EB differentiation. As a control (Luc), empty luciferase reporter vector was transfected in parallel. The average of three independent experiments is shown. Error bars indicate the standard deviation. *P* calculated by Student's two-tailed, unpaired *t* test, for comparison of fold activation of *Xist*-2p-Luc normalised to empty vector (Luc) in *Rif1*^{+/+} versus *Rif1*^{-/-} cells. ***P* \leq 0.01. See Appendix Material and Methods for details about the normalisation.
- D Association of RIF1 with *Xist* P2 in the Fa2L cells (black) and a wild-type female mESC line (grey), also harbouring one *castaneus* and one 129 X chromosome. Allele-specific ChIP-qPCR primers were used, *cast* indicates association with the *castaneus* *Xist* P2 and 129 indicates association with the 129 *Xist* P2. Enrichments are presented relative to input DNA. Mean \pm standard deviation from three independent experiments. *P* calculated by Student's two-tailed, paired *t* test comparing RIF1 association with the *castaneus* and with the 129 X chromosome *Xist* P2, ****P* \leq 0.001. Below is the schematic of the *Xist*/*Tsix* alleles in the Fa2L undifferentiated cells.
- E Association of RIF1 with *Xist* P2 in the Fa2L cells (black) upon differentiation. The analysis was performed as in (D). Mean \pm standard deviation from three independent experiments. *P* calculated by Student's two-tailed, paired *t* test comparing RIF1 association with the *castaneus* and with the 129 X chromosome *Xist* P2. **P* \leq 0.05. Below is the schematic of the *Xist*/*Tsix* alleles in the Fa2L differentiated cells.
- F Quantification by ChIP-qPCR of RIF1 association with the indicated regions in the Fa2L cell line, following treatment with DMSO only (black) or flavopiridol (grey). Primers as in (E).
- G Same as in (F) but for a wild-type female mESC line. All enrichments are presented relative to input DNA. Mean \pm standard deviation from three (F) and two (G) independent experiments are presented. Statistical significance was determined using Student's two-tailed, paired *t* test (**P* \leq 0.05, ***P* \leq 0.01 and ns = not significant).

have taken advantage of the Fa2L cell line, in which: 1. the two X chromosomes can be discriminated, as one originates from *Mus castaneus* (*cast*) and the other from *Mus musculus 129/SvJ* (129) mouse strains; 2. Xa (*cast*) and Xi (129) are pre-determined, as the 129 *Tsix* allele carries a transcriptional stop signal, approximately 4 kb downstream from the *Tsix* major promoter (Fig 3D, scheme and (Luikenhuis *et al*, 2001)). *Xist* is, therefore, preferentially up-regulated from the 129-derived X chromosome. We have analysed the association of RIF1 with *Xist* P2 promoter of the future Xa and Xi by allele-specific ChIP-qPCR (Appendix Fig S3B) and found that RIF1 is preferentially associated with the *Xist* P2 promoter of the 129 *Xist* allele (future Xi) in both mESCs (Fig 3D) and upon differentiation (Fig 3E). Importantly, in control wild-type mESCs (bi-allelically expressed *Tsix*), also carrying one *cast* and one 129 X chromosome, RIF1 is equally distributed on both P2 promoters (Fig 3D). This suggests that the asymmetric association of RIF1 with the future Xi is concomitant with/follows the switch from bi- to mono-allelic *Tsix* expression that accompanies the choice and allows *Xist* monoallelic up-regulation. As in the case of RIF1 conditional cells, depletion of RIF1 in Fa2L cells (Appendix Fig S3C) also compromises *Xist* up-regulation (Appendix Fig S3D). These data show that RIF1's asymmetric association with the future Xi parallels the choice and that it is essential for *Xist* up-regulation.

RIF1 asymmetric localisation on the future Xi is driven by *Tsix* expression

How is the transition from bi- to mono-allelic RIF1 association with *Xist* promoter regulated? While this would generally be triggered by differentiation, in undifferentiated Fa2L cells it is pre-determined and RIF1 is preferentially associated with the X chromosome that does not express full-length *Tsix* transcript (Fig 3D and E). This suggests that *Tsix* RNA and/or transcription could destabilise RIF1 association with the *Xist* promoter. In agreement with this hypothesis, we found that blocking *Tsix* expression by treating mESCs with

the CDK9-inhibitor flavopiridol, which inhibits transcriptional elongation (Chao & Price, 2001) (Fig EV3A) or, briefly, with triptolide, an inhibitor of transcription initiation (Fig EV3B), is sufficient to revert RIF1 preferential association with the future Xi in the Fa2L cells to a symmetric mode of binding (Figs 3F and EV3C). In addition, flavopiridol treatment of wild-type mESCs also leads to an increased P2 association of RIF1 (Fig 3G), indicating that this is not an effect specific to the Fa2L cells. Finally, while this work was under review, RIF1 has been found associated with *Tsix* RNA in mESCs (Aeby *et al*, 2020), supporting the hypothesis that *Tsix* RNA can compete for RIF1 association with *Xist* P2 in the genome.

KAP1 is important for the Xa/Xi choice

With the aim of understanding the molecular mechanism by which RIF1 regulates *Xist* expression, we have investigated whether some of the known transcriptional regulators associated with RIF1 (Sukackaite *et al*, 2017) are also required for XCI. We focused in particular on KAP1, as KAP1 and RIF1 have already been shown to regulate overlapping targets, such as *Dux* and *MERVLs* (Maksakova *et al*, 2013; Li *et al*, 2017; Percharde *et al*, 2018). We found that knock down of *Kap1* (Appendix Fig S4A and B) impairs *Xist* up-regulation (Figs 4A and EV4A), similarly to the knockout of *Rif1*. This is not due to compromised exit from pluripotency (Fig EV4B), impaired activation of the differentiation transcriptional program (Fig EV4C) or reduced RIF1 levels (Fig EV4D), suggesting that diminished *Xist* activation is not a consequence of an overall impaired cell differentiation. In addition, the dynamics of expression of *RNF12* appear comparable between control and *Kap1* knock down cells (Fig EV4E). However, in contrast to the depletion of *Rif1*, depletion of *Kap1* in Fa2L cells (Appendix Fig S4C), where the choice is pre-determined, has no consequences for *Xist* up-regulation (Fig 4B). These data suggest that KAP1 is required prior to or at the time of the choice, while it is dispensable once *Tsix* mono-allelic expression has been established. In agreement with a

role during the choice, we found that Kap1 knock down affects *Tsix* dynamic regulation at the onset of differentiation. In wild-type cells, during the early stages of differentiation, *Tsix* levels rise transiently (at 1, or 1 and 2 days of EB differentiation respectively, depending on the culture conditions, Fig 4C and Appendix Fig S2A). The boost corresponds to an increased detection of *Tsix* RNA from both alleles (Fig 4D), suggesting that this step precedes the switch to *Tsix* mono-

allelic expression and the consequent choice of Xa/Xi. Upon Kap1 down-regulation, we found not only a failure in the temporary boost of *Tsix* levels (Fig 4C) but also a failure to evolve towards *Tsix* mono-allelic expression, as *Tsix* becomes undetectable (Fig 4D). In a situation of pre-determined choice (Fa2L cells), *Tsix* levels remain low, even upon differentiation, and Kap1 knock down has no further effect (Appendix Fig S4D).




Figure 4.

Figure 4. KAP1 regulates the Xa/Xi choice through Tsix.

- A Time course analysis of *Xist* expression by RT-qPCR during EB differentiation of female mESCs following knock down of Luciferase (Control, black) and Kap1 (Kap1 KD, grey), at the indicated timepoints. Data are presented as mean \pm standard deviation from three independent experiments. Statistical significance was determined using two-way ANOVA. *Xist* primers Xist ex3 F and Xist ex4 R were used. Normalisation was performed using a geometric mean consisting of the expression of *Rplp0*, *Ubiquitin* and *Sdha* (* $P \leq 0.05$).
- B RT-qPCR analysis of *Xist* expression levels during differentiation of the Fa2L cells, following expression of shRNA against Luciferase (Control, black) and Kap1 (Kap1 KD, grey), at the indicated timepoints. Mean \pm standard deviation from a minimum of three independent experiments is presented. Two-way ANOVA was used to determine statistical significance. ns = not significant.
- C *Tsix* RNA levels in female mESCs infected with shRNA directed against Luciferase (Control, black) and KAP1 (Kap1 KD, grey), during differentiation. Mean \pm standard deviation values from four independent experiments are shown. Statistical significance was determined using two-way ANOVA. (** $P \leq 0.01$). Values have first been normalised to a geometric mean consisting of the expression of *Rplp0*, *Ubiquitin* and *Sdha*.
- D RNA FISH analysis of *Tsix* expression during differentiation of female mESCs expressing an shRNA directed against Luciferase (control) or against Kap1 (Kap1 KD). Left: Cells with no (0, red), one (1, grey) and two or more (2, black) *Tsix* foci were counted in two independent experiments, shown averaged. Statistical significance was determined by χ^2 . A minimum of 110 cells were counted per time point for each line (** $P \leq 0.01$, **** $P \leq 0.0001$). Right: examples of cells with one (top) or two (central and bottom) *Tsix* FISH signals. Scale bars: 5 μ m.

In summary, the failure to up-regulate *Xist* caused by *Rif1* deletion and by Kap1 knock down have very different causes. While RIF1 is directly required to promote *Xist* up-regulation, KAP1's function is to drive the transient increase of *Tsix* levels that precedes the choice. The consequent failure to up-regulate *Xist* when Kap1 is knocked down could be caused, in this case, by a failure to execute the choice. The low, bi-allelic *Tsix* levels typical of mESCs instead evolve directly towards an absence of *Tsix*.

RIF1 negatively regulates KAP1 association with the *Xist* promoter/*Tsix* terminator in mESCs

KAP1 is a multifunctional protein, and a key global regulator of transcription, involved in several aspects of gene expression modulation. Through its interaction with the H3K9 histone methyltransferase SetDB1, KAP1 can promote transcriptional silencing. Alternatively, it can modulate transcriptional or transcript levels, either regulating the release of RNA polymerase II proximal pausing from the promoter (especially at genes encoding for lncRNAs (Bunch *et al*, 2016)), or as part of the 7SK complex (McNamara *et al*, 2016). This is a ribonucleoprotein

complex with roles both at the promoter and in the transcriptional termination of several genes, including several lncRNAs (Castelo-Branco *et al*, 2013).

To gain an insight into the mechanism by which KAP1 regulates *Tsix* levels, we have analysed KAP1 distribution along *Tsix* regulatory regions. Consistent with a function during the choice, we could not detect KAP1 on any of the regions examined in mESCs. Instead, we found that KAP1 was specifically recruited to *Xist* P2 promoter at the onset of differentiation, around the time when *Tsix* levels are boosted (Fig 5A). Taking advantage of the Fa2L cells, we could also determine that KAP1 associates preferentially with *Xist* P2 of the future Xa (*castaneus* allele, Figs 5B and EV4F). KAP1 and RIF1 occupy, therefore, the same region, but with complementary spatial (Xa versus Xi) and temporal dynamics (KAP1 appears on *Xist* P2 on the Xa when RIF1 leaves it). In order to understand if these events are coordinated, we have investigated whether RIF1 regulates KAP1 association with *Xist* P2. We found that *Rif1* deletion leads to KAP1 binding to *Xist* promoter, even in undifferentiated cells (Fig 5C). This is not due to a general increase of *Kap1* expression (Fig EV1A), *KAP1* protein levels (Fig EV5A) or its overall binding to chromatin (Fig EV5B). Moreover, KAP1 enrichment is specific for *Xist*

Figure 5. RIF1 negatively regulates KAP1 association with *Xist* P2.

- A ChIP-qPCR analysis of KAP1 association with the indicated sites in wild-type female mESCs (*Rif1*^{+/+}, same as used in Fig 3B but without OHT) and during early differentiation. ZFP629 is a well-characterised KAP1 associated region (positive control). Xite A and C indicate two regions within the *Tsix* enhancer *Xite*, *Tsix* region 1 indicates *Tsix* major promoter, *Tsix* region 2 indicates the Dxpas34 region, *Tsix* region 3 indicates a region slightly downstream of the Dxpas34 region. P1 and P2 indicate the two *Xist* promoters, 5' indicates a region 2 kb upstream of *Xist* TSS. Inter1 is an intergenic region. See Appendix Fig S2C for the positions of the primers within *Xist* and *Tsix*. The data are presented as mean \pm standard deviation from three (2d EB and 1d EB) and two (ESCs) independent experiments. Statistical significance was calculated by Student's two-tailed unpaired *t* test comparing RIF1 association with *Xist* P2 and P1 in 2d EB versus 1d EB (* $P \leq 0.05$ and ns = not significant). In the inset, *Tsix* RNA levels were quantified by RT-qPCR during the differentiation of wild-type female ESCs shown in Fig 4F. The average of two experiments is shown. *Tsix* values are normalised to a geometric mean consisting of the expression of *Rplp0*, *Ubiquitin* and *Sdha*. Error bars indicate standard deviations.
- B Using allele-specific primers, ChIP-qPCR was used to measure the association of KAP1 with *Xist* P2 in the Fa2L cells (black) and a wild-type female mESC line also harbouring one *castaneus* and one 129 X chromosome (grey). *cast* indicates association with the *castaneus* *Xist* P2 and 129 indicates association with the 129 *Xist* P2. Enrichments are presented relative to input DNA. Mean \pm standard deviation from a minimum of three independent experiments. Statistical significance was determined by Student's two-tailed, paired *t* test. Below is the schematic of the *Xist*/*Tsix* alleles in the Fa2L cells.
- C KAP1 association with *Xist* promoter in two independent *Rif1*^{+/+} (*Rif1*^{+/+} +OHT, black) and two *Rif1*^{-/-} (*Rif1*^{+/+} +OHT, grey) female mESC cell lines. *Ezr* is an additional region known to be associated with KAP1 in mESCs. Enrichments are presented relative to input DNA. Mean \pm standard deviation from a minimum of three independent experiments per cell line are displayed. Statistical significance was determined using Student's two-tailed, unpaired *t* test comparing the KAP1 association with *Xist* P2 and P1 in *Rif1*^{+/+} versus *Rif1*^{-/-} cells (** $P \leq 0.01$).
- D Allele-specific KAP1 association with *Xist* P2 in Fa2L cells following knock down of Luciferase (Control, black) and *Rif1* (*Rif1* KD, grey). *cast* indicates association with the *castaneus* *Xist* P2 promoter and 129 indicates association with the 129 *Xist* P2 promoter. Enrichments are presented relative to input DNA. Average \pm standard deviation of two independent experiments.




Figure 5.

promoter, as other regions known to be associated with KAP1 that we have tested, like *Zfp629* (Fig 5C and our unpublished observation) (Ding *et al.*, 2018), did not show an increased KAP1 association upon *Rif1* deletion. Importantly, the effect of RIF1 deficiency is unlikely to be due to an indirect, general remodelling of the *Xist* promoter chromatin, as the association of another P2-specific transcription factor and *Xist* activator, Yin-Yang-1 (YY1) (Makhlouf *et al.*, 2014), is unchanged in *Rif1* knockout cells (Fig EV5C). We also found that knocking down *Rif1* in undifferentiated Fa2L cells (Fig EV5D) facilitates KAP1 association with *Xist* P2 (Fig 5D)

comparably to what happens in *Rif1* conditional cells upon induction of *Rif1* deletion (Fig 5C). Specifically, KAP1 gains access to the future Xi (129 allele, carrying the truncated *Tsix* allele), where normally RIF1 is preferentially localised (Fig 3D and E). Overall, these data indicate that, in mESCs, RIF1 is symmetrically associated with *Xist* P2 on both X chromosomes, protecting P2 from the binding of KAP1. Upon triggering differentiation, the bi-allelic increase of *Tsix* levels weakens RIF1 association with DNA, facilitating the transition of RIF1 to an asymmetric association with one of the two *Xist* promoters, the future Xi, and the consequent association of




Figure 6. KAP1 recruits the 7SK complex to the *Xist* P2 promoter/*Tsix* terminator of the future Xa.

- A, B H3K9me3 association with the *Xist* promoter (P1 and P2) in two independent *Rif1*^{+/+} and *Rif1*^{-/-} cell lines, analysed by ChIP-pPCR in mESCs (A) and 2d EBs (B). Myf5 serves as a positive control region, β actin negative. Average ± standard deviation of two independent experiments.
- C HEXIM1 association with the *Xist* promoters (P1 and P2) in two independent *Rif1*^{+/+} and two *Rif1*^{-/-} mESC lines, analysed by ChIP-qPCR. Tkbp1 and Myf5 are two control regions. As in the case of KAP1 association, deletion of *Rif1* induces accumulation of HEXIM1 on *Xist* P1 and P2. Average ± standard deviation of two independent experiments. *P* values were calculated by two-tailed, unpaired, equal variance *t* test. (***P* ≤ 0.01, ****P* ≤ 0.001).
- D Western blot analysis of KAP1 levels in protein extracts from Fa2L cells after Kap1 knock down. SMC1: loading control. Quantification of KAP1 levels normalised to SMC1 and relative to Luciferase control cells are shown below.
- E Upon infection of Fa2L cells with shRNA against Kap1 or control, against Luciferase, HEXIM1 association with P2 was analysed, on both alleles, by ChIP-qPCR. As in the case of KAP1, HEXIM1 shows preferential association with *Xist* P2 on the Cast allele (future Xa, Control). The association is lost upon knock down of Kap1 (Kap1 KD). Myf5 serves as a negative and Neat1 as a positive control region. (**P* ≤ 0.05, ***P* ≤ 0.01).

KAP1 with the other *Xist* promoter, on the future Xa. This event, in turn, sustains the KAP1-dependent increase of *Tsix* levels that precedes the switch to *Tsix* mono-allelic expression and the choice, further reinforcing RIF1 exclusion from P2 on the future Xa.

KAP1 recruits the 7SK complex to *Tsix* terminator

The timing of recruitment, the RIF1-dependent regulation and the preferential enrichment on the future Xa support the idea that KAP1

functions by promoting the choice, possibly *in cis*. The association of KAP1 with *Xist* P2 promoter on the future Xa suggests that KAP1 could repress *Xist*. However, Kap1 knock down does not induce precocious up-regulation of *Xist* (Figs 4A and EV4A), nor does KAP1 early association with *Xist* P2 promoter in *Rif1* null mESCs and EBs lead to increased tri-methylation of histone H3K9 (Fig 6A and B). These observations do not support the hypothesis of KAP1 regulating the choice through *Xist* repression. An alternative hypothesis is that KAP1 could instead regulate *Tsix* either by controlling its transcriptional termination and, consequently, RNA stability (reviewed in Peck *et al*, 2019), or by promoting the formation of a terminator–promoter-positive feedback loop (Tan-Wong *et al*, 2008), to boost *Tsix* transcription. *Xist* P2 promoter, in fact, overlaps with *Tsix* transcriptional terminator. In support of either of these hypotheses, we have found that, as in the case of KAP1, the 7SK complex component HEXIM1 is also enriched on *Xist* promoter/*Tsix* terminator in *Rif1* knockout mESCs (Fig 6C), and it is associated with the future Xa in Fa2L cells, in a KAP1-dependent manner (Fig 6D and E). Overall, these data suggest that KAP1 could promote the choice of the future Xa by sustaining *in cis* the increase of *Tsix* levels that would stabilise the asymmetric RIF1 distribution.

Discussion

While marsupials have adopted an imprinted X inactivation strategy, eutherians have evolved a mechanism based on the random choice of the X chromosome to be inactivated. The latter can contribute to a higher degree of resistance of females to pathogenic X-linked mutations and increase phenotypic diversity. Despite its importance, the mechanisms guiding the random choice are still unclear, partially because of the randomness and consequent heterogeneity in the cell population, partially because of the inaccessibility of the early embryos, where the process takes place naturally and, finally, because of the inherent difficulty of identifying asymmetry involving two identical chromosomes.

Several lines of evidence suggest that *Tsix* is involved in the choice-making process. For example, introduction of a stop codon that blocks *Tsix* transcript before its overlap with *Xist* (Luikenhuis *et al*, 2001), or deletions of its major promoter (Vigneau *et al*, 2006), or of the GC-rich repeat region that immediately follows it (Dxpas34) (Lee & Lu, 1999), or insertion of a gene trap in the same region, that abolishes the production of *Tsix* RNA (Sado *et al*, 2001), result in a non-random choice, with the *Tsix*-defective chromosome as the future Xi. Moreover, monoallelic down-regulation of *Tsix* levels by deleting *Xite*, a *cis*-acting element that positively regulates *Tsix*, also skews the choice (Ogawa & Lee, 2003). Interestingly, *Xist* itself can influence the choice, in a yet-to-be-understood feedback control loop. *Xist* ectopic up-regulation can in fact skew the choice in favour of the *Xist*-overexpressing chromosome (Newall *et al*, 2001; Nesterova *et al*, 2003).

Our experiments show that RIF1 association with the *Xist* P2 promoter is negatively regulated by *Tsix* expression or RNA levels. *Tsix* could, therefore, be the determinant of the asymmetric association of RIF1 with the future Xi at the choice. We would like to propose a model (Fig 7) whereby, at the onset of differentiation, the transient, bi-allelic increase of *Tsix* levels will promote a weaker or more dynamic association of RIF1 with *Xist* P2, thus creating a window of opportunity for KAP1 stochastic association with either

allele. The KAP1-bound allele will go on to sustain higher *Tsix* steady-state levels *in cis*, thus skewing RIF1 association with the opposite allele, and initiating a self-reinforcing loop on the future Xa. On the future Xi, RIF1 will promote *Xist* up-regulation, thus establishing the inactivation. The negative effect of RIF1 on KAP1 association with *Xist* promoter in ESCs is at the heart of the mutual exclusion, reinforced by KAP1's positive effect on the levels of *Tsix*, that is, in turn, a negative regulator of RIF1 association with *Xist* promoter. How RIF1 excludes KAP1 is currently unclear, but we can envisage at least two potential mechanisms, based either on RIF1/KAP1 competition for binding to a shared site, RNA or protein partner, or through KAP1 de-phosphorylation by RIF1-associated PP1. Phosphorylation of KAP1 has indeed been shown to regulate KAP1 association with heterochromatin protein 1 (HP1) (Chang *et al*, 2008).

In support of our model, we have shown that, the association of KAP1 with the P2 region upon differentiation coincides with the detection of higher levels of *Tsix* RNA (Fig 5A), and this increase is dependent upon KAP1 (Fig 4C and D). The molecular mechanism by which KAP1 modulates *Tsix* levels is currently unknown. The data presented here suggest that KAP1 could modulate *in cis* *Tsix* transcriptional up-regulation, termination and/or RNA stability through the 7SK complex. Finally, we cannot exclude a model where KAP1 promotes *Tsix* increase *in trans*, through a yet unknown differentiation-induced factor. In this case, the association of KAP1 with *Xist* P2 could contribute *in cis* to the identification of Xa, by establishing a stable repression of *Xist* promoter, with RIF1 shielding the future Xi by excluding KAP1. Although our data do not support the hypothesis of KAP1-dependent silencing of *Xist* (Figs 4A and EV4A) through H3K9me3 (Fig 6A and B), KAP1 could promote repression through a different mechanism, for example, DNA methylation (Coluccio *et al*, 2018).

Our data show that the increase of *Tsix* that precedes and, possibly, leads to a proficient choice, requires KAP1. It has been previously shown that failure to set up the choice as a consequence of homozygous deletion of *Tsix*, leads to a mixture of cells showing either no *Xist* up-regulation or bi-allelic up-regulation during differentiation (Lee, 2002, 2005). This is different from what we observe in Kap1 knock down cells, where we detect defective *Xist* up-regulation, but not bi-allelic expression. Nonetheless, a situation where, from the start of the process in ESCs, *Tsix* is always absent, as in the case of *Tsix*^{-/-}, is clearly different from the system where *Tsix* levels remain physiological until differentiation is triggered, as in the case of Kap1 knock down (Fig 4D).

The early embryonic lethality of *Rif1*^{-/-} females described here contrasts with the milder effect of *Xist* conditional inactivation in the epiblast described previously (Yang *et al*, 2016). However, beside the technical differences between a conditional system, where the efficiency of the deletion can be lower than 100%, and a knockout, RIF1 has at least two other key roles, in the regulation of the replication timing program (Cornacchia *et al*, 2012; Hayano *et al*, 2012; Yamazaki *et al*, 2012; Foti *et al*, 2016) and replication fork protection (Buonomo *et al*, 2009; Garzon *et al*, 2019). In fact, depending on the genetic background, most or some of the male embryos also die, although later during development (this work). We cannot, therefore, exclude that the early female lethality could derive from a synthetic effect of multiple problems, added on top of the failure of X inactivation.

In summary, we propose that, during the stochastic phase of the choice of the future Xi, *Tsix*-dependent destabilisation of the




Figure 7. Model for RIF1 and KAP1-dependent bookmarking of Xi and Xa respectively.

The low bi-allelic expression of *Tsix* in mESCs allows the association of RIF1 with P2 on both *Xist* alleles. However, the presence of pluripotency-dependent inhibitors will not allow *Xist* up-regulation, despite the presence of RIF1. (1) Upon differentiation, the increase in *Tsix* levels weakens the association of RIF1 with P2. This opens the opportunity for a stochastic KAP1 binding to P2 of one of the two alleles (2). KAP1 is required for sustained high levels of *Tsix*, further reinforcing RIF1 exclusion from the KAP1-bound/*Tsix* high allele and establishing the asymmetry. It is not known whether KAP1 gains access to P2 to promote the increase of *Tsix* levels first, or whether the increase of *Tsix* levels is initially triggered by a differentiation-dependent factor. (3). The pluripotency *Xist* inhibitors have been silenced, RIF1 promotes *Xist* expression on the future Xi. A self-sustainable binary switch is thus created and it consolidates the choice of the future Xi and Xa.

symmetric association of RIF1 with *Xist* P2 promoter sets in motion the establishment of two, mutually exclusive circuits that will identify Xi and Xa. RIF1's presence on P2, inhibiting KAP1 and promoting *Xist* expression will identify the future Xi. On the other allele, KAP1's presence on P2, sustaining Tsix levels and, thus, helping to exclude RIF1, will identify the Xa. The initial stochastic binding of KAP1 will thus become a binary switch, where a bi-stable, self-sustaining circuitry on the two X chromosomes is propagated.

Materials and Methods

mESC differentiation

Wild-type ESCs were plated onto non-coated Petri dishes at a concentration of 1×10^6 cells/10 cm², in a volume of 10 ml medium lacking 2i and LIF. At day 4 of differentiation the aggregated EBs were gently transferred to gelatinised tissue culture dishes. Medium was gently changed every 48 h with minimal disruption of the EBs. EBs were grown for up to 4 or 7 days in total. In experiments where cell differentiation was combined with *Rif1* deletion, the differentiation was preceded by 48 h of 4-hydroxytamoxifen (OHT, #H7904, Sigma-Aldrich) treatment, at a concentration of 200 nM in ES medium containing LIF and 2i. Differentiation was then started with 2×10^6 cells/10 cm² dish for *Rif1*^{+/+} and 2.5×10^6 cells/10 cm² for *Rif1*^{E/E} cells in a medium lacking 2i and LIF but containing 200 nM OHT. On day 1 of differentiation, the medium was replaced with a medium without OHT. On day 4 of differentiation, the EBs were transferred to gelatinised tissue culture dishes as above.

KAP1, RIF1 and HEXIM1 ChIP

Chromatin immunoprecipitation was performed according to Bulut-Karslioglu *et al.*, 2012). Briefly, for RIF1, KAP1 and HEXIM1 ChIP, collected cells were first cross-linked using 2 mM disuccinimidyl glutarate (DSG, # BC366 Synchem UG & Co. KG) in PBS for 45 min at RT while rotating, washed twice in PBS, followed by 10 min of additional cross-linking in 1% formaldehyde (#252549, Sigma-Aldrich) in cross-linking buffer (50 mM HEPES pH 7.8, 150 mM NaCl, 1 mM EDTA and 500 μM EGTA) at RT. Cross-linking was followed by 5 min quenching in 0.125 M glycine at RT, washed twice in cold PBS and resuspended in lysis buffer (1% SDS, 10 mM EDTA, 50 mM Tris-HCl pH 8.1, supplemented with protease inhibitor cocktail, #11873580 001, Roche). Chromatin fragmentation was performed using Soniprep 150 to produce a distribution of fragments enriched between 300 and 400 bp. The lysate was pre-cleared by centrifugation at low speed 400 g for 20 min at 4°C. Chromatin was quantified using Qubit dsDNA High Sensitivity assay kit (#Q32854, Life Technologies). Immunoprecipitation was performed by incubating 100 μg of chromatin diluted in 10 volumes of Dilution buffer (1% Triton X-100, 2 mM EDTA, 167 mM NaCl, 20 mM Tris-HCl pH 8.1, including Protease Inhibitor) overnight rotating at 4°C together with either α-KAP1, α-RIF1 or α-HEXIM1 antibodies (see Appendix Table S2) or IgG only control (#sc-2026, Santa Cruz), 10% of chromatin was isolated as input control. The following day, 50 μl of Dynabeads protein G slurry (#10004D, Thermo Fisher) per ChIP sample was added and incubated rotating for another 2 h at 4°C. The beads were magnet-separated and washed twice with low

salt buffer (0.1% SDS, 1% Triton X-100, 2 mM EDTA, 150 mM NaCl, 20 mM Tris-HCl pH8.1), one time each with high salt buffer (0.1% SDS, 1% Triton X-100, 2 mM EDTA, 500 mM NaCl, 20 mM Tris-HCl pH8.1), LiCl buffer (0.25 M LiCl, 0.5% NP-40, 0.5% sodium deoxycholate, 1 mM EDTA, 10 mM Tris-HCl pH 8.1) and finally TE. Each wash was performed for 5 min. on a rotating wheel at 4°C and all buffers were supplemented with protease inhibitor cocktail (#11873580 001, Roche). Prior to elution, samples were rinsed once in TE without protease inhibitor. ChIP-DNA was eluted from the beads by rotating at RT for 1 h in elution buffer (1% SDS, 100 mM NaHCO₃). Beads were separated and the supernatants as well as input samples were subjected to RNase A (#R5250, Sigma-Aldrich) treatment (37.5 μg/sample) for 1 h at 37°C followed by de-cross-linking using Proteinase K (#P6556, Sigma-Aldrich) treatment (45 μg/sample) overnight at 60°C. The following day, ChIP-DNA and input samples were purified using ChIP DNA Clean and Concentrator kit (#D5205, Zymo Research) and the retrieved DNA as well as input DNA was quantified using Qubit dsDNA High Sensitivity assay kit (#Q32854, Life Technologies). The concentration of ChIP-DNA and input samples was adjusted to maintain a similar ratio of ChIP-DNA:INPUT between different ChIP experiments. qPCRs were performed using the SYBR Green reaction mix (#04887352001, Roche) on a LightCycler 96 Instrument (Roche), following standard protocols. Enrichments over input control were calculated for each respective primer set. Primer sequences are presented in Appendix Table S3.

RNA extraction, reverse transcription and RT-qPCR

Frozen cell pellets were lysed and homogenised using QIAshredder column (#79656, QIAGEN) followed by RNA extraction using the RNeasy kit (#74106, QIAGEN) according to the manufacturer's instructions. On-column DNase treatment was performed at 25–30°C for 20 min. using RQ1 RNase-Free DNase (#M6101, Promega). After elution, a second round of DNase treatment was performed using 8 U of DNase/sample, incubated at 37°C for 20 min. The reaction was terminated by adding 1 μl of RQ1 DNase Stop Solution and incubated at 65°C for 10 min. RNA was quantified using Nanodrop, and cDNA synthesis was performed using RevertAid H Minus First Strand cDNA kit (#K1632, Thermo Scientific) using random hexamer priming. qPCRs were performed using the SYBR Green reaction mix (#04887352001, Roche) on a LightCycler 96 Instrument, following standard protocols. Gene expression data were normalised against a geometric mean generated by RT-qPCR of either: *Gapdh*, *Ubiquitin* and *β-Actin* or *Rplp0*, *Ubiquitin* and *Sdha*. For flavopiridol- or triptolide-treated cells, gene expression levels were normalised against 18S ribosomal RNA. Primer sequences are presented in Appendix Table S4.

Additional material and method descriptions can be found in the Appendix Material and Methods.

Data availability

The RNA-seq data have been deposited in the GEO database (GSE165704) and are available at <https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE165704>.

Expanded View for this article is available online.

Acknowledgements

We acknowledge David Kelly from the COIL facility, WTCCB, University of Edinburgh; Emerald Perlas from the Histology Facility of the Epigenetics & Neurobiology Unit, EMBL Rome; Violetta Parimbeni for mouse husbandry, (Epigenetics & Neurobiology Unit, EMBL Rome). We thank Phil Avner (Epigenetics & Neurobiology Unit, EMBL Rome) for advice, reagents, support, discussions and critically reading the manuscript. Rafael Galupa (EMBL Heidelberg) and Jacqueline Mermoud (University of Marburg) are thanked for critically reading the manuscript. Titia de Lange (The Rockefeller University) is thanked for initially supporting the generation of the *Rif1* knockout mice. Joost Gribnau and Cristina Contan (Erasmus MC, University Medical Center, Rotterdam) are thanked for the Xist-luciferase reporter plasmid. Andrew Jarman and Petra zur Lage (Centre for Discovery Brain Sciences, Edinburgh) and Sally Lowell (MRC Centre for Regenerative Medicine, Edinburgh) are all thanked for providing reagents. EE received funding from the European Union's Horizon 2020 research and the Marie Skłodowska-Curie Individual Fellowship grant agreement No. 660985 and from the ERC consolidator award 726130 to SCBB. LP and LB were funded by the ERC consolidator award 726130 to SBCB. RF was funded by the EMBL Interdisciplinary Postdoc (EIPOD) fellowship under Marie Curie Actions (COFUND). GK acknowledges funding from the IMPRS-BAC. AC is funded by a Rett Syndrome Research Trust (RSRT), BARTSCHARITY grants, and intramural QMUL support.

Author contributions

EE created the cellular system, performed the majority of the experiments and co-wrote the manuscript. RF initiated the project and performed some of the early experiments, like the staining of E3.5 embryos. LMP performed some of the ChIP experiments, the triptolide treatment, and the Luciferase assay. LB, AC and NBR performed KAP1 KD, RNA FISH and its analysis. GK analysed the RNA seq data, supervised by MV. NBR was supervised by AC, who also critically read the manuscript. FC and AP isolated and stained the E5.5 embryos. SBCB conceived the project, performed some of the experiments and wrote the manuscript.

Conflict of interest

The authors declare that they have no conflict of interest.

References

- Aeby E, Lee HG, Lee YW, Kriz A, Del Rosario BC, Oh HJ, Boukhali M, Haas W, Lee JT (2020) Decapping enzyme 1A breaks X-chromosome symmetry by controlling Tsix elongation and RNA turnover. *Nat Cell Biol* 22: 1116–1129
- Alver RC, Chadha GS, Gillespie PJ, Blow JJ (2017) Reversal of DDK-mediated MCM phosphorylation by Rif1-PP1 regulates replication initiation and replisome stability independently of ATR/Chk1. *Cell Rep* 18: 2508–2520
- Avner P, Heard E (2001) X-chromosome inactivation: counting, choice and initiation. *Nat Rev Genet* 2: 59–67
- Borensztein M, Syx L, Ancelin K, Diabangouaya P, Picard C, Liu T, Liang JB, Vassilev I, Galupa R, Servant N et al (2017) Xist-dependent imprinted X inactivation and the early developmental consequences of its failure. *Nat Struct Mol Biol* 24: 226–233
- Brockdorff N, Ashworth A, Kay GF, Cooper P, Smith S, McCabe VM, Norris DP, Penny GD, Patel D, Rastan S (1991) Conservation of position and exclusive expression of mouse Xist from the inactive X chromosome. *Nature* 351: 329–331
- Brown CJ, Ballabio A, Rupert JL, Lafreniere RG, Grompe M, Tonlorenzi R, Willard HF (1991) A gene from the region of the human X inactivation centre is expressed exclusively from the inactive X chromosome. *Nature* 349: 38–44
- Bulut-Karslioglu A, Perrera V, Scaranaro M, de la Rosa-Velazquez IA, van de Nobelen S, Shukeir N, Popow J, Gerle B, Opravil S, Pagani M et al (2012) A transcription factor-based mechanism for mouse heterochromatin formation. *Nat Struct Mol Biol* 19: 1023–1030
- Bunch H, Lawney BP, Burkholder A, Ma D, Zheng X, Motola S, Fargo DC, Levine SS, Wang YE, Hu G (2016) RNA polymerase II promoter-proximal pausing in mammalian long non-coding genes. *Genomics* 108: 64–77
- Buonomo SB, Wu Y, Ferguson D, de Lange T (2009) Mammalian Rif1 contributes to replication stress survival and homology-directed repair. *J Cell Biol* 187: 385–398
- Castelo-Branco G, Amaral PP, Engstrom PG, Robson SC, Marques SC, Bertone P, Kouzarides T (2013) The non-coding snRNA 7SK controls transcriptional termination, poisoning, and bidirectionality in embryonic stem cells. *Genome Biol* 14: R98
- Chang CW, Chou HY, Lin YS, Huang KH, Chang CJ, Hsu TC, Lee SC (2008) Phosphorylation at Ser473 regulates heterochromatin protein 1 binding and corepressor function of TIF1beta/KAP1. *BMC Mol Biol* 9: 61
- Chao SH, Price DH (2001) Flavopiridol inactivates P-TEFb and blocks most RNA polymerase II transcription in vivo. *J Biol Chem* 276: 31793–31799
- Chapman JR, Barral P, Vannier JB, Borel V, Steger M, Tomas-Loba A, Sartori AA, Adams IR, Batista FD, Boulton SJ (2013) RIF1 is essential for 53BP1-dependent nonhomologous end joining and suppression of DNA double-strand break resection. *Mol Cell* 49: 858–871
- Chaumeil J, Le Baccon P, Wutz A, Heard E (2006) A novel role for Xist RNA in the formation of a repressive nuclear compartment into which genes are recruited when silenced. *Genes Dev* 20: 2223–2237
- Chen CK, Blanco M, Jackson C, Aznauryan E, Ollikainen N, Surka C, Chow A, Cerase A, McDonel P, Guttman M (2016) Xist recruits the X chromosome to the nuclear lamina to enable chromosome-wide silencing. *Science* 354: 468–472
- Chu C, Zhang QC, da Rocha ST, Flynn RA, Bharadwaj M, Calabrese JM, Magnuson T, Heard E, Chang HY (2015) Systematic discovery of Xist RNA binding proteins. *Cell* 161: 404–416
- Chureau C, Chantalat S, Romito A, Galvani A, Duret L, Avner P, Rougeulle C (2011) Ftx is a non-coding RNA which affects Xist expression and chromatin structure within the X-inactivation center region. *Hum Mol Genet* 20: 705–718
- Coluccio A, Ecco G, Duc J, Offner S, Turelli P, Trono D (2018) Individual retrotransposon integrants are differentially controlled by KZFP/KAP1-dependent histone methylation, DNA methylation and TET-mediated hydroxymethylation in naive embryonic stem cells. *Epigenetics Chromatin* 11: 7
- Cornacchia D, Dileep V, Quivy JP, Foti R, Tili F, Santarella-Melliwig R, Antony C, Almouzni G, Gilbert DM, Buonomo SB (2012) Mouse Rif1 is a key regulator of the replication-timing programme in mammalian cells. *EMBO J* 31: 3678–3690
- Dave A, Cooley C, Garg M, Bianchi A (2014) Protein phosphatase 1 recruitment by Rif1 regulates DNA replication origin firing by counteracting DDK activity. *Cell Rep* 7: 53–61
- Daxinger L, Harten SK, Oey H, Epp T, Isbel L, Huang E, Whitelaw N, Apedaile A, Sorolla A, Yong J et al (2013) An ENU mutagenesis screen identifies novel and known genes involved in epigenetic processes in the mouse. *Genome Biol* 14: R96
- Ding D, Bergmaier P, Sachs P, Klangwart M, Ruckert T, Bartels N, Demmers J, Dekker M, Poot RA, Mermoud JE (2018) The CUE1 domain of the SNF2-like chromatin remodeler SMARCD1 mediates its association with KRAB-

- associated protein 1 (KAP1) and KAP1 target genes. *J Biol Chem* 293: 2711–2724
- Doetschman TC, Eistetter H, Katz M, Schmidt W, Kemler R (1985) The in vitro development of blastocyst-derived embryonic stem cell lines: formation of visceral yolk sac, blood islands and myocardium. *J Embryol Exp Morphol* 87: 27–45
- Dossin F, Pinheiro I, Zyllicz JJ, Roensch J, Collombet S, Le Saux A, Chelmicki T, Attia M, Kapoor V, Zhan Y et al (2020) SPEN integrates transcriptional and epigenetic control of X-inactivation. *Nature* 578: 455–460
- Engreitz JM, Pandya-Jones A, McDonel P, Shishkin A, Sirokman K, Surka C, Kadri S, Xing J, Goren A, Lander ES et al (2013) The Xist lncRNA exploits three-dimensional genome architecture to spread across the X chromosome. *Science* 341: 1237973
- Foti R, Gnan S, Cornacchia D, Dileep V, Bulut-Karslioglu A, Diehl S, Bunes A, Klein FA, Huber W, Johnstone E et al (2016) Nuclear architecture organized by Rif1 underpins the replication-timing program. *Mol Cell* 61: 260–273
- Furlan G, Gutierrez Hernandez N, Huret C, Galupa R, van Bommel JG, Romito A, Heard E, Morey C, Rougeulle C (2018) The Ftx noncoding locus controls X chromosome inactivation independently of its RNA products. *Mol Cell* 70: 462–472.e8
- Garzon J, Ursich S, Lopes M, Hiraga SI, Donaldson AD (2019) Human RIF1-protein phosphatase 1 prevents degradation and breakage of nascent DNA on replication stalling. *Cell Rep* 27: 2558–2566.e4
- Gnan S, Flyamer IM, Klein KN, Castelli E, Rapp A, Maiser A, Chen N, Weber P, Enervald E, Cardoso MC et al (2021) Nuclear organisation and replication timing are coupled through RIF1-PP1 interaction. *Nat Commun* 12: 2910
- Gontan C, Achame EM, Demmers J, Barakat TS, Rentmeester E, van Ijcken W, Grootegoed JA, Gribnau J (2012) RNF12 initiates X-chromosome inactivation by targeting REX1 for degradation. *Nature* 485: 386–390
- Hayano M, Kanoh Y, Matsumoto S, Renard-Guillet C, Shirahige K, Masai H (2012) Rif1 is a global regulator of timing of replication origin firing in fission yeast. *Genes Dev* 26: 137–150
- Hiraga S, Alvino GM, Chang F, Lian HY, Sridhar A, Kubota T, Brewer BJ, Weinreich M, Raghuraman MK, Donaldson AD (2014) Rif1 controls DNA replication by directing Protein Phosphatase 1 to reverse Cdc7-mediated phosphorylation of the MCM complex. *Genes Dev* 28: 372–383
- Hiraga SI, Ly T, Garzon J, Horejsi Z, Ohkubo YN, Endo A, Obuse C, Boulton SJ, Lamond AI, Donaldson AD (2017) Human RIF1 and protein phosphatase 1 stimulate DNA replication origin licensing but suppress origin activation. *EMBO Rep* 18: 403–419
- Johnston CM, Nesterova TB, Formstone EJ, Newall AE, Duthie SM, Sheardown SA, Brockdorff N (1998) Developmentally regulated Xist promoter switch mediates initiation of X inactivation. *Cell* 94: 809–817
- Jonkers I, Barakat TS, Achame EM, Monkhorst K, Kenter A, Rentmeester E, Grosveld F, Grootegoed JA, Gribnau J (2009) RNF12 is an X-Encoded dose-dependent activator of X chromosome inactivation. *Cell* 139: 999–1011
- Kucera KS, Reddy TE, Pauli F, Gertz J, Logan JE, Myers RM, Willard HF (2011) Allele-specific distribution of RNA polymerase II on female X chromosomes. *Hum Mol Genet* 20: 3964–3973
- Lee JT (2002) Homozygous Tsix mutant mice reveal a sex-ratio distortion and revert to random X-inactivation. *Nat Genet* 32: 195–200
- Lee JT (2005) Regulation of X-chromosome counting by Tsix and Xite sequences. *Science* 309: 768–771
- Lee JT, Davidow LS, Warshawsky D (1999) Tsix, a gene antisense to Xist at the X-inactivation centre. *Nat Genet* 21: 400–404
- Lee JT, Lu N (1999) Targeted mutagenesis of Tsix leads to nonrandom X inactivation. *Cell* 99: 47–57
- Li P, Wang L, Bennett BD, Wang J, Li J, Qin Y, Takaku M, Wade PA, Wong J, Hu G (2017) Rif1 promotes a repressive chromatin state to safeguard against endogenous retrovirus activation. *Nucleic Acids Res* 45: 12723–12738
- Luikenuis S, Wutz A, Jaenisch R (2001) Antisense transcription through the Xist locus mediates Tsix function in embryonic stem cells. *Mol Cell Biol* 21: 8512–8520
- Lyon MF (1961) Gene action in the X-chromosome of the mouse (*Mus musculus* L.). *Nature* 190: 372–373
- Makhlof M, Ouimette JF, Oldfield A, Navarro P, Neuillet D, Rougeulle C (2014) A prominent and conserved role for YY1 in Xist transcriptional activation. *Nat Commun* 5: 4878
- Maksakova IA, Thompson PJ, Goyal P, Jones SJ, Singh PB, Karimi MM, Lorincz MC (2013) Distinct roles of KAP1, HP1 and G9a/GLP in silencing of the two-cell-specific retrotransposon MERVL in mouse ES cells. *Epigenetics Chromatin* 6: 15
- Marahrens Y, Panning B, Dausman J, Strauss W, Jaenisch R (1997) Xist-deficient mice are defective in dosage compensation but not spermatogenesis. *Genes Dev* 11: 156–166
- Mattarocci S, Shyian M, Lemmens L, Damay P, Altintas DM, Shi T, Bartholomew CR, Thoma NH, Hardy CF, Shore D (2014) Rif1 controls DNA replication timing in yeast through the PP1 phosphatase Glc7. *Cell Rep* 7: 62–69
- McHugh CA, Chen CK, Chow A, Surka CF, Tran C, McDonel P, Pandya-Jones A, Blanco M, Burghard C, Moradian A et al (2015) The Xist lncRNA interacts directly with SHARP to silence transcription through HDAC3. *Nature* 521: 232–236
- McNamara RP, Reeder JE, McMillan EA, Bacon CW, McCann JL, D'Orso I (2016) KAP1 recruitment of the 75K snRNP complex to promoters enables transcription elongation by RNA polymerase II. *Mol Cell* 61: 39–53
- Moindrot B, Cerase A, Coker H, Masui O, Grijzenhout A, Pintacuda G, Schermelleh L, Nesterova TB, Brockdorff N (2015) A pooled shRNA screen identifies Rbm15, Spen, and Wtap as factors required for Xist RNA-Mediated Silencing. *Cell Rep* 12: 562–572
- Monfort A, Di Minin G, Postlmayr A, Freimann R, Arieti F, Thore S, Wutz A (2015) Identification of Spen as a crucial factor for Xist function through forward genetic screening in haploid embryonic stem cells. *Cell Rep* 12: 554–561
- Monk M, Harper M (1978) X-chromosome activity in preimplantation mouse embryos from XX and XO mothers. *J Embryol Exp Morphol* 46: 53–64
- Mutzel V, Okamoto I, Dunkel I, Saitou M, Giorgetti L, Heard E, Schulz EG (2019) A symmetric toggle switch explains the onset of random X inactivation in different mammals. *Nat Struct Mol Biol* 26: 350–360
- Mutzel V, Schulz EG (2020) Dosage sensing, threshold responses, and epigenetic memory: a systems biology perspective on random X-chromosome inactivation. *BioEssays* 42: e1900163
- de Napoles M, Mermoud JE, Wakao R, Tang YA, Endoh M, Appanah R, Nesterova TB, Silva J, Otte AP, Vidal M et al (2004) Polycomb group proteins Ring1A/B link ubiquitylation of histone H2A to heritable gene silencing and X inactivation. *Dev Cell* 7: 663–676
- Navarro P, Chambers I, Karwacki-Neisius V, Chureau C, Morey C, Rougeulle C, Avner P (2008) Molecular coupling of Xist regulation and pluripotency. *Science* 321: 1693–1695
- Navarro P, Page DR, Avner P, Rougeulle C (2006) Tsix-mediated epigenetic switch of a CTCF-flanked region of the Xist promoter determines the Xist transcription program. *Genes Dev* 20: 2787–2792
- Navarro P, Pichard S, Claudio C, Avner P, Rougeulle C (2005) Tsix transcription across the Xist gene alters chromatin conformation without affecting Xist

- transcription: implications for X-chromosome inactivation. *Genes Dev* 19: 1474–1484
- Nesterova TB, Johnston CM, Appanah R, Newall AE, Godwin J, Alexiou M, Brockdorff N (2003) Skewing X chromosome choice by modulating sense transcription across the Xist locus. *Genes Dev* 17: 2177–2190
- Newall AE, Duthie S, Formstone E, Nesterova T, Alexiou M, Johnston C, Caparros ML, Brockdorff N (2001) Primary non-random X inactivation associated with disruption of Xist promoter regulation. *Hum Mol Genet* 10: 581–589
- Ogawa Y, Lee JT (2003) Xite, X-inactivation intergenic transcription elements that regulate the probability of choice. *Mol Cell* 11: 731–743
- Ohhata T, Hoki Y, Sasaki H, Sado T (2008) Crucial role of antisense transcription across the Xist promoter in Tsix-mediated Xist chromatin modification. *Development* 135: 227–235
- Peace JM, Ter-Zakarian A, Aparicio OM (2014) Rif1 regulates initiation timing of late replication origins throughout the *S. cerevisiae* genome. *PLoS One* 9: e98501
- Peck SA, Hughes KD, Victorino JF, Mosley AL (2019) Writing a wrong: coupled RNA polymerase II transcription and RNA quality control. *Wiley Interdiscip Rev RNA* 10: e1529
- Penny GD, Kay GF, Sheardown SA, Rastan S, Brockdorff N (1996) Requirement for Xist in X chromosome inactivation. *Nature* 379: 131–137
- Percharde M, Lin CJ, Yin Y, Guan J, Peixoto GA, Bulut-Karslioglu A, Biechele S, Huang B, Shen X, Ramalho-Santos M (2018) A LINE1-nucleolin partnership regulates early development and ESC identity. *Cell* 174: 391–405.e19
- Rastan S (1982) Timing of X-chromosome inactivation in postimplantation mouse embryos. *J Embryol Exp Morphol* 71: 11–24
- Sado T, Hoki Y, Sasaki H (2005) Tsix silences Xist through modification of chromatin structure. *Dev Cell* 9: 159–165
- Sado T, Wang Z, Sasaki H, Li E (2001) Regulation of imprinted X-chromosome inactivation in mice by Tsix. *Development* 128: 1275–1286
- Seller CA, O'Farrell PH (2018) Rif1 prolongs the embryonic S phase at the *Drosophila* mid-blastula transition. *PLoS Biol* 16: e2005687
- Shibata S, Lee JT (2004) Tsix transcription- versus RNA-based mechanisms in Xist repression and epigenetic choice. *Curr Biol* 14: 1747–1754
- Shin J, Wallingford MC, Gallant J, Marcho C, Jiao B, Byron M, Bossenz M, Lawrence JB, Jones SN, Mager J et al (2014) RLIM is dispensable for X-chromosome inactivation in the mouse embryonic epiblast. *Nature* 511: 86–89
- Simon MD, Pinter SF, Fang R, Sarma K, Rutenberg-Schoenberg M, Bowman SK, Kesner BA, Maier VK, Kingston RE, Lee JT (2013) High-resolution Xist binding maps reveal two-step spreading during X-chromosome inactivation. *Nature* 504: 465–469
- Sreesankar E, Bharathi V, Mishra RK, Mishra K (2015) *Drosophila* Rif1 is an essential gene and controls late developmental events by direct interaction with PP1-87B. *Sci Rep* 5: 10679
- Starmer J, Magnuson T (2009) A new model for random X chromosome inactivation. *Development* 136: 1–10
- Stavropoulos N, Lu N, Lee JT (2001) A functional role for Tsix transcription in blocking Xist RNA accumulation but not in X-chromosome choice. *Proc Natl Acad Sci USA* 98: 10232–10237
- Sukackaite R, Cornacchia D, Jensen MR, Mas PJ, Blackledge M, Enervald E, Duan G, Auchynnikava T, Kohn M, Hart DJ et al (2017) Mouse Rif1 is a regulatory subunit of protein phosphatase 1 (PP1). *Sci Rep* 7: 2119
- Sun BK, Deaton AM, Lee JT (2006) A transient heterochromatic state in Xist preempts X inactivation choice without RNA stabilization. *Mol Cell* 21: 617–628
- Takagi N, Abe K (1990) Detrimental effects of two active X chromosomes on early mouse development. *Development* 109: 189–201
- Takagi N, Sugawara O, Sasaki M (1982) Regional and temporal changes in the pattern of X-chromosome replication during the early post-implantation development of the female mouse. *Chromosoma* 85: 275–286
- Tanaka H, Muto A, Shima H, Katoh Y, Sax N, Tajima S, Brydun A, Ikura T, Yoshizawa N, Masai H et al (2016) Epigenetic regulation of the Blimp-1 Gene (Prdm1) in B cells involves Bach2 and histone deacetylase 3. *J Biol Chem* 291: 6316–6330
- Tan-Wong SM, French JD, Proudfoot NJ, Brown MA (2008) Dynamic interactions between the promoter and terminator regions of the mammalian *BRCA1* gene. *Proc Natl Acad Sci USA* 105: 5160–5165
- Tian D, Sun S, Lee JT (2010) The long noncoding RNA, Jpx, is a molecular switch for X chromosome inactivation. *Cell* 143: 390–403
- Toteva T, Mason B, Kanoh Y, Brogger P, Green D, Verhein-Hansen J, Masai H, Thon G (2017) Establishment of expression-state boundaries by Rif1 and Taz1 in fission yeast. *Proc Natl Acad Sci USA* 114: 1093–1098
- Trinkle-Mulcahy L, Andersen J, Lam YW, Moorhead G, Mann M, Lamond AI (2006) Repo-Man recruits PP1 gamma to chromatin and is essential for cell viability. *J Cell Biol* 172: 679–692
- Vigneau S, Augui S, Navarro P, Avner P, Clerc P (2006) An essential role for the DXPas34 tandem repeat and Tsix transcription in the counting process of X chromosome inactivation. *Proc Natl Acad Sci USA* 103: 7390–7395
- Wang F, McCannell KN, Boskovic A, Zhu X, Shin J, Yu J, Gallant J, Byron M, Lawrence JB, Zhu LJ et al (2017) Rlim-dependent and -independent pathways for X chromosome inactivation in female ESCs. *Cell Rep* 21: 3691–3699
- Wutz A, Jaenisch R (2000) A shift from reversible to irreversible X inactivation is triggered during ES cell differentiation. *Mol Cell* 5: 695–705
- Wutz A, Rasmussen TP, Jaenisch R (2002) Chromosomal silencing and localization are mediated by different domains of Xist RNA. *Nat Genet* 30: 167–174
- Yamazaki S, Ishii A, Kanoh Y, Oda M, Nishito Y, Masai H (2012) Rif1 regulates the replication timing domains on the human genome. *EMBO J* 31: 3667–3677
- Yang L, Kirby JE, Sunwoo H, Lee JT (2016) Female mice lacking Xist RNA show partial dosage compensation and survive to term. *Genes Dev* 30: 1747–1760
- Zhao J, Sun BK, Erwin JA, Song JJ, Lee JT (2008) Polycomb proteins targeted by a short repeat RNA to the mouse X chromosome. *Science* 322: 750–756
- Zofall M, Smith DR, Mizuguchi T, Dhakshnamoorthy J, Grewal SIS (2016) Taz1-shelterin promotes facultative heterochromatin assembly at chromosome-internal sites containing late replication origins. *Mol Cell* 62: 862–874