

## Ion and Electron Momentum Distributions from Single and Double Ionization of Helium Induced by Compton Scattering

M. Kircher<sup>1,\*</sup>, F. Trinter<sup>1,2</sup>, S. Grundmann,<sup>1</sup> G. Kastirke,<sup>1</sup> M. Weller,<sup>1</sup> I. Vela-Perez,<sup>1</sup> A. Khan<sup>1,†</sup>, C. Janke,<sup>1</sup> M. Waitz,<sup>1</sup>  
S. Zeller,<sup>1</sup> T. Mletzko,<sup>1</sup> D. Kirchner<sup>1</sup>, V. Honkimäki<sup>3</sup>, S. Houamer<sup>4</sup>, O. Chuluunbaatar,<sup>5,6</sup> Yu. V. Popov<sup>7,5</sup>  
I. P. Volobuev<sup>7</sup>, M. S. Schöffler,<sup>1</sup> L. Ph. H. Schmidt,<sup>1</sup> T. Jahnke,<sup>1,8</sup> and R. Dörner<sup>1,‡</sup>

<sup>1</sup>*Institut für Kernphysik, Goethe-Universität, Max-von-Laue-Straße 1, D-60438 Frankfurt, Germany*

<sup>2</sup>*Molecular Physics, Fritz-Haber-Institut der Max-Planck-Gesellschaft, Faradayweg 4-6, D-14195 Berlin, Germany*

<sup>3</sup>*ESRF, 6 Rue Jules Horowitz, BP 220, 38043 Grenoble Cedex 9, France*

<sup>4</sup>*LPQSD, Department of Physics, Faculty of Science, University Sétif-1, 19000 Sétif, Algeria*

<sup>5</sup>*Joint Institute for Nuclear Research, Dubna, Moscow Region 141980, Russia*

<sup>6</sup>*Institute of Mathematics and Digital Technology, Mongolian Academy of Sciences, 13330 Ulaanbaatar, Mongolia*

<sup>7</sup>*Skobel'syn Institute of Nuclear Physics, Lomonosov Moscow State University, Moscow 119991, Russia*

<sup>8</sup>*European XFEL, Holzkoppel 4, 22869 Schenefeld, Germany*

 (Received 20 October 2021; revised 14 December 2021; accepted 6 January 2022; published 31 January 2022)

We present the momentum distributions of the nucleus and of the electrons from double ionization of the helium atom by Compton scattering of photons with  $h\nu = 40$  keV. We find that the doubly charged ion momentum distribution is very close to the Compton profile of the nucleus in the ground state of the helium atom, and the momentum distribution of the singly charged ion to give a precise image of the electron Compton profile. To reproduce these results, nonrelativistic calculations require the use of highly correlated initial- and final-state wave functions.

DOI: [10.1103/PhysRevLett.128.053001](https://doi.org/10.1103/PhysRevLett.128.053001)

Compton scattering is one of the fundamental interaction processes of light with matter. In his seminal paper reporting the discovery of the process, A. Compton described it as a binary collision between a photon and one single quasifree electron at rest. Soon after, initial momenta of the kicked electron were included into the description [1] leading to what is known today as the impulse approximation. This has established Compton scattering as a tool for studying momentum distributions of single active electrons in bound states, often referred to as Compton profiles (see [2] for a review). In this quasifree electron approximation, the kinematics are easily derived from momentum and energy conservation, and the cross section is given by the Klein-Nishina equation [3]. If the binding energy of the electron plays a role, Compton scattering becomes more intriguing. Comprehensive experiments for more such general conditions are scarce even today due to the very small cross section. Coincidence experiments on single electron processes have been reported recently in Ref. [4].

The next frontier for studies of Compton scattering are situations where the single-active-electron approximation fails, and more electrons are actively involved. These events are mediated by electron-electron interaction and, therefore, have been discussed as an experimental approach for exploring correlated bound states. Such two-electron Compton events are usually dissected in two steps: the Compton event occurring at only one electron which is kicked by the photon and the second electron is ionized by

electron-electron interaction. Depending on the order of these interactions, the process is referred to as knockoff or shakeoff [5–7]. For knockoff, the Compton event occurs first, and the Compton electron is pictured as “kicking” the other electron on its way out, transferring part of its energy and momentum. For shakeoff, the electron-electron interaction is part of the ground state, and the sudden removal of one electron leads to a shakeoff of the second one to the continuum. These correlation-driven processes also occur for photoabsorption, electron, or ion impact (see, e.g., [8] and references therein). The probabilities and, even more so, the differential cross sections for these processes, however, are very different in all cases. The main reasons for the differences between the ionization schemes are angular-momentum and parity selection rules, and the fact that different parts of the bound-state wave function are affected by different types of interactions. Angular-momentum selection rules, which are strict for single-photon absorption, strongly shape the angular distribution [9], masking fingerprints of electron correlation in the final-state momenta of the emitted electrons. Furthermore, single-photon absorption at high energies is selective to high-momentum components in the initial state, thus, probes only a very specific component of the wave function. In these respects, Compton scattering clearly stands out, as at high energies, the Compton scattering probability does not depend on the initial position or momentum of the electron while it is bound. Also, there

are no angular-momentum or parity selection rules for Compton scattering. Despite this fundamental nature, however, ejection of two electrons by Compton scattering has been experimentally studied only on the level of total cross sections [10,11]. There, it has been established that the ratio of double to single ionization of helium by Compton scattering approaches  $R_c^\infty = 0.86\%$  for high photon energies, as compared to  $R_\gamma^\infty = 1.66\%$  for photoabsorption [12] and  $R_{\text{charged}}^\infty = 0.26\%$  for charged particle impact [8]. This asymptotic value is, however, approached very slowly with increasing energy for Compton scattering [13].

In the present Letter, we present the first differential study of two-electron Compton scattering and compare its momentum balance with the case of single ionization. We choose helium double ionization as the cleanest possible process of this kind. For single ionization, we find that the left-behind ion shows a very clear fingerprint of the single-electron bound-state momentum profile including its modification by initial-state correlation, and, for Compton-scattering-driven double ionization, the continuum momentum distribution of the ion mimics the momentum distribution of the nucleus in the helium bound state.

In our experiment, we induce helium single or double ionization with an x-ray photon of an energy  $h\nu = 40$  keV

$$h\nu + \text{He} \rightarrow h\nu' + \text{He}^{1+} + e^-, \quad (1)$$

$$h\nu + \text{He} \rightarrow h\nu' + \text{He}^{2+} + e_1^- + e_2^-. \quad (2)$$

The momenta of the respective particles of the reactions (1) and (2) are given by

$$\vec{k}_\gamma + \vec{p}_{\text{He}} = \vec{k}_{\gamma'} + \vec{p}_{\text{He}^{1+}} + \vec{p}_e, \quad (3)$$

$$\vec{k}_\gamma + \vec{p}_{\text{He}} = \vec{k}_{\gamma'} + \vec{p}_{\text{He}^{2+}} + \vec{p}_{e1} + \vec{p}_{e2}, \quad (4)$$

respectively, and are related by momentum conservation. Single and double ionization are induced by the momentum transfer  $\vec{Q} = \vec{k}_\gamma - \vec{k}_{\gamma'}$ . In our experiment, we detect the ion charge state and the 3D momentum  $\vec{p}_{\text{He}^{q+}}$  of the ion and, in case of double ionization, the 3D momentum  $\vec{p}_{e2}$  of one of the two electrons. We detect only electrons with a momentum magnitude  $p_{e2} < 1.1$  a.u. Since, for single ionization, the majority of electrons have momentum magnitudes larger than 1.1 a.u., we do not detect the electron for single ionization. For double ionization, we only detect slow electrons. We do not detect the scattered photon. Because of technical limitations, the ion-momentum magnitude is restricted to a maximum of 1.8 a.u.

The cross section for ionization of helium by Compton scattering at 40 keV photon energy is only on the order of  $10^{-24}$  cm<sup>2</sup> [14]. The cross section for double ionization is less than 1% of this. Therefore, performing such an experiment in the gas phase requires the combination of highly efficient detection methods and high-intensity light

sources. The experiment reported here was performed at beam line ID31 of the European Synchrotron Radiation Facility (ESRF) in Grenoble, France, using a cold target recoil ion momentum spectroscopy (COLTRIMS) reaction microscope [15]. A supersonic helium gas jet was crossed with linearly polarized synchrotron light at right angle within a COLTRIMS spectrometer. A pinhole monochromator [16] was used to select the photon energy of  $h\nu = 40$  keV. An argon filter unit after the undulator removes low-energy photons from the beam. The overlap between gas jet and photon beam defines a localized reaction region of approximately  $0.4 \times 0.1 \times 1.0$  mm<sup>3</sup>. The synchrotron machine operated with 16 electron bunches in the storage ring at 5.68 MHz bunch rate, with a photon flux of  $8.4 \times 10^{14}$  photons/s at  $\Delta E/E = 1.1\%$ . Electric (6.5 V/cm) and magnetic (6.4 Gs) fields within the spectrometer guide the charged reaction fragments onto two position- and time-sensitive microchannel plate detectors with delay-line anodes [17]. The electron side of the spectrometer had a total length of 31.2 cm, divided in acceleration and drift regions with a length ratio of 1:2 (time-of-flight focusing). The ion side had a total length of 146 cm and included an electrostatic lens to compensate for the finite size of the reaction region [18]. We utilized He<sup>2+</sup> ion beams of 20 to 50 keV energy to calibrate our experiment. An electron capture process  $\text{He}^{2+} + \text{He} \rightarrow \text{He}^{1+} + \text{He}^{1+}$  was used to calibrate our ion detector. Electrons with well defined energies produced by the autoionization channel of the reaction  $\text{He}^{2+} + \text{Ne} \rightarrow \text{He}^{2+} + \text{Ne}^{1+} + e^-$  have been measured to calibrate the electron detector and the electric and magnetic fields of the spectrometer (see [15] for the kinematics of ion collisions). The uncertainty in the He<sup>2+</sup> projectile velocity yields a systematic uncertainty of our measured momenta of about 1.5%.

We compare our results for double ionization with non-relativistic calculations using the  $A^2$  approximation. Within the  $A^2$  approximation, only the quadratic term of the vector potential  $\vec{A}$  of the full interaction Hamiltonian describing the interaction between an electron and a photon is kept. The transition matrix element  $M = M(\vec{p}_{e1}, \vec{p}_{e2}; \vec{Q})$  is given by

$$M(\vec{p}_{e1}, \vec{p}_{e2}; \vec{Q}) = (\vec{\epsilon} \cdot \vec{\epsilon}') \langle \Phi_f | e^{i\vec{Q} \cdot \vec{r}_1} + e^{i\vec{Q} \cdot \vec{r}_2} | \Phi_i \rangle. \quad (5)$$

$\vec{r}_1$  and  $\vec{r}_2$  are the positions of the electrons, and  $\vec{\epsilon}$  and  $\vec{\epsilon}'$  are the polarization vectors of the incoming and outgoing photons, respectively.  $\Phi_{f,i}$  denotes the final- and initial-state wave function. A more detailed description of the theoretical model can be found in Refs. [19,20].

The  $A^2$  approximation for single ionization describes ionization by Compton scattering as the product of scattering of the photon at a free electron, described by the Klein-Nishina cross section leading to momentum transfer  $\vec{Q}$ , times the overlap of the bound-state wave function shifted

in momentum space by  $Q$ , with the continuum. Analogously, Eq. (5) describes double ionization in which electron 1 and electron 2 are kicked coherently with momentum transfer  $Q$ , and the overlap of the momentum-shifted two-electron ground state with the two-electron continuum is evaluated.

First, we discuss the result for the well understood case of single ionization. Figure 1 shows the momentum distribution of the  $\text{He}^{1+}$  ions. The lines show the predicted bound-state momentum distribution of one electron for three different helium ground-state wave functions. The solid blue line corresponds to a highly correlated trial wave function (CF) [21] for which the calculated binding energy equals the true value to seven significant digits, the dash-dotted orange line corresponds to a single-parameter Hylleraas wave function (Hy) [22] for  $Z = 27/16$ , resulting in a binding energy of 2.85 a.u. (as compared to the true value of 2.903 724 a.u.), and the dotted magenta line to the most simple configuration-interaction wave function (SPM, first discussed by Silverman, Platas, and Matsen [23]) which includes only a small amount of  $(2p)^2$  with a binding energy of 2.895 227 8 a.u. The experimental  $\text{He}^{1+}$  ion momentum distribution is within our calibration accuracy in excellent agreement with the electron momentum distribution from the best wave function (that is, CF, the wave function that includes the highest degree of correlations). We note that the horizontal error bars in the histogram do not reflect statistical errors, but give the systematical uncertainty of our momentum calibration of 1.5%, thus, showing the effect of an overall stretch or compression of the horizontal axis. The close match of the ion momentum distribution with the bound-electron Compton profile confirms the validity of the impulse




FIG. 1. Momentum distribution of  $\text{He}^{1+}$  ions produced by Compton scattering of photons with  $h\nu = 40$  keV. The experimental yield is normalized to the integral of the solid blue line. The horizontal error bars of the data points give the systematic accuracy of our momentum calibration. The vertical error bars are the standard statistical error. The lines give the bound-state electron momentum distribution (Compton profile) for different helium ground-state wave functions, namely Hylleraas (Hy), correlated trial (CF), and configuration-interaction (SPM) wave functions. See text for a discussion thereof.

approximation (compare [24] for theory). Compton scattering transfers the momentum  $\vec{Q}$  to the electron. If the corresponding energy  $Q^2/(2m_e)$  is large compared to the binding energy, the ionization probability becomes independent of the bound momentum of the electron (compare [4] for the other extreme), and the Coulomb potential can be neglected for the kinematics. In this case, as described by the impulse approximation, for a single active electron the ion momentum distribution is the exact mirror image of the bound-electron momentum distribution. The present experimental conditions come close to this ideal situation. According to the Klein-Nishina cross section, at 40 keV photon energy, only 0.7% of all Compton events correspond to an energy transfer below the helium binding energy of 24.6 eV. Furthermore, two-electron effects play only a minor role for the data in Fig. 1, since the  $\text{He}^{1+}$  momentum distribution contains all ground- and excited-state ions, and only a very small fraction of about 1% of the Compton events are not included as they lead to double ionization.

In the remainder of this Letter, we discuss the most interesting channel of double ionization. The corresponding momentum distribution of the  $\text{He}^{2+}$  nucleus [Fig. 2(a)] is much broader than the  $\text{He}^{1+}$  momentum distribution. As Fig. 2(a) shows, the measured distribution matches the momentum distribution (Compton profile) of the nucleus (solid blue line) in the bound helium atom prior to the photon impact. This indicates the validity of a kind of sudden approximation for the electron-pair removal. For this, two ingredients are necessary: (i) two-electron Compton scattering needs to remove both electrons from the initial state without altering the nucleus momentum, as it is assumed in the sudden approximation for one electron; and (ii) the two-electron removal must provide an unbiased sampling of the full wave function without selecting specific regions in momentum space. Ingredient (i) suggests that, in the Feynman diagrams shown in Fig. 2(c), the photon-electron vertex, as well as the electron-electron vertex, describe interactions to which the nucleus is a spectator only and is not affected in its momentum. Note that the inclusion of the Feynman diagrams is for visualization of the process only, they are not the basis of our theoretical calculations. The passive role of the nucleus in the electron-electron interaction is also underlined by the similarity between nuclear momentum distributions from the different quality bound states shown by the lines in Fig. 2(a). Despite the significantly different amount of electron-electron correlations, which lead to the differences in the single-electron momentum distributions as shown in Fig. 1, the nuclear Compton profile in Fig. 2(a) is very similar for all the wave functions, showing that the electron-electron interaction is quite decoupled from the nucleus. Ingredient (ii)—the unbiased sampling of the initial state—is a finding which is even more surprising in the light of previous literature. For example, it is known that the shakeoff




FIG. 2. Momentum distribution of  $\text{He}^{2+}$  ions produced by Compton scattering of photons at  $h\nu = 40$  keV. Panels (a) and (b) show the same experimental data points. The data points are normalized to the integral of the solid blue line and differ in each panel, respectively. The horizontal error bars correspond to the systematic accuracy of our momentum calibration. The vertical error bars are the standard statistical error. (a) Comparison of the nuclear Compton profile (the momentum distribution of the nucleus in bound helium). The lines show the different initial ground-state wave functions. (b) Comparison of our data with the calculated singly differential cross section (SDCS) for different sets of initial ground-state (CF, Hy, SPM) and final-state (BBK, CW, PW) wave functions (see text for a comprehensive explanation of all lines). (c) Feynman diagrams describing double ionization by Compton scattering (adapted from [28]). In (c), two Feynman diagrams with permutations of lines  $\vec{p}_{e1}$  and  $\vec{p}_{e2}$  were omitted.

probability depends on the initial-state momentum of the electron before its removal from the two-electron ground state [25]. This gives rise to the different high-energy asymptotes of ratios of double to single ionization for Compton scattering and photoabsorption [26,27] and their relation to charged-particle impact.

Figure 2(b) compares the measured momentum distributions with our nonrelativistic calculation. We have evaluated the matrix element in Eq. (5) with four sets of

initial and final states. For the initial state, we have used wave functions with different degrees of correlation, as shown in Figs. 1 and 2(a). For the final state, we have used the fully correlated Brauner-Briggs-Klar (BBK) wave function [29] which accounts for the electron-nucleus and the electron-electron interaction on equal footing [30]. For calculations of double ionization by photoabsorption [31] and for  $(e,2e)$  processes [29], this fully correlated final-state wave function has also been shown to yield extremely good results for very correlation-sensitive observables, such as the distribution of the mutual angle between the electrons. Our “best” calculation using the highly correlated initial state and the BBK wave function for the final state [solid blue line in Fig. 2(b)] yields extremely good agreement with our experimental data. All the other calculations (labeled CF + PW, Hy + CW, SPM + CW in the figure legend), where we have used final states which do not account for electron-electron repulsion—namely, a Coulomb wave with  $Z = 2$  for electron 2 and either a Coulomb wave with  $Z = 1$  (CW) or a plane wave (PW) for electron 1—are off. In terms of the Feynman diagrams [Fig. 2(c)], using final states without an electron-electron interaction term corresponds to neglecting the knockoff process. This underlines that at the present photon energy, the shakeoff limit is not yet reached.

To further elucidate the role of electron correlations on the observables, now, we inspect the momentum and angular distribution of the emitted electrons (Fig. 3). Comparison with the electron momentum distribution in the ground state [see Fig. 1, shown in Fig. 3(a) as dash-dot-dotted black line] shows that the continuum electrons have significantly smaller momenta. This is expected as, e.g., for the shakeoff as well as for the knockoff mechanism, a small energy transfer to the secondary electron is favored. The experimental electron-momentum distribution [Fig. 3(a)] is very well described by our calculation using the fully correlated initial and final states. When we remove the knockoff process by using a plane wave for the final state while keeping the fully correlated initial state (dashed red line labeled CF + PW) we find large discrepancies at higher electron momenta, which is in agreement with the findings in Ref. [7]. In Fig. 3(b), we find the electron angular distribution to be almost isotropic with a tendency toward forward angles. We note that this angular distribution is integrated over all electron momenta from 0.1 to 1.1 a.u., where the lower bound is due to our experimental momentum resolution. For photon scattering angles below  $15^\circ$ , corresponding to 3% of the Klein-Nishina cross section, the energy transfer  $Q^2/(2m_e)$  is below 111 eV [which is the sum of the double-ionization threshold of 79 eV and continuum energies of  $16 \text{ eV} = (1.1 \text{ a.u.})^2/(2m_e)$  per electron]. For those cases, the Compton scattered electron and the secondary electron are indistinguishable, and both contribute to the events in Figs. 3(a) and 3(b). In these cases,  $\vec{Q}$  is almost perpendicular to the photon




FIG. 3. Double ionization by Compton scattering of photons at  $h\nu = 40$  keV. Respectively, the single differential cross sections for electron momenta (a) and the emission angle  $\theta_e$  between the electron and incident photon beam direction (b) are shown. The data are integrated over 0.1–1.1 a.u. electron momentum and normalized to the integral of the solid blue line and differ in each panel, respectively. The error bars are the standard statistical error. Colored lines are calculations using different initial and final states [same as in Fig. 2(b)]. The dash-dot-dotted black line is the momentum distribution of the bound electron.

propagation direction. The peak visible at  $\cos(\vartheta) \simeq 0.1$  in the calculation using a plane wave for the faster electron (CF + PW) is due to these low-momentum-transfer double-ionization events. In reality, the Coulomb interaction with the nucleus and, even more, the electron-electron interaction redistribute these electrons in angle. This is in accordance with, e.g., the findings from  $(\gamma, 2e)$  experiments where, at low energies, the electrons are not directed strongly along the polarization axis, and the electrons show a  $\beta$  parameter around zero [32,33].

In conclusion, we have presented a differential measurement of a one-electron and a two-electron Compton-scattering process occurring at a free atom in the gas phase. Our results for one-electron Compton scattering show that single-electron momentum distributions can be measured with a precision that allows us to identify the modification of these distributions due to electron correlations in the bound state. Such precision measurements of Compton profiles can be extended to small and medium-sized molecules. In this case, our experimental technique will allow us to measure ionic molecular fragments in coincidence with the Compton electron. From these fragment

ions, the orientation of the molecule in the laboratory frame, as well as its dissociative state, can be inferred giving access to momentum profiles of molecular orbitals in the molecular frame of reference. We envision chiral molecules to be a particularly intriguing species for such studies [34]. Our second finding on double ionization by Compton scattering shows that the sum momentum of the ejected electrons is preserved upon ejection of the two electrons, and that this quantity even remains unperturbed by the actual double-ionization mechanisms (i.e., shakeoff and knockoff). For bound states of atoms and small molecules, this is a novel observable, as the momenta of such electron pairs have not been accessible by other means today. While, in the present study, we have used the measured ion momentum to access the electron sum momentum, this can also be done in the future for fragmenting molecular systems by detection of both electrons. Interesting candidates for such studies of electron-pair momenta in molecular systems are aromatic rings for which Cooper-pair-like pairing mechanisms have been speculated to be a possible reason for the anomalous stabilization energy associated with aromaticity [35].

This work is supported by DFG (Deutsche Forschungsgemeinschaft). We thank H. Isern, J. Drnec, and F. Russello from beam line ID31 at ESRF for excellent support during the beam time. The calculations were performed on the basis of the heterogeneous computing platform HybriLIT on supercomputer “Govorun” (LIT, JINR). The work was partially supported by the Hulubei-Meshcheryakov JINR Program, the Heisenberg-Landau Program, RFBR and Ministry of Education, Culture, Science and Sports of Mongolia Grant No. 20-51-4400, Foundation of Science and Technology of Mongolia Grant No. SST 18/2018, and the RFBR Grant No. 19-02-00014a. S.H. thanks the DGRSDT-Algeria Foundation for support. A. K. acknowledges support by the Alexander von Humboldt Foundation.

\*kircher@atom.uni-frankfurt.de

†Present address: Institut für Ionenphysik und Angewandte Physik, Universität Innsbruck, Technikerstraße 25/3, A-6020 Innsbruck, Austria.

‡doerner@atom.uni-frankfurt.de

- [1] J. W. M. Du Mond, *Phys. Rev.* **33**, 643 (1929).
- [2] M. J. Cooper, *Rep. Prog. Phys.* **48**, 415 (1985).
- [3] O. Klein and Y. Nishina, *Z. Phys.* **52**, 853 (1929).
- [4] M. Kircher, F. Trinter, S. Grundmann, I. Vela-Perez, S. Brennecke, N. Eicke, J. Rist, S. Eckart, S. Houamer, O. Chuluunbaatar *et al.*, *Nat. Phys.* **16**, 756 (2020).
- [5] J. H. McGuire, *Electron Correlation Dynamics in Atomic Collisions* (Cambridge University Press, Cambridge, England, 1997).
- [6] J. A. R. Samson, *Phys. Rev. Lett.* **65**, 2861 (1990).
- [7] A. Knapp, A. Kheifets, I. Bray, Th. Weber, A. L. Landers, S. Schössler, T. Jahnke, J. Nickles, S. Kammer, O. Jagutzki, L. Ph. H. Schmidt, T. Osipov, J. Rösch, M. H. Prior,

- H. Schmidt-Böcking, C.L. Cocke, and R. Dörner, *Phys. Rev. Lett.* **89**, 033004 (2002).
- [8] J.H. McGuire, N. Berrah, R.J. Barlett, J.A.R. Samson, J.A. Tanis, C.L. Cocke, and A.S. Schlachter, *J. Phys. B* **28**, 913 (1995).
- [9] F. Maulbetsch and J.S. Briggs, *J. Phys. B* **28**, 551 (1995).
- [10] L. Spielberger, O. Jagutzki, B. Krässig, U. Meyer, Kh. Khayyat, V. Mergel, Th. Tschentscher, Th. Buslaps, H. Bräuning, R. Dörner, T. Vogt, M. Achler, J. Ullrich, D.S. Gemmell, and H. Schmidt-Böcking, *Phys. Rev. Lett.* **76**, 4685 (1996).
- [11] B. Krässig, R.W. Dunford, D.S. Gemmell, S. Hasegawa, E.P. Kanter, H. Schmidt-Böcking, W. Schmitt, S.H. Southworth, Th. Weber, and L. Young, *Phys. Rev. Lett.* **83**, 53 (1999).
- [12] L.R. Andersson and J. Burgdörfer, *Phys. Rev. Lett.* **71**, 50 (1993).
- [13] L. Spielberger, H. Bräuning, A. Muthig, J.Z. Tang, J. Wang, Y. Qiu, R. Dörner, O. Jagutzki, Th. Tschentscher, V. Honkimäki, V. Mergel, M. Achler, Th. Weber, Kh. Khayyat, J. Burgdörfer, J. McGuire, and H. Schmidt-Böcking, *Phys. Rev. A* **59**, 371 (1999).
- [14] J.A.R. Samson, Z.X. He, R.J. Bartlett, and M. Sagurton, *Phys. Rev. Lett.* **72**, 3329 (1994).
- [15] J. Ullrich, R. Moshhammer, A. Dorn, R. Dörner, L. Ph. H. Schmidt, and H. Schmidt-Böcking, *Rep. Prog. Phys.* **66**, 1463 (2003).
- [16] G.B.M. Vaughan, J.P. Wright, A. Bytchkov, M. Rossat, H. Gleyzolle, I. Snigireva, and A. Snigirev, *J. Synchrotron Radiat.* **18**, 125 (2011).
- [17] O. Jagutzki, A. Cerezo, A. Czasch, R. Dörner, M. Hattas, M. Huang, V. Mergel, U. Spillmann, K. Ullmann-Pfleger, T. Weber *et al.*, *IEEE Trans. Nucl. Sci.* **49**, 2477 (2002).
- [18] R. Dörner, V. Mergel, L. Spielberger, M. Achler, Kh. Khayyat, T. Vogt, H. Bräuning, O. Jagutzki, T. Weber, J. Ullrich *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. B* **124**, 225 (1997).
- [19] O. Chuluunbaatar, S. Houamer, Yu. V. Popov, I. P. Volobuev, M. Kircher, and R. Dörner, *J. Quant. Spectrosc. Radiat. Transfer* **272**, 107820 (2021).
- [20] O. Chuluunbaatar, S. Houamer, Yu. V. Popov, I. P. Volobuev, M. Kircher, and R. Dörner, *J. Quant. Spectrosc. Radiat. Transfer* **278**, 108020 (2022).
- [21] O. Chuluunbaatar, I. V. Puzynin, P.S. Vinitzky, Yu. V. Popov, K. A. Kouzakov, and C. Dal Cappello, *Phys. Rev. A* **74**, 014703 (2006).
- [22] E. A. Hylleraas, *Z. Phys.* **48**, 469 (1928).
- [23] J. N. Silverman, O. Platas, and F. A. Matsen, *J. Chem. Phys.* **32**, 1402 (1960).
- [24] Z. Kaliman and K. Pisk, *Radiat. Phys. Chem.* **71**, 633 (2004).
- [25] T. Y. Shi and C. D. Lin, *Phys. Rev. Lett.* **89**, 163202 (2002).
- [26] T. Åberg, *Phys. Rev. A* **2**, 1726 (1970).
- [27] L. Spielberger, O. Jagutzki, R. Dörner, J. Ullrich, U. Meyer, V. Mergel, M. Unverzagt, M. Damrau, T. Vogt, I. Ali, Kh. Khayyat, D. Bahr, H.G. Schmidt, R. Frahm, and H. Schmidt-Böcking, *Phys. Rev. Lett.* **74**, 4615 (1995).
- [28] M. Ya. Amusia and A. I. Mikhailov, *J. Phys. B* **28**, 1723 (1995).
- [29] M. Brauner, J. S. Briggs, and H. Klar, *J. Phys. B* **22**, 2265 (1989).
- [30] J. S. Briggs, *J. Phys. B* **49**, 170501 (2016).
- [31] J. S. Briggs and V. Schmidt, *J. Phys. B* **33**, R1 (2000).
- [32] R. Wehlitz, F. Heiser, O. Hemmers, B. Langer, A. Menzel, and U. Becker, *Phys. Rev. Lett.* **67**, 3764 (1991).
- [33] A. Knapp, M. Walter, Th. Weber, A.L. Landers, S. Schössler, T. Jahnke, M. Schöffler, J. Nickles, S. Kammer, and O. Jagutzki, *J. Phys. B* **35**, L521 (2002).
- [34] A. F. Ordóñez and O. Smirnova, *Phys. Rev. A* **99**, 043416 (2019).
- [35] R. Wehlitz, P. N. Juranić, K. Collins, B. Reilly, E. Makoutz, T. Hartman, N. Appathurai, and S. B. Whitfield, *Phys. Rev. Lett.* **109**, 193001 (2012).