



Geometry & Topology

Volume 27 (2023)

Embedded surfaces with infinite cyclic knot group

ANTHONY CONWAY

MARK POWELL



Embedded surfaces with infinite cyclic knot group

ANTHONY CONWAY

MARK POWELL

We study locally flat, compact, oriented surfaces in 4–manifolds whose exteriors have infinite cyclic fundamental group. We give algebraic topological criteria for two such surfaces, with the same genus g , to be related by an ambient homeomorphism, and further criteria that imply they are ambiently isotopic. Along the way, we provide a classification of a subset of the topological 4–manifolds with infinite cyclic fundamental group, and we apply our results to rim surgery.

57K40, 57N35

1 Introduction

We study locally flat embeddings of compact, orientable surfaces in compact, oriented, simply connected topological 4–manifolds, where the complement of the surface has infinite cyclic fundamental group. Extending the terminology for knotted spheres, we call this group the *knot group*, so we shall study knotted surfaces with knot group \mathbb{Z} , or \mathbb{Z} –surfaces.

We will present algebraic criteria for pairs of \mathbb{Z} –surfaces to be ambiently isotopic. As part of the proof we obtain an algebraic classification of a certain subset of the 4–manifolds with boundary and fundamental group \mathbb{Z} , simultaneously generalising work of Freedman and Quinn [15] on the closed case, and of Boyer [5] on simply connected 4–manifolds with nonempty boundary; see Section 1.7. We apply our results to show that in simply connected 4–manifolds, 1–twisted rim surgery on a surface with knot group \mathbb{Z} yields a topologically ambiently isotopic surface, extending results of Kim and Ruberman [31] and Juhász, Miller and Zemke [24]; see Section 1.5.

1.1 Surfaces in S^4 and D^4

We start with our main results on surfaces in S^4 and D^4 as the most important special cases, before going on to explain more general statements for surfaces in any simply connected 4–manifold.

Theorem 1.1 *Any two locally flat, embedded, closed, orientable and genus $g \neq 1, 2$ surfaces $\Sigma_0, \Sigma_1 \subseteq S^4$ with $\pi_1(S^4 \setminus \Sigma_i) \cong \mathbb{Z}$ for $i = 0, 1$ are topologically ambiently isotopic.*

Theorem 1.2 *Let K be an Alexander polynomial one knot in S^3 . Any two locally flat, embedded, compact, orientable genus $g \neq 1, 2$ surfaces $\Sigma_0, \Sigma_1 \subseteq D^4$ with boundary K and $\pi_1(D^4 \setminus \Sigma_i) \cong \mathbb{Z}$ for $i = 0, 1$ are topologically ambiently isotopic rel boundary.*

In a previous paper, we proved Theorem 1.2 in the genus-zero case [10, Theorem 1.2]. With regards to Theorem 1.1, a genus g surface $\Sigma \subseteq S^4$ is *unknotted* if it bounds a locally flat, embedded handlebody in S^4 . The *unknotting conjecture* for locally flat surfaces posits that a locally flat, embedded, closed, oriented surface $\Sigma \subseteq S^4$ is unknotted if and only if $\pi_1(S^4 \setminus \nu\Sigma) = \mathbb{Z}$. The forwards direction holds because any two embeddings of a handlebody in S^4 are ambiently isotopic, so an unknotted surface is ambiently isotopic to a standard embedding. For the reverse direction, Freedman and Quinn [15, Theorem 11.7A] proved the $g = 0$ case of 2–knots, when $\Sigma \cong S^2$, while Hillman and Kawauchi [22] claimed it for $g \geq 1$. Theorem 1.1 offers a new proof for surfaces of genus $g \geq 3$. We also give new proofs for the $g = 0$ cases, but it should be noted that this specialisation produces somewhat similar proofs to those of Freedman and Quinn and our previous work [10].

Our proof of Theorem 1.1 differs from that of Hillman and Kawauchi in [22], who do not have any genus restrictions. In particular, in a key step in the proof, one considers a closed 4–manifold M built from the two surface exteriors, and shows that M is homeomorphic to $S^1 \times S^3 \# \#_{i=1}^{2g} S^2 \times S^2$. To see this, we control the $\mathbb{Z}[\mathbb{Z}]$ –valued intersection form of M , whereas [22, Proof of Lemma 3.1] just calculates the \mathbb{Z} –valued intersection form, and appeals to separate work of Kawauchi [27], in which it was claimed that every closed, spin 4–manifold with fundamental group \mathbb{Z} splits as $S^1 \times S^3 \# X$, where X is a closed, simply connected 4–manifold. It would follow from this claim that computing the \mathbb{Z} –valued form suffices. A mistake in [27] was found by Hambleton and Teichner [21]. Although Kawauchi later updated his

theorem [28] to include the hypothesis that the \mathbb{Z} -intersection form be indefinite, which is the case for the 4-manifolds arising in [22], the consensus in the community seems to be that an independent account is also desirable.

1.2 Surfaces in simply connected 4-manifolds with boundary S^3

To fix our terminology, throughout the article, by a 4-manifold we shall always mean a compact, connected, oriented, topological 4-manifold with connected boundary. Surfaces will always be compact, connected, and orientable. We will consider locally flat, embedded, closed, oriented surfaces $\Sigma \subseteq X$ with X a closed 4-manifold, and locally flat properly embedded surfaces $\Sigma \subseteq N$, where N is a 4-manifold with boundary S^3 and $\partial\Sigma = K \subseteq S^3$ is a fixed knot. The exteriors of such surfaces will be denoted by X_Σ and N_Σ respectively.

Next we formulate our most general statement, for the nonempty boundary case, pairs of surfaces $\Sigma_i \subseteq N$, where $\partial\Sigma_i \subseteq \partial N = S^3$ is a fixed knot. Theorem 1.2 is an immediate consequence of Theorem 1.3 below, which we state after introducing the prerequisites. One quickly deduces consequences for closed surfaces in closed 4-manifolds X , in particular Theorem 1.1, by removing an unknotted (D^4, D^2) pair from (X, Σ) , as we will explain in Section 1.3.

Given a compact, oriented 4-manifold M with $\pi_1(M) \cong \mathbb{Z}$, we write $\pi_1(M) = \mathbb{Z}$ to mean that we have chosen an isomorphism between $\pi_1(M)$ and \mathbb{Z} . In the case of the exterior X_Σ or N_Σ of an oriented surface Σ , such an identification is determined by the orientations on Σ and on X or N respectively. Set $\Lambda := \mathbb{Z}[t^{\pm 1}]$ and write $H_*(M; \Lambda)$ for the homology of the infinite cyclic cover \tilde{M} of M , considered as a Λ -module. Taking signed intersections in \tilde{M} endows the homology Λ -module $H_2(M; \Lambda)$ with a sesquilinear Hermitian intersection form

$$\lambda_M : H_2(M; \Lambda) \times H_2(M; \Lambda) \rightarrow \Lambda.$$

The adjoint of λ_M is the Λ -module homomorphism

$$\hat{\lambda}_M : H_2(M; \Lambda) \rightarrow \text{Hom}_\Lambda(H_2(M; \Lambda), \Lambda) =: H_2(M; \Lambda)^*.$$

If two 4-manifolds M_0, M_1 with infinite cyclic fundamental group are orientation-preserving homeomorphic, then their Λ -intersection forms are isometric. That is, there exists an isomorphism $F : H_2(M_0; \Lambda) \xrightarrow{\cong} H_2(M_1; \Lambda)$ such that $\hat{\lambda}_{M_0} = F^* \hat{\lambda}_{M_1} F$. We write $F : \lambda_{M_0} \cong \lambda_{M_1}$ and call F an *isometry*.

Let Y be a compact oriented 3–manifold, and let $\varphi: \pi_1(Y) \twoheadrightarrow \mathbb{Z}$ be an epimorphism. Associated with this data, there is a Λ –module $H_1(Y; \Lambda)$, called the *Alexander module*, which is the first homology group of the infinite cyclic cover associated to $\ker(\varphi)$. If this module is torsion over $\Lambda = \mathbb{Z}[t^{\pm 1}]$, then it is endowed with a sesquilinear Hermitian *Blanchfield form*

$$\text{Bl}_Y: H_1(Y; \Lambda) \times H_1(Y; \Lambda) \rightarrow \mathbb{Q}(t)/\Lambda.$$

More details on this pairing appear in Section 3.1, but we note that Bl_Y should be thought of as the analogue of the linking pairing of a \mathbb{Q} –homology sphere on the level of infinite cyclic covers.

As we recall in Sections 2 and 3, if M_0, M_1 are 4–manifolds with $\pi_1(M_i) = \mathbb{Z}$ whose boundaries ∂M_i have torsion Alexander modules, and if $\pi_1(\partial M_i) \rightarrow \pi_1(M_i)$ is surjective for $i = 0, 1$, then an isometry $F: \lambda_{M_0} \cong \lambda_{M_1}$ of the Λ –intersection forms induces an isometry of the Blanchfield forms of the boundary,

$$\partial F: (H_1(\partial M_0; \Lambda), \text{Bl}_{\partial M_0}) \xrightarrow{\cong} (H_1(\partial M_1; \Lambda), \text{Bl}_{\partial M_1}).$$

Here is the construction: via universal coefficients, Poincaré–Lefschetz duality, and the long exact sequence of a pair, the composition

$$H_2(M_i; \Lambda)^* \cong H^2(M_i; \Lambda) \cong H_2(M_i, \partial M_i; \Lambda) \xrightarrow{\delta} H_1(\partial M_i; \Lambda)$$

induces an identification $\text{coker}(\widehat{\lambda}_{M_i}) \cong H_1(\partial M_i; \Lambda)$, and the map

$$F^{-*} := (F^*)^{-1}: H_2(M_0; \Lambda)^* \rightarrow H_2(M_1; \Lambda)^*$$

induces $\partial F: \text{coker}(\widehat{\lambda}_{M_0}) \rightarrow \text{coker}(\widehat{\lambda}_{M_1})$.

Now we focus on the case where $M_i = N_{\Sigma_i}$ are surface exteriors, with N a compact, simply connected 4–manifold with boundary S^3 . The boundary ∂N_{Σ_i} is homeomorphic to

$$M_{K,g} := E_K \cup_{\partial} (\Sigma_{g,1} \times S^1),$$

where $E_K = S^3 \setminus \nu(K)$ is the knot exterior and $\Sigma_{g,1}$ denotes the (abstract) genus g surface with one boundary component. In Proposition 5.7, we show that every automorphism h of the Blanchfield pairing $\text{Bl}_{M_{K,g}}$ decomposes as $h_K \oplus h_{\Sigma}$, where h_K is an automorphism of $\text{Bl}_K := \text{Bl}_{E_K}$ and h_{Σ} is an automorphism of $\text{Bl}_{\Sigma_{g,1} \times S^1}$.

Let $f_K: E_K \rightarrow E_K$ be an orientation-preserving homeomorphism that is the identity on ∂E_K . Extend f_K via the identity on νK to an orientation-preserving self-homeomorphism of S^3 . The mapping class group of S^3 is trivial, so there is an

isotopy $\Theta(f_K): S^3 \times [0, 1] \rightarrow S^3$ between the extension and the identity, such that $\Theta(f_K)|_{S^3 \times \{0\}} = \text{Id}$ and $\Theta(f_K)|_{E_K \times \{1\}} = f_K$.

Here is the central theorem of the paper on embedded surfaces with knot group \mathbb{Z} .

Theorem 1.3 *Let X be a closed, simply connected and oriented 4–manifold, let $N = X \setminus \overset{\circ}{D}^4$ be a punctured X , and let $K \subseteq S^3 = \partial N$ be a knot. Let $\Sigma_0, \Sigma_1 \subseteq N$ be two locally flat, embedded, compact, oriented genus g surfaces with the same oriented boundary K and $\pi_1(N_{\Sigma_i}) = \mathbb{Z}$ for $i = 0, 1$. Suppose there is an isometry $F: \lambda_{N_{\Sigma_0}} \cong \lambda_{N_{\Sigma_1}}$ and write $\partial F = h_K \oplus h_\Sigma$.*

- *If h_K is induced by an orientation-preserving homeomorphism $f_K: E_K \rightarrow E_K$ that is the identity on ∂E_K , then f_K extends to an orientation-preserving homeomorphism of pairs*

$$(N, \Sigma_0) \xrightarrow{\cong} (N, \Sigma_1)$$

inducing the given isometry $F: H_2(N_{\Sigma_0}; \Lambda) \cong H_2(N_{\Sigma_1}; \Lambda)$.

- *If in addition $N = D^4$, then for any choice of isotopy $\Theta(f_K)$, the surfaces Σ_0 and Σ_1 are topologically ambiently isotopic via an ambient isotopy of D^4 extending $\Theta(f_K)$.*

In particular note that if $h_K = \text{Id}$, then we can take $f_K = \text{Id}$ and Θ_{f_K} the constant isotopy, so that the homeomorphism of pairs in the first item can be assumed to fix the boundary pointwise, and the ambient isotopy in the second item can be assumed to be rel boundary.

In general, $\lambda_{N_{\Sigma_0}}$ and $\lambda_{N_{\Sigma_1}}$ need not be isometric, even for $N = D^4$, as shown by examples due to Oba [43].

The deduction of the last item uses Alexander’s coning trick, which shows that every homeomorphism of D^4 that restricts to the identity on ∂D^4 is topologically isotopic to the identity. So in D^4 , a homeomorphism of pairs can be upgraded to a topological ambient isotopy. See Section 5.3 for details.

As explained in Section 1.5 below, we will apply Theorem 1.3 to 1–twisted rim surgery, a method which has been effective at producing exotic embeddings of surfaces. In Section 1.6, we also apply Theorem 1.3 to study Seifert surfaces that are pushed in to D^4 . Further applications can be obtained by finding classes of knots K for which every automorphism of the Blanchfield pairing is realised by a symmetry of the knot exterior E_K .

1.3 Ambient isotopy in closed 4–manifolds

For closed surfaces embedded in closed, simply connected 4–manifolds, by applying the classification of self-homeomorphisms due to Kreck [36], Perron [44] and Quinn [45], we can potentially upgrade a homeomorphism of pairs obtained from Theorem 1.3 to an ambient isotopy. The theorem (see [45] or [15, Theorem 10.1]) is that two self-homeomorphisms of a closed, simply connected 4–manifold are isotopic if and only if they induce the same self-isomorphism on second homology. An analogous classification of homeomorphisms for simply connected 4–manifolds with boundary has not yet been proven.

Theorem 1.4 *Let X be a closed, simply connected, oriented 4–manifold. Let $\Sigma_0, \Sigma_1 \subseteq X$ be two locally flat, embedded, closed, oriented genus g surfaces with $\pi_1(X_{\Sigma_i}) = \mathbb{Z}$ for $i = 0, 1$.*

- (i) *If the intersection forms $\lambda_{X_{\Sigma_0}}$ and $\lambda_{X_{\Sigma_1}}$ are isometric via an isometry F , then there is an orientation-preserving homeomorphism of pairs*

$$\Phi: (X, \Sigma_0) \xrightarrow{\cong} (X, \Sigma_1)$$

inducing the given isometry $\Phi_ = F: H_2(X_{\Sigma_0}; \Lambda) \cong H_2(X_{\Sigma_1}; \Lambda)$.*

- (ii) *The isometry F induces an isometry $F_{\mathbb{Z}}: H_2(X) \rightarrow H_2(X)$ of the standard intersection form Q_X of X . The surfaces Σ_0 and Σ_1 are topologically ambiently isotopic if and only if $F_{\mathbb{Z}} = \text{Id}$.*

With regards to (ii), Lemma 5.10 furnishes details of how the isometry $F: \lambda_{\Sigma_0} \cong \lambda_{\Sigma_1}$ induces an isometry $F_{\mathbb{Z}}$ of the standard intersection form Q_X . Theorem 1.4 also has applications to rim surgery, as we explain in Section 1.5.

Theorem 1.4(i) follows from Theorem 1.3. Here is a short outline; see Section 5.4 for details. After an isotopy, we assume Σ_0 and Σ_1 coincide on a disc D^2 . Remove an open neighbourhood of this common disc ($\mathring{D}^4, \mathring{D}^2$) from (X, Σ_i) to obtain a pair $(N, \tilde{\Sigma}_i)$, with $\partial N \cong S^3$ and $\partial \tilde{\Sigma}_i$ an unknot K . The exterior of Σ_i in X equals the exterior of $\tilde{\Sigma}_i$ in N , so the assumptions of Theorem 1.3 hold. Here, since the unknot has trivial Alexander module, $h_K = \text{Id}$ and so we take $f_K = \text{Id}$. Then Theorem 1.3 produces a homeomorphism of pairs $(N, \tilde{\Sigma}_0) \cong (N, \tilde{\Sigma}_1)$ rel boundary, which we complete with the identity on the missing 4–ball to prove (i). Deducing (ii) uses the classification of homeomorphisms from [45] and [15, Theorem 10.1] mentioned above.

Remark 1.5 It is worth contrasting Theorem 1.4 with a result of Sunukjian. Indeed, [49, Theorem 7.2] shows that closed surfaces $\Sigma_0, \Sigma_1 \subseteq X$ of the same genus with knot group \mathbb{Z} are topologically isotopic provided $b_2(X) \geq |\sigma(X)| + 2$. Thus, when X is “big enough”, restrictions on the equivariant intersection form are not needed to establish isotopy.

1.4 Deducing Theorems 1.1 and 1.2 from Theorems 1.3 and 1.4

In the closed case, with $X = S^4$, if we have an isometry F of the intersection form of $X_{\Sigma_0} = S^4_{\Sigma_0}$ and $X_{\Sigma_1} = S^4_{\Sigma_1}$ then the map $F_{\mathbb{Z}}: H_2(S^4) \rightarrow H_2(S^4)$ is automatically the identity automorphism.

For the case with nonempty boundary, with $N = D^4$, Alexander polynomial one implies that $H_1(E_K; \Lambda) = 0$, so $h_K = \text{Id}$ automatically, and we may take $f_K: E_K \rightarrow E_K$ also to be the identity.

Therefore, in both cases, it suffices to find an isometry between the intersection forms of the surface exteriors. It is an open question whether this is true in general. For genus zero, ie for discs and spheres, this is automatically the case since the Λ -coefficient second homology of the surface exterior vanishes. Therefore the genus-zero cases of Theorems 1.1 and 1.2, due to [15, Theorem 11.7A] and [10, Theorem 1.2], respectively, follow from Theorems 1.3 and 1.4.

For genus at least three, our strategy to show that the intersection forms are isometric is as follows. As explained above, the exterior of a closed surface in S^4 can be considered as the exterior of a properly embedded surface in D^4 , so we discuss only the latter case.

By [2, Theorem 5] (see also Theorem A.1), Σ_0 and Σ_1 are stably equivalent, meaning that they become isotopic after adding some number of trivial tubes. Therefore the intersection forms of $\lambda_{D^4_{\Sigma_0}}$ and $\lambda_{D^4_{\Sigma_1}}$ satisfy $\lambda_{D^4_{\Sigma_0}} \oplus \mathcal{H}_2^{\oplus n} \cong \lambda_{D^4_{\Sigma_1}} \oplus \mathcal{H}_2^{\oplus n}$ for some $n \geq 0$, where

$$\mathcal{H}_2 := \begin{pmatrix} 0 & t-1 \\ t^{-1}-1 & 0 \end{pmatrix}.$$

Then provided g is at least three we are able to leverage algebraic cancellation results for hyperbolic forms from Bass [1] (see also Crowley and Sixt [12], Hambleton and Teichner [21], Khan [29] and Magurn, van der Kallen and Vaserstein [41]) to improve such a stable isometry to an isometry. While we refer to Section 7.1 for details, we record one of the aforementioned intermediate results as it might be of independent interest.

Proposition 1.6 *Let X be a closed, simply connected, oriented 4–manifold. Let $\Sigma_0, \Sigma_1 \subseteq N = X \setminus \overset{\circ}{D}^4$ be two locally flat, properly embedded, compact, oriented genus g surfaces with boundary the same oriented knot $K \subseteq S^3$, and $\pi_1(N_{\Sigma_i}) = \mathbb{Z}$ for $i = 0, 1$. There exists an integer $n \geq 0$ and an isometry*

$$\lambda_{N_{\Sigma_0}} \oplus \mathcal{H}_2^{\oplus n} \cong \lambda_{N_{\Sigma_1}} \oplus \mathcal{H}_2^{\oplus n}.$$

Whereas Theorems 1.1 and 1.2 hold for $g \neq 1, 2$, Proposition 1.6 leads to some results for arbitrary g . Indeed, Corollary 6.6 shows that for any genus g surface $\Sigma \subseteq N = X \setminus \overset{\circ}{D}^4$ with knot group \mathbb{Z} and $\partial\Sigma$ an Alexander polynomial one knot, $\lambda_{N_\Sigma} \oplus \mathcal{H}_2^{\oplus n} \cong Q_X \oplus \mathcal{H}_2^{\oplus(g+n)}$ for some $n \geq 0$, where Q_X denotes the standard intersection form of X . The same result holds for closed surfaces $\Sigma \subseteq X$ with knot group \mathbb{Z} .

1.5 Application to rim surgery

Rim surgery is an effective way to produce smoothly inequivalent embeddings of surfaces; see Fintushel and Stern [14], Juhász, Miller and Zemke [24], Kim [30], Kim and Ruberman [31] and Mark [42]. Given a locally flat embedded closed oriented surface Σ in a closed 4–manifold X , a knot $J \subseteq S^3$ and simple closed curve $\alpha \subseteq \Sigma$, n –roll m –twist rim surgery outputs another locally flat embedded closed oriented surface $\Sigma_n^m(\alpha, J) \subseteq X$. The case where $m = 0, n = 0$ was first introduced by Fintushel and Stern [14], while m –twisted rim surgery was introduced by Kim [30], and n –roll m –twist rim surgery first appeared in Kim and Ruberman [32].

Kim and Ruberman [32, Theorem 1.3 and Proposition 3.1] showed that if X is a simply connected 4–manifold, $\pi_1(X_\Sigma) = \mathbb{Z}_d$ is a finite cyclic group and $(m, d) = 1$, then $\Sigma_n^m(\alpha, J)$ and Σ are topologically isotopic. In order to extend this to infinite cyclic fundamental groups, we consider n –roll 1–twist rim surgery, and set $\Sigma_n(\alpha, J) := \Sigma_n^1(\alpha, J)$. In this case if $\pi_1(X_\Sigma) = \mathbb{Z}$, then also $\pi_1(X_{\Sigma_n(\alpha, J)}) = \mathbb{Z}$. Our first result on rim surgery then reads as follows.

Theorem 1.7 *Let X be a closed, simply connected 4–manifold, let $\Sigma \subseteq X$ be a locally flat, embedded, orientable surface with knot group \mathbb{Z} , let $\alpha \subseteq \Sigma$ be a simple closed curve, let J be a knot, and let $n \in \mathbb{Z}$. Then the surfaces Σ and $\Sigma_n(\alpha, J)$ are topologically ambiently isotopic.*

For good fundamental groups $\pi_1(X \setminus \Sigma)$ and J a slice knot, Kim and Ruberman showed in [31, Theorem 4.5] that there is a homeomorphism of pairs $(X, \Sigma) \cong (X, \Sigma_0(\alpha, J))$, so for $\pi_1(X \setminus \Sigma) \cong \mathbb{Z}$ our result is an extension.

Rim surgery is also defined for a properly embedded surface in a 4-manifold N with boundary. Juhász, Miller and Zemke [24, Corollary 2.7] showed that if $\Sigma \subseteq N$ is a properly embedded locally flat oriented surface, and if α bounds a locally flat disc in N_Σ , then $\Sigma_0(\alpha, J)$ is topologically isotopic to Σ . In the case where Σ has knot group \mathbb{Z} , we generalise this result as follows.

Theorem 1.8 *Let X be a closed, simply connected 4-manifold, and let $N := X \setminus \mathring{D}^4$ be a punctured X . Let $\Sigma \subseteq N$ be a locally flat, properly embedded, orientable surface with knot group \mathbb{Z} , let $\alpha \subseteq \Sigma$ be a simple closed curve, let J be a knot, and let $n \in \mathbb{Z}$ be an integer. There is a rel boundary orientation-preserving homeomorphism of pairs*

$$(N, \Sigma) \xrightarrow{\cong} (N, \Sigma_n(\alpha, J)).$$

If $N = D^4$, then the surfaces Σ and $\Sigma_n(\alpha, J)$ are topologically ambiently isotopic rel boundary.

Note that in both of the previous two results, we have no restrictions on the genera of the surfaces. This is because we show that the Λ -intersection forms of $N_{\Sigma_n(\alpha, J)}$ and N_Σ coincide, allowing us to apply the first item of Theorem 1.3, for which there is no genus restriction.

For the case $N = D^4$, Theorem 1.8 means that the construction of exotic surfaces in [24] applies more generally than realised in that article, because the condition that α bounds a locally flat embedded disc in N_Σ is not needed.

1.6 Application to pushed-in Seifert surfaces

Pushing a Seifert surface for a knot K into D^4 yields a surface with knot group \mathbb{Z} . It is then intriguing to wonder whether any two Seifert surfaces of the same genus for a knot K become isotopic once pushed into D^4 ; see for instance Problem 1.20(C) of Kirby [33] and Section 6 of Livingston [40]. In Theorem 7.11, relying on Theorem 1.3, we show that this is the case for Alexander polynomial one knots.

Theorem 1.9 *If $F_0, F_1 \subseteq D^4$ are genus g pushed-in Seifert surfaces for an Alexander polynomial one knot K , then they are topologically ambiently isotopic rel boundary.*

This result provides another setting where we are able to obtain results for $g = 1, 2$. In particular, it rules out the most naive potential counter examples to the conjecture that Theorem 1.2 holds for \mathbb{Z} -surfaces of any genus.

1.7 A classification result for topological 4-manifolds

Several of the steps from the proof of Theorem 1.3 fit in a classification scheme that applies to a wider class of 4-manifolds than surface exteriors. We state the resulting theorem, which is analogous to Boyer’s classification [5] for the simply connected case.

We say that a compact, oriented, topological 4-manifold M with nonempty connected boundary has *ribbon boundary* if the inclusion induced map $\iota: \pi_1(\partial M) \rightarrow \pi_1(M)$ is surjective. For example, in simply connected 4-manifolds, (connected) surface exteriors with infinite cyclic fundamental group have ribbon boundaries. Set $\Lambda := \mathbb{Z}[t^{\pm 1}]$. If we have an identification $\pi_1(M) = \mathbb{Z}$, then we have infinite cyclic covers $\tilde{M} \rightarrow M$ and $\partial \tilde{M} \rightarrow \partial M$ and Λ -modules $H_*(M; \Lambda) = H_*(\tilde{M})$ and $H_*(\partial M; \Lambda) = H_*(\partial \tilde{M})$.

Let M_0 and M_1 be two 4-manifolds with infinite cyclic fundamental group, whose boundaries are ribbon and have Λ -torsion Alexander modules. Fix an orientation-preserving homeomorphism $f: \partial M_0 \rightarrow \partial M_1$ that intertwines the inclusion induced maps $\varphi_i: \pi_1(\partial M_i) \twoheadrightarrow \mathbb{Z}$, and an isometry $F: (H_2(M_0; \Lambda), \lambda_{M_0}) \rightarrow (H_2(M_1; \Lambda), \lambda_{M_1})$ of the Λ -valued intersection forms λ_{M_0} and λ_{M_1} . As described in Section 3.4 below, f and F induce isometries of the boundary Blanchfield forms:

$$f_*, \partial F: (H_1(\partial M_0; \Lambda), \text{Bl}_{\partial M_0}) \rightarrow (H_1(\partial M_1; \Lambda), \text{Bl}_{\partial M_1}).$$

We call (f, F) a *compatible pair* if f and F induce the same isometry. For conciseness, we write

$$\text{Homeo}_\varphi^+(\partial M_0, \partial M_1) := \{f \in \text{Homeo}^+(\partial M_0, \partial M_1) \mid \varphi_1 \circ f_* = \varphi_0\}$$

for the set of orientation-preserving homeomorphisms of the boundaries that intertwine the maps to \mathbb{Z} .

Recall that the Kirby–Siebenmann invariant $\text{ks}(M_i) \in \mathbb{Z}/2$ is the unique obstruction for the stable tangent bundle $M_i \rightarrow \text{BTOP}$ to lift to $M_i \rightarrow \text{BPL}$, or equivalently for M_i to be smoothable after adding copies of $S^2 \times S^2$; see Freedman and Quinn [15, Theorem 8.6] and Friedl, Nagel, Orson and Powell [17, Section 8]. This is relevant in the next theorem for the nonspin case; note that for 4-manifolds with fundamental group \mathbb{Z} , whether or not they are spin is determined by the intersection pairing.

Theorem 1.10 Suppose M_0 and M_1 are two compact, oriented 4–manifolds with $\pi_1(M_i) = \mathbb{Z}$ for $i = 0, 1$, whose boundaries are ribbon and have Λ –torsion Alexander modules. Let $f \in \text{Homeo}_\varphi^+(\partial M_0, \partial M_1)$ be a homeomorphism and let $F: \lambda_{M_0} \cong \lambda_{M_1}$ be an isometry. If M_0 and M_1 are not spin, assume that the Kirby–Siebenmann invariants satisfy $\text{ks}(M_0) = \text{ks}(M_1) \in \mathbb{Z}/2$. Then the following assertions are equivalent:

- (i) The pair (f, F) is compatible.
- (ii) The homeomorphism f extends to an orientation-preserving homeomorphism

$$\Phi: M_0 \xrightarrow{\cong} M_1$$

inducing the given isometry $F: H_2(M_0; \Lambda) \cong H_2(M_1; \Lambda)$.

Remark 1.11 Theorem 1.10 should be compared with both Boyer’s classification [5] of simply connected 4–manifolds with boundary and with Freedman and Quinn’s classification [15] of closed 4–manifolds with infinite cyclic fundamental group. Boyer uses a notion, similar to our compatible pairs, which he calls *morphisms*. In a nutshell, given simply connected 4–manifolds M_0, M_1 with rational homology spheres as their boundaries, Boyer [5, Theorem 0.7 and Proposition 0.8] shows that a morphism (f, F) can be extended to a homeomorphism $M_0 \rightarrow M_1$. Note that Boyer’s methods differ from ours, as he does not use the union of forms. On the other hand, Freedman and Quinn [15, Theorem 10.7A] show that closed 4–manifolds with infinite cyclic fundamental groups are classified by their Λ –intersection form and their Kirby–Siebenmann invariant; see also Stong and Wang [48] and Hambleton, Kreck and Teichner [20].

A major step in the proof of Theorem 1.10 is the following intermediate result, which might be of independent interest; a more detailed statement and a proof can be found in Theorem 3.12.

Theorem 1.12 Suppose M_0 and M_1 are two compact, oriented 4–manifolds with $\pi_1(M_i) = \mathbb{Z}$ for $i = 0, 1$, whose boundaries are ribbon and have Λ –torsion Alexander modules. Let (f, F) be a compatible pair. If M_0 and M_1 are not spin, assume that the Kirby–Siebenmann invariants satisfy $\text{ks}(M_0) = \text{ks}(M_1) \in \mathbb{Z}/2$. Then there is an orientation-preserving homeomorphism

$$M_0 \cup_f -M_1 \cong S^1 \times S^3 \# \#_{i=1}^a S^2 \times S^2 \# \#_{j=1}^b S^2 \tilde{\times} S^2$$

for some a, b with $a + b = b_2(M_0)$. If M_0 and M_1 are spin, then $b = 0$.

Remark 1.13 We compare Theorem 1.12 with a particular case of a result due to Hambleton and Teichner [21]. Hambleton and Teichner show that if M is a closed, oriented, topological 4–manifold with infinite cyclic fundamental group and with $b_2(M) - |\sigma(M)| \geq 6$, then M is homeomorphic to the connected sum of $S^1 \times S^3$ with a unique closed, simply connected 4–manifold [21, Corollary 3]. In most cases, this result is stronger than Theorem 1.12. However, there are some situations (such as the union of two genus one surface exteriors $X_{\Sigma_0}, X_{\Sigma_1} \subseteq S^4$ with $\pi_1(X_{\Sigma_i}) = \mathbb{Z}$ for $i = 0, 1$) where Theorem 1.12 applies while [21, Corollary 3] does not. Finally, note that Sunukjian [49, Theorem 7.2] has used these results to study closed surfaces $\Sigma \subseteq X$ with $\pi_1(X \setminus \nu\Sigma) = \mathbb{Z}$ in closed 4–manifolds that satisfy $b_2(X) \geq |\sigma(X)| + 6$.

1.8 Isometries of the Blanchfield form

Theorem 1.10 provides a criterion for extending a homeomorphism $f : \partial M_0 \rightarrow \partial M_1$ to a homeomorphism $M_0 \rightarrow M_1$: one must fit f into a compatible pair. However, in practice, finding compatible pairs is difficult and so we provide some sufficient conditions for their existence.

Recall that if V is a 4–manifold with $\pi_1(V) = \mathbb{Z}$ whose boundary $Y = \partial V$ is ribbon and has torsion Alexander module, then an isometry F of the Λ –intersection form λ_V induces an isometry

$$\partial F : (H_1(Y; \Lambda), \text{Bl}_Y) \xrightarrow{\cong} (H_1(Y; \Lambda), \text{Bl}_Y).$$

We write $\text{Aut}(\lambda_V)$ and $\text{Aut}(\text{Bl}_Y)$ for the groups of isometries of λ_V and Bl_Y , and note that there is a left action of $\text{Aut}(\lambda_V)$ on $\text{Aut}(\text{Bl}_Y)$ given by $F \cdot h = h \circ \partial F^{-1}$. We will be interested in a quotient of the orbit set

$$\text{Aut}(\text{Bl}_Y) / \text{Aut}(\lambda_V).$$

In order to find compatible pairs, we need to know when an isometry of the Blanchfield form is induced by a homeomorphism. As we have already alluded to, we note in Proposition 3.7 that any homeomorphism $f \in \text{Homeo}_\varphi^+(Y)$ induces a Λ –isometry f_* of Bl_Y by lifting f to the infinite cyclic covers. Here, $\text{Homeo}_\varphi^+(Y) := \text{Homeo}_\varphi^+(Y, Y)$ denotes the set of orientation-preserving self-homeomorphisms of Y that intertwine the map $\pi_1(Y) \rightarrow \mathbb{Z}$. The assignment $(f, F) \cdot h = f_* \circ h \circ \partial F^{-1}$ then gives rise to a left action

$$\text{Homeo}_\varphi^+(Y) \times \text{Aut}(\lambda_V) \curvearrowright \text{Aut}(\text{Bl}_Y).$$

We return to our 4–manifolds M_0 and M_1 , whose boundaries are ribbon and have torsion Alexander module. If M_0 and M_1 are orientation-preserving homeomorphic, then a compatible pair exists; recall Theorem 1.10. Assuming that we are given a homeomorphism $f' \in \text{Homeo}_\varphi^+(\partial M_0, \partial M_1)$ and an isometry $F': \lambda_0 \cong \lambda_1$, Proposition 3.14 notes that (f', F') gives rise to a compatible pair (f, F) if the composition $f'_* \circ \partial F'^{-1}$ is trivial in $\text{Aut}(\text{Bl}_{\partial M_1}) / \text{Homeo}_\varphi^+(\partial M_1) \times \text{Aut}(\lambda_{M_1})$. Using this result, we obtain the following consequence of Theorem 1.10.

Theorem 1.14 *Suppose M_0 and M_1 are two compact, oriented 4–manifolds with $\pi_1(M_i) = \mathbb{Z}$ for $i = 0, 1$, whose boundaries are ribbon and have Λ –torsion Alexander modules. Assume M_0 and M_1 have orientation-preserving homeomorphic boundaries, via a homeomorphism that intertwines the maps to \mathbb{Z} , and isometric intersection forms. If the orbit set*

$$\text{Aut}(\text{Bl}_{\partial M_1}) / \text{Homeo}_\varphi^+(\partial M_1) \times \text{Aut}(\lambda_{M_1})$$

is trivial, then there is an orientation-preserving homeomorphism $M_0 \cong M_1$.

Theorem 1.14 is helpful to summarise a programme to prove that two 4–manifolds M_0 and M_1 (with infinite cyclic fundamental group and identical ribbon boundaries with torsion Alexander modules) are homeomorphic:

- (i) Decide whether the Λ –intersection forms λ_{M_0} and λ_{M_1} are isometric.
- (ii) Show that the orbit set $\text{Aut}(\text{Bl}_{\partial M_1}) / \text{Homeo}_\varphi^+(\partial M_1) \times \text{Aut}(\lambda_{M_1})$ is trivial.

To illustrate this paradigm, we explain how Theorem 1.3 can be applied to classify \mathbb{Z} –surfaces for some knots purely in terms of their intersection pairings. For any knot $K \subseteq S^3$, multiplication by a monomial $\pm t^k$ gives rise to an automorphism of its Blanchfield pairing Bl_{E_K} on the exterior of the knot. For some knots K , these are the only automorphisms, and in this case we obtain a simpler classification of \mathbb{Z} –surfaces, ie locally flat, oriented surfaces Σ in N with $\pi_1(N_\Sigma) \cong \mathbb{Z}$, with boundary K .

Corollary 1.15 *Let X be a closed, simply connected, oriented 4–manifold, and let $N := X \setminus \mathring{D}^4$ be a punctured X . Let Σ_0 and Σ_1 be two \mathbb{Z} –surfaces in N with the same genus and oriented boundary, so $\partial \Sigma_0 = \partial \Sigma_1 = K \subseteq S^3$. Suppose that $\text{Aut}(\text{Bl}_{E_K}) \subseteq \{\pm t^k \mid k \in \mathbb{Z}\}$.*

- (i) *There is a rel boundary homeomorphism of pairs $(N, \Sigma_0) \cong (N, \Sigma_1)$ if and only if there is an isometry $\lambda_{N_{\Sigma_0}} \cong \lambda_{N_{\Sigma_1}}$.*
- (ii) *If $N = D^4$, then Σ_0 and Σ_1 are ambiently isotopic rel boundary if and only if there is an isometry $\lambda_{D^4_{\Sigma_0}} \cong \lambda_{D^4_{\Sigma_1}}$.*

Proof Let $F: \lambda_{N_{\Sigma_0}} \xrightarrow{\cong} \lambda_{N_{\Sigma_1}}$ be an isometry. Consider the isomorphism $g := \partial F|_{H_1(E_K; \Lambda)}$ induced on the Alexander module of K . It is an automorphism of the Blanchfield form and therefore by the hypothesis on $\text{Aut}(\text{Bl}_{E_K})$, g is multiplication by $\pm t^k$ for some $k \in \mathbb{Z}$. Precompose F with multiplication by $\pm t^{-k}$, to obtain a new isometry inducing Id on $H_1(E_K; \Lambda)$. Then apply Theorem 1.3 with $f_K = \text{Id}$. \square

Remark 1.16 If the Alexander module $H_1(E_K; \Lambda)$ is cyclic with order Δ_K , then the automorphisms of the Blanchfield pairing can be computed directly as follows. Suppose that $\text{Bl}_K(1, 1) = b/\Delta_K \in \mathbb{Q}(t)/\Lambda$ for some $b \in \Lambda$. Then

$$\text{Aut}(\text{Bl}_{E_K}) = \{[p] \in \Lambda/\Delta\Lambda \mid [p \cdot \bar{p} \cdot b] = [b] \in \Lambda/\Delta\Lambda\}.$$

In particular, for $K = \pm T_{2,3}$ a trefoil, it is not too hard to compute that $\text{Aut}(\text{Bl}_{E_K}) = \{\pm t^k \mid k = -1, 0, 1\}$, so that Corollary 1.15 applies to classify all \mathbb{Z} -surfaces with boundary a trefoil in terms of the isometry type of the intersection pairing.

Organisation

In Section 2, we review some notions on linking forms and define the union of Hermitian forms over Λ . In Section 3, we show how this algebraic union can be used to express the intersection form of a union of two 4-manifolds; we also prove Theorem 1.12. In Section 4, we prove Theorem 1.10, our partial classification result for 4-manifolds with infinite cyclic fundamental group. In Section 5, we prove Theorems 1.3 and 1.4, our main results on properly embedded surfaces in simply connected 4-manifolds. In Section 6, we discuss the equivariant intersection forms for surfaces with knot group \mathbb{Z} , before focusing on surfaces in S^4 and D^4 in Section 7, where we prove Theorems 1.1 and 1.2. In Section 8, we prove Theorems 1.7 and 1.8 on rim surgery. In the appendix, we adapt the work of Baykur and Sunukjian to properly embedded surfaces in 4-manifolds with boundary.

Acknowledgements We thank Peter Teichner for pointing out a mistake in a previous version of this work and for stressing the importance of $(-t)$ -Hermitian forms in the proof of Theorem 7.4. We are indebted to anonymous referees for several very useful suggestions, which helped us to improve the exposition and simplify some proofs. We thank Lisa Piccirillo for helpful discussions, and İnanç Baykur and Nathan Sunukjian for discussions on their stabilisation theorem, which helped us when writing the appendix. Part of this work was completed while the authors were visitors at the Max Planck Institute for Mathematics in Bonn. Powell was partially supported by EPSRC New Investigator grant EP/T028335/1 and EPSRC New Horizons grant EP/V04821X/1.

Conventions

- (i) From now on, all manifolds are assumed to be compact, connected, based and oriented; if a manifold has a nonempty boundary, then the basepoint is assumed to be in the boundary. We work in the topological category with locally flat embeddings unless otherwise stated.
- (ii) X will always denote a closed, simply connected, oriented 4–manifold and we will write $N := X \setminus \overset{\circ}{D}^4$ for the complement of an open ball $\overset{\circ}{D}^4 \subseteq X$.
- (iii) A \mathbb{Z} –surface Σ will always refer to a compact, connected, oriented, locally flat, embedded surface in a 4–manifold, whose knot group is isomorphic to \mathbb{Z} ; a surface $\Sigma \subseteq X$ is understood to be closed, while a surface $\Sigma \subseteq N$ is understood to be properly embedded.
- (iv) If P is manifold and $Q \subseteq P$ is a submanifold with tubular neighbourhood $\nu Q \subseteq P$, then $P_Q := P \setminus \nu Q$ will always denote the exterior of Q in P . The only exception to this is that the exterior of a knot K in S^3 will be denoted by E_K instead of S_K^3 .
- (v) Throughout the article, we set $\Lambda := \mathbb{Z}[t^{\pm 1}]$ for the ring of Laurent polynomials and $\mathbb{Q} := \mathbb{Q}(t)$ for its field of fractions.
- (vi) For a manifold X , we write $\pi_1(X) = \mathbb{Z}$ to mean that there is an isomorphism $\pi_1(X) \cong \mathbb{Z}$, and that we have fixed a choice of such an isomorphism. For the exterior of a \mathbb{Z} –surface, the orientations on X and Σ determine an identification of the fundamental group with \mathbb{Z} . The choice of map $\pi_1(X) = \mathbb{Z}$ determines an isomorphism $\mathbb{Z}[\pi_1(X)] \rightarrow \Lambda$, which we use to define homology and cohomology with Λ coefficients.
- (vii) We write $p \mapsto \bar{p}$ for the involution on Λ induced by sending $t \mapsto t^{-1}$ and extending linearly. Given a Λ –module H , we write \bar{H} for the Λ –module whose underlying abelian group is H but with module structure given by $p \cdot h = \bar{p}h$ for $h \in H$ and $p \in \Lambda$.
- (viii) We write $H^* := \overline{\text{Hom}_\Lambda(H, \Lambda)}$.
- (ix) For any ring R , elements of R^n are considered as column vectors.
- (x) Let $f : X \rightarrow Y$ be a morphism in a category \mathcal{C} , and let $F : \mathcal{C} \rightarrow \mathcal{D}$ be a contravariant functor. Assume that $F(f)$ is invertible. We shall often write the induced morphism in \mathcal{D} as $f^* = F(f)$, and denote its inverse by $f^{-*} := (f^*)^{-1} = F(f)^{-1}$.

2 The union of forms along isometries of their boundary linking forms

We develop the theory required for computing the intersection form of a closed 4–manifold obtained as the union of two compact 4–manifolds along their common boundary. Here is a summary of this section. In Section 2.1, we review the boundary linking form of a Hermitian form. In Section 2.2, we discuss isometries of Hermitian forms and linking forms. In Section 2.3, we study the union of two Hermitian forms over $\Lambda := \mathbb{Z}[t^{\pm 1}]$. In Section 2.4, we provide a condition for this union to admit a metaboliser.

2.1 The boundary of Hermitian forms and their isometries

We fix our terminology on Hermitian forms, linking forms and their boundaries. References include [47, Section 3.4] and [12, Section 6].

A *Hermitian form* over Λ is a pair (H, λ) , where H is a free Λ –module and

$$\lambda: H \times H \rightarrow \Lambda$$

is a sesquilinear Hermitian pairing. Here by sesquilinear, we mean that $\lambda(px, qy) = p\lambda(x, y)\bar{q}$ for all $x, y \in H$ and all $p, q \in \Lambda$. By Hermitian, we mean that $\lambda(y, x) = \overline{\lambda(x, y)}$ for all $x, y \in H$. The *adjoint* of λ is the Λ –linear map

$$\hat{\lambda}: H \rightarrow \overline{\text{Hom}_{\Lambda}(H, \Lambda)} =: H^*$$

such that $\hat{\lambda}(y)(x) = \lambda(x, y)$. A Hermitian form is *nondegenerate* if its adjoint is injective and *nonsingular* if its adjoint is an isomorphism. The *standard hyperbolic form* is

$$H^+(\Lambda) := \left(\Lambda \oplus \Lambda, \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \right).$$

Remark 2.1 If (H, λ) is a nondegenerate Hermitian form, then $\text{coker}(\hat{\lambda})$ is a torsion Λ –module. This can be seen by tensoring the following exact sequence with the field of fractions $Q := \mathbb{Q}(t)$ of Λ :

$$0 \rightarrow H \xrightarrow{\hat{\lambda}} H^* \rightarrow \text{coker}(\hat{\lambda}) \rightarrow 0.$$

A *linking form* (T, β) over Λ consists of a torsion Λ –module T together with a sesquilinear Hermitian form $\beta: T \times T \rightarrow Q/\Lambda$. The *adjoint* of β is the Λ –linear map $\hat{\beta}: T \rightarrow \overline{\text{Hom}_{\Lambda}(T, Q/\Lambda)}$ such that $\hat{\beta}(y)(x) = \beta(x, y)$. A linking form is *nondegenerate* if its adjoint is injective and *nonsingular* if its adjoint is an isomorphism.

Definition 2.2 The *boundary linking form* of a nondegenerate Hermitian form (H, λ) over Λ is the linking form $(\text{coker}(\hat{\lambda}), \partial\lambda)$, where $\partial\lambda$ is defined as

$$\partial\lambda: \text{coker}(\hat{\lambda}) \times \text{coker}(\hat{\lambda}) \rightarrow Q/\Lambda, \quad ([x], [y]) \mapsto \frac{1}{s}(y(z)),$$

where, since $\text{coker}(\hat{\lambda})$ is Λ -torsion, there exists an $s \in \Lambda$ and a $z \in H$ such that $sx = \hat{\lambda}(z)$.

It is not difficult to show that $\partial\lambda$ is independent of the choices involved, and is sesquilinear and Hermitian. We conclude with two remarks that we will use throughout this section.

Remark 2.3 Let (H, λ) be a Hermitian form, and set $H_Q := H \otimes_{\Lambda} Q$. Since H is free, we can identify $\overline{\text{Hom}}_{\Lambda}(H, \Lambda) \otimes_{\Lambda} Q$ with $H_Q^* := \overline{\text{Hom}}_Q(H_Q, Q)$. As stated in Remark 2.1, if (H, λ) is nondegenerate, then $\text{coker}(\hat{\lambda})$ is a torsion Λ -module and therefore $\hat{\lambda}_Q := \hat{\lambda} \otimes_{\Lambda} \text{Id}_Q: H_Q \rightarrow H_Q^*$ is a nonsingular Hermitian form over Q .

Using this remark, we describe an equivalent definition of the boundary linking form.

Remark 2.4 The boundary linking form of a nondegenerate Hermitian form (H, λ) can be described as $\partial\lambda([x], [y]) = y(\hat{\lambda}_Q^{-1}(x))$ for $[x], [y] \in \text{coker}(\hat{\lambda})$. Choose a basis $\mathbf{b} = (e_i)_{i=1}^n$ for H and endow $\overline{\text{Hom}}_{\Lambda}(H, \Lambda)$ with the dual basis \mathbf{b}^* . If $A_{ij} = \lambda(e_i, e_j)$ is a Hermitian matrix representing λ , then \bar{A} is a matrix for $\hat{\lambda}$ and $\partial\lambda([x], [y]) = y(\hat{\lambda}_Q^{-1}(x)) = (\bar{A}^{-1}x)^T \bar{y} = x^T A^{-1} \bar{y}$. For future reference, note that since A is Hermitian, we have $\bar{A} = A^T$.

2.2 Isometries of forms

We discuss isometries of Hermitian forms and linking forms. References include [47, Section 3.4] and [12, Section 6].

Let (H_0, λ_0) and (H_1, λ_1) be Hermitian forms over Λ . A Λ -linear isomorphism $F: H_0 \rightarrow H_1$ is an *isometry* if $\lambda_1(F(x), F(y)) = \lambda_0(x, y)$ for all $x, y \in H_0$. Let $\text{Iso}(\lambda_0, \lambda_1)$ denote the set of isometries between the Hermitian forms (H_0, λ_0) and (H_1, λ_1) , and let $\text{Aut}(\lambda) := \text{Iso}(\lambda, \lambda)$ denote the group of self-isometries of a Hermitian form (H, λ) .

We make the analogous definitions for linking forms. Let (T_0, β_0) and (T_1, β_1) be two linking forms over Λ . A Λ -linear isomorphism $h: T_0 \rightarrow T_1$ is an *isometry*

if $\beta_1(h(x), h(y)) = \beta_0(x, y)$ for all $x, y \in T_0$. We let $\text{Iso}(\beta_0, \beta_1)$ denote the set of isometries between the linking forms (T_0, β_0) and (T_1, β_1) , and let $\text{Aut}(\beta) := \text{Iso}(\beta, \beta)$ denote the group of self-isometries of a linking form (T, β) .

An isomorphism $F: H_0 \rightarrow H_1$ induces an isomorphism $F^{-*} := (F^*)^{-1}: H_0^* \rightarrow H_1^*$. If additionally, the isomorphism $F \in \text{Iso}(\lambda_0, \lambda_1)$ is an isometry, then F^{-*} descends to an isomorphism

$$\partial F := F^{-*}: \text{coker}(\widehat{\lambda}_0) \rightarrow \text{coker}(\widehat{\lambda}_1).$$

For later use, we note that $F: (H_0, \lambda_0) \rightarrow (H_1, \lambda_1)$ is an isometry if and only if $\widehat{\lambda}_0 = F^* \widehat{\lambda}_1 F$. Next, we verify that ∂F is an isometry of the boundary linking forms.

Lemma 2.5 *If $F: (H_0, \lambda_0) \rightarrow (H_1, \lambda_1)$ is an isometry of nondegenerate Hermitian forms, then ∂F is an isometry of linking forms:*

$$\partial F: (\text{coker}(\widehat{\lambda}_0), \partial\lambda_0) \rightarrow (\text{coker}(\widehat{\lambda}_1), \partial\lambda_1).$$

This construction provides a map

$$\partial: \text{Iso}(\lambda_0, \lambda_1) \rightarrow \text{Iso}(\partial\lambda_0, \partial\lambda_1),$$

which is a homomorphism on automorphism groups.

Proof Given $[x], [y] \in \text{coker}(\widehat{\lambda}_0)$, there exist $s \in \Lambda$ and $z \in H_0$ such that $sx = \widehat{\lambda}_0(z)$. Since F is an isometry, it follows that $sF^{-*}(x) = F^{-*}\widehat{\lambda}_0(z) = \widehat{\lambda}_1 F(z)$. We can now conclude that ∂F is an isometry because

$$\begin{aligned} \partial\lambda_1(\partial F([x]), \partial F([y])) &= \partial\lambda_1(F^{-*}([x]), F^{-*}([y])) \\ &= \frac{1}{s} F^{-*}(y) F(z) \\ &= \frac{1}{s} y(z) \\ &= \partial\lambda_0([x], [y]). \end{aligned}$$

The last assertion follows from the equality $(G \circ F)^{-*} = G^{-*} \circ F^{-*}$. □

Using Lemma 2.5, we are led to the following definition.

Definition 2.6 *The boundary of an isometry $F: (H_0, \lambda_0) \rightarrow (H_1, \lambda_1)$ of nondegenerate Hermitian forms is the isometry of linking forms*

$$\partial F: \partial(H_0, \lambda_0) \rightarrow \partial(H_1, \lambda_1).$$

2.3 The union of nondegenerate Hermitian forms

We define the union of two Hermitian forms (H_0, λ_0) and (H_1, λ_1) over Λ along an isometry of their boundary linking forms. The definition is inspired by [11, Chapter 3], which was concerned with forms over the integers.

We describe the main construction of this subsection. In what follows, for $i = 0, 1$, we use $\pi_i : H_i^* \rightarrow \text{coker}(\widehat{\lambda}_i)$ to denote the canonical projections.

Construction 2.7 Let (H_0, λ_0) and (H_1, λ_1) be two nondegenerate Hermitian forms over Λ , and let $h : (\text{coker}(\widehat{\lambda}_0), \partial\lambda_0) \rightarrow (\text{coker}(\widehat{\lambda}_1), \partial\lambda_1)$ be an isometry of their boundary linking forms. Consider the pair (H, λ) with

$$H := \ker(h\pi_0 - \pi_1 : H_0^* \oplus H_1^* \rightarrow \text{coker}(\widehat{\lambda}_1)),$$

$$\lambda\left(\begin{pmatrix} x_0 \\ x_1 \end{pmatrix}, \begin{pmatrix} y_0 \\ y_1 \end{pmatrix}\right) = \frac{1}{s_0}y_0(z_0) - \frac{1}{s_1}y_1(z_1) \in \mathcal{Q},$$

where since $\text{coker}(\widehat{\lambda}_i)$ is torsion, there exist $s_i \in \Lambda$ and $z_i \in H_i$ such that $s_i x_i = \widehat{\lambda}_i(z_i)$. Since the Hermitian forms λ_0 and λ_1 are nondegenerate, it is not difficult to prove that the pairing λ does not depend on the choice of s_0, s_1, z_0, z_1 .

The next proposition establishes some facts about the pairing (H, λ) .

Proposition 2.8 *The pair (H, λ) from Construction 2.7 has the following properties.*

- (i) For $(x_0, x_1), (y_0, y_1) \in H$, the pairing λ can equivalently be defined as

$$\lambda\left(\begin{pmatrix} x_0 \\ x_1 \end{pmatrix}, \begin{pmatrix} y_0 \\ y_1 \end{pmatrix}\right) = y_0(\widehat{\lambda}_{0,Q}^{-1}(x_0)) - y_1(\widehat{\lambda}_{1,Q}^{-1}(x_1)).$$

- (ii) If we choose bases for H_0, H_1 and dual bases for H_0^*, H_1^* and let A_0, A_1 be Hermitian matrices representing λ_0, λ_1 , then

$$\lambda\left(\begin{pmatrix} x_0 \\ x_1 \end{pmatrix}, \begin{pmatrix} y_0 \\ y_1 \end{pmatrix}\right) = x_0^T A_0^{-1} \bar{y}_0 - x_1^T A_1^{-1} \bar{y}_1.$$

- (iii) The pairing λ is sesquilinear, Hermitian and takes values in Λ .
- (iv) The following two maps, which we abridge by $\widehat{\lambda}_0, \widehat{\lambda}_1$, are injective:

$$(H_0, \lambda_0) \xrightarrow{\widehat{\lambda}_0} (H, \lambda) \xleftarrow{\widehat{\lambda}_1} (H_1, -\lambda_1).$$

Furthermore, these maps satisfy $\widehat{\lambda}_0(H_0)^\perp = \widehat{\lambda}_1(H_1)$.

Proof To prove the first assertion, notice that since $\lambda_{i,Q}$ is a nonsingular pairing for $i = 0, 1$, we can write $\widehat{\lambda}_{i,Q}^{-1}(x_i) = z_i/s_i$ for some $z_i \in H$ and some $s_i \in \Lambda$. Consequently, $\widehat{\lambda}_i(z_i) = s_i x_i$ and we have $(1/s_i)y_i(z_i) = y_i(\widehat{\lambda}_{i,Q}^{-1}(x_i))$, as desired. This proves the first assertion. The second assertion now follows as in Remark 2.4.

We prove the third assertion. The fact that λ is sesquilinear and Hermitian follows from the second assertion and the fact that A_0^{-1} and A_1^{-1} are Hermitian. Next, using the definition of the boundary linking forms $\partial\lambda_0$ and $\partial\lambda_1$ from Definition 2.2, we have

$$\lambda\left(\begin{pmatrix} x_0 \\ x_1 \end{pmatrix}, \begin{pmatrix} y_0 \\ y_1 \end{pmatrix}\right) = \partial\lambda_0([x_0], [y_0]) - \partial\lambda_1([x_1], [y_1]) \pmod{\Lambda}.$$

By definition of H , we have $h[x_0] = x_1$ and $h[y_0] = y_1$. Then since h is an isometry, this latter expression vanishes:

$$\begin{aligned} \partial\lambda_0([x_0], [y_0]) - \partial\lambda_1([x_1], [y_1]) &\equiv \partial\lambda_0([x_0], [y_0]) - \partial\lambda_1(h[x_0], h[y_0]) \\ &\equiv \partial\lambda_0([x_0], [y_0]) - \partial\lambda_0([x_0], [y_0]) \equiv 0 \in Q/\Lambda. \end{aligned}$$

It follows that λ takes values in Λ , concluding the proof of the third assertion. We now prove the fourth assertion. First, we check that $\widehat{\lambda}_0$ is an isometric embedding; the proof for $\widehat{\lambda}_1$ is identical. Since the pairings are nondegenerate, the maps are injective and given $z_0, z'_0 \in H_0$, we have

$$\lambda\left(\begin{pmatrix} \widehat{\lambda}_0(z_0) \\ 0 \end{pmatrix}, \begin{pmatrix} \widehat{\lambda}_0(z'_0) \\ 0 \end{pmatrix}\right) = \widehat{\lambda}(z'_0)(z_0) = \lambda_0(z_0, z'_0).$$

It remains to check that $\widehat{\lambda}_0(H_0)^\perp = \widehat{\lambda}_1(H_1)$. The inclusion $\widehat{\lambda}_0(H_0)^\perp \supseteq \widehat{\lambda}_1(H_1)$ is clear and so we prove the inclusion $\widehat{\lambda}_0(H_0)^\perp \subseteq \widehat{\lambda}_1(H_1)$. Assume that $(x_0, x_1) \in H$ (with $sx_0 = \widehat{\lambda}_0(z_0)$) satisfies, for all $a_0 \in H_0$,

$$0 = \lambda\left(\begin{pmatrix} x_0 \\ x_1 \end{pmatrix}, \begin{pmatrix} \widehat{\lambda}_0(a_0) \\ 0 \end{pmatrix}\right) = \frac{1}{s}\widehat{\lambda}_0(a_0)(z_0) = \frac{1}{s}\lambda_0(z_0, a_0) \in \Lambda.$$

As λ_0 is nondegenerate, this implies that $z_0 = 0$ and therefore $x_0 = 0$. But since $(x_0, x_1) \in H$, we also have $[x_1] = h[x_0] = 0$ and so $x_1 = \widehat{\lambda}_1(z_1)$ for some $z_1 \in H_1$. We therefore conclude that $\widehat{\lambda}_0(H_0)^\perp = \widehat{\lambda}_1(H_1)$, establishing the proposition. \square

Note that Proposition 2.8 does not contain a statement about the Λ -module H being free. Since we defined Hermitian forms over free Λ -modules, a *Hermitian module* (H, λ) will refer to a pair consisting of a finitely generated Λ -module H and a sesquilinear Hermitian pairing $\lambda: H \times H \rightarrow \Lambda$; we drop the requirement that H be free.

Definition 2.9 Let (H_0, λ_0) and (H_1, λ_1) be two nondegenerate Hermitian forms over Λ , and let $h : (\text{coker}(\widehat{\lambda}_0), \partial\lambda_0) \rightarrow (\text{coker}(\widehat{\lambda}_1), \partial\lambda_1)$ be an isometry of their boundary linking forms. The *union* of (H_0, λ_0) and (H_1, λ_1) along h is the Hermitian module $(H_0 \cup_h H_1, \lambda_0 \cup_h -\lambda_1)$ described in Construction 2.7 and Proposition 2.8:

$$H_0 \cup_h H_1 := \ker(h\pi_0 - \pi_1 : H_0^* \oplus H_1^* \rightarrow \text{coker}(\widehat{\lambda}_1)),$$

$$\lambda_0 \cup_h -\lambda_1 \left(\begin{pmatrix} x_0 \\ x_1 \end{pmatrix}, \begin{pmatrix} y_0 \\ y_1 \end{pmatrix} \right) = \frac{1}{s_0} y_0(z_0) - \frac{1}{s_1} y_1(z_1) \in \Lambda,$$

where, since $\text{coker}(\widehat{\lambda}_i)$ is torsion, there exist $s_i \in \Lambda$ and $z_i \in H_i$ such that $s_i x_i = \widehat{\lambda}_i(z_i)$.

Remark 2.10 We briefly discuss the nonsingularity of $(H_0 \cup_h H_1, \lambda_0 \cup_h -\lambda_1)$. Proposition 2.8 contains no statement about $(H, \lambda) := (H_0 \cup_h H_1, \lambda_0 \cup_h -\lambda_1)$ being nonsingular. While this holds over the integers [11, Lemma 3.6] and in the topological setting of Proposition 3.9, we will not prove it in the algebraic generality of this section. If $\widehat{\lambda} : H \xrightarrow{\cong} \overline{\text{Hom}_\Lambda(H, \Lambda)} =: H^*$ is nonsingular, then H must be free, since for any finitely generated Λ -module H , the dual H^* is free [3, Lemma 2.1].

Given a Λ -module H , we set $H^* := \overline{\text{Hom}_\Lambda(H, \Lambda)}$. If H is free, then there is a Λ -linear evaluation isomorphism $\text{ev} : H \rightarrow H^{**}$ given by $\text{ev}_x(\varphi) = \overline{\varphi(x)}$, where $x \in H$ and $\varphi \in H^*$. The next remark uses ev to describe the adjoint of $\lambda_0 \cup_h -\lambda_1$.

Remark 2.11 Let (H_0, λ_0) and (H_1, λ_1) be two nondegenerate Hermitian forms over Λ , and let $h \in \text{Iso}(\partial\lambda_0, \partial\lambda_1)$ be an isometry. The adjoint of $\lambda := \lambda_0 \cup_h -\lambda_1$ is given by

$$\widehat{\lambda} : H_0 \cup_h H_1 \rightarrow (H_0 \cup_h H_1)^*, \quad \begin{pmatrix} y_0 \\ y_1 \end{pmatrix} \mapsto (\text{ev} \ \text{ev}) \begin{pmatrix} \widehat{\lambda}_{0,Q}^{-1} & 0 \\ 0 & -\widehat{\lambda}_{1,Q}^{-1} \end{pmatrix} \begin{pmatrix} y_0 \\ y_1 \end{pmatrix}.$$

This follows by combining the definition of ev with the second and third items of Proposition 2.8. Indeed, for $(x_0, y_0) \in H_0 \cup_h H_1$, we have

$$\begin{aligned} \lambda \left(\begin{pmatrix} x_0 \\ x_1 \end{pmatrix}, \begin{pmatrix} y_0 \\ y_1 \end{pmatrix} \right) &= \overline{\lambda \left(\begin{pmatrix} y_0 \\ y_1 \end{pmatrix}, \begin{pmatrix} x_0 \\ x_1 \end{pmatrix} \right)} = \overline{x_0(\widehat{\lambda}_{0,Q}^{-1}(y_0)) - x_1(\widehat{\lambda}_{1,Q}^{-1}(y_1))} \\ &= (\text{ev} \widehat{\lambda}_{0,Q}^{-1}(y_0) \ \text{ev} - \widehat{\lambda}_{1,Q}^{-1}(y_1)) \begin{pmatrix} x_0 \\ x_1 \end{pmatrix} = \widehat{\lambda} \begin{pmatrix} y_0 \\ y_1 \end{pmatrix} \begin{pmatrix} x_0 \\ x_1 \end{pmatrix}. \end{aligned}$$

We conclude this section by describing how the union interacts with boundary isometries. The proof follows immediately from the definitions and is left to the reader.

Proposition 2.12 *Let $(H_0, \lambda_0), (H_1, \lambda_1)$ be Hermitian forms, and $h \in \text{Iso}(\partial\lambda_0, \partial\lambda_1)$ be an isometry. If $F \in \text{Iso}(\lambda_0, \lambda_1)$ is an isometry, then $F^{-*} \oplus \text{Id}$ induces an isometry*

$$(H_0, \lambda_0) \cup_h (H_1, -\lambda_1) \rightarrow (H_1, \lambda_1) \cup_{h \circ \partial F^{-1}} (H_1, -\lambda_1).$$

2.4 Lagrangian complements

This subsection provides a criterion for a union of Hermitian forms to be metabolic. It motivates the notion of a compatible pair, which will be introduced in the next subsection. The idea for this criterion stems from work of Kreck [37, Proposition 8] and Crowley and Sixt [12, Theorem 5.11], and their work on the monoid $\ell_{2q+1}(\mathbb{Z}[\pi])$.

A *Lagrangian* for a nonsingular Hermitian form (H, λ) is a direct summand $L \subseteq H$ such that $L^\perp = L$. A nonsingular Hermitian form that admits a Lagrangian is called *metabolic*. A *Lagrangian complement* for a half-rank direct summand $V \subseteq (H, \lambda)$ is a Lagrangian of (H, λ) such that $L \oplus V = H$.

In the next proposition, for an isomorphism $F: H_0 \rightarrow H_1$ we consider the graph

$$\Gamma_{F^{-*}} = \{(x, F^{-*}(x)) \mid x \in H_0^*\} \subseteq H_0^* \oplus H_1^*.$$

Since, by definition of ∂F , we have $\partial F(x) - F^{-*}(x) = F^{-*}(x) - F^{-*}(x) = 0$, we deduce the inclusions $\Gamma_{F^{-*}} \subseteq \ker(\partial F \circ \pi_0 - \pi_1) = H_0 \cup_{\partial F} H_1 \subseteq H_0^* \oplus H_1^*$.

Proposition 2.13 *Let $F: (H_0, \lambda_0) \rightarrow (H_1, \lambda_1)$ be a Λ -isometry such that $\lambda_0 \cup_{\partial F} -\lambda_1$ is nonsingular. The graph $\Gamma_{F^{-*}} \subseteq H_0 \cup_{\partial F} H_1$ is a Lagrangian complement for $\hat{\lambda}_0(H_0)$:*

$$H_0 \cup_{\partial F} H_1 = \hat{\lambda}_0(H_0) \oplus \Gamma_{F^{-*}}.$$

Proof Set $\lambda := \lambda_0 \cup_{\partial F} -\lambda_1$. We show that $\Gamma_{F^{-*}}$ is a Lagrangian. To see that $\Gamma_{F^{-*}} \subseteq \Gamma_{F^{-*}}^\perp$, we must show that for all $x_0, y_0 \in H_0^*$,

$$\lambda\left(\left(\begin{matrix} x_0 \\ F^{-*}(x_0) \end{matrix}\right), \left(\begin{matrix} y_0 \\ F^{-*}(y_0) \end{matrix}\right)\right) = 0.$$

Pick $s_0 \in \Lambda$ and $z_0 \in H_0$ so that $s_0 x_0 = \hat{\lambda}_0(z_0)$. We apply F^{-*} to both sides of this equation. Since F is an isometry, we have $F^* \hat{\lambda}_1 F = \hat{\lambda}_0$, and we therefore obtain $s_0 F^{-*}(x_0) = \hat{\lambda}_1(F(z_0))$. Using the definition of λ , we now obtain the desired conclusion:

$$\lambda\left(\left(\begin{matrix} x_0 \\ F^{-*}(x_0) \end{matrix}\right), \left(\begin{matrix} y_0 \\ F^{-*}(y_0) \end{matrix}\right)\right) = \frac{1}{s_0} y_0(z_0) - \frac{1}{s_0} F^{-*}(y_0)(F(z_0)) = 0.$$

Next, we show that $\Gamma_{F^{-*}}^\perp \subseteq \Gamma_{F^{-*}}$. We therefore assume that $(x_0, x_1) \in H_0 \cup_{\partial F} H_1$ satisfies, for all $y_0 \in H_0^*$, the equation

$$\lambda\left(\begin{pmatrix} x_0 \\ x_1 \end{pmatrix}, \begin{pmatrix} y_0 \\ F^{-*}(y_0) \end{pmatrix}\right) = 0,$$

and we must show that $x_1 = F^{-*}(x_0)$. Pick $s_0, s_1 \in \Lambda$, $z_0 \in H_0$ and $z_1 \in H_1$ so that $s_0 x_0 = \hat{\lambda}_0(z_0)$ and $s_1 x_1 = \hat{\lambda}_1(z_1)$. We have $s_0 F^{-*}(x_0) = \hat{\lambda}_1(F(z_0))$. Use consecutively the definition of λ and the definition of pullback to obtain

$$0 = \lambda\left(\begin{pmatrix} x_0 \\ x_1 \end{pmatrix}, \begin{pmatrix} y_0 \\ F^{-*}(y_0) \end{pmatrix}\right) = y_0 \begin{pmatrix} z_0 \\ s_0 \end{pmatrix} - F^{-*}(y_0) \begin{pmatrix} z_1 \\ s_1 \end{pmatrix} = y_0 \begin{pmatrix} z_0 \\ s_0 \end{pmatrix} - F^{-1} \begin{pmatrix} z_1 \\ s_1 \end{pmatrix}.$$

Since this equation holds for all y_0 , we deduce that

$$\frac{z_0}{s_0} = F^{-1} \begin{pmatrix} z_1 \\ s_1 \end{pmatrix}.$$

Using that F is an isometry, as well as the definitions of s_0, s_1, z_0 and z_1 , we obtain the desired equation:

$$x_0 = \hat{\lambda}_0 \left(\frac{z_0}{s_0} \right) = \hat{\lambda}_0 \left(F^{-1} \begin{pmatrix} z_1 \\ s_1 \end{pmatrix} \right) = F^* \left(\hat{\lambda}_1 \left(\frac{z_1}{s_1} \right) \right) = F^*(x_1).$$

To conclude the proof that $\Gamma_{F^{-*}}$ is a Lagrangian, it remains to show that it is a direct summand of $H := \ker(\partial F \circ \pi_0 - \pi_1)$. In fact, we will show that

$$\hat{\lambda}_0(H_0) \oplus \Gamma_{F^{-*}} = H.$$

First we establish the inclusion $\hat{\lambda}_0(H_0) + \Gamma_{F^{-*}} \subseteq H$. To do so, we must prove that the sum of arbitrary elements $(\hat{\lambda}_0(z_0), 0) \in \hat{\lambda}_0(H_0)$ and $(y, F^{-*}(y)) \in \Gamma_{F^{-*}}$ belongs to H . In other words, we must show that $\partial F[\hat{\lambda}_0(z_0) + y] = [F^{-*}(y)]$. This uses the fact that F^{-*} induces ∂F (by Definition 2.6) and the fact that, since F is an isometry, $F^{-*} \circ \hat{\lambda}_0(z_0) = \hat{\lambda}_1(F(z_0))$ belongs to $\text{im}(\hat{\lambda}_1)$ and so the class $[F^{-*} \circ \hat{\lambda}_0(z_0)]$ must vanish in the quotient $\text{coker}(\hat{\lambda}_1)$:

$$\partial F[\hat{\lambda}_0(z_0) + y] = [F^{-*} \hat{\lambda}_0(z_0)] + [F^{-*}(y)] = [F^{-*}(y)].$$

Next, we prove the reverse inclusion, namely that $H \subseteq \hat{\lambda}_0(H_0) + \Gamma_{F^{-*}}$. To show this, we must write $(x, y) \in H$ as a sum of an element in $\hat{\lambda}_0(H_0)$ with an element in $\Gamma_{F^{-*}}$. Consider the decomposition

$$(x, y) = (x - F^*(y), 0) + (F^*(y), y).$$

The second term certainly lies in $\Gamma_{F^{-*}}$, so we need to argue that the first term lies in $\hat{\lambda}_0(H_0)$. Since $(x, y) \in H$, we know that $\partial F[x] = [y]$ in $\text{coker}(\hat{\lambda}_1)$. It follows that

$F^{-*}(x) - y \in H_1^*$ belongs to $\text{im}(\widehat{\lambda}_1)$. Since F is an isometry, this is equivalent to saying that $x - F^*(y) \in \text{im}(\widehat{\lambda}_0)$, as desired.

Having proved that $\widehat{\lambda}_0(H_0) + \Gamma_{F^{-*}} = H$, it remains to check that $\widehat{\lambda}_0(H_0) \cap \Gamma_{F^{-*}} = 0$ (so that the sum is direct): if $(\widehat{\lambda}_0(x_0), 0) = (x, F^{-*}(x))$ belongs to this intersection, then $F^{-*}(x) = 0$ and therefore $0 = x = \widehat{\lambda}_0(x_0)$. This proves Proposition 2.13. \square

3 The intersection form of the union of two 4–manifolds

We start our study of 4–manifolds with infinite cyclic fundamental group and of their Λ –intersection forms. First, recall the following definition from the introduction.

Definition 3.1 A 4–manifold M has *ribbon boundary* if ∂M is nonempty and path-connected, and the map $\pi_1(\partial M) \rightarrow \pi_1(M)$ induced by the inclusion is surjective.

Here is a summary of Section 3. In Section 3.1, we show that if M is a 4–manifold with infinite cyclic fundamental group whose boundary is ribbon and has torsion Alexander module, then the boundary linking form $\partial\lambda_M$ of λ_M is isometric to minus the Blanchfield form of ∂M . In Section 3.2, we describe when homeomorphisms of 3–manifolds induce isometries of the Blanchfield pairing. In Section 3.3 we show that the Λ –intersection form of a union of two manifolds can be expressed using the algebraic union from Section 2.3. In Section 3.4, we introduce the notion of a compatible pair and prove Theorem 1.12 from the introduction.

3.1 The boundary of the intersection form

If a 4–manifold M with $\pi_1(M) = \mathbb{Z}$ has ribbon boundary and Λ –torsion Alexander module, then ∂M is endowed with a sesquilinear Hermitian nonsingular linking form over Λ , namely the Blanchfield form

$$\text{Bl}_{\partial M}: H_1(\partial M; \Lambda) \times H_1(\partial M; \Lambda) \rightarrow Q/\Lambda.$$

This subsection establishes that the boundary linking form of the Λ –intersection form of M is isometric to minus the Blanchfield form of ∂M . Since results of this form are known (see eg [4, Theorem 2.6]), we only outline the argument so as to fix some notation for later use.

We collect some homological facts about 4–manifolds with infinite cyclic fundamental group.

Lemma 3.2 *Let M be a 4–manifold with $\pi_1(M) = \mathbb{Z}$. When ∂M is nonempty, we assume it is ribbon and has torsion Alexander module $H_1(\partial M; \Lambda)$. The following assertions hold:*

- (i) $H_0(M; \Lambda) \cong \mathbb{Z}$ and, if $\partial M \neq \emptyset$, then $H_2(\partial M; \Lambda) \cong \mathbb{Z}$.
- (ii) The Λ –modules $H_1(M; \Lambda)$, $H_1(M, \partial M; \Lambda)$ and $H_3(M; \Lambda)$ all vanish.
- (iii) The Λ –modules $H_2(M; \Lambda)$ and $H_2(M, \partial M; \Lambda)$ are free.
- (iv) When $\partial M = \emptyset$, the Λ –intersection form λ_M is nonsingular, whereas for $\partial M \neq \emptyset$, λ_M is nondegenerate, and any matrix representing it presents the Alexander module of ∂M .

Proof Throughout this proof we use that since the manifold has fundamental group \mathbb{Z} , homology with Λ –coefficients can be computed as the homology of the universal cover. Since universal covers are 1–connected, we immediately deduce that $H_0(M; \Lambda) \cong \mathbb{Z}$ and $H_1(M; \Lambda) = 0$. If the boundary is nonempty, the long exact sequence of the pair shows that $H_1(M, \partial M; \Lambda) = 0$; here we used that ∂M is connected, as assumed in Definition 3.1. For later use, we also note that $H_0(M, \partial M; \Lambda) = 0$, also owing to the fact that ∂M is connected. As $H_1(\partial M; \Lambda)$ is torsion, we have that $\text{Hom}_\Lambda(H_1(\partial M; \Lambda), \Lambda) = 0$, and thus $H_2(\partial M; \Lambda) \cong \text{Ext}_\Lambda^1(H_0(\partial M; \Lambda), \Lambda) \cong \mathbb{Z}$ by duality and the universal coefficient spectral sequence [39, Theorem 2.3]. Since $H_i(M, \partial M; \Lambda) = 0$ for $i = 0, 1$, duality and the universal coefficient spectral sequence (UCSS for short) also show that $H_3(M; \Lambda) = H^1(M, \partial M; \Lambda) = 0$. Thus we have proved the first two assertions; we now prove the third. Poincaré duality and the UCSS imply that

$$H_2(M; \Lambda) \cong H^2(M, \partial M; \Lambda) \cong \overline{\text{Hom}_\Lambda(H_2(M, \partial M; \Lambda), \Lambda)},$$

$$H_2(M, \partial M; \Lambda) \cong H^2(M; \Lambda) \cong \overline{\text{Hom}_\Lambda(H_2(M; \Lambda), \Lambda)}.$$

Since the dual of a finitely generated Λ –module is free [3, Lemma 2.1], we deduce that these second homology modules are free over Λ . This proves the third assertion and also establishes the fourth, namely that in the closed case, the Λ –intersection form is nonsingular. Finally, when the boundary is nonempty, the map

$$\mathbb{Z} \cong H_2(\partial M; \Lambda) \rightarrow H_2(M; \Lambda)$$

is the zero map (since $H_2(M; \Lambda)$ is free) and therefore the intersection form is nondegenerate; it presents $H_1(\partial M; \Lambda)$ because we established that $H_1(M; \Lambda) = 0$. □

In the case with nonempty connected boundary, we fix bases for the free Λ –modules $H_2(M; \Lambda)$ and $H_2(M, \partial M; \Lambda)$.

Remark 3.3 Let M be a 4–manifold with $\pi_1(M) = \mathbb{Z}$ whose boundary is ribbon and has torsion Alexander module. Fix a basis \mathbf{b} for $H := H_2(M; \Lambda)$, endow the dual $H^* := \overline{\text{Hom}}_\Lambda(H, \Lambda)$ with the dual basis \mathbf{b}^* , and equip $H_2(M, \partial M; \Lambda)$ with the basis $\text{PD} \circ \text{ev}^{-1}(\mathbf{b}^*)$ coming from the isomorphisms $\text{ev}: H^2(M; \Lambda) \rightarrow H^*$ and $\text{PD}: H^2(M; \Lambda) \rightarrow H_2(M, \partial M; \Lambda)$. A short computation shows that if the matrix A is defined as $A_{ij} := \lambda_M(b_i, b_j)$, then $A^T = \bar{A}$ is a matrix for $\hat{\lambda}_M: H \rightarrow H^*$; recall Remark 2.1. The same conclusion holds for the map $H \rightarrow H_2(M, \partial M; \Lambda)$ induced by the inclusion [9, Section 5.2]. As mentioned in the fourth item of Lemma 3.2, the connecting homomorphism δ in the long exact sequence of the pair $(M, \partial M)$, together with Poincaré duality and the evaluation map, determines a map

$$D_M : \text{coker}(\hat{\lambda}_M) = \text{coker}(A^T) \xrightarrow{\cong} H_1(\partial M; \Lambda), \quad [x] \mapsto \delta \circ \text{PD} \circ \text{ev}^{-1}(x),$$

where $x \in H_2(M; \Lambda)^*$. The map D_M is well-defined and an isomorphism. That it is well-defined follows from a diagram chase in the next diagram, in which all homology has Λ coefficients. The left two vertical maps are isomorphisms, and therefore by the five lemma so is D_M . To help the reader to parse the basis choices above, we also indicate the matrices representing the maps on the left of the diagram:

$$(1) \quad \begin{array}{ccccccc} 0 & \longrightarrow & H_2(M) & \xrightarrow[A^T]{\hat{\lambda}_M} & H_2(M)^* & \longrightarrow & \text{coker}(\hat{\lambda}_M) \longrightarrow 0 \\ & & \downarrow I \text{ Id} & & \downarrow I \text{ PD} \circ \text{ev}^{-1} & & \downarrow D_M \\ 0 & \longrightarrow & H_2(M) & \xrightarrow[A^T]{j} & H_2(M, \partial M) & \xrightarrow{\delta} & H_1(\partial M) \longrightarrow 0 \end{array}$$

Next, we briefly recall the definition of the Blanchfield pairing, referring to [9; 18] as references in which the conventions are identical to ours.

Definition 3.4 Let N be a closed 3–manifold with an epimorphism $\varphi: \pi_1(N) \twoheadrightarrow \mathbb{Z}$ such that the resulting Alexander module is Λ –torsion. The adjoint of the Blanchfield pairing is defined by the composition

$$H_1(N; \Lambda) \xrightarrow[\text{PD}^{-1}]{\cong} H^2(N; \Lambda) \xrightarrow[\text{BS}^{-1}]{\cong} H^1(N; Q/\Lambda) \xrightarrow[\text{ev}]{\cong} \overline{\text{Hom}}_\Lambda(H_1(N; \Lambda), Q/\Lambda)$$

of the inverse of Poincaré duality, the inverse of a Bockstein homomorphism, and the evaluation map. The Blanchfield pairing is a nonsingular linking form, so in particular it is sesquilinear and Hermitian.

Now we can prove the main result of this subsection.

Proposition 3.5 *Let M be a 4-manifold with $\pi_1(M) = \mathbb{Z}$ whose boundary is ribbon and has torsion Alexander module. The map D_M defined in Remark 3.3 induces an isometry*

$$D_M : \partial(H_2(M; \Lambda), \lambda_M) = (\text{coker}(\widehat{\lambda}_M), \partial\lambda_M) \xrightarrow{\cong} (H_1(\partial M; \Lambda), -\text{Bl}_{\partial M}).$$

Proof Choose bases for $H_2(M; \Lambda)$ and $H_2(M, \partial M; \Lambda)$ as in Remark 3.3, and use the notation of that remark. The same argument as in [9, Section 5] shows that the following diagram (with exact columns) commutes:

$$\begin{array}{ccc}
 0 & & 0 \\
 \downarrow & & \downarrow \\
 H_2(M; \Lambda) \times H_2(M; \Lambda) & \xrightarrow{(x,y) \mapsto -x^T A \bar{y}} & \Lambda \\
 \downarrow A^T \times A^T & & \downarrow \\
 H_2(M, \partial M; \Lambda) \times H_2(M, \partial M; \Lambda) & \xrightarrow{(x,y) \mapsto -x^T A^{-1} \bar{y}} & Q \\
 \downarrow \delta \times \delta & & \downarrow \\
 H_1(\partial M; \Lambda) \times H_1(\partial M; \Lambda) & \xrightarrow{\text{Bl}_{\partial M}} & Q/\Lambda \\
 \downarrow & & \downarrow \\
 0 & & 0
 \end{array}$$

In particular, the isomorphism

$$D_M : \text{coker}(\widehat{\lambda}_M) = \text{coker}(A^T) \xrightarrow{\cong} H_1(\partial M; \Lambda)$$

induced by δ and discussed in Remark 3.3 is an isometry between the linking forms $(\text{coker}(\widehat{\lambda}_M), \partial\lambda_M)$ and $(H_1(\partial M; \Lambda), -\text{Bl}_{\partial M})$. Here we are using the characterisation of $\partial\lambda_M$ from Remark 2.4. □

The next remark records that the presence of a minus sign in Proposition 3.5 is immaterial once one passes to isometry groups, and defines the bijection $D^\#$ which identifies the two isometry groups.

Remark 3.6 Given linking forms (H_0, β_0) and (H_1, β_1) , observe that canonically we have $\text{Iso}(\beta_0, \beta_1) = \text{Iso}(-\beta_0, -\beta_1)$; the presence of the minus sign has no effect. In particular if M_0 and M_1 are 4-manifolds with $\pi_1(M_i) = \mathbb{Z}$ for $i = 0, 1$, whose boundaries are ribbon and have torsion Alexander modules, then we can use the maps $D_i := D_{M_i}$ from Proposition 3.5 to obtain an identification

$$D^\# : \text{Iso}(\partial\lambda_{M_0}, \partial\lambda_{M_1}) \xrightarrow{\cong} \text{Iso}(\text{Bl}_{\partial M_0}, \text{Bl}_{\partial M_1}), \quad \tilde{f} \mapsto D_1 \circ \tilde{f} \circ D_0^{-1}.$$

3.2 Homeomorphisms and isometries

In this short subsection, we provide a condition for a homeomorphism between 3–manifolds to induce an isometry of Blanchfield forms.

Proposition 3.7 *Suppose that Y_0^3, Y_1^3 are 3–manifolds equipped with epimorphisms $\varphi_i: \pi_1(Y_i) \twoheadrightarrow \mathbb{Z}$ and assume that the resulting Alexander modules $H_1(Y_i; \Lambda)$ are Λ –torsion for $i = 0, 1$. If an orientation–preserving homotopy equivalence $f: Y_0^3 \rightarrow Y_1^3$ satisfies $\varphi_1 \circ f_* = \varphi_0$ on $\pi_1(Y_0)$, then it induces an isometry between the Blanchfield forms:*

$$f_*: (H_1(Y_0; \Lambda), \text{Bl}_{Y_0}) \rightarrow (H_1(Y_1; \Lambda), \text{Bl}_{Y_1}).$$

Proof Since we assumed that $\varphi_1 \circ f_* = \varphi_0$, the homotopy equivalence f induces Λ –isomorphisms $f_*: H_*(Y_0; \Lambda) \rightarrow H_*(Y_1; \Lambda)$ (and similarly on cohomology). The fact that f has degree one, together with the naturality of the UCSS and the Bockstein homomorphism, ensures that f_* intertwines the Blanchfield pairings, concluding the proof of the proposition. □

For pairs (Y_0, φ_0) and (Y_1, φ_1) as in the statement of Proposition 3.7, we consider those orientation–preserving homeomorphisms that intertwine the φ_i :

$$\text{Homeo}_\varphi^+(Y_0, Y_1) = \{f \in \text{Homeo}^+(Y_0, Y_1) \mid \varphi_1 \circ f_* = \varphi_0\}.$$

In the case where $(Y_0, \varphi_0) = (Y_1, \varphi_1)$, we simply write $\text{Homeo}_\varphi^+(Y_0)$ for the resulting group.

3.3 Algebraic unions and topological unions

In this subsection, we show that under favourable conditions, the Λ –intersection form of a union of two 4–manifolds can be expressed using the algebraic union of Hermitian forms from Section 2.3. For that however, given two manifolds M_0, M_1 with $\pi_1(M_i) = \mathbb{Z}$, we need to verify that $\pi_1(M_0 \cup_f M_1) = \mathbb{Z}$. Here and from now on, when $\pi_i(M_i) = \mathbb{Z}$, we take $\varphi_i: \pi_1(\partial M_i) \rightarrow \pi_1(M_i) = \mathbb{Z}$ to be the map induced by the inclusion. Thus, when we write $\text{Homeo}_\varphi^+(\partial M_0, \partial M_1)$, it is with respect to these inclusion induced maps.

Lemma 3.8 *If $f \in \text{Homeo}_\varphi^+(\partial M_0, \partial M_1)$ is a homeomorphism where M_0, M_1 are 4–manifolds with ribbon boundaries and $\pi_1(M_i) = \mathbb{Z}$, then $\pi_1(M_0 \cup_f M_1) = \mathbb{Z}$.*

Proof In order to apply van Kampen’s theorem, for $i = 0, 1$ we write $\pi_1(M_i) = \mathbb{Z}\langle t_i \rangle$, so that $\pi_1(M_0 \cup_f -M_1)$ is generated by t_0, t_1 . Consider the commutative diagram

$$\begin{array}{ccccc}
 \pi_1(\partial M_0) & \xrightarrow{f_*, \cong} & & \xrightarrow{\quad} & \pi_1(\partial M_1) \\
 \downarrow \iota_0 & \searrow \varphi_0 & & \swarrow \varphi_1 & \downarrow \iota_1 \\
 \pi_1(M_0) \cong \mathbb{Z}\langle t_0 \rangle & \xrightarrow{\psi_0, \cong} & \mathbb{Z}\langle t \rangle & \xleftarrow{\psi_1, \cong} & \mathbb{Z}\langle t_1 \rangle \cong \pi_1(M_1)
 \end{array}$$

Here, for $j = 0, 1$, $\iota_j : \pi_1(\partial M_j) \rightarrow \pi_1(M_j)$ is the inclusion induced map and $\psi_j : t_j \mapsto t$. In the group $\pi_1(M_0 \cup_f -M_1)$, the relations identify $\iota_1 f_*(g)$ with $\iota_0(g)$ for every $g \in \pi_1(\partial M_0)$. Since ι_0 is surjective, there is a $\mu \in \pi_1(\partial M_0)$ with $\iota_0(\mu) = t_0$. We deduce that in $\pi_1(M_0 \cup_f -M_1)$, the relations identify $t_0 = \varphi_0(\mu)$ with $\iota_1(f_*(\mu)) = t_1^n$ for some n . Under the identification given by the ψ_i and using that $\varphi_0(\mu) = \varphi_1(f_*(\mu))$, we deduce that $n = 1$. Thus $\pi_1(M_0 \cup_f -M_1)$ is cyclic. For every $g \in \pi_1(\partial M_0)$, we have $\iota_0(g) = t_0^m$ for some m , and the same reasoning as above shows that $\iota_0(g) = t_0^m$ is identified with $\iota_1(f_*(g)) = t_1^m$ in $\pi_1(M_0 \cup_f -M_1)$. It follows that $\pi_1(M_0 \cup_f -M_1)$ is isomorphic to $\langle t_0, t_1 \mid t_0 = t_1 \rangle \cong \mathbb{Z}$. \square

Next, we prove the main result of this section, after setting up some notation. Continuing with the setting of Lemma 3.8, we write $M := M_0 \cup_f -M_1$. Note that $\hat{\lambda}_M : H_2(M; \Lambda) \rightarrow H_2(M; \Lambda)^*$ is an isomorphism because $\partial M = \emptyset$; recall the fourth item of Lemma 3.2. This endows $H_2(M; \Lambda)^*$ with a Hermitian form, which we shall denote by λ_M^{-1} .

For $i = 0, 1$, we set $\hat{\lambda}_i := \hat{\lambda}_{M_i}$, and as above we write $D_i := D_{M_i}$. Combining Propositions 3.5 and 3.7, we have an isometry

$$\tilde{f}_* : \text{coker}(\hat{\lambda}_0) \xrightarrow{D_0} H_1(\partial M_0; \Lambda) \xrightarrow{f_*} H_1(\partial M_1; \Lambda) \xrightarrow{D_1^{-1}} \text{coker}(\hat{\lambda}_1),$$

which satisfies $D^\#(\tilde{f}_*) = f_*$.

Proposition 3.9 *Let M_0, M_1 be two 4–manifolds with $\pi_1(M_i) = \mathbb{Z}$ for $i = 0, 1$, whose boundaries are ribbon and have torsion Alexander modules. Given a homeomorphism $f \in \text{Homeo}_\varphi^+(\partial M_0, \partial M_1)$, set $M := M_0 \cup_f -M_1$, and let $\iota_j : M_i \rightarrow M$ be the inclusion. The map*

$$\begin{pmatrix} \iota_0^* \\ \iota_1^* \end{pmatrix} : H_2(M; \Lambda)^* \rightarrow H_2(M_0; \Lambda)^* \oplus H_2(M_1; \Lambda)^*$$

induces an isometry

$$(2) \quad (H_2(M; \Lambda)^*, \lambda_M^{-1}) \xrightarrow{\cong} (H_2(M_0; \Lambda) \cup_{\tilde{f}_*} H_2(M_1; \Lambda), \lambda_{M_0} \cup_{\tilde{f}_*} -\lambda_{M_1}).$$

Furthermore, the map $\widehat{\lambda}_M$ induces an isometry $(H_2(M; \Lambda), \lambda_M) \rightarrow (H_2(M; \Lambda)^*, \lambda_M^{-1})$, which, using the identification from (2), makes the following diagram commute:

$$(3) \quad \begin{array}{ccccc} (H_2(M_0; \Lambda), \lambda_{M_0}) & \xrightarrow{\iota_0} & (H_2(M; \Lambda), \lambda_M) & \xleftarrow{\iota_1} & (H_2(M_1; \Lambda), -\lambda_{M_1}) \\ & \searrow \widehat{\lambda}_{M_0} & \downarrow \cong \widehat{\lambda}_M & \swarrow \widehat{\lambda}_{M_1} & \\ & & (H_2(M_0; \Lambda) \cup_{\tilde{f}_*} H_2(M_1; \Lambda), \lambda_{M_0} \cup_{\tilde{f}_*} -\lambda_{M_1}) & & \end{array}$$

In particular, $H_2(M_0; \Lambda) \cup_{\tilde{f}_*} H_2(M_1; \Lambda)$ is a free Λ -module and $\lambda_{M_0} \cup_{\tilde{f}_*} -\lambda_{M_1}$ is nonsingular.

Proof Lemma 3.8 implies that $\pi_1(M) = \mathbb{Z}$. Consider the following commutative diagram in which the first and third rows are exact and Λ coefficients are understood:

$$(4) \quad \begin{array}{ccccccc} 0 & \longrightarrow & H_2(M_0) \oplus H_2(M_1) & \xrightarrow{(\iota_0 \ \iota_1)} & H_2(M) & \longrightarrow & H_1(\partial M_1) \longrightarrow 0 \\ & & \downarrow j_0 \oplus j_1 & & \downarrow \text{PD}_M^{-1} & & \\ & & H_2(M_0, \partial M_0) \oplus H_2(M_1, \partial M_1) & & & & \\ & & \downarrow \text{PD}_{M_0}^{-1} \oplus -\text{PD}_{M_1}^{-1} & & \downarrow \begin{pmatrix} \iota_0^* \\ \iota_1^* \end{pmatrix} & & \\ H^2(\partial M_1) & \xleftarrow{f^{-*} \text{incl}_0^* - \text{incl}_1^*} & H^2(M_0) \oplus H^2(M_1) & \xleftarrow{\begin{pmatrix} \iota_0^* \\ \iota_1^* \end{pmatrix}} & H^2(M) & \xleftarrow{\quad} & 0 \\ & & \downarrow \cong \text{ev} \oplus \text{ev} & & \downarrow \cong \text{ev} & & \\ & & H_2(M_0)^* \oplus H_2(M_1)^* & \xleftarrow{\begin{pmatrix} \iota_0^* \\ \iota_1^* \end{pmatrix}} & H_2(M)^* & & \end{array}$$

We justify the three zeros that appear. In the first row, the rightmost zero comes from the fact that $H_1(M_i; \Lambda) = 0$ for $i = 0, 1$ by the second item of Lemma 3.2. For the leftmost zero of the first row, the first and third items of Lemma 3.2 respectively imply that $H_2(\partial M_i; \Lambda) = \mathbb{Z}$ is Λ -torsion and that $H_2(M_i; \Lambda)$ is free. It follows that the inclusion induced homomorphism $H_2(\partial M_i; \Lambda) \rightarrow H_2(M_i; \Lambda)$ is zero. The same argument explains the appearance of the zero in the third row: $H^2(M; \Lambda)$ is free and $H^1(\partial M_1; \Lambda)$ is Λ -torsion. The commutativity of the middle square follows by applying [7, Lemma 8.2] with $K = M_0$ and $L = M_1$, as well as applying excision to the pairs $(M, M \setminus M_i)$.

Our next aim is to simplify the diagram in (4). We write $\pi_i: H_2(M_i; \Lambda)^* \rightarrow \text{coker}(\widehat{\lambda}_i)$ for the canonical projection.

Claim 1 The isometry $\tilde{f}_* : \text{coker}(\hat{\lambda}_0) \xrightarrow{\cong} \text{coker}(\hat{\lambda}_1)$ fits in the diagram

$$(5) \quad \begin{array}{ccccccc} 0 & \longrightarrow & H_2(M_0; \Lambda) \oplus H_2(M_1; \Lambda) & \xrightarrow{(\iota_0 \ \iota_1)} & H_2(M; \Lambda) & \longrightarrow & H_1(\partial M_1; \Lambda) \\ & & \downarrow \hat{\lambda}_0 \oplus -\hat{\lambda}_1 & & \cong \downarrow \hat{\lambda}_M & & \\ \text{coker}(\hat{\lambda}_1) & \xleftarrow{\tilde{f}_* \pi_0 - \pi_1} & H_2(M_0; \Lambda)^* \oplus H_2(M_1; \Lambda)^* & \xleftarrow{\begin{pmatrix} \iota_0^* \\ \iota_1^* \end{pmatrix}} & H_2(M; \Lambda)^* & \xleftarrow{\quad} & 0 \end{array}$$

where the rows are exact and the square is commutative.

Proof The two vertical compositions in (4) are by definition $\hat{\lambda}_0 \oplus -\hat{\lambda}_1$ and $\hat{\lambda}_M$, respectively. Thus the exactness of the top row and the commutativity of the square follow from (4). It therefore remains to establish the exactness of the bottom row in (5). A second look at (4) shows a similar exact sequence with cohomology instead of duals and $H^2(\partial M_1; \Lambda)$ instead of $\text{coker}(\hat{\lambda}_1)$. We shall produce the commuting square

$$(6) \quad \begin{array}{ccc} H^2(\partial M_1; \Lambda) & \xleftarrow{f^{-*} \text{incl}_0^* - \text{incl}_1^*} & H^2(M_0; \Lambda) \oplus H^2(M_1; \Lambda) \\ \cong \downarrow G_1 & & \cong \downarrow \text{ev} \oplus \text{ev} \\ \text{coker}(\hat{\lambda}_1) & \xleftarrow{\tilde{f}_* \pi_0 - \pi_1} & H_2(M_0; \Lambda)^* \oplus H_2(M_1; \Lambda)^* \end{array}$$

that can be appended to the bottom left of (4). The claim will then follow immediately.

We define isomorphisms G_i , for $i = 0, 1$, as the composition

$$G_i := D_i^{-1} \circ \text{PD} : H^2(\partial M_i; \Lambda) \xrightarrow{\cong} H_1(\partial M_i; \Lambda) \xrightarrow{\cong} \text{coker}(\hat{\lambda}_i).$$

The map G_i agrees with the rightmost vertical downwards composition (inverting D_i) in the next diagram. Again Λ coefficients are understood, and the rows are exact:

$$(7) \quad \begin{array}{ccccccccc} 0 & \longrightarrow & H^2(M_i, \partial M_i) & \xrightarrow{j_i^*} & H^2(M_i) & \xrightarrow{\text{incl}_i^*} & H^2(\partial M_i) & \longrightarrow & 0 \\ & & \text{PD} \downarrow \cong & & \text{PD} \downarrow \cong & & \text{PD} \downarrow \cong & & \\ 0 & \longrightarrow & H_2(M_i) & \xrightarrow{j_i} & H_2(M_i, \partial M_i) & \xrightarrow{\delta_i} & H_1(\partial M_i) & \longrightarrow & 0 \\ & & \parallel \text{Id} & & \cong \uparrow \text{PD} \circ \text{ev}^{-1} & & \cong \uparrow D_i & & \\ 0 & \longrightarrow & H_2(M_i) & \xrightarrow{\hat{\lambda}_i} & H_2(M_i)^* & \xrightarrow{\pi_i} & \text{coker}(\hat{\lambda}_i) & \longrightarrow & 0 \end{array}$$

The diagram commutes by the definition of $\hat{\lambda}_i$ as $\text{ev} \circ \text{PD}^{-1} \circ j_i$, and using standard naturality of Poincaré duality; for the bottom right square, one uses the definition of D_i from Remark 3.3. Via G_i and ev , the concatenation of the two squares on the right identifies the maps π_i and incl_i^* . That is, for $i = 0, 1$,

$$(8) \quad G_i \circ \text{incl}_i^* = \pi_i \circ \text{ev}: H^2(M_i; \Lambda) \rightarrow \text{coker}(\hat{\lambda}_i).$$

This shows that the bottom left square in the following diagram commutes. This diagram again has Λ coefficients and vertical maps isomorphisms:

$$(9) \quad \begin{array}{ccccc} & & H_1(\partial M_0; \Lambda) & \xrightarrow[\cong]{f_*} & H_1(\partial M_1; \Lambda) \\ & & \cong \downarrow \text{PD}^{-1} & & \cong \downarrow \text{PD}^{-1} \\ H^2(M_0) & \xrightarrow{\text{incl}_0^*} & H^2(\partial M_0) & \xrightarrow[\cong]{f^{-*}} & H^2(\partial M_1) \\ \text{ev} \downarrow \cong & & G_0 \downarrow \cong & & G_1 \downarrow \cong \\ H_2(M_0)^* & \xrightarrow{\pi_0} & \text{coker}(\hat{\lambda}_0) & \xrightarrow[\cong]{\tilde{f}_*} & \text{coker}(\hat{\lambda}_1) \end{array}$$

Commutativity of the top right square follows from naturality of Poincaré duality and the fact that all four maps are isomorphisms. Using this and that $D_i = \text{PD} \circ G_i^{-1}$ for $i = 0, 1$, starting with the definition of $\tilde{f}_*: \text{coker}(\hat{\lambda}_0) \rightarrow \text{coker}(\hat{\lambda}_1)$ we obtain

$$\tilde{f}_* = D_1^{-1} \circ f_* \circ D_0 = G_1 \circ \text{PD}^{-1} \circ f_* \circ \text{PD} \circ G_0^{-1} = G_1 \circ f^{-*} \circ G_0^{-1}.$$

In other words, the bottom right square of (9) commutes.

Therefore the bottom of diagram (9) yields $G_1 \circ f^{-*} \circ \text{incl}_0^* = \tilde{f}_* \circ \pi_0 \circ \text{ev}$. Subtracting (8) with $i = 1$, we obtain the desired commutative square (6). The diagram in (5) then follows from the combination of (4) and (6), completing the proof of the claim. \square

Now we write $T := H_1(\partial M_1; \Lambda)$, as well as $H_i := H_2(M_i; \Lambda)$ for $i = 0, 1$. We also set $V := H_2(M; \Lambda)^*$ and identify $H_2(M; \Lambda)$ with the double dual $H_2(M; \Lambda)^{**} = V^*$ via the evaluation isomorphism ev . With this notation, the diagram from Claim 1 leads to the following commutative diagram in which both rows are exact:

$$(10) \quad \begin{array}{ccccccc} 0 & \longrightarrow & H_0 \oplus H_1 & \xrightarrow{(\text{ev} \ \text{ev})} & V^* & \longrightarrow & T \\ & & \downarrow \hat{\lambda}_0 \oplus -\hat{\lambda}_1 & & \cong \downarrow \hat{\lambda}_M & & \\ \text{coker}(\hat{\lambda}_1) & \xleftarrow{\tilde{f}_* \pi_0 - \pi_1} & H_0^* \oplus H_1^* & \xleftarrow{\quad} & V & \xleftarrow{\quad} & 0 \end{array}$$

Identify $V = H_2(M; \Lambda)^*$ with $\ker(\tilde{f}_*\pi_0 - \pi_1)$ and view it as a subspace of $H_0^* \oplus H_1^*$. The diagram in (10) therefore yields the equation

$$(11) \quad \begin{pmatrix} \hat{\lambda}_0 & 0 \\ 0 & -\hat{\lambda}_1 \end{pmatrix} = \hat{\lambda}_M \circ (\text{ev} \text{ ev}).$$

Since Lemma 3.2 implies that $H_2(M; \Lambda)$ is a free Λ -module and $H_1(M; \Lambda) = 0$, we may identify $H^2(M; Q)$ with $H^2(M; \Lambda) \otimes_{\Lambda} Q$, and similarly for

$$H^2(M_i, \partial M_i; Q) \cong H^2(M_i, \partial M_i; \Lambda) \otimes_{\Lambda} Q \quad \text{and} \quad H^2(M_i; Q) \cong H^2(M_i; \Lambda) \otimes_{\Lambda} Q$$

for $i = 0, 1$. As a consequence, we may identify the tensored up intersection pairings $\lambda_M \otimes_{\Lambda} \text{Id}_Q$ and $\lambda_i \otimes_{\Lambda} \text{Id}_Q$ with the nonsingular Q -valued intersections pairings on $H_2(M; Q)$ and $H_2(M_i; Q)$. Since $\hat{\lambda}_M$ is invertible by Lemma 3.2, we deduce from (11) that

$$\hat{\lambda}_M^{-1} = (\text{ev} \text{ ev}) \circ \begin{pmatrix} \hat{\lambda}_{0,Q}^{-1} & 0 \\ 0 & -\hat{\lambda}_{1,Q}^{-1} \end{pmatrix}.$$

As noted in Remark 2.11, this is precisely the adjoint of $\lambda_0 \cup_{\tilde{f}_*} -\lambda_1$.

We have thus proved that the inclusions ι_0 and ι_1 induce an isometry

$$(H_2(M; \Lambda)^*, \lambda_M^{-1}) \cong (H_0 \cup_{\tilde{f}_*} H_1, \lambda_0 \cup_{\tilde{f}_*} -\lambda_1).$$

Now, since $\lambda_M^* = \lambda_M$, a quick verification shows that the isomorphism

$$\hat{\lambda}_M: H_2(M; \Lambda) \rightarrow H_2(M; \Lambda)^*$$

induces an isometry between the forms $(H_2(M; \Lambda), \lambda_M)$ and $(H_2(M; \Lambda)^*, \lambda_M^{-1})$.

Indeed, writing $H := H_2(M; \Lambda)$, this follows from the following commutative diagram:

$$\begin{array}{ccc} H & \xrightarrow{\hat{\lambda}_M} & H^* \\ \hat{\lambda}_M \downarrow & \cong & \downarrow \hat{\lambda}_M^{-1} \\ H^* & \xleftarrow{\cong} & H \\ & \hat{\lambda}_M^* = \hat{\lambda}_M & \end{array}$$

The commutativity of the diagram in (3) follows from the commutativity of (10): we have shown that $(H_0 \cup_{\tilde{f}_*} H_1, \lambda_0 \cup_{\tilde{f}_*} -\lambda_1) \cong (H_2(M; \Lambda), \lambda_M)$ and, recalling that $(\text{ev}, \text{ev}) = (\iota_0, \iota_1)$, the diagram in (10) gives $\hat{\lambda}_M \circ \iota_0(x_0, 0) = \hat{\lambda}_0(x_0, 0)$, leading to the commutativity of the left-hand triangle of (3); the reasoning for the right-hand triangle is identical. □

3.4 Compatible pairs

Throughout this section, M_0 and M_1 are 4-manifolds with $\pi_1(M_i) = \mathbb{Z}$ for $i = 0, 1$, whose boundaries are ribbon and have torsion Alexander modules. This section introduces a notion of compatibility for an isometry $F: H_2(M_0; \Lambda) \rightarrow H_2(M_1; \Lambda)$ and a homeomorphism $f: \partial M_0 \rightarrow \partial M_1$. We then prove Theorem 1.12 from the introduction.

Since the boundaries are ribbon, and we have identifications $\pi_1(M_i) = \mathbb{Z}$, the inclusions $\partial M_i \hookrightarrow M_i$ induce surjections $\varphi_i: \pi_1(M_i) \twoheadrightarrow \mathbb{Z}$ for $i = 0, 1$. Recall from Section 3.2 that we set

$$\text{Homeo}_\varphi^+(\partial M_0, \partial M_1) = \{f \in \text{Homeo}^+(\partial M_0, \partial M_1) \mid \varphi_1 \circ f_* = \varphi_0\}.$$

Proposition 3.7 ensures that a homeomorphism $f \in \text{Homeo}_\varphi^+(\partial M_0, \partial M_1)$ induces an isometry $f_* \in \text{Iso}(\text{Bl}_{\partial M_0}, \text{Bl}_{\partial M_1})$ of the Blanchfield forms. On the other hand, recall from Lemma 2.5 that an isometry $F \in \text{Iso}(\lambda_{M_0}, \lambda_{M_1})$ of the Λ -intersection forms of M_0 and M_1 induces an isometry $\partial F = F^{-*}$ of the boundary linking forms. Using Proposition 3.5, we identify these boundary linking forms with minus the corresponding Blanchfield forms, so that using Remark 3.6 we may consider $D^\#(\partial F) \in \text{Iso}(\text{Bl}_{\partial M_0}, \text{Bl}_{\partial M_1})$.

Definition 3.10 An orientation-preserving homeomorphism $f \in \text{Homeo}_\varphi^+(\partial M_0, \partial M_1)$ is *compatible* with an isometry $F \in \text{Iso}(\lambda_{M_0}, \lambda_{M_1})$ if, using the identification $D^\#$ from Remark 3.6, we have

$$D^\#(\partial F) = f_*: (H_1(\partial M_0; \Lambda), \text{Bl}_{\partial M_0}) \rightarrow (H_1(\partial M_1; \Lambda), \text{Bl}_{\partial M_1}).$$

In this case, the pair (f, F) is called a *compatible pair*.

The next proposition shows that the existence of a compatible pair is a necessary condition for a homeomorphism $f \in \text{Homeo}_\varphi^+(\partial M_0, \partial M_1)$ to extend to a homeomorphism $M_0 \rightarrow M_1$.

Proposition 3.11 Let $f \in \text{Homeo}_\varphi^+(\partial M_0, \partial M_1)$ be a homeomorphism, and let $F \in \text{Iso}(\lambda_{M_0}, \lambda_{M_1})$ be an isometry. If f extends to an orientation-preserving homeomorphism $\Phi: M_0 \xrightarrow{\cong} M_1$ that induces F , then (f, F) is a compatible pair.

Proof Write $\lambda_i := \lambda_{M_i}$ and $D_i := D_{M_i}$ for $i = 0, 1$. The homeomorphism Φ induces an isomorphism $\Phi_*: \pi_1(M_0) \rightarrow \pi_1(M_1)$. By the assumption on f , the isomorphisms $\pi_1(M_i) = \mathbb{Z}$ are such that Φ_* intertwines them. This follows because in the next

diagram both triangles, the top square, and the large outside square commute:

$$\begin{array}{ccc}
 \pi_1(\partial M_0) & \xrightarrow{f_*} & \pi_1(\partial M_1) \\
 \downarrow & & \downarrow \\
 \pi_1(M_0) & \xrightarrow{\Phi_*} & \pi_1(M_1) \\
 \parallel & & \parallel \\
 \mathbb{Z} & \xlongequal{\quad} & \mathbb{Z}
 \end{array}
 \begin{array}{c}
 \varphi_0 \curvearrowright \\
 \varphi_1 \curvearrowleft
 \end{array}$$

It follows that the bottom square commutes too. By assumption Φ induces the maps

$$F: H_2(M_0; \Lambda) \xrightarrow{\cong} H_2(M_1; \Lambda) \quad \text{and} \quad F^{-*}: H_2(M_0; \Lambda)^* \xrightarrow{\cong} H_2(M_1; \Lambda)^*.$$

By Lemma 2.5 the map F^{-*} induces an isometry

$$\partial F: (\text{coker}(\hat{\lambda}_0), \partial\lambda_0) \xrightarrow{\cong} (\text{coker}(\hat{\lambda}_1), \partial\lambda_1).$$

We assert that ∂F fits into a commuting diagram of isometries

$$\begin{array}{ccc}
 \text{coker}(\hat{\lambda}_0) & \xrightarrow[\cong]{\partial F} & \text{coker}(\hat{\lambda}_1) \\
 \cong \downarrow D_0 & & \cong \downarrow D_1 \\
 H_1(\partial M_0; \Lambda) & \xrightarrow[\cong]{f_*} & H_1(\partial M_1; \Lambda)
 \end{array}
 \tag{12}$$

The maps D_i are isometries by Proposition 3.5, and the map f_* in the bottom row is an isometry by Proposition 3.7. By definition of $D^\#$, commutativity of the diagram (12) is equivalent to $D^\#(\partial F) = f_*$, ie to the fact that (f, F) is a compatible pair. To show that (12) commutes, we use the next diagram, in which all homology groups have Λ coefficients and the rightmost horizontal maps are all surjections:

$$\begin{array}{ccccccc}
 H_2(M_0) & \xrightarrow{\hat{\lambda}_0} & H_2(M_0)^* & \twoheadrightarrow & \text{coker}(\hat{\lambda}_0) & & \\
 \swarrow F_* \cong & \parallel \text{Id} & \swarrow F^{-*} \cong & \cong \downarrow \text{PD} \circ \text{ev}^{-1} & \swarrow \partial F \cong & \downarrow D_0 & \\
 H_2(M_1) & \xrightarrow{\hat{\lambda}_1} & H_2(M_1)^* & \twoheadrightarrow & \text{coker}(\hat{\lambda}_1) & & \\
 \downarrow \text{Id} & \parallel & \downarrow \text{PD} \circ \text{ev}^{-1} & \cong \downarrow & \downarrow D_1 & & \\
 H_2(M_0) & \xrightarrow{j_0} & H_2(M_0, \partial M_0) & \twoheadrightarrow & H_1(\partial M_0) & & \\
 \swarrow F_* \cong & \parallel & \swarrow F_* \cong & \cong \downarrow & \swarrow f_* \cong & & \\
 H_2(M_1) & \xrightarrow{j_1} & H_2(M_1, \partial M_1) & \twoheadrightarrow & H_1(\partial M_1) & &
 \end{array}$$

The map $\partial F: \text{coker}(\hat{\lambda}_0) \rightarrow \text{coker}(\hat{\lambda}_1)$ is by definition the map induced by the upper part of the diagram. The squares in the leftmost cube commute either trivially, by

the definition of $\widehat{\lambda}_i$, or by naturality of Poincaré duality. The potentially contentious point is the latter justification: naturality in fact says that $F_* \circ \text{PD}_{M_0} \circ F^* = \text{PD}_{M_1}$. But since F^* is an isomorphism, with inverse F^{-*} , it follows that the middle vertical square commutes as shown. A straightforward diagram chase now shows, as desired, that the rightmost vertical square, which is equivalent to (12), also commutes. This completes the proof that (f, F) is a compatible pair. \square

The next result shows that the existence of a compatible pair (f, F) imposes strong restrictions on the topology of $M := M_0 \cup_f -M_1$. This is Theorem 1.12 from the introduction.

Theorem 3.12 *Let (f, F) be a compatible pair. If M_0 and M_1 are not spin, assume that the Kirby–Siebenmann invariants satisfy $\text{ks}(M_0) = \text{ks}(M_1) \in \mathbb{Z}/2$. Then there is an orientation-preserving homeomorphism*

$$M := M_0 \cup_f -M_1 \cong S^1 \times S^3 \# \#_{i=1}^a S^2 \times S^2 \# \#_{j=1}^b S^2 \widetilde{\times} S^2$$

for some a, b with $a + b = b_2(M_0)$. If M_0 and M_1 are spin, then $b = 0$.

Moreover, it can be assumed that the Λ -isometry induced by this homeomorphism takes the Lagrangian $\widehat{\lambda}_M^{-1}(\Gamma_{F^{-*}}) \subseteq H_2(M; \Lambda)$ to the Lagrangian that is generated by $\{[\text{pt}] \times S^2\}_{i=1}^{b_2(M_0)}$.

Proof Lemma 3.8 establishes that $\pi_1(M) = \mathbb{Z}$. Proposition 3.9 proves that λ_M is isometric to $\lambda_{M_0} \cup_{\widetilde{f}_*} -\lambda_{M_1}$. We have $\lambda_{M_0} \cup_{\widetilde{f}_*} -\lambda_{M_1} = \lambda_{M_0} \cup_{\partial F} -\lambda_{M_1}$ because the pair (f, F) is compatible. It follows from Proposition 2.13 that $\lambda_{M_0} \cup_{\partial F} -\lambda_{M_1}$ is metabolic with $\Gamma_{F^{-*}}$ as a Lagrangian. We deduce that M is a closed 4-manifold with $\pi_1(M) = \mathbb{Z}$ and metabolic Λ -intersection form. The (nonsingular) intersection form of M (which can be obtained from λ_M by setting $t = 1$) is therefore isometric to

$$\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}^{\oplus a} \oplus \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}^{\oplus b}$$

for some a, b . As $\text{ks}(M_0) = \text{ks}(M_1)$, by additivity of the Kirby–Siebenmann invariant [17, Theorem 8.2] under gluing it follows that $\text{ks}(M) = 0$. Apart from the statement that $b = 0$ for M_i spin, the theorem now follows from the classification of closed 4-manifolds with infinite cyclic fundamental group due to [15, Theorem 10.7A(2)] and [48]: every isometry between the Λ -intersection forms of two closed, oriented 4-manifolds with fundamental group \mathbb{Z} can be realised by a homeomorphism.

Now assume that M_0 and M_1 are spin. To show that $b = 0$, it is sufficient to show that λ_M is hyperbolic, as the result will then once again follow from [15, Theorem 10.7A(2)] and [48]. We already argued that λ_M is isometric to $\lambda_{M_0} \cup_{\partial F} -\lambda_{M_1}$ because the pair (f, F) is compatible. Applying Proposition 2.12 to the isometry $h = \partial F$, this latter form is isometric to $\lambda_{M_1} \cup_{\text{Id}} -\lambda_{M_1}$. The identity certainly belongs to $\text{Homeo}_\phi^+(\partial M_1, \partial M_1)$ and so a second application of Proposition 3.9 ensures that $\lambda_{M_1} \cup_{\text{Id}} -\lambda_{M_1}$ is isometric to $\lambda_{M_1 \cup_{\text{Id}} -M_1}$. Summarising, we have the following sequence of isometries:

$$(13) \quad \lambda_M \cong \lambda_{M_0 \cup_{\tilde{f}^*} -M_1} \cong \lambda_{M_0} \cup_{\partial F} -\lambda_{M_1} \cong \lambda_{M_1} \cup_{\text{Id}} -\lambda_{M_1} \cong \lambda_{M_1 \cup_{\text{Id}} -M_1}.$$

Since M_1 is spin, the double $DM_1 := M_1 \cup_{\text{Id}} -M_1$ is also spin. As $\pi_2(DM_1) = H_2(DM_1; \Lambda)$ is free (by the third item of Lemma 3.2 and Lemma 3.8) and DM_1 is spin, it follows from [26, Remark 1.7] that λ_{DM_1} is even, meaning that $\lambda_{DM_1} = q + \bar{q}$ for some sesquilinear form q . Since λ_{DM_1} and λ_M are isometric by (13), it follows that λ_M is even. Now since λ_M is an even metabolic nonsingular Hermitian form on a finitely generated free Λ -module, it is hyperbolic by [34, Corollary 3.7.3] with Γ_{F-*} as a Lagrangian. □

We make the Lagrangian $\widehat{\lambda}_M^{-1}(\Gamma_{F-*}) \subseteq H_2(M; \Lambda)$ somewhat more concrete. In the setting of Theorem 3.12, combine the inclusion induced maps as

$$\iota = (\iota_0 \ \iota_1): H_2(M_0; \Lambda) \oplus H_2(M_1; \Lambda) \rightarrow H_2(M; \Lambda).$$

We record the following fact; compare with [5, Proposition 4.2] in the simply connected case.

Lemma 3.13 *The Lagrangian $\widehat{\lambda}_M^{-1}(\Gamma_{F-*}) \subseteq H_2(M; \Lambda)$ contains the graph Γ_{-F} of the isometry $-F: H_2(M_0; \Lambda) \rightarrow H_2(M_1; \Lambda)$.*

Proof Given $x_0 \in H_2(M_0; \Lambda)$, we have

$$\begin{aligned} \iota \left(\begin{pmatrix} x_0 \\ -F(x_0) \end{pmatrix} \right) &= \widehat{\lambda}_M^{-1} \left(\begin{pmatrix} \widehat{\lambda}_{M_0}(x_0) \\ \widehat{\lambda}_{M_1}(F(x_0)) \end{pmatrix} \right) \\ &= \widehat{\lambda}_M^{-1} \left(\begin{pmatrix} \widehat{\lambda}_0(x_0) \\ F^{-*}(\widehat{\lambda}_{M_0}(x_0)) \end{pmatrix} \right). \end{aligned}$$

Here, the first equality holds by the commutativity of the diagram in (10), while the second holds because F is an isometry. It follows that $\Gamma_{-F} \subseteq \widehat{\lambda}_M^{-1}(\Gamma_{F-*})$, as desired. □

We introduce some notation needed to provide a sufficient criterion to produce compatible pairs. Under the identification $\text{Aut}(\partial\lambda_{M_1}) = \text{Aut}(\text{Bl}_{\partial M_1})$ as in Remark 3.6, there is a left action

$$\begin{aligned} &(\text{Homeo}_\varphi^+(\partial M_1) \times \text{Aut}(\lambda_{M_1})) \times \text{Aut}(\text{Bl}_{\partial M_1}) \rightarrow \text{Aut}(\text{Bl}_{\partial M_1}), \\ &(f, F) \cdot h := f_* \circ h \circ \partial F^{-1}. \end{aligned}$$

The next proposition gives a criterion to find a compatible pair.

Proposition 3.14 *For M_0 and M_1 as above, if there exist $f \in \text{Homeo}_\varphi^+(\partial M_0, \partial M_1)$ and $F \in \text{Iso}(\lambda_{M_0}, \lambda_{M_1})$, and if the orbit set*

$$\text{Aut}(\text{Bl}_{\partial M_1}) / \text{Homeo}_\varphi^+(\partial M_1) \times \text{Aut}(\lambda_{M_1})$$

is trivial, then a compatible pair (f', F') exists.

Proof The composition $f_* \circ \partial F^{-1}$ is an isometry of $\partial\lambda_{M_1} \cong \text{Bl}_{\partial M_1}$. By the assumption, there exist a homeomorphism $g \in \text{Homeo}_\varphi^+(\partial M_1)$ and an isometry $G \in \text{Aut}(\lambda_{M_1})$ such that we have $f_* \circ \partial F^{-1} = g_*^{-1} \circ \partial G$. Equivalently, we can write $(g \circ f)_* = \partial(G \circ F)$ and consequently the pair $(f', F') := (g \circ f, G \circ F)$ is compatible. \square

4 Partial classification of compact 4–manifolds with fundamental group \mathbb{Z}

In this section, we prove Theorem 1.10 from the introduction. The proof of this result was inspired by [5] and [37].

Theorem 4.1 *Let M_0 and M_1 be two 4–manifolds with $\pi_1(M_i) = \mathbb{Z}$ for $i = 0, 1$, whose boundaries are ribbon and have torsion Alexander modules. Suppose that $f \in \text{Homeo}_\varphi^+(\partial M_0, \partial M_1)$ is an orientation-preserving homeomorphism, and suppose that $F \in \text{Iso}(\lambda_{M_0}, \lambda_{M_1})$ is an isometry. If M_0 and M_1 are not spin, assume that the Kirby–Siebenmann invariants satisfy $\text{ks}(M_0) = \text{ks}(M_1) \in \mathbb{Z}/2$. Then the following assertions are equivalent:*

- (i) *The pair (f, F) is compatible.*
- (ii) *The homeomorphism f extends to an orientation-preserving homeomorphism*

$$\Phi: M_0 \xrightarrow{\cong} M_1$$

inducing the given isometry $F: H_2(M_0; \Lambda) \cong H_2(M_1; \Lambda)$.

Proof Proposition 3.11 shows that (ii) implies (i). To prove the converse, given a compatible pair (f, F) , we must therefore establish the existence of a homeomorphism Φ . Under this assumption, and in the nonspin case assuming that the Kirby–Siebenmann invariants coincide, we saw in Theorem 3.12 that for some a, b such that $a + b = b_2(M_0)$, there is a homeomorphism

$$M_0 \cup_f -M_1 \cong S^1 \times S^3 \# \#_{i=1}^a S^2 \times S^2 \# \#_{i=1}^b S^2 \tilde{\times} S^2.$$

Define $M := M_0 \cup_f -M_1$. With this notation, Theorem 3.12 additionally states that this homeomorphism takes the Lagrangian $\hat{\lambda}_M^{-1}(\Gamma_{F-*}) \subseteq H_2(M; \Lambda)$ to the Lagrangian generated by the spheres $\{[\text{pt}] \times S^2\}_{i=1}^{b_2(M_0)}$. Consider the 5–dimensional null-bordism for M given by

$$W := S^1 \times D^4 \natural \natural_{i=1}^a S^2 \times D^3 \natural \natural_{i=1}^b S^2 \tilde{\times} D^3,$$

where $S^2 \tilde{\times} D^3$ denotes the twisted linear D^3 bundle over S^2 . We prove that W is an h –cobordism; it will then automatically be an s –cobordism as $\text{Wh}(\mathbb{Z}) = 0$. As $H_i(W, M_0; \Lambda) = 0$ for $i \neq 2, 3$, we must show that $H_2(W, M_0; \Lambda) = 0$ for $i = 2, 3$.

Use $D_{1/2}^3 \subseteq D^3$ to denote a 3–ball of smaller radius inside D^3 . Choosing the obvious set of generators for $H_2(W; \Lambda)$ and representing them by embedded spheres, we obtain

$$U := \natural_{i=1}^a S^2 \times D_{1/2}^3 \natural \natural_{i=1}^b S^2 \tilde{\times} D_{1/2}^3 \subseteq W.$$

Note that $H_*(U; \Lambda) = H_*(W; \Lambda)$ and $H_*(\partial U; \Lambda) = H_*(M; \Lambda)$. In particular, under this identification, we have

$$(14) \quad \ker(H_2(M; \Lambda) \xrightarrow{j} H_2(W; \Lambda)) = \text{im}(H_3(U, \partial U; \Lambda) \rightarrow H_2(\partial U; \Lambda)) \\ = \hat{\lambda}_M^{-1}(\Gamma_{F-*}).$$

In what follows, we identify $H_3(U, \partial U; \Lambda)$ with its image in $H_2(\partial U; \Lambda)$ allowing ourselves for instance to write $H_3(U, \partial U; \Lambda) \subseteq H_2(\partial U; \Lambda)$. We also think of the adjoint $\hat{\lambda}_M$ of the intersection form on M as having $H_2(\partial U; \Lambda)$ as its domain.

Claim 2 *The connecting homomorphism*

$$\partial_0: H_3(W \setminus \overset{\circ}{U}, M_0 \sqcup \partial U; \Lambda) \rightarrow H_2(M_0; \Lambda),$$

which arises from the long exact sequence of the triple $(W \setminus \overset{\circ}{U}, M_0 \sqcup \partial U, \partial U)$, is an isomorphism.

Proof of Claim 2 It suffices to note that by excision and by our choice of U and W , we have $H_i(W \setminus \overset{\circ}{U}, \partial U; \Lambda) \cong H_i(W, U; \Lambda) = 0$ for $i = 2, 3$. \square

Claim 3 *The following diagram commutes:*

$$\begin{array}{ccccc}
 H_2(M_0; \Lambda) & \xleftarrow{\hat{\lambda}_0} & (H_2(M_0; \Lambda) \cup_{\partial F} H_2(M_1; \Lambda), \lambda_0 \cup_{\partial F} \lambda_1) & \xleftarrow{\Gamma_{F^*}} & \\
 \cong \uparrow \partial_0 & & \cong \uparrow \hat{\lambda}_M & & \cong \uparrow \hat{\lambda}_M \\
 H_3(W \setminus \overset{\circ}{U}, M_0 \sqcup \partial U; \Lambda) & \xleftarrow{\quad} & (H_2(\partial U; \Lambda), \lambda_{\partial U}) & \xleftarrow{\quad} & H_3(U, \partial U; \Lambda)
 \end{array}$$

Proof of Claim 3 The commutativity of the right square follows from (14), and so we focus on the left square. Consider the long exact sequence of the triples

$$(W \setminus \overset{\circ}{U}, M_0 \sqcup \partial U, \partial U) \quad \text{and} \quad (W \setminus \overset{\circ}{U}, M_0 \sqcup \partial U, M_0).$$

The portions of interest can be seen in the two horizontal rows of the following diagram, where Λ coefficients are understood:

$$\begin{array}{ccccccc}
 H_3(W \setminus \overset{\circ}{U}, \partial U) = 0 & \rightarrow & H_3(W \setminus \overset{\circ}{U}, M_0 \sqcup \partial U) & \xrightarrow{\partial_0} & H_2(M_0) & \rightarrow & 0 = H_2(W \setminus \overset{\circ}{U}, \partial U) \\
 & & \parallel & & \downarrow \iota_0 & \searrow \hat{\lambda}_0 & \\
 (15) & & & & H_2(M) & \xrightarrow{\cong} & H_2(M_0) \\
 & & & & \parallel & \searrow \hat{\lambda}_M & \cup_{\partial F} \\
 & & & & & & H_2(M_1) \\
 & & & & \parallel & \nearrow \hat{\lambda}_M & \\
 H_3(W \setminus \overset{\circ}{U}, M_0) = 0 & \rightarrow & H_3(W \setminus \overset{\circ}{U}, M_0 \sqcup \partial U) & \rightarrow & H_2(\partial U) & \rightarrow & H_2(W \setminus \overset{\circ}{U}, M_0)
 \end{array}$$

The claim will follow from the central portion of the diagram, once we explain all its features and its commutativity. The zeros in the first row are a consequence of Claim 2. The bottom leftmost zero is stated in Kreck’s work [37, page 734], but we outline a proof in this setting. The exact sequence of the triple $(W \setminus \overset{\circ}{U}, M_1 \sqcup \partial U, M_1)$, together with a Mayer–Vietoris argument, give rise to an isomorphism

$$H_1(W \setminus \overset{\circ}{U}, M_1 \sqcup \partial U; \Lambda) \cong H_0(\partial U; \Lambda) = \Lambda.$$

Similarly, using the long exact sequence of the triple $(W \setminus \overset{\circ}{U}, M_1 \sqcup \partial U, \partial U)$, we deduce that $H_2(W \setminus \overset{\circ}{U}, M_1 \sqcup \partial U; \Lambda) = 0$. Since we also have $H_0(W \setminus \overset{\circ}{U}, M_1 \sqcup \partial U; \Lambda) = 0$, Poincaré duality and the UCSS imply that

$$H_3(W \setminus \overset{\circ}{U}, M_0; \Lambda) = H^2(W \setminus \overset{\circ}{U}, M_1 \sqcup \partial U; \Lambda) = 0.$$

To establish the claim, it only remains to show that the diagram in (15) is commutative. The middle square clearly commutes. Proposition 3.9 establishes the commutativity of the triangles but with $H_2(M_0) \cup_{\tilde{f}_*} H_2(M_1)$ in place of $H_2(M_0) \cup_{\partial F} H_2(M_1)$. However, since (f, F) is a compatible pair, these two modules are isomorphic. This concludes our explanation of the diagram in (15) and concludes the proof of Claim 3. \square

Write A and B for the images of $H_3(U, \partial U; \Lambda)$ and $H_3(W \setminus \overset{\circ}{U}; M_0 \sqcup \partial U; \Lambda)$ in $H_2(\partial U; \Lambda)$. As Proposition 2.13 implies that

$$H_2(M_0; \Lambda) \cup_{\partial F} H_2(M_1; \Lambda) = \Gamma_{F^{-*}} \oplus H_2(M_0; \Lambda),$$

we deduce from Claim 3 that

$$H_2(\partial U; \Lambda) = A \oplus B.$$

Assemble the long exact sequence of the triple $(W \setminus \overset{\circ}{U}, M_0 \sqcup \partial U, M_0)$ and the long exact sequence of the triple $(W, W \setminus \overset{\circ}{U}, M_0)$ into the following diagram, based on that from [37, page 738], in which Λ -coefficients are understood:

$$\begin{array}{ccccccc}
 & & & & 0 & & \\
 & & & & \downarrow & & \\
 & & & & B \cong H_3(W \setminus \overset{\circ}{U}, M_0 \sqcup \partial U) & & \\
 & & & & \downarrow & & \\
 A \cong H_3(U, \partial U) & \longrightarrow & H_2(\partial U) & & & & \\
 \downarrow \cong & & \downarrow & & & & \\
 0 \longrightarrow H_3(W, M_0) \longrightarrow H_3(W, W \setminus \overset{\circ}{U}) \longrightarrow H_2(W \setminus \overset{\circ}{U}, M_0) \longrightarrow H_2(W, M_0) \longrightarrow 0 & & & & & & \\
 & & & & \downarrow & & \\
 & & & & 0 & &
 \end{array}$$

Here, the left vertical isomorphism comes from excision. Since $H_2(\partial U; \Lambda) = A \oplus B$, the right-down composition $A \rightarrow H_2(W \setminus \overset{\circ}{U}, M_0; \Lambda)$ is an isomorphism. It follows that the central map in the long row is an isomorphism, and therefore $H_i(W, M_0; \Lambda) = 0$ for $i = 2, 3$. Thus W is a relative h -cobordism, and therefore an s -cobordism, as desired.

Since \mathbb{Z} is a good group, by the topological s -cobordism theorem [15, Theorem 7.1A] M_0 and M_1 are homeomorphic via a homeomorphism Φ that extends f . It remains to show that Φ induces the isometry F on $H_2(-; \Lambda)$. The inclusions of M_0, M_1 into $M = M_0 \cup_f M_1 \subseteq W$ give rise to the homomorphism

$$H_2(M_0) \oplus H_2(M_1) \xrightarrow{\iota=(\iota_0 \ \iota_1)} H_2(M) \xrightarrow{j} H_2(W).$$

Set $j_i = j \circ \iota_i$; these are isomorphisms because W is an s -cobordism. By definition $j_1^{-1} j_0$ is the isometry induced by the homeomorphism Φ . We noted that $H_2(U; \Lambda) = H_2(W; \Lambda)$ as well as $H_2(\partial U; \Lambda) = H_2(M; \Lambda)$. We also noted in (14) that $\ker(j) = \widehat{\lambda}_M^{-1}(\Gamma_{F-*})$. Using Lemma 3.13, we deduce that $\iota(\Gamma_{-F}) \subseteq \ker(j)$. This inclusion implies that for all $x_0 \in H_2(M_0; \Lambda)$,

$$0 = j \iota \begin{pmatrix} x_0 \\ -F(x_0) \end{pmatrix} = j \iota_0(x_0) - j \iota_1(F(x_0)) = j_0(x_0) - j_1(F(x_0)).$$

Since the j_i are isomorphisms, we obtain the desired conclusion: $j_1^{-1} j_0(x_0) = F(x_0)$. This concludes the proof of Theorem 4.1. □

5 Knotted surfaces in simply connected 4-manifolds

The goal of this section is to use Theorem 4.1 to prove Theorems 1.3 and 1.4 from the introduction. Recall from our conventions that X refers to a closed, simply connected, oriented 4-manifold, while $N = X \setminus \mathring{D}^4$, and Σ always denotes a \mathbb{Z} -surface of genus g either embedded in X or properly embedded in N (recall that a \mathbb{Z} -surface refers to an oriented surface whose knot group is \mathbb{Z}).

This section is organised as follows. In Section 5.1, we collect some initial facts concerning \mathbb{Z} -surfaces. Section 5.2 then shows that any isometry of $\text{Bl}_{\Sigma_{g,1} \times S^1}$ can be realised by a homeomorphism of $\Sigma_{g,1} \times S^1$ (allowing us to construct a compatible pair between \mathbb{Z} -surfaces exteriors N_{Σ_0} and N_{Σ_1}). In Sections 5.3 and 5.4, respectively, we prove Theorems 1.3 and 1.4 from the introduction.

5.1 Facts about \mathbb{Z} -surfaces

Lemma 5.1 *If $\Sigma \subseteq N$ is a \mathbb{Z} -surface, then it is null-homologous.*

Proof We must show that $[\Sigma, \partial\Sigma] = 0 \in H_2(N, \partial N)$. The intersection form Q_N pairs $H_2(N, \partial N)$ with $H_2(N)$ nonsingularly because N is simply connected. Thus, it is equivalent to show that $Q_N([\Sigma, \partial\Sigma], x) = 0$ for every class $x \in H_2(N)$. Represent such an $x \in H_2(N)$ by a closed surface $S \subseteq N$ that intersects Σ transversely in points p_1, \dots, p_n , so that $Q_N([\Sigma, \partial\Sigma], x) = \sum_{i=1}^n \varepsilon(p_i)$, where $\varepsilon(p_i) = \pm 1$. Now the intersection $S \cap N_\Sigma$ is a properly embedded surface in N_Σ with oriented boundary (homologous to) $\sum_{i=1}^n \varepsilon(p_i) \mu_\Sigma$. This implies that $\sum_{i=1}^n \varepsilon(p_i) \mu_\Sigma = 0 \in H_1(N_\Sigma)$. But now, since the homology group $H_1(N_\Sigma) = \pi_1(N_\Sigma) = \mathbb{Z} \mu_\Sigma$ is torsion-free, we therefore deduce that $Q_N([\Sigma, \partial\Sigma], x) = \sum_{i=1}^n \varepsilon(p_i) = 0$, establishing that Σ is null-homologous. □

It follows easily that a closed \mathbb{Z} -surface $\Sigma \subseteq X$ is also null-homologous in $H_2(X)$. For the next lemma, recall that orientable 2-plane bundles over compact, orientable surfaces are classified up to isomorphism by their (relative) Euler number.

Lemma 5.2 *Every \mathbb{Z} -surface has trivial normal bundle. In the case of nonempty boundary this holds, moreover, relative to the Seifert framing on the boundary.*

Proof In the case that the boundary is nonempty, the surface is oriented and therefore the normal bundle is trivial. However, we need to show that this holds relative to the Seifert framing, and we need to show that the normal bundle is likewise trivial in the closed case. Continuing with the nonempty boundary case, cap off the \mathbb{Z} -surface Σ with a Seifert surface F in S^3 . Form a push-off by pushing F off itself in S^3 and extending this to a normal push-off in N along a generic section that agrees with the push-off of F along $\partial F = \partial \Sigma$.

Now the argument is the same in the closed case and in the nonempty boundary case (in the closed case take a push-off of Σ using a generic section of the normal bundle). Now in both cases, by the proof of Lemma 5.1, the push-off intersects Σ algebraically trivially. Therefore the normal bundle of Σ has vanishing Euler number, relative to the Seifert framing in the case of nonempty boundary. Thus the normal bundle of Σ is trivial, again relative to the Seifert framing in the case of nonempty boundary. \square

The boundary of the exterior of N_Σ is homeomorphic to $M_{K,g} := E_K \cup_\partial (\Sigma_{g,1} \times S^1)$, where $\Sigma_{g,1}$ is the orientable genus g surface with one boundary component and $E_K := S^3 \setminus \nu(K)$. We give more details on this identification as we will then make it implicitly throughout the remainder of the paper.

Remark 5.3 If $\Sigma \subseteq X$ is a closed oriented surface with $\pi_1(X_\Sigma) = \mathbb{Z}$, then the homotopy class of the meridian of $\Sigma \subseteq X$ is the unique nontrivial primitive class in $\pi_1(\partial X_\Sigma)$ that bounds a disc in $\Sigma \times D^2$ and maps to $1 \in \mathbb{Z}$. Fix a framing of the normal bundle of Σ , ie an identification $\nu \Sigma \cong \Sigma \times \mathbb{R}^2$ compatible with the orientation, with the property that for each simple closed curve $\gamma_k \subseteq \Sigma$, we have that $\gamma_k \times \{e_1\} \subseteq X \setminus \Sigma$ is null-homologous in $H_1(X \setminus \nu \Sigma) \cong \mathbb{Z}$. Use a choice of an identification $\Sigma \cong \Sigma_g$ and this condition to fix an identification of the boundary of the exterior with $\Sigma_g \times S^1$. Any two choices now differ by an element of the mapping class group of Σ_g . Similarly, if $\Sigma \subseteq N$ is a properly embedded, oriented surface, then we can identify ∂N_Σ with $E_K \cup_\partial (\Sigma_{g,1} \times S^1)$.

Lemma 5.4 *The exterior of a \mathbb{Z} -surface Σ has ribbon boundary.*

Proof Since N is simply connected, we have

$$\{1\} = \pi_1(N_\Sigma) *_{\pi_1(\Sigma \times S^1)} \pi_1(\Sigma \times D^2),$$

which implies that $\pi_1(\Sigma \times S^1) \rightarrow \pi_1(N_\Sigma)$ is onto. In fact, by the parametrisation of $\Sigma \times S^1$ described in Remark 5.3, we have that $\pi_1(\Sigma)$ maps trivially to $\pi_1(N_\Sigma)$, so the fundamental group of N_Σ is generated by a meridian of Σ . The closed case follows, since the exterior of a closed surface in X can be thought of as the exterior of a surface in $N = X \setminus \mathring{D}^4$. □

Lemma 5.5 *The boundary of the exterior of a \mathbb{Z} -surface Σ has Λ -torsion Alexander module. More precisely, the following hold.*

(1) *If $\Sigma \subseteq X$ is closed, then*

$$H_1(\partial X_\Sigma; \Lambda) = H_1(\Sigma_g \times S^1; \Lambda) = (\Lambda/(t-1))^{\oplus 2g}.$$

(2) *If $\Sigma \subseteq N$ has boundary a knot K , then the inclusion induces an isomorphism*

$$H_1(E_K; \Lambda) \oplus H_1(\Sigma_{g,1} \times S^1; \Lambda) \cong H_1(\partial N_\Sigma; \Lambda).$$

In particular, if K has Alexander polynomial one, then

$$H_1(\partial N_\Sigma; \Lambda) = (\Lambda/(t-1))^{\oplus 2g}.$$

Proof Using infinite cyclic covers, the assertion in the closed case is immediate:

$$\begin{aligned} H_1(\partial X_\Sigma; \Lambda) &= H_1(\Sigma_g \times S^1; \Lambda) = H_1(\widetilde{\Sigma_g \times S^1}; \mathbb{Z}) = H_1(\Sigma_g \times \mathbb{R}) \\ &= \mathbb{Z}^{2g} = (\Lambda/(t-1))^{\oplus 2g}. \end{aligned}$$

We prove the second assertion. Consider the Mayer-Vietoris sequence for the decomposition $M_{K,g} = E_K \cup_{\partial} (\Sigma_{g,1} \times S^1)$ with Λ coefficients,

$$\dots \rightarrow H_1(S^1 \times S^1; \Lambda) \rightarrow H_1(E_K; \Lambda) \oplus H_1(\Sigma_{g,1} \times S^1; \Lambda) \rightarrow H_1(M_{K,g}; \Lambda) \rightarrow 0.$$

We have

$$\begin{aligned} H_1(S^1 \times S^1; \Lambda) &= H_1(S^1 \times \mathbb{R}) = \mathbb{Z}, \\ H_1(\Sigma_{g,1} \times S^1; \Lambda) &= H_1(\Sigma_{g,1} \times \mathbb{R}) = \mathbb{Z}^{2g}. \end{aligned}$$

The map $H_1(S^1 \times S^1; \Lambda) \rightarrow H_1(E_K; \Lambda)$ is the zero map: it sends the generator to the lift of the longitude of E_K ; this is null-homologous in the infinite cyclic cover E_K^∞ of E_K . Next, the homomorphism $H_1(S^1 \times S^1; \Lambda) \rightarrow H_1(\Sigma_{g,1} \times S^1; \Lambda)$ is also the zero map: it coincides with the map $H_1(S^1) \rightarrow H_1(\Sigma_{g,1})$ sending the generator to $[\partial \Sigma_{g,1}]$, which vanishes in $H_1(\Sigma_{g,1})$. This establishes the second assertion. □

5.2 Isometries of the Blanchfield pairing

Given an orientable genus g surface $\Sigma_{g,1}$ with one boundary component, we determine the isometries of the Blanchfield pairing on $\Sigma_{g,1} \times S^1$ and show that they can be realised by orientation-preserving homeomorphisms. We then describe the Blanchfield pairing of

$$M_{K,g} := E_K \cup (\Sigma_{g,1} \times S^1),$$

where $E_K := S^3 \setminus \nu(K)$ is the knot exterior.

To describe the Blanchfield pairing $\text{Bl}_{\Sigma_{g,1} \times S^1}$, we fix some notation on the symplectic group:

$$J_g := \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}^{\oplus g}, \quad \text{Sp}_{2g}(\mathbb{Z}) := \{A \in M_{2g \times 2g}(\mathbb{Z}) \mid A^T J_g A = J_g\}.$$

Given a genus g surface $\Sigma_{g,1}$ with one boundary component, we call a basis of $H_1(\Sigma_{g,1})$ *symplectic* if the intersection form on $\Sigma_{g,1}$ with respect to this basis is represented by J_g . We also consider the map $\varphi: \pi_1(\Sigma_{g,1} \times S^1) \twoheadrightarrow \mathbb{Z}$ induced by the projection on the second coordinate and recall that

$$\text{Homeo}_\varphi^+(\Sigma_{g,1} \times S^1) = \{f \in \text{Homeo}^+(\Sigma_{g,1} \times S^1) \mid f_* \circ \varphi = \varphi\}.$$

The next proposition identifies the isometries of $\text{Bl}_{\Sigma_{g,1} \times S^1}$ with $\text{Sp}_{2g}(\mathbb{Z})$ and uses this fact to show that the orbit set $\text{Aut}(\text{Bl}_{\Sigma_{g,1} \times S^1})/\text{Homeo}_\varphi^+(\Sigma_{g,1} \times S^1)$ is trivial.

Proposition 5.6 *A choice of a symplectic basis for $H_1(\Sigma_{g,1})$ yields an identification*

$$\text{Aut}(\text{Bl}_{\Sigma_{g,1} \times S^1}) = \text{Sp}_{2g}(\mathbb{Z}).$$

Every element of the group $\text{Aut}(\text{Bl}_{\Sigma_{g,1} \times S^1})$ can be realised by a homeomorphism in $\text{Homeo}_\varphi^+(\Sigma_{g,1} \times S^1)$:

$$\text{Aut}(\text{Bl}_{\Sigma_{g,1} \times S^1})/\text{Homeo}_\varphi^+(\Sigma_{g,1} \times S^1) = \{\text{Id}\}.$$

Any such homeomorphism can be assumed to be of the form $j \times \text{Id}_{S^1}$ for some $j \in \text{Homeo}^+(\Sigma_{g,1})$ that fixes the boundary of $\Sigma_{g,1}$ pointwise.

Proof Let h be an isometry of the Blanchfield pairing $\text{Bl}_{\Sigma_{g,1} \times S^1}$. Since we saw in Lemma 5.5 that $H_1(\Sigma_{g,1} \times S^1; \Lambda) = H_1(\Sigma_{g,1})$, the isometry h can be thought of as a \mathbb{Z} -linear map. We assert that it preserves the intersection form on the surface $\Sigma_{g,1}$. Pick a symplectic basis for $H_1(\Sigma_{g,1})$ so that its intersection form is represented by J_g .

By [18, Corollary 1.2], the Blanchfield form on the fibred manifold $\Sigma_{g,1} \times S^1$ is given by

$$(v, w) \mapsto v^T (t^{-1} - 1)^{-1} J_g w \in \Lambda_S / \Lambda.$$

Here, $\Lambda_S = \mathbb{Z}[t^{\pm 1}, (t - 1)^{-1}]$ denotes the ring obtained from $\Lambda = \mathbb{Z}[t^{\pm 1}]$ by inverting $(t - 1)$. Since h can be thought of as a \mathbb{Z} -linear map, we represent it by a matrix H with coefficients in \mathbb{Z} . As h is an isometry, we have the following equalities of matrices with coefficients in Λ_S / Λ :

$$(t^{-1} - 1)^{-1} H^T J_g H = J_g (t^{-1} - 1)^{-1}.$$

Since H and J_g take values in \mathbb{Z} , the coefficients of the matrix $H^T J_g H - J_g$ take values in $(t - 1)\Lambda$. On the other hand, $H^T J_g H - J_g$ has coefficients in \mathbb{Z} , so $H^T J_g H - J_g = 0$, and we deduce that $H^T J_g H = J_g$. This shows that H is a symplectomorphism, proving the first assertion.

We prove the second assertion. We may realise any $h \in \text{Sp}_{2g}(\mathbb{Z})$ as the map on $H_1(\Sigma_{g,1})$ induced by an orientation-preserving homeomorphism $j: \Sigma_{g,1} \rightarrow \Sigma_{g,1}$ that fixes the boundary pointwise [13, Section 2.1 and the discussion following Theorem 6.4]. Cross j with the identity on S^1 to obtain an orientation-preserving homeomorphism $j \times \text{Id}: \Sigma_{g,1} \times S^1 \rightarrow \Sigma_{g,1} \times S^1$ that belongs to $\text{Homeo}_\varphi^+(\Sigma_{g,1} \times S^1)$. □

The next proposition describes the Blanchfield pairing of $M_{K,g} = E_K \cup (\Sigma_{g,1} \times S^1)$ and its automorphism group; here recall that $E_K = S^3 \setminus \nu(K)$ is the knot exterior.

Proposition 5.7 *For a knot $K \subseteq S^3$ there is an isometry $\text{Bl}_{M_{K,g}} \cong \text{Bl}_{E_K} \oplus \text{Bl}_{\Sigma_{g,1} \times S^1}$, and*

$$\text{Aut}(\text{Bl}_{M_{K,g}}) \cong \text{Aut}(\text{Bl}_{E_K}) \oplus \text{Aut}(\text{Bl}_{\Sigma_{g,1} \times S^1}).$$

Proof We proved in Lemma 5.5 that the inclusion induces a Λ -isomorphism

$$H_1(E_K; \Lambda) \oplus H_1(\Sigma_{g,1} \times S^1; \Lambda) \cong H_1(M_{K,g}; \Lambda).$$

The isometry $\text{Bl}_{M_{K,g}} \cong \text{Bl}_{E_K} \oplus \text{Bl}_{\Sigma_{g,1} \times S^1}$ now follows from [16, Theorem 1.1].

We must now show that $\text{Aut}(\text{Bl}_{E_K} \oplus \text{Bl}_{\Sigma_{g,1} \times S^1}) = \text{Aut}(\text{Bl}_{E_K}) \oplus \text{Aut}(\text{Bl}_{\Sigma_{g,1} \times S^1})$. The order of $H_1(M_K; \Lambda)$ is the Alexander polynomial Δ_K (which is not divisible by $(t - 1)$) and the order of $H_1(\Sigma_{g,1} \times S^1; \Lambda)$ is $(t - 1)^{2g}$. An automorphism h in $\text{Aut}(\text{Bl}_{E_K}) \oplus \text{Aut}(\text{Bl}_{\Sigma_{g,1} \times S^1})$ can be written as

$$h = \begin{pmatrix} h_{11} & h_{12} \\ h_{21} & h_{22} \end{pmatrix}.$$

Since the orders of $H_1(M_K; \Lambda)$ and $H_1(\Sigma_{g,1} \times S^1; \Lambda)$ are coprime, we deduce that $h_{12} = h_{21} = 0$. □

5.3 Surfaces in manifolds with boundary

The aim of this subsection is to prove Theorem 1.3 from the introduction. Given a knot $K \subseteq S^3$, thanks to Proposition 5.7, we can write automorphisms of $\text{Bl}_{M_{K,g}}$ as $h = h_K \oplus h_\Sigma$ with $h_K \in \text{Aut}(\text{Bl}_{E_K})$ and $h_\Sigma \in \text{Aut}(\text{Bl}_{\Sigma_{g,1} \times S^1})$.

Let us recall the construction of an isotopy, as introduced in Section 1.3. Suppose that $f_K : E_K \rightarrow E_K$ is an orientation-preserving homeomorphism that is the identity on ∂E_K . Extend f_K via the identity on νK to an orientation-preserving self-homeomorphism \tilde{f}_K of S^3 . The mapping class group of S^3 is trivial, so there is an isotopy $\Theta(f_K)_t : S^3 \rightarrow S^3$ between the extension and the identity, such that $\Theta(f_K)_0 = \text{Id}$ and $\Theta(f_K)_1 = \tilde{f}_K$. We can now prove Theorem 1.3.

Theorem 5.8 *Let $\Sigma_0, \Sigma_1 \subseteq N$ be two \mathbb{Z} -surfaces of genus g with boundary K . Suppose there is an isometry $F : \lambda_{N_{\Sigma_0}} \cong \lambda_{N_{\Sigma_1}}$, and write $\partial F = h_K \oplus h_\Sigma$.*

- *If h_K is induced by an orientation-preserving homeomorphism $f_K : E_K \rightarrow E_K$ that is the identity on ∂E_K , then f_K extends to an orientation-preserving homeomorphism of pairs*

$$(N, \Sigma_0) \xrightarrow{\cong} (N, \Sigma_1)$$

inducing the given isometry $F : H_2(N_{\Sigma_0}; \Lambda) \cong H_2(N_{\Sigma_1}; \Lambda)$.

- *If, in addition, $N = D^4$, then for any choice of isotopy $\Theta(f_K)$, the surfaces Σ_0 and Σ_1 are topologically ambiently isotopic via an ambient isotopy of D^4 extending $\Theta(f_K)$.*

Proof By assumption, the boundary of the isometry F is $\partial F = h_K \oplus h_\Sigma$ for some $h_K \in \text{Aut}(\text{Bl}_{E_K})$ and some $h_\Sigma \in \text{Aut}(\text{Bl}_{\Sigma_{g,1} \times S^1})$. By Proposition 5.6, we can realise h_Σ by an orientation-preserving homeomorphism of the form $j \times \text{Id}_{S^1}$, with $j : \Sigma_{g,1} \rightarrow \Sigma_{g,1}$ an orientation-preserving homeomorphism that fixes the boundary pointwise. Moreover, h_K is realised by an orientation-preserving homeomorphism $f_K : E_K \rightarrow E_K$ that fixes the boundary ∂E_K . Gluing these two homeomorphisms together, we obtain a homeomorphism

$$f := f_K \cup (j \times \text{Id}_{S^1}) : M_{K,g} \rightarrow M_{K,g}.$$

Since f is the identity on $\partial E_K \subseteq M_{K,g}$, we deduce that $f \in \text{Homeo}_\phi^+(M_{K,g})$. It follows that (f, F) is a compatible pair. The N_{Σ_i} are 4-manifolds whose boundaries are ribbon (recall Lemma 5.4) and have torsion Alexander modules for $i = 0, 1$ (by Lemma 5.5). Since $\Sigma_i \times D^2$ is a smooth manifold, $\text{ks}(\Sigma_i \times D^2) = 0$. Thus for $i = 0, 1$,

by additivity of ks under gluing we have $\text{ks}(N_{\Sigma_i}) = \text{ks}(N_{\Sigma_i}) + \text{ks}(\Sigma_i \times D^2) = \text{ks}(N)$. In particular, $\text{ks}(N_{\Sigma_0}) = \text{ks}(N_{\Sigma_1})$. We may therefore apply Theorem 4.1 to extend f to an orientation-preserving homeomorphism $\Phi: N_{\Sigma_0} \rightarrow N_{\Sigma_1}$ that induces F . As a consequence, we obtain the required homeomorphism of pairs:

$$\Phi' := \Phi \cup (j \times \text{Id}_{D^2}): (N, \Sigma_0) \rightarrow (N, \Sigma_1).$$

This establishes the first assertion.

For the second assertion, let $N = D^4$ and let $\Theta(f_K)_t: S^3 \rightarrow S^3$ be an isotopy of self-homeomorphisms of S^3 , with $\Theta(f_K)_0 = \text{Id}_{S^3}$ and $\Theta(f_K)_1$ equal to the extension \tilde{f}_K of f_K by the identity to all of S^3 . Cone $\Theta(f_K)$ to obtain an isotopy $C(\Theta(f_K)_t): D^4 \rightarrow D^4$ with $C(\Theta(f_K)_0) = \text{Id}_{D^4}$ and $C(\Theta(f_K)_1)$ equal to the cone $C(\tilde{f}_K)$ of \tilde{f}_K .

Next, note that $C(\tilde{f}_K)$ and $\Phi': D^4 \rightarrow D^4$ are two homeomorphisms of D^4 that restrict to the same homeomorphism of S^3 on the boundary. Therefore $C(\tilde{f}_K)^{-1} \circ \Phi': D^4 \rightarrow D^4$ is a homeomorphism restricting to Id_{S^3} on the boundary. By the Alexander trick, $C(\tilde{f}_K)^{-1} \circ \Phi'$ is isotopic rel boundary to the identity, via an isotopy $G_t: D^4 \rightarrow D^4$ with $G_0 = \text{Id}_{D^4}$ and $G_1 = C(\tilde{f}_K)^{-1} \circ \Phi'$. Note that $G_t|_{S^3} = \text{Id}_{S^3}$ for all $t \in [0, 1]$. Then let

$$H_t := C(\Theta(f_K)_t) \circ G_t: D^4 \rightarrow D^4.$$

This is an isotopy with

$$H_0 = \text{Id}_{D^4} \circ \text{Id}_{D^4} = \text{Id}_{D^4} \quad \text{and} \quad H_1 = C(\tilde{f}_K) \circ C(\tilde{f}_K)^{-1} \circ \Phi' = \Phi'.$$

In addition, for every $t \in [0, 1]$ we have that $H_t|_{S^3} = \Theta(f_K)_t \circ G_t|_{S^3} = \Theta(f_K)_t$. Since $\text{Id}_{D^4}(\Sigma_0) = \Sigma_0$ and $\Phi'(\Sigma_0) = \Sigma_1$, the isotopy H_t is a topological ambient isotopy between Σ_0 and Σ_1 , and it extends $\Theta(f_K)$, as required. \square

We also note the following corollary, in which we give a slightly relaxed criterion, but without precise control on the isotopy of K . It is sometimes easier to construct homeomorphisms of E_K that do not fix the boundary ∂E_K , but rather only fix the basepoint, so this could be a useful variation.

Corollary 5.9 *Let $K \subseteq S^3$ be a knot and fix a basepoint in ∂E_K . Suppose that $f: S^3 \rightarrow S^3$ is an orientation-preserving, basepoint-preserving homeomorphism with $f(K) = K$ as oriented knots, so also $f(E_K) = E_K$. Then f induces an isomorphism $h: H_1(E_K; \Lambda) \rightarrow H_1(E_K; \Lambda)$. Let Σ_0, Σ_1 be two \mathbb{Z} -surfaces in $N = X \setminus \mathring{D}^4$ with boundary $K \subseteq S^3 = \partial N$, and let $F: \lambda_{N_{\Sigma_0}} \cong \lambda_{N_{\Sigma_1}}$ be an isometry with $\partial F = h$. Then Σ_0 and Σ_1 are related by a homeomorphism of pairs, and if $N = D^4$, then they are ambiently isotopic.*

Proof Since f is orientation-preserving as a map from S^3 to S^3 , and as a map $K \rightarrow f(K)$, it restricts to an orientation-preserving map on ∂E_K that preserves the homotopy class of the zero-framed longitude and therefore preserves the orientation of the meridian. Thus f commutes with the map $\pi_1(E_K) \rightarrow \mathbb{Z}$ inducing the Λ -coefficients, and therefore induces a map $h: H_1(E_K; \Lambda) \rightarrow H_1(E_K; \Lambda)$ as claimed; recall Proposition 3.7. Let $\theta_t: S^3 \rightarrow S^3$ be an isotopy from the identity of S^3 to f . Let $S^3 \times [0, 1] \hookrightarrow N$ be a collar on the boundary of N with $S^3 \times \{0\}$ mapping to ∂N . Let $\Theta_t: N \rightarrow N$ be the isotopy that is the identity outside $S^3 \times [0, 1]$ and defined by $\Theta_t(x, s) = \theta_{t(1-s)}(x)$. This performs θ_t on the boundary, and tapers it in the collar. Let $\Sigma'_0 := \Theta_1(\Sigma_0)$, and note that Σ'_0 is ambiently isotopic (not necessarily rel boundary) to Σ_0 . In addition, $\partial(\Theta_1)_* = \partial F = h$ as isomorphisms of $H_1(E_K; \Lambda)$. Therefore $G := F \circ (\Theta_1)_*^{-1}: \lambda_{N_{\Sigma'_0}} \xrightarrow{\cong} \lambda_{N_{\Sigma_1}}$ is an isometry with $\partial G = \text{Id}$. By Theorem 1.3, Σ'_0 and Σ_1 are related by a homeomorphism of pairs restricting to the identity on the boundary, and if $N = D^4$ then they are ambiently isotopic rel boundary. The corollary now follows immediately. \square

5.4 Surfaces in closed manifolds

Now we recover the statements in the closed setting, proving Theorem 1.4 from the introduction. As discussed in the introduction, for a closed, simply connected 4-manifold X there is a classification of self-homeomorphisms of X in terms of their action on $H_2(X)$. The hypothesis of Theorem 1.4 is in terms of an isometry between the Λ -intersection forms of the surface exteriors X_{Σ_0} and X_{Σ_1} . In order to carefully state Theorem 1.4, first we need to understand the relationship between such an isometry and isomorphisms of the second homology $H_2(X)$ of the ambient 4-manifold.

Lemma 5.10 *Let $\Sigma_0, \Sigma_1 \subseteq X$ be closed \mathbb{Z} -surfaces of the same genus.*

- (i) *An isometry $F: \lambda_{X_{\Sigma_0}} \xrightarrow{\cong} \lambda_{X_{\Sigma_1}}$ induces an isomorphism $F_{\mathbb{Z}}: H_2(X) \xrightarrow{\cong} H_2(X)$.*
- (ii) *If the isometry F is induced by a homeomorphism $\Phi': (X, \Sigma_0) \rightarrow (X, \Sigma_1)$ of pairs, then $F_{\mathbb{Z}} = \Phi'_*$; in particular, $F_{\mathbb{Z}}$ is an isometry of the standard intersection form Q_X .*

Proof We claim that $H_2(X_{\Sigma_i}; \Lambda) \otimes_{\Lambda} \mathbb{Z} \cong H_2(X_{\Sigma_i})$. To prove this, we use the universal coefficient spectral sequence with second page $E_{p,q}^2 = \text{Tor}_p^{\Lambda}(H_q(X_{\Sigma_i}; \Lambda), \mathbb{Z})$ and which converges to $H_*(X_{\Sigma_i})$. The claim now follows promptly from the fact that $H_1(X_{\Sigma_i}; \Lambda) = 0$ and

$$\text{Tor}_2^{\Lambda}(H_0(X_{\Sigma_i}; \Lambda), \mathbb{Z}) = \text{Tor}_2^{\mathbb{Z}[\mathbb{Z}]}(\mathbb{Z}, \mathbb{Z}) = H_2(\mathbb{Z}; \mathbb{Z}) = H_2(S^1) = 0.$$

The claim implies that the intersection form on $H_2(X_{\Sigma_i})$ is obtained from the intersection form on $H_2(X_{\Sigma_i}; \Lambda)$ by tensoring down. Therefore F induces an isometry $F \otimes_{\Lambda} \text{Id}_{\mathbb{Z}}: H_2(X_{\Sigma_0}) \rightarrow H_2(X_{\Sigma_1})$. A Mayer–Vietoris argument yields the commutative diagram

$$(16) \quad \begin{array}{ccccccc} 0 & \longrightarrow & \mathbb{Z}^{2g} & \longrightarrow & H_2(X_{\Sigma_0}) & \xrightarrow{p_0} & H_2(X) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ & & (F \otimes_{\Lambda} \text{Id}_{\mathbb{Z}}) & & F \otimes_{\Lambda} \text{Id}_{\mathbb{Z}} & & F_{\mathbb{Z}} \\ 0 & \longrightarrow & \mathbb{Z}^{2g} & \longrightarrow & H_2(X_{\Sigma_1}) & \xrightarrow{p_1} & H_2(X) \longrightarrow 0 \end{array}$$

where the dotted map labelled $F_{\mathbb{Z}}$ is defined as the unique homomorphism that makes the right-hand square commute (both existence and uniqueness follow from a short diagram chase).

The second assertion is now immediate: the homeomorphism Φ' induces an isometry of $(H_2(X), Q_X)$ that satisfies $\Phi'_* \circ p_0 = p_1 \circ (F \otimes_{\Lambda} \text{Id}_{\mathbb{Z}})$, so $F_{\mathbb{Z}} = \Phi'_*$ is also an isometry. □

We move on to the proof of Theorem 1.4, whose statement we recall here for the benefit of the reader.

Theorem 5.11 *Let $\Sigma_0, \Sigma_1 \subseteq X$ be two closed \mathbb{Z} –surfaces of the same genus.*

- (i) *If the intersection forms $\lambda_{X_{\Sigma_0}}$ and $\lambda_{X_{\Sigma_1}}$ are isometric via an isometry F , then there is an orientation-preserving homeomorphism of pairs*

$$\Phi: (X, \Sigma_0) \xrightarrow{\cong} (X, \Sigma_1)$$

inducing the given isometry $\Phi_ = F: H_2(X_{\Sigma_0}; \Lambda) \xrightarrow{\cong} H_2(X_{\Sigma_1}; \Lambda)$.*

- (ii) *The isometry F induces an isometry $F_{\mathbb{Z}}: H_2(X) \rightarrow H_2(X)$ of the standard intersection form Q_X of X by Lemma 5.10. The surfaces Σ_0 and Σ_1 are topologically ambiently isotopic if and only if $F_{\mathbb{Z}} = \text{Id}$.*

Proof After an ambient isotopy, assume that Σ_0 and Σ_1 coincide on a disc $D^2 \subseteq \Sigma_0 \cap \Sigma_1$. Assume that the normal bundles also coincide over this D^2 . Consider the preimage $\mathring{D}^2 \times \mathbb{R}^2 \subseteq \nu \Sigma_i$. This is homeomorphic to an open 4–ball \mathring{D}^4 . Remove this $(\mathring{D}^4, \mathring{D}^2)$ from (X, Σ_i) to obtain $(N, \tilde{\Sigma}_i)$, with $\partial \tilde{\Sigma}_i = \Sigma_i \cap \partial N$ the unknot K in S^3 . Then the exterior of Σ_i in X equals the exterior of $\tilde{\Sigma}_i$ in N .

Since $X_{\Sigma_i} = N_{\tilde{\Sigma}_i}$, the Λ –intersection forms are unchanged and the isometry F also induces an isometry $F: \lambda_{N_{\tilde{\Sigma}_0}} \xrightarrow{\cong} \lambda_{N_{\tilde{\Sigma}_1}}$. Write $\partial F = h_K \oplus h_{\Sigma}$ as in Theorem 5.8.

The unknot $K \subseteq S^3$ has trivial Alexander module, so $h_K = \text{Id}$ and so h_K is realised by the homeomorphism $f_K = \text{Id}: E_K \rightarrow E_K$. Theorem 5.8 provides a rel boundary homeomorphism of pairs $\Phi': (N, \tilde{\Sigma}_0) \rightarrow (N, \tilde{\Sigma}_1)$ that induces the isometry F on the Λ -homology of the surface exteriors. We recover the required homeomorphism of pairs $\Phi: (X, \Sigma_0) \rightarrow (X, \Sigma_1)$ by gluing Φ' with the identity homeomorphism $(D^4, D^2) \rightarrow (D^4, D^2)$.

Lemma 5.10 implies that the isometry F induces an isometry $F_{\mathbb{Z}}$ of the standard intersection form Q_X . In particular, Lemma 5.10(ii) specifies that $F_{\mathbb{Z}}$ is induced by the homeomorphism Φ . If $F_{\mathbb{Z}} = \text{Id}$, then Φ is a self-homeomorphism of X inducing the identity on $H_2(X)$, so by [45, Theorem 1.1] and [15, Theorem 10.1], Φ is isotopic to the identity. It follows that Σ_0 and Σ_1 are topologically ambiently isotopic.

On the other hand, if Σ_0 and Σ_1 are topologically ambiently isotopic, then the induced homeomorphism between their exteriors extends to a homeomorphism from X to itself that is isotopic to the identity, and so certainly induces the identity map on $H_2(X)$. \square

6 Equivariant intersection forms of surface exteriors

The goal of this section is to collect some results about the intersection forms of \mathbb{Z} -surface exteriors. The main result of Section 6.1 shows that up to direct summands with the Hermitian form

$$\mathcal{H}_2 := \left(\Lambda^2, \begin{pmatrix} 0 & t-1 \\ t^{-1}-1 & 0 \end{pmatrix} \right),$$

the equivariant intersection forms of any two \mathbb{Z} -surface exteriors are isometric; this will be Proposition 6.3. Section 6.2 will then focus on the case of \mathbb{Z} -surfaces $\Sigma \subseteq N = X \setminus \overset{\circ}{D}^4$ with boundary an Alexander polynomial one knot $K \subseteq S^3$. The main result is Corollary 6.6 which shows that $\lambda_{N\Sigma}$ becomes isometric to $Q_X \oplus \mathcal{H}_2^{\oplus g}$ after adding sufficiently many copies of \mathcal{H}_2 , where Q_X is the \mathbb{Z} -valued intersection form of X (note that $Q_X = Q_N$).

6.1 Equivariant intersection forms of \mathbb{Z} -surface exteriors

The goal of this section is to study the Λ -intersection form of \mathbb{Z} -surface exteriors up to stabilisations by \mathcal{H}_2 .

We start by describing the Λ -intersection form of unknotted surfaces in S^4 .




Figure 1: Two handle diagrams for the exterior of a standardly embedded genus g surface $F \subseteq D^4$ with boundary the unknot in S^3 .

Lemma 6.1 *If $\Sigma \subseteq S^4$ is an unknotted oriented closed genus g surface, then the Λ -intersection form of S^4_Σ is isometric to $\mathcal{H}_2^{\oplus g}$.*

Proof We carry out the computation for the standardly embedded genus g surface $\Sigma \subseteq S^4$. Use $U \subseteq S^3$ to denote the unknot. Slice (S^4, Σ) along an equatorial (S^3, U) to obtain a decomposition $(S^4, \Sigma) = (D^4, \overset{\circ}{\Sigma}) \cup (D^4, D)$, with $D \subseteq D^4$ a disc bounding U and $\overset{\circ}{\Sigma} \subseteq D^4$ a punctured unknotted surface in D^4 .

Note that S^4_Σ and $D^4_{\Sigma^\circ} := D^4 \setminus \nu\overset{\circ}{\Sigma}$ are homeomorphic, since we can assume that the removed $\overset{\circ}{D}^4$ lies in the regular neighbourhood $\nu\Sigma$ removed from S^4 to form S^4_Σ . It follows that $H_2(S^4_\Sigma; \Lambda) = H_2(D^4_{\Sigma^\circ}; \Lambda)$ and that the Λ -intersection forms agree.

Thus it remains to compute the Λ -intersection form of the exterior D^4_F of a properly embedded unknotted surface $F \subseteq D^4$. A handle diagram with a single one-handle and $2g$ two-handles for D^4_F appears in the left-hand side of Figure 1, produced using the formalism of [19, Section 6.2]. It can then be isotoped as in the right-hand side of Figure 1, leading to a handle diagram for the infinite cyclic cover of D^4_F depicted in Figure 2. From this diagram, by taking the union of the cores of the 2-handles with null-homotopies of their attaching curves in the 4-ball, we obtain generators of $\pi_2(D^4_F) = H_2(D^4_F; \Lambda) = \Lambda^{2g}$. The Λ -intersection form can be computed via (equivariant) linking numbers, yielding the required result. □




Figure 2: A handle diagram for the infinite cyclic cover of D^4_F , where $F \subseteq D^4$ is an unknotted punctured surface.

We recall the concept of a 1–handle stabilisation for a surface in a 4–manifold. The following definition was motivated by [25]. Let $\Sigma \subseteq V$ be a locally flat (connected) surface embedded in a 4–manifold V . Let B be an embedding of D^4 into V such that ∂B intersects Σ transversely in a 2–component unlink L , and B intersects Σ transversely in two discs D_0 and D_1 , which can be simultaneously isotoped within B to lie in ∂B . Suppose that, for $i = 0, 1$, a 3–dimensional 1–handle $D^2 \times [0, 1]$ is embedded in the interior of B such that $D^2 \times \{i\} = D_i$. The resulting 1–handle stabilisation of Σ is defined as

$$\Sigma' = (\Sigma \cap (S^4 \setminus B)) \cup_L (S^1 \times [0, 1]).$$

If there exists a path γ in Σ between $(1, 0)$ and $(1, 1)$ in $S^1 \times [0, 1]$ such that $\gamma \cup (\{1\} \times [0, 1])$ is a null-homotopic loop in V , via a null-homotopy $h: D^2 \rightarrow V$ with $h(\partial D^2) \subseteq V \setminus \Sigma$, then we call the stabilisation *trivial*.

A detailed discussion of this construction in the locally flat setting can be found in [17, Proposition 9.1]. The next result describes the effect of trivial 1–handle stabilisation on the Λ –intersection form of surface exteriors.

Lemma 6.2 *If $\Sigma \subseteq N$ is a \mathbb{Z} –surface and $\Sigma' \subseteq N$ is obtained from Σ by a trivial 1–handle stabilisation, then*

$$\lambda_{N_{\Sigma'}} \cong \lambda_{N_{\Sigma}} \oplus \mathcal{H}_2.$$

Proof Since the 1–handle stabilisation is trivial, one can write

$$(N, \Sigma') = (N, \Sigma) \# (S^4, T^2),$$

where $T^2 \subseteq S^4$ denotes a standardly embedded torus, and $\#$ denotes the interior connect sum. It follows that $N_{\Sigma'} = N_{\Sigma} \cup S_{T^2}^4$, where the identification takes place along thickened meridians: $\mu_{\Sigma} \times D^2$ is identified with $\mu_{T^2} \times D^2$. One thus deduces the isomorphism $\pi_1(N_{\Sigma'}) \cong \pi_1(N_{\Sigma}) = \mathbb{Z}$. As the coefficient system maps these meridians to $1 \in \mathbb{Z}$, a straightforward Mayer–Vietoris argument shows that

$$H_2(N_{\Sigma'}; \Lambda) = H_2(N_{\Sigma}; \Lambda) \oplus H_2(S_{T^2}^4; \Lambda),$$

noting that $H_1(\mu_{\Sigma}; \Lambda) = 0$. It then follows that $\lambda_{N_{\Sigma'}} = \lambda_{N_{\Sigma}} \oplus \lambda_{S_{T^2}^4}$. The result is now a consequence of Lemma 6.1. □

The next proposition shows that the Λ –intersection forms of any two \mathbb{Z} –surfaces exteriors become isometric after adding sufficiently many \mathcal{H}_2 summands.

Proposition 6.3 For \mathbb{Z} -surfaces $\Sigma_0, \Sigma_1 \subseteq N$ of the same genus with boundary K , there exists an integer $n \geq 0$ and an isometry

$$\lambda_{N_{\Sigma_0}} \oplus \mathcal{H}_2^{\oplus n} \cong \lambda_{N_{\Sigma_1}} \oplus \mathcal{H}_2^{\oplus n}.$$

Proof Any two null-homologous surfaces in a 4-manifold can be made isotopic by enough 1-handle stabilisations to each surface [2, Theorem 5]. While Baykur and Sunukjian prove this result in the smooth category for surfaces in closed manifolds, it also applies in the topological category for properly embedded surfaces in 4-manifolds with boundary; see Theorem A.1 for a detailed proof in the case at hand.

We apply this to Σ_0, Σ_1 . This is possible because \mathbb{Z} -surfaces are null-homologous by Lemma 5.1. Since $\pi_1(\partial(\nu\Sigma_i)) \rightarrow \pi_1(N_{\Sigma_i}) = \mathbb{Z}$ is surjective, all 1-handle stabilisations can be taken to be trivial [2, Lemma 3]. We deduce that after sufficiently many (say n) trivial 1-handle stabilisations, Σ_0 becomes isotopic to Σ_1 , also stabilised n times. Using Lemma 6.2, each such stabilisation adds an \mathcal{H}_2 -summand to the Λ -intersection form of the corresponding surface exterior, from which the result follows. \square

6.2 \mathbb{Z} -surfaces for Alexander polynomial one knots

We now restrict our attention to \mathbb{Z} -surfaces with boundary Alexander polynomial one knots. An Alexander polynomial one knot $K \subseteq S^3$ bounds a disc $D \subseteq D^4$ with $\pi_1(D^4 \setminus D) = \mathbb{Z}$; see [15]. We call this disc a \mathbb{Z} -slice disc of K . When $S^3 = \partial N$ with $N = X \setminus \mathring{D}^4$, we arrange that this disc D belong to a collar neighbourhood $S^3 \times [0, 1] \subseteq N$ of $S^3 = \partial N$. We use this disc to build a genus g surface with boundary K .

Definition 6.4 A genus g target surface Σ_g^t for an Alexander polynomial one knot K is an embedded surface obtained from a \mathbb{Z} -disc D of K by g trivial 1-handle stabilisations.

Although we do not need this fact in the sequel, an Alexander polynomial one knot $K \subseteq S^3$ bounds a *unique* \mathbb{Z} -disc: this follows either from [10, Theorem 1.2] or from Theorem 5.8 above, because $H_2(D_D^4; \Lambda) = 0$. This is why we do not keep track of the \mathbb{Z} -disc D in the notation for target surfaces.

The next lemma describes the Λ -intersection form of a target surface exterior.

Lemma 6.5 *If $D \subseteq S^3 \times [0, 1] \subseteq N$ is a \mathbb{Z} -disc, then $\lambda_{N_D} = Q_X$. If Σ_g^t is a target surface, then $\lambda_{N_{\Sigma_g^t}} \cong Q_X \oplus \mathcal{H}_2^{\oplus g}$.*

Proof Since D is properly embedded in a collar neighbourhood of ∂N , N_D is homeomorphic to the interior connected sum $N_D \cong D_D^4 \# X$. The exterior D_D^4 of a \mathbb{Z} -slice disc is aspherical, and therefore we have $H_i(D_D^4; \Lambda) = 0$ for $i > 0$ [10, Lemma 2.1]. The first assertion now follows from a straightforward Mayer–Vietoris argument. Since a target surface is obtained from a \mathbb{Z} -slice disc by g handle additions, the second assertion follows from the first and Lemma 6.2. \square

Finally, we describe the Λ -intersection form of a \mathbb{Z} -surface exterior for an Alexander polynomial one knot K .

Corollary 6.6 *If $\Sigma \subseteq N = X \setminus \mathring{D}^4$ is a genus g \mathbb{Z} -surface for an Alexander polynomial one knot K , then $\lambda_{N_\Sigma} \oplus \mathcal{H}_2^{\oplus n} \cong Q_X \oplus \mathcal{H}_2^{\oplus (n+g)}$ for some integer $n \geq 0$.*

Proof Since K has Alexander polynomial one, it bounds a genus g target surface Σ_g^t . Proposition 6.3 ensures that $\lambda_{N_\Sigma} \oplus \mathcal{H}_2^{\oplus n}$ is isometric to $\lambda_{N_{\Sigma_g^t}} \oplus \mathcal{H}_2^{\oplus n}$ for some $n \geq 0$. The result now follows from Lemma 6.5, thanks to which $\lambda_{N_{\Sigma_g^t}} = Q_X \oplus \mathcal{H}_2^{\oplus g}$. \square

7 Knotted surfaces in S^4 and D^4

As we outlined in Section 1.4, our results about surfaces in D^4 and S^4 can be deduced from Theorem 5.8 once we prove that the surface exteriors have isometric equivariant intersection forms. In Section 7.1, we show that if $\Sigma \subseteq D^4$ is a genus $g \geq 3$ \mathbb{Z} -surface, with boundary an Alexander polynomial one knot K , then $\lambda_{D_\Sigma^4} \cong \mathcal{H}_2^{\oplus g}$. In Section 7.2 we then combine this result with Theorem 5.8 to deduce our results about surfaces in D^4 and S^4 , in particular proving Theorems 1.1 and 1.2. In Section 7.3, we apply these results to study Seifert surfaces pushed into D^4 .

7.1 Intersection forms in D^4

The aim of this subsection is to show that if $\Sigma \subseteq D^4$ is a \mathbb{Z} -surface of genus $g \geq 3$ with $\Delta_{\partial\Sigma} \doteq 1$, then $\lambda_{D_\Sigma^4} \cong \mathcal{H}_2^{\oplus g}$.

Before we continue, we need to recall some general theory on ε -Hermitian forms, and on their ε -quadratic counterparts; further details can be found in [47, Section 1.1]

and [12, Section 2], where the terminology “ ε -symmetric” is used, as is customary in L -theory. Let R be a ring with involution and let $\varepsilon \in R$ be a central unit with $\varepsilon\bar{\varepsilon} = 1$. Let M be a finitely generated free left R -module. Let $\text{Sesq}(M)$ denote the abelian group of R -sesquilinear forms on M , meaning that $b(rx, sy) = rb(x, y)\bar{s}$ for $b \in \text{Sesq}(M)$, $x, y \in M$, and $r, s \in R$. Define an involution

$$T_\varepsilon: \text{Sesq}(M) \rightarrow \text{Sesq}(M), \quad (T_\varepsilon b)(y, x) = \varepsilon \overline{b(x, y)}.$$

A quick check shows that $(T_\varepsilon^2 b)(x, y) = \varepsilon b(x, y)\bar{\varepsilon} = \varepsilon\bar{\varepsilon}b(x, y) = b(x, y)$, so the conditions that ε is central and that $\varepsilon\bar{\varepsilon} = 1$ are crucial for T_ε to be an involution. This enables us to define the *symmetric Q -group* $Q^\varepsilon(M)$ of M and the *quadratic Q -group* $Q_\varepsilon(M)$ of M via the exact sequence

$$0 \rightarrow Q^\varepsilon(M) \rightarrow \text{Sesq}(M) \xrightarrow{1-T_\varepsilon} \text{Sesq}(M) \rightarrow Q_\varepsilon(M) \rightarrow 0.$$

In other words,

$$Q^\varepsilon(M) := \ker(1 - T_\varepsilon) \quad \text{and} \quad Q_\varepsilon(M) := \text{coker}(1 - T_\varepsilon).$$

A sesquilinear form $b: M \times M \rightarrow R$ is called ε -Hermitian if it belongs to $Q^\varepsilon(M)$, that is if $b(y, x) = \varepsilon \overline{b(x, y)}$ for every $x, y \in M$. A $(+1)$ -Hermitian form is a Hermitian form in the usual sense, while a (-1) -Hermitian form is a skew-Hermitian form in the usual sense. A ε -Hermitian form is called *hyperbolic* if it is isometric to

$$H^\varepsilon(R)^{\oplus g} = \left(R^{2g}, \begin{pmatrix} 0 & 1 \\ \varepsilon & 0 \end{pmatrix}^{\oplus g} \right) \quad \text{for some } g \geq 0.$$

The reason for introducing this terminology is that we will shortly be concerned with $(-t)$ -Hermitian forms over Λ , and their quadratic analogues.

An element $\psi \in Q_\varepsilon(M)$ is called an ε -quadratic form. To an ε -quadratic form in $Q_\varepsilon(M)$ is associated its *symmetrisation* $\varphi := (1 + T_\varepsilon)(\psi) \in Q^\varepsilon(M)$. Given $\varphi \in Q^\varepsilon(M)$, a quadratic form ψ with $(1 + T_\varepsilon)(\psi) = \varphi$ is called a *quadratic refinement of φ* . The symmetrisation is well-defined on equivalence classes in $Q_\varepsilon(M)$ because $(1 + T_\varepsilon)(1 - T_\varepsilon) = 1 - T_\varepsilon^2 = 0$. Note that quadratic forms are considered up to addition of forms in the image of $1 - T_\varepsilon$. A quadratic form over R is *hyperbolic* if it is isometric to $H_\varepsilon(R)^{\oplus g} = \left(R^{2g}, \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}^{\oplus g} \right)$ for some $g \geq 0$. Let us emphasise that subscripts denote quadratic forms, while superscripts denote ε -Hermitian forms.

Here is the key relationship between $(-t)$ -Hermitian and $(-t)$ -quadratic forms that we shall exploit in the proof of Theorem 7.4 below.

Lemma 7.1 *Let M be a finitely generated free Λ -module. Then the map*

$$(1 + T_{-t}): Q_{-t}(M) \rightarrow Q^{-t}(M)$$

is injective.

Proof To prove the lemma we will show the following facts:

- (i) For $\varphi \in Q^{-t}(M)$, the assignment $\varphi \mapsto (t^{-1} - 1)\varphi$ determines a map

$$\Omega: Q^{-t}(M) \rightarrow Q^{+1}(M).$$

- (ii) For $\psi \in \text{Sesq}(M)$, the assignment $\psi \mapsto (t^{-1} - 1)\psi$ determines an injective map

$$\Xi: Q_{-t}(M) \rightarrow Q_{+1}(M).$$

- (iii) The map $(1 + T_{+1}): Q_{+1}(M) \rightarrow Q^{+1}(M)$ is injective.

- (iv) The diagram

$$\begin{array}{ccc} Q_{-t}(M) & \xrightarrow{\Xi} & Q_{+1}(M) \\ \downarrow 1+T_{-t} & & \downarrow 1+T_{+1} \\ Q^{-t}(M) & \xrightarrow{\Omega} & Q^{+1}(M) \end{array}$$

commutes.

Then, since the right-down composition is injective, it follows that the left vertical map is injective, as desired. So it suffices to prove these four assertions.

For (i), suppose that $\varphi \in Q^{-t}(M) = \ker(1 - T_{-t})$, that is $\varphi = -t\bar{\varphi}^T$. Then

$$\overline{(t^{-1} - 1)\varphi}^T = (t - 1)\bar{\varphi}^T = (t^{-1} - 1)(-t)\bar{\varphi}^T = (t^{-1} - 1)\varphi.$$

Therefore $\Omega(\varphi) = (t^{-1} - 1)\varphi \in \ker(1 - T_{+1}) = Q^{+1}(M)$, as desired.

To prove (ii), first we show that $\Xi: Q_{-t}(M) \rightarrow Q_{+1}(M)$ is well-defined. We need to see that for every sesquilinear form $\theta \in \text{Sesq}(M)$, the element $(1 - T_{-t})(\theta) = \theta - (-t)\bar{\theta}^T = \theta + t\bar{\theta}^T$ maps to the trivial element in $Q_{+1}(M)$. This is indeed the case since

$$(t^{-1} - 1)(\theta + t\bar{\theta}^T) = (t^{-1} - 1)\theta - (t - 1)\bar{\theta}^T = (1 - T_{+1})((t^{-1} - 1)\theta)$$

is trivial in $\text{coker}(1 - T_{+1}) = Q_{+1}(M)$. Therefore Ξ is well-defined.

Now we prove that $\Xi: Q_{-t}(M) \rightarrow Q_{+1}(M)$ is injective. Let B and C be two $(-t)$ -quadratic forms in $Q_{-t}(M)$, and suppose that $Z := (t^{-1} - 1)(B - C) = 0 \in Q_{+1}(M)$. That is, $Z \in \text{im}(1 - T_{+1})$, so, choosing a basis for $M \cong \Lambda^n$, there exists a matrix X over Λ such that

$$(t^{-1} - 1)(B - C) = Z = X - \bar{X}^T.$$

We assert that there exists a matrix Y over Λ such that

$$Z = (t^{-1} - 1)Y - (t - 1)\bar{Y}^T.$$

Write the entries of the matrix $Z = X - \bar{X}^T$ as z_{ij} . Since $X - \bar{X}^T$ is skew-Hermitian, we have $\bar{z}_{ij} = -z_{ji}$. Also, z_{ij} is divisible by $t^{-1} - 1$ for every i, j . For $i < j$, define the entries y_{ij} of Y by the formula $y_{ij} := z_{ij}/(t^{-1} - 1)$. For $i > j$ define $y_{ij} := 0$. This way $(t^{-1} - 1)y_{ij} - (t - 1)\bar{y}_{ji} = z_{ij}$ for $i \neq j$. It remains to define the diagonal entries of Y . For each i , since the Laurent polynomial z_{ii} satisfies $z_{ii} = -\bar{z}_{ii}$, we have $z_{ii}(-1) = -z_{ii}(-1)$, and so $z_{ii}(-1) = 0$. Therefore z_{ii} is divisible by $t + 1$. Since $t^{-1} - 1$ and $(t + 1)$ are coprime, we deduce that $z_{ii} = (t + 1)(t^{-1} - 1)q$ for some polynomial q that satisfies $q = \bar{q}$. Now set $y_{ii} := q$. We compute

$$(t^{-1} - 1)y_{ii} - (t - 1)\bar{y}_{ii} = (t^{-1} - 1 - (t - 1))q = (t^{-1} - t)q = (t + 1)(t^{-1} - 1)q = z_{ii}.$$

This concludes the proof of the assertion that $Z = (t^{-1} - 1)Y - (t - 1)\bar{Y}^T$ for some matrix Y . Using this assertion, as well as the definitions of X and Y , we obtain

$$Z = (t^{-1} - 1)(B - C) = X - \bar{X}^T = (t^{-1} - 1)Y - (t - 1)\bar{Y}^T = (t^{-1} - 1)(Y + t\bar{Y}^T).$$

This implies that $B - C = Y + t\bar{Y}^T = Y - (-t)\bar{Y}^T$, which is zero in $Q_{-t}(M)$. Therefore the map $\Xi: Q_{-t}(M) \rightarrow Q_{+1}(M)$ is injective, as desired. This completes the proof of (ii).

Next we prove (iii), that $1 + T_{+1}: Q_{+1}(M) \rightarrow Q^{+1}(M)$ is injective. For an arbitrary finitely generated free module M , the map $1 + T_\varepsilon$ fits into the exact sequence

$$0 \rightarrow \hat{Q}^{-\varepsilon}(M) \rightarrow Q_\varepsilon(M) \xrightarrow{1+T_\varepsilon} Q^\varepsilon(M).$$

The group $\hat{Q}^{-\varepsilon}(M) := \ker(1 + T_\varepsilon)$ is a *hyperquadratic Q -group*; its purpose is to measure the difference between Hermitian and quadratic forms [47]. We have

$$(17) \quad \hat{Q}^{-\varepsilon}(M) = \ker(1 + T_\varepsilon: Q_\varepsilon(M) \rightarrow Q^\varepsilon(M)) = \frac{\ker(1 + T_\varepsilon)}{\text{im}(1 - T_\varepsilon)} = \frac{\ker(1 - T_{-\varepsilon})}{\text{im}(1 + T_{-\varepsilon})}.$$

We shall use the penultimate description, but include the last equation to show why the $-\varepsilon$ appears in the notation for the group $\hat{Q}^{-\varepsilon}(M)$. Writing $M = \Lambda^n$, there is an isomorphism $\hat{Q}^{-\varepsilon}(M) = \hat{Q}^{-\varepsilon}(\Lambda^n) \cong \hat{Q}^{-\varepsilon}(\Lambda)^n$ [12, Remark 3.4]. By (17) with $\varepsilon = +1$ and $M = \Lambda$, we have $\hat{Q}^{-1}(\Lambda) = \ker(1 + T_{+1})/\text{im}(1 - T_{+1})$. For $p(t) \in \Lambda$, if $p(t) + p(t^{-1}) = 0$, then $p(t)$ must be of the form $p(t) = r(t) - r(t^{-1})$ for some $r(t) \in \Lambda$. It follows that $\hat{Q}^{-1}(\Lambda) = 0$. Thus $\ker(1 + T_{+1}: Q_{+1}(M) \rightarrow Q^{+1}(M)) = 0$ and so $1 + T_{+1}$ is indeed injective. This concludes the proof of the third item of the lemma.

For (iv), let $\psi \in Q_{-t}(M)$. Then

$$\begin{aligned} (1 + T_{+1}) \circ \Xi(\psi) &= (1 + T_{+1})(t^{-1} - 1)\psi = (t^{-1} - 1)\psi + (t - 1)\bar{\psi}^T \\ &= (t^{-1} - 1)(\psi - t\bar{\psi}^T) = \Omega \circ (1 + T_{-t})(\psi), \end{aligned}$$

so the diagram commutes, which proves (iv) and completes the proof that the map $(1 + T_{-t}): Q_{-t}(M) \rightarrow Q^{-t}(M)$ is injective. □

Next we state the algebraic cancellation result that we will use. The idea behind the proof of Theorem 7.4 below will be to engineer a situation in which we can apply cancellation.

Definition 7.2 The Witt index $\text{ind}(H, \theta)$ of an ε -quadratic form is the largest integer k such that a subform of (H, θ) is isometric to $H_\varepsilon(R)^{\oplus k}$.

Proposition 7.3 Let $\varepsilon \in \Lambda$ be a central unit with $\varepsilon\bar{\varepsilon} = 1$, and let (H, θ) and (H', θ') be ε -quadratic forms over Λ . Assume that for some $n \geq 0$ there is an isometry

$$(18) \quad (H, \theta) \oplus H_\varepsilon(\Lambda)^{\oplus n} \cong (H', \theta') \oplus H_\varepsilon(\Lambda)^{\oplus n}.$$

If $\text{ind}(H', \theta') \geq 3$, then there is an isometry

$$(19) \quad (H, \theta) \cong (H', \theta').$$

Proof We apply a result due to Bass [1, Corollary IV.3.6]. Given a ring R with involution, we write $R_0 = \{\sum_i x_i \bar{x}_i \mid x_i \in R\}$ for the norm subring of R , as well as $\text{maxspec}(R_0)$ for the set of all maximal ideals of R_0 under the Zariski topology, and $d_R := \dim \text{maxspec}(R_0)$. Detailed definitions of these notions are irrelevant: we need only know that $d_\Lambda = 2$; see eg [29, Proposition 2.2] or [21, page 439]. Let (H, θ) and (H', θ') be quadratic forms over R . If there is an isometry

$$(H, \theta) \oplus H_\varepsilon(R)^{\oplus n} \cong (H', \theta') \oplus H_\varepsilon(R)^{\oplus n},$$

and if $\text{ind}((H', \theta') \oplus H_\varepsilon(R)) \geq d_R + 2$, then there is an isometry $(H, \theta) \cong (H', \theta')$ [1, Corollary IV.3.6]. In particular, this cancellation result holds if $\text{ind}(H', \theta') \geq d_R + 1$. As mentioned above, for $R = \Lambda$, we have $d_\Lambda = 2$. Since

$$\text{ind}((H', \theta') \oplus H_\varepsilon(\Lambda)) \geq 3 + 1 = d_\Lambda + 2,$$

the result of Bass says that (18) implies (19). □

The next theorem is the main result of this subsection.

Theorem 7.4 *If $\Sigma \subseteq D^4$ is a \mathbb{Z} -surface of genus $g \geq 3$ whose boundary is an Alexander polynomial one knot $K \subseteq S^3$, then*

$$\lambda_{D^4_\Sigma} \cong \mathcal{H}_2^{\oplus g}.$$

Proof We outline the strategy of the proof. Since the knot K has Alexander polynomial one and $Q_{D^4} = 0$, by Corollary 6.6, there exists an integer $n \geq 0$ and an isometry

$$(20) \quad \lambda_{D^4_\Sigma} \oplus \mathcal{H}_2^{\oplus n} \cong \mathcal{H}_2^{\oplus g+n}.$$

We cannot apply the cancellation result of Proposition 7.3 directly, since $\mathcal{H}_2^{\oplus n}$ is not hyperbolic over Λ . However, by switching the basis elements \mathcal{H}_2 is isometric to a multiple of the $(-t)$ -hyperbolic form:

$$\begin{pmatrix} 0 & t^{-1} - 1 \\ t - 1 & 0 \end{pmatrix} = (t^{-1} - 1) \cdot \begin{pmatrix} 0 & 1 \\ -t & 0 \end{pmatrix} = (t^{-1} - 1)\mathcal{H}^{-t}(\Lambda).$$

If we could remove the $(t^{-1} - 1)$ factor, we could apply Proposition 7.3. The idea for this is to use the relative intersection pairing

$$\lambda_{D^4_\Sigma}^\partial : H_2(D^4_\Sigma; \Lambda) \times H_2(D^4_\Sigma, \partial D^4_\Sigma; \Lambda) \rightarrow \Lambda$$

instead of the ‘‘absolute’’ intersection pairing $\lambda_{D^4_\Sigma}$. We will see, using (20) and appropriate bases, that the pairing $\lambda_{D^4_\Sigma}^\partial$ is represented by a $(-t)$ -Hermitian matrix over Λ . We will then apply the cancellation result of Proposition 7.3 with $\varepsilon = -t$ to these forms, before deducing the desired conclusion on the original absolute pairings. Note that the adjoints of the relative and absolute pairings fit into the commutative diagram

$$(21) \quad \begin{array}{ccc} H_2(D^4_\Sigma; \Lambda) & \xrightarrow{\hat{\lambda}} & \overline{\text{Hom}_\Lambda(H_2(D^4_\Sigma; \Lambda), \Lambda)} \\ & \searrow q & \nearrow \hat{\lambda}^\partial \\ & & H_2(D^4_\Sigma, \partial D^4_\Sigma; \Lambda) \end{array}$$

For the first step of the proof we will choose bases for $H_2(D^4_\Sigma; \Lambda)$ and $H_2(D^4_\Sigma, \partial D^4_\Sigma; \Lambda)$ and describe the map q with respect to these bases. First, a short rank computation:

Claim 4 $H_2(D^4_\Sigma; \Lambda) \cong \Lambda^{2g}.$

Proof Since $H_2(D^4_\Sigma; \Lambda)$ is free (by Lemma 3.2), we need only show that its rank is $b_1(\Sigma) = 2g$. Using $H_1(D^4_\Sigma) = \mathbb{Z}$, a Mayer–Vietoris argument shows that $b_2(D^4_\Sigma) = b_1(\Sigma)$ as well as $b_3(D^4_\Sigma) = 0$. Since we also have $b_4(D^4_\Sigma) = 0$ and $b_0(D^4_\Sigma) = 1$,

we get $\chi(D_\Sigma^4) = b_1(\Sigma)$. The Euler characteristic can also be computed with $Q = \mathbb{Q}(t)$ coefficients. By Lemma 3.2, we have $b_i^Q(D_\Sigma^4) = 0$ for $i \neq 2$, and therefore

$$\text{rk}_\Lambda H_2(D_\Sigma^4; \Lambda) = \chi^Q(D_\Sigma^4) = b_1(\Sigma),$$

as asserted. □

Claim 5 *There is a choice of bases with respect to which the homomorphism*

$$q: H_2(D_\Sigma^4; \Lambda) \rightarrow H_2(D_\Sigma^4, \partial D_\Sigma^4; \Lambda)$$

is represented by the matrix $(t - 1)$ Id.

Proof We first base $H_2(D_\Sigma^4, \partial D_\Sigma^4; \Lambda)$ using the fact that the Alexander polynomial of K is one. We have an exact sequence as part of the long exact sequence of the pair $(D_\Sigma^4, \partial D_\Sigma^4)$:

$$H_2(D_\Sigma^4; \Lambda) \xrightarrow{q} H_2(D_\Sigma^4, \partial D_\Sigma^4; \Lambda) \xrightarrow{\delta} H_1(\partial D_\Sigma^4; \Lambda) \rightarrow 0.$$

Here $H_1(D_\Sigma^4; \Lambda) = 0$ because $\pi_1(D_\Sigma^4) \cong \mathbb{Z}$. In particular, the map q presents the Λ -module $H_1(\partial D_\Sigma^4; \Lambda)$. Use Lemmas 3.2 and 5.5, Claim 4 and the fact that $\Delta_K \doteq 1$ to deduce that

$$H_2(D_\Sigma^4; \Lambda) \cong \Lambda^{2g} \cong H_2(D_\Sigma^4, \partial D_\Sigma^4; \Lambda) \quad \text{and} \quad H_1(\partial D_\Sigma^4; \Lambda) \cong \bigoplus_{i=1}^{2g} \Lambda/(t-1).$$

Choose a set of generators $\{\gamma_k\} \subseteq H_1(\partial D_\Sigma^4; \Lambda)$, represented in the form

$$g_k \times \{\text{pt}\} \subseteq \Sigma \times \{\text{pt}\} \subseteq \Sigma \times S^1 \subseteq \partial D_\Sigma^4,$$

where $\{g_k\}$ is a symplectic basis of curves on Σ . Then choose a basis $\{x_i\}$ for $H_2(D_\Sigma^4, \partial D_\Sigma^4; \Lambda)$ such that $\delta(x_i) = \gamma_i$ for $1 \leq i \leq 2g$. The homology classes x_k can be represented by embedded surfaces $S_i \subseteq D_\Sigma^4$ with $\partial S_i = \gamma_i$ for $1 \leq i \leq 2g$ [17, Section 10.3].

Next, we base $H_2(D_\Sigma^4; \Lambda)$. For each $i = 1, \dots, 2g$, define a closed surface representing a class $y_i \in H_2(D_\Sigma^4; \Lambda)$, as follows. Consider the torus $g_i \times S^1 = \partial(\bar{\nu}_\Sigma|_{g_i})$, and surger it along γ_i using the surface S_i . That is, remove an annular neighbourhood $g_i \times (p, p') \subseteq g_i \times S^1$ of $\gamma_i = g_i \times \{\text{pt}\}$, leaving $B := g_i \times (S^1 \setminus (p, p'))$. Then glue $-S_i$ to $g_i \times \{p\}$, and a push-off S'_i of S_i to $g_i \times \{p'\}$. Call the resulting surface T_i , and set $y_i := [T_i] \in H_2(D_\Sigma^4; \Lambda)$ for $1 \leq i \leq 2g$. Then, for $i = 1, \dots, 2g$, in $H_2(D_\Sigma^4, \partial D_\Sigma^4; \Lambda)$ we have

$$(22) \quad q(y_i) = q([T_i]) = [-S_i] + [B \cup S'_i] = -x_i + tx_i = (t - 1)x_i.$$




Figure 3: Representing $y_i \in H_2(D_\Sigma^4; \Lambda)$ by an immersed surface T_i in D_Σ^4 obtained by surgering the torus $g_i \times S^1$, using a surface S_i representing the relative homology class x_i . The top left picture shows a neighbourhood of a portion of g_i , intersected with a carefully chosen 3-dimensional subspace that contains the intersections of Σ and S_i with this neighbourhood. Bottom, a 4-dimensional picture of T_i , consisting of three 3-dimensional slices at certain special values of the fourth coordinate (time), in which Σ , $-S_i$, S'_i and most of B appear. The annulus B joins $-S_i$ and S'_i together; in intermediate time values only a line of B appears, including in the middle slice. The top right picture shows a cross section of T_i , looking along g_i .

Here the t arises because the annulus B wraps around a meridian of Σ . This process is illustrated in Figure 3.

Consider the diagram of short exact sequences

$$\begin{array}{ccccccc}
 0 & \longrightarrow & \Lambda^{\oplus 2g} & \xrightarrow{(t-1)\text{Id}} & \Lambda^{\oplus 2g} & \xrightarrow{\text{proj}} & (\Lambda/(t-1))^{\oplus 2g} \longrightarrow 0 \\
 & & \downarrow y & & \downarrow x & & \downarrow \gamma \\
 0 & \longrightarrow & H_2(D_\Sigma^4; \Lambda) & \xrightarrow{q} & H_2(D_\Sigma^4, \partial D_\Sigma^4; \Lambda) & \longrightarrow & H_1(\partial D_\Sigma^4; \Lambda) \longrightarrow 0
 \end{array}$$

where the maps y and x send the canonical basis $\{e_i\}_{i=1}^{2g}$ of Λ^{2g} to $\{y_i\}_{i=1}^{2g}$ and $\{x_i\}_{i=1}^{2g}$, respectively, and γ maps the $[e_i]$ to the γ_i . This diagram commutes thanks to (22). By construction x and γ are isomorphisms, and therefore so is y by the five lemma. Thus $\{y_i\}_{i=1}^{2g}$ forms a basis of $H_2(D_\Sigma^4; \Lambda)$ and q is represented by $(t-1)\text{Id}$. This completes the proof of Claim 5. □

Recall from (20) that there is an isometry $\alpha: \lambda_{D_\Sigma^4} \oplus \mathcal{H}_2^{\oplus n} \cong \mathcal{H}_2^{\oplus(g+n)}$. Denote the matrices of $\lambda_{D_\Sigma^4}$ and $\lambda_{D_\Sigma^4}^\partial$ with respect to the previously described bases by L and L^∂ , respectively. Also represent the isometry α by a matrix A with respect to these bases. Therefore we have a congruence

$$(23) \quad A^T \left(L \oplus \begin{pmatrix} 0 & t-1 \\ t^{-1}-1 & 0 \end{pmatrix}^{\oplus n} \right) \bar{A} = \begin{pmatrix} 0 & t-1 \\ t^{-1}-1 & 0 \end{pmatrix}^{\oplus(g+n)}.$$

Our aim is now to factor out a $(t^{-1} - 1)$ from both sides of this equation. Recall from (21) that the adjoints of the pairings $\lambda_{D_\Sigma^4}$ and $\lambda_{D_\Sigma^4}^\partial$ are related by $\widehat{\lambda}_{D_\Sigma^4}^\partial \circ q = \widehat{\lambda}_{D_\Sigma^4}$. With respect to our bases (and recalling (22)), for every $x, y \in H_2(D_\Sigma^4; \Lambda)$ we therefore obtain

$$\begin{aligned} x^T L^\partial (t^{-1} - 1) \bar{y} &= x^T L^\partial \overline{(t^{-1} - 1)y} = \lambda_{D_\Sigma^4}^\partial(x, q(y)) = \widehat{\lambda}^\partial(q(y))(x) \\ &= \widehat{\lambda}(y)(x) = \lambda(x, y) \\ &= x^T L \bar{y}. \end{aligned}$$

It follows that $L^\partial(t^{-1} - 1) = L$. Combine this with

$$\mathcal{H}_2 = \begin{pmatrix} 0 & t-1 \\ t^{-1}-1 & 0 \end{pmatrix} = \begin{pmatrix} 0 & -t \\ 1 & 0 \end{pmatrix} (t^{-1} - 1)$$

to rewrite (23) as

$$A^T \left(L^\partial (t^{-1} - 1) \oplus \begin{pmatrix} 0 & -t \\ 1 & 0 \end{pmatrix} (t^{-1} - 1) \right) \bar{A} = \begin{pmatrix} 0 & -t \\ 1 & 0 \end{pmatrix}^{\oplus(g+n)} (t^{-1} - 1).$$

As $(t^{-1} - 1) \cdot \text{Id}$ is diagonal with constant diagonal coefficients, it is central and thus we obtain

$$(t^{-1} - 1) \cdot A^T \left(L^\partial \oplus \begin{pmatrix} 0 & -t \\ 1 & 0 \end{pmatrix}^{\oplus n} \right) \bar{A} = (t^{-1} - 1) \begin{pmatrix} 0 & -t \\ 1 & 0 \end{pmatrix}^{\oplus(g+n)}.$$

As $(t^{-1} - 1) \cdot \text{Id}$ is nondegenerate over Λ , it follows that

$$(24) \quad A^T \left(L^\partial \oplus \begin{pmatrix} 0 & -t \\ 1 & 0 \end{pmatrix}^{\oplus n} \right) \bar{A} = \begin{pmatrix} 0 & -t \\ 1 & 0 \end{pmatrix}^{\oplus(g+n)}.$$

In particular L^∂ stabilises, via $(-t)$ -Hermitian matrices, to a $(-t)$ -Hermitian form, so it follows that L^∂ is itself $(-t)$ -Hermitian.

In order to apply cancellation, we refine (24) to a statement about quadratic forms. Set

$$\Psi := \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}.$$

Then $(1 + T_{-t})\Psi = \begin{pmatrix} 0 & -t \\ 1 & 0 \end{pmatrix}$, so Ψ is a quadratic refinement for $\begin{pmatrix} 0 & -t \\ 1 & 0 \end{pmatrix}$. Therefore $\Psi^{\oplus(g+n)}$ is a quadratic refinement for $\begin{pmatrix} 0 & -t \\ 1 & 0 \end{pmatrix}^{\oplus(g+n)}$, which is the right-hand side of (24). Thus $\Psi^{\oplus(g+n)}$ is a quadratic refinement for

$$A^T \left(L^\partial \oplus \begin{pmatrix} 0 & -t \\ 1 & 0 \end{pmatrix}^{\oplus n} \right) \bar{A}.$$

Rearranging, we deduce that $A^{-T} \Psi^{\oplus(g+n)} \bar{A}^{-1}$ is a quadratic refinement for the $(-t)$ -Hermitian matrix $L^\partial \oplus \begin{pmatrix} 0 & -t \\ 1 & 0 \end{pmatrix}^{\oplus n}$. Now, corresponding to this decomposition, write

$$A^{-T} \Psi^{\oplus(g+n)} \bar{A}^{-1} = \begin{pmatrix} \Psi^\partial & U \\ V & \Psi_{\text{ind}} \end{pmatrix},$$

where this defines the block entries on the right-hand side. Since this is a quadratic refinement of the block diagonal $(-t)$ -Hermitian matrix

$$L^\partial \oplus \begin{pmatrix} 0 & -t \\ 1 & 0 \end{pmatrix}^{\oplus n},$$

we deduce that $V - t\bar{U}^T = 0$ and therefore

$$(25) \quad A^{-T} \Psi^{\oplus(g+n)} \bar{A}^{-1} = \begin{pmatrix} \Psi^\partial & 0 \\ 0 & \Psi_{\text{ind}} \end{pmatrix} + \begin{pmatrix} 0 & U \\ t\bar{U}^T & 0 \end{pmatrix} = \Psi^\partial \oplus \Psi_{\text{ind}} \in Q_{-t}(\Lambda^{\oplus 2(g+n)}).$$

This procedure therefore endows both the $(-t)$ -Hermitian matrices L^∂ and $\begin{pmatrix} 0 & -t \\ 1 & 0 \end{pmatrix}^{\oplus n}$ with quadratic refinements, which we denoted by Ψ^∂ and Ψ_{ind} respectively. Rearranging (25) we obtain the following isometry of quadratic forms:

$$(26) \quad A^T (\Psi^\partial \oplus \Psi_{\text{ind}}) \bar{A} = \Psi^{\oplus(g+n)} \in Q_{-t}(\Lambda^{\oplus 2(g+n)}).$$

In order to apply cancellation, we need Ψ_{ind} to agree with the quadratic refinement $\Psi^{\oplus n}$ of $\begin{pmatrix} 0 & -t \\ 1 & 0 \end{pmatrix}^{\oplus n}$ in the Q -group $Q_{-t}(\Lambda^{2n})$. The fact that both are quadratic refinements of the form $\begin{pmatrix} 0 & -t \\ 1 & 0 \end{pmatrix}^{\oplus n}$ in $Q^{-t}(\Lambda^{2n})$ can be restated as

$$(1 + T_{-t})(\Psi_{\text{ind}}) = (1 + T_{-t})(\Psi^{\oplus n}) \in Q^{-t}(\Lambda^{2n}).$$

By Lemma 7.1, $(1 + T_{-t})$ is injective, and so $\Psi_{\text{ind}} = \Psi^{\oplus n}$ in $Q_{-t}(\Lambda^{2n})$, as desired.

We deduce from (26) that

$$(27) \quad A^T (\Psi^\partial \oplus \Psi^{\oplus n}) \bar{A} = \Psi^{\oplus(g+n)} \in Q_{-t}(\Lambda^{2(g+n)}).$$

Since

$$\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix},$$

we see that $\begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$ is congruent to the hyperbolic $(-t)$ -quadratic form $H_{-t}(\Lambda)$. We have established the following isometry of $(-t)$ -quadratic forms:

$$(28) \quad \Psi^\partial \oplus H_{-t}(\Lambda)^{\oplus n} \cong H_{-t}(\Lambda)^{\oplus g} \oplus H_{-t}(\Lambda)^{\oplus n} \in \mathcal{Q}_{-t}(\Lambda^{2(g+n)}).$$

Since $g \geq 3$, we now apply the cancellation result of Proposition 7.3 to (28) and symmetrise the result to deduce that L^∂ is congruent to $H^{-t}(\Lambda)^{\oplus g}$, and therefore to $\begin{pmatrix} 0 & -t \\ 1 & 0 \end{pmatrix}^{\oplus g}$. Multiplying both sides of this congruence by $t^{-1} - 1$ shows that L is congruent to $\mathcal{H}_2^{\oplus g}$. This concludes the proof of Theorem 7.4. \square

7.2 Surfaces in S^4 and D^4

We collect and prove our results concerning (rel boundary) ambient isotopy of surfaces in D^4 and S^4 . Since the proofs of these results were already outlined in the introduction or elsewhere, we proceed swiftly.

Theorem 7.5 *Let $\Sigma_0, \Sigma_1 \subseteq D^4$ be \mathbb{Z} -surfaces of genus g with boundary K .*

- (i) *Suppose there is an isometry $F \in \text{Iso}(\lambda_{D_{\Sigma_0}^4}, \lambda_{D_{\Sigma_1}^4})$ and write $\partial F = h_K \oplus h_\Sigma$. If h_K is represented by an orientation-preserving homeomorphism that is the identity on ∂E_K , then Σ_0 and Σ_1 are topologically ambiently isotopic rel boundary.*
- (ii) *If $\Delta_K = 1$, $g = 1, 2$ and $\lambda_{D_{\Sigma_0}^4} \cong \lambda_{D_{\Sigma_1}^4}$, then Σ_0 and Σ_1 are topologically ambiently isotopic rel boundary.*
- (iii) *If $\Delta_K = 1$ and $g \neq 1, 2$, then Σ_0 and Σ_1 are topologically ambiently isotopic rel boundary.*

Proof The first assertion is the second item of Theorem 5.8. When $\Delta_K = 1$, we have $H_1(E_K; \Lambda) = 0$, so $h_K = \text{Id}$ is automatic, and the second assertion follows from the first. For the third assertion, additionally use that $\lambda_{D_{\Sigma_0}^4} \cong \lambda_{D_{\Sigma_1}^4}$ when $g \neq 1, 2$ by Theorem 7.4. \square

Theorem 7.6 *Let $\Sigma_0, \Sigma_1 \subseteq S^4$ be closed \mathbb{Z} -surfaces of genus g .*

- (i) *For $g = 1, 2$, if $\lambda_{\Sigma_0} \cong \lambda_{\Sigma_1}$, then Σ_0 and Σ_1 are topologically ambiently isotopic.*
- (ii) *For $g \neq 1, 2$, the surfaces Σ_0 and Σ_1 are topologically ambiently isotopic.*

Proof Both assertions follow from Theorem 7.5 by removing a (D^4, D^2) -pair from (S^4, Σ_i) . \square

7.3 Pushed-in Seifert surfaces for Alexander polynomial one knots

We prove Theorem 1.9 from the introduction, which states that any two pushed-in Seifert surfaces of the same genus for an Alexander polynomial one knot are topologically ambiently isotopic in D^4 . Recall that a *sublagrangian* for a nonsingular ε -Hermitian form (H, λ) is a direct summand $M \subseteq H$ such that $\lambda(M \times M) = 0$, ie $M \subseteq M^\perp$. A sublagrangian is a *Lagrangian* if $M = M^\perp$. For a nonsingular ε -quadratic form (H, λ, μ) , the definitions are identical, with the additional requirement that $\mu(M) = 0$. The following lemma is known to those familiar with L-theory. For instance, given a sublagrangian $i : M \hookrightarrow H$, the result follows quite promptly from [46, Proposition 2.2], provided one assumes that $i^*\hat{\lambda} : H \rightarrow M^*$ is surjective. Since our argument is elementary (Ranicki’s proof is more involved because his statement is more general) and since most of our work goes into establishing that $i^*\hat{\lambda}$ is surjective, we include a proof for the readers’ convenience.

Lemma 7.7 *Let (H, λ) be a nonsingular ε -Hermitian form on a free Λ -module H . If $M \subseteq H$ is a half-rank sublagrangian, then M is a Lagrangian.*

Proof We know that $M \subseteq M^\perp$ and must show that $M = M^\perp$. Equivalently, we must show that $M/M^\perp = 0$. We will show that M/M^\perp is both torsion and torsion-free.

Claim 6 *The inclusion $M^\perp \hookrightarrow H$ is split and $M^\perp \subseteq H$ is a half-rank summand.*

Proof Consider the exact sequence $0 \rightarrow M^\perp \rightarrow H \xrightarrow{i^*\hat{\lambda}} M^*$. It now suffices to show that $i^*\hat{\lambda}$ is surjective: since M^* is free and $\text{rk}_\Lambda(M^*) = \text{rk}_\Lambda(M) = \frac{1}{2} \text{rk}_\Lambda(H)$, the result would then follow. Since λ is nonsingular, $\hat{\lambda}$ is surjective and so it suffices to prove that i^* is injective. This occurs if and only if $\text{Ext}_\Lambda^1(H/M, \Lambda) = 0$. But since M is a summand, it follows that $H \cong M \oplus H/M$. Since H is free, H/M is projective over Λ and therefore free over Λ [38, Chapter V, Corollary 4.12]. Thus $\text{Ext}_\Lambda^1(H/M, \Lambda) = 0$, so $i^*\hat{\lambda}$ is surjective and thus $M^\perp \hookrightarrow H$ is a split injection, as claimed. □

The fact that M/M^\perp is torsion follows from the claim because $M^\perp \subseteq M$ and $\text{rk}_\Lambda(M^\perp) = \text{rk}_\Lambda(M)$. We now show that M/M^\perp is torsion-free. Since submodules of torsion-free modules are torsion-free, it suffices to show that $H/M^\perp \supseteq M/M^\perp$ is torsion-free. But now by the claim, we know that H/M^\perp is (isomorphic to) a submodule of the free module $H \cong M^\perp \oplus H/M^\perp$ and so it is indeed torsion-free. This concludes the proof of the lemma. □

Given a Seifert matrix A for a genus g Seifert surface S , we write $(H, A) := (\mathbb{Z}^{2g}, A)$. Use $F \subseteq D^4$ to denote the result of pushing S into D^4 . After some choice of bases, the equivariant intersection form $(H_2(D_F^4; \Lambda), \lambda_{D_F^4})$ is isometric to

$$(H \otimes_{\mathbb{Z}} \Lambda, (1-t)A^T + (1-t^{-1})A).$$

See [35, Section 3], also [8, proof of Lemma 5.4].

Remark 7.8 $(H \otimes_{\mathbb{Z}} \Lambda, A - tA^T, A)$ defines a $(-t)$ -quadratic form. Furthermore, if A is a Seifert matrix for an Alexander polynomial one knot K , then this quadratic form is nonsingular since $\det(A - tA^T) \doteq \Delta_K(t) \doteq 1$.

Recall that a *metaboliser* for a Seifert matrix (\mathbb{Z}^{2g}, A) is a half-rank direct summand $L \subseteq \mathbb{Z}^{2g}$ such that $x^T Ay = 0$ for all $x, y \in L$. Since Alexander polynomial one knots are (in particular) algebraically slice, their Seifert matrices admit metabolisers.

Proposition 7.9 *Let $(H, A) := (\mathbb{Z}^{2g}, A)$ be a Seifert matrix for an Alexander polynomial one knot K . If $L \subseteq H$ is a metaboliser for (H, A) , then $L \otimes_{\mathbb{Z}} \Lambda$ is a Lagrangian for $(H \otimes_{\mathbb{Z}} \Lambda, A - tA^T, A)$, and so this latter $(-t)$ -quadratic form is hyperbolic.*

Proof We verify the three points in the definition of a Lagrangian. Since A vanishes on L , the extended form (also denoted by A) vanishes on $M := L \otimes_{\mathbb{Z}} \Lambda$. It follows that the $(-t)$ -Hermitian form $A - tA^T$ vanishes on $M \times M$. Since $L \subseteq H$ is a summand, so is $M \subseteq H \otimes_{\mathbb{Z}} \Lambda$. Thus $M \subseteq (H \otimes_{\mathbb{Z}} \Lambda, A - tA^T)$ is a sublagrangian of a $(-t)$ -Hermitian form, with $H \otimes_{\mathbb{Z}} \Lambda$ a free module. We showed in Lemma 7.7 that since M is half-rank, this forces $M^\perp = M$.

The last sentence of the proposition follows because it is known that if a nonsingular ε -quadratic form (H, λ, μ) admits a Lagrangian, then it is isometric to the standard hyperbolic form [46, Proposition 2.2]. □

Proposition 7.10 *If A is a Seifert matrix for an Alexander polynomial one knot K , then*

$$(H \otimes_{\mathbb{Z}} \Lambda, (1-t)A^T + (1-t^{-1})A) \cong \mathcal{H}_2^{\oplus g}.$$

Proof Using Proposition 7.9, we deduce the following sequence of isometries:

$$\begin{aligned} (H \otimes_{\mathbb{Z}} \Lambda, A - tA^T, A) &\cong H_{-t}(\Lambda)^{\oplus g} = \left(\Lambda^{2g}, \begin{pmatrix} 0 & 1 \\ -t & 0 \end{pmatrix}^{\oplus g}, \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}^{\oplus g} \right) \\ &\cong \left(\Lambda^{2g}, \begin{pmatrix} 0 & -t \\ 1 & 0 \end{pmatrix}^{\oplus g}, \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}^{\oplus g} \right). \end{aligned}$$

Multiplying both sides by $(1-t^{-1})$ then gives the assertion. □

We are now ready to prove Theorem 1.9 from the introduction.

Theorem 7.11 *If $F_0, F_1 \subseteq D^4$ are genus g pushed-in Seifert surfaces for an Alexander polynomial one knot K , then they are topologically ambiently isotopic rel boundary.*

Proof Let A_0 and A_1 be Seifert matrices for F_0 and F_1 . As mentioned above, the equivariant intersection form $(H_2(D_{F_i}^4; \Lambda), \lambda_{D_{F_i}^4})$ is isometric to

$$(H \otimes_{\mathbb{Z}} \Lambda, (1-t)A_i^T + (1-t^{-1})A_i).$$

By Proposition 7.10, both forms are isometric to $\mathcal{H}_2^{\oplus g}$. Since K is an Alexander polynomial one knot, the result now follows from Theorem 7.5. □

8 Rim surgery on surfaces with knot group \mathbb{Z}

We prove Theorems 1.7 and 1.8, which concern rim surgery on surfaces in 4-manifolds whose knot group is \mathbb{Z} . In Section 8.1, we review the definition of rim surgery; in Sections 8.2 and 8.3 we prove the main results.

8.1 Knot surgery and twist rim surgery

We review some facts about knot surgery and rim surgery. For references see for instance [14; 30; 32; 2].

Let Z be a compact, oriented 4-manifold containing a locally flat embedded torus T with trivial normal bundle, and let $J \subseteq S^3$ be a knot. Use μ_T to denote the meridian of $T \subseteq Z$, and μ_J and λ_J respectively for the meridian and 0-framed longitude of J . Let $\varphi: \partial v(T) \rightarrow S^1 \times \partial E_J$ be any diffeomorphism such that $\varphi_*(\mu_T) = \lambda_J$. The 4-manifold obtained by *knot surgery* along J and φ is defined as

$$Z_J(\varphi) = (Z \setminus v(T)) \cup_{\varphi} (S^1 \times E_J).$$

Given a compact 4-manifold W and a locally flat embedded orientable surface $\Sigma \subseteq W$, we assume either that W and Σ are closed or that $\Sigma \subseteq W$ is a properly embedded. As we now describe, rim surgery arises from a particular type of knot surgery on the surface exterior $W_{\Sigma} := W \setminus v(\Sigma)$. Choose a simple closed curve $\alpha \subseteq \Sigma$ and a trivialisation of the normal bundles over α and Σ so that $(v(\Sigma), v(\alpha)) = (\Sigma \times D^2, \alpha \times I \times D^2)$; in other words, the normal bundle of α inside Σ is $\alpha \times I$. This way, it is understood that $\mu_{\Sigma} = \{\text{pt}\} \times \partial D^2$, and the *rim torus* T is $\partial(v(\Sigma)|_{\alpha}) = \alpha \times \mu_{\Sigma}$. The rim torus is framed: if we write $v(\Sigma)|_{\alpha \times I} = \alpha \times I \times D^2$, then a framing of vT is given by the I -direction and the radial direction in the polar coordinates of D^2 .

In order to perform a knot surgery on W_Σ along J using the rim torus T , for integers m and n we consider the homeomorphism $\varphi_n^m: \partial\bar{v}(T) \rightarrow S^1 \times \partial E_J$ determined by

$$\varphi_n^m(\alpha) = m\mu_J + n\lambda_J + S^1, \quad \varphi_n^m(\mu_\Sigma) = \mu_J, \quad \varphi_n^m(\mu_T) = \lambda_J.$$

Gluing $\Sigma \times D^2$ back into the result of this knot surgery produces a manifold $W_J(\varphi_n^m)$ that is homeomorphic to W , and in fact if W is smooth then $W_J(\varphi_n^m)$ is diffeomorphic to W ; see [30, Lemma 2.4], in which a specific diffeomorphism is constructed. Thus, we obtain a new embedding $\Sigma_n^m(\alpha, J) \subseteq W$.

Definition 8.1 Let $\alpha \subseteq \Sigma$ be a simple closed curve, let J be a knot, and let $m, n \in \mathbb{Z}$ be integers. The n -roll m -twist rim surgery of a locally flat, properly embedded, orientable surface $\Sigma \subseteq W$ is the image $\Sigma_n^m(\alpha, J)$ of Σ under the homeomorphism $W \cong W_J(\varphi_n^m)$ mentioned above. If W is smooth and Σ is smoothly embedded, then since $W \cong W_J(\varphi_n^m)$ is a diffeomorphism, $\Sigma_n^m(\alpha, J)$ is smoothly embedded.

The exterior $W_{\Sigma_n^m(\alpha, J)}$ of the rim surgery $\Sigma_n^m(\alpha, J)$ is the knot surgery on W_Σ along the homeomorphism φ_n^m . The proof of the following lemma can be found in [30, Proposition 3.3].

Lemma 8.2 For a compact 4-manifold W and a locally flat, embedded, orientable surface $\Sigma \subseteq W$, we assume either that W and Σ are closed or that $\Sigma \subseteq W$ is properly embedded. Let $\alpha \subseteq \Sigma$ be a simple closed curve, let J be a knot, and let $m, n \in \mathbb{Z}$. If $\pi_1(W_\Sigma) = \mathbb{Z}_d$ with $d = \pm 1 \pmod m$, then

$$\pi_1(W_\Sigma) = \pi_1(W_{\Sigma_n^m(\alpha, J)}) = \mathbb{Z}_d.$$

In our case, the knot group is infinite cyclic, so we take $d = 0$ and restrict to n -roll 1-twist surgeries. We write $\Sigma_n(\alpha, J) := \Sigma_n^1(\alpha, J)$ and $\varphi_n := \varphi_n^1$.

Remark 8.3 If $J = U$ is the unknot then $\Sigma_n(\alpha, U) = \Sigma$, by [32, Lemma 2.2].

8.2 The Λ -intersection form

Now we focus on the case where the ambient manifold N has a boundary, and show that the Λ -intersection form of $N_{\Sigma_n(\alpha, J)}$ agrees with that of N_Σ .

We start with an observation concerning the rim torus.

Lemma 8.4 *Let $\Sigma \subseteq N = X \setminus \overset{\circ}{D}^4$ be a \mathbb{Z} -surface. The exterior of a rim torus T satisfies*

$$H_1(N_\Sigma \setminus \nu(T)) = \mathbb{Z}\langle \mu_\Sigma \rangle \oplus \mathbb{Z}\langle \mu_T \rangle.$$

Proof Consider the Mayer–Vietoris sequence of $N_\Sigma = (N_\Sigma \setminus \nu(T)) \cup \bar{\nu}(T)$:

$$(29) \quad \cdots \rightarrow H_2(N_\Sigma) \xrightarrow{\partial} H_1(T \times S^1) \xrightarrow{\theta} H_1(\bar{\nu}(T)) \oplus H_1(N_\Sigma \setminus \nu(T)) \rightarrow H_1(N_\Sigma) \rightarrow 0.$$

We show that $\partial = 0$. Note that

$$H_1(T \times S^1) \cong H_1(T) \oplus H_1(S^1) \xrightarrow{\theta} H_1(\bar{\nu}(T)) \cong H_1(T)$$

is projection onto the first factor, so for $x \neq 0$ in $H_1(T)$, $\theta(x, n) \neq 0$. Therefore in the image of ∂ , the first coordinate vanishes. Now for a closed surface $S \subseteq N$, the definition of the connecting homomorphism implies that

$$\partial([S]) = (0, Q_N([T], [S])) \in H_1(T \times S^1) \cong H_1(T) \oplus H_1(S^1).$$

Since T is isotopic to a torus in $\Sigma \times S^1 \subseteq \partial N_\Sigma$, we deduce that $Q_N([T], [S]) = 0$. It follows that $\partial = 0$ as asserted. The Mayer–Vietoris sequence in (29) then reduces to

$$0 \rightarrow \mathbb{Z}\langle \mu_T \rangle \rightarrow H_1(N_\Sigma \setminus \nu(T)) \rightarrow H_1(N_\Sigma) \rightarrow 0.$$

Since $H_1(N_\Sigma) = \mathbb{Z}\langle \mu_\Sigma \rangle$ is free, this sequence splits, establishing the lemma. □

Next, we extend the coefficient system on $\pi_1(N_\Sigma)$ over $\pi_1(N_{\Sigma_n(\alpha, J)})$.

Remark 8.5 The map $\pi_1(N_\Sigma) \rightarrow \mathbb{Z}$ restricts to a map $\pi_1(N_\Sigma \setminus \nu(T)) \rightarrow \mathbb{Z}$ that sends μ_Σ to 1 and μ_T to 0; recall from Lemma 8.4 that $H_1(N_\Sigma \setminus \nu(T)) \cong \mathbb{Z}\langle \mu_\Sigma \rangle \oplus \mathbb{Z}\langle \mu_T \rangle$. Consider the map $\pi_1(S^1 \times E_J) \rightarrow \mathbb{Z}$ that sends the class of the S^1 -factor to -1 and that is the abelianisation on $\pi_1(E_J)$. We verify that these maps extend to a map $\pi_1(N_{\Sigma_n(\alpha, J)}) \rightarrow \mathbb{Z}$. We are identifying the 3-torus $\partial \bar{\nu}(T) \subseteq \partial N_\Sigma \setminus \nu(T)$ with $S^1 \times \partial E_J$ via the homeomorphism φ_n . We have the identifications $[\alpha] \sim [S^1][\mu_J][\lambda_J]^n$, $[\mu_\Sigma] \sim [\mu_J]$ and $[\mu_T] \sim [\lambda_J]$. The loop $\alpha \subseteq \Sigma$ is mapped to zero under $\pi_1(N_\Sigma) \rightarrow \mathbb{Z}$; the same is true for $[S^1][\mu_J][\lambda_J]^n$ under the map $\pi_1(S^1 \times E_J) \rightarrow \mathbb{Z}$. Similarly, both μ_Σ and μ_J (resp μ_T and λ_J) are sent to 1 (resp. 0) under the respective maps to \mathbb{Z} .

Arguing as in [32, Proof of Lemma 2.3], we construct a degree one map

$$\Psi: N_{\Sigma_n(\alpha, J)} \rightarrow N_\Sigma.$$

Using obstruction theory, one can construct a degree one map $E_J \rightarrow E_U$, where $U \subseteq S^3$ is the unknot, that takes meridian to meridian and longitude to longitude. Cross it with the identity to obtain a degree one map $S^1 \times E_J \rightarrow S^1 \times E_U$. This can be glued to the identity map $\text{Id}_{N_\Sigma \setminus \nu(T)}$ to obtain a degree one map $N_{\Sigma_n(\alpha, J)} \rightarrow N_{\Sigma_n(\alpha, U)}$. By [32, Lemma 2.2], there is a homeomorphism $N_{\Sigma_n(\alpha, U)} \cong N_\Sigma$ and we have therefore obtained a degree one map

$$\Psi: N_{\Sigma_n(\alpha, J)} \rightarrow N_{\Sigma_n(\alpha, U)} \cong N_\Sigma.$$

By construction, Ψ is the identity on ∂N_Σ . Furthermore, since the fundamental groups of both $N_{\Sigma_n(\alpha, J)}$ and N_Σ are generated by the meridians of the surface, Ψ induces an isomorphism on fundamental groups: indeed by construction Ψ takes meridian to meridian. In particular Ψ induces homomorphisms on the Λ -homology groups.

In order to prove that Ψ in fact induces an isometry of the equivariant intersection forms, we compute the Λ -homology of $S^1 \times E_J$.

Lemma 8.6 *Let $J \subseteq S^3$ be a knot. With respect to the coefficient system on $S^1 \times E_J$ introduced in Remark 8.5, we have $H_2(S^1 \times E_J; \Lambda) = 0$ and $H_1(S^1 \times E_J; \Lambda) = \mathbb{Z}$. Additionally, the degree one map Ψ induces an isometry*

$$\Psi_*: (H_2(N_{\Sigma_n(\alpha, J)}; \Lambda), \lambda_{N_{\Sigma_n(\alpha, J)}}) \xrightarrow{\cong} (H_2(N_\Sigma; \Lambda), \lambda_{N_\Sigma}).$$

Proof We use the Künneth spectral sequence with

$$E_{p,q}^2 = \bigoplus_{q_1+q_2=q} \text{Tor}_p(H_{q_1}(S^1; \Lambda), H_{q_2}(E_J; \Lambda)),$$

which converges to $H_*(S^1 \times E_J; \Lambda \otimes_\Lambda \Lambda) \cong H_*(S^1 \times E_J; \Lambda)$, where $\Lambda \otimes_\Lambda \Lambda$ is a right $\mathbb{Z}[\pi_1(E_K) \times \pi_1(S^1)]$ -module via the diagonal action induced by $p \otimes q \cdot (e, s) = pe \otimes qs$ for $p, q \in \Lambda$ and $(e, s) \in \pi_1(E_K) \times \pi_1(S^1)$. Write $\varepsilon: \Lambda \rightarrow \mathbb{Z}$, $p(t) \mapsto p(1)$ for the augmentation map and \mathbb{Z}_ε for the resulting Λ -module structure on \mathbb{Z} . By definition of the coefficient system in Remark 8.5, we have $H_1(S^1; \Lambda) = 0$ and $H_0(S^1; \Lambda) = \Lambda/(t-1) = \mathbb{Z}_\varepsilon$. We also have $H_0(E_J; \Lambda) = \mathbb{Z}$ and $H_i(E_J; \Lambda) = 0$ for $i \geq 2$. This implies that $E_{0,2}^2 = 0$. In fact, we also deduce that $E_{1,0}^2 = \mathbb{Z}$ and $E_{2,0}^2 = 0$ because

$$E_{i,0}^2 = \text{Tor}_i^{\mathbb{Z}[\mathbb{Z}]}(\mathbb{Z}, \mathbb{Z}) = H_i(\mathbb{Z}; \mathbb{Z}) = H_i(S^1).$$

We claim that $\text{Tor}_i^\Lambda(\mathbb{Z}_\varepsilon, H_1(E_J; \Lambda)) = 0$ for $i = 0, 1$. These Tor groups are computed as the homology of the complex obtained tensoring the resolution $0 \rightarrow \Lambda \xrightarrow{t-1} \Lambda \rightarrow \mathbb{Z}_\varepsilon \rightarrow 0$ with the Alexander module $H_1(E_J; \Lambda)$. Since multiplication by $t-1$ induces an

isomorphism on the Alexander module [39, Proposition 1.2], these Tor groups vanish and the claim is proved.

Using the claim, we deduce that $E_{0,1}^2 = 0$ and $E_{1,1}^2 = 0$. It follows that $d_{2,0} = 0$ and now the first assertion is a consequence of a standard spectral sequence computation.

We now assert that Ψ induces an isomorphism on $H_2(-; \Lambda)$. Recall the decompositions

$$N_{\Sigma_n(\alpha, J)} = (N_\Sigma \setminus \nu(T)) \cup_{\varphi_n} (S^1 \times E_J),$$

$$N_\Sigma \cong N_{\Sigma_n(\alpha, U)} = (N_\Sigma \setminus \nu(T)) \cup_{\varphi_n} (S^1 \times E_U).$$

By construction, Ψ restricts to the identity on $N_\Sigma \setminus \nu(T)$ and $\partial\bar{\nu}(T)$ and therefore induces the identity on the Λ -homology of these spaces. Next, $\Psi_*: H_1(S^1 \times E_J; \Lambda) \rightarrow H_1(S^1 \times E_U; \Lambda)$ is also an isomorphism: indeed, the first assertion shows that both modules are isomorphic to $\mathbb{Z} = \Lambda/(t-1)$, and this term comes from $H_0(E_J; \Lambda) = \mathbb{Z}_\varepsilon$, on which Ψ does indeed induce an isomorphism. We showed that $H_2(S^1 \times E_J; \Lambda) = 0$, so the assertion now follows from the five lemma applied to the following commutative diagram, where Λ -coefficients are understood:

$$\begin{array}{ccccccccc}
 H_2(\partial\bar{\nu}(T)) & \longrightarrow & H_2(N_\Sigma \setminus \nu(T)) & \longrightarrow & H_2(N_{\Sigma_n(\alpha, J)}) & \longrightarrow & H_1(\partial\bar{\nu}(T)) & \longrightarrow & H_1(N_\Sigma \setminus \nu(T)) & \longrightarrow & 0 \\
 \Psi_* \parallel & & \Psi_* \parallel & & \Psi_* \downarrow & & \Psi_* \parallel & & \Psi_* \downarrow \cong & & \\
 H_2(\partial\bar{\nu}(T)) & \longrightarrow & H_2(N_\Sigma \setminus \nu(T)) & \longrightarrow & H_2(N_\Sigma) & \longrightarrow & H_1(\partial\bar{\nu}(T)) & \longrightarrow & H_1(N_\Sigma \setminus \nu(T)) & \longrightarrow & 0 \\
 & & & & & & & & \oplus & & \\
 & & & & & & & & \mathbb{Z} & & \\
 & & & & & & & & \downarrow & & \\
 & & & & & & & & \mathbb{Z} & &
 \end{array}$$

Since Ψ induces a Λ -isomorphism on $H_2(-; \Lambda)$, it induces one on $H_2(-; \Lambda)^*$, and therefore on $H^2(-; \Lambda)$ (recall the UCSS argument from Lemma 3.2) and thus on the second relative Λ -homology groups (because $\deg(\Psi) = 1$). We conclude that Ψ_* in fact induces an isometry of the Λ -intersection forms. □

8.3 Topological triviality of rim surgery on surfaces with knot group \mathbb{Z}

Next we prove Theorem 1.8 from the introduction.

Theorem 8.7 *Let $\Sigma \subseteq N = X \setminus \overset{\circ}{D}^4$ be a \mathbb{Z} -surface, let $\alpha \subseteq \Sigma$ be a simple closed curve, let J be a knot, and let $n \in \mathbb{Z}$ be an integer. There is a rel boundary orientation-preserving homeomorphism of pairs*

$$\Phi: (N, \Sigma_n(\alpha, J)) \xrightarrow{\cong} (N, \Sigma)$$

that induces the same isometry as the degree one map Ψ on the equivariant intersection forms of the surface exteriors. If $N = D^4$, then the surfaces are topologically isotopic rel boundary.

Proof By Lemma 8.2 both surfaces have knot group \mathbb{Z} . Lemma 8.6 states that the degree one map $\Psi: N_{\Sigma_n(\alpha, J)} \rightarrow N_\Sigma$ induces an isometry $\Psi_*: \lambda_{N_{\Sigma_n(\alpha, J)}} \cong \lambda_{N_\Sigma}$. As Ψ is the identity on ∂N_Σ , we see that $\partial\Psi$ induces the identity isometry between the boundary Blanchfield forms. The result now follows from Theorem 5.8. \square

The remaining result we owe a proof of is Theorem 1.7 from the introduction. To prove this result, which involves obtaining an isotopy between closed surfaces, we need to control the map $F_{\mathbb{Z}}$ discussed in Lemma 5.10. This will rely on the following result.

Lemma 8.8 *Let $\Sigma \subseteq N$ be a \mathbb{Z} -surface, let $\alpha \subseteq \Sigma$ be a simple closed curve with associated rim torus T , and let J be a knot. The following sequences of inclusion induced maps are exact:*

$$\begin{aligned} 0 \rightarrow \mathbb{Z}\langle \alpha \times \mu_T \rangle &\rightarrow H_2(N_\Sigma \setminus \nu(T); \Lambda) \rightarrow H_2(N_{\Sigma_n(\alpha, J)}; \Lambda) \rightarrow 0, \\ 0 \rightarrow \mathbb{Z}\langle \alpha \times \mu_T \rangle &\rightarrow H_2(N_\Sigma \setminus \nu(T); \Lambda) \rightarrow H_2(N_\Sigma; \Lambda) \rightarrow 0. \end{aligned}$$

Proof As we noted in Lemma 8.2 that $\pi_1(N_{\Sigma_n(\alpha, J)}) = \mathbb{Z}$ and since $N_{\Sigma_n(\alpha, J)}$ has ribbon boundary, we know that $H_1(N_{\Sigma_n(\alpha, J)}; \Lambda) = 0$ as well as $H_3(N_{\Sigma_n(\alpha, J)}; \Lambda) = 0$, again by Lemma 3.2. Consequently, the Mayer–Vietoris sequence for $N_{\Sigma_n(\alpha, J)} = N_\Sigma \setminus \nu(T) \cup_{\varphi_n} (S^1 \times E_J)$ with Λ coefficients gives

$$\begin{aligned} 0 \rightarrow H_2(\partial\bar{\nu}(T); \Lambda) &\rightarrow H_2(N_\Sigma \setminus \nu(T); \Lambda) \oplus H_2(S^1 \times E_J; \Lambda) \rightarrow H_2(N_{\Sigma_n(\alpha, J)}; \Lambda) \\ &\rightarrow H_1(\partial\bar{\nu}(T); \Lambda) \rightarrow H_1(N_\Sigma \setminus \nu(T); \Lambda) \oplus H_1(S^1 \times E_J; \Lambda) \rightarrow 0. \end{aligned}$$

Since there is a homotopy equivalence $\bar{\nu}(T) \simeq \alpha \times \mu_\Sigma$, the discussion of coefficient systems from Remark 8.5 implies that

$$H_i(\partial\bar{\nu}(T); \Lambda) = H_i(\alpha \times \mu_\Sigma \times \mu_T; \Lambda) = H_i(\alpha \times \mu_T) \quad \text{for } i = 1, 2.$$

We saw in Lemma 8.6 that $H_2(E_J \times S^1; \Lambda) = 0$ and $H_1(E_J \times S^1; \Lambda) = \mathbb{Z}$. The previous sequence therefore reduces to

$$(30) \quad 0 \rightarrow \mathbb{Z}\langle \alpha \times \mu_T \rangle \rightarrow H_2(N_\Sigma \setminus \nu(T); \Lambda) \rightarrow H_2(N_{\Sigma_n(\alpha, J)}; \Lambda) \rightarrow \mathbb{Z}\langle \alpha \rangle \oplus \mathbb{Z}\langle \mu_T \rangle \rightarrow H_1(N_\Sigma \setminus \nu(T); \Lambda) \oplus \mathbb{Z} \rightarrow 0.$$

When $J = U$ is the unknot, $H_1(S^1 \times E_J; \Lambda) = H_1(\alpha) = \mathbb{Z}\langle \alpha \rangle$, and therefore $\mathbb{Z}\langle \mu_T \rangle$ surjects onto $H_1(N_\Sigma \setminus \nu(T); \Lambda)$. In particular, as an abelian group, $H_1(N_\Sigma \setminus \nu(T); \Lambda)$ is cyclic.

Claim 7 The map $\mathbb{Z}\langle\mu_T\rangle \xrightarrow{\iota} H_1(N_\Sigma \setminus \nu(T); \Lambda)$ is an isomorphism.

Proof We already noted that, as an abelian group, $H_1(N_\Sigma \setminus \nu(T); \Lambda)$ is cyclic, so it suffices to show that $H_1(N_\Sigma \setminus \nu(T); \Lambda) \otimes_{\mathbb{Z}} \mathbb{Q} \neq 0$. Equivalently, we must establish that

$$H := H_1(N_\Sigma \setminus \nu(T); \mathbb{Q}[t^{\pm 1}]) \neq 0.$$

By way of contradiction, assume that $H = 0$. Let $\varepsilon: \Lambda \rightarrow \mathbb{Z}$ again denote the augmentation map and write \mathbb{Z}_ε and \mathbb{Q}_ε for the resulting Λ -module structures. Since $\text{Tor}_2^\Lambda(\mathbb{Z}_\varepsilon, \mathbb{Q}_\varepsilon) = 0$, the universal coefficient spectral sequence applied to

$$H_1(N_\Sigma \setminus \nu(T); \mathbb{Q}) = H_1(N_\Sigma \setminus \nu(T); \mathbb{Q}[t^{\pm 1}] \otimes_\Lambda \mathbb{Q}_\varepsilon)$$

produces the short exact sequence

$$0 \rightarrow H \otimes_\Lambda \mathbb{Q}_\varepsilon \rightarrow H_1(N_\Sigma \setminus \nu(T); \mathbb{Q}) \rightarrow \text{Tor}_1^\Lambda(\mathbb{Z}_\varepsilon, \mathbb{Q}_\varepsilon) \rightarrow 0.$$

We assumed that $H = 0$, so $H \otimes_\Lambda \mathbb{Q}_\varepsilon = 0$, and thus $H_1(N_\Sigma \setminus \nu(T); \mathbb{Q}) \cong \text{Tor}_1^\Lambda(\mathbb{Z}, \mathbb{Q})$. Using Lemma 8.4, $H_1(N_\Sigma \setminus \nu(T); \mathbb{Q}) = \mathbb{Q}^2$. On the other hand, $\text{Tor}_1^\Lambda(\mathbb{Z}_\varepsilon, \mathbb{Q}_\varepsilon) = H_1(S^1; \mathbb{Q}) = \mathbb{Q}$. This is a contradiction and the claim is established. \square

Using Claim 7, the penultimate map in (30) is a surjection $\mathbb{Z}^2 \rightarrow \mathbb{Z}^2$ and therefore an isomorphism. The first short exact sequence in the statement of the lemma now follows from the one displayed in (30). The exactness of the second sequence follows from exactness of the first by taking $J = U$ and recalling that $\Sigma_n(\alpha, U) = \Sigma$ by Remark 8.3. \square

Theorem 8.9 Let $\Sigma \subseteq X$ be a closed \mathbb{Z} -surface, let $\alpha \subseteq \Sigma$ be a simple closed curve, let J be a knot, and let $n \in \mathbb{Z}$. The surfaces Σ and $\Sigma_n(\alpha, J)$ are topologically ambiently isotopic.

Proof After an ambient isotopy, we may assume that the surfaces $\Sigma_0 := \Sigma_n(\alpha, J)$ and $\Sigma_1 := \Sigma$ coincide on a disc $D^2 \subseteq \Sigma \cap \Sigma_n(\alpha, J)$ that does not intersect $\alpha \subseteq \Sigma$. Assume that the normal bundles also coincide over this D^2 . Consider the preimage $\mathring{D}^2 \times \mathbb{R}^2 \subseteq \nu\Sigma_j$. This is homeomorphic to an open 4-ball \mathring{D}^4 . Remove this $(\mathring{D}^4, \mathring{D}^2)$ from (X, Σ) and $(X, \Sigma_n(\alpha, J))$ to obtain properly embedded surfaces $\tilde{\Sigma} \subseteq N$ and $\tilde{\Sigma}_n(\alpha, J) \subseteq N$ with the unknot as a common boundary. By construction, $\tilde{\Sigma}_n(\alpha, J)$ is obtained by rim surgery on $\tilde{\Sigma}$. Apply Theorem 8.7 to obtain a rel boundary homeomorphism of pairs $\Phi: (N, \tilde{\Sigma}_n(\alpha, J)) \rightarrow (N, \tilde{\Sigma})$. Recall that on $H_2(-; \Lambda)$, we have $\Phi_* = \Psi_*$, where Ψ is the degree one map described above Lemma 8.6. Construct

a homeomorphism of pairs $(X, \Sigma_n(\alpha, J)) \rightarrow (X, \Sigma)$ by gluing Φ with the identity homeomorphism $(D^4, D^2) \rightarrow (D^4, D^2)$.

We refine this argument to obtain the required ambient isotopy. Use F to denote the isometry induced by these homeomorphisms on $H_2(-; \Lambda)$ of the surface exteriors. In Lemma 8.6, we argued that this isometry fits into the following commutative diagram with exact rows (here we also used Lemma 8.8 to simplify the Mayer–Vietoris sequences):

$$(31) \quad \begin{array}{ccccccc} 0 & \longrightarrow & \mathbb{Z}\langle \alpha \times \mu_T \rangle & \longrightarrow & H_2(N_{\tilde{\Sigma}} \setminus \nu(T); \Lambda) & \longrightarrow & H_2(N_{\tilde{\Sigma}_n(\alpha, J)}; \Lambda) \longrightarrow 0 \\ & & \parallel \text{Id} & & \parallel \text{Id} & & \downarrow F = \Psi_* \\ 0 & \longrightarrow & \mathbb{Z}\langle \alpha \times \mu_T \rangle & \longrightarrow & H_2(N_{\tilde{\Sigma}} \setminus \nu(T); \Lambda) & \longrightarrow & H_2(N_{\tilde{\Sigma}}; \Lambda) \longrightarrow 0 \end{array}$$

Recall that $\Sigma_0 := \Sigma_n(\alpha, J)$ and $\Sigma_1 := \Sigma$, so $X_{\Sigma_0} \cong N_{\tilde{\Sigma}_n(\alpha, J)}$ and $X_{\Sigma_1} \cong N_{\tilde{\Sigma}}$. In Lemma 5.10, $F_{\mathbb{Z}}$ was defined by the following commutative diagram with exact rows:

$$(32) \quad \begin{array}{ccccccc} 0 & \longrightarrow & \mathbb{Z}^{2g} & \longrightarrow & H_2(X_{\Sigma_0}) & \xrightarrow{p_0} & H_2(X) \longrightarrow 0 \\ & & \downarrow (F \otimes_{\Lambda} \text{Id}_{\mathbb{Z}}) | & & \downarrow F \otimes_{\Lambda} \text{Id}_{\mathbb{Z}} & & \downarrow F_{\mathbb{Z}} \\ 0 & \longrightarrow & \mathbb{Z}^{2g} & \longrightarrow & H_2(X_{\Sigma_1}) & \xrightarrow{p_1} & H_2(X) \longrightarrow 0 \end{array}$$

By (32), every element x of $H_2(X)$ can be represented by a class in $H_2(X_{\Sigma_0})$, and by (31) every class here can be represented by a surface S in $N_{\tilde{\Sigma}_n(\alpha, J)} \setminus \nu(T) \subseteq N_{\tilde{\Sigma}_n(\alpha, J)} \cong X_{\Sigma_0}$. Since F is induced by the degree one map Ψ , mapping our surface to the same surface S in $N_{\tilde{\Sigma}} \setminus \nu(T) \subseteq N_{\tilde{\Sigma}} \cong X_{\Sigma_1}$ yields $(F \otimes_{\Lambda} \text{Id}_{\mathbb{Z}})([S]) \in H_2(X_{\Sigma_1})$. The inclusion induced map $p_1: H_2(X_{\Sigma_1}) \rightarrow H_2(X)$ then sends $(F \otimes_{\Lambda} \text{Id}_{\mathbb{Z}})([S])$ to x . Therefore $F_{\mathbb{Z}}: H_2(X) \rightarrow H_2(X)$ is indeed the identity, so the second item of Theorem 5.11 implies that Σ_0 and Σ_1 are topologically ambiently isotopic. \square

Appendix Stable isotopy for surfaces in a topological 4–manifold with boundary

The next result is an extension of [2, Theorem 5] to the topological case, and allowing nonempty boundary, but restricting the ambient 4–manifolds somewhat. Since we need the given extension, we provide details of the proof. The main ideas are due to [2]. We fill in some details in their argument for constructing a map to S^1 in the course of the proof. The case with nonempty boundary was also stated as [25, Proposition 2.13] in the smooth category.

Theorem A.1 *Let N be an oriented, connected, simply connected, compact topological 4–manifold with boundary S^3 . Let Σ be a compact, connected, orientable surface with one boundary component. Let $\Sigma_0, \Sigma_1 \subseteq N$ be two properly embedded, locally flat, oriented surfaces homeomorphic to Σ , and suppose that $\partial\Sigma_0 = \partial\Sigma_1$. Assume that for each i , $\pi_1(N \setminus \Sigma_i) \cong \mathbb{Z}$ is infinite cyclic generated by an oriented meridian to Σ_i . Then some finite number of trivial one-handle stabilisations results in ambiently isotopic surfaces Σ'_0 and Σ'_1 , with $\partial\Sigma_i = \partial\Sigma'_i$ for $i = 0, 1$.*

We begin with a lemma on framings, some of which is a recollection from Section 5. Note that any locally flat embedded surface in a 4–manifold admits a normal bundle with linear structure group [15, Theorem 9.3].

Lemma A.2 *For $i = 0, 1$ the normal bundle of Σ_i is trivial and for any choice of framing there is a well-defined induced (homotopy class of) framing on the normal bundle of $\partial\Sigma_i$. The surface Σ_i is null-homologous in $H_2(N, \partial N)$, and the induced framing on $\partial\Sigma_i$ equals the Seifert framing.*

Proof Since the surface Σ_i has nonempty boundary and is oriented, its normal bundle is trivial. Two choices of framings on Σ_i differ by a map $\Sigma_i \rightarrow \text{SO}(2)$. The difference between the two framings on the boundary $\partial\Sigma_i$ is governed by the homotopy class of the composite

$$S^1 \xrightarrow{\cong} \partial\Sigma_i \rightarrow \Sigma_i \rightarrow \text{SO}(2) \cong S^1.$$

Since the boundary determines a commutator in $\pi_1(\Sigma_i)$, the displayed map is null-homotopic. It follows that the two framings induced on $\partial\Sigma_i$ agree up to homotopy. This proves that the induced framing is well-defined. The second sentence was already shown in Lemmas 5.1 and 5.2. □

Proof of Theorem A.1 Push the boundary of Σ_1 off the boundary of Σ_0 , using the Seifert framing, to arrange that $\partial\Sigma_0$ and $\partial\Sigma_1$ in $\partial N \cong S^3$ are parallel circles. We may assume that there is a collar $\partial N \times I$ with $\partial N \times \{0\} = \partial N$, and an embedding $g: S^1 \times I \times I \hookrightarrow \partial N \times I$ corresponding to the Seifert framing push-off, with

$$\begin{aligned} g(S^1 \times I \times \{i\}) &= \Sigma_i \cap (\partial N \times I) \quad \text{for } i = 0, 1, \\ g(S^1 \times \{t\} \times I) &\subseteq \partial N \times \{t\} \quad \text{for all } t \in I. \end{aligned}$$

Let $A \subseteq \partial N \cong S^3$ be the annulus

$$A := g(S^1 \times \{0\} \times I)$$




Figure 4: A schematic diagram indicating the relation of $g(S^1 \times I \times I) \subseteq \partial N \times I$ to $\partial \Sigma_0$, $\partial \Sigma_1$, and the annulus $A \subseteq S^3$ that joins them.

connecting $\partial \Sigma_0$ and $\partial \Sigma_1$, arising from the trace of the push. Note that A induces the Seifert framing on $\partial \Sigma_i$ for $i = 0, 1$, and that by Lemma A.2 this equals the framing induced by some choice of framing of the normal bundle of Σ_i .

By an isotopy, arrange further that Σ_0 and Σ_1 intersect transversely in their interiors [15, Theorem 9.5], and since Σ is compact we may also assume there are finitely many intersection points.

Cap off each $\partial \Sigma_i$ with a Seifert surface, and by a small isotopy of the capped-off Σ_1 , arrange that the capped-off surfaces are disjoint in $\partial N \times I$. By Lemma A.2 (and since $H_2(\partial N) = H_2(S^3) = 0$), the capped-off surfaces are null-homologous in $H_2(N)$, and so intersect algebraically zero times. By tubing Σ_0 to itself, that is by 1–handle stabilisations, arrange that Σ_0 and Σ_1 are disjoint. To achieve this, pair up points with opposite signs in $\Sigma_0 \pitchfork \Sigma_1$ and for each pair $\{p, q\}$ choose a path γ in Σ_1 connecting the two intersection points and away from the other intersection points. We can choose these paths to be mutually disjoint, but this is not obligatory. For each pair of points $\{p, q\}$, remove two open discs from Σ_0 , one for each pair of points, and add a tube, coming from the normal circle bundle of Σ_1 restricted to γ , as shown in Figure 5. In case different paths on Σ_1 intersect, vary the radii of the tubes to keep them disjoint. This stabilises Σ_0 to a surface disjoint from Σ_1 .

Perform the same number of trivial 1–handle stabilisations to Σ_1 , so that the surfaces are still abstractly homeomorphic. We will abuse notation and still refer to the resulting surfaces as Σ_0 and Σ_1 . Let

$$N_{\Sigma_0, \Sigma_1} := N \setminus (\nu \Sigma_0 \cup \nu \Sigma_1).$$

We will construct an oriented 3–dimensional relative Seifert manifold $Y \subseteq N_{\Sigma_0, \Sigma_1}$, that is a rel boundary cobordism between Σ_0 and Σ_1 , embedded in N and with $\partial Y = S$, where

$$S := \Sigma_0 \cup A \cup -\Sigma_1.$$




Figure 5: Stabilising Σ_0 to Σ'_0 by adding a 1–handle in a neighbourhood of an arc γ in Σ_1 .

Claim *There is a locally flat, embedded, compact, orientable 3–manifold Y with $\partial Y = S$ and corners at $\Sigma_i \cap A$, as described in the preceding paragraph.*

Proof The strategy to construct Y is as follows: define a suitable map $\partial N_{\Sigma_0, \Sigma_1} \rightarrow S^1$, extend it to a map $N_{\Sigma_0, \Sigma_1} \rightarrow S^1$ while controlling the restriction to $\partial N_{\Sigma_0, \Sigma_1}$, and then take Y to be the inverse image of a transverse regular point in S^1 .

The first step is to construct a map $\alpha: N_{\Sigma_0, \Sigma_1} \rightarrow S^1$. Recall that $N_{\Sigma_i} := N \setminus \nu \Sigma_i$. We can express $N = N_{\Sigma_0} \cup N_{\Sigma_1}$, with $N_{\Sigma_0, \Sigma_1} = N_{\Sigma_0} \cap N_{\Sigma_1}$. A portion of the resulting Mayer–Vietoris sequence (with \mathbb{Z} coefficients) is

$$H_2(N) \xrightarrow{\delta} H_1(N_{\Sigma_0, \Sigma_1}) \rightarrow H_1(N_{\Sigma_0}) \oplus H_1(N_{\Sigma_1}) \rightarrow H_1(N).$$

By hypothesis $H_1(N) = 0$ and $H_1(N_{\Sigma_i}) \cong \mathbb{Z}$, so that we obtain a short exact sequence

$$0 \rightarrow \text{im } \delta \rightarrow H_1(N_{\Sigma_0, \Sigma_1}) \rightarrow \mathbb{Z} \oplus \mathbb{Z} \rightarrow 0.$$

Since $\mathbb{Z} \oplus \mathbb{Z}$ is free abelian, this splits and we have that

$$H_1(N_{\Sigma_0, \Sigma_1}) \cong \mathbb{Z} \oplus \mathbb{Z} \oplus \text{im } \delta.$$

By the hypotheses of Theorem A.1, the first two summands are generated by meridians μ_{Σ_0} and μ_{Σ_1} to the surfaces Σ_0 and Σ_1 , respectively. Consider the dual element

$$\alpha := \mu_{\Sigma_0}^* - \mu_{\Sigma_1}^* \in \text{Hom}(H_1(N_{\Sigma_0, \Sigma_1}), \mathbb{Z}) \cong H^1(N_{\Sigma_0, \Sigma_1}) \cong [N_{\Sigma_0, \Sigma_1}, S^1],$$

sending the $\text{im } \delta$ summand identically to 0. We will also write $\alpha: N_{\Sigma_0, \Sigma_1} \rightarrow S^1$ for a corresponding representing map. We use that since N_{Σ_0, Σ_1} is a topological 4–manifold, it is homotopy equivalent to a CW–complex [17, Theorem 4.5], and therefore we may identify $H^1(N_{\Sigma_0, \Sigma_1}) \cong [N_{\Sigma_0, \Sigma_1}, S^1]$.

Now we consider the restriction to $\partial N_{\Sigma_0, \Sigma_1}$ under the inclusion-induced map

$$\iota^*: H^1(N_{\Sigma_0, \Sigma_1}) \rightarrow H^1(\partial N_{\Sigma_0, \Sigma_1}) \cong [\partial N_{\Sigma_0, \Sigma_1}, S^1],$$

where $\iota: \partial N_{\Sigma_0, \Sigma_1} \rightarrow N_{\Sigma_0, \Sigma_1}$ is the inclusion map. We have a decomposition

$$\partial N_{\Sigma_0, \Sigma_1} = \Sigma_0 \times S^1 \cup \partial_0 N_{\Sigma_0, \Sigma_1} \cup \Sigma_1 \times S^1.$$

An elementary Mayer–Vietoris argument yields that

$$H^1(\partial N_{\Sigma_0, \Sigma_1}) \cong \mathbb{Z} \oplus \mathbb{Z} \oplus \mathbb{Z}^{2g} \oplus \mathbb{Z}^{2g}.$$

The first two summands are generated by duals to the meridians, $\mu_{\Sigma_0}^*$ and $\mu_{\Sigma_1}^*$. By our work with framings above, for $i = 0, 1$ there is a framing of the normal bundle of Σ_i , determining an identification of the tubular neighbourhood $\bar{\nu}\Sigma_i$ with $\Sigma_i \times D^2$, that agrees on $\partial\Sigma_i$ with a corresponding identification induced from the Seifert framing on $\partial\Sigma_i$. This determines an identification

$$\partial\bar{\nu}\Sigma_i \setminus \nu\partial\Sigma_i \cong \Sigma_i \times S^1.$$

The two \mathbb{Z}^{2g} summands of $H^1(\partial N_{\Sigma_0, \Sigma_1})$ are generated by dual classes to curves of the form $\gamma_k \times \{-1\}$, where γ_k is a simple closed curve forming part of a symplectic basis for $H_1(\Sigma_i)$, for some i . Here we use the chosen framing of the normal bundle of Σ_i to fix representatives for the \mathbb{Z}^{2g} summands.

The restriction $\iota^*\alpha \in H^1(\partial N_{\Sigma_0, \Sigma_1})$ is $(1, -1, x, y)$, for some $x, y \in \mathbb{Z}^{2g}$. But by changing the choice of framing of $\nu\Sigma_i$ along the curves γ_k , we can arrange that $x = y = 0$. By changing the framing of the normal bundle of Σ_i by some number of full twists along some basis curve γ_k in $H_1(\Sigma_i)$, we change the meaning of $\gamma_k \times \{-1\}$ in the previous paragraph.

Let us explain the operation of “changing the framing” in more detail. Such a change is occasioned by the action of $[\bigcup_k \gamma_k, \text{SO}(2)]$ on the set of framings of the normal bundle, to alter the given framing on the union of the curves $\{\gamma_k\}$ by some number of full twists for each curve. Any such alteration automatically extends over the 2–skeleton since the attaching map of the 2–cell of Σ_i is a commutator in the γ_k times $\partial\Sigma_i$. Any two choices of extension over the 2–cells are homotopic, since $\pi_2(\text{SO}(2)) = 0$. Therefore we have a well-defined notion of altering the framing along the curves γ_k . This changes the entry of (x, y) corresponding to γ_k , since the map $\alpha: N_{\Sigma_0, \Sigma_1} \rightarrow S^1$ now sends $\gamma_k \times \{-1\}$ to a curve representing a different element of $H_1(S^1)$. By Lemma A.2, changing the choice of framing on a basis element for $H_1(\Sigma_i)$ does not change the induced framing on the boundary.

Now we define a map $f: \partial N_{\Sigma_0, \Sigma_1} \rightarrow S^1$. On $\Sigma_i \times S^1 \subseteq \partial N_{\Sigma_0, \Sigma_1}$, define the map to S^1 by the projection $f|_{\Sigma_i \times S^1}: \Sigma_i \times S^1 \rightarrow S^1$ onto the second factor. On the remainder

of $\partial N_{\Sigma_0, \Sigma_1}$, namely

$$\partial_0 N_{\Sigma_0, \Sigma_1} := \partial N \setminus (\nu \Sigma_0 \cup \nu \Sigma_1),$$

or in other words the link exterior $S^3 \setminus (\nu \partial \Sigma_0 \cup \nu \partial \Sigma_1)$, define a Pontryagin–Thom-style collapse map $f|_{\partial_0 N_{\Sigma_0, \Sigma_1}} : \partial_0 N_{\Sigma_0, \Sigma_1} \rightarrow S^1$ by choosing a tubular neighbourhood $A \times [-1, 1] \subseteq \partial_0 N_{\Sigma_0, \Sigma_1}$ and sending $(a, x) \mapsto e^{\pi i x} \in S^1$ for $a \in A$ and $x \in [-1, 1]$, then sending $\partial_0 N_{\Sigma_0, \Sigma_1} \setminus (A \times [-1, 1])$ to $-1 \in S^1$. Then $f|_{\partial_0 N_{\Sigma_0, \Sigma_1}}^{-1}(\{1\}) = A \cap \partial_0 N_{\Sigma_0, \Sigma_1}$. Note that we may assume that our maps to S^1 agree on the torus overlaps $\partial \Sigma_i \times S^1$, since the framings on $\partial \Sigma_i$ all agree up to homotopy. This completes the construction of a map $f : \partial N_{\Sigma_0, \Sigma_1} \rightarrow S^1$. Note that $f : \partial N_{\Sigma_0, \Sigma_1} \rightarrow S^1$ corresponds to the element

$$(1, -1, 0, 0) \in \mathbb{Z} \oplus \mathbb{Z} \oplus \mathbb{Z}^{2g} \oplus \mathbb{Z}^{2g}.$$

Therefore the cohomology classes $\iota^* \alpha$ and f agree in $H^1(\partial N_{\Sigma_0, \Sigma_1}) \cong [\partial N_{\Sigma_0, \Sigma_1}, S^1]$, and so by a homotopy of α supported in a collar of $\partial N_{\Sigma_0, \Sigma_1}$ we obtain a map $F : N_{\Sigma_0, \Sigma_1} \rightarrow S^1$ with $F|_{\partial N_{\Sigma_0, \Sigma_1}} = f : \partial N_{\Sigma_0, \Sigma_1} \rightarrow S^1$.

The inverse image under F of a transverse regular point in S^1 yields a 3–dimensional relative Seifert manifold, locally flatly embedded in $N_{\Sigma_0, \Sigma_1} = N \setminus (\nu \Sigma_0 \cup \nu \Sigma_1)$. See [17, Section 10.2] for information on map transversality in the topological category. Add collars $\Sigma_0 \times I$ and $\Sigma_1 \times I$ in $\nu \Sigma_0$ and $\nu \Sigma_1$, respectively, to obtain the 3–manifold Y that we seek. This completes the proof of the claim. □

Morse theory on Y gives rise to a Heegaard decomposition relative to the annulus A . The Heegaard surface can be obtained from both Σ_0 and Σ_1 by 1–handle stabilisations and ambient isotopy.

Since by our assumptions $\pi_1(N_{\Sigma_i})$ is generated by a meridian to Σ_i , Boyle’s [6] proof shows that every 1–handle stabilisation is a trivial stabilisation. He applied [23, Theorem 4] of Hudson for the statement that D^1 cores of handle additions that are homotopic rel endpoints are in fact smoothly ambiently isotopic fixing the endpoints. To apply Boyle’s work to topologically embedded surfaces in a compact 4–manifold N , remove a point from N and smooth $N \setminus \{\text{pt}\}$ in such a way that Σ is smoothly embedded. Then Boyle’s application of Hudson’s result yields a smooth ambient isotopy, which gives rise to a topological ambient isotopy once the point is added back to N . We note that Boyle works in S^4 , but his proof applies to any oriented ambient 4–manifold.

We therefore have that after finitely many trivial stabilisations, Σ_0 and Σ_1 are ambiently isotopic in N relative to the constant isotopy on the boundary, as desired. □

References

- [1] **H Bass**, *Unitary algebraic K -theory*, from “Algebraic K -theory, III: Hermitian K -theory and geometric applications” (H Bass, editor), Lecture Notes in Math. 343, Springer (1973) 57–265 MR Zbl
- [2] **R İ Baykur, N Sunukjian**, *Knotted surfaces in 4-manifolds and stabilizations*, J. Topol. 9 (2016) 215–231 MR Zbl
- [3] **M Borodzik, S Friedl**, *On the algebraic unknotting number*, Trans. London Math. Soc. 1 (2014) 57–84 MR Zbl
- [4] **M Borodzik, S Friedl**, *The unknotting number and classical invariants, I*, Algebr. Geom. Topol. 15 (2015) 85–135 MR Zbl
- [5] **S Boyer**, *Simply-connected 4-manifolds with a given boundary*, Trans. Amer. Math. Soc. 298 (1986) 331–357 MR Zbl
- [6] **J Boyle**, *Classifying 1-handles attached to knotted surfaces*, Trans. Amer. Math. Soc. 306 (1988) 475–487 MR Zbl
- [7] **G E Bredon**, *Topology and geometry*, Graduate Texts in Math. 139, Springer (1993) MR Zbl
- [8] **T D Cochran, K E Orr, P Teichner**, *Structure in the classical knot concordance group*, Comment. Math. Helv. 79 (2004) 105–123 MR Zbl
- [9] **A Conway, S Friedl, E Toffoli**, *The Blanchfield pairing of colored links*, Indiana Univ. Math. J. 67 (2018) 2151–2180 MR Zbl
- [10] **A Conway, M Powell**, *Characterisation of homotopy ribbon discs*, Adv. Math. 391 (2021) art. id. 107960 MR Zbl
- [11] **D J Crowley**, *The classification of highly connected manifolds in dimensions 7 and 15*, PhD thesis, Indiana University (2002) MR Available at <https://www.proquest.com/docview/305508775>
- [12] **D Crowley, J Sixt**, *Stably diffeomorphic manifolds and $l_{2q+1}(\mathbb{Z}[\pi])$* , Forum Math. 23 (2011) 483–538 MR Zbl
- [13] **B Farb, D Margalit**, *A primer on mapping class groups*, Princeton Mathematical Series 49, Princeton Univ. Press (2012) MR Zbl
- [14] **R Fintushel, R J Stern**, *Surfaces in 4-manifolds*, Math. Res. Lett. 4 (1997) 907–914 MR Zbl
- [15] **M H Freedman, F Quinn**, *Topology of 4-manifolds*, Princeton Mathematical Series 39, Princeton Univ. Press (1990) MR Zbl
- [16] **S Friedl, C Leidy, M Nagel, M Powell**, *Twisted Blanchfield pairings and decompositions of 3-manifolds*, Homology Homotopy Appl. 19 (2017) 275–287 MR Zbl
- [17] **S Friedl, M Nagel, P Orson, M Powell**, *A survey of the foundations of four-manifold theory in the topological category*, preprint (2019) arXiv 1910.07372

- [18] **S Friedl, M Powell**, *A calculation of Blanchfield pairings of 3–manifolds and knots*, Mosc. Math. J. 17 (2017) 59–77 MR Zbl
- [19] **RE Gompf, AI Stipsicz**, *4–Manifolds and Kirby calculus*, Graduate Studies in Math. 20, Amer. Math. Soc. (1999) MR Zbl
- [20] **I Hambleton, M Kreck, P Teichner**, *Topological 4–manifolds with geometrically two-dimensional fundamental groups*, J. Topol. Anal. 1 (2009) 123–151 MR Zbl
- [21] **I Hambleton, P Teichner**, *A non-extended Hermitian form over $\mathbb{Z}[\mathbb{Z}]$* , Manuscripta Math. 93 (1997) 435–442 MR Zbl
- [22] **JA Hillman, A Kawauchi**, *Unknotting orientable surfaces in the 4–sphere*, J. Knot Theory Ramifications 4 (1995) 213–224 MR Zbl
- [23] **JFP Hudson**, *Embeddings of bounded manifolds*, Proc. Cambridge Philos. Soc. 72 (1972) 11–20 MR Zbl
- [24] **A Juhász, M Miller, I Zemke**, *Transverse invariants and exotic surfaces in the 4–ball*, Geom. Topol. 25 (2021) 2963–3012 MR Zbl
- [25] **A Juhász, I Zemke**, *Stabilization distance bounds from link Floer homology*, preprint (2018) arXiv 1810.09158
- [26] **D Kasprowski, M Powell, P Teichner**, *Algebraic criteria for stable diffeomorphism of spin 4–manifolds*, preprint (2020) arXiv 2006.06127
- [27] **A Kawauchi**, *Splitting a 4–manifold with infinite cyclic fundamental group*, Osaka J. Math. 31 (1994) 489–495 MR Zbl
- [28] **A Kawauchi**, *Splitting a 4–manifold with infinite cyclic fundamental group, revised*, J. Knot Theory Ramifications 22 (2013) art. id. 1350081 MR Zbl
- [29] **Q Khan**, *Cancellation for 4–manifolds with virtually abelian fundamental group*, Topology Appl. 220 (2017) 14–30 MR Zbl
- [30] **HJ Kim**, *Modifying surfaces in 4–manifolds by twist spinning*, Geom. Topol. 10 (2006) 27–56 MR Zbl
- [31] **HJ Kim, D Ruberman**, *Smooth surfaces with non-simply-connected complements*, Algebr. Geom. Topol. 8 (2008) 2263–2287 MR Zbl
- [32] **HJ Kim, D Ruberman**, *Topological triviality of smoothly knotted surfaces in 4–manifolds*, Trans. Amer. Math. Soc. 360 (2008) 5869–5881 MR Zbl
- [33] **R Kirby**, *Problems in low dimensional manifold theory*, from “Algebraic and geometric topology, II” (RJ Milgram, editor), Proc. Sympos. Pure Math. 32, Amer. Math. Soc., Providence, RI (1978) 273–312 MR Zbl
- [34] **M-A Knus**, *Quadratic and Hermitian forms over rings*, Grundle. Math. Wissen. 294, Springer (1991) MR Zbl
- [35] **KH Ko**, *A Seifert-matrix interpretation of Cappell and Shaneson’s approach to link cobordisms*, Math. Proc. Cambridge Philos. Soc. 106 (1989) 531–545 MR Zbl

- [36] **M Kreck**, *Isotopy classes of diffeomorphisms of $(k-1)$ -connected almost-parallelizable $2k$ -manifolds*, from “Algebraic topology” (JL Dupont, IH Madsen, editors), Lecture Notes in Math. 763, Springer (1979) 643–663 MR Zbl
- [37] **M Kreck**, *Surgery and duality*, Ann. of Math. 149 (1999) 707–754 MR Zbl
- [38] **T Y Lam**, *Serre’s problem on projective modules*, Springer (2006) MR Zbl
- [39] **J Levine**, *Knot modules, I*, Trans. Amer. Math. Soc. 229 (1977) 1–50 MR Zbl
- [40] **C Livingston**, *Surfaces bounding the unlink*, Michigan Math. J. 29 (1982) 289–298 MR Zbl
- [41] **B A Magurn, W van der Kallen, LN Vaserstein**, *Absolute stable rank and Witt cancellation for noncommutative rings*, Invent. Math. 91 (1988) 525–542 MR Zbl
- [42] **T E Mark**, *Knotted surfaces in 4-manifolds*, Forum Math. 25 (2013) 597–637 MR Zbl
- [43] **T Oba**, *Surfaces in the 4-disk with the same boundary and fundamental group*, Math. Res. Lett. 27 (2020) 265–279 MR Zbl
- [44] **B Perron**, *Pseudo-isotopies et isotopies en dimension quatre dans la catégorie topologique*, Topology 25 (1986) 381–397 MR Zbl
- [45] **F Quinn**, *Isotopy of 4-manifolds*, J. Differential Geom. 24 (1986) 343–372 MR Zbl
- [46] **A Ranicki**, *The algebraic theory of surgery, I: Foundations*, Proc. London Math. Soc. 40 (1980) 87–192 MR Zbl
- [47] **A Ranicki**, *Exact sequences in the algebraic theory of surgery*, Mathematical Notes 26, Princeton Univ. Press (1981) MR Zbl
- [48] **R Stong, Z Wang**, *Self-homeomorphisms of 4-manifolds with fundamental group \mathbb{Z}* , Topology Appl. 106 (2000) 49–56 MR Zbl
- [49] **NS Sunukjian**, *Surfaces in 4-manifolds: concordance, isotopy, and surgery*, Int. Math. Res. Not. 2015 (2015) 7950–7978 MR Zbl

Department of Mathematics, Massachusetts Institute of Technology
Cambridge, MA, United States

School of Mathematics and Statistics, University of Glasgow
Glasgow, United Kingdom

anthonyconway@gmail.com, mark.powell@glasgow.ac.uk

Proposed: András I Stipsicz

Received: 11 November 2020

Seconded: Ciprian Manolescu, David Gabai

Revised: 5 August 2021

GEOMETRY & TOPOLOGY

msp.org/gt

MANAGING EDITOR

András I. Stipsicz Alfréd Rényi Institute of Mathematics
stipsicz@renyi.hu

BOARD OF EDITORS

Dan Abramovich	Brown University dan_abramovich@brown.edu	Mark Gross	University of Cambridge mgross@dpms.cam.ac.uk
Ian Agol	University of California, Berkeley ianagol@math.berkeley.edu	Rob Kirby	University of California, Berkeley kirby@math.berkeley.edu
Mark Behrens	Massachusetts Institute of Technology mbehrens@math.mit.edu	Frances Kirwan	University of Oxford frances.kirwan@balliol.oxford.ac.uk
Mladen Bestvina	Imperial College, London bestvina@math.utah.edu	Bruce Kleiner	NYU, Courant Institute bkleiner@cims.nyu.edu
Martin R. Bridson	Imperial College, London m.bridson@ic.ac.uk	Urs Lang	ETH Zürich urs.lang@math.ethz.ch
Jim Bryan	University of British Columbia jbryan@math.ubc.ca	Marc Levine	Universität Duisburg-Essen marc.levine@uni-due.de
Dmitri Burago	Pennsylvania State University burago@math.psu.edu	John Lott	University of California, Berkeley lott@math.berkeley.edu
Ralph Cohen	Stanford University ralph@math.stanford.edu	Ciprian Manolescu	University of California, Los Angeles cm@math.ucla.edu
Tobias H. Colding	Massachusetts Institute of Technology colding@math.mit.edu	Haynes Miller	Massachusetts Institute of Technology hrm@math.mit.edu
Simon Donaldson	Imperial College, London s.donaldson@ic.ac.uk	Tom Mrowka	Massachusetts Institute of Technology mrowka@math.mit.edu
Yasha Eliashberg	Stanford University eliash-gt@math.stanford.edu	Walter Neumann	Columbia University neumann@math.columbia.edu
Benson Farb	University of Chicago farb@math.uchicago.edu	Jean-Pierre Otal	Université d'Orleans jean-pierre.otal@univ-orleans.fr
Steve Ferry	Rutgers University sferry@math.rutgers.edu	Peter Ozsváth	Columbia University oszvath@math.columbia.edu
Ron Fintushel	Michigan State University ronfint@math.msu.edu	Leonid Polterovich	Tel Aviv University polterov@post.tau.ac.il
David M. Fisher	Rice University davidfisher@rice.edu	Colin Rourke	University of Warwick gt@maths.warwick.ac.uk
Mike Freedman	Microsoft Research michaelf@microsoft.com	Stefan Schwede	Universität Bonn schwede@math.uni-bonn.de
David Gabai	Princeton University gabai@princeton.edu	Peter Teichner	University of California, Berkeley teichner@math.berkeley.edu
Stavros Garoufalidis	Southern U. of Sci. and Tech., China stavros@mpim-bonn.mpg.de	Richard P. Thomas	Imperial College, London richard.thomas@imperial.ac.uk
Cameron Gordon	University of Texas gordon@math.utexas.edu	Gang Tian	Massachusetts Institute of Technology tian@math.mit.edu
Lothar Götsche	Abdus Salam Int. Centre for Th. Physics gotsche@ictp.trieste.it	Ulrike Tillmann	Oxford University tillmann@maths.ox.ac.uk
Jesper Grodal	University of Copenhagen jg@math.ku.dk	Nathalie Wahl	University of Copenhagen wahl@math.ku.dk
Misha Gromov	IHÉS and NYU, Courant Institute gromov@ihes.fr	Anna Wienhard	Universität Heidelberg wienhard@mathi.uni-heidelberg.de


See inside back cover or msp.org/gt for submission instructions.

The subscription price for 2023 is US \$740/year for the electronic version, and \$1030/year (+ \$70, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP. Geometry & Topology is indexed by Mathematical Reviews, Zentralblatt MATH, Current Mathematical Publications and the Science Citation Index.

Geometry & Topology (ISSN 1465-3060 printed, 1364-0380 electronic) is published 9 times per year and continuously online, by Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840. Periodical rate postage paid at Oakland, CA 94615-9651, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840.

GT peer review and production are managed by EditFLOW[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2023 Mathematical Sciences Publishers

GEOMETRY & TOPOLOGY

Volume 27 Issue 2 (pages 417–821) 2023

Convex cocompact actions of relatively hyperbolic groups	417
MITUL ISLAM and ANDREW ZIMMER	
Anosov groups: local mixing, counting and equidistribution	513
SAMUEL EDWARDS, MINJU LEE and HEE OH	
On cubulated relatively hyperbolic groups	575
EDUARDO REYES	
Tautological classes of definite 4–manifolds	641
DAVID BARAGLIA	
Prime-localized Weinstein subdomains	699
OLEG LAZAREV and ZACHARY SYLVAN	
Embedded surfaces with infinite cyclic knot group	739
ANTHONY CONWAY and MARK POWELL	