



Towards an understanding of multimodal traits of female reproduction in chimpanzees

Marlen Kücklich^{1,2} · Susann Jänig^{1,2} · Lars Kulik^{1,2} · Claudia Birkemeyer³ · Brigitte M. Weiß^{1,2} · Anja Widdig^{1,2}

Received: 3 December 2021 / Accepted: 7 June 2022
© The Author(s) 2022

Abstract

Although primates have long been regarded as microsmatic, recent studies indicate that olfaction is an important sensory mode of primate communication, e.g., in the context of reproduction. However, large gaps remain in our understanding of primate olfactory traits, especially in the great apes. Female chimpanzees (*Pan troglodytes*) possess an exaggerated sexual swelling, which is an imprecise signal of fertility that is thought to serve to confuse paternity. However, some high-ranking males that copulate most frequently on the days when females are most fertile seem to have more precise information on the timing of ovulation, which suggests the existence of an olfactory fertility trait. In order to examine, and provide evidence for, fertility-related chemical information in female chimpanzees, we used gas chromatography—mass spectrometry to analyze the chemical composition of female body odor collected across the menstrual cycle during various stages of sexual swelling (97 samples of six females). The chemical composition was significantly affected by swelling stage, and eight substances were detected that were strongly related to the latter. The existence of an additional, olfactory, fertility trait may help males to fine-tune their sexual behavior or allow females to strengthen concealment of the exact timing of ovulation, and needs to be further investigated in follow-up studies. The results of our study provide much-needed evidence for the existence of an olfactory cue related to reproduction in chimpanzees, and form a basis for future studies on the interplay between visual and olfactory information on female fertility.

Keywords Olfactory communication · Olfaction · Gas chromatography—mass spectrometry · Chemical composition · Menstrual cycle · Anogenital swelling

Marlen Kücklich and Susann Jänig contributed equally to this study as first authors, and Brigitte M. Weiß and Anja Widdig contributed equally to this study as senior authors.

✉ Marlen Kücklich
marlen_kuecklich@eva.mpg.de

¹ Behavioural Ecology Research Group, Faculty of Life Sciences, University of Leipzig, Talstraße 33, 04103 Leipzig, Germany

² Primate Behavioural Ecology Research Group, Department of Human Behavior, Ecology and Culture, Max Planck Institute for Evolutionary Anthropology, Deutscher Platz 6, 04103 Leipzig, Germany

³ Mass Spectrometry Research Group, Institute of Analytical Chemistry, University of Leipzig, Linnéstraße 3, 04103 Leipzig, Germany

Introduction

Primates utilize various sensory modalities, including vision, acoustics and olfaction, for social communication, e.g., to convey information regarding kinship [olfaction in *Lemur catta* (Boulet et al. 2009); vision and acoustics in *Macaca mulatta* (Pfefferle et al. 2014a, b)] or dominance rank [vision in *Mandrillus leucophaeus* (Marty et al. 2009); acoustics in *Macaca nigra* (Neumann et al. 2010); olfaction in *Mandrillus sphinx* (Vaglio et al. 2016)]. Sensory modalities involved in the transfer of specific information may act either alone or as multimodal traits, whereby different modalities transmit either redundant or complementary information (Partan and Marler 1999). The fertile phase of female primates, e.g., is frequently indicated by a female's behavior, such as approaching males or reducing aggression, by visual traits such as sexual swellings, acoustic traits such as advertisement or copulation calls, as well as olfactory

traits such as scent-marking behavior, in various combinations [e.g., *Cebuella pygmaea* (Converse et al. 1995); *Microcebus murinus* (Buesching et al. 1998); *Papio ursinus* (Clarke et al. 2009); *Papio anubis* (Rigaill et al. 2013)].

Whereas visual and acoustic traits of female fertility have been relatively well studied across primates, studies on olfactory traits of fertility are comparatively scarce (Drea 2015). However, over the years, it has become evident that olfaction may also play an important role in the transmission of social information in primates (e.g., Smith and Bhatnagar 2004; Drea et al. 2013). Studies in several primate species indicate that females convey olfactory information regarding their reproductive state through whole body odor, vaginal secretions, urine and feces (Drea et al. 2015). Furthermore, olfactory inspection of the female anogenital region by males increases during the fertile period [*P. ursinus* (Clarke et al. 2009); *P. anubis* (Rigaill et al. 2013)], and males react with increased exploratory behavior and sexual activity [*Macaca mulatta* (Michael et al. 1971); *Galago crassicaudatus* (Clark 1982); *Saguinus oedipus* (Ziegler et al. 1993); *Callithrix jacchus* (Smith and Abbott 1998); *Macaca arctoides* (Cerdá-Molina et al. 2006b)] as well as a rise in testosterone levels and activation of brain areas linked to sexual behavior [*C. jacchus* (Ziegler et al. 2005; Snowdon et al. 2006); *M. arctoides* (Cerdá-Molina et al. 2006a)] when presented with vaginal secretions from fertile females.

Specific substances related to the menstrual cycle of females have recently been identified, and males were observed to distinguish between female odor from different phases of the cycle in *C. jacchus* [an aromatic hydrocarbon, a heterocyclic compound, an ester and an alcohol (Kücklich et al. 2019)]. Certain substances were also found to be fertility related and to sexually stimulate male Old World monkeys [fatty acids in *M. mulatta* (Michael et al. 1971)], although the results could not be reproduced in a subsequent study [*M. mulatta* (Goldfoot et al. 1976)]. These types of fertility indicators could constitute “signals”, i.e., traits that have specifically evolved to provide information on fertility, or “cues”, which constitute traits that have not evolved to deliver fertility-related information to conspecifics, e.g., being by-products of changes in hormone levels (Wyatt 2014).

Chimpanzees (*Pan troglodytes*) are prime candidates for the study of multimodal traits of fertility, as female chimpanzees have several traits that may be indicative of their reproductive state during their menstrual cycle, which is around 36 days long (Tutin 1979; Deschner et al. 2003; Emery and Whitten 2003). They make copulation calls, although these seem not to vary with fertility (Townsend et al. 2011), and show proceptive as well as mating resistance behavior that varies between their fertile and non-fertile periods (Stumpf and Boesch 2006). In addition, they have exaggerated sexual swellings, which vary in size across their cycle, although

these are not a precise signal of fertility (Emery Thompson 2005). The phase of increased swelling lasts for approximately 10–12 days (Emery and Whitten 2003), and indicates approaching ovulation, which occurs during the period of maximal swelling (Deschner et al. 2004). However, maximal swelling persists for up to 4 days after ovulation (Deschner et al. 2004), and is thus only an imprecise signal of the latter. This imprecise information on the actual timing of ovulation allows females to confuse paternity and reduce the risk of infanticide by attracting a high number of mating partners (Wrangham 2002; Kappeler and van Schaik 2004; Stumpf and Boesch 2006). Males compete intensely with each other over their access to females (Goodall 1986; Müller 2002), which restricts their mating opportunities [priority of access model (Altmann 1962)]. The potential for males to monopolize females is further reduced since several females of a group will show exaggerated sexual swellings synchronously (Goodall 1986; Ostner et al. 2008). As males should focus their mating efforts on the days when females are most fertile in order to mate with as many fertile females as possible, they would benefit from using precise indicators of female fertility, if available. Indeed, male chimpanzees alter their sexual behavior when female anogenital swellings increase in size and throughout the period of maximal swelling (Klinkova et al. 2005), and mating seems to be correlated with the period of peak fertility of females (Deschner et al. 2004; Emery Thompson 2005). Alpha males, in particular, were found to copulate more frequently during the fertile period compared to the other days of the maximal swelling phase (Deschner et al. 2004). These observations suggest that high-ranking males may benefit from additional information, e.g., a multimodal cue of fertility. Rigaill et al. (2013) hypothesized that the visual signal in *P. anubis* could provide a first sign of fertility to all males within visual range, whereas high-ranking males that can approach females closely could potentially gather more precise information from an olfactory inspection of females with sexual swellings. Also, females could benefit from a more precise indicator of ovulation that is only sensed by selected, competitive males, to obtain the best genes and protection for their infants (Kappeler and van Schaik 2004).

Observations of increased copulation rates at the time of highest fertility in females suggest that male chimpanzees may indeed rely on olfactory traits in a sexual context (Emery Thompson and Wrangham 2008). Certain fatty acids that are thought to be associated with female fertility in rhesus macaques (*Macaca mulatta*) (Michael et al. 1971) were also found in the vaginal secretions of female chimpanzees, but it remains unclear if these vary during the menstrual cycle (Fox 1982; Matsumoto-Oda et al. 2003). Notably, fertility-related changes in the odor of female chimpanzees could occur in genital odor and/or in the scent of the whole body (hereafter “body odor”).

Several other individual attributes have been detected in the body odor of chimpanzees and other great apes [e.g., indicating individual identity (Hepper and Wells 2010); arousal level (Klailova and Lee 2014); species identity and age (Jänig et al. 2019)]. Chimpanzees sniff other individuals (i.e., genitalia, but also other body parts), and males specifically have been observed to sniff their conspecifics more often than females do (Matsumoto-Oda et al. 2007; Jänig et al. 2018). Accordingly, we expect information about reproductive state to be provided by the body odor of female chimpanzees in general, and not only by their genital odor, which may constitute a multimodal trait of fertility in combination with the visual signal of fertility, i.e., the sexual swelling.

Given the lack of studies demonstrating the existence of olfactory fertility traits in chimpanzees, our overarching aim was to provide evidence, based on chemical analyses, that olfactory information exists that is related to reproduction. Chimpanzees live in large, multi-male/multi-female groups, which split up regularly into smaller parties of variable composition (Nishida 1968; Goodall 1986). Males preferably join parties which include females with maximal sexual swellings (Matsumoto-Oda 1999). If an olfactory fertility cue exists that is even more precise than the visual signal, this could potentially help males to specifically join parties with females close to ovulation. Cycle-dependent olfactory traits could be subtle changes that arise as a result of physiological changes, such as the breakdown of hormones involved in the menstrual cycle, e.g., luteinizing hormone, follicle-stimulating hormone, progesterone and estradiol (Nadler et al. 1985). Sexual swelling appears to be induced by a rise in the levels of estrogens, and detumescence is correlated with a rise in progesterone after ovulation (Graham et al. 1972). Hence, olfactory changes could occur as by-products of these hormonal changes that accordingly follow the timing of sexual swellings, and thus act as similarly imprecise fertility traits as the visual signal. Ovulation can be assumed to occur 1 day before the increase in the level of progesterone (Deschner et al. 2004), and is related to pre-ovulatory peaks of luteinizing hormone and follicle-stimulating hormone which are preceded by an estradiol peak mid-cycle (Nadler et al. 1985). Thus, a hormone-related olfactory change could also occur at the time of ovulation. If this were true, olfactory information on fertility would be more precise than that provided by the visual signal. On the other hand, an imprecise olfactory cue could strengthen concealment of the actual timing of ovulation, and thus support the visual signal and allow females to confuse paternity. A more precise olfactory cue, however, would allow high-ranking males in particular to monitor females with maximal swelling to pinpoint the exact time of ovulation. Either way, the composition of female body odor would be expected, at the very least, to vary between swelling stages.

In this study, we related the chemical profiles of body odor samples of regularly cycling female chimpanzees to their stage of sexual swelling. We investigated whether chemical composition varies synchronously with the visual signal, and predicted a change in the former across the sexual cycle. If this prediction were true, it would suggest that anogenital swelling and body odor potentially comprise multimodal traits of fertility. For the purposes of this study, we used data collected as part of a larger study on various traits of great ape body odor [species identity, age and sex (Jänig et al. 2019)]. However, at the time of data collection, we were not able to obtain hormone samples from females in a reliable manner that would have enabled us to determine the exact timing of ovulation. Hence, given the lack of hormonal data, we cannot show whether an olfactory cue exists that is a more precise indicator of fertility than the visual signal. Our results nonetheless provide information on whether olfaction is related to reproduction at all, and thus provide a foundation for future studies that include an investigation of hormones in this context.

Materials and methods

Subjects

We conducted our study on captive chimpanzees (*Pan troglodytes verus*) at the Wolfgang Köhler Primate Research Centre (WKPRC) at Leipzig Zoo, Germany, in 2013. We analyzed samples from six adult females (aged between 13 and 20 years) from two different social groups (for details on housing and group composition, see supplementary Online Resource), which had a regular menstrual cycle, i.e., did not receive contraceptives (for at least 12 months prior to the study) and were not lactating.

Sexual swelling stages

The sexual swelling stage of each female was routinely recorded by the zookeepers and categorized as one of the following: flat, increasing in size, maximal tumescence, decreasing in size. Mean cycle length was 35.6 ± 9.1 days and thus within the ranges previously published for chimpanzees [captive (Nadler et al. 1985); wild (Tutin 1979; Wallis 1997; Emery Thompson 2005)]. On average, the observed females were detumescent for 16.7 ± 5.4 days, the swellings increased in size for 4.3 ± 5.5 days, remained at maximal tumescence for 12.3 ± 3.4 days, and decreased in size for 2.4 ± 1.5 days, similar to data reported for wild chimpanzees (Deschner et al. 2003).

Sample collection and preparation

The examined individuals are well trained in participating in behavioral experiments by the WKPRC. Samples of skin odor were collected in the morning prior to behavioral experiments. For the purpose of our study, individuals were trained by positive reinforcement (using food items) to come to the grid of the cage to participate voluntarily in the sampling by presenting a self-selected body part. Clean Lilibe cotton swabs (60% cotton wool, 25% polyester microfiber, 15% polyester), heated for 30 min at 130 °C before use (Birkemeyer et al. 2016), were held with sterilized metal tweezers and rubbed repeatedly over the skin and fur for approximately 20 s (similar to Célérier et al. 2010; Stoffel et al. 2015). The cotton swabs were immediately placed in pre-cleaned (washed with methanol and diethyl-ether) 4-mL glass vials (Rotilabo) and stored at -80 °C until gas chromatography—mass spectrometry (GC–MS) analysis.

Given that the individual chimpanzees had full control over which body part they presented, the samples included odor from the arms, belly, back and legs, but not the genitals, as these were not readily presented by the animals. Hence, we sampled body parts that have the same types of apocrine and eccrine sweat glands (Ellis and Montagna 1962; Montagna and Yun 1963) representing general body odor rather than genital odor. Body and genital odor should be similarly affected by hormonal changes across the menstrual cycle, as previously shown in humans (Michael et al. 1974, 1975; Gildersleeve et al. 2012). Furthermore, behavioral observations of both chimpanzee groups at the WKPRC revealed that when males sniff females they focus on the genitals in just over half of cases and other body parts in the remaining cases (Jänig et al., in review). We controlled for the different body parts in our statistical analysis. Samples were not collected from the mouth, hands or feet to reduce contamination with odor from food or the environment. Furthermore, diet was likely to have only a minor impact on body odor variance within and between individuals, since animals were fed daily with similar food.

In total, 97 samples [16.2 ± 2.3 samples per individual female covering from five to eight (mean 7.2) menstrual cycles each] were collected from the animals (for details see supplementary Online Resource Table 1). In addition, 42 control samples (pure cotton swabs not rubbed over the skin or fur but otherwise handled like the animal samples; one per sampling day) were collected to identify chemical substances that did not originate from the chimpanzees.

Chemical analysis

We extracted the chemical substances for GC–MS analysis by adding 1.2 mL of *n*-hexane (Sigma Aldrich, Steinheim, Germany) to the cotton swab in a glass vial (Birkemeyer

et al. 2016). The extract was concentrated stepwise to a volume of circa 60 μ L, and 4 μ L of this solution was injected into the GC (HP6890 Series GC System, Agilent, Waldbronn, Germany, with a HP5973 MSD Mass Selective Detector in electron-impact ionization mode at 70 eV) using splitless injection, for 2 min. The GC was equipped with a J&W Fisher DB35-MS column (30 m length, 0.25 mm inner diameter, 0.25 μ m film; Agilent), with the inlet temperature set to 250 °C. Helium was used as the carrier gas at a flow rate of 1.7 mL/min. The temperature program ran for 40.5 min, starting at 35 °C for 2 min, followed by a heating step of 10 °C/min until 320 °C, which was held for 10 min. The solvent delay was set to 7 min. The ion source operated at 250 °C and the scan range was set to m/z 50–550.

Data processing

From the GC–MS data we identified peaks, their retention times (RTs) and areas (intensity) using AMDIS v. 2.65 (Stein 1999). To determine substances that were repeatedly detected in our samples, we grouped consecutive RTs to one RT range (see also Weiß et al. 2018) using a self-written script performed in R version 3.2.3 (R Core Team 2015). We verified by manual inspection that mass spectral patterns of the peaks within a given RT range were consistent, i.e., each RT range was assumed to reflect one substance, resulting in 152 RT ranges. In the following, we thus use the term “substance” rather than “RT range.” At that stage of the study, the substances had not yet been chemically identified by using a library search, and were described only by their RT and specific patterns of signals with certain mass-to-charge-ratios (m/z).

We used a non-targeted approach to find substances that might be related to female fertility. We excluded 15 substances from the statistical analysis which had the same or a higher abundance in the control than in the animal samples (i.e., data on 137 of the 152 substances remained for analysis). For all the substances which were found to be potentially related to female fertility according to the statistical analysis, mass spectral comparisons using the National Institute of Standards and Technologies (NIST) Mass Spectral Library (NIST 14 software; NIST, Gaithersburg, MD) were conducted for their tentative identification. We report the best library hit per substance when (1) the top library hit was consistent over samples, (2) the match had a probability > 80 (NIST 14 value), and (3) the tentatively identified compound could be reasonably expected to elute at the given RT. Alternatively, where no library hit allowed tentative identification, we propose a structural classification based on interpretation of the spectrum. The mass spectra of the substances discussed in this article are provided in supplementary Online Resource Fig. 2.

Statistical analysis

First, we conducted an analysis of similarities (ANOSIM) to test whether chemical profiles of females at the same swelling stage were more similar to each other than profiles of females at different swelling stages. Following common practice (e.g., Stoffel et al. 2015), the ANOSIM was based on Bray–Curtis indices calculated from the log-transformed, standardized intensities of substances (intensity of substance divided by the summed intensity of all 137 substances \times 100) for each combination of sample dyads. To control for repeated measurements per female, we used a customized ANOSIM (R script written by LK) that computed *P*-values by permuting swelling stages within individual females only. The α -level was set to 0.05 for all statistical analyses.

Second, we implemented two two-tailed generalized linear mixed models in R v. 3.2.3 (R Core Team 2015) using the package lme4 v. 1.1.11 (Bates et al. 2015) to determine if specific substances are associated with a particular swelling stage. Chemical datasets of animals may present several analytical challenges, such as (1) substances of potential biological relevance may have very high, or very low intensities; (2) the total intensities of samples can change due to environmental factors; and (3) the intensity of certain substances (e.g., those considered contaminants, such as plasticizers) can vary strongly, and thus affect the relative composition of odor profiles (van den Berg et al. 2006). To overcome these problems, we fitted the two models using two different responses: *standardized* (intensity of substance divided by the summed intensity of all 137 substances per sample \times 100) and *transformed* [arcsine and $\log(x + 0.01)$] peak areas; *standardized* peak areas *centered* (to a mean of 0) and *scaled* (to a SD of 1). In both cases, standardization was used to correct for changing total sample intensities. In the first approach, transformation was applied to achieve a normal distribution and to reduce the relative impact of large substances, while differences in the relative abundances of the various substances were maintained. In the second approach, centering and scaling was applied to adjust all the substances to the same size and to give all of them equal weight in the analyses (van den Berg et al. 2006). The responses were vectorized from a multivariate data matrix (Jamil et al. 2013) of samples ($n = 97$) and substances ($n = 137$). Swelling stage (dummy coded and centered) was fitted as fixed effect test predictor, and age (*z*-transformed) of the females was fitted as control predictor. Sample number and substance identity (ID) were included as random effects to prevent pseudo-replication as well as heteroscedastic variance due to the vectorized data matrix (Jamil et al. 2013). Other random effects were the day of observation, the sampled body part, as well as ID and group of the female. We included the random slope of the fixed effects test predictor (swelling stage) within substance as the actual test predictor [for detailed

information, see Weiß et al. (2018)]. Additionally, swelling stage was fitted as a random slope within ID of the female to achieve more accurate estimates of the test predictor.

The check for normal distribution and homogeneity of the residuals by inspection of a *qq*-plot suggested no violation of assumptions. Plotting the residuals against fitted values revealed a slight bottom effect, but model stability indicated no influential cases. We determined variance inflation factors (VIF) to check for potential collinearity (Quinn and Keough 2002; Field 2005) calculated with the function *vif* of the package *car* (Fox and Weisberg 2011) and found no indication of collinearity ($VIF_{\max} = 1.01$).

We compared the full model to a null model excluding the random slope within substance by using a likelihood ratio test (LRT) (Barr et al. 2013) to determine the significance of the full model (Forstmeier and Schielzeth 2011). For the significant full model, the slope estimates of all substances were extracted from the model results. To determine the substances whose levels changed most with swelling stage, we compared slope estimates to the average slope estimate and focused on those substances whose absolute slope estimate was higher than the average absolute slope estimate + 2SDs (Weiß et al. 2018).

Results

Chemical profiles of females at the same swelling stage were significantly more similar than profiles of females at different swelling stages (ANOSIM, $R = 0.07$, $P = 0.026$), although the effect was rather small when whole profiles were considered.

The chemical composition of substances with log-transformed abundances was affected by swelling stage (LRT, $\chi^2 = 10.77$, $P = 0.013$). Thus, the relation of these affected substances to the rest of the respective odor profile changed depending on the swelling stage. This effect was most pronounced for four substances (i.e., their absolute slope estimate was higher than the absolute slope estimate + 2SDs). These substances were all tentatively identified as steroids (Table 1), and were least abundant in samples collected during the maximal swelling as well as the flat phase, and most abundant during the increase and decrease phases of swelling (Fig. 1).

For substances with centered and scaled abundances, chemical composition was also affected by swelling stage (LRT, $\chi^2 = 19.28$, $P < 0.001$). In these cases, the substance-specific abundances changed between the profiles of different swelling stages. The swelling stage effect was most pronounced for four of the substances (i.e., their absolute slope estimate was higher than the absolute slope estimate + 2SDs), which were different from the most affected substances of the log-transformed model. One of these


Table 1 Substances most affected by swelling stage with *log-transformed* abundances, including retention times (RTs), tentative identification [National Institute of Standards and Technologies (NIST) Mass Spectral Library; NIST 14] with probability (best hit) or structural

classification, substance class and the largest differences between the absolute slope estimates of swelling stages (average absolute slope estimate \pm SD, 0.096 ± 0.08)

RTs	Best library hit/structural classification	Probability	Substance class	Slope estimates
42.40	Unknown steroid (cholestadiene-like)	–	Steroid	0.59
43.49	Cholesta-2,4-diene	94	Steroid	0.32
44.39	Cholesta-3,5-diene	94	Steroid	0.38
48.46	Cholesta-5-en-3-ol (3.beta.)-, acetate	98	Steroid	0.38

For more details, see Fig. 1 and supplementary Online Resource Table 2; for mass spectra, see supplementary Online Resource Fig. 2

Fig. 1 a–d Log-transformed intensity (%) of the four substances most influenced by swelling stage (see also Table 1). Boxplots show medians and first and third quartiles. Retention times (RTs) and standardized peak area means \pm SD are as follows: **a** RT 42.40, 1.97 ± 2.46 ; **b** RT 43.49, 0.48 ± 0.43 ; **c** RT 44.39, 13.71 ± 8.45 ; **d** RT 48.46, 0.55 ± 0.56



substances was tentatively identified by the library search as a wax ester (Fig. 2a), whereas two substances had spectra with m/z patterns typical of long-chain alkylic structures (Fig. 2b, d), and one substance was thought to feature a phenyl substructure (Fig. 2c) based on its structural classification (see Table 2). Three of the substances were most abundant in samples collected during maximal swelling compared to the other swelling stages, while one of the substances with an alkylic (sub)structure was most abundant during the flat phase and least abundant in samples collected during maximal swelling (see Fig. 2).

Discussion

The results of the present study support the presumption that olfaction is related to reproduction in chimpanzees, as the chemical profiles of the female chimpanzees varied significantly with sexual swelling stage. Initial evidence was provided by the slightly greater similarities of the whole chemical profiles of samples taken during the same swelling stage compared to those taken at different swelling stages. These findings were corroborated by a

Fig. 2 a–d Centered and scaled intensity (%) of the four substances most influenced by swelling stage (see also Table 2). Boxplots show medians and first and third quartiles. RTs and standardized peak area means \pm SD are as follows: **a** RT 32.26, 0.26 ± 1.05 ; **b** RT 34.65, 0.12 ± 0.25 ; **c** RT 40.36, 0.03 ± 0.02 ; **d** RT 48.80, 0.35 ± 0.31




Table 2 Substances most affected by swelling stages with *centered* and *scaled* abundances, including RTs, tentative identification (NIST 14 library) with probability (best hit) or structural classifica-

tion, substance class and the largest differences between the absolute slope estimates of swelling stages (average absolute slope estimate \pm SD, 0.273 ± 0.20)

RT	Best library hit/structural classification	Probability	Substance class	Slope estimates
32.26	Dodecanoic acid, isoocetyl ester	91	Ester	0.87
34.65	Unknown [long-chain alkylic (sub)structure]	–	Unknown	0.91
40.36	Unknown phenol	–	Phenol	0.67
48.80	Unknown [long-chain alkylic (sub)structure]	–	Unknown	0.86

For more details, see Fig. 2 as well as supplementary Online Resource Table 2; for mass spectra, see supplementary Online Resource Fig. 2. For abbreviations, see Table 1

significant change in chemical composition across swelling stages that was most pronounced with respect to eight substances. Thus, the results reveal that olfactory changes exist that mirror changes in fertility at least as closely as the visual signal of sexual swelling. However, our methodological approach did not allow us to address whether olfactory cues provide more precise information which could allow males to pinpoint the time of ovulation more precisely. Hence, further studies that include hormonal measurements are needed to unravel the temporal dynamics of olfactory cues in the context of reproduction.

The effect of swelling stage on chemical composition was tested with two different approaches. Peak areas were either

log transformed or centered and scaled when used in the models. In the case of log-transformed peak areas, affected substances differ in contrast to the remaining chemical profile. Hence, information on the state of fertility seems to be provided by the status quo of the chemical composition, and comparisons over time are thus not necessary. When peak areas are centered and scaled, the analysis is focused on differences between, instead of within, the samples (van den Berg et al. 2006). Thus, particular substances that are found to be affected when peak areas are centered and scaled differ in their levels between samples (i.e., over time or swelling stages). In this scenario, it seems to be the variation (or differences) over the course of the menstrual cycle that

provides information on fertility stage. A male would thus need to regularly check a female's odor over time to unravel this information. As expected, the two approaches revealed different chemical substances that were most affected by swelling stage.

Four chemical substances were found to be affected by swelling stage when abundances were log-transformed. All of them were cholesterol derived, and were more abundant during the increasing and decreasing swelling stages compared to the maximal swelling and flat stage. Cholesterol-derived substances are ubiquitous in the chemical profiles of mammals (Charpentier et al. 2012) and are precursors for smaller molecules produced by bacterial degradation (Ezenwa and Williams 2014). Specifically, cholesterol is the precursor for steroid hormones such as estradiol and progesterone (Hu et al. 2010). Swelling of the anogenital skin is estrogen dependent, and detumescence occurs after a rise in progesterone, which inhibits the effects of estrogen (Gillman and Stein 1941; Graham et al. 1972; Emery and Whitten 2003). Tumescence of the sexual swelling occurs due to the intracellular accumulation of water in the anogenital tissue, which is lost during detumescence (Krohn and Zuckerman 1937; Clarke 1940). The consistent pattern of the four most affected substances, i.e., that their levels were highest during tumescence and detumescence, leads to the assumption that these chemical changes arise due to physiological changes that occur during swelling. This chemical information being consistent with the visual signal of increasing or decreasing swelling, could enable males to recognize the beginning and end of the female fertile period, allowing them to focus their monopolization and mating efforts on females which are most fertile.

Moreover, four substances with centered and scaled abundances were found to be related to swelling stage. Similar types of substances, such as long-chain fatty acid esters, substances with alkylic structures, and phenols, were previously detected, e.g., in skin emanations [humans (Bernier et al. 2000)] and in genital secretions [*L. catta* (with the exception of phenols) Boulet et al. 2009]. These four substances were related to the maximal swelling stage, as three of them were most abundant and the other one was least abundant during this phase compared to the other swelling stages. Overall, our results suggest that the chemical composition of female chimpanzee odor changes over the menstrual cycle in accordance with sexual swelling stage. Thus, body odor can be considered at least as a proxy of fertility similar to swelling stage, which has been well established as an imprecise fertility signal.

An olfactory trait related to reproduction is expected to be present consistently within the chemical profile of female odor to allow males to infer female fertility from it. Thus, we checked whether the most affected substances as shown by the two models were continuously present across

all the females and swelling stages. All of the substances were found for all six females as well as at all of the swelling stages, and most of them were present in at least 80% of the samples for at least two swelling stages. However, two of the substances indicated by the centered and scaled model (an alkylic compound with an RT of 34.65 min and an unknown phenol with an RT of 40.36 min) were only present in a maximum of 43% of samples per swelling stage (maximal swelling and increasing/decreasing swelling stages). If substances are not always present in the chemical profile of female odor, they are not very suitable for use by males as reliable indicators of female reproductive state. Future studies that include a higher number of samples of female odor will help researchers to examine this subject in more detail.

While our results indicate that body odor does indeed change over a female's reproductive cycle, our methodology did not allow us to assess whether olfactory changes are as precise as the visual signal, or potentially even more precise. In fact, in the context of an evolutionary arms race, we would expect males to strive to pinpoint the exact timing of ovulation by using a fertility trait, and thus that they would benefit from an olfactory cue that is more precise than the visual cue of sexual swelling. Females, on the other hand, would profit from generally confusing paternity by providing the rather imprecise visual signal of fertility which could be strengthened by an additional, imprecise olfactory cue. However, high-ranking male chimpanzees in particular have been observed to start copulating only in the last 3–4 days of maximal swelling (Deschner et al. 2004). As the duration of maximal swelling is variable and its termination cannot be deduced according to the point at which it starts, such observations suggest that high-ranking male chimpanzees in particular, which are able to approach females to within a very close distance, may rely on additional and more precise indicators of ovulation, i.e., presumably olfactory traits. In humans, odor has been proposed as a more precise indicator for (approaching) ovulation than visual signs (Singh and Bronstad 2001; Havlíček et al. 2006; Haselton and Gildersleeve 2011; Gildersleeve et al. 2012). Relying on more than one of the senses for the inspection of a female is likely to be advantageous for male chimpanzees too. The visual signal could give initial information about the state of fertility of a female at short/medium distance, which may then especially attract high-ranking males with priority of access, allowing them to undertake an olfactory inspection at close proximity to gain additional information (Matsumoto-Oda et al. 2007). Such olfactory information could potentially be non-redundant to the visual information (see also Higham and Hebets 2013), but our study design did not allow us to determine whether the olfactory cue is redundant or not.

Thus, an important next step building on our current results, that show that chemical composition changes over the menstrual cycle in accordance with stages of sexual

swelling, would be investigating whether chemical profiles indeed provide more precise information than sexual swellings. To investigate this in more detail, analyses of cycle-related hormones are needed to determine the exact timing of ovulation, which could not be accomplished at the time that this study was conducted. This could be undertaken by using non-invasive hormonal measurements, as applied in previous studies to assess the timing of ovulation in non-human primates (Deschner et al. 2004; Engelhardt et al. 2006; Dubuc et al. 2012; Young et al. 2013; Douglas et al. 2016).

Moreover, recently adapted sampling techniques now allow a wider range of chemical substances to be sampled (Kücklich et al. 2017). The traditionally used cotton swab, which was also used in this study, primarily captures semi-to non-volatile substances (Birkemeyer et al. 2016). These types of larger molecules are then degraded by skin bacteria into smaller, volatile molecules that constitute the odor profile of an individual (Ezenwa and Williams 2014) that can be perceived by the main olfactory system (Dulac and Torello 2003). Future studies will certainly benefit from these recent methodological developments for the direct sampling of volatile substances, and for the correlation of their abundances to individual traits of animals.

In conclusion, this study adds much-needed information on the role of olfaction in great apes by showing that the chemical composition of body odor is associated with a visual signal of fertility in chimpanzees. Our results indicate that olfactory cues related to reproduction exist in a great ape species with visual fertility advertisement, and provide a valuable basis for further research on more specific questions on reproduction in chimpanzees and olfaction in great apes in general. In fission–fusion societies, where group composition varies and visual information is short lived, olfaction may allow males to fine-tune their sociosexual behavior and to optimize their reproductive investment.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s10329-022-00995-1>.

Acknowledgements We thank Josep Call and the curators of the Leipzig Zoo for permission to sample the chimpanzees at the WKPRC, and are most grateful for the help of the keepers at Leipzig Zoo and the staff of the WKPRC. Furthermore, we also acknowledge the help of Susan Billig and Andreas Kiontke for technical assistance with the chemical analysis. We thank Ruth Thomsen for logistic support during sample collection, and Roman Wittig and the anonymous reviewers for comments on earlier versions of the manuscript. Finally, we thank the Max Planck Institute for Evolutionary Anthropology and the University of Leipzig for their logistic support.

Funding Open Access funding enabled and organized by Projekt DEAL. This work was supported by the German Research Foundation within the Emmy-Noether program (grant no. WI 1808/3-1 awarded to AW) and by the University of Leipzig. In addition, graduate funding

from the University of Leipzig was awarded to SJ (DFPL R00061) and MK (DFPL R00017).

Data availability The dataset generated and analyzed during the current study is available on reasonable request.

Declarations

Conflict of interest The authors declare that they have no conflict of interest.

Ethical approval This study was carried out in accordance with the legal requirements of Germany and all national and institutional guidelines for the care and use of animals, and was approved by the ethics commission of the Department of Psychology of the Max Planck Institute for Evolutionary Anthropology and Leipzig Zoo.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Altmann SA (1962) A field study of the sociobiology of rhesus monkeys, *Macaca mulatta*. Ann NY Acad Sci 102:338–435. <https://doi.org/10.1111/j.1749-6632.1962.tb13650.x>
- Barr DJ, Levy R, Scheepers C, Tily HJ (2013) Random effects structure for confirmatory hypothesis testing: keep it maximal. J Mem Lang 68:255–278. <https://doi.org/10.1016/j.jml.2012.11.001>
- Bates D, Maechler M, Bolker B, Walker S (2015) lme4: linear mixed-effects models using Eigen and S4, R package. Version 1.0–8. <http://CRAN.R-project.org/package=lme4>. Accessed 20 Jan 2021
- Bernier UR, Kline DL, Barnard DR et al (2000) Analysis of human skin emanations by gas chromatography/mass spectrometry. 2. Identification of volatile compounds that are candidate attractants for the yellow fever mosquito (*Aedes aegypti*). Anal Chem 72:747–756. <https://doi.org/10.1021/ac990963k>
- Birkemeyer C, Thomsen R, Jänig S et al (2016) The sampling of body odour of primates: cotton swabs sample semi-volatiles rather than volatiles. Chem Sens 41:525–535. <https://doi.org/10.1093/chemse/bjw056>
- Boulet M, Charpentier M, Drea C (2009) Decoding an olfactory mechanism of kin recognition and inbreeding avoidance in a primate. BMC Evol Biol 9:281. <https://doi.org/10.1186/1471-2148-9-281>
- Buesching CD, Heistermann M, Hodges JK, Zimmermann E (1998) Multimodal oestrus advertisement in a small nocturnal prosimian, *Microcebus murinus*. Folia Primatol 69:295–308. <https://doi.org/10.1159/000052718>

- C el erier A, Huchard E, Alvergne A et al (2010) Detective mice assess relatedness in baboons using olfactory cues. *J Exp Biol* 213:1399–1405. <https://doi.org/10.1242/jeb.038349>
- Cerda-Molina AL, Hernandez-Lopez L, Chavira R et al (2006a) Endocrine changes in male stump-tailed macaques (*Macaca arctoides*) as a response to odor stimulation with vaginal secretions. *Horm Behav* 49:81–87. <https://doi.org/10.1016/j.yhbeh.2005.04.014>
- Cerda-Molina AL, Hernandez-Lopez L, Rojas-Maya S et al (2006b) Male-induced sociosexual behavior by vaginal secretions in *Macaca arctoides*. *Int J Primatol* 27:791–807. <https://doi.org/10.1007/s10764-006-9045-0>
- Charpentier MJE, Barthes N, Proffit M et al (2012) Critical thinking in the chemical ecology of mammalian communication: roadmap for future studies. *Funct Ecol* 26:769–774. <https://doi.org/10.1111/j.1365-2435.2012.01998.x>
- Clark AB (1982) Scent marks as social signals in *Galago crassicaudatus*. I. Sex and reproductive status as factors in signals and responses. *J Chem Ecol* 8:1133–1151. <https://doi.org/10.1007/BF00986984>
- Clarke RW (1940) Water distribution and sexual skin of the baboon. *Am J Physiol Legacy Content* 131:325–330. <https://doi.org/10.1152/ajplegacy.1940.131.2.325>
- Clarke PMR, Barrett L, Henzi SP (2009) What role do olfactory cues play in chacma baboon mating? *Am J Primatol* 71:493–502. <https://doi.org/10.1002/ajp.20678>
- Converse LJ, Carlson AA, Ziegler TE, Snowdon CT (1995) Communication of ovulatory state to mates by female pygmy marmosets, *Cebuella pygmaea*. *Anim Behav* 49:615–621. [https://doi.org/10.1016/0003-3472\(95\)80194-4](https://doi.org/10.1016/0003-3472(95)80194-4)
- Deschner T, Heistermann M, Hodges K, Boesch C (2003) Timing and probability of ovulation in relation to sex skin swelling in wild West African chimpanzees, *Pan troglodytes verus*. *Anim Behav* 66:551–560. <https://doi.org/10.1006/anbe.2003.2210>
- Deschner T, Heistermann M, Hodges K, Boesch C (2004) Female sexual swelling size, timing of ovulation, and male behavior in wild West African chimpanzees. *Horm Behav* 46:204–215. <https://doi.org/10.1016/j.yhbeh.2004.03.013>
- Douglas PH, Hohmann G, Murtagh R et al (2016) Mixed messages: wild female bonobos show high variability in the timing of ovulation in relation to sexual swelling patterns. *BMC Evol Biol* 16:1–17. <https://doi.org/10.1186/s12862-016-0691-3>
- Drea CM (2015) D'scent of man: a comparative survey of primate chemosignaling in relation to sex. *Horm Behav* 68:117–133. <https://doi.org/10.1016/j.yhbeh.2014.08.001>
- Drea CM, Boulet M, Delbarco-Trillo J et al (2013) The “secret” in secretions: methodological considerations in deciphering primate olfactory communication. *Am J Primatol* 75:621–642. <https://doi.org/10.1002/ajp.22143>
- Dubuc C, Muniz L, Heistermann M et al (2012) Do males time their mate-guarding effort with the fertile phase in order to secure fertilisation in Cayo Santiago rhesus macaques? *Horm Behav* 61:696–705. <https://doi.org/10.1016/j.yhbeh.2012.03.003>
- Dulac C, Torello AT (2003) Molecular detection of pheromone signals in mammals: from genes to behaviour. *Nat Rev Neurosci* 4:551–562. <https://doi.org/10.1038/nrn1140>
- Ellis RA, Montagna W (1962) The skin of primates. VI. The skin of the gorilla (*Gorilla gorilla*). *Am J Phys Anthropol* 20:79–93. <https://doi.org/10.1002/ajpa.1330200210>
- Emery Thompson M (2005) Reproductive endocrinology of wild female chimpanzees (*Pan troglodytes schweinfurthii*): methodological considerations and the role of hormones in sex and conception. *Am J Primatol* 67:137–158. <https://doi.org/10.1002/ajp.20174>
- Emery Thompson M, Wrangham RW (2008) Male mating interest varies with female fecundity in *Pan troglodytes schweinfurthii* of Kanyawara, Kibale National Park. *Int J Primatol* 29:885–905. <https://doi.org/10.1007/s10764-008-9286-1>
- Emery MA, Whitten PL (2003) Size of sexual swellings reflects ovarian function in chimpanzees (*Pan troglodytes*). *Behav Ecol Sociobiol* 54:340–351. <https://doi.org/10.1007/s00265-003-0648-6>
- Engelhardt A, Heistermann M, Hodges JK et al (2006) Determinants of male reproductive success in wild long-tailed macaques (*Macaca fascicularis*)—male monopolisation, female mate choice or post-copulatory mechanisms? *Behav Ecol Sociobiol* 59:740–752. <https://doi.org/10.1007/s00265-005-0104-x>
- Ezenwa VO, Williams AE (2014) Microbes and animal olfactory communication: where do we go from here? *BioEssays* 36:847–854. <https://doi.org/10.1002/bies.201400016>
- Field A (2005) *Discovering statistics using SPSS*. Sage, London
- Forstmeier W, Schielzeth H (2011) Cryptic multiple hypotheses testing in linear models: overestimated effect sizes and the winner's curse. *Behav Ecol Sociobiol* 65:47–55. <https://doi.org/10.1007/s00265-010-1038-5>
- Fox GJ (1982) Potentials for pheromones in chimpanzee vaginal fatty acids. *Folia Primatol* 37:255–266. <https://doi.org/10.1159/000156036>
- Fox J, Weisberg S (2011) *An {R} companion to applied regression*, second edition. Sage, Thousand Oaks, CA. <http://socserv.socsci.mcmaster.ca/jfox/Books/Companion>. Accessed 20 Jan 2021
- Gildersleeve KA, Haselton MG, Larson CM, Pillsworth EG (2012) Body odor attractiveness as a cue of impending ovulation in women: evidence from a study using hormone-confirmed ovulation. *Horm Behav* 61:157–166. <https://doi.org/10.1016/j.yhbeh.2011.11.005>
- Gillman J, Stein HB (1941) A quantitative study of the inhibition of estradiol benzoate by progesterone in the baboon (*Papio porcarius*). *Endocrinology* 28:274–282. <https://doi.org/10.1210/endo-28-2-274>
- Goldfoot DA, Kravetz MA, Goy RW, Freeman SK (1976) Lack of effect of vaginal lavages and aliphatic acids on ejaculatory responses in rhesus monkeys: behavioral and chemical analyses. *Horm Behav* 7:1–27. [https://doi.org/10.1016/0018-506X\(76\)90001-5](https://doi.org/10.1016/0018-506X(76)90001-5)
- Goodall J (1986) *The chimpanzees of Gombe. Patterns of behavior*. Belknap, Harvard University, Cambridge, MA
- Graham CE, Collins DC, Robinson H, Preedy JRK (1972) Urinary levels of estrogens and pregnanediol and plasma levels of progesterone during the menstrual cycle of the chimpanzee: relationship to the sexual swelling. *Endocrinology* 91:13–24. <https://doi.org/10.1210/endo-91-1-13>
- Haselton MG, Gildersleeve K (2011) Can men detect ovulation? *Curr Dir Psychol Sci* 20:87–92. <https://doi.org/10.1177/0963721411402668>
- Havli ek J, Dvorakova R, Barto  L, Flegr J (2006) Non-advertized does not mean concealed: body odour changes across the human menstrual cycle. *Ethology* 112:81–90. <https://doi.org/10.1111/j.1439-0310.2006.01125.x>
- Hepper PG, Wells DL (2010) Individually identifiable body odors are produced by the gorilla and discriminated by humans. *Chem Senses* 35:263–268. <https://doi.org/10.1093/chemse/bjq015>
- Higham JP, Hebets EA (2013) An introduction to multimodal communication. *Behav Ecol Sociobiol* 67:1381–1388. <https://doi.org/10.1007/s00265-013-1590-x>
- Hu J, Zhang Z, Shen W-J, Azhar S (2010) Cellular cholesterol delivery, intracellular processing and utilization for biosynthesis of steroid hormones. *Nutr Metab (Lond)* 7:47. <https://doi.org/10.1186/1743-7075-7-47>
- Jamil T, Ozinga WA, Kleyer M, ter Braak CJF (2013) Selecting traits that explain species–environment relationships: a generalized

- linear mixed model approach. *J Veg Sci* 24:988–1000. <https://doi.org/10.1111/j.1654-1103.2012.12036.x>
- Jänig S, Weiß BM, Widdig A (2018) Comparing the sniffing behavior of great apes. *Am J Primatol* 80:e22872. <https://doi.org/10.1002/ajp.22872>
- Jänig S, Weiß BM, Birkemeyer C, Widdig A (2019) Comparative chemical analysis of body odor in great apes. *Am J Primatol*. <https://doi.org/10.1002/ajp.22976>
- Kappeler PM, van Schaik C (2004) Sexual selection in primates—new and comparative perspectives. Cambridge University Press, Cambridge
- Klailova M, Lee PC (2014) Wild western lowland gorillas signal selectively using odor. *PLoS ONE* 9:e99554. <https://doi.org/10.1371/journal.pone.0099554>
- Klinkova E, Hodges JK, Fuhrmann K et al (2005) Male dominance rank, female mate choice and male mating and reproductive success in captive chimpanzees. *Int J Primatol* 26:357–484. <https://doi.org/10.1007/s10764-005-2929-6>
- Krohn PL, Zuckerman S (1937) Water metabolism in relation to the menstrual cycle. *J Physiol* 88:369–387
- Kücklich M, Möller M, Marcillo A et al (2017) Different methods for volatile sampling in mammals. *PLoS ONE* 12:e0183440. <https://doi.org/10.1371/journal.pone.0183440>
- Kücklich M, Weiß BM, Birkemeyer C et al (2019) Chemical cues of female fertility states in a non-human primate. *Sci Rep* 9:1–12. <https://doi.org/10.1038/s41598-019-50063-w>
- Marty JS, Higham JP, Gadsby EL, Ross C (2009) Dominance, coloration, and social and sexual behavior in male drills *Mandrillus leucophaeus*. *Int J Primatol* 30:807–823. <https://doi.org/10.1007/s10764-009-9382-x>
- Matsumoto-Oda A (1999) Mahale chimpanzees: grouping patterns and cycling females. *Am J Primatol* 47:197–207. [https://doi.org/10.1002/\(SICI\)1098-2345\(1999\)47:3%3c197::AID-AJP2%3e3.0.CO;2-3](https://doi.org/10.1002/(SICI)1098-2345(1999)47:3%3c197::AID-AJP2%3e3.0.CO;2-3)
- Matsumoto-Oda A, Oda R, Hayashi Y et al (2003) Vaginal fatty acids produced by chimpanzees during menstrual cycles. *Folia Primatol* 74:75–79. <https://doi.org/10.1159/000070000>
- Matsumoto-Oda A, Kutsukake N, Hosaka K, Matsusaka T (2007) Sniffing behaviors in Mahale chimpanzees. *Primates* 48:81–85. <https://doi.org/10.1007/s10329-006-0006-1>
- Michael RP, Keverne EB, Bonsall RW (1971) Pheromones: isolation of male sex attractants from a female primate. *Science* 172:964–966. <https://doi.org/10.1126/science.172.3986.964>
- Michael RP, Bonsall RW, Warner P (1974) Human vaginal secretions: volatile fatty acid content. *Science* 186:1217–1219. [https://doi.org/10.1016/0306-4530\(75\)90007-4](https://doi.org/10.1016/0306-4530(75)90007-4)
- Michael RP, Bonsall RW, Kutner M (1975) Volatile fatty acids, “copulins”, in human vaginal secretions. *Psychoneuroendocrinology* 1:153–163. [https://doi.org/10.1016/0306-4530\(75\)90007-4](https://doi.org/10.1016/0306-4530(75)90007-4)
- Montagna W, Yun JS (1963) The skin of primates. XV. The skin of the chimpanzee (*Pan satyrus*). *Am J Phys Anthropol* 21:189–203. <https://doi.org/10.1002/ajpa.1330210211>
- Muller MN (2002) Agonistic relations among Kanyawara chimpanzees. In: Boesch C, Hohmann G, Marchant L (eds) Behavioural diversity in chimpanzees and bonobos. Cambridge University Press, pp 112–124
- Nadler RD, Graham CE, Gosselin RE, Collins DC (1985) Serum levels of gonadotropins and gonadal steroids, including testosterone, during the menstrual cycle of the chimpanzee (*Pan troglodytes*). *Am J Primatol* 9:273–284. <https://doi.org/10.1002/ajp.1350090404>
- Neumann C, Assahad G, Hammerschmidt K et al (2010) Loud calls in male crested macaques, *Macaca nigra*: a signal of dominance in a tolerant species. *Anim Behav* 79:187–193. <https://doi.org/10.1016/j.anbehav.2009.10.026>
- Nishida T (1968) The social group of wild chimpanzees in the Mahali Mountains. *Primates* 9:167–224. <https://doi.org/10.1007/BF01730971>
- Ostner J, Nunn CL, Schülke O (2008) Female reproductive synchrony predicts skewed paternity across primates. *Behav Ecol* 19:1150–1158. <https://doi.org/10.1093/beheco/arn093>
- Partan S, Marler P (1999) Communication goes multimodal. *Science* 283:72–73. <https://doi.org/10.1126/science.283.5406.1272>
- Pfefferle D, Kazem AJN, Brockhausen RR et al (2014a) Monkeys spontaneously discriminate their unfamiliar paternal kin under natural conditions using facial cues. *Curr Biol* 24:1806–1810. <https://doi.org/10.1016/j.cub.2014.06.058>
- Pfefferle D, Ruiz-Lambides AV, Widdig A (2014b) Female rhesus macaques discriminate unfamiliar paternal sisters in playback experiments—support for acoustic phenotype matching. *Proc R Soc B* 281:1–9
- Quinn GP, Keough MJ (2002) Experimental design and data analysis for biologists, 1st edn. Cambridge University Press, Cambridge
- R Core Team (2015) A language and environment for statistical computing. Version 3.2.3. R Foundation for Statistical Computing, Vienna. <http://www.R-project.org/>. Accessed 20 Jan 2021
- Rigaill L, Higham JP, Lee PC et al (2013) Multimodal sexual signaling and mating behavior in olive baboons (*Papio anubis*). *Am J Primatol* 75:774–787. <https://doi.org/10.1002/ajp.22154>
- Singh D, Bronstad PM (2001) Female body odour is a potential cue to ovulation. *Proc R Soc B* 268:797–801. <https://doi.org/10.1098/rspb.2001.1589>
- Smith TE, Abbott DH (1998) Behavioral discrimination between circumgenital odor from peri-ovulatory dominant and anovulatory female common marmosets (*Callithrix jacchus*). *Am J Primatol* 46:265–284. [https://doi.org/10.1002/\(SICI\)1098-2345\(1998\)46:4%3c265::AID-AJP1%3e3.0.CO;2-D](https://doi.org/10.1002/(SICI)1098-2345(1998)46:4%3c265::AID-AJP1%3e3.0.CO;2-D)
- Smith TD, Bhatnagar KP (2004) Microsmatic primates: reconsidering how and when size matters. *Anat Rec B* 279B:24–31. <https://doi.org/10.1002/ar.b.20026>
- Snowdon CT, Ziegler TE, Schultz-Darken NJ, Ferris CF (2006) Social odours, sexual arousal and pairbonding in primates. *Philos Trans R Soc B* 361:2079–2089. <https://doi.org/10.1098/rstb.2006.1932>
- Stein SE (1999) An integrated method for spectrum extraction and compound identification from gas chromatography/mass spectrometry data. *J Am Soc Mass Spectrom* 10:770–781. [https://doi.org/10.1016/S1044-0305\(99\)00047-1](https://doi.org/10.1016/S1044-0305(99)00047-1)
- Stoffel MA, Caspers BA, Forcada J et al (2015) Chemical fingerprints encode mother–offspring similarity, colony membership, relatedness, and genetic quality in fur seals. *PNAS* 112:E5005–E5012. <https://doi.org/10.1073/pnas.1506076112>
- Stumpf RM, Boesch C (2006) The efficacy of female choice in chimpanzees of the Tai Forest, Cote d’Ivoire. *Behav Ecol Sociobiol* 60:749–765. <https://doi.org/10.1007/s00265-006-0219-8>
- Townsend SW, Deschner T, Zuberbühler K (2011) Copulation calls in female chimpanzees (*Pan troglodytes schweinfurthii*) convey identity but do not accurately reflect fertility. *Int J Primatol* 32:914–923. <https://doi.org/10.1007/s10764-011-9510-2>
- Tutin CEG (1979) Mating patterns and reproductive strategies in a community of wild chimpanzees (*Pan troglodytes schweinfurthii*). *Behav Ecol Sociobiol* 6:29–38. <https://doi.org/10.1007/BF00293242>
- Vaglio S, Minicozzi P, Romoli R et al (2016) Sternal gland scent-marking signals sex, age, rank, and group identity in captive mandrills. *Chem Senses* 41:177–186. <https://doi.org/10.1093/chemse/bjv077>
- van den Berg RA, Hoefsloot HC, Westerhuis JA et al (2006) Centering, scaling, and transformations: improving the biological information content of metabolomics data. *BMC Genom* 7:142. <https://doi.org/10.1186/1471-2164-7-142>

- Wallis J (1997) A survey of reproductive parameters in the free-ranging chimpanzees of Gombe National Park. *J Reprod Fertil* 109:297–307
- Weiß BM, Marcillo A, Manser M et al (2018) A non-invasive method for sampling the body odour of mammals. *Methods Ecol Evol* 9:420–429. <https://doi.org/10.1111/2041-210X.12888>
- Wrangham RW (2002) The cost of sexual attraction: is there a trade-off in female *Pan* between sex appeal and received coercion? In: Boesch C, Hohmann G, Marchant L (eds) *Behavioural diversity in chimpanzees and bonobos*. Cambridge University Press, Cambridge, pp 204–215
- Wyatt TD (2014) *Pheromones and animal behavior: chemical signals and signatures*. Cambridge University Press
- Young C, Majolo B, Heistermann M et al (2013) Male mating behaviour in relation to female sexual swellings, socio-sexual behaviour and hormonal changes in wild Barbary macaques. *Horm Behav* 63:32–39. <https://doi.org/10.1016/j.yhbeh.2012.11.004>
- Ziegler TE, Epple G, Snowdon CT et al (1993) Detection of the chemical signals of ovulation in the cotton-top tamarin, *Saguinus oedipus*. *Anim Behav* 45:313–322. <https://doi.org/10.1006/anbe.1993.1036>
- Ziegler TE, Schultz-Darken NJ, Scott JJ et al (2005) Neuroendocrine response to female ovulatory odors depends upon social condition in male common marmosets, *Callithrix jacchus*. *Horm Behav* 47:56–64. <https://doi.org/10.1016/j.yhbeh.2004.08.009>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.