

# Kähler geometry of black holes and gravitational instantons

Steffen Aksteiner\* and Bernardo Araneda†

Max Planck Institute for Gravitational Physics (Albert Einstein Institute),  
Am Mühlenberg 1, D-14476 Potsdam, Germany

(Dated: July 21, 2022)

We obtain a closed formula for the Kähler potential of a broad class of four-dimensional Lorentzian or Euclidean conformal “Kähler” geometries, including the Plebański-Demiański class and various gravitational instantons such as Fubini-Study and Chen-Teo. We show that the Kähler potentials of Schwarzschild and Kerr are related by a Newman-Janis shift. Our method also shows that a class of supergravity black holes, including the Kerr-Sen spacetime, is Hermitian (but not conformal Kähler). We finally show that the integrability conditions of complex structures lead naturally to the (non-linear) Weyl double copy, and we give new vacuum and non-vacuum examples of this relation.

## INTRODUCTION

Complex methods as a tool to investigate space-time structure in General Relativity (GR) have a long and fruitful history of remarkable developments. Profound constructions, pioneered by Penrose, Newman, Plebański, Robinson, Trautman, [1–6] among others, include twistor theory and heavenly structures, but there are also simple yet intriguing results such as the “Newman-Janis shift” relating special solutions via complex coordinate transformations.

An important insight regarding complex structures in GR is provided by Flaherty [7], who showed that type D vacuum and Einstein-Maxwell spacetimes possess an analogue of the Hermitian structures of Riemannian geometry. In Lorentz signature, a Hermitian structure must necessarily be complex-valued, so its integrability properties are more subtle than in the Euclidean case. Flaherty gave a comprehensive analysis of such properties [8, 9], and he found that the above classes of type D spacetimes are not only Hermitian but also satisfy the Lorentzian analogue of the conformal Kähler condition.

In Riemannian geometry, Kähler metrics are encoded in “generating functions” or scalar Kähler potentials. An analogous feature in GR occurs in perturbation theory, where perturbative fields are generated by scalar Debye potentials. These potentials are instrumental for modern studies of e.g. black hole stability and gravitational wave physics. The increasing interest in nonperturbative structures for gravitational wave science [10], together with the importance of scalar potentials for perturbation theory, motivate the question of whether there are “Debye potentials” for exact, astrophysically relevant solutions of GR. Moreover, the recently discovered applications of the Newman-Janis shift [11, 12] suggest that complex structures in GR may play an important role in the understanding of such nonperturbative structures.

Motivated by the above considerations, in this paper we develop a method to find the Kähler potentials of a broad class of geometries, including black holes and gravitational instantons, and we show intimate connections of this approach with other theoretical structures

of modern interest for gravitational wave physics such as the Newman-Janis shift and the double copy relation between gauge and gravity theories.

As an example, consider the Kerr metric  $g$  with parameters  $M, a$  for mass and angular momentum per mass, respectively. In Boyer-Lindquist coordinates  $(t, r, \theta, \phi)$ , the metric is block diagonal in  $(dt, d\phi)$  and  $(dr, d\theta)$ , with components in the first block given by

$$\begin{aligned} g_{tt} &= (\Delta - a^2 \sin^2 \theta) \Sigma^{-1}, \\ g_{t\phi} &= -a(\Delta - (r^2 + a^2))\Sigma^{-1} \sin^2 \theta, \\ g_{\phi\phi} &= (a^2 \Delta \sin^2 \theta - (r^2 + a^2)^2)\Sigma^{-1} \sin^2 \theta, \end{aligned}$$

where  $\Sigma = r^2 + a^2 \cos^2 \theta$ ,  $\Delta = r^2 - 2Mr + a^2$ . Following Flaherty [8], one can find four complex scalar fields  $(z^0, z^1, \tilde{z}^0, \tilde{z}^1)$ , defined by

$$\begin{aligned} dz^0 &= dt - (a^2 + r^2)\Delta^{-1}dr - ia \sin \theta d\theta, \\ dz^1 &= d\phi - a\Delta^{-1}dr - i \csc \theta d\theta, \\ d\tilde{z}^0 &= dt + (a^2 + r^2)\Delta^{-1}dr + ia \sin \theta d\theta, \\ d\tilde{z}^1 &= d\phi + a\Delta^{-1}dr + i \csc \theta d\theta, \end{aligned}$$

such that the Kerr metric is

$$g = g_{tt} dz^0 d\tilde{z}^0 + g_{t\phi} (dz^0 d\tilde{z}^1 + dz^1 d\tilde{z}^0) + g_{\phi\phi} dz^1 d\tilde{z}^1.$$

In addition, letting  $\Omega^{-2} = (r - ia \cos \theta)^2$ , there must exist a scalar  $K$  such that  $g_{tt} = \Omega^{-2} \partial_{z^0} \partial_{\tilde{z}^0} K$ ,  $g_{t\phi} = \Omega^{-2} \partial_{z^0} \partial_{\tilde{z}^1} K$ ,  $g_{\phi\phi} = \Omega^{-2} \partial_{z^1} \partial_{\tilde{z}^1} K$ . However, expressions for  $K$  do not seem to have been obtained in the literature.

The method developed in this paper computes the generating function/Kähler potential of Kerr to be

$$K = 4 \int r \Delta^{-1} dr + 4 \log \sin \theta.$$

As a function  $K = K(z, \tilde{z})$ , the potential fully generates the spacetime geometry. Moreover, using Kähler transformations, we shall show that the Kähler potentials for Kerr and Schwarzschild are simply related by a Newman-Janis shift. More generally, the geometries studied in this paper include the general Plebański-Demiański class [13]

as well as the Chen-Teo family [14] of gravitational instantons. Our method also allows us to prove that a general class of supergravity black holes [15], including the Kerr-Sen spacetime [16], has a Hermitian (not conformal Kähler) structure.

In addition, we shall show that the integrability of complex structures leads to the existence of special scalar and massless free fields associated to the geometry, that can be combined to give a unified geometric description of the ‘Weyl double copy’ [17–19]. Our results contain not only the type D and N double copies, but also provide new examples of this relation for both vacuum and non-vacuum geometries, including e.g. the general Einstein-Maxwell Plebański-Demiański class and the Fubini-Study and Chen-Teo instantons.

Importantly, we shall not impose any field equations: the conformal Kähler property of the geometries we study does not depend on a particular field theory. This means that the conformal factor does not, in principle, play a role in our construction, but we shall nevertheless include it since it arises naturally in GR, where the Einstein and Kähler metrics are conformally related.

## COMPLEXIFIED KÄHLER GEOMETRY

Given a 4-dimensional complex geometry  $(M, g)$ , we define an almost-Hermitian structure [31] as a  $(1, 1)$  tensor field  $J$  such that  $J^2 = -\mathbb{I}$  and  $g(J\cdot, J\cdot) = g(\cdot, \cdot)$ . The tangent bundle decomposes as  $TM = T^+ \oplus T^-$ , where  $T^\pm$  corresponds to vectors with eigenvalue  $\pm i$  under  $J$ . We say that the almost-Hermitian structure is integrable, and is thus a Hermitian structure, if  $[T^\pm, T^\pm] \subset T^\pm$  (for both signs), where  $[\cdot, \cdot]$  is the Lie bracket of vector fields. One can show that a Hermitian structure implies that there are four complex scalars  $(z^i, \tilde{z}^i)$  such that

$$g = g_{i\tilde{j}} dz^i d\tilde{z}^{\tilde{j}}, \quad (1)$$

where  $g_{i\tilde{j}} = g(\partial_{z^i}, \partial_{\tilde{z}^{\tilde{j}}})$ , with  $i = 0, 1, \tilde{j} = \tilde{0}, \tilde{1}$ .

The fundamental 2-form is defined by  $\kappa(\cdot, \cdot) := g(J\cdot, \cdot)$ . We say that a Hermitian geometry is Kähler if  $d\kappa = 0$ , and conformal Kähler if there is a scalar field  $\Omega^2$  such that  $d\hat{\kappa} = 0$ , where  $\hat{\kappa} = \Omega^2\kappa$ . By the complex version of the Poincaré Lemma, if  $d\hat{\kappa} = 0$  then there exists, locally, a complex scalar  $K$  such that

$$\hat{g}_{i\tilde{j}} = \partial_{z^i} \partial_{\tilde{z}^{\tilde{j}}} K \quad (2)$$

where  $\hat{g}_{i\tilde{j}} = \Omega^2 g_{i\tilde{j}}$ , cf. [8, Theorem IX.8]. We say that  $K$  is a Kähler potential. It is not unique: one has the freedom to perform *Kähler transformations*

$$K \rightarrow K + F(z^i) + \tilde{F}(\tilde{z}^{\tilde{i}}). \quad (3)$$

The Kähler potential can be found by integrating (2). Define  $p_j := \partial K / \partial \tilde{z}^{\tilde{j}}$ , then  $p_j = \int \hat{g}_{i\tilde{j}} dz^i$ . Integrating once again, the potential is  $K = \int p_i d\tilde{z}^i$ .

In this work we shall study geometries whose metric has the block-diagonal form

$$g = a_{ij} d\sigma^i d\sigma^j + b_{IJ} dx^I dx^J \quad (4)$$

for some coordinates  $\sigma^i = (\tau, \phi)$  and  $x^I = (x, y)$ , and known functions  $a_{ij}, b_{IJ}$ . Introduce an orthonormal coframe  $e^1 = c_i^1 d\sigma^i$ ,  $e^2 = c_I^2 dx^I$ ,  $e^3 = c_I^3 dx^I$ ,  $e^4 = c_i^4 d\sigma^i$ , for some functions  $c_i^1, c_I^2, c_I^3, c_i^4$ ; such that  $g = e^1 \otimes e^1 + \dots + e^4 \otimes e^4$ . Define now a null coframe by

$$\begin{aligned} \ell &= \frac{1}{\sqrt{2}}(e^1 + ie^2), & n &= \frac{1}{\sqrt{2}}(e^1 - ie^2), \\ m &= \frac{1}{\sqrt{2}}(e^3 + ie^4), & \tilde{m} &= \frac{1}{\sqrt{2}}(-e^3 + ie^4). \end{aligned} \quad (5)$$

The metric is  $g = 2(\ell \odot n - m \odot \tilde{m})$ . We shall consider almost-Hermitian structures whose fundamental 2-forms are  $\kappa_\pm = i(\ell \wedge n \pm m \wedge \tilde{m})$ . For concreteness, let us focus on  $\kappa_- \equiv \kappa$ .

Let  $V^i = (-\partial_\phi, \partial_\tau)$ . We define the 1-forms

$$\omega^i := \mu V^i \lrcorner (\ell \wedge m), \quad \tilde{\omega}^i := \tilde{\mu} V^i \lrcorner (n \wedge \tilde{m}) \quad (6)$$

where  $\mu^{-1} = (\ell \wedge m)(\partial_\tau, \partial_\phi)$  and  $\tilde{\mu}^{-1} = (n \wedge \tilde{m})(\partial_\tau, \partial_\phi)$ . A calculation shows that

$$\omega^i = d\sigma^i + E_I^i dx^I, \quad \tilde{\omega}^i = d\sigma^i - E_I^i dx^I, \quad (7)$$

for some functions  $E_I^i = \omega^i(\partial_I)$ , where  $\partial_I = \partial/\partial x^I$ . In addition, the metric and fundamental 2-form are

$$g = g_{i\tilde{j}} \omega^i \odot \tilde{\omega}^{\tilde{j}}, \quad \kappa = \frac{i}{2} g_{i\tilde{j}} \omega^i \wedge \tilde{\omega}^{\tilde{j}} \quad (8)$$

where  $g_{i\tilde{j}} = g(\partial_{\sigma^i}, \partial_{\sigma^{\tilde{j}}})$ . Note that this implies that  $g_{0\tilde{0}} = g_{\tau\tau}$ ,  $g_{0\tilde{1}} = g_{1\tilde{0}} = g_{\tau\phi}$ ,  $g_{1\tilde{1}} = g_{\phi\phi}$ .

The almost-Hermitian structure is integrable iff  $d\omega^i = 0 = d\tilde{\omega}^{\tilde{i}}$ : if this is satisfied, then there will be (locally)  $z^i, \tilde{z}^{\tilde{i}}$  such that  $\omega^i = dz^i$  and  $\tilde{\omega}^{\tilde{i}} = d\tilde{z}^{\tilde{i}}$ , and from the first equation in (8) we see that the metric (4) will have the Hermitian expression (1). Using (7), this integrability condition has a simple form:  $d\omega^i = 0 = d\tilde{\omega}^{\tilde{i}}$  iff

$$E_I^i = E_I^i(x^J) \quad \text{and} \quad \partial_I E_J^i = 0. \quad (9)$$

The second equation implies that locally, there are functions  $\psi^0(x^I), \psi^1(x^I)$  such that  $E_I^i = \partial_I \psi^i$ . The  $(z^i, \tilde{z}^{\tilde{i}})$  coordinates will then be given by

$$z^i = \sigma^i + \psi^i, \quad \tilde{z}^{\tilde{i}} = \sigma^{\tilde{i}} - \psi^{\tilde{i}}. \quad (10)$$

The associated vector fields are  $\partial_{z^i} = \frac{1}{2}(\partial_{\sigma^i} + \partial_{\psi^i})$ ,  $\partial_{\tilde{z}^{\tilde{i}}} = \frac{1}{2}(\partial_{\sigma^{\tilde{i}}} - \partial_{\psi^{\tilde{i}}})$ . In terms of  $x^I$ , we have  $\partial_{\psi^i} = E_I^i \partial_I$ , where  $E_I^i$  is the inverse of  $E_I^i$  (thought of as a  $2 \times 2$  matrix).

We shall now assume that  $\partial_{\sigma^1} = \partial_\tau$  and  $\partial_{\sigma^2} = \partial_\phi$  are Killing vectors. This includes all of the examples studied in this paper. Using  $\kappa = -ig_{i\tilde{j}} d\sigma^i \wedge d\psi^{\tilde{j}}$  and  $\hat{g}_{i\tilde{j}} := \Omega^2 g_{i\tilde{j}}$ , a short calculation shows that the conformal Kähler condition  $d(\Omega^2 \kappa) = 0$  is

$$\partial_{\psi^0} \hat{g}_{\tau\phi} - \partial_{\psi^1} \hat{g}_{\tau\tau} = 0 = \partial_{\psi^0} \hat{g}_{\phi\phi} - \partial_{\psi^1} \hat{g}_{\tau\phi}. \quad (11)$$

Assuming the above conditions, the formula  $K = \int p_i d\bar{z}^i$  for the Kähler potential can be rewritten as follows. From Cartan's formula  $\mathcal{L}_v \hat{\kappa} = d(v \lrcorner \hat{\kappa}) + v \lrcorner d\hat{\kappa}$ , we deduce that the Killing fields have Hamiltonians, i.e. functions  $H_0, H_1$  such that  $dH_i = \partial_{\sigma^i} \lrcorner \hat{\kappa} = -i\hat{g}_{i\bar{j}} d\psi^{\bar{j}}$ , where the second equality follows from the expression of  $\hat{\kappa}$  in terms of  $\sigma^i, \psi^i$ . Choosing  $K$  to be independent of  $\sigma^i$ , we get

$$K = -4i \int H_i d\psi^i. \quad (12)$$

The integration in  $\psi^i$  can be replaced by an integration in  $x^I$  by using  $d\psi^i = E_I^i dx^I$ .

To recover real metrics with different signatures, we impose reality conditions on the null coframe  $(\ell, n, m, \bar{m})$ . Euclidean signature  $(++++)$  corresponds to requiring  $n = \bar{\ell}$  and  $\bar{m} = -\bar{m}$ . The functions  $E_I^i$  in (7) are then purely imaginary, so  $\tilde{\omega}^i = \bar{\omega}^i$  and  $\tilde{z}^i = \bar{z}^i$ . Lorentzian signature  $(+---)$  corresponds to  $\ell, n$  real and  $\bar{m} = \bar{m}$ . The functions  $E_I^i$  in (7) are generally complex, so  $z^i$  and  $\bar{z}^i$  in (10) are not complex conjugates.

## BLACK HOLES AND INSTANTONS

*Diagonal metrics.* Consider the special case of (4) where  $g = g_{\tau\tau} d\tau^2 + g_{\phi\phi} d\phi^2 + g_{xx} dx^2 + g_{yy} dy^2$ . We choose the frame such that the functions  $E_I^i$  in (7) are  $E_x^\tau = i\sqrt{g_{xx}/g_{\tau\tau}}$ ,  $E_y^\tau = 0$ ,  $E_x^\phi = 0$ ,  $E_y^\phi = -i\sqrt{g_{yy}/g_{\phi\phi}}$ . The Hermitian condition is equivalent to  $\partial_y(g_{xx}/g_{\tau\tau}) = 0$ ,  $\partial_x(g_{yy}/g_{\phi\phi}) = 0$ , and the conformal Kähler condition is  $\partial_x(\Omega^2 g_{\tau\tau}) = 0$ ,  $\partial_y(\Omega^2 g_{\phi\phi}) = 0$ .

A simple example is an arbitrary static, spherically symmetric spacetime  $g = f(r)d\tau^2 - h(r)dr^2 - r^2(d\theta^2 + \sin^2\theta d\phi^2)$ . Using  $\Omega^2 = 1/r^2$ , and *regardless* of the form of  $f(r), h(r)$ , the geometry is conformal Kähler. This includes not only the well-known spherical black hole spacetimes but also solutions from the Einstein-scalar field system such as the Janis-Newman-Winicour wormhole [20]. In the special case  $h = f^{-1}$ , the Kähler potential is given by  $K = 4 \int [rf(r)]^{-1} dr + \log \sin \theta$ .

*The Plebański-Demiański class.* Consider the metric (4) with

$$\begin{aligned} g_{\tau\tau} &= [\Delta_r - a^2 \Delta_x]/(\Pi\Sigma), \\ g_{\tau\phi} &= a[(r^2 + a^2)\Delta_x - (1 - x^2)\Delta_r]/(\Pi\Sigma), \\ g_{\phi\phi} &= [a^2(1 - x^2)^2 \Delta_r - (r^2 + a^2)^2 \Delta_x]/(\Pi\Sigma), \\ g_{xx} &= -\Sigma/(\Pi\Delta_x), \quad g_{xy} = 0, \quad g_{yy} = -\Sigma/(\Pi\Delta_r) \end{aligned} \quad (13)$$

where  $y \equiv r$ ,  $\Sigma = r^2 + a^2 x^2$ ,  $\alpha$  and  $a$  are constants, and  $\Pi = \Pi(r, x)$ ,  $\Delta_x = \Delta_x(x)$ ,  $\Delta_r = \Delta_r(r)$  are arbitrary functions of their arguments. We find that, *regardless* of the specific form of  $\Pi, \Delta_r, \Delta_x$ , the geometry is conformal

Kähler, with complex coordinates

$$\begin{aligned} z^0 &= \tau - (r^* - ia x^*), & z^1 &= \phi - (ar^\# - ix^\#), \\ \bar{z}^0 &= \tau + (r^* - ia x^*), & \bar{z}^1 &= \phi + (ar^\# - ix^\#), \end{aligned} \quad (14)$$

where  $r^*, x^*, r^\#, x^\#$  are defined by

$$\begin{aligned} dr^* &= (r^2 + a^2)\Delta_r^{-1} dr, & dx^* &= (1 - x^2)\Delta_x^{-1} dx, \\ dr^\# &= \Delta_r^{-1} dr, & dx^\# &= \Delta_x^{-1} dx, \end{aligned} \quad (15)$$

and the conformal factor is

$$\Omega^2 = \Pi/(r - ia x)^2. \quad (16)$$

The Kähler form  $\hat{\kappa} = \Omega^2 \kappa$  is given by

$$\begin{aligned} \hat{\kappa} &= \frac{i}{(r - ia x)^2} \{ -d\phi \wedge [a(1 - x^2)dr - i(r^2 + a^2)dx] \\ &\quad + d\tau \wedge (dr - iadx) \}. \end{aligned} \quad (17)$$

Notice that this is independent of  $\Delta_r, \Delta_x$ . The Hamiltonians are  $H_0 = -i/(r - ia x)$  and  $H_1 = i(a + irx)/(r - ia x)$ , hence, using (12), we find that the Kähler potential is

$$K = 4 \left[ \int \frac{r}{\Delta_r} dr - \int \frac{x}{\Delta_x} dx \right]. \quad (18)$$

We stress that the existence of this potential is independent of the explicit form of the functions  $\Delta_r, \Delta_x$ .

The Plebański-Demiański family [13, 21] is (13) with  $\Pi = (1 - \alpha r x)^2$  and

$$\begin{aligned} \Delta_x &= 1 + \frac{2N}{a}x - x^2 + 2\alpha Mx^3 - \left(\frac{\lambda}{3}a^2 + \alpha^2(Q^2 + a^2)\right)x^4, \\ \Delta_r &= Q^2 + a^2 - 2Mr + r^2 - \frac{2\alpha N}{a}r^3 - \left(\alpha^2 + \frac{1}{3}\lambda\right)r^4, \end{aligned}$$

where  $Q^2 = q_e^2 + q_m^2$ , and  $\lambda, q_e, q_m$  correspond, respectively, to cosmological constant and electric and magnetic charges. The rest of the parameters can be related to mass, angular momentum, acceleration, and NUT charge, cf. [21] for details. This is the general type D solution (assuming non-null orbits of the isometry group) of the Einstein equations with an aligned electromagnetic field.

We note that, for the case  $Q = 0$ , the transformation  $(r, M) \leftrightarrow \pm(iax, iN)$  leaves the Kähler potential and the metric invariant, and the coordinates (14) change according to  $z^i \leftrightarrow \bar{z}^i$  for  $+$  and are invariant for  $-$ . A detailed analysis of this and other dualities will be given in a separate work [22].

*Newman-Janis shifts.* For the Schwarzschild and Kerr spacetimes (putting  $x = \cos \theta$ ), we find the Kähler potentials to be

$$K_{\text{schw}} = 4 \log((r - 2M) \sin \theta), \quad (19a)$$

$$\begin{aligned} K_{\text{kerr}} &= 4 \left[ \frac{1}{2} \log(r^2 - 2Mr + a^2) + \log \sin \theta \right. \\ &\quad \left. - \frac{M}{\sqrt{M^2 - a^2}} \tanh^{-1} \left( \frac{r - M}{\sqrt{M^2 - a^2}} \right) \right], \end{aligned} \quad (19b)$$

where we assume the non-extreme case  $M^2 > a^2$ . Using (14) and Kähler transformations (3), a calculation shows that (19a) and (19b) are equivalent to

$$K_{\text{schw}} = 4 \left[ -\frac{r}{2M} + \log \sin \theta \right], \quad (20a)$$

$$K_{\text{kerr}} = 4 \left[ -\frac{(r - ia \cos \theta)}{2M} + \log \sin \theta \right] \quad (20b)$$

where we assume  $M \neq 0$ . Thus, the Kähler potentials are related by a Newman-Janis shift  $r \rightarrow r - ia \cos \theta$  [5], although it is not at all obvious from (19).

For  $M = 0$ , which corresponds (locally) to flat space-time, we can see the Newman-Janis shift as follows. Consider complexified Minkowski space, in complexified spherical coordinates  $(r_c, \theta_c, \phi_c)$ . In terms of complexified inertial coordinates  $(t_c, x_c, y_c, z_c)$ , we have the usual relations  $x_c^2 + y_c^2 = r_c^2 \sin^2 \theta_c$ ,  $z_c = r_c \cos \theta_c$ . The Kähler potential can be shown to be  $K = 4 \log(r_c \sin \theta_c)$ . Consider first the real slice  $\mathbb{M}$  given by  $\{t_c = t, x_c = x, y_c = y, z_c = z\}$ , where  $t, x, y, z$  are real. Then  $(r_c, \theta_c, \phi_c)$  become ordinary real spherical coordinates, and the Kähler potential is

$$K|_{\mathbb{M}} = 4 \log(r \sin \theta). \quad (21)$$

Now consider a different real slice  $\mathbb{M}'$  given by a Newman-Janis shift [23]:  $\{t_c = t, x_c = x, y_c = y, z_c = z - ia\}$ , where  $a$  is a real constant. Choosing the complex radius to be  $r_c = r - ia \cos \theta$ , a calculation gives  $x^2 + y^2 = (r^2 + a^2) \sin^2 \theta$ , so

$$K|_{\mathbb{M}'} = 4 \left[ \frac{1}{2} \log(r^2 + a^2) + \log \sin \theta \right]. \quad (22)$$

Eqs. (21) and (22) correspond, respectively, to the  $M \rightarrow 0$  limits in (19a) and (19b).

*Supergravity black holes.* Consider the metric (4) with

$$\begin{aligned} g_{\tau\tau} &= (R - U)/W, & g_{\tau\phi} &= (RW_u + UW_r)/W, \\ g_{\phi\phi} &= (RW_u^2 - UW_r^2)/W, & & \\ g_{xx} &= -W/R, & g_{xy} &= 0, & g_{yy} &= -W/U, \end{aligned} \quad (23)$$

where  $x \equiv r$ ,  $y \equiv u$ ,  $(R, W_r)$  and  $(U, W_u)$  are arbitrary functions of  $r$  and  $u$  respectively, and  $W = a(W_r + W_u)$ , with real constant  $a$ . The metric (23) includes a general class of black hole solutions of supergravity [15], in particular the Kerr-Sen black hole [16].

Using the almost-Hermitian structure associated to the frame given in [15, Eq. (4.79)], our method shows that the geometry (23) is Hermitian, with complex coordinates (10), where  $\psi^0 = r^* + iu^*$ ,  $\psi^1 = r^\# - iu^\#$ , and  $dr^* = a(W_r/R)dr$ ,  $du^* = a(W_u/U)du$ ,  $dr^\# = (a/R)dr$ ,  $du^\# = (a/U)du$ . However, the conformal Kähler condition (11) does not hold for this Hermitian structure.

*Gravitational instantons.* We now specialize to Euclidean signature. Consider first the metric (4) with

$$\begin{aligned} g_{\tau\tau} &= \frac{a^2 x^2}{4(1+x^2)^2}, & g_{\phi\phi} &= g_{\tau\tau}(1+x^2 \sin^2 y), & g_{xy} &= 0, \\ g_{\tau\phi} &= g_{\tau\tau} \cos y, & g_{xx} &= \frac{4}{a^2 x^2} g_{\tau\tau}, & g_{yy} &= (1+x^2)g_{\tau\tau}, \end{aligned}$$

where  $a$  is an arbitrary constant. Using “ $(\mp)$ ” to denote quantities associated to  $\kappa_{\mp}$ , one can choose frames such that the functions in (7) are  $E_{(\mp)x}^\tau = 2i/(ax)$ ,  $E_{(\mp)y}^\tau = \pm i \cot y$ ,  $E_{(\mp)x}^\phi = 0$ ,  $E_{(\mp)y}^\phi = \mp i \csc y$ . Then a calculation shows that the geometry is conformal Kähler w.r.t. both sides, with  $\Omega_{\mp}^2 = [(1+x^2)/x^2]^{1 \mp 1/a}$ . For  $a = \pm 1$ , one side becomes Kähler and the metric is Einstein: this is the Fubini-Study metric in  $\mathbb{C}\mathbb{P}^2$ .

A new family of gravitational instantons was discovered by Chen and Teo [14]. This is a toric, Ricci-flat geometry of the form (4) that depends on seven parameters  $k, \nu, a_0 \dots a_4$ . The non-trivial metric components are  $g_{\tau\tau}, g_{\tau\phi}, g_{\phi\phi}, g_{xx}, g_{yy}$ , and depend on functions  $F, G, H, X, Y$  given explicitly in [14, Eq. (2.1)]. The family contains other known instantons such as Eguchi-Hanson and Euclidean Plebański-Demiański. It was recently shown [24] that the Chen-Teo geometry is one-sided type D, and thus (from Ricci-flatness) conformal Kähler, with  $\Omega^2 = (x-y)^2/(\nu x + y)^2$ . Our method computes the complex coordinates to be  $dz^0 = d\tau + d\psi^0$ ,  $dz^1 = d\phi + d\psi^1$  (cf. (10)), where  $d\psi^0 = E_x^\tau dx + E_y^\tau dy$ ,  $d\psi^1 = E_x^\phi dx + E_y^\phi dy$ , and

$$\begin{aligned} E_x^\tau &= i \frac{\sqrt{k}}{F} \left( \frac{Gx}{X} + \frac{Hy}{(x-y)} \right), & E_x^\phi &= \frac{-i\sqrt{k}x}{X}, \\ E_y^\tau &= -i \frac{\sqrt{k}}{F} \left( \frac{Gy}{Y} + \frac{Hx}{(x-y)} \right), & E_y^\phi &= \frac{i\sqrt{k}y}{Y}. \end{aligned} \quad (24)$$

The Hermitian condition (9) reduces to  $\partial_y E_x^\tau - \partial_x E_y^\tau = 0$ , which provides an interpretation for eq. (3.49) in [24]. The Hamiltonians are

$$H_0 = \frac{\sqrt{k}(x-y)}{(1+\nu)(\nu x + y)}, \quad H_1 = \frac{\sqrt{k}f(x,y)}{(\nu x + y)(x-y)},$$

where  $f(x,y) = (\nu-1)(a_0 + a_4 x^2 y^2) - a_2 x(\nu(x-2y) + y) + (a_1 + a_3 xy)(\nu x - y)$ . The Kähler potential can now be computed using (12):

$$K = \frac{4k}{1+\nu} \left[ 4(1-\nu) \log(x-y) - \int \frac{h_1}{X} dx + \int \frac{h_2}{Y} dy \right]$$

where  $h_1(x) = a_1 + a_2(1-2\nu)x + a_3(2-\nu)x^2 + 2a_4(1-\nu)x^3$  and  $h_2(y) = a_1\nu - a_2y - a_3(1-2\nu)y^2 - 2a_4(1-\nu)y^3$ .

## DOUBLE COPY STRUCTURES

In string theory, the KLT relations [25] imply that gravitational amplitudes are closely related to the square of Yang-Mills amplitudes. The extension of these relations to field theory is known as the ‘double copy’. At the classical level, a recent formulation is the ‘curved Weyl double copy’ [18], which asserts that for some vacuum gravity solutions, the Weyl curvature spinor is  $\Psi_{ABCD} = \frac{1}{S} \Phi_{(AB} \Phi_{CD)}$  for some scalar field  $S$  (“zeroth copy”) and symmetric spinor field  $\Phi_{AB}$  (“single



copy”), where  $S$  satisfies a wave equation and  $\Phi_{AB}$  satisfies Maxwell’s equations. (We refer to [26, 27] for background on the 2-spinor formalism.) The relation has been proven for vacuum type D and type N spacetimes [18, 19]. We shall now show that the integrability conditions of complex structures give automatically this sort of relations among scalar, Maxwell, and gravitational fields.

Consider a conformal Kähler geometry, with conformal factor  $\Omega$ , Kähler form  $\hat{\kappa}_{ab} = \varphi_{AB}\epsilon_{A'B'}$ , Weyl spinor  $\Psi_{ABCD}$  and Ricci spinor  $\Phi_{ABA'B'}$ . Then one can show the following identities:

$$(\square + 2\Psi_2 + R/6)\Omega = 0, \quad (25a)$$

$$d\hat{\kappa} = 0 = d^*\hat{\kappa}, \quad (25b)$$

$$\Psi_{ABCD} = \Psi_2\Omega^{-4}\varphi_{(AB}\varphi_{CD)}, \quad (25c)$$

$$\Phi_{ABA'B'} = \Phi_{11}|\Omega|^{-4}\varphi_{AB}\bar{\varphi}_{A'B'}. \quad (25d)$$

Eq. (25a) follows by first noticing that the Lee form  $f_a$  (defined by  $d\kappa = -2f \wedge \kappa$ ) is  $f_a = \partial_a \log \Omega$ , then taking a divergence and using the identity  $\nabla^a f_a + f^a f_a = -(2\Psi_2 + R/6)$  (which can be proved using the Newman-Penrose formalism). Eq. (25b) follows from the conformal Kähler condition. Finally, equations (25c)-(25d) follow from the integrability conditions of a conformal Kähler structure, where for (25d) we assume Lorentz signature (and that the Ricci tensor satisfies  $R(J\cdot, J\cdot) = R(\cdot, \cdot)$ ).

From (25) we see that any conformal Kähler geometry combines scalar  $\Omega$ , Maxwell  $\hat{\kappa}$ , and gravitational fields in a double copy-like structure, without assuming any field equations. For Einstein manifolds ( $\Phi_{ABA'B'} = 0$ ), Bianchi identities imply  $\Omega = \Psi_2^{1/3}$ , so we recover the type D double copy [18] (extended to non-trivial cosmological constant). More generally, all of the conformal Kähler examples of the previous sections have the structure (25), so they represent double copy relations. New examples include the Fubini-Study and Chen-Teo instantons, but also the whole (non-vacuum) Plebański-Demiański class.

Furthermore, in the Plebański-Demiański case, the fields  $\Omega$  and  $\hat{\kappa}$  solve flat spacetime equations. More precisely, we see from (16) that  $\Omega$  is independent of  $\{M, N, q_e, q_m, \lambda\}$ , and since the case in which these parameters vanish corresponds to Minkowski, we immediately get  $\eta^{ab}\partial_a\partial_b\Omega = 0$ . In addition, from (17) we see that the Kähler form  $\hat{\kappa}$  depends only on  $a$ , so it must solve Maxwell’s equations in Minkowski.

The type N double copy [19] is not included in the above construction, but it also arises from the integrability of complex structures. First, consider a Petrov type II spacetime whose repeated principal spinor  $o_A$  satisfies

$$o^A o^B \nabla_{AA'} o_B = 0. \quad (26)$$

Eq. (26) is the condition for “half-integrability” of a complex structure (see [28, Section 2.4]). One can show [28, Prop. 2.6] that there is a scalar  $\Omega$  such that the Lee form satisfies  $o^A f_{AA'} = o^A \nabla_{AA'} \log \Omega$ . Applying  $\iota^A \nabla_{AA'}$  to this

equation (where  $o_A \iota^A = 1$ ), after some computations we again find that  $\Omega$  satisfies (25a). Notice that  $\Omega$  is not unique: we have the freedom to add  $\Omega \rightarrow \Omega + \nu$ , where  $\nu$  is any function such that  $o^A \nabla_{AA'} \nu = 0$ .

In addition, from [27, Lemma (7.3.15)], eq. (26) implies that there are two complex scalars  $z^i = (z^0, z^1)$  such that  $dz^i = o_A Z_{A'}^i dx^{AA'}$ , for some spinors  $Z_{A'}^i$ . It follows that the 2-form  $dz^0 \wedge dz^1$  is anti-self-dual and closed, so it is a Maxwell field. Note that  $F(z^0, z^1) dz^0 \wedge dz^1$  is also a Maxwell field for any function  $F(z^0, z^1)$ .

Finally, the conditions on  $o_A$  imply that there is a scalar  $\lambda$  such that  $o^A \nabla_{AA'} (\lambda o_B) = 0$ . This leads to  $\nabla^{AA'} (\lambda \Omega o_A) = 0$ , which in turn implies that  $\varphi_{A_1 \dots A_n} = \Omega \lambda^{n-1} o_{A_1} \dots o_{A_n}$  is a massless free field:  $\nabla^{A_1 A_1} \varphi_{A_1 \dots A_n} = 0$ . For  $n = 2$  and  $n = 4$ , we get the spin 1 and 2 fields  $\varphi_{AB} = \varphi_2 o_A o_B$  and  $\psi_{ABCD} = \psi_4 o_A o_B o_C o_D$ , with  $\varphi_2 = \Omega \lambda^2$  and  $\psi_4 = \Omega \lambda^4$ . These are related by

$$\psi_4 = \frac{1}{\Omega} (\varphi_2)^2. \quad (27)$$

In the special case in which the spacetime is type N,  $\psi_{ABCD}$  can be chosen to be the Weyl curvature spinor, and (27) is the type N double copy relation [19, Eq. (6)]. The non-uniqueness noticed in [19] is due to the freedom to include the functions  $\nu(z^0, z^1)$ ,  $F(z^0, z^1)$  mentioned before.

## DISCUSSION

A general expression for the Kähler potential  $K$  for the class of conformal Kähler (Lorentzian or Euclidean) geometries of the form (4) with two Killing fields is given by eq. (12). This includes the Plebański-Demiański and Chen-Teo families. The potential  $K$  generates not only the metric, but also the Maxwell field  $\hat{\kappa}$ . Notice that this electromagnetic field is exactly the Coulomb field of the Schwarzschild solution, or the  $\sqrt{K}$  Kerr / magic field of the Kerr solution [11, 29].

In  $(z, \tilde{z})$  coordinates,  $K$  is not necessarily expressible in terms of elementary functions. For example, while a potential for Minkowski is

$$K(z, \tilde{z}) = 4 \log \left[ \frac{\tilde{z}^0 - z^0}{1 + e^{i(z^1 - \tilde{z}^1)}} \right],$$

in the Schwarzschild case (20a),  $r$  and  $(z, \tilde{z})$  are related by  $r + 2M \log(r - 2M) = (\tilde{z}^0 - z^0)/2$ , which can be solved in terms of the Lambert W function.

Nevertheless, since the Kähler potential contains (locally and up to Kähler transformations) all the information of the geometry, it represents a fully non-linear version of the Debye potentials of perturbation theory in GR. As such, it is of intrinsic interest for the investigation of nonperturbative results for gravitational wave physics, further supported by the intriguing manifestation of the

Newman-Janis shift in  $K$  found in this paper, and by the fact that, as we showed, Kähler and complex geometry in GR contain naturally the known instances of the Weyl double copy.

The general framework and results obtained in this paper motivate applications to a variety of exciting problems in different areas of interest. In mathematical GR, potential applications include the analysis of waves on black hole spacetimes, analytic compactifications, and possible generalizations of the Chen-Teo instanton. In gravitational wave science, it would be interesting to make explicit connections to modern techniques used in scattering amplitudes and quantum field theory [10, 30]. The relation between Kähler potentials and the Newman-Janis shift motivates further investigation into the geometric origin of this trick, together with connections with its interpretation as a generation of intrinsic spin [8, Chapter X], see also [11, 12]. A detailed description of dualities in the Plebański-Demiański family will appear elsewhere [22].

*Acknowledgements.* We are grateful to Lars Andersson for his interest in this work and for pointing out relations to the Lambert  $W$  function as well as analytic compactifications. BA acknowledges the support of the Alexander von Humboldt Foundation.

---

\* [steffen.aksteiner@aei.mpg.de](mailto:steffen.aksteiner@aei.mpg.de)

† [bernardo.araneda@aei.mpg.de](mailto:bernardo.araneda@aei.mpg.de)

- [1] R. Penrose, *Nonlinear Gravitons and Curved Twistor Theory*, *Gen. Rel. Grav.* **7**, 31 (1976).
- [2] E. T. Newman, *Heaven and Its Properties*, *Gen. Rel. Grav.* **7** (1976), 107-111
- [3] J. F. Plebanski, *Some solutions of complex Einstein equations*, *J. Math. Phys.* **16** (1975), 2395-2402
- [4] J. F. Plebanski and I. Robinson, *Left-Degenerate Vacuum Metrics*, *Phys. Rev. Lett.* **37**, 493 (1976).
- [5] E. T. Newman and A. I. Janis, *Note on the Kerr spinning particle metric*, *J. Math. Phys.* **6** (1965), 915-917
- [6] A. Trautman, *Analytic solutions of Lorentz-invariant linear equations*, *Proc. Roy. Soc. Lond. A* **270** (1962), 326-328
- [7] E. J. Flaherty Jr, *An integrable structure for type D spacetimes*, *Physics Letters A*, 46(6), 391-392 (1974)
- [8] E. J. Flaherty Jr., *Hermitian and Kählerian Geometry in Relativity*, Springer Lecture Notes in Physics, Vol. 46 (Springer-Verlag, New York, 1976)
- [9] E. J. Flaherty Jr., *Complex Variables in Relativity*, in *General Relativity and Gravitation*. Vol. 2. One hundred years after the birth of Albert Einstein. Edited by A. Held. New York, NY: Plenum Press, p.207, 1980
- [10] A. Buonanno, M. Khalil, D. O'Connell, R. Roiban, M. P. Solon and M. Zeng, *Snowmass White Paper: Gravitational Waves and Scattering Amplitudes*, [arXiv:2204.05194 [hep-th]].
- [11] N. Arkani-Hamed, Y. t. Huang and D. O'Connell, *Kerr black holes as elementary particles*, *JHEP* **01** (2020), 046 [arXiv:1906.10100 [hep-th]].
- [12] A. Guevara, B. Maybee, A. Ochirov, D. O'Connell and J. Vines, *A worldsheet for Kerr*, *JHEP* **03** (2021), 201 [arXiv:2012.11570 [hep-th]].
- [13] J. F. Plebański and M. Demiański, *Rotating, charged, and uniformly accelerating mass in general relativity*, *Annals of Physics*, 98(1), 98-127 (1976)
- [14] Y. Chen and E. Teo, *Five-parameter class of solutions to the vacuum Einstein equations*, *Phys. Rev. D* **91** (2015) no.12, 124005 [arXiv:1504.01235 [gr-qc]].
- [15] D. D. K. Chow and G. Compère, *Dyonic AdS black holes in maximal gauged supergravity*, *Phys. Rev. D* **89** (2014) no.6, 065003 [arXiv:1311.1204 [hep-th]].
- [16] A. Sen, *Rotating charged black hole solution in heterotic string theory*, *Phys. Rev. Lett.* **69** (1992), 1006-1009 [arXiv:hep-th/9204046 [hep-th]].
- [17] R. Monteiro, D. O'Connell and C. D. White, *Black holes and the double copy*, *JHEP* **12** (2014), 056 [arXiv:1410.0239 [hep-th]].
- [18] A. Luna, R. Monteiro, I. Nicholson and D. O'Connell, *Type D Spacetimes and the Weyl Double Copy*, *Class. Quant. Grav.* **36** (2019), 065003 [arXiv:1810.08183 [hep-th]].
- [19] H. Godazgar, M. Godazgar, R. Monteiro, D. Peinador Veiga and C. N. Pope, *Weyl Double Copy for Gravitational Waves*, *Phys. Rev. Lett.* **126** (2021) no.10, 101103 [arXiv:2010.02925 [hep-th]].
- [20] A. I. Janis, E. T. Newman and J. Winicour, *Reality of the Schwarzschild Singularity*, *Phys. Rev. Lett.* **20** (1968), 878-880
- [21] J. B. Griffiths and J. Podolsky, *A New look at the Plebanski-Demiański family of solutions*, *Int. J. Mod. Phys. D* **15** (2006), 335-370 [arXiv:gr-qc/0511091 [gr-qc]].
- [22] S. Aksteiner and B. Araneda, to appear
- [23] E. T. Newman, *On a classical, geometric origin of magnetic moments, spin angular momentum and the Dirac gyromagnetic ratio*, *Phys. Rev. D* **65** (2002), 104005 [arXiv:gr-qc/0201055 [gr-qc]].
- [24] S. Aksteiner and L. Andersson, *Gravitational Instantons and special geometry*, [arXiv:2112.11863 [gr-qc]].
- [25] H. Kawai, D. C. Lewellen and S. H. H. Tye, *A Relation Between Tree Amplitudes of Closed and Open Strings*, *Nucl. Phys. B* **269** (1986), 1-23
- [26] R. Penrose and W. Rindler, *Spinors and space-time: Volume 1, Two-spinor calculus and relativistic fields*, volume 1. Cambridge University Press, 1984.
- [27] R. Penrose and W. Rindler, *Spinors and space-time: Volume 2, Spinor and twistor methods in space-time geometry*, volume 2. Cambridge University Press, 1986.
- [28] B. Araneda, *Conformal geometry and half-integrable spacetimes*, [arXiv:2110.06167 [gr-qc]].
- [29] D. Lynden-Bell, *Electromagnetic magic: The Relativistically rotating disk*, *Phys. Rev. D* **70** (2004), 105017 [arXiv:gr-qc/0410109 [gr-qc]].
- [30] G. Travaglini, A. Brandhuber, P. Dorey, T. McLoughlin, S. Abreu, Z. Bern, N. E. J. Bjerrum-Bohr, J. Blümlein, R. Britto and J. J. M. Carrasco, *et al. The SAGEX Review on Scattering Amplitudes*, [arXiv:2203.13011 [hep-th]].
- [31] The standard definition requires  $J$  to be real-valued, but we allow  $J$  to be complex-valued since we also include metrics with Lorentzian signature. We shall continue to use terminology, such as Hermitian, Kaehler, etc., as in the Riemannian case.