

GALOIS REPRESENTATIONS, (φ, Γ) -MODULES AND PRISMATIC F-CRYSTALS

ZHIYOU WU

ABSTRACT. We prove that both local Galois representations and (φ, Γ) -modules can be recovered from prismatic F-crystals, from which we obtain a new proof of the equivalence of Galois representations and (φ, Γ) -modules.

1. INTRODUCTION

The theory of (φ, Γ) -modules was developed by Fontaine ([1]) to study local Galois representations. It plays an important role in the study of families of Galois representations and p -adic Langlands correspondence.

The main result of the theory is that (φ, Γ) -modules are equivalent to Galois representations. The rough idea is to encode the difficult deeply ramified part of Galois theory into complicated rings. In other words, representations of complicated groups with simple coefficients are traded with representations of simple groups but with complicated coefficients.

The key part of the theory is then the construction of these complicated coefficient rings. It is based on the theory of fields of norms, which is a machine to switch between characteristic 0 and characteristic p worlds. There is another, probably more well-known, theory that serves the same purpose, namely perfectoid fields. Indeed, theory of fields of norms can be viewed as a deperfection of perfectoid fields. The coefficient rings appearing in (φ, Γ) -modules are certain infinitesimal lifting of fields of norms along the p -direction, in technical terms, they are Cohen rings of fields of norms.

The insight of this work is to put these rings in a world in which they naturally live, namely the framework of prisms as developed by Bhatt and Scholze ([2]). More precisely, these mysterious rings have natural integral structures which can be viewed as prisms, and the rings themselves are viewed as a structure sheaf on the prismatic site. Then (φ, Γ) -modules are vector bundles (with extra structures) on the prismatic site. This perspective is useful since prismatic sites are very rich. In particular, we can encode infinitesimal lifting of perfectoid fields into the prismatic world as well, which links with Galois representations. We can deduce the classical equivalence of Galois representations and (φ, Γ) -modules from this perspective, namely they are both equivalent to a third, arguably more fundamental object, the prismatic F-crystals. In summary, we have

Theorem 1.1. *Let k be a perfect field of characteristic p , and K be a finite extension of $W(k)[\frac{1}{p}]$. Then (φ, Γ) -modules over \mathbf{A}_K are equivalent to prismatic F-crystals in $\mathcal{O}_{\Delta}[\frac{1}{T}]_p^\wedge$ -modules over $(\mathcal{O}_K)_{\Delta}$. Moreover, continuous finite free \mathbb{Z}_p -representations of the absolute Galois group G_K are also equivalent to prismatic F-crystals in $\mathcal{O}_{\Delta}[\frac{1}{T}]_p^\wedge$ -modules over $(\mathcal{O}_K)_{\Delta}$.*

Corollary 1.2. *The category of continuous finite free \mathbb{Z}_p -representations of the absolute Galois group G_K is equivalent to the category of (φ, Γ) -modules over \mathbf{A}_K .*

See the main text for explanations of the notation.

The equivalence of Galois representations and prismatic F-crystals in $\mathcal{O}_{\Delta}[\frac{1}{T}]_p^\wedge$ -modules is also contained in the work of Bhatt and Scholze [3]. The proof for (φ, Γ) -modules follows the same line as the proof for Galois representations.

Acknowledgments. I would like to thank Peter Scholze for helpful discussions and encouragement. I would like to thank Heng Du, Koji Shimizu, Yu Min and Liang Xiao for discussions on the initial draft. I would also like to thank the anonymous referee for many suggestions and corrections. I am grateful to Max Planck Institute for Mathematics in Bonn for its hospitality and financial support.

Convention. We follow the notation of [2]. Fixing a prime p , a δ -ring is a $\mathbb{Z}_{(p)}$ -algebra R equipped with a map $\delta : R \rightarrow R$ such that $\delta(0) = \delta(1) = 0$ satisfying

$$\begin{aligned}\delta(x + y) &= \delta(x) + \delta(y) + \frac{x^p + y^p - (x + y)^p}{p} \\ \delta(xy) &= x^p\delta(y) + y^p\delta(x) + p\delta(x)\delta(y)\end{aligned}$$

for any $x, y \in R$. We write

$$\phi(x) := x^p + p\delta(x)$$

which is a ring homomorphism lifting the Frobenius. An element x of a δ -ring R is called distinguished if $\delta(x)$ is a unit.

Following the notation of Scholze, for an integral perfectoid ring R , we denote by $R^b := \lim_{\phi} R/p$ the tilt of R . We can also identify R^b with $\lim_{x \rightarrow x^p} R$ as a multiplicative monoid, then there exists a natural monoid map $R^b \rightarrow R$ given by

$$x = (x_0, x_1, \dots) \rightarrow x^\sharp := x_0,$$

where $x_{i+1}^p = x_i$, so (x_0, x_1, \dots) represents an element of $\lim_{x \rightarrow x^p} R$. As a standard notation in p -adic Hodge theory, we denote

$$A_{\text{inf}}(R) := W(R^b).$$

There exists a canonical surjection of rings

$$\theta : W(R^b) \rightarrow R$$

characterized by $\theta([x]) = x^\sharp$. Moreover, we know that $\text{Ker}(\theta)$ is principal and is generated by any distinguished element in the kernel, see [2] lemma 3.8, lemma 2.33 and lemma 2.24 for example. In particular, if R contains a compatible system of p -power roots of unit ζ_{p^n} , then

$$\epsilon := (1, \zeta_p, \zeta_{p^2}, \dots) \in R^b$$

and $1 + [\epsilon]^{\frac{1}{p}} + \dots + [\epsilon]^{\frac{p-1}{p}}$ generates $\text{Ker}(\theta)$.

We will use the theory of diamonds as developed in [4] and [5] in a rudimentary way. Recall that a diamond is a sheaf on the pro-étale site of characteristic p perfectoid spaces which can be written

as the quotient of a representable sheaf by a pro-étale equivalence relation. There is a functor from analytic adic spaces over $\mathrm{Spa}(\mathbb{Z}_p)$ into diamonds, which sends

$$X \longrightarrow X^\diamond$$

where X^\diamond is the sheaf whose value on a characteristic p perfectoid space S is the set of untilts S^\sharp of S together with a map $S^\sharp \rightarrow X$ of adic spaces. When $X = \mathrm{Spa}(R, R^+)$ is affinoid, we sometimes denote

$$\mathrm{Spd}(R, R^+) := \mathrm{Spa}(R, R^+)^\diamond.$$

When $A = R \otimes_S R$ and M is an R -module, we will write

$$M \otimes_R A$$

when A is viewed as an R -algebra with respect to the first factor, i.e. the R -algebra structure is $R \rightarrow R \otimes_S R$ is $x \rightarrow x \otimes 1$. Similarly, we write

$$A \otimes_R M$$

when A is equipped with the R -algebra structure with respect to the second factor. The convention applies also to other situations when A has two structure maps, such as $A = R \hat{\otimes}_S R$.

2. PRISMS

We recall the basic theory of prisms as developed in [2], and introduce the primary examples that will be relevant to us.

2.1. Definitions and Examples.

Definition 2.1. *A prism is a pair (A, I) , where A is a δ -ring, and I is an ideal of A such that A is derived (p, I) -complete, I defines a Cartier divisor on $\mathrm{Spec}(A)$, and $p \in I + \phi(I)A$. The category of prisms has objects the prisms, and the arrows are δ -ring maps preserving the given ideals.*

A prism (A, I) is called bounded if $A/I[p^\infty] = A/I[p^n]$ for some n .

Definition 2.2. *Let X be a p -adic formal scheme, then the (absolute) prismatic site X_Δ of X has objects bounded prisms (A, I) together with a map of formal schemes $\mathrm{Spf}(A/I) \rightarrow X$. The arrows are morphisms of prisms preserving the structure map to X . An arrow $(A, I) \rightarrow (B, J)$ is a cover if B is (p, I) -completely flat over A .*

When $X = \mathrm{Spf}(R)$ is affine, we simplify the notation by writing $R_\Delta := X_\Delta$.

Example 2.3. *Let k be a perfect field of characteristic p , then*

$$(W(k)[[q-1]], ([p]_q))$$

is a prism in $W(k)_\Delta$, where $[p]_q := \frac{q^p-1}{q-1}$, and the δ -structure is given by the usual δ -structure on $W(k)$ and $\delta(q) = 0$. It is clearly $(p, q-1)$ -complete, which is equivalent to being $(p, [p]_q)$ -complete as $[p]_q \equiv p \pmod{q-1}$ and $[p]_q \equiv (q-1)^{p-1} \pmod{p}$. Moreover, from $[p]_q \equiv p \pmod{q-1}$ we have

$$p = [p]_q + (q-1)\alpha$$

for some $\alpha \in W(k)[[q-1]]$. Applying ϕ to both sides, we have

$$p = \phi([p]_q) + (q^p-1)\phi(\alpha) = \phi([p]_q) + [p]_q(q-1)\phi(\alpha),$$

proving $p \in ([p]_q, \phi([p]_q))$.

Example 2.4. Let \mathbb{C} be the completion of algebraic closure of $W(k)$, and $\mathcal{O}_{\mathbb{C}}$ its ring of integers. Then $(W(\mathcal{O}_{\mathbb{C}}^b), \text{Ker}(\theta))$ is a prism, where $\theta : W(\mathcal{O}_{\mathbb{C}}^b) \rightarrow \mathcal{O}_{\mathbb{C}}$ is the canonical map characterized by $\theta([\alpha]) = \alpha^\sharp$. We choose a compatible system $\{\zeta_{p^n}\}$ of p -power roots in \mathbb{C} , and let

$$\epsilon := (1, \zeta_p, \zeta_{p^2}, \dots) \in \mathcal{O}_{\mathbb{C}}^b,$$

then it is well-known that $1 + [\epsilon^{\frac{1}{p}}] + [\epsilon^{\frac{2}{p}}] + \dots + [\epsilon^{\frac{p-1}{p}}]$ generates $\text{Ker}(\theta)$, and we have a map of δ -rings

$$W(k)[[q-1]] \rightarrow W(\mathcal{O}_{\mathbb{C}}^b)$$

by sending q to $[\epsilon^{\frac{1}{p}}]$, then $[p]_q$ is mapped to $1 + [\epsilon^{\frac{1}{p}}] + [\epsilon^{\frac{2}{p}}] + \dots + [\epsilon^{\frac{p-1}{p}}]$, and the condition $p \in (\text{Ker}(\theta), \phi(\text{Ker}(\theta)))$ follows from the same condition for the prism $(W(k)[[q-1]], ([p]_q))$.

Moreover, $W(\mathcal{O}_{\mathbb{C}}^b)$ is obviously $(p, [\varpi])$ -complete, where $\varpi \in \mathcal{O}_{\mathbb{C}}^b$ is any nonzero topologically nilpotent element of $\mathcal{O}_{\mathbb{C}}^b$. This implies $(p, \text{Ker}(\theta))$ -completeness since $1 + [\epsilon^{\frac{1}{p}}] + [\epsilon^{\frac{2}{p}}] + \dots + [\epsilon^{\frac{p-1}{p}}] \bmod p$ is such a ϖ .

We made use of an embedding $W(k)[[q-1]] \rightarrow W(\mathcal{O}_{\mathbb{C}}^b)$ sending q to $[\epsilon^{\frac{1}{p}}]$ in the previous example. This is not standard, and from now on we view $W(k)[[q-1]]$ as embedded into $W(\mathcal{O}_{\mathbb{C}}^b)$ using the embedding

$$W(k)[[q-1]] \rightarrow W(\mathcal{O}_{\mathbb{C}}^b)$$

which sends q to $[\epsilon]$. It is the ϕ -twist of the previous embedding.

We now recall the theory of fields of norms, see [6] for details. Let K be a finite totally ramified extension of $W(k)[\frac{1}{p}]$ contained in a fixed completed algebraically closure \mathbb{C} of $W(k)[\frac{1}{p}]$, then we can associate the cyclotomic tower

$$K \subset K(\zeta_p) \subset K(\zeta_{p^2}) \subset \dots$$

the field of norms \mathbf{E}_K , whose ring of integers \mathbf{E}_K^+ can be characterized as a subring of $\mathcal{O}_{K(\zeta_{p^\infty})}^b$, namely

$$\mathbf{E}_K^+ = \{(\alpha_n)_n \in \lim \mathcal{O}_{K(\zeta_{p^n})}/p \mid \alpha_n \in \mathcal{O}_{K(\zeta_{p^n})}/p \text{ for } n \text{ sufficiently large}\} \subset \mathcal{O}_{K(\zeta_{p^\infty})}^b.$$

Then \mathbf{E}_K^+ is a complete discrete valuation ring of characteristic p , which by construction contains $\mathbf{E}_{W(k)[\frac{1}{p}]}$. Moreover, we know that \mathbf{E}_K is a finite separable extension of $\mathbf{E}_{W(k)[\frac{1}{p}]}$.

We can compute $\mathbf{E}_{W(k)[\frac{1}{p}]}^+$ explicitly as

$$\mathbf{E}_{W(k)[\frac{1}{p}]}^+ = k[[\epsilon - 1]] \subset \mathcal{O}_{W(k)[\frac{1}{p}](\zeta_{p^\infty})}^b \subset \mathcal{O}_{\mathbb{C}}^b.$$

We observe that

$$W(k)((q-1)) := W(k)[[q-1]]\left[\frac{1}{q-1}\right]_p^\wedge \subset W(\mathbb{C}^b)$$

is a Cohen ring of $\mathbf{E}_{W(k)[\frac{1}{p}]}$. By the henselian property of $W(k)((q-1))$, the extension

$$\mathbf{E}_{W(k)[\frac{1}{p}]} \subset \mathbf{E}_K \subset \mathbb{C}^b$$

lifts canonically to

$$W(k)((q-1)) \subset \mathbf{A}_K \subset W(\mathbb{C}^b)$$

for a Cohen ring \mathbf{A}_K of \mathbf{E}_K . Let

$$\mathbf{A}_K^+ := \mathbf{A}_K \cap W(\mathcal{O}_{\mathbb{C}}^b),$$

then as $W(k)((q-1)) \subset W(\mathbb{C}^b)$ is invariant under the lift of Frobenius on $W(\mathbb{C}^b)$, so is \mathbf{A}_K^+ by naturality of \mathbf{A}_K being extension of $W(k)((q-1))$ inside $W(\mathbb{C}^b)$. This equips \mathbf{A}_K^+ with a δ -ring structure such that the embedding $\mathbf{A}_K^+ \subset W(\mathcal{O}_{\mathbb{C}}^b)$ is a δ -ring map. The inclusion $W(k)[[q-1]] \subset \mathbf{A}_K^+$ enables us to view $[p]_q$ as elements of \mathbf{A}_K^+ . We have the following lemma.

Lemma 2.5. *For every $n \in \mathbb{N}$, $(\mathbf{A}_K^+, (\phi^n([p]_q)))$ is a prism.*

Proof. Being a subring of $W(\mathcal{O}_{\mathbb{C}}^b)$, \mathbf{A}_K^+ is an integral domain, so $\phi^n([p]_q)$ is a non-zero divisor. Since $(W(k)[[q-1]], ([p]_q))$ is a prism, we have $p = a[p]_q + b\phi([p]_q)$ for some $a, b \in W(k)[[q-1]]$. Applying ϕ^n to it, we have $p = \phi^n(a)\phi^n([p]_q) + \phi^n(b)\phi^{n+1}([p]_q)$, proving $p \in (\phi^n([p]_q), \phi(\phi^n([p]_q)))$. Lastly, \mathbf{A}_K^+ is p -complete by construction and $\mathbf{A}_K^+/p \cong \mathbf{E}_K^+$, which is a complete DVR. We have $\phi^n([p]_q) \equiv (q-1)^{p^n(p-1)} \pmod{p}$, which is a pseudouniformizer in $\mathbf{E}_{W(k)[\frac{1}{p}]}^+ \cong k[[q-1]]$, hence a pseudouniformizer in \mathbf{E}_K^+ . This proves that \mathbf{A}_K^+ is $(p, \phi^n([p]_q))$ -complete. \square

We record some simple algebraic properties of the prism $(\mathbf{A}_K^+, (\phi^n([p]_q)))$.

Lemma 2.6. *The canonical inclusions $\mathbf{A}_K^+ \rightarrow W(\mathcal{O}_{\mathbb{C}}^b)$ and $W(k)[[q-1]] \rightarrow \mathbf{A}_K^+$ are flat.*

Proof. \mathbf{A}_K^+ is noetherian as it is of finite type over $W(k)[[q-1]]$. By remark 4.31 of [7], it is enough to prove flatness of the maps after reduction mod p , which are $\mathbf{E}_K^+ \rightarrow \mathcal{O}_{\mathbb{C}}^b$ and $k[[q-1]] \rightarrow \mathbf{E}_K^+$. They are injective maps from DVRs to integral domains, hence flat. \square

Corollary 2.7. *$\mathbf{A}_K^+/\phi^n([p]_q)$ is p -torsionfree, so the prism $(\mathbf{A}_K^+, (\phi^n([p]_q)))$ is bounded.*

Proof. We have that

$$W(k)[[q-1]]/\phi^n([p]_q) \rightarrow \mathbf{A}_K^+/\phi^n([p]_q)$$

is flat since it is the base change of $W(k)[[q-1]] \rightarrow \mathbf{A}_K^+$. We observe that

$$W(k)[[q-1]]/\phi^n([p]_q) \cong W(k)[\zeta_{p^n}],$$

which is p -torsionfree. Then the flatness tells us that $\mathbf{A}_K^+/\phi^n([p]_q)$ is also p -torsionfree. \square

Lemma 2.8. *The map $\phi : \mathbf{A}_K^+ \rightarrow \mathbf{A}_K^+$ is flat.*

Proof. Again by remark 4.31 of [7], it is enough to show the flatness mod p , which is the Frobenius $\phi : \mathbf{E}_K^+ \rightarrow \mathbf{E}_K^+$. It is an injective map of DVRs, so flat. \square

2.2. Perfect prisms and perfectoid rings. There is an important class of prisms that has close connection with perfectoid rings.

Definition 2.9. *A prism (A, I) is called perfect if ϕ is an automorphism of A .*

We have a natural perfection functor for prisms.

Proposition 2.10. *([2] lemma 3.8) Let (A, I) be a prism, and*

$$A_{perf} := (\operatorname{colim}_{\phi} A)_{(p, I)}^{\wedge},$$

then (A_{perf}, IA_{perf}) is a perfect prism. Moreover, it is the universal perfect prism over (A, I) .

Perfect prisms are canonically equivalent to integral perfectoid rings as defined in [7]. We recall the definition of integral perfectoid rings first.

Definition 2.11. *A ring R is integral perfectoid if it is π -adically complete for some $\pi \in R$ such that π^p divides p , the Frobenius on R/p is surjective, and the canonical map $\theta : W(R^{\flat}) \rightarrow R$ has principal kernel.*

The desired equivalence with perfect prisms is the following theorem.

Theorem 2.12. *([2] theorem 3.9) The category of perfect prisms is equivalent to the category of integral perfectoid rings. The equivalence functors are $(A, I) \rightarrow A/I$, and $R \rightarrow (W(R^{\flat}), \operatorname{Ker}(\theta))$.*

There is another notion of perfectoid rings used in the theory of perfectoid spaces. We recall the definition and compare it with integral perfectoid rings. Recall that a complete Tate ring is a complete Huber ring that contains a topological nilpotent unit. In more concrete terms, it is a complete topological ring R which contains an open subring R^+ whose topology is π -adic for some element $\pi \in R^+$, and $R = R^+[\frac{1}{\pi}]$. For any Huber ring R , we denote by R° the subring of power bounded elements.

Definition 2.13. *A perfectoid Tate ring is a uniform complete Tate ring R , i.e. R° is bounded, such that there exists a topological nilpotent unit $\pi \in R^{\circ}$ such that π^p divides p , and Frobenius is surjective on R°/π^p .*

We have the following comparison between the two notions of perfectoid rings. Recall that a ring of integral elements of a Huber ring R is an open and integrally closed subring R^+ of R° .

Proposition 2.14. *([7] lemma 3.20) Let R be a complete Tate ring, and $R^+ \subset R$ be a ring of integral elements. Then R is a perfectoid Tate ring if and only if R^+ is bounded in R and integral perfectoid.*

The proposition characterizes integral perfectoid subrings of a perfectoid Tate ring. We can also build a perfectoid Tate ring from an integral perfectoid ring as in the following proposition.

Proposition 2.15. *Let R be an integral perfectoid ring and $\pi \in R$ be an element such that R is π -adically complete and π^p divides p , then $R/\operatorname{Ann}(\pi)$ is integral perfectoid, and $R[\frac{1}{\pi}]$ is a perfectoid Tate ring with ring of definition $R/\operatorname{Ann}(\pi)$.*

Similarly, for R integral perfectoid, then $R/\operatorname{Ann}(p)$ is integral perfectoid and $R[\frac{1}{p}]$ is a perfectoid Tate ring with ring of definition $R/\operatorname{Ann}(p)$.

Proof. By [7] lemma 3.9, there are units u, v of R such that both πu and pv has compatible systems of p -power roots in R . Then by [8] 16.3.69, $R/\text{Ann}_R(\pi u)$, resp. $R/\text{Ann}_R(pv)$, is integral perfectoid without π -, resp. p -, torsion (the definition of perfectoid in [8] is the same as being integral perfectoid, as [7] remark 3.8 shows). Now $R[\frac{1}{\pi}]$, resp. $R[\frac{1}{p}]$, is a perfectoid Tate ring with ring of definition $R/\text{Ann}_R(\pi u)$, resp. $R/\text{Ann}_R(pv)$, by [7] lemma 3.21. \square

Remark 2.16. *We have not excluded the zero ring in the proposition. For example, any perfect ring of characteristic p is integral perfectoid with $\pi = 0$, then the rings produced in the proposition are all zero. This tells us that in some sense the integral perfectoid rings are more general than being perfectoid Tate. For example, finite fields are integral perfectoid, but can not be nonzero (ring of integers of) perfectoid Tate in any way.*

We can compute the perfection of the prism $(\mathbf{A}_K^+, (\phi^n([p]_q)))$.

Lemma 2.17. *We have*

$$(\mathbf{A}_K^+)_{\text{perf}} \cong W(\mathcal{O}_{K(\zeta_{p^\infty})}^\flat)^\wedge.$$

Proof. Let $I = (\phi^n([p]_q))$. By [2] corollary 2.31, we have

$$(\mathbf{A}_K^+)_{\text{perf}} \cong W((\mathbf{A}_K^+)_{\text{perf}}/p) \cong W((\text{colim}_\phi \mathbf{A}_K^+)_I^\wedge/p) \cong W((\text{colim}_\phi \mathbf{A}_K^+/p)_I^\wedge) \cong W((\text{colim}_\phi \mathbf{E}_K^+)_I^\wedge)$$

where we use lemma 10.96.1 (1) of Stacks project, and commutation of colimit with tensoring with \mathbb{Z}/p , in the third equality. From the theory of fields of norms, we know that

$$(\text{colim}_\phi \mathbf{E}_K^+)_I^\wedge = \mathcal{O}_{K(\zeta_{p^\infty})}^\flat,$$

where the completion is with respect to the natural valuation. We know that $\phi^n([p]_q) \equiv (q-1)^{p^n(p-1)} \pmod{p}$ is a pseudouniformizer in $k[[q-1]] \subset \mathbf{E}_K^+$, hence the completion is the same as completion with respect to I . Then we have

$$(\mathbf{A}_K^+)_{\text{perf}} \cong W(\mathcal{O}_{K(\zeta_{p^\infty})}^\flat)$$

as desired. \square

Corollary 2.18. *The automorphism group of $(\mathbf{A}_K^+, (\phi^n([p]_q)))$ in the category $\text{Spf}(\mathcal{O}_K)_\Delta$ is $\Gamma = \text{Gal}(K(\zeta_{p^\infty})/K)$.*

Proof. Let γ be an automorphism of $(\mathbf{A}_K^+, (\phi^n([p]_q)))$, then γ by definition is a δ -ring morphism, and is continuous with respect to $(p, \phi^n([p]_q))$ -topology, hence γ extends to an automorphism of $(\mathbf{A}_K^+, (\phi^n([p]_q)))_{\text{perf}}$. By theorem 3.10 of [2], the automorphism group of $(\mathbf{A}_K^+, (\phi^n([p]_q)))_{\text{perf}}$ as an abstract prism is the same as the automorphism group of the corresponding integral perfectoid ring, which is $\mathcal{O}_{K(\zeta_{p^\infty})}^\flat$ by the proposition. The automorphism of $(\mathbf{A}_K^+, (\phi^n([p]_q)))_{\text{perf}}$ as objects of $\text{Spf}(\mathcal{O}_K)_\Delta$ is then the \mathcal{O}_K -algebra automorphism of $\mathcal{O}_{K(\zeta_{p^\infty})}^\flat$, so

$$\gamma \in \text{Aut}(\mathcal{O}_{K(\zeta_{p^\infty})}^\flat/\mathcal{O}_K) = \Gamma.$$

But we observe that Γ already acts on $(\mathbf{A}_K^+, (\phi^n([p]_q)))$. The action on \mathbf{A}_K^+ is clear from its construction, and we need to check that it preserves the ideal $(\phi^n([p]_q))$.

First,

$$\gamma(q) = q^\alpha = (1+q-1)^\alpha = \sum_{i=0}^{\infty} \binom{\alpha}{i} (q-1)^i$$

for some $\alpha \in \mathbb{Z}_p^\times$, where $\binom{\alpha}{i} := \frac{\alpha(\alpha-1)\cdots(\alpha-i+1)}{i!}$. This follows from the definition $q = [\epsilon]$, and α is the value at γ of the cyclotomic character. Then

$$\gamma(\phi^n([p]_q)) = \frac{q^{\alpha p^{n+1}} - 1}{q^{\alpha p^n} - 1} = \frac{q^{p^n} - 1}{q^{\alpha p^n} - 1} \times \frac{q^{\alpha p^{n+1}} - 1}{q^{p^{n+1}} - 1} \times \phi^n([p]_q)$$

and we claim that $\frac{q^{p^n} - 1}{q^{\alpha p^n} - 1} \times \frac{q^{\alpha p^{n+1}} - 1}{q^{p^{n+1}} - 1}$ is a unit, which is what we want to prove. This follows from the computation

$$\begin{aligned} \frac{q^{p^n} - 1}{q^{\alpha p^n} - 1} &= \left(\alpha + \sum_{i=1}^{\infty} \binom{\alpha}{i+1} (q^{p^n} - 1)^i \right)^{-1} = \alpha^{-1} + (q-1)(\cdots) \\ \frac{q^{\alpha p^{n+1}} - 1}{q^{p^{n+1}} - 1} &= \alpha + \sum_{i=1}^{\infty} \binom{\alpha}{i+1} (q^{p^{n+1}} - 1)^i = \alpha + (q-1)(\cdots) \end{aligned}$$

which are units in $W(k)[[q-1]]$ as $\alpha \in \mathbb{Z}_p^\times$.

Now the corollary follows from the fact that $\mathbf{A}_K^+ \rightarrow (\mathbf{A}_K^+)_{\text{perf}}$ is injective (ϕ is injective as \mathbf{A}_K^+ is a δ -subring of the perfect δ -ring $W(\mathcal{O}_{\mathbb{C}}^b)$). \square

Moreover, we have the following proposition.

Proposition 2.19. *There exists $n \in \mathbb{N}$ such that*

$$(\mathbf{A}_K^+, (\phi^n([p]_q))) \in (\mathcal{O}_K)_\Delta,$$

i.e. there exists a map $\mathcal{O}_K \rightarrow \mathbf{A}_K^+ / \phi^n([p]_q)$.

Remark 2.20. \mathbf{A}_K^+ depends only on $K(\zeta_{p^\infty})$. In other words, \mathbf{A}_K^+ cannot see the difference between $K(\zeta_{p^k})$ and K , and this is taken care by the choice of $\phi^n([p]_q)$.

Proof. Let $I = ([p]_q)$, we have by lemma 2.17

$$(\mathbf{A}_K^+)_{\text{perf}} / I \cong W(\mathcal{O}_{K(\zeta_{p^\infty})}^b) / I \cong \mathcal{O}_{K(\zeta_{p^\infty})}^\wedge,$$

where the last isomorphism follows from ϕ being an automorphism and $\text{Ker}(\theta) = (\phi^{-1}([p]_q))$, for $\theta : W(\mathcal{O}_{K(\zeta_{p^\infty})}^b) \rightarrow \mathcal{O}_{K(\zeta_{p^\infty})}^\wedge$. On the other hand, we also have

$$(\mathbf{A}_K^+)_{\text{perf}} / I \cong (\text{colim}_\phi \mathbf{A}_K^+)_{(p,I)}^\wedge / I \cong (\text{colim}_\phi \mathbf{A}_K^+)^\wedge_p / I \cong (\text{colim}_\phi \mathbf{A}_K^+ / \phi^i(I))^\wedge_p$$

by definition of $(\mathbf{A}_K^+)_{\text{perf}}$, hence

$$(\text{colim}_\phi \mathbf{A}_K^+ / \phi^i(I))^\wedge_p \cong \mathcal{O}_{K(\zeta_{p^\infty})}^\wedge,$$

and we claim that this implies that

$$\text{colim}_\phi \mathbf{A}_K^+ / \phi^i(I) \cong \mathcal{O}_{K(\zeta_{p^\infty})}. \quad (1)$$

Observe that $\mathbf{A}_K^+/\phi^i(I)$ is integral over $W(k)$, so $\operatorname{colim}_{\phi} \mathbf{A}_K^+/\phi^i(I)$ is also integral over $W(k)$. Since there is no non-trivial integral extension of $\mathcal{O}_{K(\zeta_{p^\infty})}$ in its completion $\mathcal{O}_{K(\zeta_{p^\infty})}^\wedge$, $\operatorname{colim}_{\phi} \mathbf{A}_K^+/\phi^i(I)$, being integral over a subring of $\mathcal{O}_{K(\zeta_{p^\infty})}$, has to be contained in $\mathcal{O}_{K(\zeta_{p^\infty})}$. We look at the short exact sequence of $(\operatorname{colim}_{\phi} \mathbf{A}_K^+/\phi^i(I))$ -modules

$$0 \longrightarrow \operatorname{colim}_{\phi} \mathbf{A}_K^+/\phi^i(I) \longrightarrow \mathcal{O}_{K(\zeta_{p^\infty})} \longrightarrow \mathcal{O}_{K(\zeta_{p^\infty})}/\operatorname{colim}_{\phi} \mathbf{A}_K^+/\phi^i(I) =: M \longrightarrow 0.$$

From

$$(\operatorname{colim}_{\phi} \mathbf{A}_K^+/\phi^i(I))_p^\wedge \cong \mathcal{O}_{K(\zeta_{p^\infty})}^\wedge$$

and Stacks project lemma 10.96.1, we have $M_p^\wedge = 0$. The identification after completion also implies that $\operatorname{colim}_{\phi} \mathbf{A}_K^+/\phi^i(I)$ and $\mathcal{O}_{K(\zeta_{p^\infty})}$ have the same fraction field, in other words, M is p -torsion. Further, we observe that $\operatorname{colim}_{\phi} \mathbf{A}_K^+/\phi^i(I)$ contains

$$\operatorname{colim}_{\phi} W(k)[[q-1]]/\phi^i([p]_q) \cong W(k)[\zeta_{p^\infty}]$$

over which $\mathcal{O}_{K(\zeta_{p^\infty})}$ is finite. This implies that M is a finitely generated $(\operatorname{colim}_{\phi} \mathbf{A}_K^+/\phi^i(I))$ -module, which, together with being p -torsion, tells us that M is killed by p^k for some k , so M is p -adically complete. Then we have $M = M_p^\wedge = 0$, proving the claim.

Next, we show that the transition map $\phi : \mathbf{A}_K^+/\phi^i(I) \rightarrow \mathbf{A}_K^+/\phi^{i+1}(I)$ is injective. It is flat as it is base change of $\phi : \mathbf{A}_K^+ \rightarrow \mathbf{A}_K^+$, which is flat by lemma 2.8. The claim follows once we have shown that $\mathbf{A}_K^+/\phi^i(I)$ is an integral domain (flat maps preserves regular elements, so kernel consists of zero divisors, which can only be 0 in an integral domain). Let $J = \phi^i([p]_q)\mathbf{A}_K^+$, we have by definition $\mathbf{A}_K^+ \subset W(\mathcal{O}_{\mathbb{C}}^\flat)$, and we know that $W(\mathcal{O}_{\mathbb{C}}^\flat)/JW(\mathcal{O}_{\mathbb{C}}^\flat) \cong \mathcal{O}_{\mathbb{C}}$ which is an integral domain. Thus $JW(\mathcal{O}_{\mathbb{C}}^\flat)$ is a prime ideal in $W(\mathcal{O}_{\mathbb{C}}^\flat)$, so $JW(\mathcal{O}_{\mathbb{C}}^\flat) \cap \mathbf{A}_K^+$ is a prime ideal in \mathbf{A}_K^+ , and it remains to prove $JW(\mathcal{O}_{\mathbb{C}}^\flat) \cap \mathbf{A}_K^+ = J$. Let $x \in JW(\mathcal{O}_{\mathbb{C}}^\flat) \cap \mathbf{A}_K^+$, so $x = \phi^i([p]_q)y$ for some $y \in W(\mathcal{O}_{\mathbb{C}}^\flat)$. Then $y = \frac{x}{\phi^i([p]_q)} \in \mathbf{A}_K \cap W(\mathcal{O}_{\mathbb{C}}^\flat)$, which is \mathbf{A}_K^+ by definition, proving $x \in J$. Note that $\phi^i([p]_q)$ is a unit in \mathbf{A}_K since $\phi^i([p]_q) \equiv (q-1)^{p^i(p-1)} \pmod{p}$ is a unit in $\mathbf{A}_K/p = \mathbf{E}_K$.

Now as $\mathcal{O}_K \subset \mathcal{O}_{K(\zeta_{p^\infty})}$ is finite over $W(k)$, (1) together with injectiveness of the transition maps imply that $\mathcal{O}_K \subset \mathbf{A}_K^+/\phi^n([p]_q)$ for some n . \square

Remark 2.21. *We have used freely in the above proof the fact that there is no algebraic extension of nonarchimedean fields in their completion. More precisely, let L be an algebraic extension of $W(k)[\frac{1}{p}]$, then there is no nontrivial algebraic extension in its completion L^\wedge . Equivalently, completion induces an equivalence between algebraic extensions M/L of L and algebraic extension M^\wedge/L^\wedge of L^\wedge . This follows immediately from the fact that $\mathbb{C}^{\operatorname{Gal}(\bar{L}/M)} = M^\wedge$, where \mathbb{C} is a completed algebraic closure of L .*

3. PRISMATIC F-CRYSTALS

We recall the definition of prismatic F-crystals and make explicit an example that is relevant for us. Recall that we have a natural structure sheaf of δ -rings \mathcal{O}_Δ on X_Δ , together with an ideal sheaf I .

Definition 3.1. *Let X be a p -adic formal scheme, and \mathcal{R} be $\mathcal{O}_\Delta[\frac{1}{I}]_p^\wedge$, the p -adic completion of the structure sheaf \mathcal{O}_Δ with (locally) a generator of I inverted. A prismatic F-crystal on X in \mathcal{R} -modules is a finite locally free \mathcal{R} -module \mathcal{M} over X_Δ such that*

$$\mathcal{M}(A, I) \otimes_{\mathcal{R}(A, I)} \mathcal{R}(B, IB) \cong \mathcal{M}(B, IB)$$

for any arrow $(A, I) \rightarrow (B, IB)$ in X_Δ , together with an isomorphism

$$F : \phi^* \mathcal{M}[\frac{1}{I}] \cong \mathcal{M}[\frac{1}{I}]$$

of \mathcal{R} -modules.

A concrete way to work with prismatic F-crystals in $\mathcal{O}_\Delta[\frac{1}{I}]_p^\wedge$ -modules is to choose a cover (A, I) of the final object $*$ of the topos X_Δ , suppose that $(A, I) \times_* (A, I)$ is representable, and

$$(B, J) := (A, I) \times_* (A, I),$$

then a prismatic F-crystal in $\mathcal{O}_\Delta[\frac{1}{I}]_p^\wedge$ -modules is a finite projective φ - $A[\frac{1}{I}]_p^\wedge$ -module M together with an isomorphism

$$\beta : M \otimes_{A[\frac{1}{I}]_p^\wedge} B[\frac{1}{J}]_p^\wedge \cong B[\frac{1}{J}]_p^\wedge \otimes_{A[\frac{1}{I}]_p^\wedge} M$$

of φ - $B[\frac{1}{J}]_p^\wedge$ -modules satisfying cocycle conditions. Indeed, it is obvious that we can obtain such an object from a prismatic F-crystal. Conversely, if we are given such data, we can build a prismatic F-crystal \mathcal{M} as follows. Given a prism $(C, K) \in X_\Delta$, since (A, I) covers the final object, there exists a cover (C', K') of (C, K) , which also lies over (A, I) . Then we can define

$$\mathcal{M}(C, K) := \text{Eq}(M \otimes_{A[\frac{1}{I}]_p^\wedge} C'[\frac{1}{K'}]_p^\wedge \rightrightarrows M \otimes_{A[\frac{1}{I}]_p^\wedge} C''[\frac{1}{K''}]_p^\wedge)$$

where $(C'', K'') := (C', K') \times_{(C, K)} (C', K')$, which lives over (B, J) , and the two arrows comes from the base change descent data $\beta \otimes_{B[\frac{1}{J}]_p^\wedge} C''[\frac{1}{K''}]_p^\wedge$. Now by the next proposition, the descent data is effective on finite projective modules, so $\mathcal{M}(C, K)$ is a finite projective module over $C[\frac{1}{K}]_p^\wedge$.

Proposition 3.2. *Let $(A, I) \rightarrow (B, IB)$ be a cover in the category of bounded prisms, and (B^\bullet, IB^\bullet) be the corresponding Čech nerve, then we have an equivalence of categories*

$$\text{Vect}(A[\frac{1}{I}]_p^\wedge) \xrightarrow{\sim} \lim_{\leftarrow} (\text{Vect}(B[\frac{1}{I}]_p^\wedge) \rightrightarrows \text{Vect}(B^2[\frac{1}{I}]_p^\wedge) \rightrightarrows \cdots),$$

where $\text{Vect}(R)$ denotes the category of finite projective modules over the ring R .

Proof. Since $A[\frac{1}{I}]_p^\wedge$ (resp. $B^\bullet[\frac{1}{I}]_p^\wedge$) is p -complete, by [9] tag 0D4B, $\text{Vect}(A[\frac{1}{I}]_p^\wedge)$ (resp. $\text{Vect}(B^\bullet[\frac{1}{I}]_p^\wedge)$) is equivalent to $\lim_n \text{Vect}(A[\frac{1}{I}]/p^n)$ (resp. $\lim_n \text{Vect}(B^\bullet[\frac{1}{I}]/p^n)$). Thus it suffices to prove the equivalence

$$\text{Vect}(A/p^n[\frac{1}{I}]) \xrightarrow{\sim} \lim_{\leftarrow} (\text{Vect}(B/p^n[\frac{1}{I}]) \rightrightarrows \text{Vect}(B^2/p^n[\frac{1}{I}]) \rightrightarrows \cdots). \quad (2)$$

We know from the proof of [2] corollary 3.12 that B^2 is the derived (p, I) -completion of $B \otimes_A^{\mathbb{L}} B$, which is proved in *loc. cit.* to be discrete and classically (p, I) -complete. Then lemma 3.3 shows that

$$B^2 = (B \otimes_A B)_{(p, I)}^{\wedge},$$

i.e. it is the classical (p, I) -completion of $B \otimes_A B$, and similarly for B^\bullet . Indeed, by the discreteness and classical (p, I) -completeness of $\widehat{B \otimes_A^{\mathbb{L}} B}$, the derived (p, I) -completion of $B \otimes_A^{\mathbb{L}} B$, we have

$$B^2 = H^0(\widehat{B \otimes_A^{\mathbb{L}} B}) = H^0(\widehat{B \otimes_A^{\mathbb{L}} B})_{(p, I)}^{\wedge} = H^0(B \otimes_A^{\mathbb{L}} B)_{(p, I)}^{\wedge} = (B \otimes_A B)_{(p, I)}^{\wedge},$$

where the third equality follows from lemma 3.3. Then we have

$$B^2/p^n = (B/p^n \otimes_{A/p^n} B/p^n)_I^{\wedge}$$

$$B^3/p^n = (B/p^n \otimes_{A/p^n} B/p^n \otimes_{A/p^n} B/p^n)_I^{\wedge}$$

and similarly for B^\bullet/p^n . Since $A \rightarrow B$ is assumed to be (p, I) -completely faithfully flat, $A/p^n \rightarrow B/p^n$ is I -completely faithfully flat, and (2) follows from [10] theorem 7.8. \square

Lemma 3.3. *Let R be a ring and I be a finitely generated ideal of R . Let $K \in D^{\leq 0}(R)$ be a complex of R -modules with zero cohomology in positive degrees, and \hat{K} its derived I -completion, then we have a canonical identification*

$$H^0(\hat{K})_I^{\wedge} \cong H^0(K)_I^{\wedge},$$

in other words, the classical I -completion of $H^0(\hat{K})$ is the same as the classical I -completion of $H^0(K)$.

Proof. For any R -module M , we denote \hat{M} its derived I -completion by viewing M as a complex concentrated in zero degree. Recall that there is a natural map $M \rightarrow H^0(\hat{M})$ which is initial among maps $M \rightarrow N$ with N a derived I -complete R -module, see [10] definition 2.27 for example. By the spectral sequence in [9] tag 0BKE, we have

$$H^0(\hat{K}) = H^0(\widehat{H^0(K)}),$$

so we see that

$$H^0(K) \rightarrow H^0(\hat{K})$$

is the initial map among $H^0(K) \rightarrow N$ with N a derived I -complete R -module. We now claim that the composite map

$$H^0(K) \rightarrow H^0(\hat{K}) \rightarrow H^0(\hat{K})_I^{\wedge}$$

is initial among $H^0(K) \rightarrow N'$ where N' is classical I -complete. This is exactly the universal property of the classical I -completion of $H^0(K)$, whence $H^0(\hat{K})_I^{\wedge} = H^0(K)_I^{\wedge}$.

We now prove the claim. Let $H^0(K) \rightarrow N'$ where N' be as given, since classical I -completeness implies derived I -completeness ([9] tag 091T), we have a unique factorization

$$H^0(K) \rightarrow H^0(\hat{K}) \rightarrow N',$$

which further factors as

$$H^0(K) \rightarrow H^0(\hat{K}) \rightarrow H^0(\hat{K})_I^{\wedge} \rightarrow N'$$

by the universal property of $H^0(\hat{K})_I^{\wedge}$. \square

Example 3.4. Let k be a perfect field of characteristic p , K be a finite totally ramified extension of $W(k)[\frac{1}{p}]$, and $X = \mathrm{Spf}(\mathcal{O}_K)$. Let $n \in \mathbb{N}$ be chosen as in proposition 2.19 such that

$$(\mathbf{A}_K^+, (\phi^n([p]_q))) \in X_{\Delta},$$

then it is a cover of the final object by the following lemma 3.5. Moreover, we have by lemma 3.6

$$(\mathbf{A}_K^+, (\phi^n([p]_q))) \times_* (\mathbf{A}_K^+, (\phi^n([p]_q))) = (C, J),$$

where C is $(p, \phi^n([p]_q) \otimes 1)$ -completion of $\mathbf{A}_K^+ \hat{\otimes}_{W(k)} \mathbf{A}_K^+ \{ \frac{\omega \otimes 1 - 1 \otimes \omega}{\phi^n([p]_q) \otimes 1}, \frac{1 \otimes \phi^n([p]_q)}{\phi^n([p]_q) \otimes 1} \}$, which is freely adjoining $\frac{\omega \otimes 1 - 1 \otimes \omega}{\phi^n([p]_q) \otimes 1}$ and $\frac{1 \otimes \phi^n([p]_q)}{\phi^n([p]_q) \otimes 1}$ to $\mathbf{A}_K^+ \hat{\otimes}_{W(k)} \mathbf{A}_K^+$ as δ -rings, and $J = (\phi^n([p]_q) \otimes 1)$.

Thus a prismatic F -crystal in $\mathcal{O}_{\Delta}[\frac{1}{I}]_p^\wedge$ -modules over X is a \mathbf{A}_K -module M together with an isomorphism

$$M \otimes_{\mathbf{A}_K} C[\frac{1}{J}]_p^\wedge \cong C[\frac{1}{J}]_p^\wedge \otimes_{\mathbf{A}_K} M$$

satisfying cocycle conditions. The ring C is difficult to understand explicitly, we will see below how we can bypass this difficulty by passing to perfection.

Lemma 3.5. Let K be a finite totally ramified extension of $W(k)[\frac{1}{p}]$, $X = \mathrm{Spf}(\mathcal{O}_K)$, and $n \in \mathbb{N}$ be chosen as in proposition 2.19 such that

$$(\mathbf{A}_K^+, (\phi^n([p]_q))) \in X_{\Delta},$$

then it covers the final object in X_{Δ} .

Proof. Let (A, I) be an object of X_{Δ} , then A/I is a \mathcal{O}_K -algebra. We have a quasisyntomic cover $\mathcal{O}_K[\zeta_{p^\infty}]_p^\wedge$ of \mathcal{O}_K , hence

$$A/I \hat{\otimes}_{\mathcal{O}_K} \mathcal{O}_K[\zeta_{p^\infty}]_p^\wedge$$

is a quasisyntomic cover of A/I . By [2] proposition 7.11, we can find a prism (C, J) that covers (A, I) such that there is a morphism

$$A/I \hat{\otimes}_{\mathcal{O}_K} \mathcal{O}_K[\zeta_{p^\infty}]_p^\wedge \rightarrow C/J.$$

Now as $\mathcal{O}_K[\zeta_{p^\infty}]_p^\wedge$ is integral perfectoid, the composition

$$\mathcal{O}_K[\zeta_{p^\infty}]_p^\wedge \rightarrow A/I \hat{\otimes}_{\mathcal{O}_K} \mathcal{O}_K[\zeta_{p^\infty}]_p^\wedge \rightarrow C/J$$

lifts to a map of prisms

$$(A_{\mathrm{inf}}(\mathcal{O}_K[\zeta_{p^\infty}]_p^\wedge), \mathrm{Ker}(\theta)) \rightarrow (C, J)$$

by [2] lemma 4.7. We have that

$$(\mathbf{A}_K^+, (\phi^n([p]_q)))_{\mathrm{perf}} = (A_{\mathrm{inf}}(\mathcal{O}_K[\zeta_{p^\infty}]_p^\wedge), \phi^{n+1}(\mathrm{Ker}(\theta)))$$

by lemma 2.17, so we have a map

$$(\mathbf{A}_K^+, (\phi^n([p]_q))) \rightarrow (A_{\mathrm{inf}}(\mathcal{O}_K[\zeta_{p^\infty}]_p^\wedge), \phi^{n+1}(\mathrm{Ker}(\theta))) \xrightarrow{\phi^{-n-1}} (A_{\mathrm{inf}}(\mathcal{O}_K[\zeta_{p^\infty}]_p^\wedge), \mathrm{Ker}(\theta)) \rightarrow (C, J)$$

of prisms. As (C, J) covers (A, I) , we have finished the proof. \square

Lemma 3.6. *Let k be a perfect field of characteristic p , K be a finite totally ramified extension of $W(k)[\frac{1}{p}]$, and $\omega \in \mathbf{A}_K^+$ be an element which modulo $\phi^n([p]_q)$ becomes a uniformizer of \mathcal{O}_K under the inclusion $\mathcal{O}_K \subset \mathbf{A}_K^+/\phi^n([p]_q)$. Then as objects of $\mathrm{Spf}(\mathcal{O}_K)_\Delta$, we have*

$$(\mathbf{A}_K^+, (\phi^n([p]_q))) \times_* (\mathbf{A}_K^+, (\phi^n([p]_q))) = (C, J),$$

where

$$(C, J) := (\mathbf{A}_K^+ \hat{\otimes}_{W(k)} \mathbf{A}_K^+ \left\{ \frac{\omega \otimes 1 - 1 \otimes \omega}{\phi^n([p]_q) \otimes 1}, \frac{1 \otimes \phi^n([p]_q)}{\phi^n([p]_q) \otimes 1} \right\}^\wedge, (\phi^n([p]_q) \otimes 1)).$$

The ring displayed is $(p, \phi^n([p]_q) \otimes 1)$ -completion of $\mathbf{A}_K^+ \hat{\otimes}_{W(k)} \mathbf{A}_K^+ \left\{ \frac{\omega \otimes 1 - 1 \otimes \omega}{\phi^n([p]_q) \otimes 1}, \frac{1 \otimes \phi^n([p]_q)}{\phi^n([p]_q) \otimes 1} \right\}$, which is freely adjoining $\frac{\omega \otimes 1 - 1 \otimes \omega}{\phi^n([p]_q) \otimes 1}$ and $\frac{1 \otimes \phi^n([p]_q)}{\phi^n([p]_q) \otimes 1}$ to $\mathbf{A}_K^+ \hat{\otimes}_{W(k)} \mathbf{A}_K^+$ as δ -rings.

Proof. By definition, an object of X_Δ is a prism (A, I) equipped with a morphism $\mathcal{O}_K \rightarrow A/I$. The \mathcal{O}_K -algebra structure does not necessarily lift to A , but the corresponding $W(k)$ -algebra structure does. Indeed, by deformation theory of perfect rings, $W(k) \subset \mathcal{O}_K \rightarrow A/I$ lifts canonically to a morphism $W(k) \rightarrow A$, hence objects of X_Δ are naturally equipped with $W(k)$ -algebra structures, and all arrows in X_Δ are $W(k)$ -algebra morphisms.

Now given such an (A, I) , together with two arrows $(\mathbf{A}_K^+, (\phi^n([p]_q))) \rightarrow (A, I)$ in X_Δ , they give rise canonically to a δ -ring morphism

$$f : \mathbf{A}_K^+ \hat{\otimes}_{W(k)} \mathbf{A}_K^+ \longrightarrow A.$$

We know from properties of prisms that $I = (f(\phi^n([p]_q) \otimes 1)) = (f(1 \otimes \phi^n([p]_q)))$. Moreover, being arrows in X_Δ , the two arrows are \mathcal{O}_K -algebra morphisms mod $\phi^n([p]_q)$, which means that f maps $1 \otimes \omega - \omega \otimes 1 \bmod \phi^n([p]_q) \otimes 1$ to 0 in A/I . Thus f factors through $\mathbf{A}_K^+ \hat{\otimes}_{W(k)} \mathbf{A}_K^+ \left\{ \frac{\omega \otimes 1 - 1 \otimes \omega}{\phi^n([p]_q) \otimes 1}, \frac{1 \otimes \phi^n([p]_q)}{\phi^n([p]_q) \otimes 1} \right\}^\wedge$.

It remains to show that (C, J) is a prism. First we observe that $\frac{1 \otimes \phi^n([p]_q)}{\phi^n([p]_q) \otimes 1}$ is a unit in C , whence $(1 \otimes \phi^n([p]_q)) = (\phi^n([p]_q) \otimes 1)$. This follows from the equation

$$1 \otimes \phi^n([p]_q) = \phi^n([p]_q) \otimes 1 \cdot \frac{1 \otimes \phi^n([p]_q)}{\phi^n([p]_q) \otimes 1}$$

and [2] lemma 2.24.

We now claim that $\{1 \otimes \phi^n([p]_q), \omega \otimes 1 - 1 \otimes \omega\}$ is a regular sequence of $\mathbf{A}_K^+ \hat{\otimes}_{W(k)} \mathbf{A}_K^+$. Viewing $\mathbf{A}_K^+ \hat{\otimes}_{W(k)} \mathbf{A}_K^+$ as a \mathbf{A}_K^+ -algebra along the first factor, then it follows from [2] proposition 3.13 that (C, J) is a prism. It remains to prove the claim, which puts us into the setting of [2] proposition 3.13.

Observe that $1 \otimes \phi^n([p]_q)$ is regular since $\phi^n([p]_q)$ is regular in \mathbf{A}_K^+ (being an integral domain), and $\mathbf{A}_K^+ \hat{\otimes}_{W(k)} \mathbf{A}_K^+$ is flat over \mathbf{A}_K^+ . We want to show that $\omega \otimes 1 - 1 \otimes \omega$ is regular in

$$\mathbf{A}_K^+ \hat{\otimes}_{W(k)} \mathbf{A}_K^+ / 1 \otimes \phi^n([p]_q) \cong \mathbf{A}_K^+ \hat{\otimes}_{W(k)} (\mathbf{A}_K^+ / \phi^n([p]_q)).$$

We note that

$$\omega \otimes 1 - 1 \otimes \omega \in \mathbf{A}_K^+ \hat{\otimes}_{W(k)} \mathcal{O}_K \subset \mathbf{A}_K^+ \hat{\otimes}_{W(k)} (\mathbf{A}_K^+ / \phi^n([p]_q))$$

by definition of ω . By our assumption on k , \mathcal{O}_K is totally ramified over $W(k)$, so $\mathcal{O}_K \cong W(k)[x]/E$ for an Eisenstein polynomial $E \in W(k)[x]$. Then

$$\mathbf{A}_K^+ \hat{\otimes}_{W(k)} \mathcal{O}_K \cong \mathbf{A}_K^+[x]/E,$$

and by Eisenstein criterion, E is irreducible in $\mathbf{A}_K^+[x]$. Note that \mathbf{A}_K^+ is a regular local ring, whence a UFD by Auslander–Buchsbaum theorem. Then Gauss’s lemma tells us $\mathbf{A}_K^+[x]$ is a UFD as well, so irreducible polynomials are prime. Then $\mathbf{A}_K^+[x]/E$ is an integral domain, and $\omega \otimes 1 - 1 \otimes \omega$ is regular in $\mathbf{A}_K^+ \hat{\otimes}_{W(k)} \mathcal{O}_K$. We have seen in the proof of proposition 2.19 that $\mathbf{A}_K^+/\phi^n([p]_q)$ is an integral domain, so $\mathcal{O}_K \subset \mathbf{A}_K^+/\phi^n([p]_q)$ is flat, from which it follows that $\omega \otimes 1 - 1 \otimes \omega$ is regular in $\mathbf{A}_K^+ \hat{\otimes}_{W(k)} (\mathbf{A}_K^+/\phi^n([p]_q))$. \square

4. ÉTALE φ -MODULES

In this section, we prove that étale φ -modules on prisms does not change by passing to perfections. We first recall the definition of étale φ -modules.

Definition 4.1. *Let R be a ring equipped with a ring morphism $\varphi : R \rightarrow R$, an étale φ -modules over R is a finite projective R -module M equipped with an R -module isomorphism*

$$F : \varphi^* M = M \otimes_{R, \varphi} R \xrightarrow{\sim} M.$$

The morphisms between étale φ -modules are R -module morphisms preserving F . We denote by $\mathbf{ÉM}/_R$ the category of étale φ -modules over R .

First, we observe that by passing to naive perfections, the category does not change.

Proposition 4.2. *There is an equivalence of categories*

$$\mathbf{ÉM}/_R \xrightarrow{\sim} \mathbf{ÉM}/_{\underset{\varphi}{\text{colim}} R}$$

induced by base changing to $\underset{\varphi}{\text{colim}} R$.

Proof. For notational convenience, we index the rings R in the relevant system by R_n , i.e. the ring $\underset{\varphi}{\text{colim}} R$ is the colimit of the cofiltered system

$$R_0 \xrightarrow{\varphi} R_1 \xrightarrow{\varphi} R_2 \xrightarrow{\varphi} \dots$$

with each $R_i = R$. As the data of an étale φ -module is finite in nature, an étale φ -module over $\underset{\varphi}{\text{colim}} R$ comes via base change from an étale φ -module over R_n for some n . We need to show that it has further descent to R_0 . Now let M be an étale φ -module over R_n . Since $R_n = R = R_0$, we can view M as an étale φ -module over R_0 , and we claim that $M \otimes_{R_0} R_n = (\varphi^n)^* M$ is isomorphic to M as étale φ -modules over R_n . Iterate the φ -module structure $F : \varphi^* M \cong M$, we obtain an R -module isomorphism

$$G := F \circ \varphi^* F \circ \dots \circ (\varphi^{n-1})^* F : (\varphi^n)^* M \xrightarrow{\sim} M$$

we need to check that this is an étale φ -module isomorphism, i.e. $G \circ (\varphi^n)^* F = F \circ \varphi^* G$, but this is clear. This proves essential surjectivity.

For fully faithfulness, we need to show that for $(M, F_M), (N, F_N) \in \hat{\mathbf{E}}\mathbf{M}/_{R_0}$, an arrow $M \otimes_{R_0} \text{colim}_{\varphi} R \rightarrow N \otimes_{R_0} \text{colim}_{\varphi} R$ in $\hat{\mathbf{E}}\mathbf{M}/_{\text{colim}_{\varphi} R}$ comes uniquely from an arrow in $\hat{\mathbf{E}}\mathbf{M}/_{R_0}$ via base change. The arrow comes from an arrow in $\hat{\mathbf{E}}\mathbf{M}/_{R_n}$ for some n by standard finiteness argument, and we are reduced to showing that any arrow

$$f : M \otimes_{R_0} R_n \longrightarrow N \otimes_{R_0} R_n$$

in $\hat{\mathbf{E}}\mathbf{M}/_{R_n}$ has a unique descent to R_0 . We show that it descends uniquely to R_{n-1} , which proves the claim by iteration. Now we can assume $n = 1$, the previous paragraph shows that

$$F_M : \varphi^* M \xrightarrow{\sim} M$$

$$F_N : \varphi^* N \xrightarrow{\sim} N$$

are isomorphisms as étale φ -modules over R (being denoted by G in the previous paragraph). Let

$$g := F_N \circ f \circ F_M^{-1} : M \longrightarrow N$$

then g is an arrow in $\hat{\mathbf{E}}\mathbf{M}/_{R_0}$, and we claim that $\varphi^* g = f$. Since f is an arrow in $\hat{\mathbf{E}}\mathbf{M}/_{R_1}$, we have

$$\varphi^* F_N \circ \varphi^* f = f \circ \varphi^* F_M$$

and then

$$\varphi^* g = \varphi^* F_N \circ \varphi^* f \circ \varphi^* F_M^{-1} = f \circ \varphi^* F_M \circ \varphi^* F_M^{-1} = f,$$

proving the existence of the descent. It is unique since any descent h of f satisfies the relation $\varphi^* h = f$ by definition, and h satisfies the relation $h \circ F_M = F_N \circ \varphi^* h$ as it is in $\hat{\mathbf{E}}\mathbf{M}/_{R_0}$, combining the two we have

$$h \circ F_M = F_N \circ \varphi^* h = F_N \circ f$$

proving

$$h = F_N \circ f \circ F_M^{-1} = g.$$

□

Next we show that p -adic completion of the perfection does not lose information of étale φ -modules over p -complete rings.

Lemma 4.3. *Let R be a p -adically complete ring equipped with a ring morphism $\varphi : R \rightarrow R$, then base change induces an equivalence of categories*

$$\hat{\mathbf{E}}\mathbf{M}/_R \xrightarrow{\sim} \hat{\mathbf{E}}\mathbf{M}/_{(\text{colim}_{\varphi} R)_p^\wedge}.$$

Proof. By p -completeness of $(\text{colim}_{\varphi} R)_p^\wedge$, we have

$$\begin{aligned} \hat{\mathbf{E}}\mathbf{M}/_{(\text{colim}_{\varphi} R)_p^\wedge} &= \lim_n \hat{\mathbf{E}}\mathbf{M}/_{(\text{colim}_{\varphi} R)_p^\wedge / p^n} = \lim_n \hat{\mathbf{E}}\mathbf{M}/_{(\text{colim}_{\varphi} R) / p^n} = \\ &= \lim_n \hat{\mathbf{E}}\mathbf{M}/_{\text{colim}_{\varphi} (R/p^n)} = \lim_n \hat{\mathbf{E}}\mathbf{M}/_{(R/p^n)} = \hat{\mathbf{E}}\mathbf{M}/_R \end{aligned}$$

where we use the commutativity of colimit with tensoring with \mathbb{Z}/p^n in the third equality, proposition 4.2 in the fourth equality and p -completeness in the last one. □

We now specialize to the case of prisms and closely related rings. Let (A, I) be a bounded prism, we want to study étale φ -modules over $A[\frac{1}{I}]_p^\wedge$. There are two natural ways to form a perfection of the ring. The first is take the perfection directly and then p -complete it, namely

$$(\operatorname{colim}_\phi A[\frac{1}{I}]_p^\wedge)_p^\wedge,$$

while the second is to take the perfection of the prism (A, I) first, then inverting I and p -complete, i.e.

$$((\operatorname{colim}_\phi A)_{(p,I)}^\wedge[\frac{1}{I}])_p^\wedge.$$

It is the second one that will ultimately help us, and we want to understand étale φ -modules over it. Note that we have already understand étale φ -modules over the first ring, namely lemma 4.3 tells us that étale φ -modules over $(\operatorname{colim}_\phi A[\frac{1}{I}]_p^\wedge)_p^\wedge$ is the same as étale φ -modules over $A[\frac{1}{I}]_p^\wedge$. Observe that we have a natural morphism

$$(\operatorname{colim}_\phi A[\frac{1}{I}]_p^\wedge)_p^\wedge \longrightarrow ((\operatorname{colim}_\phi A)_{(p,I)}^\wedge[\frac{1}{I}])_p^\wedge$$

and we have the following theorem characterizing étale φ -modules over $((\operatorname{colim}_\phi A)_{(p,I)}^\wedge[\frac{1}{I}])_p^\wedge$.

Theorem 4.4. *Let (A, I) be a bounded prism such that $\phi(I) \bmod p$ is generated by a non-zero divisor in A/p , then we have an equivalence of categories*

$$\mathbf{EM}_{/(\operatorname{colim}_\phi A[\frac{1}{I}]_p^\wedge)_p^\wedge} \xrightarrow{\sim} \mathbf{EM}_{/((\operatorname{colim}_\phi A)_{(p,I)}^\wedge[\frac{1}{I}])_p^\wedge}$$

induced by base change.

Proof. We compute $(\operatorname{colim}_\phi A[\frac{1}{I}]_p^\wedge)_p^\wedge$ first. Being a p -complete perfect δ -ring, we know from [2] corollary 2.31 that

$$(\operatorname{colim}_\phi A[\frac{1}{I}]_p^\wedge)_p^\wedge = W((\operatorname{colim}_\phi A[\frac{1}{I}]_p^\wedge)/p) =$$

$$W((\operatorname{colim}_\phi A[\frac{1}{I}]_p^\wedge)/p) = W(\operatorname{colim}_\phi (A[\frac{1}{I}]_p^\wedge/p)) = W(\operatorname{colim}_\phi (A/p[\frac{1}{I}]))$$

where we use again the commutation of colimit with tensoring with \mathbb{Z}/p . A is (p, I) -complete as (A, I) is a bounded prism, so A/p is I -adically complete. By [2] lemma 3.6, $\phi(I)A$ is principal, so $\phi(I) \equiv I^p \bmod p$ is principal which is generated by a non-zero divisor by assumption. It follows that $A/p[\frac{1}{I}]$ is a Tate ring with ring of definition A/p .

On the other hand,

$$((\operatorname{colim}_\phi A)_{(p,I)}^\wedge[\frac{1}{I}])_p^\wedge = W((\operatorname{colim}_\phi A)_{(p,I)}^\wedge/p[\frac{1}{I}]) = W((\operatorname{colim}_\phi A/p)^\wedge[\frac{1}{I}])$$

by [2] corollary 2.31 again.

We know that étale φ -modules over $W(R)$ are equivalent to lisse sheaves on R for a perfect ring R by [11] proposition 3.2.7, hence it is enough to compare the finite étale sites of $(\operatorname{colim}_{\phi} A/p)_{\Gamma}^{\wedge}[\frac{1}{l}]$ and $\operatorname{colim}_{\phi} (A/p[\frac{1}{l}])$.

We note that $(\operatorname{colim}_{\phi} A/p)_{\Gamma}^{\wedge}[\frac{1}{l}]$ is the completed perfection of the Tate ring $A/p[\frac{1}{l}]$, and by the following lemma the finite étale site of $(\operatorname{colim}_{\phi} A/p)_{\Gamma}^{\wedge}[\frac{1}{l}]$ is the same as that of $A/p[\frac{1}{l}]$. But perfection also does not change finite étale site (see [11] theorem 3.1.15(a)), hence finite étale site of $\operatorname{colim}_{\phi} (A/p[\frac{1}{l}])$ is also identified with $A/p[\frac{1}{l}]$. This proves that the finite étale sites of $(\operatorname{colim}_{\phi} A/p)_{\Gamma}^{\wedge}[\frac{1}{l}]$ and $\operatorname{colim}_{\phi} (A/p[\frac{1}{l}])$ are equivalent via base change (as all intermediate equivalences here are through base change). \square

Lemma 4.5. *Let R be a Banach ring of characteristic p (in the sense of [11] definition 2.2.1), then the finite étale site of $(\operatorname{colim}_{\phi} R)^{\wedge}$ is equivalent to that of R via base change.*

Proof. Let R^u be the uniformization of R as defined in [11] definition 2.8.13, then by [11] proposition 2.8.16, the finite étale site of R^u is equivalent to that of R under base change. Moreover, by [11] theorem 3.1.15 (b), the finite étale sites of $(\operatorname{colim}_{\phi} R^u)^{\wedge}$ and R^u are equivalent, so we have the comparison between finite étale sites of R and $(\operatorname{colim}_{\phi} R^u)^{\wedge}$. We claim that $(\operatorname{colim}_{\phi} R^u)^{\wedge} = (\operatorname{colim}_{\phi} R)^{\wedge}$. By [11] lemma 2.6.2, there exists an affonid system R_i (see [11] definition 2.6.1 for definition) such that $R = (\operatorname{colim} R_i)^{\wedge}$, then $R^u = (\operatorname{colim} R_i^{\operatorname{red}})^{\wedge}$ by [11] corollary 2.5.6. Let $R_{\operatorname{perf}} := \operatorname{colim}_{\phi} R$, then

$$(R_{\operatorname{perf}})^{\wedge} = ((\operatorname{colim} R_i)_{\operatorname{perf}})^{\wedge} = (\operatorname{colim} (R_i)_{\operatorname{perf}})^{\wedge} = (\operatorname{colim} (R_i^{\operatorname{red}})_{\operatorname{perf}})^{\wedge} = (R_{\operatorname{perf}}^u)^{\wedge}$$

where we use that $T_{\operatorname{perf}}^{\operatorname{red}} = T_{\operatorname{perf}}$ for any ring T of characteristic p , which can be checked directly. \square

Combining all the equivalences we have established, we have the following theorem.

Theorem 4.6. *With assumptions as in theorem 4.4, we have an equivalence of categories*

$$\mathbf{EM}_{/A[\frac{1}{l}]_p^{\wedge}} \xrightarrow{\sim} \mathbf{EM}_{/((\operatorname{colim} A)_{(p,l)}^{\wedge}[\frac{1}{l}])_p^{\wedge}}$$

induced by base change.

5. (φ, Γ) -MODULES AND PRISMATIC F -CRYSTALS

In this section, we interpret (φ, Γ) -modules in terms of prismatic F -crystals. We recover the equivalence between Galois representations and (φ, Γ) -modules using the new interpretation.

Let us first recall the definition of (φ, Γ) -modules. Let K be a finite totally ramified extension of $W(k)[\frac{1}{p}]$, and \mathbf{A}_K^+ be as in section 2. Recall that $\mathbf{A}_K \subset W(\mathbb{C}^b)$ is stable by the canonical Frobenius lifting ϕ and the action of the Galois group $\operatorname{Gal}(\bar{K}/K)$ on $W(\mathcal{O}_{\mathbb{C}}^b)$. Moreover, the action factorizes through

$$\Gamma := \operatorname{Gal}(K(\zeta_{p^\infty})/K).$$

Definition 5.1. A (φ, Γ) -module over \mathbf{A}_K is an étale φ -module M over \mathbf{A}_K (with respect to the ϕ -structure on \mathbf{A}_K), i.e. a finite projective \mathbf{A}_K -module M equipped with an isomorphism

$$F : \phi^* M \cong M,$$

together with an action of Γ on M that commutes with F , and semilinear with respect to the action of Γ on \mathbf{A}_K .

We have the following theorem.

Theorem 5.2. The category of prismatic F -crystals in $\mathcal{O}_{\Delta}[\frac{1}{I}]_p^\wedge$ -modules over $\mathrm{Spf}(\mathcal{O}_K)$ is equivalent to the category of (φ, Γ) -modules over \mathbf{A}_K .

Proof. By example 3.4, we see that a prismatic F -crystal in $\mathcal{O}_{\Delta}[\frac{1}{I}]_p^\wedge$ -modules over $\mathrm{Spf}(\mathcal{O}_K)$ is an étale φ -module over \mathbf{A}_K together with an isomorphism

$$M \otimes_{\mathbf{A}_K} C[\frac{1}{J}]_p^\wedge \cong C[\frac{1}{J}]_p^\wedge \otimes_{\mathbf{A}_K} M$$

of étale φ -modules over $C[\frac{1}{J}]_p^\wedge$ satisfying cocycle conditions, with

$$(C, J) = (\mathbf{A}_K^+ \hat{\otimes}_{W(k)} \mathbf{A}_K^+ \left\{ \frac{\omega \otimes 1 - 1 \otimes \omega}{\phi^n([p]_q) \otimes 1}, \frac{1 \otimes \phi^n([p]_q)}{\phi^n([p]_q) \otimes 1} \right\}^\wedge, (\phi^n([p]_q) \otimes 1))$$

as in example 3.4. Then theorem 4.6 implies that this is equivalent (via base change) to an étale φ -module \mathcal{M} over

$$(\mathbf{A}_K^+)_{\mathrm{perf}}[\frac{1}{J}]_p^\wedge \cong A_{\mathrm{inf}}(\mathcal{O}_{K(\zeta_{p^\infty})_p^\wedge})[\frac{1}{J}]_p^\wedge \cong W((K(\zeta_{p^\infty})_p^\wedge)^b)$$

together with an isomorphism

$$\mathcal{M} \otimes_{W((K(\zeta_{p^\infty})_p^\wedge)^b)} B[\frac{1}{I}]_p^\wedge \cong B[\frac{1}{I}]_p^\wedge \otimes_{W((K(\zeta_{p^\infty})_p^\wedge)^b)} \mathcal{M}$$

as étale φ -modules over $B[\frac{1}{I}]_p^\wedge$, where $(B, I) = (C, J)_{\mathrm{perf}}$. Then the lemma below tells us that the descent data is equivalent to an action of Γ on \mathcal{M} that is semilinear with respect to the action of Γ on $W((K(\zeta_{p^\infty})_p^\wedge)^b)$. As the action of Γ is already defined on \mathbf{A}_K , theorem 4.6 tells us that this is equivalent to an action of Γ on M that is semilinear with respect to the action of Γ on \mathbf{A}_K , which is exactly a (φ, Γ) -module over \mathbf{A}_K . \square

Lemma 5.3. Let (B, I) be the perfection of the prism (C, J) in lemma 3.6, then

$$B[\frac{1}{I}]_p^\wedge \cong C^0(\Gamma, W((K(\zeta_{p^\infty})_p^\wedge)^b))$$

where $C^0(\Gamma, W((K(\zeta_{p^\infty})_p^\wedge)^b))$ is the ring of continuous functions on Γ with values in $W((K(\zeta_{p^\infty})_p^\wedge)^b)$. Moreover, the two structure maps from $W((K(\zeta_{p^\infty})_p^\wedge)^b)$ to $C^0(\Gamma, W((K(\zeta_{p^\infty})_p^\wedge)^b))$ are the obvious constant function map, and the one sending $x \in W((K(\zeta_{p^\infty})_p^\wedge)^b)$ to $\{\gamma \rightarrow \gamma(x)\}$.

Proof. Let

$$(F, F^+) := (K(\zeta_{p^\infty})_p^\wedge, \mathcal{O}_{K(\zeta_{p^\infty})_p^\wedge}).$$

By lemma 3.6, (B, I) is the initial perfect prism in $\mathrm{Spf}(\mathcal{O}_K)_\Delta$ equipped with two arrows from $(\mathbf{A}_K^+, (\phi^n([p]_q)))$ into it. This is the same as the initial perfect prism in $\mathrm{Spf}(\mathcal{O}_K)_\Delta$ with two arrows from

$$(\mathbf{A}_K^+, (\phi^n([p]_q)))_{\mathrm{perf}} = (A_{\mathrm{inf}}(F^+), \mathrm{Ker}(\theta)),$$

where we use lemma 2.17. Since perfect prisms are equivalent to integral perfectoid rings, we see that B/I is the initial integral perfectoid \mathcal{O}_K -algebra with two maps (as \mathcal{O}_K -algebras) from F^+ . We claim that $B/I[\frac{1}{p}]$ is the initial perfectoid Tate K -algebra with two maps from F . This follows immediately from proposition 2.15 and proposition 2.14.

Now for any perfectoid Tate K -algebra A with two maps from F , we obtain two maps from $\mathrm{Spa}(A, A^\circ)$ to $\mathrm{Spa}(F, F^+)$, as perfectoid spaces over $\mathrm{Spa}(K, \mathcal{O}_K)$. We view them as diamonds over $\mathrm{Spd}(K, \mathcal{O}_K)$. Since diamonds is determined by their values on (affinoid) perfectoid test objects, the previous paragraph shows that

$$\mathrm{Spa}(B/I[\frac{1}{p}], B/I[\frac{1}{p}]^\circ) = \mathrm{Spa}(F, F^+)^\circ \times_{\mathrm{Spd}(K, \mathcal{O}_K)} \mathrm{Spa}(F, F^+)^\circ$$

for some ring of definition $B/I[\frac{1}{p}]^\circ$, which can be identified as the image of B/I in $B/I[\frac{1}{p}]$, providing we know the right hand side is affinoid perfectoid. But it is well-known that

$$\mathrm{Spd}(K, \mathcal{O}_K) = \mathrm{Spa}(F, F^+)^\circ / \Gamma,$$

see [5] lemma 10.1.7 for example, so

$$\mathrm{Spa}(F, F^+)^\circ \times_{\mathrm{Spd}(K, \mathcal{O}_K)} \mathrm{Spa}(F, F^+)^\circ \cong \mathrm{Spa}(F, F^+)^\circ \times \underline{\Gamma} \cong \mathrm{Spa}(C^0(\Gamma, F), C^0(\Gamma, F^+)^\circ)$$

where $C^0(\Gamma, F)$, resp. $C^0(\Gamma, F^+)^\circ$, is the ring of continuous functions on Γ with values in F , resp. F^+ , see [4] example 11.12 for the last isomorphism. We then have

$$B/I[\frac{1}{p}] \cong C^0(\Gamma, F).$$

as the functor from perfectoid spaces over $\mathrm{Spa}(K, \mathcal{O}_K)$ to diamonds over $\mathrm{Spd}(K, \mathcal{O}_K)$ is fully faithful. The two structure maps from F to $C^0(\Gamma, F)$ is then the constant function map, and the one sending $x \in F$ to $\{\gamma \rightarrow \gamma(x)\}$, which is easy to see by chasing through the above canonical isomorphisms. Now

$$B[\frac{1}{I}]_p^\wedge = W((B/I)^\flat[\frac{1}{\omega}]) = W((B/I[\frac{1}{p}])^\flat)$$

for a uniformizer $\varpi = (a_0, a_1, \dots) \in (B/I)^\flat$ which we can choose so that a_0 divides p in B/I . Thus

$$B[\frac{1}{I}]_p^\wedge \cong W(C^0(\Gamma, F)^\flat) \cong W(C^0(\Gamma, F^\flat)) \cong C^0(\Gamma, W(F^\flat))$$

where the second isomorphism follows directly from the description $F^\flat = \lim_{x \rightarrow x^p} F$, and the same description for $C^0(\Gamma, F)^\flat$. The last isomorphism follows from the concrete description of the Witt vector, namely $W(F^\flat) = (F^\flat)^\mathbb{N}$. This clearly commutes with taking continuous functions. Moreover, the ring structure is defined by polynomial equations that is independent of the input ring, whence the last isomorphism. The description of the two structure maps is straightforward by chasing through the various canonical isomorphisms. \square

Remark 5.4. *The action of Γ can be directly detected as follows. For a prismatic F-crystal M over \mathcal{O}_K , and $\gamma \in \Gamma$. The action of γ on $M((\mathbf{A}_K^+, (\phi^n([p]_q))))$ is induced by the base change isomorphism (the crystal structure)*

$$M((\mathbf{A}_K^+, (\phi^n([p]_q)))) \otimes_{\mathbf{A}_K, \gamma} \mathbf{A}_K \cong M((\mathbf{A}_K^+, (\phi^n([p]_q))))$$

corresponding to the arrow

$$\gamma : (\mathbf{A}_K^+, (\phi^n([p]_q))) \xrightarrow{\sim} (\mathbf{A}_K^+, (\phi^n([p]_q)))$$

in $(\mathcal{O}_K)_\Delta$ as described in the proof of corollary 2.18.

Remark 5.5. *The proof also shows that prismatic F-crystals in $\mathcal{O}_\Delta[\frac{1}{I}]_p^\wedge$ -modules over $\mathrm{Spf}(\mathcal{O}_K)$ are equivalent to F-crystals in $\mathcal{O}_\Delta[\frac{1}{I}]_p^\wedge$ -modules over $\mathrm{Spf}(\mathcal{O}_K)_\Delta^{\mathrm{perf}}$, i.e. the site of perfect prisms over $\mathrm{Spf}(\mathcal{O}_K)$, and the equivalence is induced by the obvious restriction functor. Indeed, prismatic F-crystals in $\mathcal{O}_\Delta[\frac{1}{I}]_p^\wedge$ -modules over $\mathrm{Spf}(\mathcal{O}_K)$ (resp. $\mathrm{Spf}(\mathcal{O}_K)_\Delta^{\mathrm{perf}}$) are equivalent to φ -modules over $\mathbf{A}_K^+[\frac{1}{I}]_p^\wedge$ together with descent data over $C[\frac{1}{I}]_p^\wedge$ (resp. φ -modules over $(\mathbf{A}_K^+)_{\mathrm{perf}}[\frac{1}{I}]_p^\wedge$ together with descent data over $C_{\mathrm{perf}}[\frac{1}{I}]_p^\wedge$), and the proof shows that the latter objects are equivalent.*

With exactly the same idea, we can recover Galois representations from prismatic F-crystals in $\mathcal{O}_\Delta[\frac{1}{I}]_p^\wedge$ -modules. As it is along the same reasoning as above, we only sketch the argument.

Theorem 5.6. *The category of prismatic F-crystals in $\mathcal{O}_\Delta[\frac{1}{I}]_p^\wedge$ -modules over $\mathrm{Spf}(\mathcal{O}_K)$ is equivalent to the category of finite free continuous \mathbb{Z}_p -representations of $G := \mathrm{Gal}(\overline{K}/K)$.*

Proof. Let $\mathbb{C} := \widehat{\overline{K}}$ and $\mathcal{O}_\mathbb{C}$ be the ring of integers of it. By remark 5.5, it enough to work in $\mathrm{Spf}(\mathcal{O}_K)_\Delta^{\mathrm{perf}}$. We can evaluate a prismatic F-crystal at $(A_{\mathrm{inf}}(\mathcal{O}_\mathbb{C}), \mathrm{Ker}(\theta))$. Let (B, J) be the product

$$(A_{\mathrm{inf}}(\mathcal{O}_\mathbb{C}), \mathrm{Ker}(\theta)) \times (A_{\mathrm{inf}}(\mathcal{O}_\mathbb{C}), \mathrm{Ker}(\theta))$$

in $\mathrm{Spf}(\mathcal{O}_K)_\Delta^{\mathrm{perf}}$, and we need to compute

$$B[\frac{1}{J}]_p^\wedge = W((B/J[\frac{1}{p}])^b).$$

We know that $B/J[\frac{1}{p}]$ is a perfectoid Tate K -algebra and can be interpreted as

$$\mathrm{Spa}(B/J[\frac{1}{p}], B/J[\frac{1}{p}]^+)^\diamond = \mathrm{Spd}(\mathbb{C}, \mathcal{O}_\mathbb{C}) \times_{\mathrm{Spd}(K, \mathcal{O}_K)} \mathrm{Spd}(\mathbb{C}, \mathcal{O}_\mathbb{C}).$$

We know that

$$\mathrm{Spd}(\mathbb{C}, \mathcal{O}_\mathbb{C}) \times_{\mathrm{Spd}(K, \mathcal{O}_K)} \mathrm{Spd}(\mathbb{C}, \mathcal{O}_\mathbb{C}) \cong \mathrm{Spd}(\mathbb{C}, \mathcal{O}_\mathbb{C}) \times \underline{G} = \mathrm{Spd}(C^0(G, \mathbb{C}), C^0(G, \mathcal{O}_\mathbb{C})),$$

so

$$B[\frac{1}{J}]_p^\wedge \cong C^0(G, W(\mathbb{C}^b)).$$

With the help of theorem 4.6, this proves that a prismatic F-crystal is the same as an étale φ -module over $W(\mathbb{C}^b)$ together with a G -action which is semilinear with respect to the action of G on $W(\mathbb{C}^b)$. Now as \mathbb{C}^b is algebraically closed, it is well-known that the category of étale φ -modules

over $W(\mathbb{C}^b)$ is equivalent to the category of finite free \mathbb{Z}_p -modules via taking F -invariants, see [11] proposition 3.2.7 for example. This shows that the prismatic F-crystals are equivalent to finite free \mathbb{Z}_p -representations of G . \square

Remark 5.7. *Similarly as before, for a prismatic F-crystal M , the G -action on $M((A_{\text{inf}}(\mathcal{O}_{\mathbb{C}}), \text{Ker}(\theta)))$ is induced by the base change isomorphism (the crystal structure)*

$$M((A_{\text{inf}}(\mathcal{O}_{\mathbb{C}}), \text{Ker}(\theta))) \otimes_{W(\mathbb{C}^b), g} W(\mathbb{C}^b) \cong M((A_{\text{inf}}(\mathcal{O}_{\mathbb{C}}), \text{Ker}(\theta)))$$

with respect to the arrow

$$g : (A_{\text{inf}}(\mathcal{O}_{\mathbb{C}}), \text{Ker}(\theta)) \xrightarrow{\sim} (A_{\text{inf}}(\mathcal{O}_{\mathbb{C}}), \text{Ker}(\theta))$$

in $(\mathcal{O}_K)_{\Delta}$.

Corollary 5.8. *The category of (φ, Γ) -modules over \mathbf{A}_K is equivalent to the category of finite free \mathbb{Z}_p -representations of $G = \text{Gal}(\overline{K}/K)$. The equivalence functors are*

$$\begin{aligned} \mathcal{M} &\rightarrow (\mathcal{M} \otimes_{\mathbf{A}_K} W(\mathbb{C}^b))^{F \otimes \phi = 1} \\ T &\rightarrow (T \otimes_{\mathbb{Z}_p} W(\mathbb{C}^b))^{H=1} \xrightarrow{\text{deperfection}} (\cdot) \end{aligned}$$

where \mathcal{M} is a (φ, Γ) -module over \mathbf{A}_K , T is a finite free \mathbb{Z}_p -representation of G , and

$$H := \text{Gal}(\overline{K}/K(\zeta_{p^\infty})) \subset G.$$

The action of G is diagonal on both $T \otimes_{\mathbb{Z}_p} W(\mathbb{C}^b)$ and $\mathcal{M} \otimes_{\mathbf{A}_K} W(\mathbb{C}^b)$, where G acts on \mathcal{M} through the canonical quotient $G \rightarrow \Gamma$. The φ -structure on $T \otimes_{\mathbb{Z}_p} W(\mathbb{C}^b)$ is defined by ϕ on the second factor. Moreover, the deperfection functor is the equivalence from the category of (φ, Γ) -modules over $W(K(\zeta_{p^\infty})^b)$ to the category of (φ, Γ) -modules over \mathbf{A}_K , as induced from theorem 4.6.

Proof. Both categories are equivalent to prismatic F-crystals in $\mathcal{O}_{\Delta}[\frac{1}{\Gamma}]_p^\wedge$ -modules. We check the equivalence functors are given by the stated ones. Given a prismatic F-crystal M , the associated (φ, Γ) -module is $M((\mathbf{A}_K^+, (\phi^n([p]_q))))$, while the associated Galois representation is $M((A_{\text{inf}}(\mathcal{O}_{\mathbb{C}}), \text{Ker}(\theta)))^{\varphi=1}$. M being a crystal, we have a canonical identification

$$M((A_{\text{inf}}(\mathcal{O}_{\mathbb{C}}), \text{Ker}(\theta))) \cong M((\mathbf{A}_K^+, (\phi^n([p]_q)))) \otimes_{\mathbf{A}_K} W(\mathbb{C}^b) \otimes_{W(\mathbb{C}^b), \phi^{-n-1}} W(\mathbb{C}^b) \quad (3)$$

using the arrow

$$(\mathbf{A}_K^+, (\phi^n([p]_q))) \rightarrow (A_{\text{inf}}(\mathcal{O}_{\mathbb{C}}), (\phi^n([p]_q))) \xrightarrow{\phi^{-n-1}} (A_{\text{inf}}(\mathcal{O}_{\mathbb{C}}), \text{Ker}(\theta))$$

in $\text{Spf}(\mathcal{O}_K)_{\Delta}$. We have seen in the proof of proposition 4.2 that the base change along ϕ^{n+1} does not affect étale φ -modules, namely for any étale φ -module N over A with respect to $\phi : A \rightarrow A$, there is a canonical identification $N \otimes_{A, \phi^{n+1}} A \cong N$ of étale φ -modules over A . Hence we have a canonical identification

$$M((A_{\text{inf}}(\mathcal{O}_{\mathbb{C}}), \text{Ker}(\theta))) \cong M((\mathbf{A}_K^+, (\phi^n([p]_q)))) \otimes_{\mathbf{A}_K} W(\mathbb{C}^b) \quad (4)$$

of étale φ -modules over $W(\mathbb{C}^b)$. Thus the functor from (φ, Γ) -modules to Galois representations is of the expected form, we still need to identify the Galois action. This follows easily from remarks 5.7 and 5.4, as the Galois action on both sides of (3) are induced by base change, while the

identification (3) itself is also induced by base change. An easy base change computation together with the observation that the action of G on \mathbf{A}_K^+ through the inclusion $\mathbf{A}_K^+ \rightarrow A_{\text{inf}}(\mathcal{O}_{\mathbb{C}})$ is via the quotient $G \rightarrow \Gamma$ proves the compatibility of Galois action in (3). Then the naturality of the identification $N \otimes_{A, \phi^{n+1}} A \cong N$ gives us the desired description of Galois action on both sides of (4).

On the other hand, we have a canonical map

$$(A_{\text{inf}}(\mathcal{O}_{K(\zeta_{p^\infty})_p^\wedge}), \text{Ker}(\theta)) \longrightarrow (A_{\text{inf}}(\mathcal{O}_{\mathbb{C}}), \text{Ker}(\theta))/H$$

in the site $(\mathcal{O}_K)_{\Delta}^{\text{perf}}$, which is not necessarily an isomorphism, but becomes so on the structure sheaf $\mathcal{O}_{\Delta}[\frac{1}{I}]_p^\wedge$. Indeed, this follows from that \mathbb{C} has (continuous) Galois group H over $K(\zeta_p^\infty)_p^\wedge$ and the tilting equivalence of perfectoid fields. This tells us that

$$M((A_{\text{inf}}(\mathcal{O}_{K(\zeta_{p^\infty})_p^\wedge}), \text{Ker}(\theta))) \cong M((A_{\text{inf}}(\mathcal{O}_{\mathbb{C}}), \text{Ker}(\theta)))^{H=1}.$$

Moreover, by looking at the descent data of the F -crystal, the Γ -action on the right hand side is carried to the descent data of the left hand side. Then theorem 4.6 applied to \mathbf{A}_K gives the other direction. Note that the categorical equivalence in theorem 4.6 implies that base change preserves not only the φ -structure, but also Γ -action, thereby inducing an equivalence of (φ, Γ) -modules. \square

Remark 5.9. *The fact that twist by ϕ^{n+1} does not change étale φ -modules and has all the expected functoriality is used secretly throughout the above proof. For example, it is needed in checking the two functors are quasi-inverse to each other. All that says is that we can ignore the issue caused by twisting by ϕ .*

The equivalence functors in the above corollary may look different from the treatment one usually finds in the literature. We now check that they are equivalent.

Let \mathbf{A} be the p -adic completion of the maximal unramified extension of \mathbf{A}_K inside $W(\mathbb{C}^b)$, i.e. \mathbf{A} is the Cohen ring of $\mathbf{E}_K^{\text{sep}}$, the separable closure of \mathbf{E}_K , which lies inside $W(\mathbb{C}^b)$ and extends \mathbf{A}_K . It is stable by ϕ and the action of the Galois group G . If we can write \mathbf{A} as $A[\frac{1}{I}]_p^\wedge$ for some prism (A, I) , then we can repeat the above argument with $W(\mathbb{C}^b)$ replaced by \mathbf{A} , and deduce the usual description of equivalence functors between Galois representations and (φ, Γ) -modules. However, this is not possible since $\mathbf{E}_K^{\text{sep}}$ is not complete. Instead, we proceed with the following lemma.

Lemma 5.10. *There is an equivalence of categories*

$$\mathbf{EM}/_{\mathbf{A}} \xrightarrow{\sim} \mathbf{EM}/_{W(\mathbb{C}^b)}$$

induced by base change.

Proof. By [2] corollary 2.31,

$$(\text{colim}_{\phi} \mathbf{A})_p^\wedge \cong W(\text{colim}_{\phi} \mathbf{A}/p) \cong W(\text{colim}_{\phi} \mathbf{E}_K^{\text{sep}}) \cong W(\mathbf{E}_K^{\text{alg}}),$$

then lemma 4.3 implies that

$$\mathbf{EM}/_{\mathbf{A}} \xrightarrow{\sim} \mathbf{EM}/_{W(\mathbf{E}_K^{\text{alg}})}.$$

As both $\mathbf{E}_K^{\text{alg}}$ and \mathbb{C}^b are algebraically closed, étale φ -modules over $W(\mathbf{E}_K^{\text{alg}})$ and $W(\mathbb{C}^b)$ are both equivalent to finite free \mathbb{Z}_p -modules by taking F -invariants, we have the equivalence

$$\mathbf{EM}_{/W(\mathbf{E}_K^{\text{alg}})} \xrightarrow{\sim} \mathbf{EM}_{/W(\mathbb{C}^b)}.$$

Combining the two we have the desired equivalence. \square

Theorem 5.11. *The category of (φ, Γ) -modules over \mathbf{A}_K is equivalent to the category of finite free \mathbb{Z}_p -representations of $G = \text{Gal}(\overline{K}/K)$. The equivalence functors are*

$$\mathcal{M} \rightarrow (\mathcal{M} \otimes_{\mathbf{A}_K} \mathbf{A})^{F \otimes \phi = 1}$$

$$T \rightarrow (T \otimes_{\mathbb{Z}_p} \mathbf{A})^{H=1}$$

where \mathcal{M} is a (φ, Γ) -module over \mathbf{A}_K , T is a finite free \mathbb{Z}_p -representation of G , and

$$H := \text{Gal}(\overline{K}/K(\zeta_{p^\infty})) \subset G.$$

The action of G is diagonal on both $T \otimes_{\mathbb{Z}_p} \mathbf{A}$ and $\mathcal{M} \otimes_{\mathbf{A}_K} \mathbf{A}$, where G acts on \mathcal{M} through the canonical quotient $G \rightarrow \Gamma$. The φ -structure on $T \otimes_{\mathbb{Z}_p} \mathbf{A}$ is defined by ϕ on the second factor.

Proof. By lemma 5.10, the functor

$$\mathcal{M} \rightarrow (\mathcal{M} \otimes_{\mathbf{A}_K} W(\mathbb{C}^b))^{F \otimes \phi = 1}$$

in corollary 5.8 is the same as

$$\mathcal{M} \rightarrow (\mathcal{M} \otimes_{\mathbf{A}_K} \mathbf{A})^{F \otimes \phi = 1}.$$

Indeed, as étale φ -modules over $W(\mathbb{C}^b)$ are all isomorphic to the trivial ones $(W(\mathbb{C}^b)^n, \phi)$, lemma 5.10 tells us that étale φ -modules over \mathbf{A} are also of the form (\mathbf{A}^n, ϕ) , so taking F -invariants produces the same finite free \mathbb{Z}_p -modules, namely

$$(N \otimes_{\mathbf{A}} W(\mathbb{C}^b))^{F=1} \cong N^{F=1}$$

for any étale φ -module N over \mathbf{A} . This gives the identification of the \mathbb{Z}_p -modules, the identification of Galois actions also follows since the equivalence in lemma 5.10 is a categorical one, so arrows corresponding to Galois action are also preserved.

Conversely, let \mathcal{N} be the image of the functor

$$T \rightarrow (T \otimes_{\mathbb{Z}_p} W(\mathbb{C}^b))^{H=1} \xrightarrow{\text{deperfection}} (\cdot)$$

in corollary 5.8. Then by definition, \mathcal{N} is a (φ, Γ) -module over \mathbf{A}_K such that

$$\mathcal{N} \otimes_{\mathbf{A}_K} W((K(\zeta_{p^\infty})^\wedge)^b) \cong (T \otimes_{\mathbb{Z}_p} W(\mathbb{C}^b))^{H=1}.$$

Since the functor is the quasi-inverse of $\mathcal{M} \rightarrow (\mathcal{M} \otimes_{\mathbf{A}_K} W(\mathbb{C}^b))^{F \otimes \phi = 1}$, we have

$$\mathcal{N} \otimes_{\mathbf{A}_K} W(\mathbb{C}^b) \cong T \otimes_{\mathbb{Z}_p} W(\mathbb{C}^b),$$

whence

$$(\mathcal{N} \otimes_{\mathbf{A}_K} \mathbf{A}) \otimes_{\mathbf{A}} W(\mathbb{C}^b) \cong (T \otimes_{\mathbb{Z}_p} \mathbf{A}) \otimes_{\mathbf{A}} W(\mathbb{C}^b).$$

Then lemma 5.10 gives us

$$\mathcal{N} \otimes_{\mathbf{A}_K} \mathbf{A} \cong T \otimes_{\mathbb{Z}_p} \mathbf{A},$$

so

$$\mathcal{N} \cong (T \otimes_{\mathbb{Z}_p} \mathbf{A})^{H=1}.$$

We have now proved that the functors in corollary 5.8 is the same as the ones described in the statement of the theorem, so we can conclude using corollary 5.8. \square

REFERENCES

- [1] Jean-Marc Fontaine. *Représentations p-adiques des corps locaux (1ère partie)*, pages 249–309. Birkhäuser Boston, Boston, MA, 2007.
- [2] Bhargav Bhatt and Peter Scholze. Prisms and Prismatic Cohomology. *arXiv e-prints*, page arXiv:1905.08229, May 2019.
- [3] Bhargav Bhatt and Peter Scholze. Prismatic f-crystals and crystalline galois representations, 2021.
- [4] Peter Scholze. Etale cohomology of diamonds. *arXiv e-prints*, page arXiv:1709.07343, September 2017.
- [5] Peter Scholze and Jared Weinstein. *Berkeley lectures on p-adic geometry*. Princeton University Press, Princeton, NJ, 2020.
- [6] Olivier Brinon and Brian Conrad. Cmi summer school notes on p-adic hodge theory, 2009.
- [7] Bhargav Bhatt, Matthew Morrow, and Peter Scholze. Integral p-adic hodge theory. *Publications mathématiques de l’IHÉS*, 128(1):219–397, November 2018.
- [8] Ofer Gabber and Lorenzo Ramero. Foundations for almost ring theory – Release 7.5. *arXiv Mathematics e-prints*, page math/0409584, September 2004.
- [9] The Stacks project authors. The stacks project. <https://stacks.math.columbia.edu>, 2021.
- [10] Akhil Mathew. Faithfully flat descent of almost perfect complexes in rigid geometry. *arXiv e-prints*, page arXiv:1912.10968, December 2019.
- [11] Kiran Kedlaya and Ruochuan Liu. *Relative p-adic Hodge theory : foundations*. Société mathématique de France, Paris, 2015.