

RECEIVED: November 10, 2022

REVISED: May 15, 2023

ACCEPTED: June 22, 2023

PUBLISHED: July 14, 2023

Gravitational quadratic-in-spin Hamiltonian at NNNLO in the post-Newtonian framework

Manoj K. Mandal,^a Pierpaolo Mastrolia,^{b,a} Raj Patil^{c,d,e} and Jan Steinhoff^c

^a*INFN, Sezione di Padova,*

Via Marzolo 8, Padova I-35131, Italy

^b*Dipartimento di Fisica e Astronomia, Università degli Studi di Padova,*

Via Marzolo 8, Padova I-35131, Italy

^c*Max Planck Institute for Gravitational Physics (Albert Einstein Institute),*

Am Mühlenberg 1, Potsdam 14476, Germany

^d*Institut für Physik und IRIS Adlershof, Humboldt-Universität zu Berlin,*

Zum Großen Windkanal 2, Berlin D-12489, Germany

^e*Indian Institute of Science Education and Research Bhopal,*

Bhopal Bypass Rd, Bhauri, Madhya Pradesh 462066, India

E-mail: manojkumar.mandal@pd.infn.it, pierpaolo.mastrolia@unipd.it,
raj.patil@aei.mpg.de, jan.steinhoff@aei.mpg.de

ABSTRACT: We present the result of the quadratic-in-spin interaction Hamiltonian for binary systems of rotating compact objects with generic spins, up to N³LO corrections within the post-Newtonian expansion. The calculation is performed by employing the effective field theory diagrammatic approach, and it involves Feynman integrals up to three loops, evaluated within the dimensional regularization scheme. The gauge-invariant binding energy and the scattering angle, in special kinematic regimes and spin configurations, are explicitly derived. The results extend our earlier study on the spin-orbit interaction effects.

KEYWORDS: Classical Theories of Gravity, Effective Field Theories, Scattering Amplitudes, Black Holes

ARXIV EPRINT: [2210.09176](https://arxiv.org/abs/2210.09176)

Contents

1	Introduction	1
2	EFT of spinning objects	3
2.1	Action	3
2.2	Post-Newtonian formulation of general relativity	5
3	Computational algorithm	7
4	Processing the effective Lagrangian	10
4.1	Elimination of higher-order time derivatives	10
4.2	Computation of the EFT Hamiltonians	12
4.3	Removal of the poles and logarithms	12
4.3.1	3PN non-spinning sector	13
4.3.2	Quadratic-in-spin N ³ LO sector	14
5	Results	15
6	Computation of observables with spin	22
6.1	Binding energy for circular orbits with aligned spins	23
6.2	Scattering angle with aligned spins	24
7	Conclusion	26
A	Notation and convention	27
B	Lower-order Hamiltonians	28
B.1	Spin1-spin2 sector up to N ² LO	28
B.2	Spin1-spin1 and spin2-spin2 sector up to N ² LO	29
B.3	ES ² sector up to N ² LO	30

1 Introduction

The successful detection of the gravitational waves (GW) [1] by the LIGO-Virgo-KAGRA detectors opened a new era in precision astronomy and cosmology. Since then, the LVK collaboration has detected around 90 GW events [2]. The primary source of these GWs are the compact binaries and a worldwide network of ground-based [3–8] as well as space-based GW detectors [9] are coming up to explore the dynamical evolution of these compact binaries. The compact objects in the binary may be close to maximally rotating, as seen in several recent detections [10]. Therefore, high-precision waveforms incorporating spin

contributions are essential to exploit the full potential in GW astronomy. Initial studies extending the classical techniques to include spin were done in [11, 12], which was later extended by [13–27]. The spin effects in the post-Newtonian formalism were developed using the effective field theory approach in [28–43]. Another recently developed method of using quantum scattering amplitudes involving massive particles of arbitrary spin were used to obtain classical spin corrections to the two-body effective potential [44–52]. See [53–55] for recent reviews and a more comprehensive reference to literature.

In the effective field theory approach, the current state of the art for the post-Newtonian calculation for the conservative potential without spinning degrees of freedom is the 5PN correction computed in [56–60]. Several partial results of 6PN [61, 62] are also known. For the spin-orbit sector, the leading order (LO) effective potential was first computed in [28]. The next-to-leading order (NLO) potential in [29, 30], and N²LO in [63], and N³LO in [64, 65]. Similarly, for the quadratic in spin sector, the LO effective potential was computed in [28]; the NLO in [32–35]; and the N²LO in [36–38]. Partial results for N³LO were reported in [39, 40], and very recently the complete N³LO contribution has been reported in [66]. The computation for the cubic and higher orders in the spin variables can be found in [41–43], whereas the finite size effects are described in detail in [35, 42, 43]. Here, for maximally rotating compact objects, the NⁿLO spin-orbit sector contributes at $(3/2 + n)$ PN order and NⁿLO spin-squared sector contributes at $(2 + n)$ PN order.

In this article, following the same strategy adopted in [65], we present the complete conservative N³LO quadratic-in-spin interaction potential, obtained by adopting the EFT approach proposed in [28, 67] and using the computational diagrammatic techniques proposed in [68]. In particular, we begin with deriving the required Feynman rules and then the Feynman diagrams, where we employ the Kaluza-Klein (KK) parametrization for the metric, for convenience. The corresponding Feynman amplitudes contain tensor integrals, which are reduced to scalar integrals by means of a suitable set of projectors built out of Lorentz invariance. The emerging scalar integrals, up to three loop, are further reduced to a minimal set of independent integrals, dubbed master integrals (MIs), using the integration-by-parts (IBP) identities [69, 70]. The analytical expression of the MIs are used to obtain the analytic expression of the contribution of each Feynman diagram to the scattering amplitude. Finally, the effective Lagrangian is obtained by taking the Fourier transform of the amplitude. The computational procedure has been performed through in-house codes, in an automated manner using `Mathematica` routines with an interface to `QGRAF` [71], generating the diagrams, `xTensor` [72], manipulating tensor algebras, `LiteRed` [73], performing the IBP decomposition, inspired by several ideas implemented in `EFTofPNG` [74]. The derived effective Lagrangian contain higher-order time derivatives of the position and the spin, which are removed by applying appropriate coordinate transformations. Then, the EFT Hamiltonian is derived by applying the Legendre transformation. However, it contains poles in the parameter of the dimensional regularization $\epsilon = d - 3$ (d being the number of the continuous space dimensions), and logarithmic pieces depending on the scale of the binary system and they are removed following the application of appropriate canonical transformations, which is the main result of the article. The novel result for the quadratic-in-spin interaction Hamiltonian are further used to compute gauge invariant

observable, specifically, the binding energy for circular orbits with aligned spins and the scattering angle with aligned spins, and they agree with the results reported in [66].

The paper is organised as follows. In section 2, we review the description of the spinning binaries within the EFT formalism. In section 3, we present the computation for the N³LO quadratic-in-spin potential employing the Feynman diagrammatic approach within the EFT framework. Then, in section 4, we describe the procedure of removing the residual divergences and logarithms to derive the EFT Hamiltonian. We provide our main result of the quadratic-in-spin Hamiltonian up to N³LO in section 5. In section 6, we compute two observable from the EFT Hamiltonian, namely, the binding energy of the binary system in circular orbits with aligned spin configuration and scattering angle for two spinning compact objects with aligned spins. We summarize our main results in section 7. In appendix A, we describe the notations used in this article and in appendix B, we provide the Hamiltonians till N²LO in the quadratic-in-spin sector.

We provide the required EFT Feynman rules in the supplementary material as `Feynman_Rules.m` and the analytic results of the quadratic-in-spin Hamiltonian till N³LO in the supplementary material as `Hamiltonian.m`.

2 EFT of spinning objects

In this section, we describe the effective action for a spinning compact object at the orbital scale by considering the degrees of freedom of the gravitational field and the degrees of freedom of the spinning compact objects, namely their center of mass and their spin. Then we describe the techniques of the Post-Newtonian formulation of GR in detail and briefly outline the procedure to compute the effective action.

2.1 Action

The effective action of the spinning compact binary can be described as the sum of the action of the gravitational field and the point particle effective action for each of the spinning compact objects as

$$S_{\text{eff}} = S_{\text{EH}} + S_{\text{pp}}. \tag{2.1}$$

Here S_{EH} is the usual Einstein-Hilbert action and it expresses the dynamics of the gravitational field ($g_{\mu\nu}$) along with a gauge fixing term,

$$S_{\text{EH}} = -\frac{c^4}{16\pi G_{\text{N}}} \int d^4x \sqrt{g} R[g_{\mu\nu}] + \frac{c^4}{32\pi G_{\text{N}}} \int d^4x \sqrt{g} g_{\mu\nu} \Gamma^\mu \Gamma^\nu, \tag{2.2}$$

where $\Gamma^\mu = \Gamma^\mu_{\rho\sigma} g^{\rho\sigma}$ (in the harmonic gauge $\Gamma^\mu = 0$), $\Gamma^\mu_{\rho\sigma}$ is the Christoffel symbol, G_{N} is the Newton's constant, R is the Ricci scalar, and g is the determinant of the $g_{\mu\nu}$.

The spinning compact objects can be described by a point-particle effective action of each of them. This point-particle action can be written as an integral along a worldline in

the following way [42],

$$S_{\text{pp}} = \sum_{a=1,2} \int d\tau \left(-m_{(a)} c \sqrt{u_{(a)}^2} - \frac{1}{2} S_{(a)\mu\nu} \Omega_{(a)}^{\mu\nu} - \frac{S_{(a)\mu\nu} u_{(a)}^\nu}{u_{(a)}^2} \frac{du_{(a)}^\mu}{d\tau} + \mathcal{L}_{(a)}^{(R)} + \mathcal{L}_{(a)}^{(R^2)} + \dots \right), \quad (2.3)$$

where $\mathcal{L}_{(a)}^{(R)}$, $\mathcal{L}_{(a)}^{(R^2)}$, etc. denote Lagrangians containing nonminimal couplings at linear, quadratic, etc. order in curvature specified below. We use the Pryce, Newton, and Wigner gauge for spin supplementarity condition (SSC) given by $S_{(a)\mu\nu} (u_{(a)}^\nu + \sqrt{u_{(a)}^2} \delta^{\nu 0}) \approx 0$. Here, $u_{(a)}^2 = g_{\mu\nu} u_{(a)}^\mu u_{(a)}^\nu$, and $u_{(a)}^\mu$ is the four velocity, defined as $u_{(a)}^\mu = dx_{(a)}^\mu/d\tau$. The worldline $x_{(a)}^\mu(\tau)$ represents the center of the spinning object and is parametrized by an affine parameter τ , which we are going to gauge-fix to the coordinate time t by $\tau = ct$. The $\Omega_{(a)}^{\mu\nu}$ denotes the angular velocity tensor of the spinning object and is defined as

$$\Omega_{(a)}^{\mu\nu} = \Lambda_{(a)A}^\mu \frac{d\Lambda_{(a)}^{A\nu}}{d\tau}, \quad (2.4)$$

where $\Lambda_{(a)A}^\mu(\tau)$ represent the tetrads along the worldlines, connecting the body-fixed frame (denoted by upper case Latin indices) of the a^{th} compact object and the general coordinate frame (denoted by Greek indices). The $S_{(a)\mu\nu}$ are the spin tensors of the spinning objects, defined as the conjugate momenta to the $\Lambda_{(a)A}^\mu$ as

$$S_{(a)\mu\nu} = -2 \frac{\partial L_{\text{pp}}}{\partial \Omega_{(a)}^{\mu\nu}}. \quad (2.5)$$

The non-minimal coupling Lagrangians proportional to the different powers of curvature tensor R in equation (2.3) can be represented as

$$\mathcal{L}_{(a)}^{(R^m)} = \mathcal{L}_{(a)}^{(R^m, S^0)} + \mathcal{L}_{(a)}^{(R^m, S^1)} + \mathcal{L}_{(a)}^{(R^m, S^2)} + \dots \quad (2.6)$$

where $m = 1, 2, \dots$ and the n -th term in this series $\mathcal{L}_{(a)}^{(R^m, S^n)}$ denotes that n powers of the spin tensor is contracted explicitly to the Riemann tensor, only. Each of the Lagrangians in this series contain a number of Wilson coefficients. Following the convention in [42], these Wilson coefficients can be expressed as a function of the invariants¹ $m_{(a)}$ and the $S_{(a)}^2$. We then expand these Wilson coefficients explicitly as a series in $S_{(a)}^2$,

$$C_{(a)} = C_{(a)}^{(0)} + C_{(a)}^{(2)} \left(\frac{S_{(a)}^2 c^2}{G_N^2 m_{(a)}^4} \right) + C_{(a)}^{(4)} \left(\frac{S_{(a)}^2 c^2}{G_N^2 m_{(a)}^4} \right)^2 + \dots, \quad (2.7)$$

where the $C_{(a)}^{(n)}$ depends only on the $m_{(a)}$.

¹Since we are only computing the conservative dynamics, the absolute value of the spin tensor is conserved, i.e. $d(|S|)/dt = 0$. Hence the Wilson coefficients, even though explicitly depend on S^2 , are time-independent.

The contribution relevant for N³LO spin-squared from the spin-induced multipole interactions linear-in-curvature starts from $\mathcal{L}_{(a)}^{(R,S^2)}$ and is given by,

$$\mathcal{L}_{(a)}^{(R,S^2)} = -\frac{1}{2m_{(a)}c} \left(C_{\text{ES}^2}^{(0)}\right)_{(a)} \frac{E_{\mu\nu}}{u_{(a)}} \left[S_{(a)}^\mu S_{(a)}^\nu\right]_{\text{STF}} + \dots, \quad (2.8)$$

where $E_{\mu\nu} = R_{\mu\alpha\nu\beta}u^\alpha u^\beta$ and the STF denotes the symmetric-tracefree part (in a 3-dimensional comoving frame), which transforms irreducibly under the little group SO(3) of massive particles and hence it makes sense to construct interactions from STF building blocks.

Moving on to quadratic order in curvature, the contribution from $\mathcal{L}_{(a)}^{(R^2,S^0)}$,² entering at 5PN order, after using the expansion (2.7) is expressed as

$$\mathcal{L}_{(a)}^{(R^2,S^0)} = \frac{1}{2} \left(C_{\text{E}^2}^{(2)}\right)_{(a)} \frac{G_{\text{N}}^2 m_{(a)}}{c^5} \frac{E_{\mu\nu} E^{\mu\nu}}{u_{(a)}^3} S_{(a)}^2 + \dots. \quad (2.9)$$

The contribution from $\mathcal{L}_{(a)}^{(R^2,S^1)}$ start beyond the 5PN order and can be neglected here (since e.g. $E_{\mu\alpha} E_\nu^\alpha S_{(a)}^{\mu\nu} \equiv 0$). However, there is a relevant term in $\mathcal{L}_{(a)}^{(R^2,S^2)}$ [39], which reads as,

$$\mathcal{L}_{(a)}^{(R^2,S^2)} = \frac{1}{2} \left(C_{\text{E}^2\text{S}^2}^{(0)}\right)_{(a)} \frac{G_{\text{N}}^2 m_{(a)}}{c^5} \frac{E_{\mu\alpha} E_\nu^\alpha}{u_{(a)}^3} \left[S_{(a)}^\mu S_{(a)}^\nu\right]_{\text{STF}} + \dots. \quad (2.10)$$

These completed the description of the relevant contributions coming from all non-minimal couplings contributing at N³LO, see (2.8), (2.9), and (2.10). We note that in order to simplify the Feynman rules, we follow the custom to drop the STF projection in (2.8) and to replace in all nonminimal couplings the Weyl tensor in $E_{\mu\nu}$, $B_{\mu\nu}$ by the Riemann tensor (being equivalent to a field redefinition). The disadvantage of this practice is that spins in the C_{ES^2} coupling are not manifestly appearing in an STF combination any more, but could be brought into this form using a field redefinition.

We employ dimensional regularization for the computation of multi-loop Feynman diagrams and write the gravitational coupling constant in d spatial dimensions as

$$G_d = G_{\text{N}} \left(\sqrt{4\pi} e^{\gamma_{\text{E}}} R_0\right)^{d-3}, \quad (2.11)$$

where, γ_{E} is the Euler-Mascheroni constant, and R_0 is an arbitrary length scale.

2.2 Post-Newtonian formulation of general relativity

The inspiral phase of the binary compact objects contains three widely separated scales, namely, the length scale related to the single compact object (R_{s}) (Schwarzschild radius), the orbital separation of the binary (r), and the wavelength of the radiated gravitational wave (λ). The scales have the following hierarchy

$$R_{\text{s}} \ll r \ll \lambda. \quad (2.12)$$

²We point out that the quadratic powers of spin ($S_{(a)}^2$) on the r.h.s. of (2.9) is a scalar quantity and appears due to the use of the expansion of the Wilson coefficients, as shown in equation (2.7). As there is no explicit spin tensors contracted with the quadratic-in-curvature tensor (R^2), in the l.h.s. this contribution is denoted by $\mathcal{L}_{(a)}^{(R^2,S^0)}$.

Our goal in this article is to compute the contribution of the quadratic-in-spin effective potential up to N³LO. For this purpose, we further decompose the effective kinetic and potential terms in the following way

$$\mathcal{K}_{\text{eff}} = \mathcal{K}_{\text{pp}} + \mathcal{K}_{\text{spin}}, \quad \mathcal{V}_{\text{eff}} = \mathcal{V}_{\text{pp}} + \mathcal{V}_{\text{SO}} + \mathcal{V}_{\text{SS}}, \quad (2.16)$$

where \mathcal{K}_{pp} and \mathcal{V}_{pp} constitutes the kinetic and potential contributions for the center of mass degrees of freedom of the point particles and $\mathcal{K}_{\text{spin}}$, \mathcal{V}_{SO} , and \mathcal{V}_{SS} constitutes the kinetic, linear in spin potential and quadratic in spin potential contributions.

The effective kinetic terms can be expressed as

$$\mathcal{K}_{\text{pp}} = \sum_{a=1,2} m_{(a)} \left[\frac{1}{2} \mathbf{v}_a^2 + \frac{1}{8} \mathbf{v}_{(a)}^4 \left(\frac{1}{c^2} \right) + \frac{1}{16} \mathbf{v}_{(a)}^6 \left(\frac{1}{c^4} \right) + \frac{5}{128} \mathbf{v}_{(a)}^8 \left(\frac{1}{c^6} \right) \right] + \mathcal{O} \left(\frac{1}{c^8} \right), \quad (2.17)$$

$$\mathcal{K}_{\text{spin}} = \sum_{a=1,2} \left\{ -\frac{1}{2} \mathbf{S}_{(a)}^{ij} \boldsymbol{\Omega}_{(a)}^{ij} \left(\frac{1}{c} \right) + \mathbf{S}_{(a)}^{ij} \mathbf{v}_{(a)}^i \mathbf{a}_{(a)}^j \left(\frac{1}{c^3} \right) \left[\frac{1}{2} + \frac{3}{8} \mathbf{v}_{(a)}^2 \left(\frac{1}{c^2} \right) + \frac{5}{16} \mathbf{v}_{(a)}^4 \left(\frac{1}{c^4} \right) + \frac{35}{128} \mathbf{v}_{(a)}^6 \left(\frac{1}{c^6} \right) \right] + \mathcal{O} \left(\frac{1}{c^{11}} \right) \right\}, \quad (2.18)$$

and the effective potential terms can be written as follows

$$\mathcal{V}_{\text{pp}} = \mathcal{V}_{\text{N}} + \left(\frac{1}{c^2} \right) \mathcal{V}_{\text{1PN}} + \left(\frac{1}{c^4} \right) \mathcal{V}_{\text{2PN}} + \left(\frac{1}{c^6} \right) \mathcal{V}_{\text{3PN}} + \mathcal{O} \left(\frac{1}{c^8} \right), \quad (2.19)$$

$$\mathcal{V}_{\text{SO}} = \left(\frac{1}{c^3} \right) \left[\mathcal{V}_{\text{LO}}^{\text{SO}} + \left(\frac{1}{c^2} \right) \mathcal{V}_{\text{NLO}}^{\text{SO}} + \left(\frac{1}{c^4} \right) \mathcal{V}_{\text{N}^2\text{LO}}^{\text{SO}} \right] + \mathcal{O} \left(\frac{1}{c^9} \right), \quad (2.20)$$

$$\mathcal{V}_{\text{SS}} = \left(\frac{1}{c^4} \right) \left[\mathcal{V}_{\text{LO}}^{\text{SS}} + \left(\frac{1}{c^2} \right) \mathcal{V}_{\text{NLO}}^{\text{SS}} + \left(\frac{1}{c^4} \right) \mathcal{V}_{\text{N}^2\text{LO}}^{\text{SS}} + \left(\frac{1}{c^6} \right) \mathcal{V}_{\text{N}^3\text{LO}}^{\text{SS}} \right] + \mathcal{O} \left(\frac{1}{c^{12}} \right). \quad (2.21)$$

Here, \mathcal{V}_{N} corresponds to the Newtonian potential; \mathcal{V}_j with $j = \{1\text{PN}, 2\text{PN}, 3\text{PN}\}$ denotes the corresponding PN correction for the non-spinning part of the potential; $\mathcal{V}_j^{\text{SO}}$ with $j = \{\text{LO}, \text{NLO}, \text{N}^2\text{LO}\}$ denotes the corresponding correction to the spin-orbit coupling of the binary system; and $\mathcal{V}_j^{\text{SS}}$ with $j = \{\text{LO}, \text{NLO}, \text{N}^2\text{LO}, \text{N}^3\text{LO}\}$ denotes the corresponding correction to the quadratic-in-spin coupling of the binary system. Each $\mathcal{V}_j^{\text{SS}}$ is then separated with respect to their origin in the effective action as; S1S2 and S² sectors corresponding to the first three terms present given in (2.3), ES² corresponding to the action given in (2.8), E² corresponding to the action given in (2.9) and E²S² corresponding to the action given in (2.10). Our goal in this article is to compute the $\mathcal{V}_{\text{N}^3\text{LO}}^{\text{SS}}$ employing the advanced techniques of multi-loop scattering amplitudes, as further discussed in the next sections.

3 Computational algorithm

We employ a Feynman diagrammatic approach following equation (2.15) to compute the effective potential. First, we generate all the relevant topologies contributing at different

Order	Diagrams	Loops	Diagrams
LO	1	0	1
NLO	7	1	3
		0	4
N ² LO	58	2	27
		1	24
		0	7
N ³ LO	553	3	125
		2	342
		1	76
		0	10

 (a) Spin1-Spin2 and Spin1² (Spin2²) sector.

Order	Diagrams	Loops	Diagrams
LO	1	0	1
NLO	4	1	1
		0	3
N ² LO	25	2	7
		1	12
		0	6
N ³ LO	168	3	15
		2	101
		1	43
		0	9

 (b) ES² sector.

Order	Loops	Diagrams
LO	1	1

 (c) E² sector.

Order	Loops	Diagrams
LO	1	1

 (d) E²S² sector.

Table 1. Number of Feynman diagrams contributing different sectors.

orders of G_N . The virial theorem implies that contributions at NⁿLO constitute all the terms proportional to G_N^l where $l = 1, 2, \dots, n + 1$, and consequently, we take into account all the topologies at $l - 1$ loops for the contributions at specific order l . So, for the evaluation of the N³LO quadratic-in-spin potential, we generate all the relevant topologies till the order G_N^4 (3-loop) employing QGRAF [71]. There is 1 topology at order G_N (tree-level), 2 topologies at order G_N^2 (one-loop), 9 topologies at G_N^3 (two-loop), and 32 topologies at order G_N^4 (three-loop). The topologies are dressed with the KK field and we use the Feynman rules, obtained from the effective action of the PN expansion, to acquire all the Feynman diagrams contributing to the given order of G_N and v , depending on the specific perturbation order. We provide the relevant Feynman rules after the KK parameterization in the supplementary material as `Feynman_Rules.m`. The number of diagrams corresponding to the particular sector of quadratic-in-spin potential at a particular order in $1/c$ and of particular loop topology are shown in table 1.³ The diagrams contributing to the non-spinning and spin-orbit sector are provided in [65]. The generated Feynman diagrams can be understood as two-point multi-loop Feynman diagrams with all internal lines mass-less and the external momentum, identified with the momentum transferred between two sources, as shown in figure 1. We perform the tensor algebra using `xTensor` [72]

³While considering the spin effects, we count only the representative Feynman diagrams, where the spin can contribute from any of the world-line graviton interaction vertex present in the diagram. Additionally, the diagrams, which can be obtained from the change in the label $1 \leftrightarrow 2$, are not counted as separate diagrams.




Figure 1. Diagrammatic correspondence between the 4-point EFT-Gravity graphs and the 2-point QFT graphs.

to translate the Feynman diagrams to their corresponding Feynman Amplitudes. Consequently, we can write the generic expression of the effective potential corresponding to any l -loop Feynman graph G as the following

$$\mathcal{V}_G^{(l)} = \underbrace{N_C^{\mu_1, \mu_2, \dots}_{\nu_1, \nu_2, \dots}(x_{(a)}, \dots, S_{(a)}, \dots)}_{\text{Coefficient that depends on orbital variables}} \underbrace{\int_p e^{ip_\mu(x_{(1)} - x_{(2)})^\mu} N_F^{\alpha_1, \alpha_2, \dots}_{\mu_1, \mu_2, \dots}(p)}_{\text{Fourier integral}} \underbrace{\prod_{i=1}^l \int_{k_i} \frac{N_M^{\nu_1, \nu_2, \dots}_{\alpha_1, \alpha_2, \dots}(k_i)}{\prod_{\sigma \in G} D_\sigma(p, k_i)}}_{\text{Multi-loop integral}}, \tag{3.1}$$

- (i) N_C stands for a tensor polynomial built out of the world-line coordinates ($x_{(a)}^\mu$), the spin tensor ($S_{(a)\mu\nu}$), and their higher-order time derivatives,
- (ii) p denotes the transfer of the momentum between the sources (Fourier momentum),
- (iii) N_F is the tensor polynomial depending on p ,
- (iv) k_i denotes the loop momentum,
- (v) D_σ is the set of denominators corresponding to the graph G ,
- (vi) N_M is a tensor polynomial depending on external momenta p and loop momenta k_i .

We employ a set of projectors to reduce the multi-loop tensor integrals to scalar integrals. The projectors are built with the Lorentz invariant external momentum (p) and the background metric by exploiting Lorentz invariance. With the application of the projectors, the numerator (N_M) of the multi-loop integral translates to

$$N_M^{\nu_1, \nu_2, \dots}_{\alpha_1, \alpha_2, \dots}(p, k_i) \longrightarrow \tilde{N}_M(p, k_i) \tilde{N}_F^{\nu_1, \nu_2, \dots}_{\alpha_1, \alpha_2, \dots}(p), \tag{3.2}$$

where, \tilde{N}_M is a polynomial, which depends on the scalar products built out of external momentum (p) and the loop momenta (k_i). Specifically, we use projectors up to rank 6 to compute the N³LO quadratic-in-spin effective potential. Following this procedure, we obtain many scalar integrals, which are not all independent. There are linear relations between these scalar Feynman integrals, thanks to dimensional regularization, known as IBP relations [69]. These IBP identities can be generated and solved algorithmically

and automatically, thanks to several publicly available automatic IBP solvers [73, 77, 78]. Specifically, we use LiteRED [73] to build these IBP identities and solve them, thus obtaining a smaller set of independent scalar integrals, known as Master integrals (MIs). For the entire computation of N³LO quadratic-in-spin effective potential, we obtain 1 MI at one loop, 2 MIs at two-loop, and 3 MIs at three-loop. The MIs till three-loop are already known in the literature and have closed analytic expressions in d dimension. The expressions of these MIs are provided in appendix B of [65].

With the evaluation of the multi-loop integrals, we apply the Fourier transform to the tensor polynomials, which have the following generic form

$$\int_p e^{p_\mu(x_{(1)}-x_{(2)})^\mu} N_F^{\alpha_1, \alpha_2, \dots}_{\mu_1, \mu_2, \dots}(p) \tilde{N}_F^{\nu_1, \nu_2, \dots}_{\alpha_1, \alpha_2, \dots}(p) f(p). \quad (3.3)$$

Here, N_F and \tilde{N}_F are the tensor polynomials depending on the Fourier momenta p and $f(p)$ is a function of the external momentum obtained from the evaluation of the multi-loop integrals. We require Fourier integrals till rank 8. After the Fourier transformation, we expand the expression as a Laurent series in ϵ around $d = 3$ and obtain the effective potential, which depends on the orbital variables, namely, $\mathbf{x}_{(a)}$, $\mathbf{S}_{(a)}$, and their higher-order time derivatives.

We perform the generation of the relevant Feynman diagrams, translation of them to their corresponding multi-loop integrands, application of IBP reduction, and the Fourier transformation via our in-house code in an automated manner, thus obtaining the full effective potential. The different steps of this procedure have been explained using a flow chart in [65].

4 Processing the effective Lagrangian

The conservative quadratic-in-spin potential at N³LO contains higher-order time derivatives of the position ($\mathbf{a}_{(a)}$, $\dot{\mathbf{a}}_{(a)}$, $\ddot{\mathbf{a}}_{(a)}$, \dots) and the spin ($\dot{\mathbf{S}}_{(a)}$, $\ddot{\mathbf{S}}_{(a)}$, \dots) of the compact objects. Specifically, we obtain the 6th-order time derivative in the position and the 5th-order time derivative in the spin. Besides, the effective potential in the non-spinning sector at 3PN, the N³LO quadratic-in-spin potential also contains poles in the dimensional regularization parameter ϵ and the logarithmic terms of the form $\log(\frac{r}{R_0})$. We must eliminate higher-order time derivatives and the divergences to obtain the EFT Hamiltonian. All the logarithms in this computation result from a poor choice of coordinates, as there are no tail effects present either at the 3PN non-spinning sector or N³LO quadratic-in-spin sector. So, we can define a set of suitable coordinate transformations to eliminate these contributions.

First, we describe the elimination procedure of the higher-order time derivatives. We then demonstrate the appropriate coordinate transformations required to remove the divergent and the logarithmic terms, thus obtaining the EFT Hamiltonian free of them.

4.1 Elimination of higher-order time derivatives

Here, we discuss the procedure for the removal of all the higher-order time derivatives, by defining appropriate coordinate transformations [37, 79–82]. Under a small arbitrary shift

in the coordinate $\mathbf{x}_{(a)} \rightarrow \mathbf{x}_{(a)} + \delta\mathbf{x}_{(a)}$, the change in the Lagrangian is

$$\delta\mathcal{L} = \left(\frac{\delta\mathcal{L}}{\delta\mathbf{x}_{(a)}^i} \right) \delta\mathbf{x}_{(a)}^i + \frac{1}{2} \left(\frac{\delta^2\mathcal{L}}{\delta\mathbf{x}_{(a)}^i \delta\mathbf{x}_{(a)}^j} \right) \delta\mathbf{x}_{(a)}^i \delta\mathbf{x}_{(a)}^j + \mathcal{O}(\delta\mathbf{x}_{(a)}^3). \quad (4.1)$$

We can choose the small arbitrary shift $\delta\mathbf{x}_{(a)}$ in such a way that $\mathcal{L} + \delta\mathcal{L}$ is free of the $\mathbf{a}_{(a)}$, assuming that the equation (4.1) is linear in $\mathbf{a}_{(a)}$ at LO. We can also apply the total time derivative on the $\delta\mathbf{x}_{(a)}$ to modify the equations in such a way that terms involving higher-order time derivatives of $\mathbf{a}_{(a)}$ are removed. Following the same strategy, we can define a small arbitrary transformation simultaneously to the rotation matrix $\mathbf{\Lambda}_{(a)}^{ij} \rightarrow \mathbf{\Lambda}_{(a)}^{ij} + \delta\mathbf{\Lambda}_{(a)}^{ij}$ and the spin $\mathbf{S}_{(a)}^{ij} \rightarrow \mathbf{S}_{(a)}^{ij} + \delta\mathbf{S}_{(a)}^{ij}$ to eliminate the higher-order time derivatives in spin. A generic rotation can be expressed in terms of a matrix exponential $e^{\omega^{(a)}}$ such that the transformation of the rotation matrix and the spin can be described as $\mathbf{\Lambda}_{(a)} \rightarrow \mathbf{\Lambda}_{(a)} e^{\omega^{(a)}}$ and $\mathbf{S}_{(a)} \rightarrow e^{-\omega^{(a)}} \mathbf{S}_{(a)} e^{\omega^{(a)}}$, correspondingly. Consequently, the small arbitrary shift in the rotation matrix becomes

$$\delta\mathbf{\Lambda}_{(a)}^{ij} = \mathbf{\Lambda}_{(a)}^{ik} \omega_{(a)}^{kj} + \mathcal{O}(\omega_{(a)}^2), \quad (4.2)$$

where the $\omega_{(a)}^{kj}$ denotes the antisymmetric generator of the rotation matrix, and similarly the shift in spin becomes

$$\delta\mathbf{S}_{(a)}^{ij} = 2\mathbf{S}_{(a)}^{k[i} \omega_{(a)}^{j]k} + \mathcal{O}(\omega_{(a)}^2). \quad (4.3)$$

Following these transformations, the change in the Lagrangian⁴ becomes

$$\delta\mathcal{L} = - \left(\frac{1}{c} \right) \frac{1}{2} \dot{\mathbf{S}}_{(a)}^{ij} \omega_{(a)}^{ij} - \left(\frac{\delta V}{\delta \mathbf{S}_{(a)}^{ij}} \right) \delta\mathbf{S}_{(a)}^{ij} + \mathcal{O}(\omega_{(a)}^2, \delta\mathbf{S}_{(a)}^2). \quad (4.4)$$

So, assuming the equation of motion is linear in $\dot{\mathbf{S}}_{(a)}$ at LO, we can build the $\omega_{(a)}^{ij}$ in such a way that $\mathcal{L} + \delta\mathcal{L}$ is free of $\dot{\mathbf{S}}_{(a)}$. We can also apply the total time derivative on the equation of motion to define the modified equations in terms of higher-order time derivatives of $\dot{\mathbf{S}}_{(a)}$, thus removing them from the new effective Lagrangian.

We use this approach iteratively for the elimination of the higher-order time derivatives from the N³LO quadratic-in-spin potential. Specifically, we perform 5 iterations, where

1. we remove the terms with $\mathbf{a}_{(a)}$ and its higher-order time derivatives from the LO and NLO spin-orbit potentials,
2. we remove $\dot{\mathbf{S}}_{(a)}$ and its higher-order time derivatives from the NLO spin-orbit potential,
3. we remove the $\mathbf{a}_{(a)}$ and its higher-order time derivatives from non-spinning 2PN and 3PN potentials as well as LO and NLO quadratic-in-spin potentials,

⁴Here $V \equiv - \left(\mathcal{L} - \left(-\frac{1}{2} \mathbf{S}_{(a)}^{ij} \mathbf{\Omega}_{(a)}^{ij} \right) \right)$.

4. we remove $\dot{\mathbf{S}}_{(a)}$ and its higher-order time derivatives from the NLO quadratic-in-spin potential,
5. we remove the $\mathbf{a}_{(a)}$, $\dot{\mathbf{S}}_{(a)}$ and their higher-order time derivatives from the N²LO spin-orbit potential as well as N²LO and N³LO quadratic-in-spin potential.

After each iteration, we obtain a new Lagrangian and compute the equation of motion again to be used in the next iteration. One can check that at each step contributions cubic in $\delta\mathbf{x}_{(a)}$ and quadratic in $\boldsymbol{\omega}_{(a)}$ are negligible (higher order in spin or the PN approximation), in contrast to an (incorrect) insertion of all higher-order time derivatives in a single step. Following these steps, we obtain the effective Lagrangian, which depends on the position, velocity, and spin only.

4.2 Computation of the EFT Hamiltonians

We derive the EFT Hamiltonian by applying the Legendre transformation on the effective Lagrangian

$$\mathcal{H}(\mathbf{x}, \mathbf{p}, \mathbf{S}) = \sum_{a=1,2} \mathbf{p}_{(a)}^i \dot{\mathbf{x}}_{(a)}^i - \mathcal{L}(\mathbf{x}, \dot{\mathbf{x}}, \mathbf{S}), \quad (4.5)$$

where \mathbf{p}^i denotes the canonical momenta and it is defined as

$$\mathbf{p}_{(a)}^i = \frac{\partial \mathcal{L}(\mathbf{x}, \dot{\mathbf{x}}, \mathbf{S})}{\partial \dot{\mathbf{x}}_{(a)}^i}. \quad (4.6)$$

We express $\dot{\mathbf{x}}_{(a)}^i$ in terms of $\mathbf{p}_{(a)}^i$ by inverting this relation in every order of $1/c$. Exploiting this equation, we obtain a relation between $\dot{\mathbf{x}}_{(a)}^i$ and $\mathbf{p}_{(a)}^i$ and using it in the equation (4.5) we obtain the required Hamiltonian $\mathcal{H}(\mathbf{x}, \mathbf{p}, \mathbf{S})$.

4.3 Removal of the poles and logarithms

Both the Hamiltonian and the Lagrangian obtained in the previous step contain divergent pieces and logarithmic terms, which can be eliminated following appropriate coordinate transformations. In the Lagrangian description, we can employ a set of total derivative terms with arbitrary coefficients, which are determined by demanding the cancellation of the divergent pieces during the elimination of the higher-order time derivatives as described in section 4.1. Another alternative method to remove such spurious divergent terms and logarithmic terms is to add counter terms [83] in the point particle action (2.3), which can then be removed by a field redefinitions, i.e. they vanish on-shell. On the other hand, in the Hamiltonian description, we can utilize the properties of the canonical transformation⁵ to remove the poles and the logarithmic terms, namely

$$\mathcal{H}' = \mathcal{H} + \{\mathcal{H}, \mathcal{G}\}, \quad (4.7)$$

⁵The Poisson bracket is defined as

$$\{A, B\} = \sum_{i=1}^N \left(\frac{\partial A}{\partial r_{(i)}} \frac{\partial B}{\partial p_{(i)}} - \frac{\partial A}{\partial p_{(i)}} \frac{\partial B}{\partial r_{(i)}} \right) + \sum_{i=1}^N \left(\mathbf{S}_{(i)} \times \frac{\partial A}{\partial \mathbf{S}_{(i)}} \cdot \frac{\partial B}{\partial \mathbf{S}_{(i)}} \right),$$

where the last term is ignored in the context of the current analysis since it contributes to higher PN orders.

where \mathcal{G} represents the infinitesimal generator of the arbitrary canonical transformation and \mathcal{H}' is free of the divergences and logarithmic terms.

Here, we follow the procedure of removal of the poles and the logarithms in the Hamiltonian description, where we derive all the necessary suitable canonical transformations. Specifically, we construct an ansatz with several arbitrary coefficients for the infinitesimal generator \mathcal{G} and build a system of linear equations in terms of those unknown coefficients by requiring the elimination of the divergent pieces in \mathcal{H}' . The solutions of this system of equations provide a set of values for the arbitrary coefficients, thus obtaining the final effective Hamiltonian free of poles and logarithmic terms. In the next sections, we illustrate the strategies for the elimination of the poles and logarithmic terms from the 3PN non-spinning sector and the quadratic-in-spin N³LO sector.

4.3.1 3PN non-spinning sector

In the non-spinning sector the 3PN corrections has been known for a long time [84]. Following [84], we pursue the same procedure, by adding a total derivative term with the complete 3PN Lagrangian, to remove the divergent pieces.

$$\mathcal{L}_{\text{TD}} = \left(\frac{1}{c^6} \right) \frac{1}{\epsilon} \frac{d}{dt} \left[\frac{G_{\text{N}}^3}{r} \left(c_1 (\mathbf{v}_{(1)} \cdot \mathbf{n}) + c_2 (\mathbf{v}_{(2)} \cdot \mathbf{n}) \right) \right], \quad (4.8)$$

with,

$$c_1 = \frac{1}{3} \left(4m_{(1)}^3 m_{(2)} - m_{(1)} m_{(2)}^3 \right), \quad c_2 = \frac{1}{3} \left(m_{(1)}^3 m_{(2)} - 4m_{(1)} m_{(2)}^3 \right). \quad (4.9)$$

We start adopting the procedures for the removal of \mathbf{a}^i and $\dot{\mathbf{S}}^{ij}$ and their higher-order time derivatives as mentioned in section 4.1 with the 3PN Lagrangian along with the total derivative term. Ultimately, the divergent pieces are eliminated, thus deriving a finite Lagrangian as well as free of any higher-order time derivatives till 3PN. We build the Hamiltonian by applying the Legendre transformation on this Lagrangian and the Hamiltonian still contains the logarithmic terms. So, we construct the ansatz needed to define an appropriate canonical transformation in the Hamiltonian description to remove these logarithmic terms. We build the following generic ansatz for the arbitrary infinitesimal generator as,

$$\mathcal{G}_{\text{3PN}} = \left(\frac{1}{c^6} \right) \frac{G_{\text{N}}^3}{r^2} \left(g_1 \frac{1}{m_{(1)}} (\mathbf{p}_{(1)} \cdot \mathbf{n}) + g_2 \frac{1}{m_{(2)}} (\mathbf{p}_{(2)} \cdot \mathbf{n}) \right), \quad (4.10)$$

with,

$$g_n \equiv g_{Ln} \log \left(\frac{r}{R_0} \right), \quad \text{for } n = 1, 2. \quad (4.11)$$

Using this generator, we obtain the canonical transformation following eq. (4.7). We construct a system of linear equations in the unknown coefficients, defined in eq. (4.11) by demanding the cancellation of the logarithmic pieces. We solve these set of equations to obtain

$$g_{L1} = m_{(1)} m_{(2)}^3 - 4m_{(1)}^3 m_{(2)}, \quad g_{L2} = 4m_{(1)} m_{(2)}^3 - m_{(1)}^3 m_{(2)}. \quad (4.12)$$

4.3.2 Quadratic-in-spin N³LO sector

In the quadratic-in-spin sector at N³LO, we eliminate both the divergent and the logarithmic pieces by finding an appropriate canonical transformation in the Hamiltonian description. For this purpose, we construct the following ansatz for the infinitesimal generator stimulated by [37],

$$\mathcal{G}_{\text{SS-N}^3\text{LO}} = \mathcal{G}_{\text{S1S2-N}^3\text{LO}} + \mathcal{G}_{\text{S}^2\text{-N}^3\text{LO}} \quad (4.13)$$

where,

$$\begin{aligned} \mathcal{G}_{\text{S1S2-N}^3\text{LO}} = & \left(\frac{1}{c^{10}} \right) \frac{G_{\text{N}}^3}{r^4} \left\{ \left(\mathbf{S}_{(1)}^{ij} \mathbf{S}_{(2)}^{ij} \right) \left(g_3 \frac{1}{m_{(1)}} \left(\mathbf{p}_{(1)} \cdot \mathbf{n} \right) + g_4 \frac{1}{m_{(2)}} \left(\mathbf{p}_{(2)} \cdot \mathbf{n} \right) \right) \right. \\ & + \left(\mathbf{S}_{(1)}^{ik} \mathbf{S}_{(2)}^{jk} \mathbf{n}^i \mathbf{n}^j \right) \left(g_5 \frac{1}{m_{(1)}} \left(\mathbf{p}_{(1)} \cdot \mathbf{n} \right) + g_6 \frac{1}{m_{(2)}} \left(\mathbf{p}_{(2)} \cdot \mathbf{n} \right) \right) \\ & + g_7 \left(\mathbf{S}_{(1)}^{ik} \mathbf{S}_{(2)}^{jk} \mathbf{p}_{(1)}^i \mathbf{n}^j \right) + g_8 \left(\mathbf{S}_{(1)}^{ik} \mathbf{S}_{(2)}^{jk} \mathbf{n}^i \mathbf{p}_{(1)}^j \right) \\ & \left. + g_9 \left(\mathbf{S}_{(1)}^{ik} \mathbf{S}_{(2)}^{jk} \mathbf{p}_{(2)}^i \mathbf{n}^j \right) + g_{10} \left(\mathbf{S}_{(1)}^{ik} \mathbf{S}_{(2)}^{jk} \mathbf{n}^i \mathbf{p}_{(2)}^j \right) \right\}, \quad (4.14) \end{aligned}$$

$$\begin{aligned} \mathcal{G}_{\text{S}^2\text{-N}^3\text{LO}} = & \left(\frac{1}{c^{10}} \right) \frac{G_{\text{N}}^3}{r^4} \left\{ \left(\mathbf{S}_{(1)}^{ij} \mathbf{S}_{(1)}^{ij} \right) \left(g_{11} \frac{1}{m_{(1)}} \left(\mathbf{p}_{(1)} \cdot \mathbf{n} \right) + g_{12} \frac{1}{m_{(2)}} \left(\mathbf{p}_{(2)} \cdot \mathbf{n} \right) \right) \right. \\ & + \left(\mathbf{S}_{(1)}^{ik} \mathbf{S}_{(1)}^{jk} \mathbf{n}^i \mathbf{n}^j \right) \left(g_{13} \frac{1}{m_{(1)}} \left(\mathbf{p}_{(1)} \cdot \mathbf{n} \right) + g_{14} \frac{1}{m_{(2)}} \left(\mathbf{p}_{(2)} \cdot \mathbf{n} \right) \right) \\ & \left. + g_{15} \left(\mathbf{S}_{(1)}^{ik} \mathbf{S}_{(1)}^{jk} \mathbf{p}_{(1)}^i \mathbf{n}^j \right) + g_{16} \left(\mathbf{S}_{(1)}^{ik} \mathbf{S}_{(1)}^{jk} \mathbf{p}_{(2)}^i \mathbf{n}^j \right) \right\} + (1 \leftrightarrow 2). \quad (4.15) \end{aligned}$$

The g_n s are the unknown coefficients and defined as

$$g_n \equiv g_{\epsilon n} \frac{1}{\epsilon} + g_{Ln} \log \left(\frac{r}{R_0} \right), \quad \text{for } n = 3, 4, \dots, 16. \quad (4.16)$$

Following eq. (4.7), we use the ansatz for the infinitesimal generator defined in eq. (4.14), (4.15) to obtain the canonical transformation and prepare a system of linear equation in terms of the unknown coefficients in eq. (4.16) by requiring the cancellation of the divergent as well as the logarithmic pieces. Next, we solve them to determine the arbitrary coefficients as following

$$\begin{aligned} g_{\epsilon 3} &= \frac{1}{10} (5m_{(2)}^2 - 61m_{(1)}^2), & g_{\epsilon 4} &= \frac{1}{10} (61m_{(2)}^2 - 5m_{(1)}^2), \\ g_{\epsilon 5} &= \frac{1}{2} (61m_{(1)}^2 - 5m_{(2)}^2), & g_{\epsilon 6} &= \frac{1}{2} (5m_{(1)}^2 - 61m_{(2)}^2), \\ g_{\epsilon 7} &= -\frac{61m_{(1)}^2 - 5m_{(2)}^2}{10m_{(1)}}, & g_{\epsilon 8} &= -\frac{61m_{(1)}^2 - 5m_{(2)}^2}{10m_{(1)}}, \\ g_{\epsilon 9} &= -\frac{5m_{(1)}^2 - 61m_{(2)}^2}{10m_{(2)}}, & g_{\epsilon 10} &= -\frac{5m_{(1)}^2 - 61m_{(2)}^2}{10m_{(2)}}, \end{aligned}$$

$$\begin{aligned}
 g_{L3} &= \frac{3}{10} (61m_{(1)}^2 - 5m_{(2)}^2), & g_{L4} &= \frac{3}{10} (5m_{(1)}^2 - 61m_{(2)}^2), \\
 g_{L5} &= \frac{1}{2} (-3) (61m_{(1)}^2 - 5m_{(2)}^2), & g_{L6} &= \frac{1}{2} (-3) (5m_{(1)}^2 - 61m_{(2)}^2), \\
 g_{L7} &= \frac{3(61m_{(1)}^2 - 5m_{(2)}^2)}{10m_{(1)}}, & g_{L8} &= \frac{3(61m_{(1)}^2 - 5m_{(2)}^2)}{10m_{(1)}}, \\
 g_{L9} &= \frac{3(5m_{(1)}^2 - 61m_{(2)}^2)}{10m_{(2)}}, & g_{L10} &= \frac{3(5m_{(1)}^2 - 61m_{(2)}^2)}{10m_{(2)}}, \\
 g_{\epsilon 11} &= \frac{1}{70} \left(2m_1 m_2 - 159 \left(C_{\text{ES}^2}^{(0)} \right)_{(1)} m_1 m_2 \right), & g_{\epsilon 12} &= - \frac{\left(C_{\text{ES}^2}^{(0)} \right)_{(1)} (-175m_2^3 + 16m_1^2 m_2) + 30m_1^2 m_2}{70m_1}, \\
 g_{\epsilon 13} &= \frac{1}{14} \left(159 \left(C_{\text{ES}^2}^{(0)} \right)_{(1)} m_1 m_2 - 2m_1 m_2 \right), & g_{\epsilon 14} &= - \frac{\left(C_{\text{ES}^2}^{(0)} \right)_{(1)} (175m_2^3 - 16m_1^2 m_2) - 30m_1^2 m_2}{14m_1}, \\
 g_{\epsilon 15} &= \frac{1}{35} \left(2m_2 - 159 \left(C_{\text{ES}^2}^{(0)} \right)_{(1)} m_2 \right), & g_{\epsilon 16} &= - \frac{\left(C_{\text{ES}^2}^{(0)} \right)_{(1)} (16m_1^2 - 175m_2^2) + 30m_1^2}{35m_1}, \\
 g_{L11} &= \frac{3}{70} \left(159 \left(C_{\text{ES}^2}^{(0)} \right)_{(1)} m_1 m_2 - 2m_1 m_2 \right), & g_{L12} &= \frac{3 \left(\left(C_{\text{ES}^2}^{(0)} \right)_{(1)} (-175m_2^3 + 16m_1^2 m_2) + 30m_1^2 m_2 \right)}{70m_1}, \\
 g_{L13} &= \frac{(-3)}{14} \left(159 \left(C_{\text{ES}^2}^{(0)} \right)_{(1)} m_1 m_2 - 2m_1 m_2 \right), & g_{L14} &= - \frac{3 \left(\left(C_{\text{ES}^2}^{(0)} \right)_{(1)} (-175m_2^3 + 16m_1^2 m_2) + 30m_1^2 m_2 \right)}{14m_1}, \\
 g_{L15} &= \frac{3}{35} \left(159 \left(C_{\text{ES}^2}^{(0)} \right)_{(1)} m_2 - 2m_2 \right), & g_{L16} &= \frac{3 \left(\left(C_{\text{ES}^2}^{(0)} \right)_{(1)} (16m_1^2 - 175m_2^2) + 30m_1^2 \right)}{35m_1}.
 \end{aligned}$$

Ultimately, following these steps, we derive the full effective Hamiltonian, which is free of the divergent and logarithmic pieces, in the quadratic-in-spin sector till $N^3\text{LO}$.

5 Results

In this section, we illustrate the results for the effective Hamiltonian till $N^3\text{LO}$. For the convenience, we introduce a set of dimensionless variables, and write the Hamiltonian in terms of these variables, thus obtaining a compact form. We define the total mass $M = m_{(1)} + m_{(2)}$, and the reduced mass of the two body system $\mu = m_{(1)}m_{(2)}/M$, the mass ratio $q = m_{(1)}/m_{(2)}$, the symmetric mass ratio $\nu = \mu/M$, and the antisymmetric mass ratio $\delta = (m_{(1)} - m_{(2)})/M$. The relations between these dimensionless variables can be expressed as

$$\nu = \frac{m_{(1)}m_{(2)}}{M^2} = \frac{\mu}{M} = \frac{q}{(1+q)^2} = \frac{(1-\delta^2)}{4}. \quad (5.1)$$

Furthermore, we choose the center of mass (COM) frame of reference and show the Hamiltonian in the COM frame, for convenience. The momentum in the COM frame is defined as $\mathbf{p} \equiv \mathbf{p}_{(1)} = -\mathbf{p}_{(2)}$, and the orbital angular momentum can be expressed as $\mathbf{L} = (\mathbf{r} \times \mathbf{p})$.

Consequently, we obtain $p^2 = \tilde{p}_r^2 + \tilde{L}^2/\tilde{r}^2$, where $\tilde{p}_r = \mathbf{p} \cdot \mathbf{n}$, $p \equiv |\mathbf{p}|$ and $L \equiv |\mathbf{L}|$. Specifically, we use the following definitions in terms of dimensionless parameters

$$\tilde{\mathbf{p}} = \frac{\mathbf{p}}{\mu c}, \quad \tilde{\mathbf{r}} = \frac{\mathbf{r}}{G_N M}, \quad \tilde{\mathbf{L}} = \frac{\mathbf{L}}{G_N M \mu}, \quad \tilde{\mathbf{S}}_{(a)} = \frac{\mathbf{S}_{(a)}}{G_N M \mu}, \quad \tilde{\mathcal{H}} = \frac{\mathcal{H}}{\mu c^2}. \quad (5.2)$$

The total EFT Hamiltonian contributing towards the quadratic-in-spin sector till N³LO is written as

$$\tilde{\mathcal{H}} = \tilde{\mathcal{H}}_{\text{pp}} + \tilde{\mathcal{H}}_{\text{SO}} + \tilde{\mathcal{H}}_{\text{SS}}, \quad (5.3)$$

where,

$$\tilde{\mathcal{H}}_{\text{pp}} = \tilde{\mathcal{H}}_{0\text{PN}} + \left(\frac{1}{c^2}\right) \tilde{\mathcal{H}}_{1\text{PN}} + \left(\frac{1}{c^4}\right) \tilde{\mathcal{H}}_{2\text{PN}} + \left(\frac{1}{c^6}\right) \tilde{\mathcal{H}}_{3\text{PN}} + \mathcal{O}\left(\frac{1}{c^8}\right), \quad (5.4)$$

$$\tilde{\mathcal{H}}_{\text{SO}} = \left(\frac{1}{c^3}\right) \tilde{\mathcal{H}}_{\text{LO}}^{\text{SO}} + \left(\frac{1}{c^5}\right) \tilde{\mathcal{H}}_{\text{NLO}}^{\text{SO}} + \left(\frac{1}{c^7}\right) \tilde{\mathcal{H}}_{\text{N}^2\text{LO}}^{\text{SO}} + \mathcal{O}\left(\frac{1}{c^9}\right), \quad (5.5)$$

$$\begin{aligned} \tilde{\mathcal{H}}_{\text{SS}} &= \left(\frac{1}{c^4}\right) \left(\tilde{\mathcal{H}}_{\text{LO}}^{\text{S1S2}} + \tilde{\mathcal{H}}_{\text{LO}}^{\text{S}^2} + \tilde{\mathcal{H}}_{\text{LO}}^{\text{ES}^2}\right) + \left(\frac{1}{c^6}\right) \left(\tilde{\mathcal{H}}_{\text{NLO}}^{\text{S1S2}} + \tilde{\mathcal{H}}_{\text{NLO}}^{\text{S}^2} + \tilde{\mathcal{H}}_{\text{NLO}}^{\text{ES}^2}\right) \\ &+ \left(\frac{1}{c^8}\right) \left(\tilde{\mathcal{H}}_{\text{N}^2\text{LO}}^{\text{S1S2}} + \tilde{\mathcal{H}}_{\text{N}^2\text{LO}}^{\text{S}^2} + \tilde{\mathcal{H}}_{\text{N}^2\text{LO}}^{\text{ES}^2}\right) \\ &+ \left(\frac{1}{c^{10}}\right) \left(\tilde{\mathcal{H}}_{\text{N}^3\text{LO}}^{\text{S1S2}} + \tilde{\mathcal{H}}_{\text{N}^3\text{LO}}^{\text{S}^2} + \tilde{\mathcal{H}}_{\text{N}^3\text{LO}}^{\text{ES}^2} + \tilde{\mathcal{H}}_{\text{LO}}^{\text{E}^2\text{S}^2} + \tilde{\mathcal{H}}_{\text{LO}}^{\text{E}^2}\right) + \mathcal{O}\left(\frac{1}{c^{12}}\right). \end{aligned} \quad (5.6)$$

In the non-spinning part, the Hamiltonian till 3PN is known in the literature [84] and in the spin-orbit sector the Hamiltonian is known till N²LO [63]. We present the novel result of the $\tilde{\mathcal{H}}_{\text{N}^3\text{LO}}^{\text{SS}}$ in the COM frame following an EFT approach in the following as

$$\begin{aligned} \tilde{\mathcal{H}}_{\text{N}^3\text{LO}}^{\text{S1S2}} &= (\tilde{\mathbf{S}}_{(1)} \cdot \tilde{\mathbf{S}}_{(2)}) \\ &\times \left\{ \tilde{L}^6 \left(-\frac{5\nu^4}{64\tilde{r}^9} + \frac{7\nu^3}{32\tilde{r}^9} - \frac{45\nu^2}{16\tilde{r}^9} + \frac{5\nu}{8\tilde{r}^9} \right) \right. \\ &+ \tilde{L}^4 \left(\tilde{p}_r^2 \left(-\frac{27\nu^4}{64\tilde{r}^7} + \frac{81\nu^3}{4\tilde{r}^7} - \frac{603\nu^2}{64\tilde{r}^7} + \frac{15\nu}{16\tilde{r}^7} \right) - \frac{11\nu^3}{8\tilde{r}^8} - \frac{449\nu^2}{32\tilde{r}^8} + \frac{29\nu}{8\tilde{r}^8} \right) \\ &+ \tilde{L}^2 \left(\tilde{p}_r^2 \left(\frac{325\nu^3}{32\tilde{r}^6} - \frac{7365\nu^2}{64\tilde{r}^6} + \frac{47\nu}{4\tilde{r}^6} \right) + \tilde{p}_r^4 \left(-\frac{69\nu^4}{64\tilde{r}^5} + \frac{2025\nu^3}{32\tilde{r}^5} - \frac{891\nu^2}{64\tilde{r}^5} \right) \right. \\ &+ \frac{57\nu^3}{16\tilde{r}^7} + \frac{(12478 - 675\pi^2)\nu^2}{1200\tilde{r}^7} + \frac{49\nu}{4\tilde{r}^7} \left. \right) \\ &+ \tilde{p}_r^2 \left(\frac{145\nu^3}{8\tilde{r}^5} + \frac{(5400\pi^2 - 325199)\nu^2}{2400\tilde{r}^5} + \frac{9\nu}{2\tilde{r}^5} \right) + \tilde{p}_r^4 \left(\frac{783\nu^3}{32\tilde{r}^4} + \frac{4887\nu^2}{64\tilde{r}^4} - \frac{31\nu}{8\tilde{r}^4} \right) \\ &+ \tilde{p}_r^6 \left(-\frac{187\nu^4}{64\tilde{r}^3} - \frac{253\nu^3}{16\tilde{r}^3} + \frac{93\nu^2}{16\tilde{r}^3} - \frac{5\nu}{16\tilde{r}^3} \right) + \frac{(5296 - 6975\pi^2)\nu^2}{1600\tilde{r}^6} + \frac{67\nu}{2\tilde{r}^6} \left. \right\} \\ &+ ((\tilde{\mathbf{S}}_{(1)} \cdot \tilde{\mathbf{r}})(\tilde{\mathbf{S}}_{(2)} \cdot \tilde{\mathbf{r}})) \end{aligned}$$

$$\begin{aligned}
 & \times \left\{ \tilde{L}^6 \left(\frac{15\nu^4}{16\tilde{r}^{11}} - \frac{15\nu^3}{8\tilde{r}^{11}} + \frac{9\nu^2}{16\tilde{r}^{11}} \right) + \tilde{L}^4 \left(\tilde{p}_r^2 \left(\frac{225\nu^4}{64\tilde{r}^9} - \frac{779\nu^3}{32\tilde{r}^9} + \frac{69\nu^2}{8\tilde{r}^9} + \frac{5\nu}{16\tilde{r}^9} \right) - \frac{63\nu^3}{8\tilde{r}^{10}} - \frac{5\nu^2}{\tilde{r}^{10}} + \frac{17\nu}{8\tilde{r}^{10}} \right) \right. \\
 & + \tilde{L}^2 \left(\tilde{p}_r^2 \left(\frac{20\nu^3}{\tilde{r}^8} + \frac{1005\nu^2}{8\tilde{r}^8} - \frac{3\nu}{\tilde{r}^8} \right) + \tilde{p}_r^4 \left(\frac{177\nu^4}{32\tilde{r}^7} - \frac{1849\nu^3}{32\tilde{r}^7} + \frac{615\nu^2}{64\tilde{r}^7} + \frac{5\nu}{8\tilde{r}^7} \right) \right. \\
 & + \frac{79\nu^3}{16\tilde{r}^9} + \frac{(270\pi^2 - 17917)\nu^2}{96\tilde{r}^9} + \frac{107\nu}{4\tilde{r}^9} \left. \right) \\
 & + \tilde{p}_r^2 \left(\frac{327\nu^3}{8\tilde{r}^7} + \frac{(494789 - 5400\pi^2)\nu^2}{800\tilde{r}^7} - \frac{83\nu}{2\tilde{r}^7} \right) + \tilde{p}_r^4 \left(-\frac{3919\nu^3}{32\tilde{r}^6} - \frac{3575\nu^2}{64\tilde{r}^6} + \frac{7\nu}{8\tilde{r}^6} \right) \\
 & + \tilde{p}_r^6 \left(\frac{699\nu^4}{64\tilde{r}^5} + \frac{125\nu^3}{16\tilde{r}^5} - \frac{69\nu^2}{16\tilde{r}^5} + \frac{5\nu}{16\tilde{r}^5} \right) + \left. \frac{(20925\pi^2 - 71168)\nu^2}{1600\tilde{r}^8} - \frac{135\nu}{2\tilde{r}^8} \right\} \\
 & + ((\tilde{\mathbf{S}}_{(1)} \cdot \tilde{\mathbf{L}})(\tilde{\mathbf{S}}_{(2)} \cdot \tilde{\mathbf{L}})) \\
 & \times \left\{ \tilde{L}^4 \left(-\frac{15\nu^4}{16\tilde{r}^9} + \frac{59\nu^3}{16\tilde{r}^9} + \frac{39\nu^2}{8\tilde{r}^9} - \frac{25\nu}{16\tilde{r}^9} \right) \right. \\
 & + \tilde{L}^2 \left(\tilde{p}_r^2 \left(-\frac{27\nu^4}{8\tilde{r}^7} - \frac{25\nu^3}{2\tilde{r}^7} + \frac{489\nu^2}{32\tilde{r}^7} - \frac{25\nu}{8\tilde{r}^7} \right) + \frac{387\nu^3}{16\tilde{r}^8} + \frac{1799\nu^2}{32\tilde{r}^8} - \frac{51\nu}{4\tilde{r}^8} \right) \\
 & + \tilde{p}_r^2 \left(-\frac{1285\nu^3}{16\tilde{r}^6} - \frac{3141\nu^2}{32\tilde{r}^6} - \frac{75\nu}{4\tilde{r}^6} \right) + \tilde{p}_r^4 \left(-\frac{69\nu^4}{16\tilde{r}^5} - \frac{677\nu^3}{8\tilde{r}^5} + \frac{933\nu^2}{32\tilde{r}^5} - \frac{25\nu}{16\tilde{r}^5} \right) \\
 & - \left. \frac{22\nu^3}{\tilde{r}^7} + \frac{(43151 - 1350\pi^2)\nu^2}{1200\tilde{r}^7} - \frac{221\nu}{4\tilde{r}^7} \right\} \\
 & + ((\tilde{\mathbf{S}}_{(1)} \cdot \tilde{\mathbf{r}})(\tilde{\mathbf{S}}_{(2)} \cdot \tilde{\mathbf{L}})) \\
 & \times \tilde{p}_r \left\{ \tilde{L}^4 \left(-\frac{27\nu^4}{64\tilde{r}^9} - \frac{149\nu^3}{32\tilde{r}^9} + \frac{15\nu^2}{4\tilde{r}^9} - \frac{5\nu}{8\tilde{r}^9} \right) \right. \\
 & + \tilde{L}^2 \left(\left(-\frac{\nu^4}{2\tilde{r}^8} + \frac{1981\nu^3}{16\tilde{r}^8} - \frac{295\nu^2}{16\tilde{r}^8} - \frac{3\nu}{4\tilde{r}^8} \right) + \tilde{p}_r^2 \left(-\frac{45\nu^4}{32\tilde{r}^7} - \frac{1009\nu^3}{32\tilde{r}^7} + \frac{819\nu^2}{64\tilde{r}^7} - \frac{5\nu}{4\tilde{r}^7} \right) \right) \\
 & - \frac{4593\nu^3}{32\tilde{r}^7} + \frac{(10800\pi^2 - 842053)\nu^2}{2400\tilde{r}^7} + \frac{41\nu}{2\tilde{r}^7} + \tilde{p}_r^2 \left(-\frac{5\nu^4}{\tilde{r}^6} - \frac{795\nu^3}{16\tilde{r}^6} + \frac{603\nu^2}{16\tilde{r}^6} - \frac{27\nu}{4\tilde{r}^6} \right) \\
 & + \tilde{p}_r^4 \left(\frac{27\nu^4}{64\tilde{r}^5} - \frac{55\nu^3}{16\tilde{r}^5} + \frac{51\nu^2}{16\tilde{r}^5} - \frac{5\nu}{8\tilde{r}^5} \right) \\
 & + \frac{1}{q} \left(\tilde{L}^4 \left(\frac{3\nu^4}{4\tilde{r}^9} + \frac{9\nu^3}{8\tilde{r}^9} + \frac{15\nu^2}{16\tilde{r}^9} \right) + \tilde{L}^2 \left(\left(-\frac{\nu^4}{2\tilde{r}^8} + \frac{3057\nu^3}{32\tilde{r}^8} + \frac{3\nu^2}{2\tilde{r}^8} \right) + \tilde{p}_r^2 \left(\frac{27\nu^4}{8\tilde{r}^7} - \frac{33\nu^3}{2\tilde{r}^7} + \frac{15\nu^2}{8\tilde{r}^7} \right) \right) \right. \\
 & + \left. \frac{121\nu^2}{4\tilde{r}^7} - \frac{3227\nu^3}{32\tilde{r}^7} + \tilde{p}_r^2 \left(-\frac{5\nu^4}{\tilde{r}^6} - \frac{4269\nu^3}{32\tilde{r}^6} + \frac{27\nu^2}{2\tilde{r}^6} \right) + \tilde{p}_r^4 \left(\frac{147\nu^4}{16\tilde{r}^5} - \frac{9\nu^3}{2\tilde{r}^5} + \frac{15\nu^2}{16\tilde{r}^5} \right) \right) \left. \right\} \\
 & + ((\tilde{\mathbf{S}}_{(1)} \cdot \tilde{\mathbf{L}})(\tilde{\mathbf{S}}_{(2)} \cdot \tilde{\mathbf{r}})) \\
 & \times \tilde{p}_r \left\{ \tilde{L}^4 \left(-\frac{123\nu^4}{64\tilde{r}^9} - \frac{197\nu^3}{32\tilde{r}^9} + \frac{3\nu^2}{\tilde{r}^9} + \frac{5\nu}{16\tilde{r}^9} \right) \right. \\
 & + \tilde{L}^2 \left(\left(\frac{\nu^4}{2\tilde{r}^8} - \frac{271\nu^3}{4\tilde{r}^8} + \frac{2371\nu^2}{32\tilde{r}^8} + \frac{3\nu}{4\tilde{r}^8} \right) + \tilde{p}_r^2 \left(-\frac{261\nu^4}{32\tilde{r}^7} + \frac{155\nu^3}{32\tilde{r}^7} - \frac{477\nu^2}{64\tilde{r}^7} + \frac{5\nu}{8\tilde{r}^7} \right) \right) \left. \right\}
 \end{aligned}$$

$$\begin{aligned}
& + \frac{1861\nu^3}{32\tilde{r}^7} + \frac{(5400\pi^2 - 614639)\nu^2}{1200\tilde{r}^7} + \frac{203\nu}{4\tilde{r}^7} + \tilde{p}_r^2 \left(\frac{5\nu^4}{\tilde{r}^6} + \frac{1697\nu^3}{8\tilde{r}^6} - \frac{3927\nu^2}{32\tilde{r}^6} + \frac{27\nu}{4\tilde{r}^6} \right) \\
& + \tilde{p}_r^4 \left(-\frac{1149\nu^4}{64\tilde{r}^5} + \frac{59\nu^3}{4\tilde{r}^5} - \frac{51\nu^2}{16\tilde{r}^5} + \frac{5\nu}{16\tilde{r}^5} \right) \\
& + \frac{1}{q} \left(\tilde{L}^4 \left(-\frac{3\nu^4}{4\tilde{r}^9} - \frac{9\nu^3}{8\tilde{r}^9} - \frac{15\nu^2}{16\tilde{r}^9} \right) + \tilde{L}^2 \left(\left(\frac{\nu^4}{2\tilde{r}^8} - \frac{3057\nu^3}{32\tilde{r}^8} - \frac{3\nu^2}{2\tilde{r}^8} \right) + \tilde{p}_r^2 \left(-\frac{27\nu^4}{8\tilde{r}^7} + \frac{33\nu^3}{2\tilde{r}^7} - \frac{15\nu^2}{8\tilde{r}^7} \right) \right) \right. \\
& \left. + \left(\frac{3227\nu^3}{32\tilde{r}^7} - \frac{121\nu^2}{4\tilde{r}^7} \right) + \tilde{p}_r^2 \left(\frac{5\nu^4}{\tilde{r}^6} + \frac{4269\nu^3}{32\tilde{r}^6} - \frac{27\nu^2}{2\tilde{r}^6} \right) + \tilde{p}_r^4 \left(-\frac{147\nu^4}{16\tilde{r}^5} + \frac{9\nu^3}{2\tilde{r}^5} - \frac{15\nu^2}{16\tilde{r}^5} \right) \right) \Bigg\}, \tag{5.7}
\end{aligned}$$

$$\tilde{\mathcal{H}}_{\text{N}^3\text{LO}}^{\text{S}^2} = (\tilde{\mathbf{S}}_{(1)} \cdot \tilde{\mathbf{S}}_{(1)})$$

$$\begin{aligned}
& \times \left\{ \tilde{L}^6 \left(\frac{67\nu^4}{32\tilde{r}^9} - \frac{499\nu^3}{128\tilde{r}^9} + \frac{47\nu^2}{64\tilde{r}^9} \right) + \tilde{L}^4 \left(\tilde{p}_r^2 \left(\frac{1113\nu^4}{128\tilde{r}^7} - \frac{21\nu^3}{16\tilde{r}^7} + \frac{99\nu^2}{128\tilde{r}^7} \right) - \frac{169\nu^4}{128\tilde{r}^8} - \frac{5027\nu^3}{128\tilde{r}^8} + \frac{947\nu^2}{128\tilde{r}^8} \right) \right. \\
& + \tilde{L}^2 \left(\tilde{p}_r^2 \left(-\frac{247\nu^4}{64\tilde{r}^6} + \frac{3179\nu^3}{64\tilde{r}^6} - \frac{15\nu^2}{64\tilde{r}^6} \right) + \tilde{p}_r^4 \left(\frac{801\nu^4}{64\tilde{r}^5} - \frac{669\nu^3}{128\tilde{r}^5} - \frac{21\nu^2}{32\tilde{r}^5} \right) + \frac{(21311 + 225\pi^2)\nu^3}{600\tilde{r}^7} + \frac{48541\nu^2}{1960\tilde{r}^7} \right) \\
& + \tilde{p}_r^2 \left(-\frac{(121757 + 25200\pi^2)\nu^3}{16800\tilde{r}^5} - \frac{25553\nu^2}{3920\tilde{r}^5} \right) + \tilde{p}_r^4 \left(\frac{587\nu^4}{128\tilde{r}^4} + \frac{5187\nu^3}{128\tilde{r}^4} - \frac{785\nu^2}{128\tilde{r}^4} \right) \\
& + \tilde{p}_r^6 \left(-\frac{293\nu^4}{128\tilde{r}^3} + \frac{235\nu^3}{64\tilde{r}^3} - \frac{89\nu^2}{128\tilde{r}^3} \right) + \frac{23913\nu^2}{3920\tilde{r}^6} - \frac{(21619 + 3150\pi^2)\nu^3}{8400\tilde{r}^6} \\
& + \frac{1}{q} \left(\tilde{L}^6 \left(\frac{253\nu^4}{128\tilde{r}^9} - \frac{75\nu^3}{8\tilde{r}^9} + \frac{861\nu^2}{128\tilde{r}^9} - \frac{47\nu}{64\tilde{r}^9} \right) \right. \\
& + \tilde{L}^4 \left(\tilde{p}_r^2 \left(\frac{261\nu^4}{32\tilde{r}^7} - \frac{831\nu^3}{64\tilde{r}^7} + \frac{165\nu^2}{64\tilde{r}^7} - \frac{99\nu}{128\tilde{r}^7} \right) - \frac{169\nu^4}{128\tilde{r}^8} - \frac{2123\nu^3}{64\tilde{r}^8} + \frac{1531\nu^2}{128\tilde{r}^8} - \frac{899\nu}{128\tilde{r}^8} \right) \\
& + \tilde{L}^2 \left(\tilde{p}_r^2 \left(-\frac{247\nu^4}{64\tilde{r}^6} + \frac{4319\nu^3}{64\tilde{r}^6} + \frac{7103\nu^2}{64\tilde{r}^6} + \frac{7\nu}{64\tilde{r}^6} \right) + \tilde{p}_r^4 \left(\frac{1479\nu^4}{128\tilde{r}^5} - \frac{9\nu^3}{8\tilde{r}^5} - \frac{1083\nu^2}{128\tilde{r}^5} + \frac{21\nu}{32\tilde{r}^5} \right) \right. \\
& + \frac{(149027 + 1575\pi^2)\nu^3}{4200\tilde{r}^7} + \frac{(13568 + 1341\pi^2)\nu^2}{6144\tilde{r}^7} - \frac{221\nu}{8\tilde{r}^7} \Bigg\} \\
& + \tilde{p}_r^2 \left(-\frac{(38207 + 25200\pi^2)\nu^3}{16800\tilde{r}^5} + \frac{(3952 - 1341\pi^2)\nu^2}{1536\tilde{r}^5} + \frac{185\nu}{16\tilde{r}^5} \right) \\
& + \tilde{p}_r^4 \left(\frac{587\nu^4}{128\tilde{r}^4} + \frac{3949\nu^3}{64\tilde{r}^4} - \frac{2101\nu^2}{128\tilde{r}^4} + \frac{721\nu}{128\tilde{r}^4} \right) + \tilde{p}_r^6 \left(-\frac{181\nu^4}{64\tilde{r}^3} + \frac{369\nu^3}{64\tilde{r}^3} - \frac{69\nu^2}{16\tilde{r}^3} + \frac{89\nu}{128\tilde{r}^3} \right) \\
& - \left. \left(\frac{(21619 + 3150\pi^2)\nu^3}{8400\tilde{r}^6} + \frac{(587 - 9\pi^2)\nu^2}{48\tilde{r}^6} + \frac{155\nu}{16\tilde{r}^6} \right) \right\} \\
& + (\tilde{\mathbf{S}}_{(1)} \cdot \tilde{\mathbf{F}})^2 \\
& \times \left\{ \tilde{L}^4 \left(\tilde{p}_r^2 \left(-\frac{133\nu^4}{128\tilde{r}^9} - \frac{265\nu^3}{64\tilde{r}^9} + \frac{41\nu^2}{128\tilde{r}^9} \right) - \frac{113\nu^4}{128\tilde{r}^{10}} - \frac{795\nu^3}{128\tilde{r}^{10}} + \frac{159\nu^2}{128\tilde{r}^{10}} \right) \right. \\
& + \tilde{L}^2 \left(\tilde{p}_r^2 \left(-\frac{93\nu^4}{32\tilde{r}^8} - \frac{1005\nu^3}{32\tilde{r}^8} + \frac{61\nu^2}{16\tilde{r}^8} \right) + \tilde{p}_r^4 \left(-\frac{145\nu^4}{64\tilde{r}^7} + \frac{35\nu^3}{32\tilde{r}^7} + \frac{65\nu^2}{64\tilde{r}^7} \right) + \frac{(1832 - 315\pi^2)\nu^3}{168\tilde{r}^9} + \frac{22779\nu^2}{784\tilde{r}^9} \right) \Bigg\}
\end{aligned}$$

$$\begin{aligned}
 & + \widetilde{p}_r^2 \left(\frac{(4927+25200\pi^2)\nu^3}{5600\widetilde{r}^7} - \frac{118151\nu^2}{3920\widetilde{r}^7} \right) + \widetilde{p}_r^4 \left(-\frac{1147\nu^4}{128\widetilde{r}^6} - \frac{2483\nu^3}{128\widetilde{r}^6} + \frac{481\nu^2}{128\widetilde{r}^6} \right) \\
 & + \widetilde{p}_r^6 \left(\frac{293\nu^4}{128\widetilde{r}^5} - \frac{235\nu^3}{64\widetilde{r}^5} + \frac{89\nu^2}{128\widetilde{r}^5} \right) + \frac{(3150\pi^2-11071)\nu^3}{2800\widetilde{r}^8} - \frac{71809\nu^2}{3920\widetilde{r}^8} \\
 & + \frac{1}{q} \left(\widetilde{L}^4 \left(\widetilde{p}_r^2 \left(-\frac{59\nu^4}{64\widetilde{r}^9} - \frac{327\nu^3}{64\widetilde{r}^9} + \frac{15\nu^2}{8\widetilde{r}^9} - \frac{41\nu}{128\widetilde{r}^9} \right) - \frac{113\nu^4}{128\widetilde{r}^{10}} - \frac{101\nu^3}{16\widetilde{r}^{10}} + \frac{267\nu^2}{128\widetilde{r}^{10}} - \frac{31\nu}{128\widetilde{r}^{10}} \right) \right. \\
 & + \widetilde{L}^2 \left(\widetilde{p}_r^2 \left(-\frac{93\nu^4}{32\widetilde{r}^8} - \frac{1633\nu^3}{64\widetilde{r}^8} - \frac{573\nu^2}{32\widetilde{r}^8} - \frac{75\nu}{16\widetilde{r}^8} \right) + \widetilde{p}_r^4 \left(-\frac{59\nu^4}{32\widetilde{r}^7} - \frac{183\nu^3}{32\widetilde{r}^7} + \frac{159\nu^2}{16\widetilde{r}^7} - \frac{65\nu}{64\widetilde{r}^7} \right) \right. \\
 & + \frac{(7841-1260\pi^2)\nu^3}{672\widetilde{r}^9} + \frac{(235904-6705\pi^2)\nu^2}{6144\widetilde{r}^9} - \frac{35\nu}{16\widetilde{r}^9} \left. \right) \\
 & + \widetilde{p}_r^2 \left(\frac{9(7053+2800\pi^2)\nu^3}{5600\widetilde{r}^7} + \frac{3(447\pi^2-2032)\nu^2}{512\widetilde{r}^7} - \frac{185\nu}{16\widetilde{r}^7} \right) \\
 & + \widetilde{p}_r^4 \left(-\frac{1147\nu^4}{128\widetilde{r}^6} - \frac{3221\nu^3}{64\widetilde{r}^6} + \frac{2101\nu^2}{128\widetilde{r}^6} - \frac{721\nu}{128\widetilde{r}^6} \right) + \widetilde{p}_r^6 \left(\frac{181\nu^4}{64\widetilde{r}^5} - \frac{369\nu^3}{64\widetilde{r}^5} + \frac{69\nu^2}{16\widetilde{r}^5} - \frac{89\nu}{128\widetilde{r}^5} \right) \\
 & + \left. \frac{(3150\pi^2-11071)\nu^3}{2800\widetilde{r}^8} + \frac{3(3\pi^2-179)\nu^2}{16\widetilde{r}^8} - \frac{155\nu}{16\widetilde{r}^8} \right\} \\
 & + (\widetilde{\mathbf{S}}_{(1)} \cdot \widetilde{\mathbf{L}})^2 \\
 & \times \left\{ \widetilde{L}^4 \left(-\frac{661\nu^4}{128\widetilde{r}^9} + \frac{1087\nu^3}{128\widetilde{r}^9} - \frac{193\nu^2}{128\widetilde{r}^9} \right) + \widetilde{L}^2 \left(\widetilde{p}_r^2 \left(-\frac{1423\nu^4}{64\widetilde{r}^7} + \frac{697\nu^3}{64\widetilde{r}^7} - \frac{169\nu^2}{64\widetilde{r}^7} \right) + \frac{297\nu^4}{64\widetilde{r}^8} + \frac{5345\nu^3}{64\widetilde{r}^8} - \frac{1217\nu^2}{64\widetilde{r}^8} \right) \right. \\
 & + \widetilde{p}_r^2 \left(\frac{1005\nu^4}{64\widetilde{r}^6} + \frac{3277\nu^3}{64\widetilde{r}^6} - \frac{1141\nu^2}{64\widetilde{r}^6} \right) + \widetilde{p}_r^4 \left(-\frac{4675\nu^4}{128\widetilde{r}^5} + \frac{2317\nu^3}{128\widetilde{r}^5} - \frac{145\nu^2}{128\widetilde{r}^5} \right) \\
 & + \frac{(12600\pi^2-1208189)\nu^3}{16800\widetilde{r}^7} - \frac{114069\nu^2}{980\widetilde{r}^7} \\
 & + \frac{1}{q} \left(\widetilde{L}^4 \left(-\frac{601\nu^4}{128\widetilde{r}^9} + \frac{1431\nu^3}{64\widetilde{r}^9} - \frac{1695\nu^2}{128\widetilde{r}^9} + \frac{193\nu}{128\widetilde{r}^9} \right) \right. \\
 & + \widetilde{L}^2 \left(\widetilde{p}_r^2 \left(-\frac{1291\nu^4}{64\widetilde{r}^7} + \frac{375\nu^3}{8\widetilde{r}^7} - \frac{1119\nu^2}{64\widetilde{r}^7} + \frac{169\nu}{64\widetilde{r}^7} \right) + \frac{297\nu^4}{64\widetilde{r}^8} + \frac{2135\nu^3}{32\widetilde{r}^8} - \frac{3005\nu^2}{64\widetilde{r}^8} + \frac{1089\nu}{64\widetilde{r}^8} \right) \\
 & + \widetilde{p}_r^2 \left(\frac{1005\nu^4}{64\widetilde{r}^6} - \frac{331\nu^3}{32\widetilde{r}^6} - \frac{21555\nu^2}{64\widetilde{r}^6} + \frac{1013\nu}{64\widetilde{r}^6} \right) + \widetilde{p}_r^4 \left(-\frac{4231\nu^4}{128\widetilde{r}^5} + \frac{1449\nu^3}{64\widetilde{r}^5} - \frac{1023\nu^2}{128\widetilde{r}^5} + \frac{145\nu}{128\widetilde{r}^5} \right) \\
 & + \left. \frac{(12600\pi^2-1175639)\nu^3}{16800\widetilde{r}^7} + \frac{(1341\pi^2-317152)\nu^2}{3072\widetilde{r}^7} + \frac{353\nu}{4\widetilde{r}^7} \right\} \\
 & + (\widetilde{\mathbf{S}}_{(1)} \cdot \widetilde{\mathbf{F}})(\widetilde{\mathbf{S}}_{(1)} \cdot \widetilde{\mathbf{L}}) \\
 & \times \widetilde{p}_r \left\{ \widetilde{L}^4 \left(-\frac{401\nu^4}{128\widetilde{r}^9} - \frac{31\nu^3}{128\widetilde{r}^9} - \frac{53\nu^2}{128\widetilde{r}^9} \right) + \widetilde{L}^2 \left(\left(\frac{65\nu^4}{16\widetilde{r}^8} + \frac{55\nu^3}{4\widetilde{r}^8} - \frac{25\nu^2}{16\widetilde{r}^8} \right) + \widetilde{p}_r^2 \left(-\frac{635\nu^4}{64\widetilde{r}^7} + \frac{419\nu^3}{64\widetilde{r}^7} - \frac{5\nu^2}{64\widetilde{r}^7} \right) \right) \right. \\
 & + \widetilde{p}_r^2 \left(\frac{185\nu^4}{16\widetilde{r}^6} - \frac{1219\nu^3}{16\widetilde{r}^6} + \frac{37\nu^2}{16\widetilde{r}^6} \right) + \widetilde{p}_r^4 \left(-\frac{1019\nu^4}{128\widetilde{r}^5} + \frac{59\nu^3}{128\widetilde{r}^5} + \frac{43\nu^2}{128\widetilde{r}^5} \right) \\
 & + \left. \frac{109993\nu^2}{3920\widetilde{r}^7} - \frac{(155101+12600\pi^2)\nu^3}{2100\widetilde{r}^7} \right\}
 \end{aligned}$$

$$\begin{aligned}
 & + \frac{1}{q} \left(\tilde{L}^4 \left(-\frac{371\nu^4}{128\tilde{r}^9} + \frac{273\nu^3}{64\tilde{r}^9} - \frac{621\nu^2}{128\tilde{r}^9} + \frac{53\nu}{128\tilde{r}^9} \right) \right. \\
 & + \tilde{L}^2 \left(\left(\frac{65\nu^4}{16\tilde{r}^8} + \frac{13\nu^3}{2\tilde{r}^8} - \frac{549\nu^2}{16\tilde{r}^8} + \frac{49\nu}{16\tilde{r}^8} \right) + \tilde{p}_r^2 \left(-\frac{581\nu^4}{64\tilde{r}^7} + \frac{99\nu^3}{8\tilde{r}^7} + \frac{351\nu^2}{64\tilde{r}^7} + \frac{5\nu}{64\tilde{r}^7} \right) \right) \\
 & + \tilde{p}_r^2 \left(\frac{185\nu^4}{16\tilde{r}^6} - \frac{1093\nu^3}{16\tilde{r}^6} - \frac{1383\nu^2}{16\tilde{r}^6} - \frac{13\nu}{16\tilde{r}^6} \right) + \tilde{p}_r^4 \left(-\frac{881\nu^4}{128\tilde{r}^5} + \frac{69\nu^3}{64\tilde{r}^5} + \frac{363\nu^2}{128\tilde{r}^5} - \frac{43\nu}{128\tilde{r}^5} \right) \\
 & \left. - \frac{(729829+50400\pi^2)\nu^3}{8400\tilde{r}^7} + \frac{(9544-1341\pi^2)\nu^2}{384\tilde{r}^7} + \frac{315\nu}{16\tilde{r}^7} \right\} \\
 & + (1 \leftrightarrow 2), \tag{5.8} \\
 \tilde{\mathcal{H}}_{\text{N}^3\text{LO}}^{\text{ES}^2} = & \left(C_{\text{ES}^2}^{(0)} \right)_{(1)} \left(\tilde{\mathbf{S}}_{(1)} \cdot \tilde{\mathbf{S}}_{(1)} \right) \\
 & \times \left\{ \tilde{L}^6 \left(-\frac{15\nu^4}{16\tilde{r}^9} + \frac{\nu^3}{\tilde{r}^9} - \frac{3\nu^2}{16\tilde{r}^9} \right) + \tilde{L}^4 \left(\tilde{p}_r^2 \left(\frac{9\nu^3}{4\tilde{r}^7} - \frac{27\nu^4}{8\tilde{r}^7} \right) + \frac{9\nu^4}{16\tilde{r}^8} - \frac{69\nu^3}{16\tilde{r}^8} - \frac{221\nu^2}{16\tilde{r}^8} \right) \right. \\
 & + \tilde{L}^2 \left(\tilde{p}_r^2 \left(\frac{9\nu^4}{8\tilde{r}^6} + \frac{259\nu^3}{4\tilde{r}^6} + \frac{76\nu^2}{\tilde{r}^6} \right) + \tilde{p}_r^4 \left(-\frac{3\nu^4}{\tilde{r}^5} - \frac{9\nu^3}{4\tilde{r}^5} + \frac{9\nu^2}{16\tilde{r}^5} \right) + \frac{447\nu^3}{56\tilde{r}^7} + \frac{125787\nu^2}{4900\tilde{r}^7} \right) \\
 & + \tilde{p}_r^2 \left(-\frac{123\nu^3}{14\tilde{r}^5} - \frac{192173\nu^2}{4900\tilde{r}^5} \right) + \tilde{p}_r^4 \left(-\frac{63\nu^4}{16\tilde{r}^4} - \frac{755\nu^3}{16\tilde{r}^4} - \frac{99\nu^2}{16\tilde{r}^4} \right) \\
 & + \tilde{p}_r^6 \left(\frac{6\nu^4}{\tilde{r}^3} - \frac{7\nu^3}{2\tilde{r}^3} + \frac{3\nu^2}{8\tilde{r}^3} \right) - \frac{69\nu^3}{28\tilde{r}^6} - \frac{68387\nu^2}{4900\tilde{r}^6} \\
 & + \frac{1}{q} \left(\tilde{L}^6 \left(-\frac{35\nu^4}{32\tilde{r}^9} - \frac{53\nu^3}{32\tilde{r}^9} + \frac{31\nu^2}{16\tilde{r}^9} - \frac{13\nu}{32\tilde{r}^9} \right) \right. \\
 & + \tilde{L}^4 \left(\tilde{p}_r^2 \left(-\frac{33\nu^4}{8\tilde{r}^7} - \frac{207\nu^3}{32\tilde{r}^7} + \frac{99\nu^2}{16\tilde{r}^7} - \frac{21\nu}{32\tilde{r}^7} \right) + \frac{9\nu^4}{16\tilde{r}^8} + \frac{27\nu^3}{8\tilde{r}^8} + \frac{53\nu^2}{16\tilde{r}^8} - \frac{9\nu}{2\tilde{r}^8} \right) \\
 & + \tilde{L}^2 \left(\tilde{p}_r^2 \left(\frac{9\nu^4}{8\tilde{r}^6} + \frac{365\nu^3}{4\tilde{r}^6} + \frac{1701\nu^2}{16\tilde{r}^6} - \frac{3\nu}{\tilde{r}^6} \right) + \tilde{p}_r^4 \left(-\frac{9\nu^4}{2\tilde{r}^5} - \frac{75\nu^3}{8\tilde{r}^5} + \frac{45\nu^2}{16\tilde{r}^5} - \frac{3\nu}{32\tilde{r}^5} \right) \right. \\
 & + \frac{499\nu^3}{112\tilde{r}^7} + \frac{(177536-2547\pi^2)\nu^2}{6144\tilde{r}^7} - \frac{205\nu}{8\tilde{r}^7} \left. \right) \\
 & + \tilde{p}_r^2 \left(-\frac{457\nu^3}{28\tilde{r}^5} + \frac{(2547\pi^2-273728)\nu^2}{1536\tilde{r}^5} + \frac{39\nu}{8\tilde{r}^5} \right) \\
 & + \tilde{p}_r^4 \left(-\frac{63\nu^4}{16\tilde{r}^4} - \frac{257\nu^3}{8\tilde{r}^4} - \frac{81\nu^2}{8\tilde{r}^4} + \frac{3\nu}{2\tilde{r}^4} \right) + \tilde{p}_r^6 \left(\frac{4\nu^4}{\tilde{r}^3} + \frac{2\nu^3}{\tilde{r}^3} - \frac{23\nu^2}{16\tilde{r}^3} + \frac{5\nu}{32\tilde{r}^3} \right) \\
 & \left. - \frac{69\nu^3}{28\tilde{r}^6} - \frac{(2888+387\pi^2)\nu^2}{192\tilde{r}^6} + \frac{21\nu}{2\tilde{r}^6} \right\} \\
 & + \left(C_{\text{ES}^2}^{(0)} \right)_{(1)} \left((\tilde{\mathbf{S}}_{(1)} \cdot \tilde{\mathbf{r}})^2 \right) \\
 & \times \left\{ \tilde{L}^4 \left(\tilde{p}_r^2 \left(-\frac{9\nu^4}{16\tilde{r}^9} - \frac{11\nu^3}{16\tilde{r}^9} + \frac{3\nu^2}{4\tilde{r}^9} \right) + \frac{15\nu^4}{16\tilde{r}^{10}} + \frac{311\nu^3}{16\tilde{r}^{10}} + \frac{33\nu^2}{16\tilde{r}^{10}} \right) \right\}
 \end{aligned}$$

$$\begin{aligned}
 & +\widetilde{L}^2 \left(\widetilde{p}_r^2 \left(\frac{35\nu^4}{8\widetilde{r}^8} + \frac{107\nu^3}{4\widetilde{r}^8} - \frac{37\nu^2}{2\widetilde{r}^8} \right) + \widetilde{p}_r^4 \left(-\frac{45\nu^4}{16\widetilde{r}^7} + \frac{17\nu^3}{4\widetilde{r}^7} - \frac{21\nu^2}{8\widetilde{r}^7} \right) - \frac{1063\nu^3}{56\widetilde{r}^9} - \frac{3069\nu^2}{49\widetilde{r}^9} \right) \\
 & +\widetilde{p}_r^2 \left(-\frac{1535\nu^3}{14\widetilde{r}^7} - \frac{19671\nu^2}{4900\widetilde{r}^7} \right) + \widetilde{p}_r^4 \left(\frac{203\nu^4}{16\widetilde{r}^6} + \frac{263\nu^3}{16\widetilde{r}^6} - \frac{457\nu^2}{16\widetilde{r}^6} \right) \\
 & +\widetilde{p}_r^6 \left(-\frac{6\nu^4}{\widetilde{r}^5} + \frac{23\nu^3}{2\widetilde{r}^5} - \frac{27\nu^2}{8\widetilde{r}^5} \right) + \frac{111851\nu^2}{4900\widetilde{r}^8} - \frac{101\nu^3}{28\widetilde{r}^8} \\
 & +\frac{1}{q} \left(\widetilde{L}^6 \left(\frac{15\nu^4}{32\widetilde{r}^{11}} + \frac{105\nu^3}{32\widetilde{r}^{11}} - \frac{51\nu^2}{16\widetilde{r}^{11}} + \frac{21\nu}{32\widetilde{r}^{11}} \right) \right. \\
 & +\widetilde{L}^4 \left(\widetilde{p}_r^2 \left(\frac{21\nu^4}{16\widetilde{r}^9} + \frac{319\nu^3}{32\widetilde{r}^9} - \frac{73\nu^2}{8\widetilde{r}^9} + \frac{39\nu}{32\widetilde{r}^9} \right) + \frac{15\nu^4}{16\widetilde{r}^{10}} + \frac{10\nu^3}{\widetilde{r}^{10}} - \frac{129\nu^2}{8\widetilde{r}^{10}} + \frac{15\nu}{2\widetilde{r}^{10}} \right) \\
 & +\widetilde{L}^2 \left(\widetilde{p}_r^2 \left(\frac{35\nu^4}{8\widetilde{r}^8} - \frac{163\nu^3}{16\widetilde{r}^8} - \frac{1931\nu^2}{16\widetilde{r}^8} + \frac{6\nu}{\widetilde{r}^8} \right) + \widetilde{p}_r^4 \left(\frac{3\nu^4}{8\widetilde{r}^7} + \frac{181\nu^3}{16\widetilde{r}^7} - \frac{73\nu^2}{16\widetilde{r}^7} + \frac{3\nu}{32\widetilde{r}^7} \right) \right. \\
 & \left. - \frac{1725\nu^3}{112\widetilde{r}^9} + \frac{(12735\pi^2 - 929536)\nu^2}{6144\widetilde{r}^9} + \frac{369\nu}{8\widetilde{r}^9} \right) \\
 & +\widetilde{p}_r^2 \left(-\frac{2493\nu^3}{28\widetilde{r}^7} - \frac{3(849\pi^2 - 59072)\nu^2}{512\widetilde{r}^7} - \frac{117\nu}{8\widetilde{r}^7} \right) \\
 & +\widetilde{p}_r^4 \left(\frac{203\nu^4}{16\widetilde{r}^6} - \frac{44\nu^3}{\widetilde{r}^6} + \frac{7\nu^2}{8\widetilde{r}^6} - \frac{9\nu}{2\widetilde{r}^6} \right) + \widetilde{p}_r^6 \left(-\frac{\nu^3}{\widetilde{r}^5} + \frac{37\nu^2}{16\widetilde{r}^5} - \frac{15\nu}{32\widetilde{r}^5} \right) \\
 & \left. - \frac{101\nu^3}{28\widetilde{r}^8} + \frac{(296 + 387\pi^2)\nu^2}{64\widetilde{r}^8} - \frac{63\nu}{2\widetilde{r}^8} \right) \Big\} \\
 & + \left(C_{\text{ES}^2}^{(0)} \right)_{(1)} \left((\widetilde{\mathbf{S}}_{(1)} \cdot \widetilde{\mathbf{L}})^2 \right) \\
 & \times \left\{ \widetilde{L}^4 \left(\frac{15\nu^4}{16\widetilde{r}^9} - \frac{7\nu^3}{2\widetilde{r}^9} + \frac{27\nu^2}{16\widetilde{r}^9} \right) + \widetilde{L}^2 \left(\widetilde{p}_r^2 \left(\frac{63\nu^4}{16\widetilde{r}^7} - \frac{97\nu^3}{16\widetilde{r}^7} - \frac{3\nu^2}{4\widetilde{r}^7} \right) - \frac{3\nu^4}{4\widetilde{r}^8} + \frac{47\nu^3}{8\widetilde{r}^8} + \frac{75\nu^2}{4\widetilde{r}^8} \right) \right. \\
 & +\widetilde{p}_r^2 \left(-\frac{2\nu^4}{\widetilde{r}^6} - \frac{351\nu^3}{4\widetilde{r}^6} - \frac{245\nu^2}{4\widetilde{r}^6} \right) + \widetilde{p}_r^4 \left(\frac{93\nu^4}{16\widetilde{r}^5} + \frac{4\nu^3}{\widetilde{r}^5} - \frac{39\nu^2}{16\widetilde{r}^5} \right) + \frac{35099\nu^2}{4900\widetilde{r}^7} - \frac{73\nu^3}{14\widetilde{r}^7} \\
 & +\frac{1}{q} \left(\widetilde{L}^4 \left(\frac{15\nu^4}{16\widetilde{r}^9} - \frac{\nu^3}{16\widetilde{r}^9} - \frac{13\nu^2}{8\widetilde{r}^9} + \frac{9\nu}{16\widetilde{r}^9} \right) \right. \\
 & +\widetilde{L}^2 \left(\widetilde{p}_r^2 \left(\frac{69\nu^4}{16\widetilde{r}^7} + \frac{115\nu^3}{16\widetilde{r}^7} - \frac{151\nu^2}{16\widetilde{r}^7} + \frac{3\nu}{4\widetilde{r}^7} \right) - \frac{3\nu^4}{4\widetilde{r}^8} - \frac{7\nu^3}{2\widetilde{r}^8} + \frac{107\nu^2}{16\widetilde{r}^8} + \frac{6\nu}{\widetilde{r}^8} \right) \\
 & +\widetilde{p}_r^2 \left(-\frac{2\nu^4}{\widetilde{r}^6} - \frac{1925\nu^3}{16\widetilde{r}^6} - \frac{621\nu^2}{4\widetilde{r}^6} + \frac{3\nu}{\widetilde{r}^6} \right) + \widetilde{p}_r^4 \left(\frac{57\nu^4}{8\widetilde{r}^5} + \frac{341\nu^3}{16\widetilde{r}^5} - \frac{55\nu^2}{8\widetilde{r}^5} + \frac{3\nu}{16\widetilde{r}^5} \right) \\
 & \left. - \frac{17\nu^3}{14\widetilde{r}^7} + \frac{(247616 - 2547\pi^2)\nu^2}{3072\widetilde{r}^7} + \frac{123\nu}{4\widetilde{r}^7} \right) \Big\} \\
 & + \left(C_{\text{ES}^2}^{(0)} \right)_{(1)} \left((\widetilde{\mathbf{S}}_{(1)} \cdot \widetilde{\mathbf{F}})(\widetilde{\mathbf{S}}_{(1)} \cdot \widetilde{\mathbf{L}}) \right) \\
 & \times \widetilde{p}_r \left\{ \widetilde{L}^4 \left(\frac{3\nu^4}{8\widetilde{r}^9} - \frac{67\nu^3}{16\widetilde{r}^9} + \frac{39\nu^2}{16\widetilde{r}^9} \right) + \widetilde{L}^2 \left(\left(-\frac{29\nu^4}{16\widetilde{r}^8} + \frac{323\nu^3}{16\widetilde{r}^8} + \frac{16\nu^2}{\widetilde{r}^8} \right) + \widetilde{p}_r^2 \left(\frac{9\nu^4}{8\widetilde{r}^7} - \frac{29\nu^3}{16\widetilde{r}^7} - \frac{27\nu^2}{8\widetilde{r}^7} \right) \right) \right\}
 \end{aligned}$$

$$\begin{aligned}
& +\tilde{p}_r^2 \left(-\frac{93\nu^4}{16\tilde{r}^6} + \frac{1611\nu^3}{16\tilde{r}^6} - \frac{72\nu^2}{\tilde{r}^6} \right) + \tilde{p}_r^4 \left(-\frac{3\nu^4}{16\tilde{r}^5} + \frac{31\nu^3}{2\tilde{r}^5} - \frac{93\nu^2}{16\tilde{r}^5} \right) - \frac{65\nu^3}{4\tilde{r}^7} - \frac{250693\nu^2}{1225\tilde{r}^7} \\
& + \frac{1}{q} \left(\tilde{L}^4 \left(-\frac{9\nu^4}{16\tilde{r}^9} - \frac{37\nu^3}{8\tilde{r}^9} + \frac{41\nu^2}{16\tilde{r}^9} - \frac{3\nu}{16\tilde{r}^9} \right) \right. \\
& + \tilde{L}^2 \left(\left(-\frac{29\nu^4}{16\tilde{r}^8} + \frac{157\nu^3}{4\tilde{r}^8} + \frac{57\nu^2}{16\tilde{r}^8} - \frac{3\nu}{\tilde{r}^8} \right) + \tilde{p}_r^2 \left(-\frac{45\nu^4}{16\tilde{r}^7} - \frac{4\nu^3}{\tilde{r}^7} + \frac{49\nu^2}{16\tilde{r}^7} - \frac{9\nu}{8\tilde{r}^7} \right) \right) \\
& + \tilde{p}_r^2 \left(-\frac{93\nu^4}{16\tilde{r}^6} + \frac{2779\nu^3}{16\tilde{r}^6} + \frac{417\nu^2}{16\tilde{r}^6} - \frac{9\nu}{\tilde{r}^6} \right) + \tilde{p}_r^4 \left(-\frac{63\nu^4}{8\tilde{r}^5} + \frac{145\nu^3}{16\tilde{r}^5} + \frac{19\nu^2}{8\tilde{r}^5} - \frac{15\nu}{16\tilde{r}^5} \right) \\
& \left. - \frac{827\nu^3}{28\tilde{r}^7} + \frac{(2547\pi^2 - 237272)\nu^2}{384\tilde{r}^7} - \frac{30\nu}{\tilde{r}^7} \right\} \\
& + (1 \leftrightarrow 2), \tag{5.9}
\end{aligned}$$

$$\tilde{\mathcal{H}}_{\text{LO}}^{\text{E}^2} = \left(C_{\text{E}^2}^{(2)} \right)_{(1)} (\tilde{\mathbf{S}}_{(1)} \cdot \tilde{\mathbf{S}}_{(1)}) \left\{ -\frac{3\nu^3}{\tilde{r}^6} - \frac{1}{q} \frac{3\nu^3}{\tilde{r}^6} \right\} + (1 \leftrightarrow 2), \tag{5.10}$$

$$\begin{aligned}
\tilde{\mathcal{H}}_{\text{LO}}^{\text{E}^2\text{S}^2} &= \left(C_{\text{E}^2\text{S}^2}^{(0)} \right)_{(1)} (\tilde{\mathbf{S}}_{(1)} \cdot \tilde{\mathbf{S}}_{(1)}) \left\{ -\frac{\nu^3}{2\tilde{r}^6} - \frac{1}{q} \frac{\nu^3}{2\tilde{r}^6} \right\} \\
&+ \left(C_{\text{E}^2\text{S}^2}^{(0)} \right)_{(1)} ((\tilde{\mathbf{S}}_{(1)} \cdot \tilde{\mathbf{r}})^2) \left\{ \frac{3\nu^3}{2\tilde{r}^8} + \frac{1}{q} \frac{3\nu^3}{2\tilde{r}^8} \right\} + (1 \leftrightarrow 2). \tag{5.11}
\end{aligned}$$

The novel computation of the $\tilde{\mathcal{H}}_{\text{N}^3\text{LO}}^{\text{SS}}$ along with the previously known lower order pieces of the Hamiltonian as well as the non-spinning and the spin-orbit Hamiltonian permits us to obtain the total quadratic-in-spin N³LO Hamiltonian adopting the EFT formalism. The known lower order Hamiltonians in the non-spinning and spin-orbit sectors are reported in the appendix C of [65], where as the quadratic-in-spin sector lower order Hamiltonians are reported in the appendix B. The total Hamiltonian in the generic frame is also provided in the supplementary material as **Hamiltonian.m**. Recently, the total Hamiltonian at N³LO has been also reported in [66]. However, as the Hamiltonians are gauge dependent quantities, we perform a comparison of the results of gauge-independent observable in the later part of the article.

6 Computation of observables with spin

The derived generic Hamiltonian is still gauge dependent, because of it's dependence on the radial coordinate. So, we can compute observable, which are gauge invariant and comparable with other results in the literature. In this section, we focus on the computation of two gauge invariant observable, namely, the binding energy, and the scattering angle.

For this purpose, we adopt the COM frame, where $\mathbf{p}_{(1)} + \mathbf{p}_{(2)} = 0$, as described in section 5. We also assume the aligned spin configurations, which implies that the spins are aligned to the direction of the orbital angular momentum of the compact binary. Such aligned spin configuration is realized by

$$\mathbf{S}_{(a)} \cdot \mathbf{r} = \mathbf{S}_{(a)} \cdot \mathbf{p} = 0 \implies \mathbf{S}_{(a)} \cdot (\mathbf{r} \times \mathbf{p}) = S_{(a)}L, \tag{6.1}$$

with, $L = |\mathbf{L}|$ and $S_{(a)} = |\mathbf{S}_{(a)}|$.

6.1 Binding energy for circular orbits with aligned spins

The gauge invariant relation between the binding energy and the orbital frequency for circular orbits is obtained by eliminating the dependence on the radial coordinate. For circular orbits we have

$$\frac{\partial \tilde{\mathcal{H}}(\tilde{r}, \tilde{L}, \tilde{S}_{(a)})}{\partial \tilde{r}} = 0. \quad (6.2)$$

We invert the above relation to express \tilde{r} as a function of \tilde{L} . Then we substitute \tilde{L} as a function of $\tilde{\omega}$, the orbital frequency defined as

$$\tilde{\omega} = \frac{\partial \tilde{\mathcal{H}}(\tilde{L}, \tilde{S}_{(a)})}{\partial \tilde{L}}. \quad (6.3)$$

Additionally, we define a gauge invariant PN parameter $x = \tilde{\omega}^{2/3}$. Following the above procedure with the Hamiltonian given in section 5 we obtain

$$E(x, \tilde{S}_{(a)}) = E_{\text{pp}}(x) + E_{\text{SO}}(x, \tilde{S}_{(a)}) + E_{\text{SS}}(x, \tilde{S}_{(a)}), \quad (6.4)$$

where E_{pp} and E_{SO} are reported in [65], and

$$E_{\text{SS}}(x, \tilde{S}_{(a)}) = E_{\text{S1S2}}(x, \tilde{S}_{(a)}) + E_{\text{S}^2}(x, \tilde{S}_{(a)}) \\ + E_{\text{ES}^2}(x, \tilde{S}_{(a)}) + E_{\text{E}^2\text{S}^2}(x, \tilde{S}_{(a)}) + E_{\text{E}^2}(x, \tilde{S}_{(a)}). \quad (6.5)$$

The individual terms in the above equation are given as,

$$E_{\text{S1S2}}(x, \tilde{S}) = \tilde{S}_{(1)}\tilde{S}_{(2)} \left\{ x^3 \{\nu\} + x^4 \left\{ \frac{5}{6}\nu + \frac{5}{18}\nu^2 \right\} + x^5 \left\{ \frac{35}{8}\nu - \frac{1001}{72}\nu^2 - \frac{371}{216}\nu^3 \right\} \right. \\ \left. + x^6 \left\{ \frac{243}{16}\nu - \left(\frac{2107}{16} - \frac{123}{32}\pi^2 \right) \nu^2 + \frac{147}{8}\nu^3 + \frac{13}{16}\nu^4 \right\} \right\}, \quad (6.6)$$

$$E_{\text{S}^2}(x, \tilde{S}) = \tilde{S}_{(1)}^2 \left\{ x^4 \left\{ \frac{25}{18}\nu^2 + \frac{1}{q} \left(-\frac{5}{2}\nu + \frac{5}{6}\nu^2 \right) \right\} \right. \\ + x^5 \left\{ \frac{10}{3}\nu^2 - \frac{749}{108}\nu^3 + \frac{1}{q} \left(-\frac{21}{4}\nu - \frac{7}{6}\nu^2 - \frac{217}{36}\nu^3 \right) \right\} \\ + x^6 \left\{ \frac{1947}{112}\nu^2 - \frac{48357}{560}\nu^3 + \frac{159}{16}\nu^4 \right. \\ \left. + \frac{1}{q} \left(-\frac{243}{16}\nu + \left(\frac{747}{16} - \frac{189\pi^2}{2048} \right) \nu^2 - \frac{13731}{280}\nu^3 + \frac{153}{16}\nu^4 \right) \right\} \\ + (1 \leftrightarrow 2), \quad (6.7)$$

$$E_{\text{ES}^2}(x, \tilde{S}) = \left(C_{\text{ES}^2}^{(0)} \right)_{(1)} \tilde{S}_{(1)}^2 \left\{ x^3 \left\{ \frac{1}{q} \frac{\nu}{2} \right\} + x^4 \left\{ \frac{5}{3}\nu^2 + \frac{1}{q} \left(\frac{5}{4}\nu + \frac{5}{4}\nu^2 \right) \right\} \right. \\ + x^5 \left\{ \frac{31}{4}\nu^2 - \frac{35}{18}\nu^3 + \frac{1}{q} \left(\frac{63}{16}\nu + \frac{77}{48}\nu^2 - \frac{91}{48}\nu^3 \right) \right\} \\ + x^6 \left\{ \frac{789}{28}\nu^2 - \frac{156}{7}\nu^3 + \frac{5}{8}\nu^4 \right. \\ \left. + \frac{1}{q} \left(\frac{405}{32}\nu + \left(\frac{3747\pi^2}{2048} - \frac{2389}{32} \right) \nu^2 - \frac{555}{56}\nu^3 + \frac{21}{32}\nu^4 \right) \right\} \\ + (1 \leftrightarrow 2), \quad (6.8)$$

$$E_{\text{E}^2}(x, \tilde{S}) = \left(C_{\text{E}^2}^{(2)} \right)_{(1)} \tilde{S}_{(1)}^2 x^6 \left\{ 9\nu^3 \left(1 + \frac{1}{q} \right) \right\} + (1 \leftrightarrow 2), \quad (6.9)$$

$$E_{\text{E}^2\text{S}^2}(x, \tilde{S}) = \left(C_{\text{E}^2\text{S}^2}^{(0)} \right)_{(1)} \tilde{S}_{(1)}^2 x^6 \left\{ \frac{3\nu^3}{2} \left(1 + \frac{1}{q} \right) \right\} + (1 \leftrightarrow 2). \quad (6.10)$$

The equation (6.6) agrees with previously known classical results of [85], and the equations (6.6), (6.7), (6.8), (6.9) and (6.10) agree⁶ with recent results given in [39, 66] derived using similar EFT techniques as presented in this paper.

6.2 Scattering angle with aligned spins

In this section, we compute the scattering angle considering an aligned spin binary system following [45]. First, we re-scale the spin variables as $a_{(a)} = S_{(a)}/m_{(a)}$. In the COM, the Hamiltonian \mathcal{H} is expressed as a function of \tilde{p}_r , L , r , and $S(a)$ and inverting that we obtain $\tilde{p}_r(\mathcal{H}, L, r, S_{(a)})$. Then the scattering angle χ is given by

$$\chi(\mathcal{H}, L, S_{(a)}) = - \int dr \frac{\partial \tilde{p}_r(\mathcal{H}, L, r, S_{(a)})}{\partial L} - \pi. \quad (6.11)$$

We invert the relation between the Lorentz factor γ and the total energy per total rest mass $\Gamma = \mathcal{H}/(Mc^2)$ given by

$$\gamma = \frac{1}{\sqrt{1 - v^2/c^2}} = 1 + \frac{\Gamma^2 - 1}{2\nu}, \quad (6.12)$$

where, $v \equiv |\dot{\mathbf{r}}|$ is the relative velocity of the compact objects, and the relation between the total angular momentum L and the impact parameter b given by

$$L = \frac{\mu\gamma vb}{\Gamma} + Mc \left(\frac{\Gamma - 1}{2} \right) \left(a_+ - \frac{\delta}{\Gamma} a_- \right), \quad (6.13)$$

where, $a_{(+)} = a_{(1)} + a_{(2)}$ and $a_{(-)} = a_{(1)} - a_{(2)}$. With the above inversions, we trade H for v and L for b . This allows us to express the scattering angle as

$$\chi(v, b, S_{(a)}) = - \frac{\gamma}{\mu\gamma v} \int dr \frac{\partial \tilde{p}_r(v, b, r, S_{(a)})}{\partial b} - \pi. \quad (6.14)$$

Now, applying the above procedure with the Hamiltonian given in section 5, we obtain the scattering angle computed in the COM for aligned spins, which can be expressed as

$$\chi(v, b, S_{(a)}) = \chi_{\text{pp}}(v, b) + \chi_{\text{SO}}(v, b, S_{(a)}) + \chi_{\text{SS}}(v, b, S_{(a)}), \quad (6.15)$$

where χ_{pp} and χ_{SO} are reported in [65], and

$$\begin{aligned} \chi_{\text{SS}}(v, b, S_{(a)}) &= \chi_{\text{S1S2}}(v, b, S_{(a)}) + \chi_{\text{S}^2}(v, b, S_{(a)}) \\ &+ \chi_{\text{ES}^2}(v, b, S_{(a)}) + \chi_{\text{E}^2\text{S}^2}(v, b, S_{(a)}) + \chi_{\text{E}^2}(v, b, S_{(a)}). \end{aligned} \quad (6.16)$$

⁶We point out that our expression of binding energy contains the expanded Wilson coefficients, which only depend on the invariant m and the corresponding results in [39, 66] are provided in terms of Wilson coefficients depending on two invariants, namely m and S^2 .

The individual terms in the above equation are given as,

$$\begin{aligned}
 \frac{\chi_{\text{S1S2}} + \chi_{\text{S2}}}{\Gamma} &= \frac{v^2}{b^2 c^2} \left[a_{(+)}^2 \quad \delta a_{(+)} a_{(-)} \quad a_{(-)}^2 \right] \cdot \left(\left(\frac{G_{\text{NM}}}{v^2 b} \right) \left\{ \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix} + \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix} \left(\frac{v^2}{c^2} \right) \right\} \right. \\
 &+ \pi \left(\frac{G_{\text{NM}}}{v^2 b} \right)^2 \left\{ \frac{3}{4} \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix} + \frac{3}{16} \begin{bmatrix} 41 \\ 10 \\ -19 \end{bmatrix} \left(\frac{v^2}{c^2} \right) + \frac{3}{128} \begin{bmatrix} 55 \\ -10 \\ -41 \end{bmatrix} \left(\frac{v^4}{c^4} \right) \right\} \\
 &+ \left(\frac{G_{\text{NM}}}{v^2 b} \right)^3 \left\{ 2 \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix} + \begin{bmatrix} 102 - 2\nu \\ 32 \\ -38 - 6\nu \end{bmatrix} \left(\frac{v^2}{c^2} \right) + \frac{1}{14} \begin{bmatrix} 2332 - 499\nu \\ 688 \\ -748 - 37\nu \end{bmatrix} \left(\frac{v^4}{c^4} \right) \right. \\
 &+ \left. \frac{1}{140} \begin{bmatrix} 1704 - 9925\nu \\ -288 \\ -1096 + 5377\nu \end{bmatrix} \left(\frac{v^6}{c^6} \right) \right\} \\
 &+ \pi \left(\frac{G_{\text{NM}}}{v^2 b} \right)^4 \left\{ \frac{15}{16} \begin{bmatrix} 63 - 2\nu \\ 22 \\ -21 - 6\nu \end{bmatrix} \left(\frac{v^2}{c^2} \right) - \frac{15}{448} \begin{bmatrix} -11063 + 2638\nu \\ -4206 + 308\nu \\ 2657 + 790\nu \end{bmatrix} \left(\frac{v^4}{c^4} \right) \right. \\
 &- \left. \frac{3}{229376} \begin{bmatrix} 19102720 + (-29293696 + 135555\pi^2)\nu \\ -256(-24640 + 11507\nu) \\ -4730880 + (8911744 - 139965\pi^2)\nu \end{bmatrix} \left(\frac{v^6}{c^6} \right) \right\} \\
 &+ \mathcal{O} \left(G_{\text{N}}^5, \frac{v^8}{c^8} \right), \tag{6.17}
 \end{aligned}$$

$$\begin{aligned}
 \frac{\chi_{\text{ES2}}}{\Gamma} &= \frac{v^2}{b^2 c^2} \left[a_{\text{ES2}(+)}^2 \quad \delta a_{\text{ES2}(-)}^2 \right] \cdot \left\{ \left(\frac{G_{\text{NM}}}{v^2 b} \right) \left\{ \begin{bmatrix} 2 \\ 0 \end{bmatrix} + \begin{bmatrix} 2 \\ 0 \end{bmatrix} \left(\frac{v^2}{c^2} \right) \right\} \right. \\
 &+ \pi \left(\frac{G_{\text{NM}}}{v^2 b} \right)^2 \left\{ \frac{1}{2} \begin{bmatrix} 3 \\ 0 \end{bmatrix} + \frac{1}{4} \begin{bmatrix} 27 \\ 3 \end{bmatrix} \left(\frac{v^2}{c^2} \right) + \frac{1}{64} \begin{bmatrix} 117 \\ 57 \end{bmatrix} \left(\frac{v^4}{c^4} \right) \right\} \\
 &+ \left(\frac{G_{\text{NM}}}{v^2 b} \right)^3 \left\{ \begin{bmatrix} 4 \\ 0 \end{bmatrix} + \begin{bmatrix} 76 - 4\nu \\ 8 \end{bmatrix} \left(\frac{v^2}{c^2} \right) + \frac{1}{7} \begin{bmatrix} 748 - 131\nu \\ 216 \end{bmatrix} \left(\frac{v^4}{c^4} \right) + \frac{1}{70} \begin{bmatrix} 1096 - 5013\nu \\ 704 \end{bmatrix} \left(\frac{v^6}{c^6} \right) \right\} \\
 &+ \pi \left(\frac{G_{\text{NM}}}{v^2 b} \right)^4 \left\{ \frac{15}{4} \begin{bmatrix} 11 - \nu \\ 1 \end{bmatrix} \left(\frac{v^2}{c^2} \right) + \frac{15}{224} \begin{bmatrix} 2867 - 708\nu \\ 627 - 28\nu \end{bmatrix} \left(\frac{v^4}{c^4} \right) \right. \\
 &+ \left. \frac{15}{114688} \begin{bmatrix} 1026816 - (2184832 + 27111\pi^2)\nu \\ 128(3262 - 395\nu) \end{bmatrix} \left(\frac{v^6}{c^6} \right) \right\} + \mathcal{O} \left(G_{\text{N}}^5, \frac{v^8}{c^8} \right), \tag{6.18}
 \end{aligned}$$

$$\frac{\chi_{\text{E2}}}{\Gamma} = \frac{v^2}{b^2 c^2} \left[a_{\text{E2}(+)}^2 \quad \delta a_{\text{E2}(-)}^2 \right] \cdot \left\{ \pi \left(\frac{G_{\text{NM}}}{v^2 b} \right)^4 \frac{45}{16} \nu \begin{bmatrix} 1 \\ 1 \end{bmatrix} \left(\frac{v^6}{c^6} \right) \right\} + \mathcal{O} \left(G_{\text{N}}^5, \frac{v^8}{c^8} \right), \tag{6.19}$$

$$\frac{\chi_{\text{E2S2}}}{\Gamma} = \frac{v^2}{b^2 c^2} \left[a_{\text{E2S2}(+)}^2 \quad \delta a_{\text{E2S2}(-)}^2 \right] \cdot \left\{ \pi \left(\frac{G_{\text{NM}}}{v^2 b} \right)^4 \frac{15}{32} \nu \begin{bmatrix} 1 \\ 1 \end{bmatrix} \left(\frac{v^6}{c^6} \right) \right\} + \mathcal{O} \left(G_{\text{N}}^5, \frac{v^8}{c^8} \right), \tag{6.20}$$

where, $a_{\text{ES2}(\pm)}^2 = \left(C_{\text{ES2}}^{(0)} \right)_{(1)} a_{(1)}^2 \pm \left(C_{\text{ES2}}^{(0)} \right)_{(2)} a_{(2)}^2$ and similarly for $a_{\text{E2S2}(\pm)}^2$ and $a_{\text{E2}(\pm)}^2$. The equation (6.17) agrees with previously known classical results of [85], and the equa-

tions (6.17), (6.18) and (6.20) agree with recent results given in [66] derived using similar EFT techniques as presented in this paper. Note that the scattering angle depends on the same combination of $C_{E^2}^{(2)}$ and $C_{E^2S^2}^{(0)}$ as the binding energy. Matching both constants independently hence requires another observable in a more asymmetric (e.g. precessing) kinematic regime.

7 Conclusion

In this work, we presented the complete evaluation of N³LO Post-Newtonian correction to the quadratic-in-spin Hamiltonian for the spinning compact binaries, within the EFT Feynman diagrammatic approach of GR. Together with our earlier study on the spin-orbit Hamiltonian [65], this is at the current state-of-the-art result for the conservative part of the spinning sectors up to quadratic-in-spin.

The necessary Feynman diagrams in momentum space were generated using Feynman rules derived from the EFT Lagrangian describing the spinning compact objects. The corresponding Feynman amplitudes were written as a linear combination of master integrals, employing the integration-by-parts identities for dimensionally regularized integrals. The contribution of each diagram to the effective potential was obtained upon Fourier transform, and Laurent series expanded around $d = 3 + \epsilon$ space dimensions. Finally, the Hamiltonian was derived by applying Legendre transform, and canonical transformations were employed to remove the non-physical divergences and spurious logarithmic terms. Furthermore, we computed two specific gauge invariant observable, namely, the binding energy with aligned spins in the case of circular orbit, and the scattering angle for the case of aligned spins. These results were found in agreement with the corresponding expressions available in literature, previously obtained using the self-force formalism as well as the EFT framework. Additionally, the obtained results in the article, particularly the scattering angle, could be used with the recently developed boundary-to-bound maps [86–88] to compute other observables like periastron advance.

The obtained results depend on three Wilson coefficients, namely, $C_{E^2}^{(2)}$, $C_{ES^2}^{(0)}$, and $C_{E^2S^2}^{(0)}$. They arise from the non-minimal couplings at linear and quadratic order in curvature in the EFT Lagrangian of the point particles. The coefficient $C_{ES^2}^{(0)}$ starts contributing from 2PN (LO S^2 sector), whereas the other two coefficients $C_{E^2}^{(2)}$ and $C_{E^2S^2}^{(0)}$ contribute for the first time at 5PN. Out of them, $C_{ES^2}^{(0)}$ is related to the spin-induced quadrupole moment of the compact object: for Kerr black holes, it is known to be 1 [89, 90]; and for neutron stars its value ranges within 2–8 [91]. The other two coefficients, $C_{E^2}^{(2)}$ and $C_{E^2S^2}^{(0)}$, encode quadrupolar deformations, due to an external field and spin-square effects. Their determination requires the matching against stationary linear perturbations of spinning compact objects — see [92–95] for crucial work in that direction.

The observable presented in this work could also be used to match a combination of $C_{E^2}^{(2)}$ and $C_{E^2S^2}^{(0)}$ to the motion of a small body within the self-force formalism. The dependence of gravitational waves from compact binaries on $C_{E^2}^{(2)}$ and $C_{E^2S^2}^{(0)}$ provides a probe of the nature of black holes and encodes information about the equation of state of neutron stars.

Acknowledgments

The work of M.K.M is supported by Fellini — Fellowship for Innovation at INFN funded by the European Union’s Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No. 754496. RP is grateful to IISER Bhopal for the fellowship. RP’s research is funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation), Projektnummer 417533893/GRK2575 “Rethinking Quantum Field Theory”.

A Notation and convention

Spacetime metric	$\eta_{\mu\nu} = (1, -1, -1, -1)$	(A.1a)
4 dimensional indices	μ, ν	(A.1b)
3 dimensional indices	i, j	(A.1c)
Compact object label	(a) where $a = \{1, 2\}$	(A.1d)
Time derivative	·	(A.1e)
Position of a^{th} object	$\mathbf{x}_{(a)}$	(A.1f)
Velocity of a^{th} object	$\mathbf{v}_{(a)} \equiv \dot{\mathbf{x}}_{(a)}$	(A.1g)
Acceleration of a^{th} object	$\mathbf{a}_{(a)} \equiv \ddot{\mathbf{x}}_{(a)}$	(A.1h)
Separation vector for binary	$\mathbf{r} \equiv \mathbf{x}_{(a)} - \mathbf{x}_{(a')}$	(A.1i)
Separation distance for binary	$r \equiv \mathbf{r} $	(A.1j)
Separation unit vector for binary	$\mathbf{n} \equiv \frac{\mathbf{r}}{r}$	(A.1k)
Angular momentum of the binary	$\mathbf{L} \equiv (\mathbf{r} \times \mathbf{p})$	(A.1l)
Spin vector of a^{th} object	$\mathbf{S}_{(a)}^i \equiv \epsilon^{ijk} \mathbf{S}_{(a)}^{jk}$	(A.1m)
	$\int_{\mathbf{p}} \int \frac{d^d p}{(2\pi)^d}$	(A.1n)
	$\int_{\mathbf{p}} \int \frac{d^3 p}{(2\pi)^3}$	(A.1o)
Center of mass coordinates	$\mathbf{p}_{(1)} + \mathbf{p}_{(2)} = 0$	(A.1p)
Circular orbits	$\tilde{p}_r \equiv \mathbf{p} \cdot \mathbf{n} = 0$ and $\dot{p}_r = 0$	(A.1q)
Aligned spins	$\mathbf{S}_{(a)} \cdot \mathbf{r} = \mathbf{S}_{(a)} \cdot \mathbf{p} = 0$	(A.1r)
Exchange of particle label ($1 \leftrightarrow 2$)	$({}_{(1)} \leftrightarrow {}_{(2)}), \left(q \leftrightarrow \frac{1}{q}\right), (\delta \leftrightarrow -\delta).$	(A.1s)

B Lower-order Hamiltonians

In this appendix we give the results for all the lower order Hamiltonians given in eq. (5.6).

B.1 Spin1-spin2 sector up to N²LO

$$\tilde{\mathcal{H}}_{\text{LO}}^{\text{S1S2}} = \left(\tilde{\mathbf{S}}_{(1)} \cdot \tilde{\mathbf{S}}_{(2)} \right) \left\{ -\frac{\nu}{\tilde{r}^3} \right\} + \left(\tilde{\mathbf{S}}_{(1)} \cdot \tilde{\mathbf{r}} \right) \left(\tilde{\mathbf{S}}_{(2)} \cdot \tilde{\mathbf{r}} \right) \left\{ \frac{3\nu}{\tilde{r}^5} \right\} \quad (\text{B.1})$$

$$\begin{aligned} \tilde{\mathcal{H}}_{\text{NLO}}^{\text{S1S2}} = & \left(\tilde{\mathbf{S}}_{(1)} \cdot \tilde{\mathbf{S}}_{(2)} \right) \left\{ \tilde{L}^2 \left(\frac{\nu}{\tilde{r}^5} - \frac{\nu^2}{4\tilde{r}^5} \right) + \tilde{p}_r^2 \left(-\frac{7\nu^2}{4\tilde{r}^3} - \frac{\nu}{2\tilde{r}^3} \right) + \frac{5\nu}{\tilde{r}^4} \right\} \\ & + \left(\tilde{\mathbf{S}}_{(1)} \cdot \tilde{\mathbf{r}} \right) \left(\tilde{\mathbf{S}}_{(2)} \cdot \tilde{\mathbf{r}} \right) \left\{ \frac{3\tilde{L}^2\nu^2}{2\tilde{r}^7} + \tilde{p}_r^2 \left(\frac{23\nu^2}{4\tilde{r}^5} + \frac{\nu}{2\tilde{r}^5} \right) - \frac{11\nu}{\tilde{r}^6} \right\} \\ & + \left(\tilde{\mathbf{S}}_{(1)} \cdot \tilde{\mathbf{L}} \right) \left(\tilde{\mathbf{S}}_{(2)} \cdot \tilde{\mathbf{L}} \right) \left\{ -\frac{\nu^2}{\tilde{r}^5} - \frac{5\nu}{2\tilde{r}^5} \right\} \\ & + \left(\tilde{\mathbf{S}}_{(1)} \cdot \tilde{\mathbf{r}} \right) \left(\tilde{\mathbf{S}}_{(2)} \cdot \tilde{\mathbf{L}} \right) \tilde{p}_r \left\{ \frac{1}{q} \frac{3\nu^2}{2\tilde{r}^5} + \left(-\frac{\nu^2}{4\tilde{r}^5} - \frac{\nu}{\tilde{r}^5} \right) \right\} \\ & + \left(\tilde{\mathbf{S}}_{(1)} \cdot \tilde{\mathbf{L}} \right) \left(\tilde{\mathbf{S}}_{(2)} \cdot \tilde{\mathbf{r}} \right) \tilde{p}_r \left\{ \left(\frac{\nu}{2\tilde{r}^5} - \frac{13\nu^2}{4\tilde{r}^5} \right) - \frac{1}{q} \frac{3\nu^2}{2\tilde{r}^5} \right\} \end{aligned} \quad (\text{B.2})$$

$$\begin{aligned} \tilde{\mathcal{H}}_{\text{N}^2\text{LO}}^{\text{S1S2}} = & \left(\tilde{\mathbf{S}}_{(1)} \cdot \tilde{\mathbf{S}}_{(2)} \right) \left\{ \tilde{L}^4 \left(-\frac{\nu^3}{8\tilde{r}^7} + \frac{9\nu^2}{8\tilde{r}^7} - \frac{3\nu}{4\tilde{r}^7} \right) + \tilde{L}^2 \left(\tilde{p}_r^2 \left(-\frac{5\nu^3}{8\tilde{r}^5} + \frac{237\nu^2}{16\tilde{r}^5} - \frac{3\nu}{8\tilde{r}^5} \right) + \frac{61\nu^2}{16\tilde{r}^6} - \frac{19\nu}{4\tilde{r}^6} \right) \right. \\ & + \tilde{p}_r^2 \left(\frac{75\nu^2}{16\tilde{r}^4} + \frac{9\nu}{4\tilde{r}^4} \right) + \tilde{p}_r^4 \left(-\frac{19\nu^3}{8\tilde{r}^3} - \frac{81\nu^2}{16\tilde{r}^3} + \frac{3\nu}{8\tilde{r}^3} \right) - \frac{31\nu^2}{8\tilde{r}^5} - \frac{27\nu}{2\tilde{r}^5} \left. \right\} \\ & + \left(\tilde{\mathbf{S}}_{(1)} \cdot \tilde{\mathbf{r}} \right) \left(\tilde{\mathbf{S}}_{(2)} \cdot \tilde{\mathbf{r}} \right) \left\{ \tilde{L}^4 \left(\frac{9\nu^3}{8\tilde{r}^9} - \frac{3\nu^2}{4\tilde{r}^9} \right) + \tilde{L}^2 \left(\tilde{p}_r^2 \left(\frac{7\nu^3}{2\tilde{r}^7} - \frac{97\nu^2}{8\tilde{r}^7} - \frac{3\nu}{8\tilde{r}^7} \right) - \frac{17\nu^2}{4\tilde{r}^8} - \frac{9\nu}{4\tilde{r}^8} \right) \right. \\ & + \tilde{p}_r^2 \left(\frac{3\nu}{4\tilde{r}^6} - \frac{723\nu^2}{16\tilde{r}^6} \right) + \tilde{p}_r^4 \left(\frac{67\nu^3}{8\tilde{r}^5} + \frac{49\nu^2}{16\tilde{r}^5} - \frac{3\nu}{8\tilde{r}^5} \right) + \frac{95\nu^2}{8\tilde{r}^7} + \frac{53\nu}{2\tilde{r}^7} \left. \right\} \\ & + \left(\tilde{\mathbf{S}}_{(1)} \cdot \tilde{\mathbf{L}} \right) \left(\tilde{\mathbf{S}}_{(2)} \cdot \tilde{\mathbf{L}} \right) \left\{ \tilde{L}^2 \left(-\frac{\nu^3}{\tilde{r}^7} - \frac{\nu^2}{2\tilde{r}^7} + \frac{15\nu}{8\tilde{r}^7} \right) + \tilde{p}_r^2 \left(-\frac{5\nu^3}{2\tilde{r}^5} - \frac{181\nu^2}{8\tilde{r}^5} + \frac{15\nu}{8\tilde{r}^5} \right) + \frac{5\nu^2}{2\tilde{r}^6} + \frac{27\nu}{2\tilde{r}^6} \right\} \\ & + \left(\tilde{\mathbf{S}}_{(1)} \cdot \tilde{\mathbf{r}} \right) \left(\tilde{\mathbf{S}}_{(2)} \cdot \tilde{\mathbf{L}} \right) \tilde{p}_r \left\{ \tilde{L}^2 \left(-\frac{5\nu^3}{8\tilde{r}^7} - \frac{43\nu^2}{8\tilde{r}^7} + \frac{3\nu}{4\tilde{r}^7} \right) + \left(-\frac{3\nu^3}{2\tilde{r}^6} + \frac{175\nu^2}{8\tilde{r}^6} + \frac{2\nu}{\tilde{r}^6} \right) \right. \\ & + \tilde{p}_r^2 \left(-\frac{\nu^3}{4\tilde{r}^5} - \frac{35\nu^2}{16\tilde{r}^5} + \frac{3\nu}{4\tilde{r}^5} \right) \\ & + \frac{1}{q} \left(\tilde{L}^2 \left(\frac{3\nu^3}{4\tilde{r}^7} - \frac{9\nu^2}{8\tilde{r}^7} \right) + \left(-\frac{3\nu^3}{2\tilde{r}^6} - \frac{7\nu^2}{\tilde{r}^6} \right) + \tilde{p}_r^2 \left(\frac{9\nu^3}{2\tilde{r}^5} - \frac{9\nu^2}{8\tilde{r}^5} \right) \right) \left. \right\} \\ & + \left(\tilde{\mathbf{S}}_{(1)} \cdot \tilde{\mathbf{L}} \right) \left(\tilde{\mathbf{S}}_{(2)} \cdot \tilde{\mathbf{r}} \right) \tilde{p}_r \left\{ \tilde{L}^2 \left(-\frac{17\nu^3}{8\tilde{r}^7} - \frac{19\nu^2}{8\tilde{r}^7} - \frac{3\nu}{8\tilde{r}^7} \right) + \left(\frac{3\nu^3}{2\tilde{r}^6} + \frac{275\nu^2}{8\tilde{r}^6} - \frac{5\nu}{\tilde{r}^6} \right) \right. \\ & + \tilde{p}_r^2 \left(-\frac{37\nu^3}{4\tilde{r}^5} + \frac{73\nu^2}{16\tilde{r}^5} - \frac{3\nu}{8\tilde{r}^5} \right) \\ & + \frac{1}{q} \left(\tilde{L}^2 \left(\frac{9\nu^2}{8\tilde{r}^7} - \frac{3\nu^3}{4\tilde{r}^7} \right) + \left(\frac{3\nu^3}{2\tilde{r}^6} + \frac{7\nu^2}{\tilde{r}^6} \right) + \tilde{p}_r^2 \left(\frac{9\nu^2}{8\tilde{r}^5} - \frac{9\nu^3}{2\tilde{r}^5} \right) \right) \left. \right\}. \end{aligned} \quad (\text{B.3})$$

B.2 Spin1-spin1 and spin2-spin2 sector up to N²LO

$$\tilde{\mathcal{H}}_{\text{LO}}^{\text{S}^2} = 0 \quad (\text{B.4})$$

$$\begin{aligned} \tilde{\mathcal{H}}_{\text{NLO}}^{\text{S}^2} = & \left(\tilde{\mathbf{S}}_{(1)} \cdot \tilde{\mathbf{S}}_{(1)} \right) \left\{ \frac{3\tilde{L}^2\nu^2}{2\tilde{r}^5} - \frac{3\nu^2\tilde{p}_r^2}{8\tilde{r}^3} + \frac{7\nu^2}{8\tilde{r}^4} + \frac{1}{q} \left(\tilde{L}^2 \left(\frac{11\nu^2}{8\tilde{r}^5} - \frac{3\nu}{2\tilde{r}^5} \right) + \tilde{p}_r^2 \left(\frac{3\nu}{8\tilde{r}^3} - \frac{\nu^2}{2\tilde{r}^3} \right) + \frac{7\nu^2}{8\tilde{r}^4} + \frac{9\nu}{8\tilde{r}^4} \right) \right\} \\ & + \left(\tilde{\mathbf{S}}_{(1)} \cdot \tilde{\mathbf{r}} \right)^2 \left\{ + \frac{3\nu^2\tilde{p}_r^2}{8\tilde{r}^5} - \frac{15\nu^2}{8\tilde{r}^6} + \frac{1}{q} \left(\tilde{p}_r^2 \left(\frac{\nu^2}{2\tilde{r}^5} - \frac{3\nu}{8\tilde{r}^5} \right) - \frac{15\nu^2}{8\tilde{r}^6} - \frac{9\nu}{8\tilde{r}^6} \right) \right\} \\ & + \left(\tilde{\mathbf{S}}_{(1)} \cdot \tilde{\mathbf{L}} \right)^2 \left\{ - \frac{33\nu^2}{8\tilde{r}^5} + \frac{1}{q} \left(\frac{33\nu}{8\tilde{r}^5} - \frac{29\nu^2}{8\tilde{r}^5} \right) \right\} \\ & + \left(\tilde{\mathbf{S}}_{(1)} \cdot \tilde{\mathbf{r}} \right) \left(\tilde{\mathbf{S}}_{(1)} \cdot \tilde{\mathbf{L}} \right) \tilde{p}_r \left\{ - \frac{9\nu^2}{8\tilde{r}^5} + \frac{1}{q} \left(\frac{9\nu}{8\tilde{r}^5} - \frac{7\nu^2}{8\tilde{r}^5} \right) \right\} \\ & + (1 \leftrightarrow 2) \end{aligned} \quad (\text{B.5})$$

$$\begin{aligned} \tilde{\mathcal{H}}_{\text{N}^2\text{LO}}^{\text{S}^2} = & \left(\tilde{\mathbf{S}}_{(1)} \cdot \tilde{\mathbf{S}}_{(1)} \right) \left\{ \tilde{L}^4 \left(\frac{15\nu^3}{8\tilde{r}^7} - \frac{7\nu^2}{8\tilde{r}^7} \right) + \tilde{L}^2 \left(\tilde{p}_r^2 \left(\frac{87\nu^3}{16\tilde{r}^5} - \frac{\nu^2}{4\tilde{r}^5} \right) - \frac{13\nu^3}{16\tilde{r}^6} - \frac{35\nu^2}{4\tilde{r}^6} \right) \right. \\ & + \tilde{p}_r^2 \left(\frac{39\nu^3}{16\tilde{r}^4} + \frac{9\nu^2}{2\tilde{r}^4} \right) + \tilde{p}_r^4 \left(\frac{5\nu^2}{8\tilde{r}^3} - \frac{9\nu^3}{8\tilde{r}^3} \right) - \frac{101\nu^2}{56\tilde{r}^5} \\ & + \frac{1}{q} \left(\tilde{L}^4 \left(\frac{7\nu^3}{4\tilde{r}^7} - \frac{67\nu^2}{16\tilde{r}^7} + \frac{7\nu}{8\tilde{r}^7} \right) + \tilde{L}^2 \left(\tilde{p}_r^2 \left(\frac{5\nu^3}{\tilde{r}^5} - \frac{41\nu^2}{16\tilde{r}^5} + \frac{\nu}{4\tilde{r}^5} \right) - \frac{13\nu^3}{16\tilde{r}^6} - \frac{37\nu^2}{8\tilde{r}^6} + \frac{17\nu}{2\tilde{r}^6} \right) \right. \\ & + \tilde{p}_r^2 \left(\frac{39\nu^3}{16\tilde{r}^4} + \frac{147\nu^2}{16\tilde{r}^4} - \frac{3\nu}{\tilde{r}^4} \right) + \tilde{p}_r^4 \left(- \frac{23\nu^3}{16\tilde{r}^3} + \frac{13\nu^2}{8\tilde{r}^3} - \frac{5\nu}{8\tilde{r}^3} \right) - \frac{5\nu^2}{2\tilde{r}^5} - \frac{33\nu}{8\tilde{r}^5} \left. \right\} \\ & + \left(\tilde{\mathbf{S}}_{(1)} \cdot \tilde{\mathbf{r}} \right)^2 \left\{ \tilde{L}^2 \left(\tilde{p}_r^2 \left(- \frac{9\nu^3}{16\tilde{r}^7} - \frac{7\nu^2}{16\tilde{r}^7} \right) - \frac{9\nu^3}{8\tilde{r}^8} - \frac{25\nu^2}{8\tilde{r}^8} \right) - \frac{79\nu^3\tilde{p}_r^2}{16\tilde{r}^6} + \tilde{p}_r^4 \left(\frac{9\nu^3}{8\tilde{r}^5} - \frac{5\nu^2}{8\tilde{r}^5} \right) + \frac{317\nu^2}{56\tilde{r}^7} \right. \\ & + \frac{1}{q} \left(\tilde{L}^2 \left(\tilde{p}_r^2 \left(- \frac{7\nu^3}{16\tilde{r}^7} - \frac{23\nu^2}{16\tilde{r}^7} + \frac{7\nu}{16\tilde{r}^7} \right) - \frac{9\nu^3}{8\tilde{r}^8} - \frac{19\nu^2}{8\tilde{r}^8} + \frac{3\nu}{8\tilde{r}^8} \right) \right. \\ & + \tilde{p}_r^2 \left(- \frac{79\nu^3}{16\tilde{r}^6} - \frac{147\nu^2}{16\tilde{r}^6} + \frac{3\nu}{\tilde{r}^6} \right) + \tilde{p}_r^4 \left(\frac{23\nu^3}{16\tilde{r}^5} - \frac{13\nu^2}{8\tilde{r}^5} + \frac{5\nu}{8\tilde{r}^5} \right) + \frac{17\nu^2}{2\tilde{r}^7} + \frac{33\nu}{8\tilde{r}^7} \left. \right\} \\ & + \left(\tilde{\mathbf{S}}_{(1)} \cdot \tilde{\mathbf{L}} \right)^2 \left\{ \tilde{L}^2 \left(\frac{2\nu^2}{\tilde{r}^7} - \frac{39\nu^3}{8\tilde{r}^7} \right) + \tilde{p}_r^2 \left(\frac{29\nu^2}{16\tilde{r}^5} - \frac{123\nu^3}{8\tilde{r}^5} \right) + \frac{31\nu^3}{8\tilde{r}^6} + \frac{235\nu^2}{8\tilde{r}^6} \right. \\ & + \frac{1}{q} \left(\tilde{L}^2 \left(- \frac{35\nu^3}{8\tilde{r}^7} + \frac{175\nu^2}{16\tilde{r}^7} - \frac{2\nu}{\tilde{r}^7} \right) + \tilde{p}_r^2 \left(- \frac{55\nu^3}{4\tilde{r}^5} + \frac{43\nu^2}{4\tilde{r}^5} - \frac{29\nu}{16\tilde{r}^5} \right) + \frac{31\nu^3}{8\tilde{r}^6} + \frac{305\nu^2}{16\tilde{r}^6} - \frac{203\nu}{8\tilde{r}^6} \right) \left. \right\} \\ & + \left(\tilde{\mathbf{S}}_{(1)} \cdot \tilde{\mathbf{r}} \right) \left(\tilde{\mathbf{S}}_{(1)} \cdot \tilde{\mathbf{L}} \right) \tilde{p}_r \left\{ \tilde{L}^2 \left(\frac{7\nu^2}{16\tilde{r}^7} - \frac{39\nu^3}{16\tilde{r}^7} \right) + \left(\frac{13\nu^3}{4\tilde{r}^6} + \frac{13\nu^2}{4\tilde{r}^6} \right) + \tilde{p}_r^2 \left(\frac{\nu^2}{16\tilde{r}^5} - \frac{15\nu^3}{4\tilde{r}^5} \right) \right. \\ & + \frac{1}{q} \left(\tilde{L}^2 \left(- \frac{35\nu^3}{16\tilde{r}^7} + \frac{11\nu^2}{4\tilde{r}^7} - \frac{7\nu}{16\tilde{r}^7} \right) + \left(\frac{13\nu^3}{4\tilde{r}^6} + \frac{7\nu^2}{\tilde{r}^6} - \frac{25\nu}{4\tilde{r}^6} \right) \right. \\ & + \tilde{p}_r^2 \left(- \frac{25\nu^3}{8\tilde{r}^5} + \frac{19\nu^2}{8\tilde{r}^5} - \frac{\nu}{16\tilde{r}^5} \right) \left. \right\} \\ & + (1 \leftrightarrow 2). \end{aligned} \quad (\text{B.6})$$

B.3 ES² sector up to N²LO

$$\tilde{\mathcal{H}}_{\text{LO}}^{\text{ES}^2} = \left(C_{\text{ES}^2}^{(0)} \right)_{(1)} \left(\tilde{\mathbf{S}}_{(1)} \cdot \tilde{\mathbf{S}}_{(1)} \right) \left\{ -\frac{1}{q} \frac{\nu}{2\tilde{r}^3} \right\} + \left(C_{\text{ES}^2}^{(0)} \right)_{(1)} \left((\tilde{\mathbf{S}}_{(1)} \cdot \tilde{\mathbf{r}})^2 \right) \left\{ \frac{1}{q} \frac{3\nu}{2\tilde{r}^5} \right\} + (1 \leftrightarrow 2) \quad (\text{B.7})$$

$$\begin{aligned} \tilde{\mathcal{H}}_{\text{NLO}}^{\text{ES}^2} = & \left(C_{\text{ES}^2}^{(0)} \right)_{(1)} \left(\tilde{\mathbf{S}}_{(1)} \cdot \tilde{\mathbf{S}}_{(1)} \right) \left\{ -\frac{\tilde{L}^2 \nu^2}{2\tilde{r}^5} + \frac{\nu^2 \tilde{p}_r^2}{\tilde{r}^3} - \frac{\nu^2}{2\tilde{r}^4} + \frac{1}{q} \left(\tilde{L}^2 \left(-\frac{3\nu^2}{4\tilde{r}^5} - \frac{5\nu}{4\tilde{r}^5} \right) + \frac{\nu \tilde{p}_r^2}{4\tilde{r}^3} - \frac{\nu^2}{2\tilde{r}^4} + \frac{2\nu}{\tilde{r}^4} \right) \right\} \\ & + \left(C_{\text{ES}^2}^{(0)} \right)_{(1)} \left((\tilde{\mathbf{S}}_{(1)} \cdot \tilde{\mathbf{r}})^2 \right) \left\{ \frac{5\nu^2}{2\tilde{r}^6} - \frac{\nu^2 \tilde{p}_r^2}{\tilde{r}^5} + \frac{1}{q} \left(\tilde{L}^2 \left(\frac{3\nu^2}{4\tilde{r}^7} + \frac{9\nu}{4\tilde{r}^7} \right) + \tilde{p}_r^2 \left(\frac{2\nu^2}{\tilde{r}^5} - \frac{3\nu}{4\tilde{r}^5} \right) + \frac{5\nu^2}{2\tilde{r}^6} - \frac{6\nu}{\tilde{r}^6} \right) \right\} \\ & + \left(C_{\text{ES}^2}^{(0)} \right)_{(1)} \left((\tilde{\mathbf{S}}_{(1)} \cdot \tilde{\mathbf{L}})^2 \right) \left\{ \frac{\nu^2}{2\tilde{r}^5} + \frac{1}{q} \left(\frac{\nu^2}{2\tilde{r}^5} + \frac{3\nu}{2\tilde{r}^5} \right) \right\} \\ & + \left(C_{\text{ES}^2}^{(0)} \right)_{(1)} \left((\tilde{\mathbf{S}}_{(1)} \cdot \tilde{\mathbf{r}}) (\tilde{\mathbf{S}}_{(1)} \cdot \tilde{\mathbf{L}}) \right) \tilde{p}_r \left\{ -\frac{\nu^2}{2\tilde{r}^5} + \frac{1}{q} \left(-\frac{2\nu^2}{\tilde{r}^5} - \frac{3\nu}{2\tilde{r}^5} \right) \right\} \\ & + (1 \leftrightarrow 2) \end{aligned} \quad (\text{B.8})$$

$$\begin{aligned} \tilde{\mathcal{H}}_{\text{N}^2\text{LO}}^{\text{ES}^2} = & \left(C_{\text{ES}^2}^{(0)} \right)_{(1)} \left(\tilde{\mathbf{S}}_{(1)} \cdot \tilde{\mathbf{S}}_{(1)} \right) \left\{ \tilde{L}^4 \left(\frac{\nu^2}{4\tilde{r}^7} - \frac{3\nu^3}{4\tilde{r}^7} \right) + \tilde{L}^2 \left(\tilde{p}_r^2 \left(-\frac{3\nu^3}{2\tilde{r}^5} - \frac{\nu^2}{4\tilde{r}^5} \right) + \frac{\nu^3}{4\tilde{r}^6} + \frac{\nu^2}{2\tilde{r}^6} \right) \right. \\ & + \tilde{p}_r^2 \left(-\frac{7\nu^3}{4\tilde{r}^4} - \frac{15\nu^2}{4\tilde{r}^4} \right) + \tilde{p}_r^4 \left(\frac{3\nu^3}{\tilde{r}^3} - \frac{\nu^2}{2\tilde{r}^3} \right) + \frac{27\nu^2}{14\tilde{r}^5} \\ & + \frac{1}{q} \left(\tilde{L}^4 \left(-\frac{15\nu^3}{16\tilde{r}^7} - \frac{7\nu^2}{4\tilde{r}^7} + \frac{9\nu}{16\tilde{r}^7} \right) + \tilde{L}^2 \left(\tilde{p}_r^2 \left(-\frac{9\nu^3}{4\tilde{r}^5} - \frac{17\nu^2}{4\tilde{r}^5} + \frac{3\nu}{8\tilde{r}^5} \right) + \frac{\nu^3}{4\tilde{r}^6} + \frac{49\nu^2}{8\tilde{r}^6} + \frac{15\nu}{2\tilde{r}^6} \right) \right. \\ & \left. + \tilde{p}_r^2 \left(-\frac{7\nu^3}{4\tilde{r}^4} + \frac{27\nu^2}{8\tilde{r}^4} - \frac{3\nu}{2\tilde{r}^4} \right) + \tilde{p}_r^4 \left(\frac{3\nu^3}{2\tilde{r}^3} + \frac{5\nu^2}{4\tilde{r}^3} - \frac{3\nu}{16\tilde{r}^3} \right) + \frac{\nu^2}{4\tilde{r}^5} - \frac{19\nu}{4\tilde{r}^5} \right\} \\ & + \left(C_{\text{ES}^2}^{(0)} \right)_{(1)} \left((\tilde{\mathbf{S}}_{(1)} \cdot \tilde{\mathbf{r}})^2 \right) \left\{ \tilde{L}^2 \left(\tilde{p}_r^2 \left(-\frac{3\nu^3}{4\tilde{r}^7} - \frac{\nu^2}{8\tilde{r}^7} \right) + \frac{5\nu^3}{4\tilde{r}^8} + \frac{17\nu^2}{2\tilde{r}^8} \right) + \tilde{p}_r^2 \left(\frac{27\nu^3}{4\tilde{r}^6} + \frac{15\nu^2}{4\tilde{r}^6} \right) \right. \\ & + \tilde{p}_r^4 \left(\frac{5\nu^2}{2\tilde{r}^5} - \frac{3\nu^3}{\tilde{r}^5} \right) - \frac{123\nu^2}{14\tilde{r}^7} \\ & + \frac{1}{q} \left(\tilde{L}^4 \left(\frac{9\nu^3}{16\tilde{r}^9} + \frac{3\nu^2}{\tilde{r}^9} - \frac{15\nu}{16\tilde{r}^9} \right) + \tilde{L}^2 \left(\tilde{p}_r^2 \left(\frac{9\nu^3}{8\tilde{r}^7} + \frac{23\nu^2}{4\tilde{r}^7} - \frac{3\nu}{4\tilde{r}^7} \right) + \frac{5\nu^3}{4\tilde{r}^8} - \frac{\nu^2}{4\tilde{r}^8} - \frac{27\nu}{2\tilde{r}^8} \right) \right. \\ & \left. + \tilde{p}_r^2 \left(\frac{27\nu^3}{4\tilde{r}^6} - \frac{177\nu^2}{8\tilde{r}^6} + \frac{9\nu}{2\tilde{r}^6} \right) + \tilde{p}_r^4 \left(\frac{3\nu^3}{2\tilde{r}^5} - \frac{7\nu^2}{4\tilde{r}^5} + \frac{9\nu}{16\tilde{r}^5} \right) + \frac{57\nu}{4\tilde{r}^7} - \frac{23\nu^2}{4\tilde{r}^7} \right\} \\ & + \left(C_{\text{ES}^2}^{(0)} \right)_{(1)} \left((\tilde{\mathbf{S}}_{(1)} \cdot \tilde{\mathbf{L}})^2 \right) \left\{ \tilde{L}^2 \left(\frac{3\nu^3}{4\tilde{r}^7} - \frac{5\nu^2}{4\tilde{r}^7} \right) + \tilde{p}_r^2 \left(\frac{9\nu^3}{4\tilde{r}^5} + \frac{11\nu^2}{8\tilde{r}^5} \right) - \frac{\nu^3}{2\tilde{r}^6} - \frac{5\nu^2}{2\tilde{r}^6} \right. \\ & \left. + \frac{1}{q} \left(\tilde{L}^2 \left(\frac{3\nu^3}{4\tilde{r}^7} + \frac{5\nu^2}{4\tilde{r}^7} - \frac{3\nu}{4\tilde{r}^7} \right) + \tilde{p}_r^2 \left(\frac{21\nu^3}{8\tilde{r}^5} + \frac{8\nu^2}{\tilde{r}^5} - \frac{3\nu}{8\tilde{r}^5} \right) - \frac{\nu^3}{2\tilde{r}^6} - \frac{65\nu^2}{8\tilde{r}^6} - \frac{9\nu}{\tilde{r}^6} \right) \right\} \end{aligned}$$

$$\begin{aligned}
 & + \left(C_{\text{ES}^2}^{(0)} \right)_{(1)} \left((\tilde{\mathbf{S}}_{(1)} \cdot \tilde{\mathbf{r}}) (\tilde{\mathbf{S}}_{(1)} \cdot \tilde{\mathbf{L}}) \tilde{p}_r \left\{ -\frac{11\tilde{L}^2\nu^2}{8\tilde{r}^7} + \left(\frac{25\nu^2}{\tilde{r}^6} - \frac{7\nu^3}{4\tilde{r}^6} \right) + \tilde{p}_r^2 \left(\frac{31\nu^2}{8\tilde{r}^5} - \frac{3\nu^3}{4\tilde{r}^5} \right) \right. \right. \\
 & + \frac{1}{q} \left(\tilde{L}^2 \left(-\frac{9\nu^3}{8\tilde{r}^7} - \frac{11\nu^2}{4\tilde{r}^7} + \frac{3\nu}{8\tilde{r}^7} \right) + \left(-\frac{7\nu^3}{4\tilde{r}^6} + \frac{42\nu^2}{\tilde{r}^6} + \frac{9\nu}{\tilde{r}^6} \right) \right. \\
 & \left. \left. + \tilde{p}_r^2 \left(-\frac{39\nu^3}{8\tilde{r}^5} - \frac{\nu^2}{2\tilde{r}^5} + \frac{9\nu}{8\tilde{r}^5} \right) \right) \right\} \\
 & + (1 \leftrightarrow 2). \tag{B.9}
 \end{aligned}$$

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] LIGO SCIENTIFIC and VIRGO collaborations, *Observation of Gravitational Waves from a Binary Black Hole Merger*, *Phys. Rev. Lett.* **116** (2016) 061102 [[arXiv:1602.03837](https://arxiv.org/abs/1602.03837)] [[INSPIRE](#)].
- [2] LIGO SCIENTIFIC, VIRGO and KAGRA collaborations, *GWTC-3: Compact Binary Coalescences Observed by LIGO and Virgo During the Second Part of the Third Observing Run*, [arXiv:2111.03606](https://arxiv.org/abs/2111.03606) [[INSPIRE](#)].
- [3] LIGO SCIENTIFIC collaboration, *Advanced LIGO*, *Class. Quant. Grav.* **32** (2015) 074001 [[arXiv:1411.4547](https://arxiv.org/abs/1411.4547)] [[INSPIRE](#)].
- [4] VIRGO collaboration, *Advanced Virgo: a second-generation interferometric gravitational wave detector*, *Class. Quant. Grav.* **32** (2015) 024001 [[arXiv:1408.3978](https://arxiv.org/abs/1408.3978)] [[INSPIRE](#)].
- [5] KAGRA collaboration, *Overview of KAGRA: Calibration, detector characterization, physical environmental monitors, and the geophysics interferometer*, *Prog. Theor. Exp. Phys.* **2021** (2021) 05A102 [[arXiv:2009.09305](https://arxiv.org/abs/2009.09305)] [[INSPIRE](#)].
- [6] M. Saleem et al., *The science case for LIGO-India*, *Class. Quant. Grav.* **39** (2022) 025004 [[arXiv:2105.01716](https://arxiv.org/abs/2105.01716)] [[INSPIRE](#)].
- [7] LIGO SCIENTIFIC collaboration, *Exploring the Sensitivity of Next Generation Gravitational Wave Detectors*, *Class. Quant. Grav.* **34** (2017) 044001 [[arXiv:1607.08697](https://arxiv.org/abs/1607.08697)] [[INSPIRE](#)].
- [8] M. Punturo et al., *The third generation of gravitational wave observatories and their science reach*, *Class. Quant. Grav.* **27** (2010) 084007 [[INSPIRE](#)].
- [9] LISA collaboration, *Laser Interferometer Space Antenna*, [arXiv:1702.00786](https://arxiv.org/abs/1702.00786) [[INSPIRE](#)].
- [10] S. Olsen, T. Venumadhav, J. Mushkin, J. Roulet, B. Zackay and M. Zaldarriaga, *New binary black hole mergers in the LIGO-Virgo O3a data*, *Phys. Rev. D* **106** (2022) 043009 [[arXiv:2201.02252](https://arxiv.org/abs/2201.02252)] [[INSPIRE](#)].
- [11] W. Tulczyjew, *Equations of motion of rotating bodies in general relativity theory*, *Acta Phys. Polon* **18** (1959) 37.
- [12] T. Damour, *Problème des deux corps et freinage de rayonnement en relativité générale*, *C. R. Acad. Sci.* **294** (1982) 1355.

- [13] H. Tagoshi, A. Ohashi and B.J. Owen, *Gravitational field and equations of motion of spinning compact binaries to 2.5 post-Newtonian order*, *Phys. Rev. D* **63** (2001) 044006 [[gr-qc/0010014](#)] [[INSPIRE](#)].
- [14] G. Faye, L. Blanchet and A. Buonanno, *Higher-order spin effects in the dynamics of compact binaries. Part I. Equations of motion*, *Phys. Rev. D* **74** (2006) 104033 [[gr-qc/0605139](#)] [[INSPIRE](#)].
- [15] T. Damour, P. Jaranowski and G. Schäfer, *Hamiltonian of two spinning compact bodies with next-to-leading order gravitational spin-orbit coupling*, *Phys. Rev. D* **77** (2008) 064032 [[arXiv:0711.1048](#)] [[INSPIRE](#)].
- [16] J. Hartung and J. Steinhoff, *Next-to-next-to-leading order post-Newtonian spin-orbit Hamiltonian for self-gravitating binaries*, *Annalen Phys.* **523** (2011) 783 [[arXiv:1104.3079](#)] [[INSPIRE](#)].
- [17] J. Hartung, J. Steinhoff and G. Schäfer, *Next-to-next-to-leading order post-Newtonian linear-in-spin binary Hamiltonians*, *Annalen Phys.* **525** (2013) 359 [[arXiv:1302.6723](#)] [[INSPIRE](#)].
- [18] J. Steinhoff, G. Schäfer and S. Hergt, *ADM canonical formalism for gravitating spinning objects*, *Phys. Rev. D* **77** (2008) 104018 [[arXiv:0805.3136](#)] [[INSPIRE](#)].
- [19] J. Steinhoff and G. Schäfer, *Canonical formulation of self-gravitating spinning-object systems*, *Europhys. Lett.* **87** (2009) 50004 [[arXiv:0907.1967](#)] [[INSPIRE](#)].
- [20] A. Bohe, S. Marsat, G. Faye and L. Blanchet, *Next-to-next-to-leading order spin-orbit effects in the near-zone metric and precession equations of compact binaries*, *Class. Quant. Grav.* **30** (2013) 075017 [[arXiv:1212.5520](#)] [[INSPIRE](#)].
- [21] R. Aoude, K. Haddad and A. Helset, *Classical Gravitational Spinning-Spinless Scattering at $\mathcal{O}(G^2 S^\infty)$* , *Phys. Rev. Lett.* **129** (2022) 141102 [[arXiv:2205.02809](#)] [[INSPIRE](#)].
- [22] F. Febres Cordero, M. Kraus, G. Lin, M.S. Ruf and M. Zeng, *Conservative Binary Dynamics with a Spinning Black Hole at $\mathcal{O}(G^3)$ from Scattering Amplitudes*, *Phys. Rev. Lett.* **130** (2023) 021601 [[arXiv:2205.07357](#)] [[INSPIRE](#)].
- [23] Z. Bern, D. Kosmopoulos, A. Luna, R. Roiban and F. Teng, *Binary Dynamics through the Fifth Power of Spin at $\mathcal{O}(G^2)$* , *Phys. Rev. Lett.* **130** (2023) 201402 [[arXiv:2203.06202](#)] [[INSPIRE](#)].
- [24] Z. Liu, R.A. Porto and Z. Yang, *Spin Effects in the Effective Field Theory Approach to Post-Minkowskian Conservative Dynamics*, *JHEP* **06** (2021) 012 [[arXiv:2102.10059](#)] [[INSPIRE](#)].
- [25] D. Kosmopoulos and A. Luna, *Quadratic-in-spin Hamiltonian at $\mathcal{O}(G^2)$ from scattering amplitudes*, *JHEP* **07** (2021) 037 [[arXiv:2102.10137](#)] [[INSPIRE](#)].
- [26] G.U. Jakobsen and G. Mogull, *Conservative and Radiative Dynamics of Spinning Bodies at Third Post-Minkowskian Order Using Worldline Quantum Field Theory*, *Phys. Rev. Lett.* **128** (2022) 141102 [[arXiv:2201.07778](#)] [[INSPIRE](#)].
- [27] G.U. Jakobsen and G. Mogull, *Linear response, Hamiltonian, and radiative spinning two-body dynamics*, *Phys. Rev. D* **107** (2023) 044033 [[arXiv:2210.06451](#)] [[INSPIRE](#)].
- [28] R.A. Porto, *Post-Newtonian corrections to the motion of spinning bodies in NRGR*, *Phys. Rev. D* **73** (2006) 104031 [[gr-qc/0511061](#)] [[INSPIRE](#)].

- [29] R.A. Porto, *Next to leading order spin-orbit effects in the motion of inspiralling compact binaries*, *Class. Quant. Grav.* **27** (2010) 205001 [[arXiv:1005.5730](#)] [[INSPIRE](#)].
- [30] M. Levi, *Next to Leading Order gravitational Spin-Orbit coupling in an Effective Field Theory approach*, *Phys. Rev. D* **82** (2010) 104004 [[arXiv:1006.4139](#)] [[INSPIRE](#)].
- [31] M. Levi, A.J. Mcleod and M. Von Hippel, *N^3LO gravitational spin-orbit coupling at order G^4* , *JHEP* **07** (2021) 115 [[arXiv:2003.02827](#)] [[INSPIRE](#)].
- [32] R.A. Porto and I.Z. Rothstein, *The Hyperfine Einstein-Infeld-Hoffmann potential*, *Phys. Rev. Lett.* **97** (2006) 021101 [[gr-qc/0604099](#)] [[INSPIRE](#)].
- [33] R.A. Porto and I.Z. Rothstein, *Spin(1)Spin(2) Effects in the Motion of Inspiralling Compact Binaries at Third Order in the Post-Newtonian Expansion*, *Phys. Rev. D* **78** (2008) 044012 [*Erratum ibid.* **81** (2010) 029904] [[arXiv:0802.0720](#)] [[INSPIRE](#)].
- [34] M. Levi, *Next to Leading Order gravitational Spin1-Spin2 coupling with Kaluza-Klein reduction*, *Phys. Rev. D* **82** (2010) 064029 [[arXiv:0802.1508](#)] [[INSPIRE](#)].
- [35] R.A. Porto and I.Z. Rothstein, *Next to Leading Order Spin(1)Spin(1) Effects in the Motion of Inspiralling Compact Binaries*, *Phys. Rev. D* **78** (2008) 044013 [*Erratum ibid.* **81** (2010) 029905] [[arXiv:0804.0260](#)] [[INSPIRE](#)].
- [36] M. Levi, *Binary dynamics from spin1-spin2 coupling at fourth post-Newtonian order*, *Phys. Rev. D* **85** (2012) 064043 [[arXiv:1107.4322](#)] [[INSPIRE](#)].
- [37] M. Levi and J. Steinhoff, *Equivalence of ADM Hamiltonian and Effective Field Theory approaches at next-to-next-to-leading order spin1-spin2 coupling of binary inspirals*, *JCAP* **12** (2014) 003 [[arXiv:1408.5762](#)] [[INSPIRE](#)].
- [38] M. Levi and J. Steinhoff, *Next-to-next-to-leading order gravitational spin-squared potential via the effective field theory for spinning objects in the post-Newtonian scheme*, *JCAP* **01** (2016) 008 [[arXiv:1506.05794](#)] [[INSPIRE](#)].
- [39] J.-W. Kim, M. Levi and Z. Yin, *Quadratic-in-spin interactions at fifth post-Newtonian order probe new physics*, *Phys. Lett. B* **834** (2022) 137410 [[arXiv:2112.01509](#)] [[INSPIRE](#)].
- [40] M. Levi, A.J. Mcleod and M. Von Hippel, *N^3LO gravitational quadratic-in-spin interactions at G^4* , *JHEP* **07** (2021) 116 [[arXiv:2003.07890](#)] [[INSPIRE](#)].
- [41] M. Levi, S. Mougiakakos and M. Vieira, *Gravitational cubic-in-spin interaction at the next-to-leading post-Newtonian order*, *JHEP* **01** (2021) 036 [[arXiv:1912.06276](#)] [[INSPIRE](#)].
- [42] M. Levi and J. Steinhoff, *Spinning gravitating objects in the effective field theory in the post-Newtonian scheme*, *JHEP* **09** (2015) 219 [[arXiv:1501.04956](#)] [[INSPIRE](#)].
- [43] M. Levi and J. Steinhoff, *Leading order finite size effects with spins for inspiralling compact binaries*, *JHEP* **06** (2015) 059 [[arXiv:1410.2601](#)] [[INSPIRE](#)].
- [44] A. Guevara, *Holomorphic Classical Limit for Spin Effects in Gravitational and Electromagnetic Scattering*, *JHEP* **04** (2019) 033 [[arXiv:1706.02314](#)] [[INSPIRE](#)].
- [45] J. Vines, J. Steinhoff and A. Buonanno, *Spinning-black-hole scattering and the test-black-hole limit at second post-Minkowskian order*, *Phys. Rev. D* **99** (2019) 064054 [[arXiv:1812.00956](#)] [[INSPIRE](#)].
- [46] A. Guevara, A. Ochirov and J. Vines, *Scattering of Spinning Black Holes from Exponentiated Soft Factors*, *JHEP* **09** (2019) 056 [[arXiv:1812.06895](#)] [[INSPIRE](#)].

- [47] M.-Z. Chung, Y.-T. Huang, J.-W. Kim and S. Lee, *The simplest massive S-matrix: from minimal coupling to Black Holes*, *JHEP* **04** (2019) 156 [[arXiv:1812.08752](#)] [[INSPIRE](#)].
- [48] A. Guevara, A. Ochirov and J. Vines, *Black-hole scattering with general spin directions from minimal-coupling amplitudes*, *Phys. Rev. D* **100** (2019) 104024 [[arXiv:1906.10071](#)] [[INSPIRE](#)].
- [49] M.-Z. Chung, Y.-T. Huang and J.-W. Kim, *Classical potential for general spinning bodies*, *JHEP* **09** (2020) 074 [[arXiv:1908.08463](#)] [[INSPIRE](#)].
- [50] N. Siemonsen and J. Vines, *Test black holes, scattering amplitudes and perturbations of Kerr spacetime*, *Phys. Rev. D* **101** (2020) 064066 [[arXiv:1909.07361](#)] [[INSPIRE](#)].
- [51] A. Guevara, B. Maybee, A. Ochirov, D. O’Connell and J. Vines, *A worldsheet for Kerr*, *JHEP* **03** (2021) 201 [[arXiv:2012.11570](#)] [[INSPIRE](#)].
- [52] N. Arkani-Hamed, Y.-T. Huang and D. O’Connell, *Kerr black holes as elementary particles*, *JHEP* **01** (2020) 046 [[arXiv:1906.10100](#)] [[INSPIRE](#)].
- [53] M. Levi, *Effective Field Theories of Post-Newtonian Gravity: A comprehensive review*, *Rept. Prog. Phys.* **83** (2020) 075901 [[arXiv:1807.01699](#)] [[INSPIRE](#)].
- [54] R.A. Porto, *The effective field theorist’s approach to gravitational dynamics*, *Phys. Rept.* **633** (2016) 1 [[arXiv:1601.04914](#)] [[INSPIRE](#)].
- [55] L. Blanchet, *Gravitational Radiation from Post-Newtonian Sources and Inspiralling Compact Binaries*, *Living Rev. Rel.* **17** (2014) 2 [[arXiv:1310.1528](#)] [[INSPIRE](#)].
- [56] S. Foffa, P. Mastrolia, R. Sturani, C. Sturm and W.J. Torres Bobadilla, *Static two-body potential at fifth post-Newtonian order*, *Phys. Rev. Lett.* **122** (2019) 241605 [[arXiv:1902.10571](#)] [[INSPIRE](#)].
- [57] J. Blümlein, A. Maier and P. Marquard, *Five-Loop Static Contribution to the Gravitational Interaction Potential of Two Point Masses*, *Phys. Lett. B* **800** (2020) 135100 [[arXiv:1902.11180](#)] [[INSPIRE](#)].
- [58] J. Blümlein, A. Maier, P. Marquard and G. Schäfer, *The fifth-order post-Newtonian Hamiltonian dynamics of two-body systems from an effective field theory approach: potential contributions*, *Nucl. Phys. B* **965** (2021) 115352 [[arXiv:2010.13672](#)] [[INSPIRE](#)].
- [59] S. Foffa, R. Sturani and W.J. Torres Bobadilla, *Efficient resummation of high post-Newtonian contributions to the binding energy*, *JHEP* **02** (2021) 165 [[arXiv:2010.13730](#)] [[INSPIRE](#)].
- [60] J. Blümlein, A. Maier, P. Marquard and G. Schäfer, *The fifth-order post-Newtonian Hamiltonian dynamics of two-body systems from an effective field theory approach*, *Nucl. Phys. B* **983** (2022) 115900 [Erratum *ibid.* **985** (2022) 115991] [[arXiv:2110.13822](#)] [[INSPIRE](#)].
- [61] J. Blümlein, A. Maier, P. Marquard and G. Schäfer, *Testing binary dynamics in gravity at the sixth post-Newtonian level*, *Phys. Lett. B* **807** (2020) 135496 [[arXiv:2003.07145](#)] [[INSPIRE](#)].
- [62] J. Blümlein, A. Maier, P. Marquard and G. Schäfer, *The 6th post-Newtonian potential terms at $O(G_N^4)$* , *Phys. Lett. B* **816** (2021) 136260 [[arXiv:2101.08630](#)] [[INSPIRE](#)].
- [63] M. Levi and J. Steinhoff, *Next-to-next-to-leading order gravitational spin-orbit coupling via the effective field theory for spinning objects in the post-Newtonian scheme*, *JCAP* **01** (2016) 011 [[arXiv:1506.05056](#)] [[INSPIRE](#)].
- [64] J.-W. Kim, M. Levi and Z. Yin, *N^3LO spin-orbit interaction via the EFT of spinning gravitating objects*, *JHEP* **05** (2023) 184 [[arXiv:2208.14949](#)] [[INSPIRE](#)].

- [65] M.K. Mandal, P. Mastrolia, R. Patil and J. Steinhoff, *Gravitational spin-orbit Hamiltonian at NNNLO in the post-Newtonian framework*, *JHEP* **03** (2023) 130 [[arXiv:2209.00611](#)] [[INSPIRE](#)].
- [66] J.-W. Kim, M. Levi and Z. Yin, *N^3LO quadratic-in-spin interactions for generic compact binaries*, *JHEP* **03** (2023) 098 [[arXiv:2209.09235](#)] [[INSPIRE](#)].
- [67] W.D. Goldberger and I.Z. Rothstein, *An Effective field theory of gravity for extended objects*, *Phys. Rev. D* **73** (2006) 104029 [[hep-th/0409156](#)] [[INSPIRE](#)].
- [68] S. Foffa, P. Mastrolia, R. Sturani and C. Sturm, *Effective field theory approach to the gravitational two-body dynamics, at fourth post-Newtonian order and quintic in the Newton constant*, *Phys. Rev. D* **95** (2017) 104009 [[arXiv:1612.00482](#)] [[INSPIRE](#)].
- [69] K.G. Chetyrkin and F.V. Tkachov, *Integration by parts: The algorithm to calculate β -functions in 4 loops*, *Nucl. Phys. B* **192** (1981) 159 [[INSPIRE](#)].
- [70] S. Laporta, *High precision calculation of multiloop Feynman integrals by difference equations*, *Int. J. Mod. Phys. A* **15** (2000) 5087 [[hep-ph/0102033](#)] [[INSPIRE](#)].
- [71] P. Nogueira, *Automatic Feynman graph generation*, *J. Comput. Phys.* **105** (1993) 279 [[INSPIRE](#)].
- [72] J.M.M. García, *xact: Efficient tensor computer algebra for mathematica*, version 1.2.0 (2021).
- [73] R.N. Lee, *LiteRed 1.4: a powerful tool for reduction of multiloop integrals*, *J. Phys. Conf. Ser.* **523** (2014) 012059 [[arXiv:1310.1145](#)] [[INSPIRE](#)].
- [74] M. Levi and J. Steinhoff, *EFTofPNG: A package for high precision computation with the Effective Field Theory of Post-Newtonian Gravity*, *Class. Quant. Grav.* **34** (2017) 244001 [[arXiv:1705.06309](#)] [[INSPIRE](#)].
- [75] B. Kol and M. Smolkin, *Non-Relativistic Gravitation: From Newton to Einstein and Back*, *Class. Quant. Grav.* **25** (2008) 145011 [[arXiv:0712.4116](#)] [[INSPIRE](#)].
- [76] B. Kol and M. Smolkin, *Classical Effective Field Theory and Caged Black Holes*, *Phys. Rev. D* **77** (2008) 064033 [[arXiv:0712.2822](#)] [[INSPIRE](#)].
- [77] A. von Manteuffel and C. Studerus, *Reduze 2 — Distributed Feynman Integral Reduction*, [arXiv:1201.4330](#) [[INSPIRE](#)].
- [78] P. Maierhöfer, J. Usovitsch and P. Uwer, *Kira — A Feynman integral reduction program*, *Comput. Phys. Commun.* **230** (2018) 99 [[arXiv:1705.05610](#)] [[INSPIRE](#)].
- [79] G. Schäfer, *Acceleration-dependent Lagrangian in general relativity*, *Phys. Lett. A* **100** (1984) 128 [[INSPIRE](#)].
- [80] T. Damour and G. Schäfer, *Redefinition of position variables and the reduction of higher order Lagrangians*, *J. Math. Phys.* **32** (1991) 127 [[INSPIRE](#)].
- [81] T. Damour and G. Schäfer, *Lagrangians for point masses at the second post-Newtonian approximation of general relativity*, *Gen. Rel. Grav.* **17** (1985) 879 [[INSPIRE](#)].
- [82] B.M. Barker and R.F. O’Connell, *Acceleration-dependent Lagrangian and equations of motion*, *Phys. Lett. A* **78** (1980) 231 [[INSPIRE](#)].
- [83] S. Foffa, R.A. Porto, I. Rothstein and R. Sturani, *Conservative dynamics of binary systems to fourth Post-Newtonian order in the EFT approach. Part II. Renormalized Lagrangian*, *Phys. Rev. D* **100** (2019) 024048 [[arXiv:1903.05118](#)] [[INSPIRE](#)].

- [84] S. Foffa and R. Sturani, *Effective field theory calculation of conservative binary dynamics at third post-Newtonian order*, *Phys. Rev. D* **84** (2011) 044031 [[arXiv:1104.1122](#)] [[INSPIRE](#)].
- [85] A. Antonelli, C. Kavanagh, M. Khalil, J. Steinhoff and J. Vines, *Gravitational spin-orbit and aligned spin₁-spin₂ couplings through third-subleading post-Newtonian orders*, *Phys. Rev. D* **102** (2020) 124024 [[arXiv:2010.02018](#)] [[INSPIRE](#)].
- [86] G. Kälin and R.A. Porto, *From Boundary Data to Bound States*, *JHEP* **01** (2020) 072 [[arXiv:1910.03008](#)] [[INSPIRE](#)].
- [87] G. Kälin and R.A. Porto, *From boundary data to bound states. Part II. Scattering angle to dynamical invariants (with twist)*, *JHEP* **02** (2020) 120 [[arXiv:1911.09130](#)] [[INSPIRE](#)].
- [88] G. Cho, G. Kälin and R.A. Porto, *From boundary data to bound states. Part III. Radiative effects*, *JHEP* **04** (2022) 154 [*Erratum ibid.* **07** (2022) 002] [[arXiv:2112.03976](#)] [[INSPIRE](#)].
- [89] P.D. D’Eath, *Interaction of two black holes in the slow-motion limit*, *Phys. Rev. D* **12** (1975) 2183 [[INSPIRE](#)].
- [90] K.S. Thorne, *Multipole Expansions of Gravitational Radiation*, *Rev. Mod. Phys.* **52** (1980) 299 [[INSPIRE](#)].
- [91] W.G. Laarakkers and E. Poisson, *Quadrupole moments of rotating neutron stars*, *Astrophys. J.* **512** (1999) 282 [[gr-qc/9709033](#)] [[INSPIRE](#)].
- [92] E. Poisson, *Tidal deformation of a slowly rotating black hole*, *Phys. Rev. D* **91** (2015) 044004 [[arXiv:1411.4711](#)] [[INSPIRE](#)].
- [93] P. Pani, L. Gualtieri, A. Maselli and V. Ferrari, *Tidal deformations of a spinning compact object*, *Phys. Rev. D* **92** (2015) 024010 [[arXiv:1503.07365](#)] [[INSPIRE](#)].
- [94] A. Le Tiec and M. Casals, *Spinning Black Holes Fall in Love*, *Phys. Rev. Lett.* **126** (2021) 131102 [[arXiv:2007.00214](#)] [[INSPIRE](#)].
- [95] M.M. Ivanov and Z. Zhou, *Vanishing of Black Hole Tidal Love Numbers from Scattering Amplitudes*, *Phys. Rev. Lett.* **130** (2023) 091403 [[arXiv:2209.14324](#)] [[INSPIRE](#)].