

PAPER • OPEN ACCESS

Feynman-Kac theory of time-integrated functionals: Itô versus functional calculus

To cite this article: Cai Dieball and Aljaž Godec 2023 *J. Phys. A: Math. Theor.* **56** 155002

View the [article online](#) for updates and enhancements.

You may also like

- [Path integrals for higher derivative actions](#)
David S Dean, Bing Miao and Rudi Podgornik
- [Brownian motion in trapping enclosures: steep potential wells, bistable wells and false bistability of induced Feynman–Kac \(well\) potentials](#)
Piotr Garbaczewski and Mariusz aba
- [Branching exponential flights: travelled lengths and collision statistics](#)
Andrea Zoia, Eric Dumonteil, Alain Mazzolo et al.

Feynman-Kac theory of time-integrated functionals: Itô versus functional calculus

Cai Dieball  and Aljaž Godec* 

Mathematical bioPhysics Group, Max Planck Institute for Multidisciplinary Sciences,
37077 Göttingen, Germany

E-mail: agodec@mpinat.mpg.de

Received 8 June 2022; revised 30 December 2022

Accepted for publication 8 March 2023

Published 20 March 2023



CrossMark

Abstract

The fluctuations of dynamical functionals such as the empirical density and current as well as heat, work and generalized currents in stochastic thermodynamics are usually studied within the Feynman-Kac tilting formalism, which in the Physics literature is typically derived by some form of Kramers-Moyal expansion, or in the Mathematical literature via the Cameron-Martin-Girsanov approach. Here we derive the Feynman-Kac theory for general additive dynamical functionals directly via Itô calculus and via functional calculus, where the latter results in fact appears to be new. Using Dyson series we then independently recapitulate recent results on steady-state (co)variances of general additive dynamical functionals derived recently in Dieball and Godec (2022 *Phys. Rev. Lett.* **129** 140601) and Dieball and Godec (2022 *Phys. Rev. Res.* **4** 033243). We hope for our work to put the different approaches to the statistics of dynamical functionals employed in the field on a common footing, and to illustrate more easily accessible ways to the tilting formalism.

Keywords: Feynman-Kac theory, Itô calculus, functional calculus, additive dynamical functionals, time-integrated density and current

1. Introduction

Dynamical functionals and diverse path-based observables [1–8], such as local and occupation times (also known as the ‘empirical density’) [9–17] as well as diverse time-integrated and time-averaged currents [18–31] are central to ‘time-average statistical mechanics’ [32–34],

* Author to whom any correspondence should be addressed.



Original Content from this work may be used under the terms of the [Creative Commons Attribution 4.0 licence](https://creativecommons.org/licenses/by/4.0/). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

large deviation theory (see e.g. [19, 22, 25, 26, 35]), macroscopic fluctuation theory [36–38], and path-wise, stochastic thermodynamics [28, 29, 39–44].

Several techniques are available for the study of dynamical functionals, presumably best known is the Lie-Trotter-Kato formalism [10, 45] that was employed by Kac in his seminal work [9]. The techniques typically employed in physics rely on an analogy to quantum mechanical problems (see e.g. [15]) or assume some form of the Kramers-Moyal expansion [13, 16, 46, 47] (see also interesting generalizations to anomalous dynamics [12, 17]).

Deriving Feynman-Kac theory [9] of such additive functionals amounts to obtaining a ‘tilted’ generator which generates the time-evolution of the observables under consideration. The tilted evolution operator can be obtained using the Cameron-Martin-Girsanov theorem [48, 49]—a well-known technical theorem often employed in the Mathematical Physics literature [21–23].

In this paper we develop the Itô [34, 50] and functional calculus [51, 52] approaches to Feynman-Kac theory, whereby we focus on the methodology and accessibility for readers that are unfamiliar with the Cameron-Martin-Girsanov approach to ‘tilting’. We thereby hope to provide two accessible alternative (but equivalent) ways to obtaining the tilted generator. While the Itô approach already exists (see e.g. [34] for the empirical density), our functional calculus approach is a generalization of the pedagogical work of Fox [51, 52] and is aimed towards readers who prefer to avoid Itô calculus. Since both methods are equivalent they yield the same tilted generator. This generator is subsequently used to re-derive recent results on the statistics of time-integrated densities and currents obtained in [30, 31] using a different, more direct, stochastic calculus approach that avoids tilting. In particular, these results illustrate the use of the tilted generator to derive the statistics of time-integrated observables for finite times, i.e. extending beyond large deviation theory.

The outline of the paper is as follows. In section 2.1 we provide the mathematical setup of the problem. In section 2.2 we derive the Feynman-Kac equation for a general dynamical functional of diffusion processes using Itô calculus. By generalizing the approach by Fox [51, 52] we derive in section 2.3 the Feynman-Kac equation using functional calculus. In section 3 we apply the formalism to compute steady-state (co)variances of general dynamical functionals using a Dyson-series approach. We conclude with a brief perspective.

2. Tilted generator

In this section, we first introduce the considered stochastic dynamics and define what we call ‘dynamical functionals’. Subsequently we derive the tilted generator (i.e. the operator generating the time-evolution of time-integrated functionals) based on Itô calculus, and finally equivalently also via functional calculus.

2.1. Set-up

We consider overdamped stochastic motion in d -dimensional space described by the stochastic differential equation

$$d\mathbf{x}_t = \mathbf{F}(\mathbf{x}_t)dt + \boldsymbol{\sigma}d\mathbf{W}_t, \quad (1)$$

where $d\mathbf{W}_t$ denotes increment of the Wiener process [50]. The corresponding diffusion constant is $\mathbf{D} = \boldsymbol{\sigma}\boldsymbol{\sigma}^T/2$. For simplicity we stick to additive noise whereas all present results generalize to multiplicative noise $\mathbf{D}(\mathbf{x})$ as described in [31]. In the physics literature equation (1) is typically written in the form of a Langevin equation

$$\dot{\mathbf{x}}_t = \mathbf{F}(\mathbf{x}_t) + \mathbf{f}(t), \quad (2)$$

with white noise amplitude $\langle \mathbf{f}(t)\mathbf{f}(t')^T \rangle = 2\mathbf{D}\delta(t-t')$. Comparing the two equations, $\mathbf{f}(t)$ corresponds to the derivative of \mathbf{W}_t , which however (with probability one) is not differentiable; more precisely, upon taking $dt \rightarrow 0$ one has $\|\mathbf{dW}_t/dt\| = \infty$ with probability one, which is why the mathematics literature prefers equation (1).

If one describes the system on the level of probability densities instead of trajectories, the above equations translate to the Fokker-Planck equation $\partial_t G(\mathbf{x}, t|\mathbf{x}_0) = \hat{L}(\mathbf{x})G(\mathbf{x}, t|\mathbf{x}_0)$ with conditional density $G(\mathbf{x}, t|\mathbf{x}_0)$ to be at \mathbf{x} at time t after starting in \mathbf{x}_0 and the Fokker-Planck operator [53, 54]

$$\hat{L}(\mathbf{x}) = -\nabla_{\mathbf{x}} \cdot \mathbf{F}(\mathbf{x}) + \nabla_{\mathbf{x}} \cdot \mathbf{D} \nabla_{\mathbf{x}} = -\nabla_{\mathbf{x}} \cdot \hat{\mathbf{j}}_{\mathbf{x}}, \quad (3)$$

where we have defined the current operator $\hat{\mathbf{j}}_{\mathbf{x}} \equiv \mathbf{F}(\mathbf{x}) - \mathbf{D} \nabla_{\mathbf{x}}$. Note that all differential operators act on all functions to the right, e.g. $\nabla_{\mathbf{x}} \cdot \mathbf{F}(\mathbf{x})g(\mathbf{x}) = g(\mathbf{x})\nabla_{\mathbf{x}} \cdot \mathbf{F}(\mathbf{x}) + \mathbf{F}(\mathbf{x}) \cdot \nabla_{\mathbf{x}}g(\mathbf{x})$. Although the approach presented here is more general, we restrict our attention to (possibly non-equilibrium) steady states where the drift $\mathbf{F}(\mathbf{x})$ is sufficiently smooth and confining to assure the existence of a steady-state (invariant) density $p_s(\mathbf{x}) = \lim_{t \rightarrow \infty} G(\mathbf{x}, t|\mathbf{x}_0)$ and steady-state current $\mathbf{j}_s(\mathbf{x}) = \hat{\mathbf{j}}_{\mathbf{x}}p_s(\mathbf{x})$. The special case $\mathbf{j}_s(\mathbf{x}) = 0$ corresponds to equilibrium steady states. For systems that eventually evolve into a steady state we can rewrite the current operator as [31] (again the differential operator in $\nabla_{\mathbf{x}}p_s^{-1}(\mathbf{x})$ also acts on functions to the right if $\hat{\mathbf{j}}_{\mathbf{x}}$ is applied to a function)

$$\hat{\mathbf{j}}_{\mathbf{x}} = \mathbf{j}_s(\mathbf{x})p_s^{-1}(\mathbf{x}) - \mathbf{D}p_s(\mathbf{x})\nabla_{\mathbf{x}}p_s^{-1}(\mathbf{x}). \quad (4)$$

We will later also restrict the treatment to systems evolving from steady-state initial conditions, i.e. the initial condition $\mathbf{x}_{t=0}$ is drawn according to the density p_s .

We define the two fundamental additive dynamical functionals—time-integrated current and density—as

$$\begin{aligned} \mathbf{J}_t &= \int_{\tau=0}^{\tau=t} U(\mathbf{x}_{\tau}) \circ d\mathbf{x}_{\tau} \\ \rho_t &= \int_0^t V(\mathbf{x}_{\tau}) d\tau, \end{aligned} \quad (5)$$

with differentiable and square-integrable (real-valued) functions $U, V: \mathbb{R}^d \rightarrow \mathbb{R}$ and \circ denoting the Stratonovich integral [50, 55, 56]. These objects depend on the whole trajectory $[\mathbf{x}_{\tau}]_{0 \leq \tau \leq t}$ and are thus random functionals with non-trivial statistics. In the following we will derive an equation for the characteristic function of the joint distribution of $\mathbf{x}_t, \rho_t, \mathbf{J}_t$ via a Feynman-Kac approach which will then yield the moments (including variances and correlations) via a Dyson series. The formalism was already applied to the time-averaged density ρ_t/t (under the term of local/occupation time fraction) [9, 34, 57]. To do so, we need to derive a tilted Fokker-Planck equation, which we first do via Itô calculus and then, equivalently, via a functional calculus. Note that the tilted generator can also be found in the literature on large deviation theory [22, 23] (in this case obtained via the Feynman-Kac-Girsanov approach).

2.2. Tilting via Itô's lemma

We first derive a tilted the Fokker-Planck equation using Itô calculus. From the Itô-Stratonovich correction term $dU(\mathbf{x}_{\tau})d\mathbf{x}_{\tau}/2$ and $d\mathbf{x}_{\tau}d\mathbf{x}_{\tau}^T = 2\mathbf{D}d\tau$ (where $\mathbf{D} = \sigma\sigma^T/2$) we obtain from equations (1) and (5) the increments (curly brackets $\{\nabla \dots\}$ throughout denote that derivatives only act inside brackets)

$$\begin{aligned} d\mathbf{J}_\tau &= U(\mathbf{x}_\tau) \circ d\mathbf{x}_\tau = U(\mathbf{x}_\tau)d\mathbf{x}_\tau + \mathbf{D}\{\nabla_{\mathbf{x}}U\}(\mathbf{x}_\tau)d\tau \\ d\rho_\tau &= V(\mathbf{x}_\tau)d\tau. \end{aligned} \quad (6)$$

We use Itô's lemma [50] in d dimensions for a twice differentiable test function $f = f(\mathbf{x}_t, \rho_t, \mathbf{J}_t)$ and equations (1) and (6), to obtain

$$\begin{aligned} df &= \sum_{i=1}^d \frac{\partial f}{\partial x_i} dx_i^i + \frac{\partial f}{\partial \rho} d\rho_t + \sum_{i=1}^d \frac{\partial f}{\partial J_i} dJ_i^i \\ &\quad + \frac{1}{2} \sum_{i,j=1}^d \left(\frac{\partial^2 f}{\partial x_i \partial x_j} dx_i^i dx_j^j + \frac{\partial^2 f}{\partial J_i \partial J_j} dJ_i^i dJ_j^j + 2 \frac{\partial^2 f}{\partial x_i \partial J_j} dx_i^i dJ_j^j \right) \\ &= [(\nabla_{\mathbf{x}}f) + (\nabla_{\mathbf{J}}f)U(\mathbf{x}_t)][\mathbf{F}(\mathbf{x}_t)dt + \boldsymbol{\sigma}d\mathbf{W}_t] + (\nabla_{\mathbf{J}}f)\mathbf{D}\{\nabla_{\mathbf{x}}U\}(\mathbf{x}_t)dt + V(\mathbf{x}_t)\partial_\rho f dt \\ &\quad + (\nabla_{\mathbf{x}}^T \mathbf{D} \nabla_{\mathbf{x}} + U(\mathbf{x}_t)^2 \nabla_{\mathbf{J}}^T \mathbf{D} \nabla_{\mathbf{J}} + 2U(\mathbf{x}_t) \nabla_{\mathbf{x}}^T \mathbf{D} \nabla_{\mathbf{J}}) f dt. \end{aligned} \quad (7)$$

For the time derivative of f this gives

$$\begin{aligned} \frac{d}{dt} f(\mathbf{x}_t, \rho_t, \mathbf{J}_t) &= \left[\left(\mathbf{F} + \boldsymbol{\sigma} \frac{d\mathbf{W}_t}{dt} \right) (\nabla_{\mathbf{x}} + U \nabla_{\mathbf{J}}) + \{\nabla_{\mathbf{x}}U\} \mathbf{D} \nabla_{\mathbf{J}} \right. \\ &\quad \left. + V \partial_\rho + \nabla_{\mathbf{x}}^T \mathbf{D} \nabla_{\mathbf{x}} + U^2 \nabla_{\mathbf{J}}^T \mathbf{D} \nabla_{\mathbf{J}} + 2U \nabla_{\mathbf{x}}^T \mathbf{D} \nabla_{\mathbf{J}} \right] f(\mathbf{x}_t, \rho_t, \mathbf{J}_t). \end{aligned} \quad (8)$$

Following this formalism, we move towards a tilted Fokker-Planck equation [9, 34]. Using the conditional probability density $Q_t(\mathbf{x}, \rho, \mathbf{J}|\mathbf{x}_0)$ we may write (omitting \mathbf{x} dependence in \mathbf{F}, U, V for brevity) the evolution equation for $\langle f(\mathbf{x}_t, \rho_t, \mathbf{J}_t) \rangle_{\mathbf{x}_0}$, i.e. the expected value of $f(\mathbf{x}_t, \rho_t, \mathbf{J}_t)$ over the ensemble of paths propagating between \mathbf{x}_0 and \mathbf{x} in time t . Using equation (8) and integration by parts, we obtain (note that non-negative functions $V \geq 0$ imply $\rho \geq 0$, such that one would restrict the ρ -integration to $\int_0^\infty d\rho$ as in [34])

$$\begin{aligned} \frac{d}{dt} \langle f(\mathbf{x}_t, \rho_t, \mathbf{J}_t) \rangle_{\mathbf{x}_0} &= \int d^d x \int_{-\infty}^{\infty} d\rho \int d^d J f(\mathbf{x}, \rho, \mathbf{J}) \partial_t Q_t(\mathbf{x}, \rho, \mathbf{J}|\mathbf{x}_0) \\ &= \int d^d x \int_{-\infty}^{\infty} d\rho \int d^d J Q_t(\mathbf{x}, \rho, \mathbf{J}|\mathbf{x}_0) [\mathbf{F}(\nabla_{\mathbf{x}} + U \nabla_{\mathbf{J}}) + \{\nabla_{\mathbf{x}}U\} \mathbf{D} \nabla_{\mathbf{J}} \\ &\quad + V \partial_\rho + \nabla_{\mathbf{x}}^T \mathbf{D} \nabla_{\mathbf{x}} + U^2 \nabla_{\mathbf{J}}^T \mathbf{D} \nabla_{\mathbf{J}} + 2U \nabla_{\mathbf{x}}^T \mathbf{D} \nabla_{\mathbf{J}}] f(\mathbf{x}, \rho, \mathbf{J}) \\ &= \int d^d x \int_{-\infty}^{\infty} d\rho \int d^d J f(\mathbf{x}, \rho, \mathbf{J}) [-\nabla_{\mathbf{x}} \mathbf{F} - U \mathbf{F} \nabla_{\mathbf{J}} - \{\nabla_{\mathbf{x}}U\} \mathbf{D} \nabla_{\mathbf{J}} - V \partial_\rho \\ &\quad + \nabla_{\mathbf{x}}^T \mathbf{D} \nabla_{\mathbf{x}} + U^2 \nabla_{\mathbf{J}}^T \mathbf{D} \nabla_{\mathbf{J}} + 2U \nabla_{\mathbf{x}}^T \mathbf{D} \nabla_{\mathbf{J}}] Q_t(\mathbf{x}, \rho, \mathbf{J}|\mathbf{x}_0). \end{aligned} \quad (9)$$

Since the test function f is an arbitrary twice differentiable function, the resulting tilted Fokker-Planck equation reads

$$\partial_t Q_t(\mathbf{x}, \rho, \mathbf{J}|\mathbf{x}_0) = \hat{\mathcal{L}}_{\mathbf{x}, \rho, \mathbf{J}} Q_t(\mathbf{x}, \rho, \mathbf{J}|\mathbf{x}_0), \quad (10)$$

with the tilted Fokker-Planck operator¹

¹ For non-negative functions $V \geq 0$ an additional boundary term appears at $\rho=0$ upon partial integration in equation (9), leading to an extra term $-V(\mathbf{x})\delta(\rho)$ in equation (10) that ensures conservation of probability (see [34]).

$$\begin{aligned}
\hat{\mathcal{L}}_{\mathbf{x},\rho,\mathbf{J}} &= -\nabla_{\mathbf{x}} \cdot \mathbf{F}(\mathbf{x}) + \nabla_{\mathbf{x}}^T \mathbf{D} \nabla_{\mathbf{x}} - V(\mathbf{x}) \partial_{\rho} - U(\mathbf{x}) \mathbf{F}(\mathbf{x}) \cdot \nabla_{\mathbf{J}} \\
&\quad - \{ \nabla_{\mathbf{x}} U(\mathbf{x}) \}^T \mathbf{D} \nabla_{\mathbf{J}} + U(\mathbf{x})^2 \nabla_{\mathbf{J}}^T \mathbf{D} \nabla_{\mathbf{J}} + 2 \nabla_{\mathbf{J}}^T \mathbf{D} \nabla_{\mathbf{x}} U(\mathbf{x}) \\
&= -[\nabla_{\mathbf{x}} + U(\mathbf{x}) \nabla_{\mathbf{J}}] \mathbf{F}(\mathbf{x}) - V(\mathbf{x}) \partial_{\rho} + [\nabla_{\mathbf{x}} + U(\mathbf{x}) \nabla_{\mathbf{J}}]^T \mathbf{D} [\nabla_{\mathbf{x}} + U(\mathbf{x}) \nabla_{\mathbf{J}}]. \quad (11)
\end{aligned}$$

We see that the ρ dependence enters in standard Feynman-Kac form [9, 34], whereas the \mathbf{J} dependence enters less trivially and shifts the gradient operator $\nabla_{\mathbf{x}} \rightarrow \nabla_{\mathbf{x}} + U(\mathbf{x}) \nabla_{\mathbf{J}}$.

2.3. Tilting via functional calculus

We now re-derive the tilted Fokker-Planck operator in equation (11) using a functional calculus approach [51, 52] instead of the Itô calculus in the previous section. This shows that both alternative approaches are equivalent, as expected. We closely follow the derivation of the Fokker-Planck equation in [51] but for d -dimensional space and we generalize the approach to include the functionals defined in equation (5). The following approach is equivalent to a Stratonovich interpretation of stochastic calculus which is manifested in the convention $\int_0^t \delta(t') dt' = \int_0^t \delta(t-t') dt' = 1/2$ [51]. The white noise term $\mathbf{f}(\tau)$ with $\langle \mathbf{f}(\tau) \mathbf{f}(\tau')^T \rangle_s = 2\mathbf{D} \delta(\tau - \tau')$ in the Langevin equation (2) can be considered to be described by a path-probability measure [51]

$$P[\mathbf{f}] = N \exp \left[-\frac{1}{2} \int_0^t \mathbf{f}(\tau)^T \mathbf{D}^{-1} \mathbf{f}(\tau) d\tau \right], \quad (12)$$

with normalization constant N which may formally be problematic but always cancels out.

We now derive a tilted Fokker-Planck equation for the joint conditional density Q of \mathbf{x}_t and the functionals \mathbf{J}_t, ρ_t , as defined in equation (5), given a deterministic initial condition \mathbf{x}_0 at time $t=0$,

$$Q_t(\mathbf{x}, \rho, \mathbf{J} | \mathbf{x}_0) \equiv \int \mathcal{D}\mathbf{f} P[\mathbf{f}] \delta(\mathbf{x} - \mathbf{x}_t) \delta(\rho - \rho_t) \delta(\mathbf{J} - \mathbf{J}_t). \quad (13)$$

Note for the time derivatives that $\dot{\mathbf{J}}_t = U(\mathbf{x}_t) \dot{\mathbf{x}}_t$ and $\dot{\rho}_t = V(\mathbf{x}_t)$ to obtain (as a generalization of the calculation in [51] to dynamical functionals)

$$\begin{aligned}
\partial_t Q_t(\mathbf{x}, \rho, \mathbf{J} | \mathbf{x}_0) &= \partial_t \int \mathcal{D}\mathbf{f} P[\mathbf{f}] \delta(\mathbf{x} - \mathbf{x}_t) \delta(\rho - \rho_t) \delta(\mathbf{J} - \mathbf{J}_t) \\
&= \int \mathcal{D}\mathbf{f} P[\mathbf{f}] [-\nabla_{\mathbf{x}} \cdot \dot{\mathbf{x}}_t - \partial_{\rho} \dot{\rho}_t - \nabla_{\mathbf{J}} \cdot \dot{\mathbf{J}}_t] \delta(\mathbf{x} - \mathbf{x}_t) \delta(\rho - \rho_t) \delta(\mathbf{J} - \mathbf{J}_t) \\
&= \int \mathcal{D}\mathbf{f} P[\mathbf{f}] [-\nabla_{\mathbf{x}} \cdot [\mathbf{F}(\mathbf{x}_t) + \mathbf{f}(t)] - V(\mathbf{x}_t) \partial_{\rho} - U(\mathbf{x}_t) [\mathbf{F}(\mathbf{x}_t) + \mathbf{f}(t)] \cdot \nabla_{\mathbf{J}}] \\
&\quad \times \delta(\mathbf{x} - \mathbf{x}_t) \delta(\rho - \rho_t) \delta(\mathbf{J} - \mathbf{J}_t) \\
&= [-\nabla_{\mathbf{x}} \mathbf{F}(\mathbf{x}) - V(\mathbf{x}) \partial_{\rho} - U(\mathbf{x}) \mathbf{F}(\mathbf{x}) \cdot \nabla_{\mathbf{J}}] Q_t(\mathbf{x}, \rho, \mathbf{J} | \mathbf{x}_0) \\
&\quad - [\nabla_{\mathbf{x}} + U(\mathbf{x}) \nabla_{\mathbf{J}}] \cdot \int \mathcal{D}\mathbf{f} P[\mathbf{f}] \mathbf{f}(t) \delta(\mathbf{x} - \mathbf{x}_t) \delta(\rho - \rho_t) \delta(\mathbf{J} - \mathbf{J}_t). \quad (14)
\end{aligned}$$

The functional derivative of equation (12) reads [51]

$$\frac{\delta P[\mathbf{f}]}{\delta \mathbf{f}(t)} = -\frac{1}{2} \mathbf{D}^{-1} \mathbf{f}(t) P[\mathbf{f}], \quad (15)$$

which we use to obtain, via an integration by parts in $\delta \mathbf{f}(t)$,

$$\begin{aligned} - \int \mathcal{D}\mathbf{f} P[\mathbf{f}|\mathbf{f}(t)] \delta(\mathbf{x} - \mathbf{x}_t) \delta(\rho - \rho_t) \delta(\mathbf{J} - \mathbf{J}_t) &= 2\mathbf{D} \int \mathcal{D}\mathbf{f} \frac{\delta P[\mathbf{f}]}{\delta \mathbf{f}(t)} \delta(\mathbf{x} - \mathbf{x}_t) \delta(\rho - \rho_t) \delta(\mathbf{J} - \mathbf{J}_t) \\ &= -2\mathbf{D} \int \mathcal{D}\mathbf{f} P[\mathbf{f}] \frac{\delta}{\delta \mathbf{f}(t)} \delta(\mathbf{x} - \mathbf{x}_t) \delta(\rho - \rho_t) \delta(\mathbf{J} - \mathbf{J}_t). \end{aligned} \quad (16)$$

As before, differentials are understood to act on all functions to the right, i.e. $\frac{\delta}{\delta \mathbf{f}(t)}$ here acts on the full product of delta functions. We obtain

$$\begin{aligned} \frac{\delta}{\delta \mathbf{f}(t)} \delta(\mathbf{x} - \mathbf{x}_t) \delta(\rho - \rho_t) \delta(\mathbf{J} - \mathbf{J}_t) \\ = \left[-\nabla_{\mathbf{x}} \frac{\delta \mathbf{x}_t}{\delta \mathbf{f}(t)} - \partial_{\rho} \frac{\delta \rho_t}{\delta \mathbf{f}(t)} - \nabla_{\mathbf{J}} \frac{\delta \mathbf{J}_t}{\delta \mathbf{f}(t)} \right] \delta(\mathbf{x} - \mathbf{x}_t) \delta(\rho - \rho_t) \delta(\mathbf{J} - \mathbf{J}_t), \end{aligned} \quad (17)$$

and we use that $\delta \rho_t / \delta \mathbf{f}(t) = \mathbf{0}$, and $\delta \mathbf{x}_t / \delta \mathbf{f}(t) = \mathbf{1}/2$ [51] which implies $\delta \mathbf{J}_t / \delta \mathbf{f}(t) = U(\mathbf{x}_t) \mathbf{1}/2$, to get

$$\frac{\delta}{\delta \mathbf{f}(t)} \delta(\mathbf{x} - \mathbf{x}_t) \delta(\rho - \rho_t) \delta(\mathbf{J} - \mathbf{J}_t) = \frac{1}{2} [-\nabla_{\mathbf{x}} - U(\mathbf{x}_t) \nabla_{\mathbf{J}}] \delta(\mathbf{x} - \mathbf{x}_t) \delta(\rho - \rho_t) \delta(\mathbf{J} - \mathbf{J}_t). \quad (18)$$

Plugging equation (18) first into equation (16) and then into equation (14) yields the tilted Fokker-Planck equation for the joint conditional density

$$\begin{aligned} \partial_t Q_t(\mathbf{x}, \rho, \mathbf{J} | \mathbf{x}_0) &= \left[-\nabla_{\mathbf{x}} \mathbf{F}(\mathbf{x}) - V(\mathbf{x}) \partial_{\rho} - U(\mathbf{x}) \mathbf{F}(\mathbf{x}) \nabla_{\mathbf{J}} \right. \\ &\quad \left. + [\nabla_{\mathbf{x}} + U(\mathbf{x}) \nabla_{\mathbf{J}}]^T \mathbf{D} [\nabla_{\mathbf{x}} + U(\mathbf{x}) \nabla_{\mathbf{J}}] \right] Q_t(\mathbf{x}, \rho, \mathbf{J} | \mathbf{x}_0). \end{aligned} \quad (19)$$

Note that equation (19) fully agrees with equation (11) derived via Itô calculus thus establishing the announced equivalence of the two approaches.

3. Steady-state covariance via dyson expansion of the tilted propagator

In this section we employ the tilted Fokker-Planck equation (19) to derive results for the mean value and (co)variances of time-integrated densities and currents. These follow as derivatives of the characteristic function evaluated at zero, and it thus suffices to treat the tilt as a perturbation of the ‘bare’ generator (see [34]). The derivation is based on a Dyson expansion of the exponential of a Fourier-transformed tilted generator (i.e. tilted Fokker-Planck operator). Therefore, consider a one-dimensional Fourier variable ν and a d -dimensional Fourier variable $\boldsymbol{\omega} = (\omega_1, \dots, \omega_d)$ and define the Fourier transform of $Q_t(\mathbf{x}, \rho, \mathbf{J} | \mathbf{x}_0)$ as

$$\tilde{Q}_t(\mathbf{x}, \nu, \boldsymbol{\omega} | \mathbf{x}_0) \equiv \int_{-\infty}^{\infty} d\rho \int d^d J Q_t(\mathbf{x}, \rho, \mathbf{J} | \mathbf{x}_0) \exp(-i\nu\rho - i\boldsymbol{\omega} \cdot \mathbf{J}). \quad (20)$$

In the case $V \geq 0$ where $\rho \geq 0$ one would instead take the Laplace transform in the ρ -coordinate, see [34]. Recall the (untilted) Fokker-Planck operator $\hat{L}(\mathbf{x}) = -\nabla_{\mathbf{x}} \cdot \hat{\mathbf{j}}_{\mathbf{x}}$ with the current operator $\hat{\mathbf{j}}_{\mathbf{x}} = \mathbf{F}(\mathbf{x}) - \mathbf{D} \nabla_{\mathbf{x}}$ from equation (3). The Fourier transform of the tilted Fokker-Planck operator in equations (11) and (19) reads

$$\begin{aligned} \hat{L}(\mathbf{x}, \nu, \boldsymbol{\omega}) &= \hat{L}(\mathbf{x}) - i\nu V(\mathbf{x}) - i\boldsymbol{\omega}^T \cdot \hat{\mathbf{L}}^U(\mathbf{x}) - U(\mathbf{x})^2 \boldsymbol{\omega}^T \mathbf{D} \boldsymbol{\omega}, \\ \hat{\mathbf{L}}^U(\mathbf{x}) &\equiv U(\mathbf{x}) \hat{\mathbf{j}}_{\mathbf{x}} - \mathbf{D} \nabla_{\mathbf{x}} U(\mathbf{x}). \end{aligned} \quad (21)$$

As always, the differential operators act on all functions to the right unless written inside curly brackets, i.e. $\nabla_{\mathbf{x}} U(\mathbf{x}) = \{\nabla_{\mathbf{x}} U(\mathbf{x})\} + U(\mathbf{x}) \nabla_{\mathbf{x}}$. Note that whereas we obtained the tilted generator directly and only subsequently Fourier transformed it, there are also approaches that directly target the Fourier image of the tilted generator (see e.g. [58]). Compared to the tilt of the density (i.e. the ν -term; see also [34]), the tilt corresponding to the current observable (ω -terms) involves more terms and even a term that is second order in ω . The second order term occurs since $(d\mathbf{W}_\tau)^2 \sim d\tau$ and therefore (in contrast to $d\tau d\mathbf{W}_\tau$ and $d\tau^2$) contributes in the tilting of the generator.

We now restrict our attention to dynamics starting in the steady state p_s and denote the average over an ensemble over paths propagating from the steady state by $\langle \cdot \rangle_s$. Extensions of the formalism to any initial distribution are straightforward and introduce additional transient terms. For the derivation of the moments of ρ_t and \mathbf{J}_t , we introduce and expand the characteristic function (also known as moment-generating function)

$$\tilde{\mathcal{P}}_t^{\rho\mathbf{J}}(\nu, \omega | p_s) \equiv \langle e^{-i\nu\rho_t - i\omega\cdot\mathbf{J}_t} \rangle_s = 1 - i\nu \langle \rho_t \rangle_s - i\omega \cdot \langle \mathbf{J}_t \rangle_s - \nu\omega \cdot \langle \rho_t \mathbf{J}_t \rangle_s + O(\omega^2, \nu^2). \quad (22)$$

This expansion in ν, ω will now be compared to the Dyson expansion of the exponential of equation (21) which yields expressions for $\langle \rho_t \rangle_s, \langle \mathbf{J}_t \rangle_s, \langle \rho_t \mathbf{J}_t \rangle_s$ by comparing individual orders.

The Dyson expansion allows to expand for small $|\nu|, |\omega|$ (see also [34])

$$\begin{aligned} e^{\hat{\mathcal{L}}(\mathbf{x}_1, \nu, \omega)t} &= 1 - i \int_0^t dt_1 e^{\hat{\mathcal{L}}(\mathbf{x}_1)(t-t_1)} \left[\nu V(\mathbf{x}_1) + \omega^T \cdot \hat{\mathbf{L}}^U(\mathbf{x}_1) \right] e^{\hat{\mathcal{L}}(\mathbf{x}_1)t_1} \\ &\quad - \int_0^t dt_2 \int_0^{t_2} dt_1 e^{\hat{\mathcal{L}}(\mathbf{x}_1)(t-t_2)} \left[\nu V(\mathbf{x}_1) + \omega^T \cdot \hat{\mathbf{L}}^U(\mathbf{x}_1) \right] e^{\hat{\mathcal{L}}(\mathbf{x}_1)(t_2-t_1)} \\ &\quad \times \left[\nu V(\mathbf{x}_1) + \omega^T \cdot \hat{\mathbf{L}}^U(\mathbf{x}_1) \right] e^{\hat{\mathcal{L}}(\mathbf{x}_1)t_1} + O(\omega^2, \nu^2). \end{aligned} \quad (23)$$

Using that the first propagation only differs from 1 by total derivatives (recall $\hat{\mathcal{L}}(\mathbf{x}) = -\nabla_{\mathbf{x}} \cdot \hat{\mathbf{j}}_{\mathbf{x}}$), and using for the last propagation term $e^{\hat{\mathcal{L}}(\mathbf{x}_1)t_1} p_s(\mathbf{x}_1) = p_s(\mathbf{x}_1)$, we obtain

$$\begin{aligned} \tilde{\mathcal{P}}_t^{\rho\mathbf{J}}(\nu, \omega | p_s) &= \int d^d x_1 e^{\hat{\mathcal{L}}(\mathbf{x}_1, \nu, \omega)t} p_s(\mathbf{x}_1) = 1 - i \int d^d x_1 \int_0^t dt_1 \left[\nu V(\mathbf{x}_1) + \omega^T \cdot \hat{\mathbf{L}}^U(\mathbf{x}_1) \right] p_s(\mathbf{x}_1) \\ &\quad - \sum_{l,m=1}^d \int d^d x_1 \int_0^t dt_2 \int_0^{t_2} dt_1 \left[\nu V(\mathbf{x}_1) + \omega^T \cdot \hat{\mathbf{L}}^U(\mathbf{x}_1) \right] e^{\hat{\mathcal{L}}(\mathbf{x}_1)(t_2-t_1)} \\ &\quad \times \left[\nu V(\mathbf{x}_1) + \omega^T \cdot \hat{\mathbf{L}}^U(\mathbf{x}_1) \right] p_s(\mathbf{x}_1) + O(\omega^2, \nu^2). \end{aligned} \quad (24)$$

We substitute the one-step propagation by the conditional density $G(\mathbf{x}_2, t | \mathbf{x}_1) = e^{\hat{\mathcal{L}}(\mathbf{x}_1)t} \delta(\mathbf{x}_2 - \mathbf{x}_1)$ [56, 59],

$$\int d^d x_1 f(\mathbf{x}_1) e^{\hat{\mathcal{L}}(\mathbf{x}_1)(t_2-t_1)} g(\mathbf{x}_1) = \int d^d x_1 \int d^d x_2 f(\mathbf{x}_2) G(\mathbf{x}_2, t_2 - t_1 | \mathbf{x}_1) g(\mathbf{x}_1), \quad (25)$$

which yields

$$\begin{aligned} \tilde{\mathcal{P}}_t^{\rho\mathbf{J}}(\nu, \omega | p_s) &= 1 - i \int d^d x_1 \int_0^t dt_1 \left[\nu V(\mathbf{x}_1) + \omega^T \cdot \hat{\mathbf{L}}^U(\mathbf{x}_1) \right] p_s(\mathbf{x}_1) \\ &\quad - \int d^d x_1 \int d^d x_2 \int_0^t dt_2 \int_0^{t_2} dt_1 \left[\nu V(\mathbf{x}_2) + \omega^T \cdot \hat{\mathbf{L}}^U(\mathbf{x}_2) \right] \\ &\quad \times G(\mathbf{x}_2, t_2 - t_1 | \mathbf{x}_1) \left[\nu V(\mathbf{x}_1) + \omega^T \cdot \hat{\mathbf{L}}^U(\mathbf{x}_1) \right] p_s(\mathbf{x}_1) + O(\omega^2, \nu^2). \end{aligned} \quad (26)$$

This concludes the expansion of the exponential of the Fourier transformed tilted generator. Now, by comparing the definition and expansion of the characteristic function equation (22) with the result equation (26) from the Dyson expansion, we obtain the moments and correlations of the functionals $\mathbf{J}_t = \int_{\tau=0}^t U(\mathbf{x}_\tau) \circ d\mathbf{x}_\tau$ and $\rho_t = \int_0^t V(\mathbf{x}_\tau) d\tau$.

Note that the first moments (i.e. the mean values for steady-state initial conditions) can also be obtained directly [18, 31] but we obtain them here by comparing the terms of order ν and ω in equations (22) and (26),

$$\begin{aligned} \langle \rho_t \rangle_s &= \int_0^t dt_1 \int d^d x_1 V(\mathbf{x}_1) p_s(\mathbf{x}_1) = t \int d^d x_1 V(\mathbf{x}_1) p_s(\mathbf{x}_1) \\ \langle \mathbf{J}_t \rangle_s &= t \int d^d x_1 [U(\mathbf{x}_1) \hat{\mathbf{j}}_{\mathbf{x}_1} - \mathbf{D} \nabla_{\mathbf{x}_1} U(\mathbf{x}_1)] p_s(\mathbf{x}_1) = t \int d^d x_1 U(\mathbf{x}_1) \mathbf{j}_s(\mathbf{x}_1), \end{aligned} \quad (27)$$

where $\nabla_{\mathbf{x}_1} U(\mathbf{x}_1) p_s(\mathbf{x}_1)$ vanishes after integration by parts and $\mathbf{j}_s(\mathbf{x}_1) \equiv \hat{\mathbf{j}}_{\mathbf{x}_1} p_s(\mathbf{x}_1)$ is the steady-state current.

By comparing the terms of order $\nu\omega$ in equations (22) and (26) we have for the steady-state expectation $\langle \mathbf{J}_t \rho_t \rangle_s$ that

$$\begin{aligned} \langle \mathbf{J}_t \rho_t \rangle_s &= \int_0^t dt_2 \int_0^{t_2} dt_1 \int d^d x_1 \int d^d x_2 \\ &\quad \times \left[\hat{\mathbf{L}}^U(\mathbf{x}_2) G(\mathbf{x}_2, t_2 - t_1 | x_1) V(\mathbf{x}_1) + V(\mathbf{x}_2) G(\mathbf{x}_2, t_2 - t_1 | x_1) \mathbf{L}_U(\mathbf{x}_1) \right] p_s(\mathbf{x}_1) \\ &= \int_0^t dt_2 \int_0^{t_2} dt_1 \int d^d x_1 \int d^d x_2 \left[U(\mathbf{x}_2) \hat{\mathbf{j}}_{\mathbf{x}_2} G(\mathbf{x}_2, t_2 - t_1 | \mathbf{x}_1) V(\mathbf{x}_1) \right. \\ &\quad \left. + V(\mathbf{x}_2) G(\mathbf{x}_2, t_2 - t_1 | \mathbf{x}_1) [U(\mathbf{x}_1) \hat{\mathbf{j}}_{\mathbf{x}_1} - \mathbf{D} \nabla_{\mathbf{x}_1} U(\mathbf{x}_1)] \right] p_s(\mathbf{x}_1). \end{aligned} \quad (28)$$

We note that for any function f the following identity holds

$$\int_0^t dt_2 \int_0^{t_2} dt_1 f(t_2 - t_1) = \int_0^t dt' (t - t') f(t'), \quad (29)$$

and further introduce the shorthand notation

$$\hat{\mathcal{I}}_{\mathbf{xy}}^t[\dots] = \int_0^t dt' (t - t') \int d^d x_1 \int d^d x_2 U(\mathbf{x}_1) V(\mathbf{x}_2) [\dots]. \quad (30)$$

Moreover, we define the joint density $P_{\mathbf{y}}(\mathbf{x}, t) \equiv G(\mathbf{x}, t | \mathbf{y}) p_s(\mathbf{y})$ and following [31] introduce the dual-reversed current operator $\hat{\mathbf{j}}_{\mathbf{x}}^\dagger \equiv \mathbf{j}_s(\mathbf{x}) / p_s(\mathbf{x}) + \mathbf{D} p_s(\mathbf{x}) \nabla_{\mathbf{x}} p_s^{-1}(\mathbf{x}) = -\hat{\mathbf{j}}_{\mathbf{x}}(\mathbf{j}_s \rightarrow -\mathbf{j}_s)$. With these notations, using integration by parts, and by relabeling $\mathbf{x}_1 \leftrightarrow \mathbf{x}_2$ in one term, we rewrite equation (28) to obtain for the correlation, reproducing the main result of [30, 31],

$$\begin{aligned} \langle \mathbf{J}_t \rho_t \rangle_s - \langle \mathbf{J}_t \rangle_s \langle \rho_t \rangle_s &= \hat{\mathcal{I}}_{\mathbf{xy}}^t \left[\hat{\mathbf{j}}_{\mathbf{x}_1} P_{\mathbf{x}_2}(\mathbf{x}_1, t') + \mathbf{j}_s(\mathbf{x}_1) p_s^{-1}(\mathbf{x}_1) P_{\mathbf{x}_1}(\mathbf{x}_2, t') \right. \\ &\quad \left. + \mathbf{D} p_s(\mathbf{x}_1) \nabla_{\mathbf{x}_1} p_s(\mathbf{x}_1)^{-1} P_{\mathbf{x}_1}(\mathbf{x}_2, t') \right] - \langle \mathbf{J}_t \rangle_s \langle \rho_t \rangle_s \\ &= \hat{\mathcal{I}}_{\mathbf{xy}}^t \left[\hat{\mathbf{j}}_{\mathbf{x}_1} P_{\mathbf{x}_2}(\mathbf{x}_1, t') + \hat{\mathbf{j}}_{\mathbf{x}_1}^\dagger P_{\mathbf{x}_1}(\mathbf{x}_2, t') - 2\mathbf{j}_s(\mathbf{x}_1) p_s(\mathbf{x}_2) \right]. \end{aligned} \quad (31)$$

We will discuss this result below, but first derive analogous results for (co)variances of densities and currents, respectively.

Instead of obtaining $\langle \rho_t^2 \rangle_s$ from the ν^2 order in equation (26) we here consider a generalization to two densities, $\rho_t = \int_0^t V(\mathbf{x}_\tau) d\tau$ and $\rho'_t = \int_0^t U(\mathbf{x}_\tau) d\tau$. The Fourier-transformed tilted generator in equation (21) with Fourier variables ν, ν' corresponding to ρ_t, ρ'_t is obtained

equivalently and gives $\hat{\mathcal{L}}(\mathbf{x}, \nu, \nu') = \hat{L}(\mathbf{x}) - i\nu V(\mathbf{x}) - i\nu' U(\mathbf{x})$. The related term in the Dyson series (by an adaption of equation (26) including $\nu' U$) becomes $[\nu V(\mathbf{x}_2) + \nu' U(\mathbf{x}_2)]G(\mathbf{x}_2, t_2 - t_1 | \mathbf{x}_1)[\nu V(\mathbf{x}_1) + \nu' U(\mathbf{x}_1)]p_s(\mathbf{x}_1)$ (see also [34]). By comparison with the characteristic function in equation (22) including ρ'_t , one obtains the known result [9, 34],

$$\langle \rho_t \rho'_t \rangle_s - \langle \rho_t \rangle_s \langle \rho'_t \rangle_s = \hat{\mathcal{I}}_{\mathbf{xy}}^t [P_{\mathbf{x}_2}(\mathbf{x}_1, t') + P_{\mathbf{x}_1}(\mathbf{x}_2, t') - 2p_s(\mathbf{x}_1)p_s(\mathbf{x}_2)]. \quad (32)$$

For $U = V$ this becomes the variance of ρ_t which can also be obtained from the order ν^2 in equations (22) and (26).

To obtain the current covariance, we accordingly require a tilted generator with two Fourier variables ω, ω' corresponding to $\mathbf{J}_t = \int_{\tau=0}^{\tau=t} U(\mathbf{x}_\tau) \circ d\mathbf{x}_\tau$ and $\mathbf{J}'_t = \int_{\tau=0}^{\tau=t} V(\mathbf{x}_\tau) \circ d\mathbf{x}_\tau$, which can, by the same formalism, be derived as

$$\begin{aligned} \hat{\mathcal{L}}(\mathbf{x}, \omega, \omega') &= \hat{L}(\mathbf{x}) - i\omega^T \cdot \hat{\mathbf{L}}^U(\mathbf{x}) - i\omega'^T \cdot \hat{\mathbf{L}}^V(\mathbf{x}) - U(\mathbf{x})^2 \omega^T \mathbf{D} \omega - V(\mathbf{x})^2 \omega'^T \mathbf{D} \omega' \\ &\quad - 2U(\mathbf{x})V(\mathbf{x})\omega^T \mathbf{D} \omega' \\ \hat{\mathbf{L}}^V(\mathbf{x}) &\equiv V(\mathbf{x})\mathbf{j}_x - \mathbf{D} \nabla_x V(\mathbf{x}). \end{aligned} \quad (33)$$

The Dyson series (by adapting equation (26)) based on $\hat{\mathcal{L}}(\mathbf{x}, \omega, \omega')$ for two currents \mathbf{J}, \mathbf{J}' reads

$$\begin{aligned} \tilde{\mathcal{P}}_{\mathbf{J}\mathbf{J}'}^t(\omega, \omega' | p_s) &= 1 - \int d^d x_1 \int_0^t dt_1 \left[i\omega^T \cdot \hat{\mathbf{L}}^U(\mathbf{x}_1) + i\omega'^T \cdot \hat{\mathbf{L}}^V(\mathbf{x}_1) + 2U(\mathbf{x}_1)V(\mathbf{x}_1)\omega^T \mathbf{D} \omega' \right] p_s(\mathbf{x}_1) \\ &\quad + \int d^d x_1 \int d^d x_2 \int_0^t dt_1 \int_0^{t_1} dt_2 \left[i\omega^T \cdot \hat{\mathbf{L}}^U(\mathbf{x}_2) + i\omega'^T \cdot \hat{\mathbf{L}}^V(\mathbf{x}_2) \right] \\ &\quad \times G(\mathbf{x}_2, t_2 - t_1 | \mathbf{x}_1) \left[i\omega^T \cdot \hat{\mathbf{L}}^U(\mathbf{x}_1) + i\omega'^T \cdot \hat{\mathbf{L}}^V(\mathbf{x}_1) \right] p_s(\mathbf{x}_1) + O(\omega^2, \omega'^2). \end{aligned} \quad (34)$$

The expectation value of the product of current components $\langle J_{t,n} J'_{t,m} \rangle_s$ is given by the terms that are linear in $\omega_n \omega'_m$, i.e. (recall $D_{nm} = D_{mn}$)

$$\begin{aligned} \langle J_{t,n} J'_{t,m} \rangle_s &= 2t D_{nm} \int d^d x_1 U(\mathbf{x}_1) V(\mathbf{x}_1) p_s(\mathbf{x}_1) + \int_0^t dt' (t - t') \int d^d x_1 \int d^d x_2 \\ &\quad \times \left[\hat{L}_n^U(\mathbf{x}_2) G(\mathbf{x}_2, t' | \mathbf{x}_1) \cdot \hat{L}_m^V(\mathbf{x}_1) p_s(\mathbf{x}_1) + \hat{L}_m^V(\mathbf{x}_2) G(\mathbf{x}_2, t' | \mathbf{x}_1) \cdot \hat{L}_n^U(\mathbf{x}_1) p_s(\mathbf{x}_1) \right]. \end{aligned} \quad (35)$$

We denote by $\hat{=}$ equality up to gradient terms that vanish upon integration to write

$$\begin{aligned} &\hat{L}_n^U(\mathbf{x}_2) G(\mathbf{x}_2, t' | \mathbf{x}_1) \cdot \hat{L}_m^V(\mathbf{x}_1) p_s(\mathbf{x}_1) \\ &\hat{=} U(\mathbf{x}_2) \hat{\mathbf{j}}_{\mathbf{x}_2, n} G(\mathbf{x}_2, t' | \mathbf{x}_1) \\ &\quad \times \left[V(\mathbf{x}_1) \mathbf{j}_s(\mathbf{x}_1) p_s^{-1}(\mathbf{x}_1) - p_s(\mathbf{x}_1) \mathbf{D} \nabla_{\mathbf{x}_1} p_s(\mathbf{x}_1)^{-1} - \mathbf{D} \nabla_{\mathbf{x}_1} V(\mathbf{x}_1) \right]_m p_s(\mathbf{x}_1) \\ &\hat{=} U(\mathbf{x}_2) V(\mathbf{x}_1) \hat{\mathbf{j}}_{\mathbf{x}_2, n} [\mathbf{j}_s(\mathbf{x}_1) p_s^{-1}(\mathbf{x}_1) + p_s(\mathbf{x}_1) \mathbf{D} \nabla_{\mathbf{x}_1} p_s^{-1}(\mathbf{x}_1)]_m G(\mathbf{x}_2, t' | \mathbf{x}_1) p_s(\mathbf{x}_1) \\ &= U(\mathbf{x}_2) V(\mathbf{x}_1) \hat{\mathbf{j}}_{\mathbf{x}_2, n} \hat{\mathbf{j}}_{\mathbf{x}_1, m}^\dagger P_{\mathbf{x}_1}(\mathbf{x}_2, t). \end{aligned} \quad (36)$$

Inserting this into equation (35), and relabeling in one term $\mathbf{x}_1 \leftrightarrow \mathbf{x}_2$ we obtain for the nm -element of the current covariance matrix

$$\begin{aligned} \langle J_{t,n} J'_{t,m} \rangle_s - \langle J_{t,n} \rangle_s \langle J'_{t,m} \rangle_s &= 2t D_{nm} \int d^d x_1 U(\mathbf{x}_1) V(\mathbf{x}_1) p_s(\mathbf{x}_1) \\ &\quad + \hat{\mathcal{I}}_{\mathbf{xy}}^t \left[\hat{\mathbf{j}}_{\mathbf{x}_1, m} \hat{\mathbf{j}}_{\mathbf{x}_2, n}^\dagger P_{\mathbf{x}_2}(\mathbf{x}_1, t') + \hat{\mathbf{j}}_{\mathbf{x}_2, n} \cdot \hat{\mathbf{j}}_{\mathbf{x}_1, m}^\dagger P_{\mathbf{x}_1}(\mathbf{x}_2, t') \right]. \end{aligned} \quad (37)$$

This reproduces and slightly generalizes the main result of [30, 31] where the diagonal elements ($m = n$) of the covariance matrix were derived. This result for the current covariance matrix and equation (31) for the current-density correlation are the natural generalizations of the density-density covariance equation (32), as described in detail in [30, 31], with the additional $2tD_{nm}$ -term in equation (37) arising from the $(d\mathbf{W}_\tau)^2$ contribution in $J_{t,n}J'_{t,m}$ manifested in the term $-2U(\mathbf{x})V(\mathbf{x})\omega^T\mathbf{D}\omega'$ in the tilted generator in equation (33). While the density-density covariance equation (32) only depends on integration over all paths from \mathbf{x}_1 to \mathbf{x}_2 (and vice versa) in time t' via $P_{\mathbf{x}_1}(\mathbf{x}_2, t')$, the current-density correlation equation (31) instead involves $\hat{\mathbf{j}}_{\mathbf{x}_1} P_{\mathbf{x}_2}(\mathbf{x}_1, t')$ and $\hat{\mathbf{j}}_{\mathbf{x}_1}^\dagger P_{\mathbf{x}_1}(\mathbf{x}_2, t')$ which describe currents at the final- and initial-points, respectively [31]. This notion is further extended in the result equation (37) where $\hat{\mathbf{j}}_{\mathbf{x}_2, n} \hat{\mathbf{j}}_{\mathbf{x}_1, m}^\dagger P_{\mathbf{x}_1}(\mathbf{x}_2, t')$ corresponds to products of components of displacements along individual trajectories from \mathbf{x}_1 to \mathbf{x}_2 [30].

4. Conclusion

We employed a Feynman-Kac approach to derive moments and correlations of dynamical functionals of diffusive paths—the time-integrated densities and currents. We presented two different but equivalent approaches to tilting the generator—Itô and functional calculus. These two approaches illustrate how one can freely choose between Itô and functional calculus to derive results on dynamical functionals. In particular, both approaches are accessible without further technical mathematical concepts such as the Cameron-Martin-Girsanov theorem that is often used in the study of tilted generators. Our methodological advance thus provides a flexible repertoire of easily accessible methods that will hopefully prove useful in future studies of related problems.

The derivation of the moments and correlations based on the tilted generator reproduces results with important implications for stochastic thermodynamics and large deviation theory, in particular for the physical and mathematical role of coarse graining [30, 31], and thereby displays how the tilted generator yields results on the statistics of dynamical functionals, even beyond the large deviation limit.

Data availability statement

No new data were created or analysed in this study.

Acknowledgments

Financial support from Studienstiftung des Deutschen Volkes (to C D) and the German Research Foundation (DFG) through the Emmy Noether Program GO 2762/1-2 (to A G) is gratefully acknowledged.

ORCID iDs

Cai Dieball  <https://orcid.org/0000-0002-0011-2358>

Aljaž Godec  <https://orcid.org/0000-0003-1888-6666>

References

- [1] Derrida B and Lebowitz J L 1998 Exact large deviation function in the asymmetric exclusion process *Phys. Rev. Lett.* **80** 209
- [2] Bodineau T and Derrida B 2004 Current fluctuations in nonequilibrium diffusive systems: an additivity principle *Phys. Rev. Lett.* **92** 180601
- [3] Lebowitz J L and Spohn H 1999 A Gallavotti-Cohen-type symmetry in the large deviation functional for stochastic dynamics *J. Stat. Phys.* **95** 333
- [4] Garrahan J P, Jack R L, Lecomte V, Pitard E, van Duijvendijk K and van Wijland F 2009 First-order dynamical phase transition in models of glasses: an approach based on ensembles of histories *J. Phys. A* **42** 075007
- [5] Bodineau T and Toninelli C 2012 Activity phase transition for constrained dynamics *Commun. Math. Phys.* **311** 357
- [6] Maes C 2020 Frenesy: time-symmetric dynamical activity in nonequilibria *Phys. Rep.* **850** 1
- [7] Jack R L, Thompson I R and Sollich P 2015 Hyperuniformity and phase separation in biased ensembles of trajectories for diffusive systems *Phys. Rev. Lett.* **114** 060601
- [8] Vanicat M, Bertin E, Lecomte V and Ragoucy E 2021 Mapping current and activity fluctuations in exclusion processes: consequences and open questions *SciPost Phys.* **10** 028
- [9] Kac M 1949 On distributions of certain Wiener functionals *Trans. Am. Math. Soc.* **65** 1
- [10] Darling D A and Kac M 1957 On occupation times for Markoff processes *Trans. Am. Math. Soc.* **84** 444
- [11] Aghion E, Kessler D A and Barkai E 2019 From non-normalizable Boltzmann-Gibbs statistics to infinite-ergodic theory *Phys. Rev. Lett.* **122** 010601
- [12] Carmi S and Barkai E 2011 Fractional Feynman-Kac equation for weak ergodicity breaking *Phys. Rev. E* **84** 061104
- [13] Majumdar S N and Comtet A 2002 Local and occupation time of a particle diffusing in a random medium *Phys. Rev. Lett.* **89** 060601
- [14] Majumdar S N and Dean D S 2002 Exact occupation time distribution in a non-Markovian sequence and its relation to spin glass models *Phys. Rev. E* **66** 041102
- [15] Majumdar S N 2005 Brownian functionals in physics and computer science *Curr. Sci.* **89** 2075
- [16] Bray A J, Majumdar S N and Schehr G 2013 Persistence and first-passage properties in nonequilibrium systems *Adv. Phys.* **62** 225
- [17] Bel G and Barkai E 2005 Weak ergodicity breaking in the continuous-time random walk *Phys. Rev. Lett.* **94** 240602
- [18] Maes C, Netočný K and Wynants B 2008 Steady state statistics of driven diffusions *Phys. A* **387** 2675
- [19] Touchette H 2009 The large deviation approach to statistical mechanics *Phys. Rep.* **478** 1
- [20] Kusuoka S, Kuwada K and Tamura Y 2009 Large deviation for stochastic line integrals as L^p - currents *Probab. Theory Relat. Fields* **147** 649
- [21] Chetrite R and Touchette H 2013 Nonequilibrium microcanonical and canonical ensembles and their equivalence *Phys. Rev. Lett.* **111** 120601
- [22] Chetrite R and Touchette H 2014 Nonequilibrium Markov processes conditioned on large deviations *Ann. Henri Poincaré* **16** 2005
- [23] Barato A C and Chetrite R 2015 A formal view on level 2.5 large deviations and fluctuation relations *J. Stat. Phys.* **160** 1154
- [24] Hoppenau J, Nickelsen D and Engel A 2016 Level 2 and level 2.5 large deviation functionals for systems with and without detailed balance *New J. Phys.* **18** 083010
- [25] Touchette H 2018 Introduction to dynamical large deviations of Markov processes *Phys. A* **504** 5
- [26] Mallmin E, du Buisson J and Touchette H 2021 Large deviations of currents in diffusions with reflective boundaries *J. Phys. A: Math. Theor.* **54** 295001
- [27] Monthus C 2021 Inference of Markov models from trajectories via large deviations at level 2.5 with applications to random walks in disordered media *J. Stat. Mech.* **063211**
- [28] Dechant A and Sasa S-I 2021 Improving thermodynamic bounds using correlations *Phys. Rev. X* **11** 041061
- [29] Dechant A and Sasa S-I 2021 Continuous time reversal and equality in the thermodynamic uncertainty relation *Phys. Rev. Res.* **3** 042012
- [30] Dieball C and Godec A 2022 Mathematical, thermodynamical and experimental necessity for coarse graining empirical densities and currents in continuous space *Phys. Rev. Lett.* **129** 140601

- [31] Dieball C and Godec A 2022 Coarse graining empirical densities and currents in continuous-space steady states *Phys. Rev. Res.* **4** 033243
- [32] Rebenshtok A and Barkai E 2008 Weakly non-ergodic statistical physics *J. Stat. Phys.* **133** 565
- [33] Burov S, Jeon J-H, Metzler R and Barkai E 2011 Single particle tracking in systems showing anomalous diffusion: the role of weak ergodicity breaking *Phys. Chem. Chem. Phys.* **13** 1800
- [34] Lapolla A, Hartich D and Godec A 2020 Spectral theory of fluctuations in time-average statistical mechanics of reversible and driven systems *Phys. Rev. Res.* **2** 043084
- [35] Coghi F, Chetrite R and Touchette H 2021 Role of current fluctuations in nonreversible samplers *Phys. Rev. E* **103** 062142
- [36] Bertini L, Sole A D, Gabrielli D, Jona-Lasinio G and Landim C 2005 Current fluctuations in stochastic lattice gases *Phys. Rev. Lett.* **94** 030601
- [37] Bertini L, Sole A D, Gabrielli D, Jona-Lasinio G and Landim C 2015 Macroscopic fluctuation theory *Rev. Mod. Phys.* **87** 593–636
- [38] Derrida B 2007 Non-equilibrium steady states: fluctuations and large deviations of the density and of the current *J. Stat. Mech.* **07023**
- [39] Seifert U 2018 Stochastic thermodynamics: from principles to the cost of precision *Phys. A* **504** 176
- [40] Koyuk T and Seifert U 2020 Thermodynamic uncertainty relation for time-dependent driving *Phys. Rev. Lett.* **125** 260604
- [41] Pietzonka P, Barato A C and Seifert U 2016 Universal bounds on current fluctuations *Phys. Rev. E* **93** 052145
- [42] Seifert U 2005 Entropy production along a stochastic trajectory and an integral fluctuation theorem *Phys. Rev. Lett.* **95** 040602
- [43] Pigolotti S, Neri I, Roldán E and Jülicher F 2017 Generic properties of stochastic entropy production *Phys. Rev. Lett.* **119** 140604
- [44] Seifert U 2012 Stochastic thermodynamics, fluctuation theorems and molecular machines *Rep. Prog. Phys.* **75** 126001
- [45] Dell’Antonio G 2016 Lecture 6: Lie-Trotter formula, Wiener process, Feynman-Kac formula *Lectures on the Mathematics of Quantum Mechanics II: Selected Topics* p 133
- [46] Ehrhardt G C M A, Majumdar S N and Bray A J 2004 Persistence exponents and the statistics of crossings and occupation times for Gaussian stationary processes *Phys. Rev. E* **69** 016106
- [47] Sabhapandit S, Majumdar S N and Comtet A 2006 Statistical properties of functionals of the paths of a particle diffusing in a one-dimensional random potential *Phys. Rev. E* **73** 051102
- [48] Cameron R H and Martin W T 1944 Transformations of weiner integrals under translations *Ann. Math.* **45** 386
- [49] Girsanov I V 1960 On transforming a certain class of stochastic processes by absolutely continuous substitution of measures *Theory Probab. Appl.* **5** 285–301
- [50] Ikeda N and Watanabe S 1981 *Stochastic Differential Equations and Diffusion Processes* 1st edn (Amsterdam: North-Holland)
- [51] Fox R F 1986 Functional-calculus approach to stochastic differential equations *Phys. Rev. A* **33** 467–76
- [52] Fox R F 1987 Stochastic calculus in physics *J. Stat. Phys.* **46** 1145–57
- [53] Risken H 1989 *The Fokker-Planck Equation* (Berlin: Springer)
- [54] Gardiner C W 1985 *Handbook of Stochastic Methods for Physics, Chemistry and the Natural Sciences* (Berlin: Springer)
- [55] Stratonovich R L 1966 A new representation for stochastic integrals and equations *SIAM J. Control* **4** 362
- [56] van Kampen N G 1992 *Stochastic Processes in Physics and Chemistry* (Amsterdam: North-Holland)
- [57] Lapolla A and Godec A 2018 Unfolding tagged particle histories in single-file diffusion: exact single- and two-tag local times beyond large deviation theory *New J. Phys.* **20** 113021
- [58] Tizón-Escamilla N, Lecomte V and Bertin E 2019 Effective driven dynamics for one-dimensional conditioned Langevin processes in the weak-noise limit *J. Stat. Mech.* **013201**
- [59] Pavliotis G A 2014 *Stochastic Processes and Applications* (New York: Springer)