




Contents lists available at ScienceDirect

Advances in Mathematics

journal homepage: www.elsevier.com/locate/aimUndistorted fillings in subsets of metric spaces[☆]Giuliano Basso^{a,b}, Stefan Wenger^{a,*}, Robert Young^c^a Department of Mathematics, University of Fribourg, Chemin du Musée 23, 1700 Fribourg, Switzerland^b Max Planck Institute for Mathematics, Vivatsgasse 7, 53111 Bonn, Germany^c Courant Institute of Mathematical Sciences, New York University, 251 Mercer St., New York, NY 10012, USA

ARTICLE INFO

Article history:

Received 17 February 2022

Received in revised form 2 April 2023

Accepted 2 April 2023

Available online 19 April 2023

Communicated by Moon Duchin

MSC:

primary 53C23

secondary 49Q15

Keywords:

Isoperimetric inequalities

Coning inequalities

Lipschitz connectivity

Isoperimetric subspace distortion

Deformation theorem

Integral currents

ABSTRACT

Lipschitz k -connectivity, Euclidean isoperimetric inequalities, and coning inequalities all measure the difficulty of filling a k -dimensional cycle in a space by a $(k+1)$ -dimensional object. In many cases, such as Banach spaces and CAT(0) spaces, it is easy to prove Lipschitz connectivity or a coning inequality, but harder to obtain a Euclidean isoperimetric inequality. We show that in spaces of finite Nagata dimension, Lipschitz connectedness implies Euclidean isoperimetric inequalities, and Euclidean isoperimetric inequalities imply coning inequalities. We show this by proving that if X has finite Nagata dimension and is Lipschitz k -connected or admits Euclidean isoperimetric inequalities up to dimension k then any isometric embedding of X into a metric space is isoperimetrically undistorted up to dimension $k+1$. Since X embeds in L_∞ , which admits a Euclidean isoperimetric inequality and a coning inequality, X admits such inequalities as well. In addition, we prove that an analog of the Federer-Fleming deformation theorem holds in such spaces X and use it to show that if X has finite Nagata dimension and is Lipschitz k -connected,

[☆] G. B. and S. W. were supported by Swiss National Science Foundation grant 182423. R. Y. was supported by National Science Foundation grant 2005609.

* Corresponding author.

E-mail addresses: basso@mpim-bonn.mpg.de (G. Basso), stefan.wenger@unifr.ch (S. Wenger), ryoung@cims.nyu.edu (R. Young).

then integral $(k + 1)$ -currents in X can be approximated by Lipschitz chains in total mass.

© 2023 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

1. Introduction

1.1. Overview

Isoperimetric inequalities measure how difficult it is to fill (Lipschitz) cycles in a given space by (Lipschitz) chains of one dimension higher. They are important in many branches of mathematics and play a crucial role in particular in asymptotic geometry and geometric group theory, where they appear as Dehn functions and higher filling functions and are quasi-isometry invariants of the underlying space or group. In this article, we study the relationship between isoperimetric inequalities and extension properties, especially Lipschitz connectivity and coning inequalities.

A space X is Lipschitz k -connected if Lipschitz maps from k -spheres to X can be extended to Lipschitz maps from $(k + 1)$ -balls. This is a key ingredient in constructing Lipschitz extensions in general; see [2], [29]. A 1-dimensional Lipschitz cycle is a sum of closed Lipschitz curves, so if X is Lipschitz 1-connected, it can be filled by filling each curve by a Lipschitz disc. Higher-dimensional Lipschitz cycles, however, can have complicated topology and it may not be possible to decompose them as sums of Lipschitz spheres. For this reason, it is often much more difficult to prove higher dimensional isoperimetric inequalities, and it is open in general whether Lipschitz k -connectedness implies a k -dimensional isoperimetric inequality of Euclidean type.

Coning inequalities bound the filling volume of a cycle in terms of its mass and diameter. They were introduced by Gromov in his seminal article [20], where it was shown that Riemannian manifolds with coning inequalities admit Euclidean isoperimetric inequalities. This result was later generalized to complete metric spaces by the second named author in [42]. Coning inequalities have also played a crucial role in recent articles on higher rank hyperbolicity [27], Morse quasiflats [24], [23], and the equivalence of flat and weak convergence of currents [43]. It is open in general whether spaces with k -dimensional Euclidean isoperimetric inequalities admit k -dimensional coning inequalities. This is clear in dimension 1, where one can fill a Lipschitz closed curve with diameter d and length L by decomposing it into L/d closed curves of length at most $3d$, but it is not clear whether higher-dimensional Lipschitz cycles can be decomposed in the same way.

In this paper, we will show that Lipschitz connectivity implies Euclidean isoperimetric inequalities and that Euclidean isoperimetric inequalities imply coning inequalities when the underlying space has finite Nagata dimension, thus establishing a partial converse to the results mentioned above. Nagata dimension can be thought of as a quantitative

metric version of topological dimension and is closely related to Gromov's asymptotic dimension [21]. These results are consequences of a more general theorem about isoperimetric subspace distortion. Isoperimetric distortion of subspaces was briefly addressed by Gromov in [21] and has recently been studied in the articles [11], [49], [31], [32] in connection with conjectures of Thurston, Gromov, and Bux-Wortman. Roughly speaking, a subspace X of a metric space Y is (isoperimetrically) undistorted up to dimension $k + 1$ if m -cycles in X with $0 \leq m \leq k$ can be filled almost as efficiently by $(m + 1)$ -chains in the subspace X as they can be filled in the ambient space Y . Our theorem about isoperimetric distortion asserts that if a quasiconvex metric space X of finite Nagata dimension is Lipschitz k -connected or has Euclidean isoperimetric inequalities up to dimension k then X is undistorted up to dimension $k + 1$ in any ambient space Y . This generalizes and strengthens a recent result of the third named author [49]. Since any metric space embeds isometrically in a Banach space and Banach spaces admit coning inequalities and hence Euclidean isoperimetric inequalities the relationships between Lipschitz connectedness, Euclidean isoperimetric inequalities, and coning inequalities described above follow.

Besides these structural results, our theorem on isoperimetric subspace distortion also allows for a Federer-Fleming type deformation theorem in quasiconvex metric spaces of finite Nagata dimension that admit Euclidean isoperimetric inequalities. The deformation theorem and its variants approximate chains or currents in Euclidean space or a simplicial complex by polyhedral chains in a cubical lattice or triangulation. Metric spaces of finite Nagata dimension need not admit a triangulation, but we show that any quasiconvex space X of finite Nagata dimension can be approximated by simplicial complexes. When X has a Euclidean isoperimetric inequality, we can use these approximations to define substitutes for polyhedral chains in X . Moreover, we obtain a Federer-Fleming type deformation theorem in X by combining the classical deformation theorem in the simplicial approximation and the isoperimetric subspace distortion theorem in X . As an application, we show that when X has finite Nagata dimension and is Lipschitz connected, integral currents in X can be approximated in total mass by Lipschitz chains.

1.2. Undistorted fillings and first applications

Given a complete metric space X and $k \geq 0$, we denote by $\mathbf{I}_k(X)$ the abelian group of k -dimensional metric integral currents in X in the sense of Ambrosio-Kirchheim [3]. See Sections 2 and 3 for the definitions and concepts used throughout this introduction. Every Lipschitz k -chain in X induces a k -dimensional metric integral current in X and we encourage the geometrically minded reader to think of integral currents as suitable limits of Lipschitz chains. The filling volume in X of $T \in \mathbf{I}_k(X)$ is defined by

$$\text{Fillvol}_X(T) = \inf \{ \mathbf{M}(S) : S \in \mathbf{I}_{k+1}(X) \text{ and } \partial S = T \},$$

where $\mathbf{M}(S)$ denotes the mass of S and ∂S is its boundary.

We say that X has (EI_k) for some $k \geq 0$ if there exists $D > 0$ such that for every $0 \leq m \leq k$ and every cycle $T \in \mathbf{I}_m(X)$ we have

$$\text{Fillvol}_X(T) \leq D \mathbf{M}(T)^{\frac{m+1}{m}}.$$

This means that X has Euclidean isoperimetric inequalities in dimensions $m = 1, \dots, k$ whenever $k \geq 1$. Condition (EI_0) is always true and is only included to shorten the statements of our theorems. We remark that (EI_k) is sometimes also denoted by (EII_k) by some authors (see e.g. [19], [24]).

We say that X has (LC_k) or that X is Lipschitz k -connected if there exists $c \geq 1$ such that for every $0 \leq m \leq k$ each L -Lipschitz map from the Euclidean m -sphere to X extends to a cL -Lipschitz map defined on the $(m+1)$ -ball. It is not difficult to see that a metric space has (LC_0) if and only if it is quasiconvex (see Section 2.1 for the precise definition). Examples of Lipschitz k -connected metric spaces are compact k -connected Riemannian manifolds, $\text{CAT}(0)$ -spaces and, more generally, spaces with a convex bicombing. This class also includes some Carnot groups such as the $(2k+1)$ -dimensional Heisenberg group.

A closed subset $X \subset Y$ of a complete metric space Y is said to be (isoperimetrically) *undistorted in Y up to dimension $k+1$* , for some $k \geq 0$, if there exists $C > 0$ such that for all cycles $T \in \mathbf{I}_m(X)$ with $0 \leq m \leq k$ one has

$$\text{Fillvol}_X(T) \leq C \text{Fillvol}_Y(T).$$

Our definition of undistorted does not involve additive error terms and thus differs slightly from the definitions introduced in [11], [49], [32]. The following theorem is one of the main results in the present paper.

Theorem 1.1. *Let Y be a complete metric space and $X \subset Y$ a closed quasiconvex subset of finite Nagata dimension. If X has (LC_k) or (EI_k) for some $k \geq 0$, then X is undistorted in Y up to dimension $k+1$. The distortion constant only depends on the data of X .*

For the definition of Nagata dimension see Section 2.3 and [29]. Examples of metric spaces of finite Nagata dimension include compact Riemannian manifolds, homogeneous or negatively pinched Hadamard manifolds, Carnot groups and, more generally, equiregular sub-Riemannian manifolds, doubling metric spaces and many more. Moreover, products, finite unions, and subsets of spaces of finite Nagata dimension have finite Nagata dimension.

In Section 5.1 we will discuss variants and generalizations of Theorem 1.1 in which we relax the condition that X have finite Nagata dimension and in addition obtain control on the support of fillings in X . Theorem 1.1 generalizes and strengthens [49, Theorem 1.3] by the third named author, which was used to prove isoperimetric inequalities for subsets of symmetric spaces. Notice that in [49] the ambient space Y is assumed to have finite Nagata dimension. In contrast, we do not pose any restrictions on Y . Since

X embeds isometrically into a Banach space and since every Banach space admits Euclidean isoperimetric inequalities and coning inequalities in all dimensions, see [42], we in particular obtain the following consequences.

Corollary 1.2. *Let X be a complete metric space of finite Nagata dimension. If X has (LC_k) for some $k \geq 1$, then X has (EI_k) .*

When $k = 1$ the corollary holds without the assumption that X have finite Nagata dimension and is much easier to prove. This is because integral 1-cycles can be decomposed into the sum of closed Lipschitz curves, see [44]. The corollary is furthermore known when X has (LC_k) and is of Nagata dimension at most k because in this case X is an absolute Lipschitz retract by [29]. Thus, the corollary is most interesting for spaces which are not Lipschitz connected up to the Nagata dimension. Typical examples with such a behavior are Carnot groups such as the higher Heisenberg groups; see the paragraph after the next corollary. We mention that the (LC_k) property is preserved under various constructions such as taking products, or passing to ultralimits or asymptotic cones. In comparison, it is not known whether the (EI_k) condition is preserved under these constructions.

In order to formulate the second consequence of the theorem, recall that a complete metric space X is said to admit coning inequalities up to dimension k , or has (CI_k) for short, if there exists $C > 0$ such that for every $0 \leq m \leq k$ and every cycle $T \in \mathbf{I}_m(X)$ of bounded support $\text{spt } T$ we have

$$\text{Fillvol}_X(T) \leq C \text{diam}(\text{spt } T) \mathbf{M}(T).$$

It is well-known that in a complete metric space (CI_k) implies (EI_k) . This was proved by Gromov [20] for Riemannian manifolds and extended to metric spaces in [42]. The following corollary gives a partial converse.

Corollary 1.3. *Let X be a complete quasiconvex metric space of finite Nagata dimension. If X has (EI_k) for some $k \geq 1$, then X has (CI_k) .*

Using a suitable variant of Theorem 1.1 established in Section 5.1 we will actually get a stronger version of the coning inequality in which we also obtain control over the diameter of the filling chain; see Corollary 5.2. Coning inequalities play an important role in the recent articles [24], [23], [27], and the stronger version of the coning inequality established in Corollary 5.2 prominently appears in [24], [23], where it is called strong coning inequality.

We now briefly discuss some examples of spaces to which Corollaries 1.2 and 1.3 apply. Compact k -connected Riemannian manifolds have (LC_k) and finite Nagata dimension by [29] and therefore have (CI_k) and (EI_k) . The n -th Heisenberg group \mathbb{H}^n of topological dimension $2n + 1$, equipped with a left-invariant sub-Riemannian or sub-Finsler metric d_c , has Nagata dimension $2n + 1$ by [30] and satisfies (LC_{n-1}) by [46]. It thus follows from

the corollaries above that (\mathbb{H}^n, d_c) has (CI_{n-1}) and (EI_{n-1}) . Previously, it was known that \mathbb{H}^n , when equipped with a left-invariant Riemannian metric, admits Euclidean isoperimetric inequalities up to dimension $n - 1$ for Lipschitz cycles, see [48]. Together with [45] this also yields that (\mathbb{H}^n, d_c) has (EI_{n-1}) for compactly supported integral currents.

We finally mention that Theorem 1.1 can be reformulated in terms of the *absolute filling volume* of a cycle $T \in \mathbf{I}_k(X)$ in a complete metric space X defined by

$$\text{Fillvol}_\infty(T) := \inf\{\text{Fillvol}_Y(T) : Y \text{ complete metric space containing } X\},$$

compare with Gromov's filling volume of abstract Riemannian manifolds in [20]. Our theorem then says that if X is a complete, quasiconvex metric space of finite Nagata dimension which has (LC_k) or (EI_k) for some $k \geq 0$ then every cycle $T \in \mathbf{I}_k(X)$ satisfies

$$\text{Fillvol}_X(T) \leq C \text{Fillvol}_\infty(T)$$

for some $C > 0$ only depending on the data of X . It is easy to see that if Y is an injective metric space containing X then $\text{Fillvol}_\infty(T) = \text{Fillvol}_Y(T)$ for every cycle $T \in \mathbf{I}_k(X)$. Particular examples of such spaces Y are the Banach space $\ell_\infty(X)$ of bounded functions on X equipped with the sup norm or the injective hull of X ; see [25].

1.3. Deformation theorem in spaces of finite Nagata dimension

The classical deformation theorem of Federer-Fleming, see [17] or [16], shows that integral k -currents in Euclidean \mathbb{R}^n can be approximated by polyhedral k -chains in the k -skeleton of the standard cubical subdivision of \mathbb{R}^n . This result can be generalized to the setting of Riemannian manifolds with a geometric group action and, more generally, to metric spaces admitting a bilipschitz triangulation. The deformation theorem and its variants have been of fundamental importance in geometric measure theory and other fields, for example in the context of Dehn functions and higher filling functions; see, for example, [15], [10], [49], [48], [1]. Metric spaces of finite Nagata dimension do not in general admit a bilipschitz triangulation. Nevertheless, using a variant of Theorem 1.1 and the methods developed in its proof, we can establish an analog of the Federer-Fleming deformation theorem in spaces of finite Nagata dimension.

We first introduce a substitute for a cubical subdivision in our setting. Given a metric simplicial complex Σ and $m \geq 0$, let $\mathcal{P}_m(\Sigma)$ be the abelian group of polyhedral m -chains in Σ , that is, finite sums of the form $\sum \theta_i \llbracket \sigma_i \rrbracket$, where $\theta_i \in \mathbb{Z}$ and $\llbracket \sigma_i \rrbracket$ is the current induced by the (oriented) m -simplex $\sigma_i \subset \Sigma$. Let X be a complete metric space and let $k \in \mathbb{N}$ and $C, \varepsilon > 0$. Suppose there exist a finite-dimensional metric simplicial complex Σ and a map $\varphi: \Sigma^{(0)} \rightarrow X$, defined on the 0-skeleton $\Sigma^{(0)}$ of Σ , with the following properties:

- (1) the metric on Σ is a length metric and each simplex in Σ is a Euclidean simplex of side length ε ;
- (2) φ is a quasi-isometry; more precisely, the image of φ is $(C\varepsilon)$ -dense in X and

$$d(z, w) - C\varepsilon \leq d(\varphi(z), \varphi(w)) \leq Cd(z, w) \quad \text{for all } z, w \in \Sigma^{(0)}.$$

Suppose also that there are homomorphisms $\Lambda_m: \mathcal{P}_m(\Sigma) \rightarrow \mathbf{I}_m(X)$, $m = 0, \dots, k$, subject to:

- (3) $\partial \circ \Lambda_{m+1} = \Lambda_m \circ \partial$ and $\Lambda_0(\llbracket z \rrbracket) = \llbracket \varphi(z) \rrbracket$ for every $z \in \Sigma^{(0)}$;
- (4) $\Lambda_m(\llbracket \sigma \rrbracket)$ has support in $B(\varphi(\sigma^{(0)}), C\varepsilon)$ and $\mathbf{M}(\Lambda_m(\llbracket \sigma \rrbracket)) \leq C\varepsilon^m$ for every m -simplex σ in Σ .

Any triple $(\Sigma, \varphi, \Lambda_*)$, where Σ , φ and $\Lambda_* = \{\Lambda_0, \dots, \Lambda_k\}$ satisfy the properties listed above, is called (k, ε) -polyhedral structure on X with constant C . If such a polyhedral structure is given we set $\mathcal{P}_m(X) := \{\Lambda_m(Q) : Q \in \mathcal{P}_m(\Sigma)\}$ and call its elements *polyhedral m -chains in X* .

Theorem 1.4. *Let X be a complete quasiconvex metric space of finite Nagata dimension which has (LC_k) or (EI_k) for some $k \geq 1$. Then there exists $C > 0$ such that for every $\varepsilon > 0$ there is a (k, ε) -polyhedral structure on X with constant C such that the following holds. For every $m = 1, \dots, k$ and every $T \in \mathbf{I}_m(X)$ there exist $P \in \mathcal{P}_m(X)$, $R \in \mathbf{I}_m(X)$, and $S \in \mathbf{I}_{m+1}(X)$ with $T = P + R + \partial S$ and such that*

$$\begin{aligned} \mathbf{M}(P) &\leq C \mathbf{M}(T), & \mathbf{M}(\partial P) &\leq C \mathbf{M}(\partial T), \\ \mathbf{M}(S) &\leq \varepsilon C \mathbf{M}(T) + \varepsilon^2 C \mathbf{M}(\partial T), & \mathbf{M}(R) &\leq \varepsilon C \mathbf{M}(\partial T), \end{aligned}$$

as well as $\text{spt } P, \text{spt } S \subset B(\text{spt } T, C\varepsilon)$ and $\text{spt } \partial P, \text{spt } R \subset B(\text{spt } \partial T, C\varepsilon)$.

The bounds on the mass of P , ∂P , and R in Theorem 1.4 are as in the deformation theorems of Simon [39] and White [47]. In particular, this gives a slightly sharper bound on the mass of P than the original version of Federer and Fleming, which only gives a bound $\mathbf{M}(P) \leq C[\mathbf{M}(T) + \varepsilon \mathbf{M}(\partial T)]$. The bound on the mass of S includes an additional error term coming from a use of Theorem 1.1. This error term disappears however when T is a cycle. If X has (LC_k) then the map φ in the polyhedral structure $(\Sigma, \varphi, \Lambda_*)$ in Theorem 1.4 is even defined on $\Sigma^{(k)}$ and the homomorphisms Λ_m are simply given by the pushforward under φ . Hence, in this case, elements of $\mathcal{P}_k(X)$ are (currents induced by) Lipschitz k -chains. It thus follows that if a metric space X has finite Nagata dimension and has (LC_k) for some $k \geq 1$ then integral k -currents in X can be approximated in the flat norm by Lipschitz chains; see Corollary 7.3. The classical deformation theorem yields a stronger approximation result in Euclidean space which asserts that integral currents

can be approximated by Lipschitz chains in total mass. Using the tools developed to prove Theorem 1.4 we can generalize this stronger approximation result as follows.

Corollary 1.5. *Let X be a complete metric space of finite Nagata dimension and let $T \in \mathbf{I}_k(X)$ for some $k \geq 1$. If X has (LC_{k-1}) then there exists a sequence of Lipschitz k -chains such that the induced integral currents T_i converge to T in total mass, that is, $\mathbf{M}(T - T_i) \rightarrow 0$ and $\mathbf{M}(\partial T - \partial T_i) \rightarrow 0$ as $i \rightarrow \infty$.*

For related approximation results in a different setting see for example [13], [18]. We do not know whether Corollary 1.5 holds without the assumption that X have finite Nagata dimension.

The proof of Theorem 1.4 crucially relies on the following approximation result which should be of independent interest and whose proof uses methods and arguments developed by Lang-Schlichenmaier [29] in the context of Lipschitz extensions.

Theorem 1.6. *Let X, Y be metric spaces such that X is quasiconvex and of Nagata dimension $\leq n$ and Y has (LC_{n-1}) and contains X . Then there exists a constant C depending only on the data of X and Y such that for every $\varepsilon > 0$ there is a metric simplicial complex Σ and C -Lipschitz maps $\psi: X \rightarrow \Sigma$ and $\varphi: \Sigma \rightarrow Y$ with the following properties:*

- (1) Σ has dimension $\leq n$, the metric on Σ is a length metric, and every simplex in Σ is a Euclidean simplex of side length ε ;
- (2) $\varphi(\Sigma^{(0)}) \subset X$, $\text{Hull}(\psi(X)) = \Sigma$ and $d(x, \varphi(\psi(x))) \leq C\varepsilon$ for all $x \in X$.

If X has (LC_{k-1}) for some $k \geq 1$, then $\varphi(\Sigma^{(k)}) \subset X$.

If X is compact then Σ is a finite simplicial complex and, in particular, the metric on Σ is a geodesic metric. The hull of a subset $A \subset \Sigma$ is the smallest subcomplex of Σ containing A and is denoted $\text{Hull}(A)$. Theorem 1.6 is used as follows to construct a (k, ε) -polyhedral structure on a metric space X as in Theorem 1.4. Let Σ, ψ, φ be as in Theorem 1.6, when applied to X and $Y = \ell_\infty(X)$. It is not difficult to see that Σ and the restriction $\varphi|_{\Sigma^{(0)}}: \Sigma^{(0)} \rightarrow X$ satisfy properties (1) and (2) in the definition of a polyhedral structure. Furthermore, using $\varphi|_{\Sigma^{(0)}}$ and the (LC_k) or (EI_k) condition on X one easily builds homomorphisms $\Lambda_m: \mathcal{P}_m(\Sigma) \rightarrow \mathbf{I}_m(X)$ for $m = 0, \dots, k$ satisfying properties (3) and (4) in the definition of a polyhedral structure. This is accomplished in Proposition 3.8. We finally mention that Theorem 1.6 has found another application in the recent article [35], where it was used to prove the existence of a quasiconformal parametrization of 2-dimensional metric surface under minimal conditions.

1.4. Outlines of proof

The idea behind the proof of Theorem 1.1 is inspired by the arguments used to prove [49, Theorem 1.3]. Let X and Y be as in the statement of the theorem. After possibly embedding Y into a bigger space (for example a Banach space) we may assume that Y has (LC_n) , where n is the Nagata dimension of X .

One of the main ingredients in the proof is a factorization theorem (see Theorem 4.1), which is established using arguments from [29] and roughly says the following. There exists a Lipschitz map $f: Y \rightarrow Y$ such that f restricts to the identity on X and $f|_{Y \setminus X}$ factors through an $(n + 1)$ -dimensional simplicial complex Σ . More precisely, on $Y \setminus X$ we have $f = h \circ g$ for suitable maps $g: Y \setminus X \rightarrow \Sigma$ and $h: \Sigma \rightarrow Y$ satisfying the following properties. Let $\varrho: Y \rightarrow \mathbb{R}$ be the distance function from the set X and let $r > 0$. Then g is Cr^{-1} -Lipschitz on the set $\{\varrho > r\}$ and h is Cr -Lipschitz on every simplex $\sigma \subset \Sigma$ for which $g^{-1}(\text{st } \sigma) \cap \{\varrho \leq r\} \neq \emptyset$, where C is independent of r and $\text{st } \sigma$ denotes the open star of σ in Σ . Moreover, h maps the 1-skeleton of Σ to X . The simplicial complex Σ is equipped with the so-called ℓ_2 -metric and every simplex is isometric to a standard Euclidean simplex.

We now outline how this factorization theorem is used to prove Theorem 1.1. Let $T \in \mathbf{I}_k(X)$ be a cycle in X and $S \in \mathbf{I}_{k+1}(Y)$ a filling of T in Y . After possibly modifying Y and S slightly we may achieve that S meets the set X only on the boundary T and not too much mass of S is concentrated near X . For suitable $r > 0$ (small) we then send the restriction $S \llcorner \{\varrho > r\}$ to an integral current $S'_r \in \mathbf{I}_{k+1}(\Sigma)$ via the map g . Notice that the mass distortion of g is proportional to $s^{-(k+1)}$ on the part of S located at a distance around s . Next, we use a refined version of the classical Federer-Fleming deformation theorem in Σ to “push” the integral current S'_r to the $(k + 1)$ -skeleton $\Sigma^{(k+1)}$ of Σ and obtain a polyhedral chain $P'_r \in \mathcal{P}_{k+1}(\Sigma)$. This refined version of the deformation theorem, a detailed proof of which is given in the Appendix, includes local mass estimates and allows us to control the mass of P'_r on every simplex. Finally, we map P'_r back to X via a chain homomorphism $\Lambda_{k+1}: \mathcal{P}_{k+1}(\Sigma) \rightarrow \mathbf{I}_{k+1}(X)$. Here, Λ_1 is the map induced by the restriction $h|_{\Sigma^{(1)}}: \Sigma^{(1)} \rightarrow X$. For $m = 2, \dots, k$, we construct Λ_{m+1} from Λ_m by applying the (LC_m) or (EI_m) property of X . For a suitable sequence $r_i \rightarrow 0$, the currents $P_i := \Lambda_{k+1}(P'_{r_i}) \in \mathbf{I}_{k+1}(X)$ converge to a filling $P \in \mathbf{I}_{k+1}(X)$ of T with the desired bound on mass, thus concluding the proof of Theorem 1.1.

The proof of Theorem 1.4 is similar but somewhat easier because all the involved maps are globally Lipschitz. Instead of using the factorization theorem sketched above we apply Theorem 1.6 with $Y = \ell_\infty(X)$ to obtain a finite dimensional metric simplicial complex Σ and Lipschitz maps $\psi: X \rightarrow \Sigma$ and $\varphi: \Sigma \rightarrow Y$ as in the theorem. As described after the theorem, this leads to a (k, ε) -polyhedral structure $(\Sigma, \varphi, \Lambda_*)$ on X . Now, let $T \in \mathbf{I}_k(X)$ and consider the pushforward current $T' = \psi_\# T \in \mathbf{I}_k(\Sigma)$. The classical Federer-Fleming deformation theorem in Σ yields a decomposition $T' = P' + R' + \partial S'$ for currents $P' \in \mathcal{P}_k(\Sigma)$ and $R' \in \mathbf{I}_k(X)$, $S' \in \mathbf{I}_{k+1}(\Sigma)$ with suitable bounds on mass. We then use the fact that $\varphi \circ \psi$ is close to the identity on X together with further properties

of Λ_m to show that T can be decomposed as $T = P + \hat{R} + \partial\hat{S}$, where $P = \Lambda_k(P')$ and where $\hat{R} \in \mathbf{I}_k(Y)$ and $\hat{S} \in \mathbf{I}_{k+1}(Y)$ are suitable currents. We finally use Theorem 1.1 to “push” \hat{R} and \hat{S} from Y to X in order to obtain a decomposition as in Theorem 1.4.

1.5. Structure of the paper

In Section 2 we introduce notation and definitions concerning concepts used in the article, including simplicial complexes, Nagata dimension, Lipschitz connectedness, and pointwise Lipschitz constants. Section 3 contains the necessary definitions from the Ambrosio-Kirchheim theory of metric currents. We furthermore prove Proposition 3.8 which allows us to construct suitable homomorphisms from the group of polyhedral chains in a simplicial complex to the group of integral currents in a metric space. This technical result will be needed in the proofs of Theorems 1.1 and 1.4. The main result in Section 4 is the factorization theorem sketched in Section 1.4 above; see Theorem 4.1. As already mentioned, this is one of the main ingredients in the proof of Theorem 1.1. In Section 5 we discuss and prove variants and generalizations of Theorem 1.1 and deduce Theorem 1.1 from them. The main aim of Section 6 is to establish Theorem 1.6, whose proof is similar to that of Theorem 4.1. Section 7 contains the proof of Theorem 1.4. Finally, in the Appendix we give a detailed proof of a refined version of the Federer-Fleming deformation theorem in metric spaces admitting a bilipschitz triangulation. Our refined version includes local mass estimates which are of particular importance in the proof of our isoperimetric subspace distortion theorem.

1.6. Acknowledgments

We thank the anonymous referee for their helpful suggestions.

2. Preliminaries

2.1. Notation and definitions

We recall standard notation and definitions from metric geometry. Suppose that $A, B \subset X$ are subsets of a metric space X . We write

$$d(A, B) := \inf\{d(a, b) : a \in A, b \in B\} \quad \text{and} \quad \text{diam}(B) := \sup\{d(x, x') : x, x' \in B\}.$$

We use the convention that $\inf \emptyset = +\infty$ and $\sup \emptyset = -\infty$. For every $\varepsilon > 0$ we let $U(A, \varepsilon)$ and $B(A, \varepsilon)$ denote the open and closed ε -neighborhood of A , respectively. We say that a subset $Z \subset X$ is an ε -net if $d(z, z') > \varepsilon$ for all distinct $z, z' \in Z$ and if the family of all balls $B(z, \varepsilon)$ with $z \in Z$ covers X .

A metric space X is called c -quasiconvex, $c \geq 1$, if all $x, x' \in X$ can be connected by a curve $\gamma: [0, 1] \rightarrow X$ such that $\ell(\gamma) \leq cd(x, x')$. For us a curve is always continuous

and we use $\ell(\gamma) \in [0, \infty]$ to denote the *length* of a curve γ . Let $k \geq 0$ and denote by \mathcal{H}^k the k -dimensional Hausdorff measure on X and let \mathcal{L}^k denote the Lebesgue measure on \mathbb{R}^k . A subset $A \subset X$ is said to be *countably \mathcal{H}^k -rectifiable* if it is \mathcal{H}^k -measurable and there exist subsets $A_i \subset \mathbb{R}^k$ and Lipschitz maps $f_i: A_i \rightarrow X$ such that $\mathcal{H}^k(A \setminus \bigcup_{i=1}^{\infty} f_i(A_i)) = 0$. A finite Borel measure μ on X is said to be concentrated on a Borel set $B \subset X$ if $\mu(X \setminus B) = 0$. The set $\text{spt } \mu$ consisting of all $x \in X$ such that $\mu(U(x, \varepsilon)) > 0$ for all $\varepsilon > 0$ is called the support of μ . If X is separable and complete then μ is concentrated on $\text{spt } \mu$ and $\text{spt } \mu$ is σ -compact. This is true more generally if the cardinality of X is an Ulam number; see [16, 2.1.6]. As is done for example in [3] we will assume throughout this paper that the cardinality of any set is an Ulam number so that our arguments apply in both the separable and non-separable cases; see [3, Lemma 2.9] and the remark following it.

2.2. Simplicial complexes

For the concepts appearing below we refer e.g. to [40]. Let Σ be a simplicial complex. We denote by $\mathcal{F}_k(\Sigma)$ the collection of closed k -simplices in Σ and by $\mathcal{F}(\Sigma)$ the collection of closed simplices of any dimension in Σ . We also write \mathcal{F}_k and \mathcal{F} if there is no danger of ambiguity. We say that Σ is n -dimensional if $\mathcal{F}_n \neq \emptyset$ and $\mathcal{F}_{n+1} = \emptyset$. The k -skeleton of Σ is denoted by $\Sigma^{(k)}$. Notice that if Σ is n -dimensional, then $\Sigma^{(m)} = \Sigma^{(n)}$ for every $m \geq n$. The smallest subcomplex of Σ containing a given subset $A \subset \Sigma$ is called the hull of A and is denoted $\text{Hull}(A)$. The open star of $\sigma \in \mathcal{F}$ is defined by

$$\text{st } \sigma := \bigcup \left\{ \text{int } \tau : \tau \in \mathcal{F} \text{ and } \sigma \subset \tau \right\},$$

where $\text{int } \tau$ is the interior of τ . We will use the following two specific choices of a metric on a given simplicial complex. For a nonempty index set I define a simplicial complex by

$$\Sigma(I) := \left\{ x \in \ell_2(I) : x_i \geq 0 \text{ and } \sum_{i \in I} x_i = 1 \right\}$$

and equip $\Sigma(I) \subset \ell_2(I)$ with the metric induced by the norm of $\ell_2(I)$. A given simplicial complex Σ can be naturally realized as a subset of $\Sigma(I)$, where $I = \mathcal{F}_0$ is the collection of vertices in Σ . The induced metric on $\Sigma \subset \Sigma(I)$ will be referred to as the ℓ_2 -metric on Σ . The diameter of Σ with respect to the ℓ_2 -metric is always at most $\sqrt{2}$. If Σ is path-connected then the associated length metric on Σ will be called the *length metric* on Σ . Notice that with respect to either metric each simplex of Σ is isometric to a standard Euclidean simplex. We close this subsection with the following lemma which collects various results that will be used later.

Lemma 2.1. *Let Σ be a finite-dimensional simplicial complex, equipped with the ℓ_2 -metric. Then the following holds:*

- (1) For every $x \in \Sigma$, there is $\varepsilon_x > 0$ such that if $y \in U(x, \varepsilon_x)$ then x and y lie in a common simplex;
- (2) If $A \subset \Sigma$ is separable, then $\Sigma' := \text{Hull}(A)$ is a countable simplicial complex, that is, the set of vertices of Σ' is countable;
- (3) If Σ is path-connected, then the length metric and the ℓ_2 -metric induce the same topology.

An analogous statement holds when the ℓ_2 -metric is replaced by the length metric.

Proof. For $x \in \Sigma$ denote by $\sigma(x)$ the unique simplex σ such that $x \in \text{int } \sigma$. Given $x = (x_i)_{i \in I}$, we put $I(x) := \{i \in X : x_i \neq 0\}$ and $\varepsilon_x := \min\{x_i : i \in I(x)\}$. Clearly, $\varepsilon_x > 0$ and if $|x - y| < \varepsilon_x$, then $y_i \neq 0$ for all $i \in I(x)$. Hence, $\sigma(x) \subset \sigma(y)$ whenever $|x - y| < \varepsilon_x$. This gives (1). Next, we show (2). Let Z be a countable dense subset of A . We claim that

$$\text{Hull}(A) = \bigcup_{z \in Z} \sigma(z). \quad (2.1)$$

Fix $x \in A$. As Z is a dense subset of A , there exists $z \in Z$ such that $|x - z| < \varepsilon_x$ and thus $\sigma(x) \subset \sigma(z)$. This yields (2.1) and (2) follows.

Finally, we prove (3). Suppose now that Σ is path-connected. By virtue of (1), the open balls $U(x, \varepsilon)$, where $\varepsilon \leq \varepsilon_x$, with respect to the length metric and the ℓ_2 -metric coincide with each other. This shows in particular that both metrics induce the same topology on Σ , as desired. \square

2.3. Nagata dimension

In the following, we recall the definition of Nagata dimension and some of its basic properties. A covering $\mathcal{B} = (B_i)_{i \in I}$ of a metric space X is said to be D -bounded, for a real number $D \geq 0$, if every set $B \in \mathcal{B}$ has diameter less than or equal to D . Let $n \geq 0$ be an integer and $s > 0$ a real number. A covering \mathcal{B} has s -multiplicity at most n if every subset $B \subset X$ with $\text{diam}(B) \leq s$ meets at most n members of \mathcal{B} .

Definition 2.2. We say that a metric space X has Nagata dimension $\leq n$ with constant c if for every real number $s > 0$, X admits a cs -bounded covering \mathcal{B}_s having s -multiplicity at most $n + 1$.

Equivalent definitions can be found in [29, Proposition 2.5]. The infimum of those $n \geq 0$ for which X has Nagata dimension $\leq n$ is denoted by $\text{dim}_N(X)$ and is called the Nagata dimension of X . Nagata dimension was introduced by Assouad in [4], building on earlier work of Nagata [36]. It can be thought of as a quantitative metric version of the topological dimension $\text{dim}(X)$ of X . Clearly, one has $\text{dim}(X) \leq \text{dim}_N(X)$ and this inequality can be strict. For example, $\text{dim}(\mathbb{Z}) = 0$ but $\text{dim}_N(\mathbb{Z}) = 1$. As a more striking

example, it can be shown that the group $(\mathbb{Z}/2\mathbb{Z}) \wr \mathbb{Z}^2$ has topological dimension zero but infinite Nagata dimension (see [8]).

However, on the positive side, if X is a compact Riemannian manifold or a Carnot group equipped with the Carnot-Carathéodory distance, then $\dim(X) = \dim_N(X)$. See [29, p. 3635] and [30, Theorem 4.2]. Moreover, any doubling metric space has finite Nagata dimension (see [29, Lemma 2.3]). Nagata dimension also enjoys many good structural properties. For example, $\dim_N X \times Y \leq \dim_N X + \dim_N Y$ and if $X = A \cup B$, then $\dim_N X = \sup \{ \dim_N(A), \dim_N(B) \}$. In [14], [30] it is proved that $\dim(Z) \leq \dim_N(X)$ whenever Z is an asymptotic cone or tangent cone of X . For further results concerning Nagata dimension we refer to the articles [5], [7], [9], [29], [30] and the references therein.

2.4. Lipschitz connectedness

Let X and Y be metric spaces. A map $f: X \rightarrow Y$ is called L -Lipschitz, where $L \geq 0$ is a real number, if $d(f(x), f(x')) \leq L d(x, x')$ for all $x, x' \in X$. The quantity

$$\text{Lip}(f) := \inf \{ L \geq 0 : f \text{ is } L\text{-Lipschitz} \}$$

is called the *Lipschitz constant* of f . We say that f is C -bilipschitz, $C \geq 1$, if

$$C^{-1}d(x, y) \leq d(f(x), f(y)) \leq Cd(x, y)$$

for all $x, y \in X$. In the following, we recall the definition of Lipschitz k -connected metric spaces and state several examples. Let $B^{m+1} := \{x \in \mathbb{R}^{m+1} : |x| \leq 1\}$, for $m \geq 0$, denote the closed $(m+1)$ -dimensional Euclidean unit ball and put $S^m := \partial B^{m+1}$. We equip S^m and B^{m+1} with the induced Euclidean metric.

Definition 2.3. A metric space X is said to be Lipschitz k -connected if there exists $c \geq 1$ such that every L -Lipschitz map $f: S^m \rightarrow X$ with $0 \leq m \leq k$ has a cL -Lipschitz extension $\bar{f}: B^{m+1} \rightarrow X$. We abbreviate this by saying that X has property (LC_k) .

In particular, Banach spaces and $\text{CAT}(0)$ spaces or, more generally, metric spaces admitting a convex bicombing have (LC_k) with constant 3; see [38, Proposition 6.2.2.]. Every k -connected compact Riemannian manifold has (LC_k) . This follows from [29, Theorem 5.1] which states that if a compact metric space X is k -connected and Lipschitz k -connected in the small, then X has (LC_k) . Moreover, certain horospheres in symmetric spaces of noncompact type of rank k or in the product of k proper $\text{CAT}(0)$ spaces have (LC_{k-2}) . This has been established in [31] and [33], respectively. The jet space Carnot groups $(J^s(\mathbb{R}^k), d_c)$ have (LC_{k-1}) ; see [46]. Examples of Carnot groups which can be realized as jet space Carnot groups include the n th Heisenberg group, the Engel group and the model filiform groups. We refer to [41] for more information on these groups. There are many examples of metric groups which have (LC_1) ; see e.g. [12], [34].

2.5. Pointwise Lipschitz constants

Let $f: X \rightarrow Y$ be a map between metric spaces X and Y . The *pointwise (lower) Lipschitz constant* of f at x is defined by

$$\text{lip } f(x) := \lim_{r \rightarrow 0^+} \ell_r f(x), \quad \text{where } \ell_r f(x) := \inf_{0 < s < r} \sup_{d(x,y) < s} \frac{d(f(x), f(y))}{s}.$$

If f is Lipschitz then the function $\text{lip } f$ is real-valued and Borel; see [26, Lemma 4.1.2].

Lemma 2.4. *Let $f: X \rightarrow Y$ be a map such that $\text{lip } f(x) \leq C$ for all $x \in X$. Then f is continuous and $\ell(f \circ \gamma) \leq C\ell(\gamma)$ for every curve γ in X . In particular, if X is λ -quasiconvex, then f is $C\lambda$ -Lipschitz.*

Proof. Let $x \in X$. Since $\text{lip } f(x) \leq C$, for every $r > 0$ there is $s \in (0, r)$ such that

$$\sup_{d(x,y) < s} d(f(x), f(y)) \leq 2Cs;$$

hence, f is continuous at x . In order to prove the second statement it is enough to show that

$$d(f(\gamma(0)), f(\gamma(l))) \leq C\ell(\gamma) \tag{2.2}$$

for every rectifiable curve $\gamma: [0, l] \rightarrow X$. Fix a rectifiable curve γ , which we may assume to be parametrized by arc-length. Let $C' > C$ and consider the set

$$A := \{t \in [0, l] : d(f(\gamma(0)), f(\gamma(t))) \leq C't\}.$$

Clearly, A is nonempty and closed, so $t := \sup A$ is contained in A . We claim that $t = l$. We argue by contradiction and assume that $t < l$. Choose $s > 0$ sufficiently small such that $t + s < l$ and

$$\sup_{d(\gamma(t), y) < s} \frac{d(f(\gamma(t)), f(y))}{s} \leq C'.$$

Since $d(\gamma(t), \gamma(t + s')) \leq \ell(\gamma|_{[t, t+s']}) = s'$ for all $0 < s' < s$, we obtain from the above and the continuity of f that $d(f(\gamma(t)), f(\gamma(t + s))) \leq C's$ and hence

$$d(f(\gamma(0)), f(\gamma(t + s))) \leq C'(t + s).$$

This shows that $t + s \in A$, which is impossible. We therefore have $t = l$ and thereby $d(f(\gamma(0)), f(\gamma(l))) \leq C'l = C'\ell(\gamma)$. Since $C' > C$ was arbitrary this shows (2.2) and proves the second statement of the lemma. \square

3. Currents in metric spaces

We recall relevant notions from Ambrosio-Kirchheim’s theory of currents of finite mass in metric spaces [3]. See [28] for a variant of this theory.

3.1. Currents of finite mass, push-forwards and restrictions

Let X be a complete metric space, $k \geq 0$ an integer, and let $\mathcal{D}^k(X)$ be the set of tuples (π_0, \dots, π_k) of Lipschitz functions $\pi_i: X \rightarrow \mathbb{R}$ with π_0 bounded.

Definition 3.1. A multi-linear functional $T: \mathcal{D}^k(X) \rightarrow \mathbb{R}$ is called metric k -current (of finite mass) in X if the following properties hold:

- (i) $T(\pi_0, \pi_1^j, \dots, \pi_k^j) \rightarrow T(\pi_0, \pi_1, \dots, \pi_k)$ whenever $\sup_{i,j} \text{Lip}(\pi_i^j) < \infty$ and π_i^j converges pointwise to π_i .
- (ii) if $\{x \in X : \pi_0(x) \neq 0\}$ is contained in the union $\bigcup_{i=1}^k B_i$ of Borel sets B_i and if π_i is constant on B_i , then $T(\pi_0, \dots, \pi_k) = 0$.
- (iii) there exists a finite Borel measure μ on X such that for all $(\pi_0, \dots, \pi_k) \in \mathcal{D}^k(X)$

$$|T(\pi_0, \dots, \pi_k)| \leq \prod_{i=1}^k \text{Lip}(\pi_i) \int_X |\pi_0(x)| d\mu(x). \tag{3.1}$$

The minimal Borel measure μ satisfying (3.1) is denoted $\|T\|$. The closed set

$$\text{spt } T := \{x \in X : \|T\|(U(x, r)) > 0 \text{ for all } r > 0\}$$

is called the support of T , and $\mathbf{M}(T) = \|T\|(X)$ is the mass of T . Let $\mathbf{M}_k(X)$ be the space of metric k -currents in X . When equipped with the mass norm $\mathbf{M}(\cdot)$, this becomes a Banach space. If $T \in \mathbf{M}_k(X)$ and $T \neq 0$, then $\dim_N(\text{spt } T) \geq k$; see [51, Proposition 2.5]. In particular, one has $\mathbf{M}_k(X) = \{0\}$ whenever $k > \dim_N(X)$.

Let Y be another complete metric space and let $f: X \rightarrow Y$ be a Lipschitz map. The push-forward of $T \in \mathbf{M}_k(X)$ under f is the element $f_{\#}T \in \mathbf{M}_k(Y)$ defined by

$$f_{\#}T(\pi_0, \dots, \pi_k) = T(\pi_0 \circ f, \dots, \pi_k \circ f)$$

for all $(\pi_0, \dots, \pi_k) \in \mathcal{D}^k(Y)$. We have $\text{spt } f_{\#}T \subset \overline{f(\text{spt } T)}$ and $\mathbf{M}(f_{\#}T) \leq \text{Lip}(f)^k \mathbf{M}(T)$. If $T \in \mathbf{M}_k(X)$ then the restriction $T \llcorner B$ to a Borel set $B \subset X$ is defined by

$$T \llcorner B(\pi_0, \dots, \pi_k) := T(\mathbb{1}_B \pi_0, \pi_1, \dots, \pi_k)$$

for all $(\pi_0, \dots, \pi_k) \in \mathcal{D}^k(X)$, where $\mathbb{1}_B$ is the characteristic function of B . This is well-defined since T admits a unique extension to a functional on $(k + 1)$ -tuples for which

the first argument is a bounded Borel function. One has $T \llcorner B \in \mathbf{M}_k(X)$ and $\|T \llcorner B\| = \|T\| \llcorner B$. Moreover, we have $(f_{\#}T) \llcorner B = f_{\#}(T \llcorner f^{-1}(B))$ whenever $f: X \rightarrow Y$ is Lipschitz and $B \subset Y$ is Borel.

A sequence $(T_i) \subset \mathbf{M}_k(X)$ is said to *converge weakly* to $T \in \mathbf{M}_k(X)$ if (T_i) converges pointwise to T , thus for all $(\pi_0, \dots, \pi_k) \in \mathcal{D}^k(X)$ we have $T_i(\pi_0, \dots, \pi_k) \rightarrow T(\pi_0, \dots, \pi_k)$ as $i \rightarrow \infty$.

3.2. Boundaries, slicings and integral currents

When $k \geq 1$, the boundary of $T \in \mathbf{M}_k(X)$ is the function $\partial T: \mathcal{D}^{k-1}(X) \rightarrow \mathbb{R}$ given by

$$\partial T(\pi_0, \dots, \pi_{k-1}) := T(1, \pi_0, \pi_1, \dots, \pi_{k-1}).$$

Any $T \in \mathbf{M}_k(X)$, $k \geq 1$, satisfying $\partial T = 0$ is called a cycle. If $T \in \mathbf{M}_0(X)$ then we say that T is a cycle provided that $T(1) = 0$. Notice that ∂T always satisfies (i) and (ii) of Definition 3.1. If $\partial T \in \mathbf{M}_{k-1}(X)$ then T is called *normal* current. For such T we have $\text{spt}(\partial T) \subset \text{spt} T$. We denote by $\mathbf{N}_k(X)$ the space of normal k -currents for $k \geq 1$, and we set $\mathbf{N}_0(X) := \mathbf{M}_0(X)$. Notice that $\partial(\partial T) = 0$ for all $T \in \mathbf{N}_k(X)$ with $k \geq 2$. If $\theta \in L^1(\mathbb{R}^k)$ then the function $[\theta]: \mathcal{D}^k(\mathbb{R}^k) \rightarrow \mathbb{R}$ given by

$$[\theta](\pi_0, \dots, \pi_k) := \int_{\mathbb{R}^k} \theta \pi_0 \det\left([\partial_i \pi_j]_{i,j=1}^k\right) d\mathcal{L}^k$$

defines an element of $\mathbf{M}_k(\mathbb{R}^k)$ and satisfies $\|[\theta]\| = |\theta| \mathcal{L}^k$. If θ has bounded variation then $[\theta] \in \mathbf{N}_k(\mathbb{R}^k)$. In particular, if $B \subset \mathbb{R}^k$ is a bounded Borel set of finite perimeter then $[B] := [\mathbb{1}_B]$ belongs to $\mathbf{N}_k(\mathbb{R}^k)$.

Definition 3.2. An element $T \in \mathbf{N}_k(X)$ with $k \geq 1$ is called an *integral current* if $\|T\|$ is concentrated on a countably \mathcal{H}^k -rectifiable set and if for any Lipschitz map $f: X \rightarrow \mathbb{R}^k$ and any open set $U \subset X$ there exists $\theta \in L^1(\mathbb{R}^k, \mathbb{Z})$ such that $f_{\#}(T \llcorner U) = [\theta]$.

Furthermore, $T \in \mathbf{N}_0(X)$ is called an integral current if there exist $x_1, \dots, x_m \in X$ and $\theta_1, \dots, \theta_m \in \mathbb{Z}$ such that $T = \sum_{i=1}^m \theta_i [x_i]$. Here, for $x \in X$ the current $[x] \in \mathbf{N}_0(X)$ is defined by $[x](\pi) = \pi(x)$ for every bounded Lipschitz function $\pi: X \rightarrow \mathbb{R}$. The family of integral k -currents in X is an additive subgroup of $\mathbf{N}_k(X)$ and is denoted by $\mathbf{I}_k(X)$. If $f: X \rightarrow Y$ is Lipschitz and $T \in \mathbf{I}_k(X)$ then $f_{\#}T \in \mathbf{I}_k(Y)$. In particular, if $\Delta \subset \mathbb{R}^k$ is a k -simplex and $\varphi: \Delta \rightarrow X$ a Lipschitz map then $\varphi_{\#}[\Delta] \in \mathbf{I}_k(X)$; consequently, every Lipschitz chain in X induces an integral current in X . The following lemma shows that the mass estimate $\mathbf{M}(f_{\#}T) \leq \text{Lip}(f)^k \mathbf{M}(T)$ can be improved if T is an integral current.

Lemma 3.3. *If $f: X \rightarrow Y$ is a Lipschitz map then*

$$\|f_{\#}T\|(B) \leq \int_{f^{-1}(B)} [\text{lip } f(x)]^k d\|T\|(x)$$

for every integral current $T \in \mathbf{I}_k(X)$ and every Borel set $B \subset Y$.

Proof. The inequality is obviously true when $k = 0$. If $k \geq 1$ then [37, Lemma 3.10] implies that

$$|f_{\#}T(\pi_0, \dots, \pi_k)| \leq \int_X |(\pi_0 \circ f)(x)| \prod_{i=1}^k \text{lip}(\pi_i \circ f)(x) d\|T\|(x)$$

for all $(\pi_0, \dots, \pi_k) \in \mathcal{D}^k(Y)$. Since $\text{lip}(\pi_i \circ f)(x) \leq \text{Lip}(\pi_i) \cdot \text{lip } f(x)$ for all $x \in X$, we have

$$|f_{\#}T(\pi_0, \dots, \pi_k)| \leq \prod_{i=1}^k \text{Lip}(\pi_i) \int_X |\pi_0 \circ f(x)| \cdot [\text{lip } f(x)]^k d\|T\|(x).$$

Hence, the claim follows from the definition of $\|f_{\#}T\|$. \square

If $T \in \mathbf{N}_k(X)$ with $k \geq 1$ and $u: X \rightarrow \mathbb{R}$ is Lipschitz then for almost every $r \in \mathbb{R}$,

$$\langle T, u, r \rangle := \partial(T \llcorner \{u \leq r\}) - (\partial T) \llcorner \{u \leq r\}$$

defines an element of $\mathbf{N}_{k-1}(X)$, which is called a slice of T . By the slicing theorem, the measure $\|\langle T, u, r \rangle\|$ is supported on $\text{spt } T \cap \{u = r\}$, and

$$\int_a^b \mathbf{M}(\langle T, u, r \rangle) dr \leq \text{Lip}(u) \|T\|(\{a < u < b\})$$

whenever $a < b$. Moreover, if $T \in \mathbf{I}_k(X)$ then $\langle T, u, r \rangle \in \mathbf{I}_{k-1}(X)$ for almost all $r \in \mathbb{R}$.

3.3. Homotopy formula and coning inequality

Let $\varepsilon > 0$ and let $[0, \varepsilon] \times X$ be equipped with the Euclidean product metric. For a function $f: [0, \varepsilon] \times X \rightarrow \mathbb{R}$ and $t \in [0, \varepsilon]$ we let $f_t: X \rightarrow \mathbb{R}$ be defined by $f_t(x) := f(t, x)$. If $T \in \mathbf{I}_k(X)$ for some $k \geq 0$ then the function on $\mathcal{D}^k([0, \varepsilon] \times X)$ given by

$$[[t]] \times T(\pi_0, \dots, \pi_k) := T(\pi_{0t}, \dots, \pi_{kt})$$

defines an element of $\mathbf{I}_k([0, \varepsilon] \times X)$. Moreover, the functional $[0, \varepsilon] \times T$ assigning

$$(\pi_0, \dots, \pi_{k+1}) \mapsto \sum_{i=1}^{k+1} (-1)^{i+1} \int_0^\varepsilon T\left(\pi_{0t} \frac{\partial \pi_{it}}{\partial t}, \pi_{1t}, \dots, \pi_{(i-1)t}, \pi_{(i+1)t}, \dots, \pi_{(k+1)t}\right) dt$$

is a multi-linear functional on $\mathcal{D}^{k+1}([0, \varepsilon] \times X)$ and has the following property.

Proposition 3.4. *If $T \in \mathbf{I}_k(X)$ then $\llbracket 0, \varepsilon \rrbracket \times T \in \mathbf{I}_{k+1}([0, \varepsilon] \times X)$ and*

$$\partial(\llbracket 0, \varepsilon \rrbracket \times T) + \llbracket 0, \varepsilon \rrbracket \times \partial T = \llbracket \varepsilon \rrbracket \times T - \llbracket 0 \rrbracket \times T. \tag{3.2}$$

For $k = 0$ the second term on the left-hand side is zero.

This is analogous to [42, Theorem 2.9] and [3]. Notice that the assumption made in [42, Theorem 2.9] that T has bounded support is not needed.

Given a map $h: [0, \varepsilon] \times X \rightarrow Y$ and $t \in [0, \varepsilon]$, $x \in X$, we let $h_t: X \rightarrow Y$ and $h_x: [0, \varepsilon] \rightarrow Y$ be the maps given by $h_t(z) := h(t, z)$ and $h_x(s) := h(s, x)$.

Lemma 3.5. *If $h: [0, \varepsilon] \times X \rightarrow Y$ is Lipschitz and $T \in \mathbf{I}_k(X)$, then*

$$\|h_\#(\llbracket 0, \varepsilon \rrbracket \times T)\|(B) \leq (k+1) \int_{h^{-1}(B)} \text{lip } h_x(t) \cdot [\text{lip } h_t(x)]^k d(\mathcal{L}^1 \times \|T\|)(t, x)$$

for every Borel set $B \subset Y$.

Proof. Let $(\pi_0, \dots, \pi_{k+1}) \in \mathcal{D}^{k+1}(Y)$. We have

$$\begin{aligned} & |h_\#(\llbracket 0, \varepsilon \rrbracket \times T)(\pi_0, \dots, \pi_{k+1})| \\ & \leq \sum_{i=1}^{k+1} \left| \int_0^\varepsilon T\left(\pi_0 \circ h_t \frac{\partial(\pi_i \circ h_t)}{\partial t}, \pi_1 \circ h_t, \dots, \pi_{(i-1)} \circ h_t, \pi_{(i+1)} \circ h_t, \dots, \pi_{(k+1)} \circ h_t\right) dt \right| \\ & \leq \sum_{i=1}^{k+1} \int_0^\varepsilon \int_X \left| \pi_0 \circ h_t \frac{\partial(\pi_i \circ h_t)}{\partial t} \right| \prod_{j=1, j \neq i}^{k+1} \text{lip}(\pi_j \circ h_t)(x) d\|T\|(x) dt, \end{aligned}$$

where the second inequality follows from the definition of $\|T\|$ when $k = 0$ and from [37, Lemma 3.10] when $k \geq 1$. Since

$$\left| \frac{\partial(\pi_i \circ h_t(x))}{\partial t} \right| = \text{lip}(\pi_i \circ h_x)(t)$$

for almost all $t \in [0, \varepsilon]$ and since $\text{lip}(\pi_j \circ h_t)(x) \leq \text{Lip}(\pi_j) \text{lip } h_t(x)$ we obtain

$$|h_\#(\llbracket 0, \varepsilon \rrbracket \times T)(\pi_0, \dots, \pi_{k+1})|$$

$$\leq (k + 1) \prod_{i=1}^{k+1} \text{Lip}(\pi_i) \int_0^1 \int_X |(\pi_0 \circ h)(t, x)| \text{lip } h_x(t) \cdot [\text{lip } h_t(x)]^k d\|T\| dt.$$

Hence, the claim follows from the definition of $\|h_{\#}(\llbracket 0, \varepsilon \rrbracket \times T)\|$. \square

With the homotopy formula from Proposition 3.4 and the mass estimate from Lemma 3.5 at hand it is not hard to see that Banach spaces admit coning inequalities for any $k \geq 1$.

Corollary 3.6. *Banach spaces have (CI_k) with constant $C = 1$ for every $k \geq 0$.*

Proof. Let X be a Banach space and $T \in \mathbf{I}_k(X)$ a cycle of bounded support for some $k \geq 0$. Fix $x_0 \in \text{spt } T$. The map $h: [0, 1] \times X \rightarrow X$ given by $h(t, x) = (1 - t)x_0 + tx$ is Lipschitz and satisfies $\text{lip } h_x(t) \leq \text{diam}(\text{spt } T)$ and $\text{lip } h_t(x) = t$ for all $t \in [0, 1]$ and all $x \in \text{spt } T$. Thus, by Proposition 3.4 and Lemma 3.5 the current defined by $S := h_{\#}(\llbracket 0, 1 \rrbracket \times T)$ belongs to $\mathbf{I}_{k+1}(X)$ and satisfies $\partial S = T$ as well as

$$\mathbf{M}(S) \leq (k + 1) \text{diam}(\text{spt } T) \int_{[0,1] \times X} t^k d(\mathcal{L}^1 \times \|T\|) = \text{diam}(\text{spt } T) \mathbf{M}(T),$$

as desired. Since this holds for every $k \geq 0$ the proof is complete. \square

We conclude this subsection with the following lemma which will be used in the proof of Proposition 3.8 and also later in the article.

Lemma 3.7. *Let X be a complete metric space and $T \in \mathbf{I}_k(X)$ a cycle, where $k \geq 1$. If X has (EI_k) , then for every $\varepsilon > 0$ there exists $S \in \mathbf{I}_{k+1}(X)$ with $\partial S = T$ and such that*

$$\mathbf{M}(S) \leq \text{Fillvol}_X(T) + \varepsilon \quad \text{and} \quad \text{spt } S \subset B(\text{spt } T, D \mathbf{M}(T)^{\frac{1}{k}}) \tag{3.3}$$

for some D only depending on k and on the constant in (EI_k) . All minimal fillings of T satisfy (3.3).

The lemma follows directly from [42, Lemma 3.4] and its proof. By choosing $\varepsilon > 0$ small enough we may assume that the filling S satisfies the isoperimetric inequality $\mathbf{M}(S) \leq D \mathbf{M}(T)^{\frac{k+1}{k}}$.

3.4. Mapping polyhedral currents to spaces with (LC_k) or (EI_k)

The aim of this subsection is to prove the technical Proposition 3.8 below which will be important in the proofs of Theorems 1.1 and 1.4.

Let Σ be a simplicial complex, equipped with the length metric or the ℓ_2 -metric. We define the set $\mathcal{P}_k(\Sigma) \subset \mathbf{I}_k(\Sigma)$ of polyhedral k -chains in Σ as follows. One has $P \in \mathcal{P}_k(\Sigma)$

if and only if there exist finitely many $\sigma_i \in \mathcal{F}_k$, $\theta_i \in \mathbb{Z}$, and isometries $\varphi_i: \Delta \rightarrow \sigma_i$, where Δ denotes the standard Euclidean k -simplex, such that

$$P = \sum \theta_i \varphi_{i\#} \llbracket \Delta \rrbracket.$$

Since Δ can be embedded isometrically into \mathbb{R}^k , the integral current $\llbracket \Delta \rrbracket$ is well-defined. By construction, $\mathcal{P}_k(\Sigma) \subset \mathbf{I}_k(\Sigma)$ is an additive subgroup with generating set $\{\llbracket \sigma \rrbracket : \sigma \in \mathcal{F}_k\}$. We use the convention that $\llbracket \sigma \rrbracket$ denotes any of the currents $\varphi_{\#} \llbracket \Delta \rrbracket$, where $\varphi: \Delta \rightarrow \sigma$ is an isometry. Notice that $\llbracket \sigma \rrbracket$ is uniquely determined up to a sign.

Given a Lipschitz map $h: \Sigma \rightarrow Y$ such that $h(\Sigma^{(0)}) \subset X$, the following proposition constructs a chain homomorphism $\Lambda_*: \mathcal{P}_*(\Sigma) \rightarrow \mathbf{I}_*(X)$ and a chain homotopy $\Gamma_*: \mathcal{P}_*(\Sigma) \rightarrow \mathbf{I}_{*+1}(X)$ from Λ_* to $h_{\#}$. This chain homotopy connects Λ_* to $h_{\#}$ in the sense that if $T \in \mathcal{P}_m(\Sigma)$ is a m -cycle, then $\Gamma_k(T)$ is an $(m + 1)$ -chain such that $\partial \Gamma_m(T) = \Lambda_m(T) - h_{\#}T$.

Proposition 3.8. *Let Y be complete metric space, $X \subset Y$ a closed quasiconvex subset and Σ a simplicial complex equipped with the length metric or ℓ_2 -metric, both rescaled by a factor $\varepsilon > 0$. Suppose that $h: \Sigma \rightarrow Y$ is a map such that $h(\Sigma^{(0)}) \subset X$ and $h|_{\sigma}$ is Lipschitz for every $\sigma \in \mathcal{F}$. Let $k \geq 0$ and suppose Y has (EI_{k+1}) and X has (LC_k) or (EI_k) . Then there exist homomorphisms $\Lambda_m: \mathcal{P}_m(\Sigma) \rightarrow \mathbf{I}_m(X)$ and $\Gamma_m: \mathcal{P}_m(\Sigma) \rightarrow \mathbf{I}_{m+1}(Y)$ for $m = 0, \dots, k + 1$ with the following properties:*

- (1) $\Lambda_0 = h_{\#}$ and $\Gamma_0 = 0$,
- (2) $\Lambda_m \circ \partial = \partial \circ \Lambda_{m+1}$ for $m = 0, \dots, k$,
- (3) $\partial \circ \Gamma_m = \Lambda_m - h_{\#} - \Gamma_{m-1} \circ \partial$ for $m = 1, \dots, k + 1$,
- (4) For each $m = 0, \dots, k + 1$ and each $\sigma \in \mathcal{F}_m$,

$$\mathbf{M}(\Lambda_m(\llbracket \sigma \rrbracket)) \leq C \cdot [\varepsilon \text{Lip}(h|_{\sigma})]^m$$

and

$$\mathbf{M}(\Gamma_m(\llbracket \sigma \rrbracket)) \leq C \cdot [\varepsilon \text{Lip}(h|_{\sigma})]^{m+1},$$

- (5) For each $m = 0, \dots, k + 1$ and $\sigma \in \mathcal{F}_m$ the currents $\Lambda_m(\llbracket \sigma \rrbracket)$ and $\Gamma_m(\llbracket \sigma \rrbracket)$ have support in $B(h(\sigma^{(0)}), C\varepsilon \text{Lip}(h|_{\sigma}))$.

The constant C depends only on k and the data of X and Y .

We use the convention that $0^0 = 1$.

Proof. By a simple scaling argument, it suffices to consider the case when $\varepsilon = 1$. We first construct the homomorphisms Λ_m . If X has (LC_k) , then there exists a map $\bar{h}: \Sigma^{(k+1)} \rightarrow X$ which agrees with h on $\Sigma^{(0)}$ and which is $C \text{Lip}(h|_{\sigma})$ -Lipschitz on every simplex

$\sigma \in \Sigma^{(k+1)}$. In this case we define $\Lambda_m(\llbracket \sigma \rrbracket) := \bar{h}_\# \llbracket \sigma \rrbracket$ for every oriented m -simplex $\sigma \in \Sigma$ with $0 \leq m \leq k + 1$ and extend Λ_m linearly to $\mathcal{P}_m(\Sigma)$. This yields a homomorphism Λ_m with the desired properties.

Next, we construct Λ_m in the case when X has (EI_k) . Set $\Lambda_0 := h_\#$. If σ is an oriented 1-simplex in Σ with vertices e_- and e_+ we let $\Lambda_1(\llbracket \sigma \rrbracket)$ be the integral current induced by a Lipschitz curve γ in X from $h(e_-)$ to $h(e_+)$ of length $\ell(\gamma) \leq C_0 d(h(e_-), h(e_+))$, where C_0 is the quasiconvexity constant of X . In particular, we have $\partial \Lambda_1(\llbracket \sigma \rrbracket) = \Lambda_0(\llbracket \partial \sigma \rrbracket)$ and

$$\mathbf{M}(\Lambda_1(\llbracket \sigma \rrbracket)) \leq \ell(\gamma) \leq C_1 \text{Lip}(h|_\sigma)$$

and $\text{spt}(\Lambda_1(\llbracket \sigma \rrbracket)) \subset B(h(\sigma^{(0)}), C_1 \text{Lip}(h|_{\sigma^{(0)}}))$, where $C_1 := \sqrt{2}C_0$. Doing this for every oriented 1-simplex σ and extending linearly we obtain the desired homomorphism $\Lambda_1: \mathcal{P}_1(\Sigma) \rightarrow \mathbf{I}_1(X)$. If $k = 0$ this finishes the construction of the homomorphisms Λ_m . If $k \geq 1$, then suppose we have defined Λ_m with properties (2), (4), (5) for some $1 \leq m \leq k$. Let σ be an oriented $(m + 1)$ -simplex in Σ . It follows from (2) that $\Lambda_m(\partial \llbracket \sigma \rrbracket)$ is a cycle in $\mathbf{I}_m(X)$. Thus, by Lemma 3.7, there exists $S \in \mathbf{I}_{m+1}(X)$ such that $\partial S = \Lambda_m(\partial \llbracket \sigma \rrbracket)$ and

$$\mathbf{M}(S) \leq D \mathbf{M}(\Lambda_m(\partial \llbracket \sigma \rrbracket))^{\frac{m+1}{m}} \leq D \left[(m + 2) C_m \text{Lip}(h|_\sigma)^m \right]^{\frac{m+1}{m}} \leq C_{m+1} \text{Lip}(h|_\sigma)^{m+1}$$

and

$$\text{spt}(S) \subset B\left(\text{spt}(\Lambda_m(\partial \llbracket \sigma \rrbracket)), D \mathbf{M}(\Lambda_m(\partial \llbracket \sigma \rrbracket))^{\frac{1}{m}}\right) \subset B(h(\sigma^{(0)}), C_{m+1} \text{Lip}(h|_\sigma)),$$

where $C_{m+1} := C_m + D((m + 2)C_m)^{\frac{m+1}{m}}$. We set $\Lambda_{m+1}(\llbracket \sigma \rrbracket) := S$ and by defining $\Lambda_{m+1}(\llbracket \sigma \rrbracket)$ like this for every oriented $(m + 1)$ -simplex and extending linearly we obtain a homomorphism $\Lambda_{m+1}: \mathcal{P}_{m+1}(\Sigma) \rightarrow \mathbf{I}_{m+1}(X)$ satisfying properties (2), (4), (5).

We proceed analogously in order to construct Γ_m . Set $\Gamma_0 := 0$ and suppose we have already constructed Γ_{m-1} with the desired properties for some $1 \leq m \leq k + 1$. Let σ be an oriented m -simplex in Σ . The current $P := \Lambda_m(\llbracket \sigma \rrbracket) - h_\# \llbracket \sigma \rrbracket - \Gamma_{m-1}(\llbracket \sigma \rrbracket)$ is a cycle and satisfies

$$\mathbf{M}(P) \leq \mathbf{M}(\Lambda_m(\llbracket \sigma \rrbracket)) + \mathbf{M}(h_\# \llbracket \sigma \rrbracket) + \mathbf{M}(\Gamma_{m-1}(\llbracket \sigma \rrbracket)) \leq C''_m \text{Lip}(h|_\sigma)^m,$$

where $C''_m := C_m + 1 + C'_{m-1}$. Therefore, by Lemma 3.7, there exists $S \in \mathbf{I}_{m+1}(Y)$ with $\partial S = P$ and

$$\mathbf{M}(S) \leq D' \mathbf{M}(P)^{\frac{m+1}{m}} \leq D' C''_m{}^{\frac{m+1}{m}} \text{Lip}(h|_\sigma)^{m+1} \leq C'_m \text{Lip}(h|_\sigma)^{m+1}$$

and

$$\text{spt}(S) \subset B(\text{spt}(P), D' \mathbf{M}(P)^{\frac{1}{m}}) \subset B(h(\sigma^{(0)}), C'_m \text{Lip}(h|_\sigma)),$$

where $C'_m := C''_m + D'C''_m \frac{m+1}{m}$. We put $\Gamma_m(\llbracket \sigma \rrbracket) := S$. Doing this for all σ and extending linearly yields the desired homomorphism Γ_m . It is clear that it satisfies all the desired properties. We set $C := \max\{C_m \cdot C'_m : m = 0, \dots, k + 1\}$. This completes the proof. \square

4. Factorization through simplicial complexes

In this section, we prove the factorization theorem sketched in Section 1.4. The theorem is obtained by a close inspection of the proofs of Lang and Schlichenmaier’s results in [29].

Theorem 4.1. *Let $Z \subset Y$ be complete metric spaces such that Z has Nagata dimension n and Y is Lipschitz n -connected for some integer $n \geq 0$. Then there exist $C \geq 1$, an $(n+1)$ -dimensional simplicial complex Σ equipped with the ℓ_2 -metric, and maps $g: Y \setminus Z \rightarrow \Sigma$ and $h: \Sigma \rightarrow Y$ with $h(\Sigma^{(0)}) \subset Z$ such that the following holds:*

- (1) $f: Y \rightarrow Y$, defined by $f = h \circ g$ on $Y \setminus Z$ and $f = \text{id}_Z$ on Z , is C -Lipschitz;
- (2) g is Cr^{-1} -Lipschitz on $\{y \in Y : d(y, Z) > r\}$ for all $r > 0$;
- (3) h is Cr_σ -Lipschitz on every $\sigma \in \mathcal{F}$, where $r_\sigma := \inf \{d(y, Z) : y \in g^{-1}(\text{st } \sigma)\}$;
- (4) for every $\sigma \in \mathcal{F}$ one has $R_\sigma \leq Cr_\sigma$, where $R_\sigma := \sup \{d(y, Z) : y \in g^{-1}(\text{st } \sigma)\}$.

Moreover, if $B \subset Y$ is a bounded subset intersecting Z , then there exists a simplicial complex $\Sigma' \subset \Sigma$ such that Σ' is $(n + 1)$ -dimensional and C -quasiconvex, $g(B \setminus Z) \subset \Sigma'$ and h is $C \text{ diam}(B)$ -Lipschitz on Σ' .

The constant C depends only on the data of Z and Y . Theorem 4.1 is a crucial component of the proof of Theorem 1.1. If $X \subset Y$ is a closed subset which contains Z and is Lipschitz k -connected for some $k \geq 0$ then we may choose h to furthermore satisfy $h(\Sigma^{(k+1)}) \subset X$. As a result, if Z is Lipschitz n -connected, then it follows that f is a C -Lipschitz retraction onto Z .

Proof. We may suppose that $Y \setminus Z \neq \emptyset$. Following [29, Theorems 1.6 and 5.2], we find an infinite index set I , a covering $(B_i)_{i \in I}$ of $Y \setminus Z$ by subsets of $Y \setminus Z$, and constants $\delta \in (0, 1)$ and $\alpha > 0$ depending only on the data of Z such that the following holds. For all $i \in I$, $\text{diam}(B_i) \leq \alpha d(B_i, Z)$ and for every $y \in Y \setminus Z$ there are at most $n + 2$ indices $i \in I$ such that $\tau_i(y) > 0$. Here, $\tau_i: Y \setminus Z \rightarrow [0, +\infty)$ is defined by

$$\tau_i(y) = \max\{\delta d(B_i, Z) - d(y, B_i), 0\}.$$

We set $\Sigma := \Sigma^{(n+1)}(I)$ and $\bar{\tau}(y) := \sum_{i \in I} \tau_i(y)$. By the above, Σ is $(n + 1)$ -dimensional and $g: Y \setminus Z \rightarrow \Sigma$ assigning $y \mapsto \bar{\tau}(y)^{-1}(\tau_i(y))_{i \in I}$ is well-defined. In the following, we prove items (1) – (4). To begin, we show that $g: Y \setminus Z \rightarrow \Sigma$ has the desired properties. Let $y, y' \in Y \setminus Z$ and denote by $K \subset I$ the set of all $i \in I$ such that $\tau_i(y) > 0$ or $\tau_i(y') > 0$. Notice that K contains at most $2(n + 2)$ elements. We estimate

$$\begin{aligned} \sum_{i \in K} \left| \frac{\tau_i(y)}{\bar{\tau}(y)} - \frac{\tau_i(y')}{\bar{\tau}(y')} \right| &= \sum_{i \in K} \frac{1}{\bar{\tau}(y)\bar{\tau}(y')} \left| \sum_{j \in K} \left(\tau_i(y)\tau_j(y') - \tau_i(y')\tau_j(y) \right) \right| \\ &\leq \frac{2}{\bar{\tau}(y)\bar{\tau}(y')} \sum_{i \in K} \sum_{j \in K} \tau_j(y') |\tau_i(y) - \tau_i(y')| \\ &\leq \frac{4(n+2)}{\bar{\tau}(y)} d(y, y'), \end{aligned}$$

where in the last inequality we used that τ_i is 1-Lipschitz. Since $(B_i)_{i \in I}$ covers $Y \setminus Z$, there exists $i \in I$ such that $y \in B_i$. As a result,

$$d(y, Z) \leq \text{diam}(B_i) + d(B_i, Z) \leq (1 + \alpha)d(B_i, Z)$$

and $\bar{\tau}(y) \geq \delta d(B_i, Z)$. Thus, $(1 + \alpha)\bar{\tau}(y) \geq (1 + \alpha)\delta d(B_i, Z) \geq \delta d(y, Z)$. By the above,

$$|g(y) - g(y')| \leq \frac{4(n+2)(1+\alpha)}{\delta} \frac{1}{d(y, Z)} d(y, y').$$

This shows that g is $C_1 r^{-1}$ -Lipschitz on $\{y \in Y : d(y, Z) > r\}$. Next, we show (3) and (4). For each $i \in I$ select $y_i \in B_i$ and $x_i \in Z$ for which $d(y_i, x_i) \leq 2d(B_i, Z)$. Let $h_0: \Sigma^{(0)} \rightarrow Z$ denote the map that sends e_i to x_i for each $i \in I$. Here, $e_i \in \ell_2(I)$ is defined by $(e_i)_j = 1$ if $j = i$ and $(e_i)_j = 0$ otherwise. Notice that $\Sigma^{(0)} = \{e_i : i \in I\}$. Using that Y is Lipschitz n -connected, we obtain a constant $C_2 > 0$ and an extension $h: \Sigma \rightarrow Y$ of h_0 such that h is $C_2 \text{Lip}(h|_{\sigma^{(0)}})$ -Lipschitz on each $\sigma \in \mathcal{F}$. Let $\sigma \in \mathcal{F}$ such that $g^{-1}(\text{st } \sigma) \neq \emptyset$. Notice that

$$r_\sigma = \inf \{d(Z, g^{-1}(\text{int } \sigma')) : \sigma' \in \mathcal{F} \text{ and } \sigma \subset \sigma'\}.$$

Thus, to show that h is $C_3 r_\sigma$ -Lipschitz on σ it suffices to show that h is $C_3 r$ -Lipschitz on each $\sigma' \in \mathcal{F}$ with $\sigma \subset \sigma'$, where $r := d(Z, g^{-1}(\text{int } \sigma'))$. Fix $\sigma' \in \mathcal{F}$ and let e_i and e_j be two vertices of σ' . We may suppose that $d(Z, B_j) \leq d(Z, B_i)$ and there exists some $y \in Y \setminus Z$ such that $g(y) \in \text{int } \sigma'$. We estimate

$$\begin{aligned} d(h(e_i), h(e_j)) &\leq d(x_i, y) + d(y, x_j) \\ &\leq 2d(Z, B_i) + d(y_i, y) + d(y_j, y) + 2d(Z, B_j). \end{aligned}$$

As $g(y) \in \text{int } \sigma'$, we infer $\tau_i(y) > 0$ and thus $d(y, B_i) \leq \delta d(Z, B_i)$. Therefore, $d(y_i, y) \leq \text{diam}(B_i) + d(y, B_i) \leq \text{diam}(B_i) + \delta d(Z, B_i)$, and we arrive at

$$d(h(e_i), h(e_j)) \leq 2(2 + \delta + \alpha)d(Z, B_i).$$

Since $(1 - \delta)d(B_i, Z) \leq d(y, Z)$, we obtain

$$d(h(e_i), h(e_j)) \leq \sqrt{2} \left(\frac{2 + \delta + \alpha}{1 - \delta} \right) d(y, Z) |e_i - e_j|.$$

As $y \in Y \setminus Z$ with $g(y) \in \text{int}(\sigma')$ was arbitrary, this implies that h is C_3r -Lipschitz on σ' for $r := d(Z, g^{-1}(\text{int} \sigma'))$, as desired. This concludes the proof of (3). If e_i is a vertex of σ , then we have

$$d(y, Z) \leq 2d(Z, B_i) + d(y_i, y) \leq (2 + \delta + \alpha)d(Z, B_i) \leq \left(\frac{2 + \delta + \alpha}{1 - \delta}\right) r_\sigma,$$

and so $R_\sigma \leq C_3r_\sigma$, as $y \in g^{-1}(\text{st} \sigma)$ was arbitrary. This yields (4).

Next, we show (1). Notice that Y is c -quasiconvex for some $c \geq 1$. We want to apply Lemma 2.4 to show that f is C_4c -Lipschitz. To this end, we prove that $\text{lip} f(y) \leq C_4$ for all $y \in Y$. For any $y \in Y \setminus Z$ there exists $i \in I$ such that $y \in B_i$ and thus using (3) we obtain $d(f(y), x_i) = d(h(g(y)), h(e_i)) \leq C_3 d(y, Z)\sqrt{2}$, and so

$$d(f(y), y) \leq d(f(y), x_i) + d(x_i, y_i) + d(y_i, y) \leq (C_3\sqrt{2} + 2 + \alpha)d(y, Z). \tag{4.1}$$

By combining $d(f(x), f(y)) \leq d(x, y) + d(y, f(y))$, where $x \in Z$, with (4.1), we find that $\text{lip} f(x) \leq (C_3\sqrt{2} + 3 + \alpha)$ for all $x \in Z$. Now, fix $y \in Y \setminus Z$ and choose $\varepsilon > 0$ such that $d(y', Z) \leq 2d(y, Z)$ for all $y' \in U_\varepsilon(y)$, and whenever $y' \in U_\varepsilon(y)$ is contained in $g^{-1}(\text{int}(\sigma))$ for some $\sigma \in \mathcal{F}$, then $y \in g^{-1}(\sigma)$. Because of (2) and (3), for all $y' \in U_\varepsilon(y)$,

$$d(f(y), f(y')) \leq C_3 d(y', Z) d(g(y), g(y')) \leq 2C_3C_1 d(y, y'),$$

and so $\text{lip} f(y) \leq 2C_3C_1$. Hence, we have shown that $\text{lip} f(y) \leq C_4$ for all $y \in Y$. By Lemma 2.4, f is C_4c -Lipschitz, as desired.

To finish the proof, we show the statements of the moreover part. Let $B \subset Y$ be a bounded subset intersecting Z and suppose that $J \subset I$ is the subset of those indices $i \in I$ for which there exists $y \in B$ such that $\tau_i(y) > 0$. By enlarging B (if necessary) we may assume that J has infinitely many elements. We set $\Sigma' := \Sigma^{(n+1)}(J)$. Clearly, $\Sigma' \subset \Sigma$ and Σ' is $(n + 1)$ -dimensional. By definition of g , we have $g(B \setminus Z) \subset \Sigma'$. Next, we claim that Σ' is C_5 -quasiconvex for some constant C_5 depending only on n . Let $\sigma_1, \sigma_2 \in \mathcal{F}(\Sigma')$. Clearly, there is $\sigma \in \mathcal{F}(\Sigma(J))$ of dimension at most $2n + 3$ such that $\sigma_1, \sigma_2 \subset \sigma^{(n+1)}$. As σ is isometric to a Euclidean standard simplex, it follows that $\sigma^{(n+1)}$ is C_5 -quasiconvex, and so, by using that $\sigma^{(n+1)} \subset \Sigma'$, we find that Σ' is C_5 -quasiconvex as well. Next, we show that h is $C_6 \text{diam}(B)$ -Lipschitz on Σ' . For each $r \in J$ choose $z_r \in B$ such that $\tau_r(z_r) > 0$. We estimate

$$\begin{aligned} d(h(e_r), h(e_s)) &\leq d(x_r, y_r) + d(y_r, z_r) + d(z_r, z_s) + d(z_s, y_s) + d(y_s, x_s) \\ &\leq \text{diam}(B) + 2(2 + \alpha + \delta) \max\{d(B_r, Z), d(B_s, Z)\}, \end{aligned}$$

for all $r, s \in J$. Fix $y_0 \in B \cap Z$. Since $(1 - \delta)d(B_s, Z) \leq d(z_s, Z)$ for all $s \in J$, it follows that

$$d(h(e_r), h(e_s)) \leq \text{diam}(B) + 2 \left(\frac{2 + \delta + \alpha}{1 - \delta}\right) \text{diam}(B);$$

thus, by construction of h , for each $\sigma \in \mathcal{F}(\Sigma')$, the map h is $C_6 \text{diam}(B)$ -Lipschitz on σ . Now, an argument as in the proof of (1) above yields $\text{lip } h(x) \leq C_6 \text{diam}(B)$ for all $x \in \Sigma'$. Hence, as Σ' is C_5 -quasiconvex, Lemma 2.4 tells us that h is $C_5 C_6 \text{diam}(B)$ -Lipschitz on Σ' , as desired. We put $C := \max\{C_i : i = 1, \dots, 6\}$. \square

If h is as in Theorem 4.1 above, then h is Cr -Lipschitz on every $\sigma \in \mathcal{F}$ for which $g^{-1}(\text{st } \sigma) \cap \{d(y, Z) \leq r\} \neq \emptyset$. Hence, the following proposition tells us that $\text{lip } h|_{\Sigma'}(x) \leq Cr$ for every $x \in \Sigma'$, where $\Sigma' := \text{Hull}(\{y \in Y : 0 < d(y, Z) \leq r\})$.

Proposition 4.2. *Let $h: \Sigma \rightarrow Y$ be a map from a simplicial complex Σ equipped with the ℓ_2 -metric to a metric space Y . Let $L > 0$ and suppose $A \subset \Sigma$ is such that h is L -Lipschitz on every $\sigma \in \mathcal{F}$ for which $\text{st } \sigma \cap A \neq \emptyset$. Then the restriction of h to $\Sigma' := \text{Hull}(A)$ satisfies $\text{lip } h|_{\Sigma'}(x) \leq L$ for every $x \in \Sigma'$.*

Proof. For $y \in \Sigma$ denote by $\sigma(y)$ be the unique simplex in Σ with $y \in \text{int } \sigma(y)$. Notice that if $y \in \Sigma'$ then $h|_{\sigma(y)}$ is L -Lipschitz because $\sigma(y) \subset \Sigma'$ and hence $\text{st } \sigma(y) \cap A \neq \emptyset$.

Now, let $x \in \Sigma'$. By Lemma 2.1 there is a real number $s_0 > 0$ such that $x \in \sigma(y)$ for all $y \in \Sigma'$ with $|x - y| < s_0$. Therefore, if $y \in \Sigma'$ satisfies $0 < |x - y| < s < s_0$ then, by the above,

$$\frac{d(h(x), h(y))}{s} \leq \frac{d(h(x), h(y))}{|x - y|} \leq L.$$

This implies that $\text{lip } h|_{\Sigma'}(x) \leq L$ for all $x \in \Sigma'$, as was to be shown. \square

Corollary 4.3. *Suppose that $h: \Sigma \rightarrow X$ is a homeomorphism between a simplicial complex Σ equipped with the ℓ_2 -metric and a metric space X . If h is D -bilipschitz on every $\sigma \in \mathcal{F}$ and X is c -quasiconvex, then $h: \Sigma \rightarrow X$ is cD -bilipschitz when Σ is equipped with the length metric.*

Proof. Proposition 4.2 tells us that $\text{lip } h(z) \leq D$ for all $z \in \Sigma$. Let $z, z' \in \Sigma$ and let γ be a curve in Σ connecting z and z' . By Lemma 2.4, it follows that

$$d(h(z), h(z')) \leq \ell(h \circ \gamma) \leq D\ell(\gamma).$$

Consequently, h is D -Lipschitz with respect to the length metric d_i on Σ . To finish the proof it remains to show that $g: X \rightarrow (\Sigma, d_i)$ given by $x \mapsto h^{-1}(x)$ is cD -Lipschitz.

For every $z \in \Sigma$ denote by $\sigma(z)$ the unique simplex $\sigma \in \mathcal{F}$ such that $z \in \text{int } \sigma$. Given $x \in X$, we abbreviate $\sigma(x) := h(\sigma(z))$, where $z = g(x)$. As h is a homeomorphism, for every $x \in X$ there exists $\varepsilon_x > 0$ such that $x \in \sigma(x')$ whenever $x' \in U(x, \varepsilon_x)$. See Lemma 2.1. Notice that g is D -Lipschitz on $\sigma(x)$ for every $x \in X$. Hence, exactly the same argument as in the proof of Proposition 4.2 shows that $\text{lip } g(x) \leq D$ for all $x \in X$. Now, by invoking Lemma 2.4 we get that g is cD -Lipschitz, as X is c -quasiconvex. This completes the proof. \square

5. Proof of Theorem 1.1 and generalizations

In this section we prove Theorem 1.1 from the introduction about isoperimetric subspace distortion. We also establish strengthenings and generalizations of this theorem which will be used in Section 7 to establish Theorem 1.4, a deformation theorem in spaces of finite Nagata dimension.

5.1. Undistorted fillings with controlled support

In this subsection, we discuss two theorems on isoperimetric subspace distortion and deduce Theorem 1.1 from them. The main difference between the following theorem and Theorem 1.1 is that we do not assume that the whole subspace X has finite Nagata dimension and that in addition we obtain control on the filling chain in the subspace X .

Theorem 5.1. *Let Y be a complete metric space and let $Z \subset X \subset Y$ be closed subsets such that Z is bounded and has finite Nagata dimension, and X is quasiconvex and has (LC_k) or (EI_k) for some $k \geq 0$. If $S \in \mathbf{I}_{k+1}(Y)$ satisfies $\text{spt}(\partial S) \subset Z$ and $\text{spt } S \subset B(Z, \eta)$ for some $\eta > 0$, then there exists $\bar{S} \in \mathbf{I}_{k+1}(Y)$ which has support in X and satisfies $\partial \bar{S} = \partial S$ as well as*

$$\mathbf{M}(\bar{S}) \leq C \mathbf{M}(S) \quad \text{and} \quad \text{spt } \bar{S} \subset B(Z, C\eta),$$

where C depends only on the data of X and the Nagata dimension and constant of Z . Moreover, if Y is a Banach space then there exists $W \in \mathbf{I}_{k+2}(Y)$ with $\partial W = \bar{S} - S$ and such that $\mathbf{M}(W) \leq \eta C \mathbf{M}(S)$ and $\text{spt } W \subset B(Z, C\eta)$.

Notice that the constant C is independent of the diameter of Z . The hypothesis that Z be bounded can be dropped if one assumes that the whole subspace X has finite Nagata dimension, see Theorem 5.3 below. Theorem 5.1 will be proved in Subsection 5.2. As a consequence of the theorem we obtain the following strengthening of Corollary 1.3.

Corollary 5.2. *Let X be a complete quasiconvex metric space of finite Nagata dimension. Suppose that X has (LC_k) or (EI_k) for some $k \geq 0$. Then every cycle $T \in \mathbf{I}_m(X)$ of bounded support with $m = 0, \dots, k$ admits a filling $S \in \mathbf{I}_{m+1}(X)$ satisfying*

$$\mathbf{M}(S) \leq C \text{diam}(\text{spt } T) \cdot \mathbf{M}(T) \quad \text{and} \quad \text{diam}(\text{spt } S) \leq C \text{diam}(\text{spt } T)$$

for some constant $C > 0$ depending only on the data of X . In particular, X has (CI_k) .

The stronger version of the coning inequality asserted in the corollary is called *strong coning inequality* in [24] and [23] and plays an important role in these papers.

Proof. Let $m \in \{0, \dots, k\}$ be an integer and suppose that $T \in \mathbf{I}_m(X)$ is a cycle of bounded support. View X as a subset of $Y := \ell_\infty(X)$ via a Kuratowski embedding. Fix $x_0 \in \text{spt } T$ and define $h: [0, 1] \times Y \rightarrow Y$ by $h(t, y) := (1 - t)x_0 + ty$. The integral current $S := h_{\#}(\llbracket 0, 1 \rrbracket \times T)$ satisfies $\partial S = T$ and $\mathbf{M}(S) \leq \text{diam}(\text{spt } T) \mathbf{M}(T)$ as well as

$$\text{spt } S \subset h(\llbracket 0, 1 \rrbracket \times \text{spt } T) \subset B(\text{spt } T, \text{diam}(\text{spt } T));$$

see Section 3.3. Applying Theorem 5.1 with $Z = \text{spt } T$, we obtain $\bar{S} \in \mathbf{I}_{m+1}(X)$ with $\partial \bar{S} = T$ and such that

$$\mathbf{M}(\bar{S}) \leq C \mathbf{M}(S) \leq C \text{diam}(\text{spt } T) \mathbf{M}(T)$$

as well as

$$\text{spt } \bar{S} \subset B(\text{spt } T, C \text{diam}(\text{spt } T))$$

for some constant C only depending on the data of X . In particular, it follows that $\text{diam}(\text{spt } \bar{S}) \leq (2C + 1) \text{diam}(\text{spt } T)$. \square

Every complete metric space with (CI_k) also has (EI_k) , as was shown in [42]. Hence, by Corollary 5.2, if X is quasiconvex, has finite Nagata dimension and satisfies (LC_k) for some $k \geq 1$, then X has (EI_k) . In the following, we use this fact in combination with Theorem 5.1, to show that when X has finite Nagata dimension, then the conclusion of Theorem 5.1 is also valid for currents $S \in \mathbf{I}_{k+1}(Y)$ for which the support of ∂S is unbounded.

Theorem 5.3. *Let Y be a complete metric space and let $X \subset Y$ be a closed quasiconvex subset of finite Nagata dimension. Suppose furthermore that X has (LC_k) or (EI_k) for some $k \geq 0$. Then for all subsets $Z \subset X$ and all $S \in \mathbf{I}_{k+1}(Y)$ with $\text{spt } \partial S \subset Z$ and $\text{spt } S \subset B(Z, \eta)$ for some $\eta > 0$, there exists $\bar{S} \in \mathbf{I}_{k+1}(Y)$ which has support in X and satisfies $\partial \bar{S} = \partial S$ as well as*

$$\mathbf{M}(\bar{S}) \leq C \mathbf{M}(S) \quad \text{and} \quad \text{spt } \bar{S} \subset B(Z, C\eta),$$

where C depends only on the data of X . Moreover, if Y is a Banach space then there exists $W \in \mathbf{I}_{k+2}(Y)$ with $\partial W = \bar{S} - S$ and such that $\mathbf{M}(W) \leq \eta C \mathbf{M}(S)$ and $\text{spt } W \subset B(Z, C\eta)$.

Proof. We first treat the case $k \geq 1$. The space X has (CI_k) by Corollary 5.2 and hence also (EI_k) by [42]. By possibly embedding Y isometrically into $\ell_\infty(Y)$ we may further assume that Y has (EI_k) . Lemma 3.7 thus implies that for every $\eta > 0$ the following holds. For every $\varepsilon > 0$ sufficiently small and every cycle $U \in \mathbf{I}_m(Y)$ with $1 \leq m \leq k$ and $\mathbf{M}(U) \leq \varepsilon$ there is a filling $V \in \mathbf{I}_{m+1}(Y)$ with $\mathbf{M}(V) \leq \varepsilon$ and $\text{spt } V \subset B(\text{spt } U, \eta)$; if $\text{spt } U \subset X$ then $\text{spt } V \subset X$.

Let $Z, S,$ and η be as in the statement of the theorem and set $T := \partial S$. Fix $y_0 \in Y$ and let $\varrho: Y \rightarrow \mathbb{R}$ be the distance function from y_0 . Notice that $\langle T, \varrho, r \rangle$ and $\langle S, \varrho, r \rangle$ are integral currents for almost every $r > 0$ and

$$\partial(T \llcorner \{\varrho \leq r\}) = \langle T, \varrho, r \rangle = -\partial \langle S, \varrho, r \rangle.$$

Let $\varepsilon \in (0, 1)$ be sufficiently small. There exists $r > 0$ arbitrarily large such that the integral currents $\langle T, \varrho, r \rangle$ and $\langle S, \varrho, r \rangle$ are supported on $\text{spt } T \cap \{\varrho = r\}$ and $\text{spt } S \cap \{\varrho = r\}$, respectively, and satisfy $\mathbf{M}(\langle T, \varrho, r \rangle) \leq \varepsilon$ and $\mathbf{M}(\langle S, \varrho, r \rangle) \leq \varepsilon$. If $r > 0$ was chosen large enough we may also assume that $\mathbf{M}(T \llcorner \{\varrho > r\}) \leq \varepsilon$ and $\mathbf{M}(S \llcorner \{\varrho > r\}) \leq \varepsilon$.

In the following, we consider the cases $k = 1$ and $k \geq 2$ separately. Assume first that $k \geq 2$. For $s, t > 0$ we abbreviate $A(s, t) := B(Z, s) \cap B(y_0, t)$. By the observation made at the beginning of the proof there exists a filling $T' \in \mathbf{I}_k(Y)$ of the cycle $\langle T, \varrho, r \rangle$ with $\mathbf{M}(T') \leq \varepsilon$ and $\text{spt } T' \subset X \cap A(\eta, r + \eta)$. Define cycles by

$$T_1 := T \llcorner \{\varrho > r\} + T' \quad \text{and} \quad T_2 := T \llcorner \{\varrho \leq r\} - T'$$

and notice that $T = T_1 + T_2$. We will find a filling of T by constructing suitable fillings of T_1 and T_2 . Since T_1 has mass at most 2ε and support in $X \cap B(Z, \eta)$ it has a filling \bar{S}_1 of mass at most 2ε and support in $X \cap B(Z, 2\eta)$, again by the observation at the beginning of the proof. Next, we construct a filling of T_2 . Notice that T_2 has support in the set $Z' := X \cap A(\eta, r + 5\eta)$. The cycle $T' + \langle S, \varrho, r \rangle$ is contained in $\mathbf{I}_k(Y)$, has mass less than or equal to 2ε and is supported in $A(\eta, r + \eta)$, so it has a filling $S' \in \mathbf{I}_{k+1}(Y)$ with $\mathbf{M}(S') \leq 2\varepsilon$ and support in $A(2\eta, r + 2\eta)$. Thus, $S_2 := S \llcorner \{\varrho \leq r\} - S'$ is a filling of T_2 satisfying $\text{spt } S_2 \subset A(2\eta, r + 2\eta) \subset B(Z', 2\eta)$ and $\mathbf{M}(S_2) \leq \mathbf{M}(S) + 2\varepsilon$. Hence, by Theorem 5.1, there exists a filling $\bar{S}_2 \in \mathbf{I}_{k+1}(Y)$ of T_2 with support in X such that

$$\mathbf{M}(\bar{S}_2) \leq C \mathbf{M}(S_2) \leq C \mathbf{M}(S) + 2C\varepsilon$$

and $\text{spt } \bar{S}_2 \subset B(Z', 2C\eta) \subset B(Z, 3C\eta)$ for some constant $C \geq 1$ only depending on the data of X . It follows that $\bar{S} := \bar{S}_1 + \bar{S}_2$ is a filling of T with support in X and satisfying $\text{spt } \bar{S} \subset B(Z, 3C\eta)$ and

$$\mathbf{M}(\bar{S}) \leq \mathbf{M}(\bar{S}_1) + \mathbf{M}(\bar{S}_2) \leq C \mathbf{M}(S) + 2(C + 1)\varepsilon \leq 2C \mathbf{M}(S),$$

where the last inequality holds provided $\varepsilon > 0$ was chosen small enough.

In order to prove the last statement in the theorem, suppose Y is a Banach space. Set $R_1 := \bar{S}_1 - S' - S \llcorner \{\varrho > r\}$ and notice that R_1 is an integral cycle of mass at most 5ε and with support in $B(Z, 2\eta)$. Therefore, if $\varepsilon > 0$ was chosen sufficiently small, then R_1 has a filling $W_1 \in \mathbf{I}_{k+2}(Y)$ with $\mathbf{M}(W_1) \leq 5\varepsilon$ and $\text{spt } W_1 \subset B(Z, 3\eta)$, compare with the observation at the beginning of the proof. Moreover, by Theorem 5.1, there exists $W_2 \in \mathbf{I}_{k+2}(Y)$ with $\partial W_2 = \bar{S}_2 - S_2$ and $\text{spt } W_2 \subset B(Z, 3C\eta)$ and such that $\mathbf{M}(W_2) \leq 3\eta C \mathbf{M}(S_2)$. Consequently, if $\varepsilon > 0$ was chosen small enough, the integral

current $W := W_1 + W_2$ satisfies $\partial W = \bar{S} - S$, has support in $B(Z, C\eta)$ and mass bounded by $\eta C \mathbf{M}(S)$ for some C only depending on the data of X . This concludes the proof of the case $k \geq 2$.

If $k = 1$, then $\mathbf{M}(\langle T, \varrho, r \rangle)$ is a non-negative integer satisfying $\mathbf{M}(\langle T, \varrho, r \rangle) < \varepsilon$. Hence, $\langle T, \varrho, r \rangle = 0$ and it follows that $T_1 = T \llcorner \{\varrho > r\}$, $T_2 = T \llcorner \{\varrho \leq r\}$ and $\langle S, \varrho, r \rangle$ are integral cycles. Now, the same reasoning as in the proof of the case $k \geq 2$ yields a filling \bar{S} of T with the desired properties and, if Y is a Banach space, a filling W of $\bar{S} - S$ with the desired properties. This completes the proof when $k \geq 1$.

We finally treat the remaining case $k = 0$. Let ϱ be as above. Similarly to the case $k = 1$ there exist arbitrarily large $r > 0$ such that $\langle S, \varrho, r \rangle = 0$. Thus, there clearly exists r such that the integral current $S' := S \llcorner \{\varrho \leq r\}$ satisfies $\partial S' = \partial S$ and has bounded support. Therefore, applying Theorem 5.1 to S' we obtain a suitable filling of ∂S in X . The moreover part can easily be proved as above. \square

We can finally prove Theorem 1.1.

Proof of Theorem 1.1. Let X and Y be as in the statement of the theorem and let $T \in \mathbf{I}_m(X)$ be a non-trivial cycle for some $m = 0, \dots, k$. By replacing Y by $\ell_\infty(Y)$ we may assume that Y is quasiconvex and has (EI_k) . In particular, $\text{Fillvol}_Y(T)$ is finite. To apply Theorem 5.3, we must find a filling $S \in \mathbf{I}_{m+1}(Y)$ of T with $\mathbf{M}(S) \leq 2 \text{Fillvol}_Y(T)$ such that $\text{spt } S \subset B(\text{spt } T, \eta)$ for some finite $\eta > 0$. When $m \geq 1$, Lemma 3.7 implies that such an S exists. When $m = 0$ then for any filling S of T with $\mathbf{M}(S) \leq 2 \text{Fillvol}_Y(T)$ one constructs a filling of T with smaller mass and bounded support by restricting S to a suitable ball as at the end of the proof of Theorem 5.3. In either case, we apply Theorem 5.3 to S to show that there exists a filling $\bar{S} \in \mathbf{I}_{m+1}(Y)$ of T with support in X such that $\mathbf{M}(\bar{S}) \leq C \mathbf{M}(S) \leq 2C \text{Fillvol}_Y(T)$ for some constant C depending only on the data of X . This completes the proof. \square

5.2. Proof of Theorem 5.1

Let Z, X, Y, S, η be as in the statement of Theorem 5.1. We may of course assume that $\partial S \neq 0$ and thus $k \leq \dim_N(Z)$; see Section 3.1. By possibly embedding Y isometrically into $\ell_\infty(Y)$ we may assume that Y itself is a Banach space. In particular, Y has (LC_n) and (EI_n) for all $n \in \mathbb{N}$, with constants only depending on n (see [42]). Let $\varrho: Y \rightarrow \mathbb{R}$ be the distance function from the set Z .

Lemma 5.4. *We may assume that*

$$\|S\|(\{\varrho \leq r\}) \leq r(k+1) \mathbf{M}(\partial S) \quad \text{and} \quad \mathbf{M}(\partial(S \llcorner \{\varrho > r\})) \leq \mathbf{M}(\partial S) \quad (5.1)$$

for almost every $r > 0$ small enough.

Proof. It is clearly enough to find a metric space \hat{Y} containing Y and a current $\hat{S} \in \mathbf{I}_{k+1}(\hat{Y})$ satisfying the properties stated in the lemma and with $\partial\hat{S} = \partial S$, $\mathbf{M}(\hat{S}) \leq 2\mathbf{M}(S)$ as well as $\text{spt } \hat{S} \subset B(Z, 2\eta)$.

Let $0 < \varepsilon < \eta$ be so small that $\varepsilon(k + 1)\mathbf{M}(\partial S) \leq \mathbf{M}(S)$. We define $\hat{Y} := [0, \varepsilon] \times Y$ and identify Y with the subset $\{0\} \times Y \subset \hat{Y}$. We set

$$\hat{S} := \llbracket \varepsilon \rrbracket \times S - \llbracket 0, \varepsilon \rrbracket \times \partial S. \tag{5.2}$$

Clearly, $\hat{S} \in \mathbf{I}_{k+1}(\hat{Y})$ and $\partial\hat{S} = \llbracket 0 \rrbracket \times \partial S$. It follows from Lemma 3.5 that

$$\mathbf{M}(\hat{S}) \leq \mathbf{M}(S) + \varepsilon(k + 1)\mathbf{M}(\partial S) \leq 2\mathbf{M}(S).$$

Since $\text{spt } \hat{S} \subset [0, \varepsilon] \times \text{spt } S$, we infer $\text{spt } \hat{S} \subset B(Z, 2\eta)$. We now show that \hat{S} satisfies (5.1). Let $\hat{\varrho}: \hat{Y} \rightarrow \mathbb{R}$ denote the distance function from Z . We claim that

$$\hat{S} \llcorner \{\hat{\varrho} \leq r\} = -\llbracket 0, r \rrbracket \times \partial S \tag{5.3}$$

for every $r \in (0, \varepsilon)$. To simplify the notation, we set $Q_r := \llbracket 0, r \rrbracket \times \partial S$. To establish (5.3), it suffices to show that $Q_\varepsilon \llcorner \{\hat{\varrho} \leq r\} = Q_r$, as $\text{spt}(\llbracket \varepsilon \rrbracket \times S) \subset \{\varepsilon\} \times \text{spt } S$ and thus $(\llbracket \varepsilon \rrbracket \times S) \llcorner \{\hat{\varrho} \leq r\} = 0$ for all $r \in (0, \varepsilon)$. Choose a sequence $\ell_j: \mathbb{R} \rightarrow [0, 1]$ of Lipschitz functions such that $\ell_j \rightarrow \mathbb{1}_{[0, r]}$ in $L^1(\mathbb{R})$ as $j \rightarrow \infty$. Letting $\hat{\varrho}_j := \ell_j \circ \hat{\varrho}$ we observe that $\hat{\varrho}_j \rightarrow \mathbb{1}_{\{\hat{\varrho} \leq r\}}$ in $L^1(\hat{Y}, \|Q_\varepsilon\|)$ as $j \rightarrow \infty$. Moreover, using that $\hat{\varrho}_{jt} = \ell_j(t)$ on Z , we find that $Q_\varepsilon(\hat{\varrho}_j \pi_0, \pi_1, \dots, \pi_{k+1})$ is equal to

$$\sum_{i=1}^{k+1} (-1)^{i+1} \int_0^\varepsilon \ell_j(t) \cdot \partial S \left(\pi_{0t} \frac{\partial \pi_{it}}{\partial t}, \pi_{1t}, \dots, \pi_{(i-1)t}, \pi_{(i+1)t}, \dots, \pi_{(k+1)t} \right) dt$$

for all $(\pi_0, \dots, \pi_{k+1}) \in \mathcal{D}^{k+1}(\hat{Y})$. Hence, by letting $j \rightarrow \infty$, we obtain $Q_\varepsilon \llcorner \{\hat{\varrho} \leq r\} = Q_r$ and (5.3) follows. Because of (5.3), we have

$$\begin{aligned} \|\hat{S}\|(\{\hat{\varrho} \leq r\}) &= \mathbf{M}(\llbracket 0, r \rrbracket \times \partial S) \leq r(k + 1)\mathbf{M}(\partial S), \\ \mathbf{M}(\partial(\hat{S} \llcorner \{\hat{\varrho} > r\})) &= \mathbf{M}(\llbracket r \rrbracket \times \partial S) = \mathbf{M}(\partial S) \end{aligned}$$

for every $r \in (0, \varepsilon)$. This completes the proof. \square

Next, denote by n the Nagata dimension of Z . By applying Theorem 4.1 with $B = B(Z, \eta)$, we obtain an $(n + 1)$ -dimensional simplicial complex Σ equipped with the ℓ_2 -metric and maps $f: Y \rightarrow Y$, $g: Y \setminus Z \rightarrow \Sigma$, and $h: \Sigma \rightarrow Y$ with $h(\Sigma^{(0)}) \subset Z$ satisfying the properties listed in the statement of that theorem. In particular, Σ is quasiconvex, $g(B \setminus Z) \subset \Sigma$ and h is Lipschitz on Σ . For the remainder of this section we denote by C the constant of Theorem 4.1. Notice that C does not depend on the diameter of Z , but the Lipschitz constant of h might.

For almost every $r > 0$ the current $S_r := S \llcorner \{\varrho > r\}$ belongs to $\mathbf{I}_{k+1}(Y)$ and $\text{spt}(\partial S_r) \subset \{\varrho = r\}$. Fix a decreasing sequence (r_i) of positive real numbers converging to zero such that $S_{r_i} \in \mathbf{I}_{k+1}(Y)$ and (5.1) holds with $r = r_i$ for all $i \geq 1$. Notice that S_{r_i} is supported on $\{\varrho \geq r_i\}$ and g is $C r_i^{-1}$ -Lipschitz on this set, so the current $S'_i := g_{\#} S_{r_i}$ is well-defined and an element of $\mathbf{I}_{k+1}(\Sigma)$. Let $P'_i \in \mathcal{P}_{k+1}(\Sigma^{(k+1)})$, $R'_i \in \mathbf{I}_{k+1}(\Sigma)$, $Q'_i \in \mathbf{I}_{k+2}(\Sigma)$ with

$$S'_i = P'_i + R'_i + \partial Q'_i \tag{5.4}$$

be the integral currents obtained by applying the deformation theorem (Theorem A.2) to S'_i . Further, let Λ_m and Γ_m be the homomorphisms from Proposition 3.8 and define

$$P_i := \Lambda_{k+1}(P'_i) \in \mathbf{I}_{k+1}(X). \tag{5.5}$$

In what follows, we will show that the sequence (P_i) converges to a filling of T which has the desired properties.

To keep track of the constants, we denote by K the maximum of the constants appearing in Theorem A.2 and Proposition 3.8. By construction, K depends only on the data of X and on the Nagata dimension and constant of Z . The next two lemmas show that the P_i 's defined in (5.5) have uniformly bounded mass and boundary mass and the sequence (∂P_i) converges weakly to ∂S .

Lemma 5.5. *For all sufficiently large $i \geq 1$ we have $\text{spt } P_i \subset B(Z, \bar{C}\eta)$ as well as*

$$\mathbf{M}(P_i) \leq \bar{C} \mathbf{M}(S) \quad \text{and} \quad \mathbf{M}(\partial P_i) \leq \bar{C} \mathbf{M}(\partial S),$$

where \bar{C} is a constant depending only on n , C and K .

Proof. For each $\sigma \in \mathcal{F}_{k+1}$ for which $P'_i \llcorner \sigma \neq 0$ we have

$$0 \neq \mathbf{M}(P'_i \llcorner \sigma) \leq K \mathbf{M}(S'_i \llcorner \text{st } \sigma) = K \mathbf{M}(g_{\#}(S_{r_i} \llcorner g^{-1}(\text{st } \sigma))).$$

In particular, $g^{-1}(\text{st } \sigma) \cap B(Z, \eta)$ is nonempty, and so $r_\sigma := d(Z, g^{-1}(\text{st } \sigma))$ satisfies $r_\sigma \leq \eta$. It furthermore follows from Theorem 4.1(4) that $r_\sigma > 0$. Since $g^{-1}(\text{st } \sigma) \subset \{\varrho \geq r_\sigma\}$ and g is $C r_\sigma^{-1}$ -Lipschitz on $\{\varrho \geq r_\sigma\}$ it follows with the above that

$$\mathbf{M}(P'_i \llcorner \sigma) \leq K C^{k+1} r_\sigma^{-(k+1)} \mathbf{M}(S_{r_i} \llcorner g^{-1}(\text{st } \sigma)).$$

Since $h|_\sigma$ is $C r_\sigma$ -Lipschitz, we conclude

$$\begin{aligned} \mathbf{M}(\Lambda_{k+1}(P'_i \llcorner \sigma)) &\leq K C^{k+1} r_\sigma^{k+1} \mathbf{M}(P'_i \llcorner \sigma) \\ &\leq K^2 C^{2(k+1)} \mathbf{M}(S_{r_i} \llcorner g^{-1}(\text{st } \sigma)). \end{aligned}$$

Summing over all $\sigma \in \mathcal{F}_{k+1}$ and observing that each open simplex in Σ is in the open star of at most $C_{n,k}$ different $(k + 1)$ -simplices we obtain

$$\mathbf{M}(P_i) \leq C_{n,k} K^2 C^{2(k+1)} \mathbf{M}(S_{r_i}),$$

and hence $\mathbf{M}(P_i) \leq \bar{C} \mathbf{M}(S)$ for all i large enough, where $\bar{C} := 2C_{n,k} K^2 C^{2(k+1)}$. The same argument as above shows that for each $\sigma \in \mathcal{F}_k$ we have

$$\mathbf{M}(\Lambda_k((\partial P'_i) \sqcup \sigma)) \leq K^2 C^{2k} \mathbf{M}((\partial S_{r_i}) \sqcup g^{-1}(\text{st } \sigma)).$$

From this one concludes as above that

$$\mathbf{M}(\partial P_i) \leq C_{n,k} K^2 C^{2k} \mathbf{M}(\partial S_{r_i}) \leq \bar{C} \mathbf{M}(\partial S).$$

Finally, for every σ for which $P'_i \sqcup \sigma \neq \emptyset$ we have $\text{Lip}(h|_\sigma) \leq Cr_\sigma \leq C\eta$ and therefore

$$\text{spt}(\Lambda_{k+1}(P'_i \sqcup \sigma)) \subset B(h(\sigma^{(0)}), KC\eta) \subset B(Z, \bar{C}\eta).$$

This shows that $\text{spt } P_i \subset B(Z, \bar{C}\eta)$ and completes the proof. \square

Lemma 5.6. *We have $\text{Fillvol}_Y(\partial S - \partial P_i) \rightarrow 0$ as $i \rightarrow \infty$. In particular, the sequence (∂P_i) converges weakly to ∂S .*

Proof. Since $\partial P'_i = \partial S'_i - \partial R'_i$, we find

$$h_\#(\partial P'_i) = \partial(h_\# S'_i) - \partial(h_\# R'_i) = \partial(f_\# S_{r_i}) - \partial(h_\# R'_i)$$

and thus, by Proposition 3.8,

$$\begin{aligned} \partial S - \partial P_i &= \partial S - \Lambda_k(\partial P'_i) = \partial S - h_\#(\partial P'_i) - \partial \Gamma_k(\partial P'_i) \\ &= \partial(f_\#(S - S_{r_i})) + \partial(h_\# R'_i) - \partial \Gamma_k(\partial P'_i) =: \partial A_i. \end{aligned}$$

It follows that

$$\text{Fillvol}_Y(\partial S - \partial P_i) \leq \mathbf{M}(A_i) \leq \mathbf{M}(f_\#(S - S_{r_i})) + \mathbf{M}(h_\# R'_i) + \mathbf{M}(\Gamma_k(\partial P'_i)).$$

We claim that there is a constant $\bar{C} > 0$ such that for every i each term on the right hand side in the inequality above is bounded by $\bar{C} r_i \mathbf{M}(\partial S)$. Clearly, we have

$$\mathbf{M}(f_\#(S - S_{r_i})) \leq C^{k+1} \mathbf{M}(S - S_{r_i}) = C^{k+1} \|S\|(\{\varrho \leq r_i\}) \leq C^{k+1} (k + 1) r_i \mathbf{M}(\partial S),$$

where we have used (5.1) in the last inequality. Next, set $M_i := \text{spt } R'_i$ and observe that

$$M_i \subset \text{Hull}(\text{spt } \partial S'_i) \subset \text{Hull}(g(\text{spt } \partial S_{r_i})) \subset \text{Hull}(g(\{\varrho = r_i\})). \tag{5.6}$$

Thanks to Theorem 4.1(3), h is Cr_i -Lipschitz on every $\sigma \in \mathcal{F}$ for which the set $\text{st } \sigma \cap g(\{\varrho = r_i\})$ is nonempty. It thus follows from (5.6) and Proposition 4.2 that $\text{lip}(h|_{M_i})(x) \leq Cr_i$ for all $x \in M_i$, and so, by Lemma 3.3, we get

$$\mathbf{M}(h_{\#}R'_i) \leq C^{k+1}r_i^{k+1} \mathbf{M}(R'_i) \leq KC^{k+1}r_i^{k+1} \mathbf{M}(\partial S'_i)$$

and hence $\mathbf{M}(h_{\#}R'_i) \leq KC^{2k+1}r_i \mathbf{M}(\partial S)$. Finally, one calculates exactly as in the proof of Lemma 5.5 that for each $\sigma \in \mathcal{F}_k$

$$\mathbf{M}(\Gamma_k((\partial P'_i) \llcorner \sigma)) \leq K^2C^{2k}r_i \mathbf{M}((\partial S_{r_i}) \llcorner g^{-1}(\text{st } \sigma))$$

and

$$\mathbf{M}(\Gamma_k(\partial P'_i)) \leq C_{n,k}K^2C^{2k}r_i \mathbf{M}(\partial S).$$

Thus, $\mathbf{M}(A_i) \rightarrow 0$ as i goes to infinity. This proves the claim and completes the proof. \square

The proof of all except the last statement of Theorem 5.1 can now easily be concluded if we can show that (P_i) has a weakly converging subsequence, which we will show in the following lemma. Indeed, if a subsequence of (P_i) converges weakly to \bar{S} , then \bar{S} belongs to $\mathbf{I}_{k+1}(X)$ and satisfies

$$\mathbf{M}(\bar{S}) \leq \bar{C} \mathbf{M}(S) \quad \text{and} \quad \text{spt } \bar{S} \subset B(Z, \bar{C}\eta),$$

as follows from Lemma 5.5, the closure theorem [3, Theorem 8.5], and the lower semi-continuity of mass. Moreover, we have $\partial \bar{S} = \partial S$ thanks to Lemma 5.6, and thus \bar{S} satisfies the properties stated in Theorem 5.1.

Lemma 5.7. *The sequence (P_i) has a weakly converging subsequence.*

Notice that when X is proper the lemma follows directly from Lemma 5.5 and the compactness theorem [3, Theorem 5.2].

Proof. Fix $r > 0$, let \mathcal{A}_r be the collection of all $\sigma \in \mathcal{F}_{k+1}$ satisfying $r_\sigma > r$, and denote by A_r the union of all simplices in \mathcal{A}_r . We first claim that the set $\{P'_i \llcorner A_r : i \geq 1\}$ contains only finitely many elements. To see this, let $\sigma \in \mathcal{A}_r$ and notice that $P'_i \llcorner \sigma = \theta_i^\sigma \llbracket \sigma \rrbracket$ for some $\theta_i^\sigma \in \mathbb{Z}$. By (A.2) we have

$$\mathcal{L}(\Delta) \cdot |\theta_i^\sigma| = \mathbf{M}(P'_i \llcorner \sigma) \leq K \mathbf{M}(S'_i \llcorner \text{st } \sigma) =: KM_i^\sigma, \tag{5.7}$$

where $\mathcal{L}(\Delta)$ denotes the Lebesgue measure of the Euclidean $(k + 1)$ -simplex Δ . By construction, there exists $i_0 \geq 1$ depending on r such that for all $\sigma \in \mathcal{A}_r$ and all $i, j \geq i_0$

we have $S'_i \llcorner \text{st } \sigma = S'_j \llcorner \text{st } \sigma$. In particular, $M_i^\sigma = M_{i_0}^\sigma =: M^\sigma$ for all $i \geq i_0$. Furthermore, since $M^\sigma \neq 0$ for at most countably many $\sigma \in \mathcal{A}_r$, we see that

$$\sum_{\sigma \in \mathcal{A}_r} M^\sigma = \sum_{\sigma \in \mathcal{A}_r} \mathbf{M}(S'_{i_0} \llcorner \text{st } \sigma) \leq C_{n,k} \sum_{\tau \in \mathcal{F}, r_\tau > r} \mathbf{M}(S'_{i_0} \llcorner \text{int } \tau) \leq C_{n,k} \mathbf{M}(S'_{i_0}) < \infty,$$

where we have used that each open simplex in Σ is in the open star of at most $C_{n,k}$ different $(k + 1)$ -simplices. Thus, the collection $\mathcal{C} := \{\sigma \in \mathcal{A}_r : KM^\sigma \geq \mathcal{L}(\Delta)\}$ is finite and letting $M := \sup_{\sigma' \in \mathcal{A}_r} M^{\sigma'} < \infty$, by virtue of (5.7) we find that for all $i \geq i_0$, $\theta_i^\sigma = 0$ if $\sigma \notin \mathcal{C}$ and $|\theta_i^\sigma| \leq KM$ for all $\sigma \in \mathcal{C}$. This implies our first claim.

Next, we claim that for almost every $r > 0$ small enough and every i sufficiently large we have

$$\sum_{\sigma \in \mathcal{F}_{k+1} \setminus \mathcal{A}_r} \mathbf{M}(\Lambda_{k+1}(P'_i \llcorner \sigma)) \leq \bar{C}r \mathbf{M}(\partial S),$$

where \bar{C} only depends on n, C , and K . Indeed, the proof of Lemma 5.5 shows that

$$\mathbf{M}(\Lambda_{k+1}(P'_i \llcorner \sigma)) \leq K^2 C^{2(k+1)} \mathbf{M}(S_{r_i} \llcorner g^{-1}(\text{st } \sigma)).$$

Summing over all $\sigma \in \mathcal{F}_{k+1} \setminus \mathcal{A}_r$ and noting that $g^{-1}(\text{st } \sigma) \subset \{\varrho \leq Cr\}$, because of Theorem 4.1(4), yields

$$\sum_{\sigma \in \mathcal{F}_{k+1} \setminus \mathcal{A}_r} \mathbf{M}(\Lambda_{k+1}(P'_i \llcorner \sigma)) \leq C_{n,k} K^2 C^{2(k+1)} \|S_{r_i}\|(\{\varrho \leq Cr\}),$$

and together with Lemma 5.4 we obtain the second claim.

Finally, it follows from the first claim and a standard diagonal sequence argument, that (P'_i) has a subsequence $(P'_{i(m)})$ with the following property. For every $r > 0$ we have

$$P'_{i(m_1)} - P'_{i(m_2)} = \sum_{\sigma \in \mathcal{F}_{k+1} \setminus \mathcal{A}_r} (P'_{i(m_1)} - P'_{i(m_2)}) \llcorner \sigma$$

for all m_1, m_2 sufficiently large (depending on r). This together with the second claim shows that for every $r > 0$ sufficiently small, the mass of $P'_{i(m_1)} - P'_{i(m_2)}$ is bounded by

$$\sum_{\sigma \in \mathcal{F}_{k+1} \setminus \mathcal{A}_r} \left[\mathbf{M}(\Lambda_{k+1}(P'_{i(m_1)} \llcorner \sigma)) + \mathbf{M}(\Lambda_{k+1}(P'_{i(m_2)} \llcorner \sigma)) \right] \leq 2\bar{C}r \mathbf{M}(\partial S)$$

whenever m_1, m_2 are sufficiently large. Consequently, $(P'_{i(m)})$ is a Cauchy sequence in $\mathbf{M}_{k+1}(X)$ and, in particular, is a weakly converging subsequence of (P_i) . \square

The following lemma proves the last statement in Theorem 5.1.

Lemma 5.8. *If Y is a Banach space then there exists $W \in \mathbf{I}_{k+2}(Y)$ with $\partial W = \bar{S} - S$ and such that*

$$\mathbf{M}(W) \leq \eta \bar{C} \mathbf{M}(S) \quad \text{and} \quad \text{spt } W \subset B(Z, \bar{C}\eta)$$

for some \bar{C} depending only on n, C and K .

Proof. Let $H: [0, 1] \times Y \rightarrow Y$ be the straight line homotopy defined by $H(t, y) = (1 - t)y + tf(y)$. Since $\text{spt } \partial S \subset Z$ and $H(t, z) = z$ for every $z \in Z$ it follows that $H_{\#}(\llbracket 0, 1 \rrbracket \times \partial S) = 0$ and thus, by Proposition 3.4, the integral current $V := H_{\#}(\llbracket 0, 1 \rrbracket \times S)$ satisfies $\partial V = f_{\#}S - S$. We have $d(y, f(y)) \leq (C + 1)\eta$ for every $y \in B(Z, \eta)$ and thus, by Lemma 3.5,

$$\mathbf{M}(V) \leq (k + 2)C^{k+1}(C + 1)\eta \mathbf{M}(S)$$

as well as

$$\text{spt } V \subset H(\llbracket 0, 1 \rrbracket \times \text{spt } S) \subset B(Z, (C + 1)C\eta).$$

Fix i sufficiently large, to be determined later. By (5.4) and Proposition 3.8 we have

$$P_i = f_{\#}S_{r_i} + \Gamma_k(\partial P'_i) - h_{\#}R'_i + \partial\Gamma_{k+1}(P'_i) - \partial h_{\#}Q'_i = f_{\#}S - A_i + \partial\Gamma_{k+1}(P'_i) - \partial h_{\#}Q'_i,$$

where A_i is as in the proof of Lemma 5.6. Hence,

$$\bar{S} - S = (\bar{S} - P_i) - A_i + \partial\Gamma_{k+1}(P'_i) - \partial h_{\#}Q'_i + \partial V. \tag{5.8}$$

Using the same arguments as in the proof of Lemma 5.5 one easily shows that $\Gamma_{k+1}(P'_i)$ and $h_{\#}Q'_i$ are supported in $B(Z, \bar{C}\eta)$ and satisfy $\mathbf{M}(h_{\#}Q'_i) \leq \eta \bar{C} \mathbf{M}(S)$ as well as $\mathbf{M}(\Gamma_{k+1}(P'_i)) \leq \bar{C}\eta \mathbf{M}(S)$ for some \bar{C} depending on n, C and K .

It follows from the proof of Lemma 5.7 that, after passing to a subsequence, P_i converges to \bar{S} in mass, so we can take $\mathbf{M}(\bar{S} - P_i)$ arbitrarily small. By the proof of Lemma 5.6, $\mathbf{M}(A_i)$ converges to zero as i tends to infinity. Moreover, (5.8) implies that $\bar{S} - P_i - A_i$ is a cycle whose support is contained in $B(Z, \bar{C}\eta)$. Hence, by Lemma 3.7 and the remark following it, we can choose an arbitrarily large i such that $\bar{S} - P_i - A_i$ has a filling $U \in \mathbf{I}_{k+2}(Y)$ satisfying $\mathbf{M}(U) \leq \eta \bar{C} \mathbf{M}(S)$ and $\text{spt } U \subset B(Z, 2\bar{C}\eta)$. Consequently, the integral current $W := U + \Gamma_{k+1}(P'_i) - h_{\#}Q'_i + V$ satisfies $\partial W = \bar{S} - S$ as well as $\mathbf{M}(W) \leq \eta \bar{C} \mathbf{M}(S)$ and $\text{spt } W \subset B(Z, \bar{C}\eta)$ for some \bar{C} depending on n, C and K . \square

6. Approximating finite-dimensional spaces by simplicial complexes

The aim of this section is to prove Theorem 1.6 from the introduction. To this end, we first establish the following approximation result:

Proposition 6.1. *Let $X \subset Y$ be metric spaces such that X has Nagata dimension $\leq n$ and Y is Lipschitz $(n - 1)$ -connected for some $n \geq 0$. Then there is a constant C such that for every $\varepsilon > 0$ there exist a simplicial complex Σ equipped with the ℓ_2 -metric and maps $\psi: X \rightarrow \Sigma$ and $\varphi: \Sigma \rightarrow Y$ with the following properties:*

- (1) $\varphi(\Sigma^{(0)}) \subset X$, $\text{Hull}(\psi(X)) = \Sigma$ and Σ has dimension $\leq n$;
- (2) ψ is $C\varepsilon^{-1}$ -Lipschitz and φ is $C\varepsilon$ -Lipschitz on every $\sigma \in \mathcal{F}$;
- (3) $\varphi \circ \psi$ is C -Lipschitz and $d(x, \varphi(\psi(x))) \leq C\varepsilon$ for all $x \in X$.

The constant C depends only on the data of X and Y .

Notice that in the case $n = 0$, we do not impose any assumption on the Lipschitz connectedness of Y . If $X \subset Y$ is a closed subset such that X is Lipschitz k -connected for some $k \leq n$, then we can choose φ such that, in addition, $\varphi(\Sigma^{(k+1)}) \subset X$. The proof of Proposition 6.1 is similar to the proof of the factorization theorem in Section 4.

Proof of Proposition 6.1. Let c denote the Nagata constant of X . Fix $\varepsilon > 0$ and let $\mathcal{B} = (B_i)_{i \in I}$ be a $c\varepsilon$ -bounded covering of X with ε -multiplicity $n + 1$. We begin by constructing a map $\psi: X \rightarrow \Sigma(I)$. For each $i \in I$ we define $\tau_i: X \rightarrow \mathbb{R}$ via $x \mapsto \max\{\frac{\varepsilon}{2} - d(x, B_i), 0\}$. If $\tau_i(x) > 0$, then there is $b_i \in B_i$ with $d(x, b_i) \leq \frac{\varepsilon}{2}$; consequently, for any $x \in X$ there are at most $(n + 1)$ indices $i \in I$ such that $\tau_i(x) > 0$, as \mathcal{B} has ε -multiplicity $n + 1$. We set $\bar{\tau}(x) = \sum_{i \in I} \tau_i(x)$. Clearly, $\bar{\tau}(x) > 0$ for every $x \in X$. By the above, $\psi: X \rightarrow \Sigma(I)$ given by $x \mapsto \bar{\tau}(x)^{-1}(\tau_i(x))_{i \in I}$ is well-defined and $\Sigma := \text{Hull}(\psi(X))$ is an n -dimensional simplicial complex. In what follows, we equip Σ with the ℓ_2 -metric and view ψ as a map from X to Σ . One calculates exactly as in the proof of Theorem 4.1 that for all $x, x' \in X$,

$$|\psi(x) - \psi(x')| \leq \frac{4(n + 1)}{\bar{\tau}(x)} d(x, x').$$

Since \mathcal{B} is a covering of X , it follows that $\bar{\tau}(x) \geq \frac{\varepsilon}{2}$ for all $x \in X$, so by the estimate above, ψ is $C_0\varepsilon^{-1}$ -Lipschitz, as desired.

Next, we construct $\varphi: \Sigma \rightarrow Y$. As Y is Lipschitz $(n - 1)$ -connected, the same argument as in the proof of Theorem 4.1 yields a map $\varphi: \Sigma \rightarrow Y$ such that $h(e_i) \in B_i$ and $\text{Lip}(\varphi|_\sigma) \leq C_1 \text{Lip}(\varphi|_{\sigma^{(0)}})$ for every $\sigma \in \mathcal{F}(\Sigma)$. Here, C_1 denotes a constant depending only on the data of X and Y . Fix $\sigma \in \mathcal{F}$ and let e_i, e_j be two vertices of σ . Clearly, there is $x \in \psi^{-1}(\text{st } \sigma)$ such that $\tau_i(x) > 0$ and $\tau_j(x) > 0$. We obtain

$$d(\varphi(e_i), \varphi(e_j)) \leq d(\varphi(e_i), x) + d(x, \varphi(e_j)) \leq 2\left(c\varepsilon + \frac{\varepsilon}{2}\right),$$

since \mathcal{B} is $c\varepsilon$ -bounded. Hence, by the above, φ is $C_2\varepsilon$ -Lipschitz on σ , where $C_2 := \sqrt{2}(c + 1)C_1$. We proceed by showing (3). Let $\delta := \varphi \circ \psi$. Fix $x \in X$ and let $\sigma \in \mathcal{F}$ be the unique simplex such that $\psi(x) \in \text{int } \sigma$. Since there exists some $i \in I$ with $\tau_i(x) > 0$, we find that

$$d(\delta(x), x) \leq d(\delta(x), \varphi(e_i)) + d(\varphi(e_i), x) \leq C_3\varepsilon, \tag{6.1}$$

as φ is $C_2\varepsilon$ -Lipschitz on σ and $d(\varphi(e_i), x) \leq c\varepsilon + \frac{\varepsilon}{2}$. In particular, if $d(x, x') \geq \frac{\varepsilon}{8}$, then (6.1) yields $d(\delta(x), \delta(x')) \leq C_4d(x, x')$, where $C_4 := 1 + 16C_3$. Now, suppose that $x, x' \in X$ satisfy $d(x, x') \leq \frac{\varepsilon}{8}$, and let $\sigma, \sigma' \in \mathcal{F}$ denote the unique simplices such that $\psi(x) \in \text{int } \sigma$ and $\psi(x') \in \text{int } \sigma'$, respectively. If $x \in B_i$ for some $i \in I$, then $\tau_i(x') > 0$; thus, $\sigma \cap \sigma' \neq \emptyset$. There is $p \in \sigma \cap \sigma'$ such that $|\psi(x) - p| + |p - \psi(x')| \leq 2n|\psi(x) - \psi(x')|$, and we can estimate

$$d(\delta(x), \delta(x')) \leq d(\delta(x), \varphi(p)) + d(\varphi(p), \delta(x')) \leq C_2\varepsilon \left(|\psi(x) - p| + |p - \psi(x')| \right).$$

Hence, $d(\delta(x), \delta(x')) \leq 2C_0C_2n d(x, x')$, for ψ is $C_0\varepsilon^{-1}$ -Lipschitz. By setting, $C_5 := 2C_0C_2C_4n$, we conclude that $\delta = \varphi \circ \psi$ is C_5 -Lipschitz, as desired. We put $C := \max\{C_i : i = 0, \dots, 5\}$, which does not depend on ε . This finishes the proof of Proposition 6.1. \square

We conclude this section with the proof of Theorem 1.6.

Proof of Theorem 1.6. By Proposition 6.1 there is $C > 0$ such that for every $\varepsilon > 0$ there exist a simplicial complex Σ of dimension $\leq n$ equipped with the ℓ_2 -metric and maps $\psi: X \rightarrow \Sigma$ and $\varphi: \Sigma \rightarrow Y$ such that statements (1) – (3) of Proposition 6.1 hold. Fix $\varepsilon > 0$. Notice that $\psi(X)$ is rectifiably connected and every $z \in \Sigma$ lies in a simplex of Σ which intersects $\psi(X)$, so Σ is also rectifiably connected. Let d_ε be the length metric on Σ , scaled by the factor $\varepsilon \cdot 2^{-1/2}$. Then d_ε is a length metric and (Σ, d_ε) satisfies (1). In the following, we show that $\psi: X \rightarrow (\Sigma, d_\varepsilon)$ and $\varphi: (\Sigma, d_\varepsilon) \rightarrow Y$ have the desired properties.

Let $x, x' \in X$ and let α be a curve in X from x to x' satisfying $\ell(\alpha) \leq cd(x, x')$, where c is the quasicvexity constant of X . Then

$$\sqrt{2}d_\varepsilon(\psi(x), \psi(x')) \leq \varepsilon \ell(\psi \circ \alpha) \leq C\ell(\alpha) \leq Cc d(x, x').$$

This shows that $\psi: X \rightarrow (\Sigma, d_\varepsilon)$ is $2^{-1/2} \cdot Cc$ -Lipschitz. Next, let $z, z' \in \Sigma$. It follows from Proposition 4.2 and Lemma 2.4 that for every curve γ in Σ connecting z and z' we have $d(\varphi(z), \varphi(z')) \leq C\varepsilon\ell(\gamma)$. By taking the infimum over all such curves γ , we conclude

$$d(\varphi(z), \varphi(z')) \leq C\sqrt{2}d_\varepsilon(z, z').$$

In particular, the map $\varphi: (\Sigma, d_\varepsilon) \rightarrow Y$ is $C\sqrt{2}$ -Lipschitz and it is clear that φ and ψ satisfy condition (2) and the remaining statements in the theorem. \square

7. Deformation theorem in spaces of finite Nagata dimension

The aim of this section is to prove Theorem 1.4. The proof will rely on Theorem 1.6 from the introduction and the deformation theorem for bilipschitz triangulated metric spaces (see Theorem A.2). We follow the proof strategy outlined in Section 1.4.

Let X be as in Theorem 1.4. We define $Y := \ell_\infty(X)$ and view X as a subset of Y . By Theorem 1.6 there is a constant $C > 0$ depending only on the data of X such that for every $\varepsilon > 0$ there exist a metric simplicial complex Σ and C -Lipschitz maps $\psi: X \rightarrow \Sigma$ and $\varphi: \Sigma \rightarrow Y$ such that (1) and (2) of Theorem 1.6 hold. Fix $\varepsilon > 0$. Let Λ_m and Γ_m be the homomorphisms from Proposition 3.8 when applied to φ . These homomorphisms, satisfy

$$\mathbf{M}(\Lambda_m(\llbracket\sigma\rrbracket)) \leq C\varepsilon^m, \quad \mathbf{M}(\Gamma_m(\llbracket\sigma\rrbracket)) \leq C\varepsilon^{m+1}$$

and $\text{spt } \Lambda_m(\llbracket\sigma\rrbracket), \text{spt } \Gamma_m(\llbracket\sigma\rrbracket) \subset B(\varphi(\sigma^{(0)}), C\varepsilon)$ for every $m = 0, \dots, k + 1$ and each m -simplex σ in Σ . The proof of Theorem 1.4 relies on the following two auxiliary lemmas.

Lemma 7.1. *For every subset $A \subset X$ we have $\varphi(\text{Hull}(\psi(A))) \subset B(A, C\varepsilon)$. Moreover, if $Q \in \mathcal{P}_m(\Sigma)$ has support in $\text{Hull}(\psi(A))$, then*

$$\text{spt}(\Lambda_m(Q)), \text{spt}(\Gamma_m(Q)) \subset B(A, C\varepsilon),$$

where C is a constant only depending on the data of X .

Proof. Let $A \subset X$ be a nonempty set and let $z \in \text{Hull}(\psi(A))$. Then there exists $x \in A$ such that $\psi(x)$ and z lie in a common simplex in $\text{Hull}(\psi(A))$. In particular, we have $d(z, \psi(x)) \leq \varepsilon$ and hence

$$d(\varphi(z), x) \leq d(\varphi(z), \varphi(\psi(x))) + d(\varphi(\psi(x)), x) \leq 2C\varepsilon.$$

This implies that $\varphi(\text{Hull}(\psi(A))) \subset B(A, 2C\varepsilon)$, as claimed. In order to prove the second statement, let $\sigma \subset \text{Hull}(\psi(A))$ be an m -simplex. Since $\varphi(\sigma^{(0)}) \subset B(A, 2C\varepsilon)$ it follows from Proposition 3.8 that

$$\text{spt}(\Lambda_m(\llbracket\sigma\rrbracket)), \text{spt}(\Gamma_m(\llbracket\sigma\rrbracket)) \subset B(A, 3C\varepsilon).$$

This implies the second statement. \square

Lemma 7.2. *The triple $(\Sigma, \varphi|_{\Sigma^{(0)}}, \Lambda_*)$, where $\Lambda_* := \{\Lambda_0, \Lambda_1, \dots, \Lambda_k\}$, defines a (k, ε) -polyhedral structure with a constant only depending on the data of X .*

Proof. We only need to show that $\varphi|_{\Sigma^{(0)}}$ has the desired properties. Let $z, w \in \Sigma^{(0)}$. Then there exist $x, y \in X$ such that z and $\psi(x)$ lie in a common simplex and w and $\psi(y)$ lie in a common simplex. In particular, we have

$$\begin{aligned} d(\varphi(z), \varphi(w)) &\geq d(\varphi(\psi(x)), \varphi(\psi(y))) - d(\varphi(z), \varphi(\psi(x))) - d(\varphi(w), \varphi(\psi(y))) \\ &\geq d(x, y) - 4C\varepsilon. \end{aligned}$$

Moreover, for every $x \in X$ we have

$$d(x, \varphi(\Sigma^{(0)})) \leq C\varepsilon + d(\varphi(\psi(x)), \varphi(\Sigma^{(0)})) \leq C\varepsilon + Cd(\psi(x), \Sigma^{(0)}) \leq 2C\varepsilon.$$

This completes the proof. \square

Now we are in a position to prove Theorem 1.4.

Proof of Theorem 1.4. In what follows, C and C' will denote constants depending only on k and the data of X and they may change from one appearance to another. Let $1 \leq m \leq k$ and let $T \in \mathbf{I}_m(X)$. Define $T' := \psi_{\#}T$ and write $T' = P' + R' + \partial S'$, where P', R', S' are as in Theorem A.2. If m is strictly larger than the dimension of Σ then $T' = 0$ by the comment after Theorem A.2 and in this case we let P', R' , and S' be the zero currents. Define $P := \Lambda_m(P')$. By Proposition 3.8 and Theorem A.2 we have that

$$\mathbf{M}(P) \leq C \mathbf{M}(T), \quad \mathbf{M}(\partial P) \leq C \mathbf{M}(\partial T).$$

Moreover, $\text{spt } P' \subset \text{Hull}(\psi(\text{spt } T))$ and $\text{spt } \partial P' \subset \text{Hull}(\psi(\text{spt } \partial T))$ and thus Lemma 7.1 implies that

$$\text{spt } P \subset B(\text{spt } T, C\varepsilon), \quad \text{spt } \partial P \subset B(\text{spt } \partial T, C\varepsilon).$$

Now, let $H: [0, 1] \times X \rightarrow Y$ be the straight line homotopy

$$H(t, x) := (1 - t)x + t\varphi(\psi(x)).$$

The integral currents in Y defined by $U := H_{\#}(\llbracket 0, 1 \rrbracket \times \partial T)$ and $V := H_{\#}(\llbracket 0, 1 \rrbracket \times T)$ satisfy

$$T - \varphi_{\#}T' = -U - \partial V$$

and, by Lemma 3.5, $\mathbf{M}(U) \leq \varepsilon C \mathbf{M}(\partial T)$ and $\mathbf{M}(V) \leq \varepsilon C \mathbf{M}(T)$. Moreover, we have

$$\text{spt } U \subset H([0, 1] \times \text{spt } \partial T) \subset B(\text{spt } \partial T, C\varepsilon)$$

and similarly $\text{spt } V \subset H([0, 1] \times \text{spt } T) \subset B(\text{spt } T, C\varepsilon)$. Proposition 3.8 yields

$$\varphi_{\#}P' = P - \partial\Gamma_m(P') - \Gamma_{m-1}(\partial P')$$

and hence

$$T = T - \varphi_{\#}T' + \varphi_{\#}T' = -U - \partial V + \varphi_{\#}P' + \varphi_{\#}R' + \partial\varphi_{\#}S' = P + \hat{R} + \partial\hat{S},$$

where we have set $\hat{R} := \varphi_{\#}R' - U - \Gamma_{m-1}(\partial P')$ and $\hat{S} := \varphi_{\#}S' - V - \Gamma_m(P')$. A direct calculation shows that

$$\mathbf{M}(\hat{R}) \leq \varepsilon C \mathbf{M}(\partial T), \quad \mathbf{M}(\hat{S}) \leq \varepsilon C \mathbf{M}(T).$$

Moreover, Lemma 7.1 implies that $\text{spt } \hat{R} \subset B(\text{spt } \partial T, C\varepsilon)$ and $\text{spt } \hat{S} \subset B(\text{spt } T, C\varepsilon)$.

Since $\partial \hat{R} = \partial(T - P)$ is supported in X , Theorem 5.3 implies the following two statements. There exists $R \in \mathbf{I}_m(X)$ with $\partial R = \partial \hat{R}$ and such that $\mathbf{M}(R) \leq C \mathbf{M}(\hat{R}) \leq \varepsilon C' \mathbf{M}(\partial T)$ as well as $\text{spt } R \subset B(\text{spt } \partial T, C\varepsilon)$. Moreover, there is $\bar{S} \in \mathbf{I}_{m+1}(Y)$ satisfying $\partial \bar{S} = \hat{R} - R$ and

$$\mathbf{M}(\bar{S}) \leq \varepsilon C \mathbf{M}(\hat{R}) \leq \varepsilon^2 C' \mathbf{M}(\partial T)$$

as well as $\text{spt } \bar{S} \subset B(\text{spt } \partial T, C'\varepsilon)$. By construction, $T = P + R + \partial(\bar{S} + \hat{S})$ and hence $\partial(\bar{S} + \hat{S})$ is supported in X and satisfies

$$\text{spt}(\bar{S} + \hat{S}) \subset B(\text{spt } T, C\varepsilon).$$

Thus, by Theorem 5.3, there exists $S \in \mathbf{I}_{m+1}(X)$ with $\partial S = \partial(\bar{S} + \hat{S})$ and

$$\mathbf{M}(S) \leq C \mathbf{M}(\bar{S} + \hat{S}) \leq \varepsilon C' \mathbf{M}(T) + \varepsilon^2 C' \mathbf{M}(\partial T)$$

as well as

$$\text{spt } S \subset B(\text{spt } T, C'\varepsilon).$$

Since $T = P + \hat{R} + \partial \hat{S} = P + R + \partial \bar{S} + \partial \hat{S} = P + R + \partial S$ we have a decomposition as in the statement of the theorem. \square

Theorem 1.4 implies the following approximation result mentioned in the introduction.

Corollary 7.3. *Suppose X is a complete metric space of finite Nagata dimension which has (LC_k) for some $k \geq 1$, and let $T \in \mathbf{I}_k(X)$. Then there exists a sequence of Lipschitz k -chains in X such that the induced integral currents T_i satisfy $\mathcal{F}_X(T - T_i) \rightarrow 0$ as $i \rightarrow \infty$ and*

$$\sup_i \left[\mathbf{M}(T_i) + \mathbf{M}(\partial T_i) \right] < \infty.$$

If T is a cycle, then the T_i are cycles and $\text{Fillvol}_X(T - T_i) \rightarrow 0$ as $i \rightarrow \infty$.

In the above, $\mathcal{F}_X(T)$ is the flat norm in X of a current $T \in \mathbf{I}_k(X)$ and is defined by

$$\mathcal{F}_X(T) := \inf \{ \mathbf{M}(U) + \mathbf{M}(V) : U \in \mathbf{I}_k(X), V \in \mathbf{I}_{k+1}(X), T = U + \partial V \}.$$

In any complete metric space, convergence with respect to the flat norm implies weak convergence. Moreover, in metric spaces with local coning inequalities the two notions of convergence are equivalent for sequences with uniformly bounded mass and boundary mass (see [43] for more information).

Proof of Corollary 7.3. Since X has (LC_k) the map φ constructed above satisfies $\varphi(\Sigma^{(k+1)}) \subset X$; see Theorem 1.6. Moreover, the homomorphisms Λ_m in the polyhedral structure $(\Sigma, \varphi|_{\Sigma^{(0)}}, \Lambda_*)$ can be chosen such that $\Lambda(\llbracket \sigma \rrbracket) = \varphi_{\#} \llbracket \sigma \rrbracket$ for every m -simplex σ . See the beginning of the proof of Proposition 3.8. It thus follows that $\mathcal{P}_k(X) = \{\Lambda_k(Q) : Q \in \mathcal{P}_k(\Sigma)\}$ consists of integral currents induced by Lipschitz chains in X . Now, the corollary follows from Theorem 1.4. \square

We finally turn to the proof of Corollary 1.5. Given $k \geq 0$ we denote the space of Lipschitz k -chains in X by $C_k^{\text{Lip}}(X)$. Precisely, $C_k^{\text{Lip}}(X)$ is the free abelian group generated by all Lipschitz maps $\varphi: \Delta^k \rightarrow X$, where Δ^k is the Euclidean standard k -simplex. We view $C_k^{\text{Lip}}(X)$ as a subcomplex of the singular chain complex of X and equip it with the singular boundary operator ∂ (see, for instance, [22, Ch. 2.1]).

As already mentioned in Section 3, Lipschitz chains induce integral currents. Indeed, if $\alpha = \sum_{i=1}^N \theta_i \varphi_i$ is a Lipschitz k -chain in X then

$$\llbracket \alpha \rrbracket := \sum_{i=1}^N \theta_i \varphi_{i\#} \llbracket \Delta^k \rrbracket$$

is a k -dimensional integral current in X . Moreover, one has that $\partial \llbracket \alpha \rrbracket = \llbracket \partial \alpha \rrbracket$ for all $\alpha \in C_k^{\text{Lip}}(X)$. This can readily be verified by combining Stokes' theorem for chains with [3, Theorem 11.1]. The following lemma proves Corollary 1.5 in the special case when ∂T is induced by a Lipschitz cycle.

Lemma 7.4. *Let X be a complete metric space of finite Nagata dimension and let $T \in \mathbf{I}_k(X)$ for some $k \geq 1$. Suppose that there is a cycle $\delta \in C_{k-1}^{\text{Lip}}(X)$ such that $\partial T = \llbracket \delta \rrbracket$. If X has (LC_{k-1}) then for every $\varepsilon > 0$ there exists $\varrho \in C_k^{\text{Lip}}(X)$ such that*

$$\partial \varrho = \delta \quad \text{and} \quad \mathbf{M}(T - \llbracket \varrho \rrbracket) \leq \varepsilon.$$

We proceed with the proof of Corollary 1.5.

Proof of Corollary 1.5. Let $\varepsilon > 0$ and $k \geq 2$. By Lemma 7.4, there exists $\varrho' \in C_{k-1}^{\text{Lip}}(X)$ with $\partial \varrho' = 0$ and $\mathbf{M}(\partial T - \llbracket \varrho' \rrbracket) \leq \varepsilon$. Using Corollary 1.2 we find that X has (EI_{k-1}) , and thus there is $S' \in \mathbf{I}_k(X)$ satisfying $\partial S' = \partial T - \llbracket \varrho' \rrbracket$ and $\mathbf{M}(S') \leq D \varepsilon^{k/(k-1)}$. Let $T' = T - S'$. Then $\partial T' = \llbracket \varrho' \rrbracket$, and so by Lemma 7.4 there exists $\varrho \in C_k^{\text{Lip}}(X)$ with $\partial \varrho = \varrho'$ and $\mathbf{M}(T' - \llbracket \varrho \rrbracket) \leq \varepsilon$. Therefore,

$$\begin{aligned} \mathbf{M}(T - \llbracket \varrho \rrbracket) &\leq \varepsilon + \mathbf{M}(S') \leq 2D\varepsilon \\ \mathbf{M}(\partial T - \partial \llbracket \varrho \rrbracket) &= \mathbf{M}(\partial T - \llbracket \varrho' \rrbracket) \leq \varepsilon. \end{aligned}$$

This completes the proof in the case when $k \geq 2$. If $k = 1$, then $\partial T \in \mathbf{I}_0(X)$ and thus there are finitely many $x_i \in X$ and $\theta_i \in \mathbb{Z}$ such that $\partial T = \sum \theta_i \llbracket x_i \rrbracket$. In particular,

$\partial T = \llbracket \delta \rrbracket$ for some cycle $\delta \in C_0^{\text{Lip}}(X)$. Hence, by Lemma 7.4, for every $\varepsilon > 0$ there exists $\varrho \in C_1^{\text{Lip}}(X)$ with $\partial \varrho = \delta$ and $\mathbf{M}(T - \llbracket \varrho \rrbracket) \leq \varepsilon$, as desired. \square

Thus, we are left to prove Lemma 7.4. The main components in its proof are Theorem 1.6 and [50, Lemma 3]. Moreover, we will need the following straightforward approximation result.

Lemma 7.5. *Let X be a complete metric space and let $T \in \mathbf{I}_k(X)$ for some $k \geq 1$. If X has (LC_{k-1}) then for every $\varepsilon > 0$ there exists $\alpha \in C_k^{\text{Lip}}(X)$ such that $\mathbf{M}(T - \llbracket \alpha \rrbracket) \leq \varepsilon$.*

Proof. Let $K \subset \mathbb{R}^k$ be a compact subset and $\varphi: K \rightarrow X$ a Lipschitz map. Due to [3, Theorem 4.5] and the inner regularity of the Lebesgue measure on \mathbb{R}^k , to prove the lemma it suffices to show the following statement: for every $\varepsilon > 0$ there exists $\alpha \in C_k^{\text{Lip}}(X)$ such that $\mathbf{M}(\varphi_{\#} \llbracket K \rrbracket - \llbracket \alpha \rrbracket) \leq \varepsilon$. Fix $\varepsilon > 0$. By a straightforward construction, we find finitely many k -cubes $Q_i \subset \mathbb{R}^k$ such that $K \subset \bigcup Q_i$, $\text{int } Q_i \cap \text{int } Q_j = \emptyset$ whenever $i \neq j$, and $\mathcal{H}^k(\bigcup Q_i) \leq \mathcal{H}^k(K) + \varepsilon$. Clearly, there exists $\gamma \in C_k^{\text{Lip}}(\mathbb{R}^k)$ with $\llbracket \gamma \rrbracket = \sum \llbracket Q_i \rrbracket$. Now, since X has (LC_{k-1}) there is a C -Lipschitz map $\hat{\varphi}: \mathbb{R}^k \rightarrow X$ extending φ , and so by letting $\alpha := \hat{\varphi}_{\#} \gamma$, we obtain

$$\mathbf{M}(\varphi_{\#} \llbracket K \rrbracket - \llbracket \alpha \rrbracket) \leq C^k \mathbf{M}(\llbracket K \rrbracket - \sum \llbracket Q_i \rrbracket) \leq C^k \varepsilon,$$

where in the first inequality we used that $\varphi_{\#} \llbracket K \rrbracket = \hat{\varphi}_{\#} \llbracket K \rrbracket$. \square

The proof of Lemma 7.4 will use the following construction. Let Σ be a simplicial complex. By choosing a total ordering on the vertices of Σ , we can construct an isomorphism from $\mathcal{P}_*(\Sigma)$ to a subcomplex of $C_*^{\text{Lip}}(\Sigma)$. Namely, for every k and every k -simplex σ of Σ , there is a unique isometry $\varphi_{\sigma}: \Delta^k \rightarrow \Sigma$ that preserves the ordering of the vertices. The ordering also lets us fix an orientation for σ so that $\llbracket \sigma \rrbracket = (\varphi_{\sigma})_{\#} \llbracket \Delta^k \rrbracket$. Let $q: \mathcal{P}_k(\Sigma) \rightarrow C_k^{\text{Lip}}(\Sigma)$ be defined by

$$q\left(\sum_{\sigma} \theta_{\sigma} \llbracket \sigma \rrbracket\right) = \sum_{\sigma} \theta_{\sigma} \varphi_{\sigma}. \tag{7.1}$$

Then q is a homomorphism of chain complexes and $\llbracket q(P) \rrbracket = P$ for any $P \in \mathcal{P}_*(\Sigma)$.

Proof of Lemma 7.4. By Theorem 1.6 there exists $C > 0$ such that for every $\varepsilon > 0$ there are a metric simplicial complex $\Sigma = \Sigma(\varepsilon)$ and C -Lipschitz maps $\psi: X \rightarrow \Sigma$ and $\varphi: \Sigma^{(k)} \rightarrow X$ satisfying the properties listed in Theorem 1.6. Fix $\varepsilon_0 > 0$. By Lemma 7.5 there exists $\tilde{\alpha} \in C_k^{\text{Lip}}(X)$ with $\mathbf{M}(T - \llbracket \tilde{\alpha} \rrbracket) \leq \varepsilon_0$. Let $\alpha := \partial \tilde{\alpha} - \delta$. Then $\partial \alpha = 0$, and thus by [50, Lemma 3] there is a $c_{\alpha} > 0$ such that for every $\varepsilon > 0$ there are $\gamma \in C_k^{\text{Lip}}(\Sigma)$ and $\lambda \in C_k^{\text{Lip}}(X)$, and a $(k - 1)$ -cycle $\alpha' \in C_{k-1}^{\text{Lip}}(\Sigma)$, such that $\llbracket \alpha' \rrbracket \in \mathcal{P}_{k-1}(\Sigma)$ and

$$\partial \gamma = \psi_{\#}(\alpha) - \alpha', \quad \mathbf{M}(\llbracket \gamma \rrbracket) \leq c_{\alpha} \varepsilon,$$

$$\partial\lambda = \alpha - \varphi_{\#}(\alpha'), \quad \mathbf{M}(\llbracket\lambda\rrbracket) \leq c_{\alpha}\varepsilon.$$

After fixing a total ordering on the vertices of Σ , we may suppose that $\alpha' = q(\llbracket\alpha'\rrbracket)$, where q is as in (7.1).

Fix $\varepsilon > 0$ small enough such that $c_{\alpha}\varepsilon \leq \varepsilon_0$. The boundary of $T' := \psi_{\#}(\llbracket\tilde{\alpha}\rrbracket - T) - \llbracket\gamma\rrbracket$ is equal to $\llbracket\alpha'\rrbracket$ and one has $\mathbf{M}(T') \leq C^k\varepsilon_0 + \varepsilon_0$. Now, on account of Theorem A.2, there exist $P' \in \mathcal{P}_k(\Sigma)$ and $S' \in \mathbf{I}_{k+1}(\Sigma)$ with $T' = P' + \partial S'$ and such that $\mathbf{M}(P') \leq K\mathbf{M}(T')$, where K is a constant depending only on the Nagata dimension of X . Let $\beta := q(P') \in C_k^{\text{Lip}}(\Sigma)$ and $\varrho := \tilde{\alpha} - \lambda - \varphi_{\#}(\beta)$. Then $\partial\beta = q(\partial P') = q(\llbracket\alpha'\rrbracket) = \alpha'$, and so $\partial\varrho = \delta$ and

$$\mathbf{M}(T - \llbracket\varrho\rrbracket) \leq \varepsilon_0 + c_{\alpha}\varepsilon + C^k\mathbf{M}(P') \leq 2\varepsilon_0 + (C^kK) \cdot \mathbf{M}(T') \leq \bar{C}\varepsilon_0,$$

where \bar{C} is a constant which is independent of ε_0 . \square

Appendix A. Deformation theorem in bilipschitz triangulated metric spaces

The aim of this appendix is to provide a detailed proof of the classical Federer-Fleming deformation theorem in the setting of metric spaces admitting a bilipschitz triangulation in the sense below. In fact, we establish a version of the deformation theorem which moreover includes local mass estimates. Such estimates are of crucial importance in the proofs of Theorems 1.1 and 1.4.

Definition A.1. A metric space X is said to admit an (n, D, ε) -triangulation, where $D \geq 1$, $n \in \mathbb{N}$, $\varepsilon > 0$, if there exist an n -dimensional simplicial complex Σ and a homeomorphism $p: \Sigma \rightarrow X$ which is D -bilipschitz on every $\sigma \in \mathcal{F}$ if Σ is equipped with the scaled ℓ_2 -metric such that each simplex has side length ε .

Notice that in a path-connected simplicial complex the ℓ_2 -metric and the length metric induce the same topology and they agree on each simplex. Thus, for path connected spaces X the definition above could equivalently be stated with the scaled ℓ_2 -metric replaced by the scaled length metric on Σ . If X is quasiconvex then the homeomorphism $p: \Sigma \rightarrow X$ in the definition above is a bilipschitz homeomorphism when Σ is equipped with the scaled length metric; see Corollary 4.3.

We will use the following notation for simplices in (n, D, ε) -triangulated spaces. A subset $\sigma \subset X$ of a metric space X admitting a triangulation $p: \Sigma \rightarrow X$ in the sense above is called k -simplex in X if $p^{-1}(\sigma) \subset \Sigma$ is a k -simplex. We denote by \mathcal{F} the family of simplices in X and by \mathcal{F}_k the collection of k -simplices in X . For every $\sigma \in \mathcal{F}$ we write $\text{int } \sigma := p(\text{int } p^{-1}(\sigma))$ to denote the relative interior of σ in X . Given $\sigma \in \mathcal{F}$, the set

$$\text{st } \sigma := \bigcup \left\{ \text{int } \tau : \tau \in \mathcal{F} \text{ and } \sigma \subset \tau \right\}$$

is called the *open star* of σ . Further, the *hull* of a subset $B \subset X$ is the union of all simplices $\sigma \in \mathcal{F}$ such that $\text{int } \sigma \cap B \neq \emptyset$ and is denoted $\text{Hull}(B)$. For $k \geq 0$ we denote by $\mathcal{P}_k(X)$ the abelian subgroup of $\mathbf{I}_k(X)$ generated by currents of the form $p_{\#}[\![\sigma]\!]$, where $\sigma \subset \Sigma$ is a k -simplex.

Theorem A.2. *Let X be a c -quasiconvex (n, D, ε) -triangulated metric space. Then there is $C = C(c, n, D) > 0$ such that for every $k = 1, \dots, n$ and every $T \in \mathbf{I}_k(X)$ there exist $P \in \mathcal{P}_k(X)$, $R \in \mathbf{I}_k(X)$, and $S \in \mathbf{I}_{k+1}(X)$ with $T = P + R + \partial S$ and*

$$\begin{aligned} \mathbf{M}(P) &\leq C \mathbf{M}(T), & \mathbf{M}(\partial P) &\leq C \mathbf{M}(\partial T), \\ \mathbf{M}(S) &\leq \varepsilon C \mathbf{M}(T), & \mathbf{M}(R) &\leq \varepsilon C \mathbf{M}(\partial T), \end{aligned} \tag{A.1}$$

as well as $\text{spt } R, \text{spt } \partial P \subset \text{Hull}(\text{spt } \partial T)$ and $\text{spt } P, \text{spt } S \subset \text{Hull}(\text{spt } T)$. Moreover,

$$\begin{aligned} \|P\|(\text{int } \sigma) &\leq C \|T\|(\text{st } \sigma), & \|\partial P\|(\text{int } \sigma) &\leq C \|\partial T\|(\text{st } \sigma), \\ \|S\|(\text{int } \sigma) &\leq \varepsilon C \|T\|(\text{st } \sigma), & \|R\|(\text{int } \sigma) &\leq \varepsilon C \|\partial T\|(\text{st } \sigma) \end{aligned} \tag{A.2}$$

for all $\sigma \in \mathcal{F}$.

If X is as in the theorem and $T \in \mathbf{I}_k(X)$ for some $k > n$ then $T = 0$; see Remark A.6. Hence, the statement of the theorem holds also in this case with P, R, S equal to zero.

If M is a smooth Riemannian manifold of dimension n admitting a geometric group action by isometries, then M admits an (n, D, ε) -triangulation for every $\varepsilon > 0$, where $D \geq 1$ depends only on M . For such manifolds a variant of Theorem A.2 without the local mass estimates was established in [15, Theorem 10.3.3]. We furthermore remark that any n -dimensional smooth Riemannian manifold of bounded geometry admits an (n, D, ε) -triangulation for every $\varepsilon > 0$, for some $D \geq 1$ depending only on n and the parameters of bounded geometry (see [6] and the references therein).

We prove Theorem A.2 by analyzing radial projections from simplices of X to their boundaries. Let Δ^m be the Euclidean standard m -simplex and denote by $o \in \text{int } \Delta^m$ and $r_m > 0$ the incenter and inradius of Δ^m , respectively. There exists $K > 0$ only depending on m with the following property. If $b \in \Delta^m \cap U(o, r_m/2)$ then the radial projection $\varrho_b: \Delta^m \setminus \{b\} \rightarrow \partial \Delta^m$ with center b is Kr^{-1} -Lipschitz on $\Delta^m \setminus U(b, r)$ for every $r > 0$. The radial projection ϱ_b is homotopic to the identity map on Δ^m by a straight-line homotopy h . Like the radial projection itself, the Lipschitz constant of h depends on the distance from b , but Proposition A.3 below gives conditions under which we can define the pushforward of T under ϱ_b and the pushforward of $[0, 1] \times T$ under h .

A.1. Limit homotopy formula

Let X be a complete metric space and let $L, \varepsilon > 0$. Suppose that $u: X \rightarrow [0, 1]$ is 1-Lipschitz and $h: [0, 1] \times \{u \neq 0\} \rightarrow X$ is a continuous map which is Lipschitz

on $[0, 1] \times \{u > r\}$ for every $r > 0$. Suppose furthermore that $\text{lip } h_t(x) \leq Lu(x)^{-1}$ and $\text{lip } h_x(t) \leq L\varepsilon$ for all $t \in [0, 1]$ and all $x \in \{u \neq 0\}$, where we have abbreviated $h_t = h(t, \cdot)$ and $h_x = h(\cdot, x)$.

Let $k \geq 1$ and $T \in \mathbf{I}_k(X)$. Then $T \llcorner \{u > r\}$ and $(\partial T) \llcorner \{u > r\}$ are integral currents for almost every $r > 0$ and thus $T_r^i := h_{i\#}(T \llcorner \{u > r\})$ for $i = 0, 1$ and

$$S_r := h_{\#}(\llbracket 0, 1 \rrbracket \times (T \llcorner \{u > r\})), \quad R_r := h_{\#}(\llbracket 0, 1 \rrbracket \times ((\partial T) \llcorner \{u > r\}))$$

are also integral currents. We will call such $r > 0$ admissible.

Proposition A.3. *Suppose $\{B_l\}_{l \in \mathbb{N}}$ is a Borel partition of X (for example, the partition of a countable simplicial complex into the interiors of its faces) such that for every $l \geq 1$,*

$$\int_{B_l} u(x)^{-k} d\|T\|(x) \leq L \|T\|(B_l), \quad \int_{B_l} u(x)^{1-k} d\|\partial T\|(x) \leq L \|\partial T\|(B_l).$$

Then there exist admissible $r_m \searrow 0$ such that $T_{r_m}^i \rightarrow T^i$, $R_{r_m} \rightarrow R$, $S_{r_m} \rightarrow S$ for some $T^0, T^1, R \in \mathbf{I}_k(X)$ and $S \in \mathbf{I}_{k+1}(X)$ with

$$\partial S + R = T^1 - T^0, \tag{A.3}$$

and the following property holds for some $C = C(L, k)$. If $B \subset X$ is Borel then

$$\|T^i\|(B) \leq C\|T\|(A), \quad \|\partial T^i\|(B) \leq C\|\partial T\|(A), \tag{A.4}$$

where $A \subset X$ is the union of all B_l such that $B_l \cap h_i^{-1}(B) \neq \emptyset$, and

$$\|S\|(B) \leq \varepsilon C\|T\|(A'), \quad \|R\|(B) \leq \varepsilon C\|\partial T\|(A'), \tag{A.5}$$

where $A' \subset X$ is the union of all B_l such that $B_l \cap h_t^{-1}(B) \neq \emptyset$ for some $t \in [0, 1]$.

Proof. By the slicing theorem we have for every $r > 0$ that

$$\frac{1}{r} \int_r^{2r} t^{1-k} \mathbf{M}(\langle T, u, t \rangle) dt \leq r^{-k} \|T\|(\{r \leq u \leq 2r\}) \leq 2^k \int_{\{u \leq 2r\}} u(x)^{-k} d\|T\|(x).$$

The last integral tends to zero with $r \rightarrow 0$, so there exist admissible $r_m \searrow 0$ such that

$$\lim_{m \rightarrow \infty} r_m^{1-k} \mathbf{M}(\langle T, u, r_m \rangle) = 0.$$

Set $T_m^i := T_{r_m}^i$ and observe that if $l > m$ then $T_l^i - T_m^i = h_{i\#}(T \llcorner \{r_l < u \leq r_m\})$ and

$$\partial T_l^i - \partial T_m^i = h_{i\#}((\partial T) \llcorner \{r_l < u \leq r_m\}) + h_{i\#}\langle T, u, r_m \rangle - h_{i\#}\langle T, u, r_l \rangle,$$

so Lemma 3.3 implies

$$\mathbf{M}(T_l^i - T_m^i) \leq L^k \int_{\{u \leq r_m\}} u(x)^{-k} d\|T\|(x)$$

and

$$\begin{aligned} \mathbf{M}(\partial T_l^i - \partial T_m^i) &\leq L^{k-1} \int_{\{u \leq r_m\}} u(x)^{1-k} d\|\partial T\|(x) \\ &\quad + L^{k-1} \left[r_m^{1-k} \mathbf{M}(\langle T, u, r_m \rangle) + r_l^{1-k} \mathbf{M}(\langle T, u, r_l \rangle) \right]. \end{aligned}$$

The right hand side in both inequalities tends to zero as both $m, l \rightarrow \infty$, thus (T_m^i) is a Cauchy sequence in $\mathbf{N}_k(X)$ and hence converges to some $T^i \in \mathbf{N}_k(X)$.

Next, set $S_m := S_{r_m}$ and notice that

$$S_l - S_m = h_{\#}(\llbracket 0, 1 \rrbracket \times (T \llcorner \{r_l < u \leq r_m\}))$$

for $l > m$ and thus, by Lemma 3.5,

$$\mathbf{M}(S_l - S_m) \leq (k + 1)L^{k+1}\varepsilon \int_{\{u \leq r_m\}} u(x)^{-k} d\|T\|(x).$$

Since the integral on the right tends to 0 as $m \rightarrow \infty$ it follows that (S_m) is a Cauchy sequence in $\mathbf{M}_{k+1}(X)$ and converges to some $S \in \mathbf{M}_{k+1}(X)$. One shows analogously that $R_m := R_{r_m}$ converges to some $R \in \mathbf{M}_k(X)$ as $m \rightarrow \infty$. Since

$$\partial S_m = T_m^1 - T_m^0 - R_m + h_{\#}(\llbracket 0, 1 \rrbracket \times \langle T, u, r_m \rangle)$$

by the homotopy formula and since

$$\mathbf{M}(h_{\#}(\llbracket 0, 1 \rrbracket \times \langle T, u, r_m \rangle)) \leq k\varepsilon L^k r_m^{1-k} \mathbf{M}(\langle T, u, r_m \rangle) \rightarrow 0$$

as $m \rightarrow \infty$, it follows that $\partial S + R = T^1 - T^0$. From this we conclude that $S \in \mathbf{N}_{k+1}(X)$ and $R \in \mathbf{N}_k(X)$. As T_m^i, S_m, R_m are integral currents for all m and the convergence is in mass it follows that T^i, S, R are also integral currents.

Finally, if $B \subset X$ is Borel and $A \subset X$ is the union of all B_l such that $B_l \cap h_i^{-1}(B) \neq \emptyset$ then, by Lemma 3.3,

$$\|T_m^i\|(B) \leq L^k \int_{h_i^{-1}(B)} u(x)^{-k} d\|T\|(x) \leq L^{k+1}\|T\|(A)$$

and

$$\|\partial T_m^i\|(B) \leq L^{k-1} \int_{h_i^{-1}(B)} u(x)^{1-k} d\|\partial T\|(x) + L^{k-1} r_m^{1-k} \mathbf{M}(\langle T, u, r_m \rangle),$$

which implies (A.4) upon taking $m \rightarrow \infty$. The inequalities in (A.5) are proved analogously using Lemma 3.5. \square

A.2. Special case of the deformation theorem

We will prove the deformation theorem by successive applications of the following proposition, which we prove by applying Proposition A.3 to a map from X to X which restricts to a radial projection on each m -simplex.

Proposition A.4. *Given $m > k \geq 1$ and $D, c \geq 1$ there is a positive constant C with the following property. Let X be a complete c -quasiconvex metric space admitting an (m, D, ε) -triangulation and $T \in \mathbf{I}_k(X)$. Then there exist $P, R \in \mathbf{I}_k(X)$ and $S \in \mathbf{I}_{k+1}(X)$ with $\text{spt } P \subset X^{(m-1)}$ and such that $T = P + R + \partial S$ as well as*

$$\begin{aligned} \mathbf{M}(P) &\leq C \mathbf{M}(T), & \mathbf{M}(\partial P) &\leq C \mathbf{M}(\partial T), \\ \mathbf{M}(S) &\leq \varepsilon C \mathbf{M}(T), & \mathbf{M}(R) &\leq \varepsilon C \mathbf{M}(\partial T) \end{aligned}$$

and $\text{spt } R, \text{spt } \partial P \subset \text{Hull}(\text{spt } \partial T)$, and $\text{spt } P, \text{spt } S \subset \text{Hull}(\text{spt } T)$. Moreover,

$$\begin{aligned} \|P\|(\text{int } \sigma) &\leq C \|T\|(\text{st } \sigma), & \|\partial P\|(\text{int } \sigma) &\leq C \|\partial T\|(\text{st } \sigma), \\ \|S\|(\text{int } \sigma) &\leq \varepsilon C \|T\|(\text{st } \sigma), & \|R\|(\text{int } \sigma) &\leq \varepsilon C \|\partial T\|(\text{st } \sigma), \end{aligned}$$

for every $\sigma \in \mathcal{F}$.

Let $o \in \text{int } \Delta^m$ and $r_m > 0$ be the incenter and inradius of Δ^m as above. As in [17], one can establish that there exists $K > 0$ only depending on m such that if μ_1, μ_2 are finite Borel measures on Δ^m then there exists $b \in \Delta^m \cap U(o, r_m/2)$ such that for $i = 1, 2$,

$$\int_{\text{int } \Delta^m} |y - b|^{1-m} d\mu_i(y) \leq K \mu_i(\text{int } \Delta^m).$$

Proof of Proposition A.4. Denote the metric on X by d . By possibly considering X equipped with the metric $\varepsilon^{-1}d$ we may assume that $\varepsilon = 1$. Let $q: X \rightarrow \Sigma$ be the inverse of an $(m, D, 1)$ -triangulation of X , and let $T \in \mathbf{I}_k(X)$. By the discussion preceding the proof there exists, for every $\sigma \in \mathcal{F}_m$, some point $a_\sigma \in \text{int } \sigma$ such that

$$\int_{\text{int } \sigma} d(x, a_\sigma)^{-k} d\|T\|(x) \leq KD^k \|T\|(\text{int } \sigma)$$

and

$$\int_{\text{int } \sigma} d(x, a_\sigma)^{1-k} d\|\partial T\|(x) \leq KD^k \|\partial T\|(\text{int } \sigma).$$

Set $Z := \{a_\sigma : \sigma \in \mathcal{F}_m\}$ and define a 1-Lipschitz function $u : X \rightarrow [0, 1]$ by

$$u(x) := \min\{1, d(x, Z)\}.$$

Set $r' := \frac{r_m}{2cD}$ and observe that $u(x) \geq r'$ for every $x \in X^{(m-1)}$ and if $\sigma \in \mathcal{F}_m$ then

$$\min\{d(x, a_\sigma), r'\} \leq u(x) \leq d(x, a_\sigma)$$

for every $x \in \sigma$. With the above it follows that for every $\tau \in \mathcal{F}$ we have

$$\int_{\text{int } \tau} u(x)^{-k} d\|T\|(x) \leq K' \|T\|(\text{int } \tau), \quad \int_{\text{int } \tau} u(x)^{1-k} d\|\partial T\|(x) \leq K' \|\partial T\|(\text{int } \tau)$$

for some $K' = K'(D, m, c)$.

Let $h : [0, 1] \times \{u \neq 0\} \rightarrow X$ be the unique map such that for each $\sigma \in \mathcal{F}_m$,

$$q(h(t, x)) = (1 - t)q(x) + t\rho_{q(a_\sigma)}(q(x))$$

for all $t \in [0, 1]$ and all $x \in \sigma$, where we have naturally identified $q(\sigma)$ with Δ^m . We have $h(t, x) = x$ for all $t \in [0, 1]$ and all $x \in X^{(m-1)}$, so this is well-defined. Then $h_0(x) = x$ and $h_1(x) \in X^{(m-1)}$ for all $x \in \{u \neq 0\}$. Moreover, if $A \subset \{u \neq 0\}$ then

$$h([0, 1] \times A) \subset \text{Hull}(A)$$

and if $\sigma, \sigma' \in \mathcal{F}$ are simplices such that $h([0, 1] \times \text{int } \sigma') \cap \text{int } \sigma \neq \emptyset$ then $\sigma' = \sigma$ or σ' is m -dimensional and contains σ .

Let $t \in [0, 1]$ and $x \in \{u \neq 0\}$. Then clearly we have $\text{lip } h_x(t) \leq D^2$. We next show that $\text{lip } h_t(x) \leq K''u(x)^{-1}$ for some $K'' = K''(D, m, c)$. For this, given $y \in X$ let $\sigma(y) \subset X$ be the unique simplex containing y in its interior. Let $x \in \{u \neq 0\}$. For all $y \in X$ sufficiently close to x we have $\sigma(x) \subset \sigma(y)$. If $\sigma(y) \subset X^{(m-1)}$ then $h_t(x) = x$ and $h_t(y) = y$ and therefore $d(h_t(x), h_t(y)) = d(x, y) \leq r'u(x)^{-1} \cdot d(x, y)$. If $\sigma(y)$ is m -dimensional then, letting $b = q(a_{\sigma(y)})$ and identifying $q(\sigma(y))$ with Δ^m , we have

$$\begin{aligned} d(h_t(x), h_t(y)) &\leq D \left((1 - t) |q(x) - q(y)| + t |\rho_b(q(x)) - \rho_b(q(y))| \right) \\ &\leq D^2 \left(1 + \frac{tKD}{\min\{d(x, a_\sigma), d(y, a_\sigma)\}} \right) d(x, y), \end{aligned}$$

which implies $d(h_t(x), h_t(y)) \leq D^2(1 + tKD \min\{u(x), u(y)\}^{-1})d(x, y)$. As a result, $\text{lip } h_t(x) \leq K''(1 + tu(x)^{-1}) \leq 2K''u(x)^{-1}$ for some $K'' = K''(D, m, c)$. Next, since

$\{u \leq r\} = \bigcup B(a_\sigma, r)$ for sufficiently small $r > 0$, it follows that $\{u > r\}$ is quasiconvex and hence, by Lemma 2.4, h is Lipschitz on $[0, 1] \times \{u > r\}$ for sufficiently small $r > 0$.

Finally, let T_r^i, S_r, R_r be defined as at the beginning of Section A.1. By the properties of h listed above, $T_r^0 = T \llcorner \{u > r\}$, $\text{spt } T_r^1 \subset X^{(m-1)}$, and $\text{spt } T_r^1, \text{spt } S_r \subset \text{Hull}(\text{spt } T)$ as well as $\text{spt } R_r \subset \text{Hull}(\text{spt } \partial T)$ for almost every $r > 0$.

By Lemma 2.1, it follows that $q(\text{Hull}(\text{spt } T))$ is a countable simplicial complex and thus $\mathcal{A} := \{\text{int } \sigma : \sigma \in \mathcal{F} \text{ and } \sigma \subset \text{Hull}(\text{spt } T)\}$ is countable. Applying Proposition A.3 with the countable Borel partition $\mathcal{A} \cup \{X \setminus \text{Hull}(\text{spt } T)\}$ shows that there exists a sequence of admissible $r_s \searrow 0$ such that the currents $T_{r_s}^i, S_{r_s}, R_{r_s}$ converge to currents T^i, S, R satisfying the properties listed in Proposition A.3.

Set $P := T^1$. It follows that $T^0 = T$, $\text{spt } P \subset X^{(m-1)}$, and $\text{spt } P, \text{spt } S \subset \text{Hull}(\text{spt } T)$ as well as $\text{spt } R \subset \text{Hull}(\text{spt } \partial T)$. Since $T = P - R - \partial S$ it moreover follows that $\text{spt } \partial P \subset \text{spt } \partial T \cup \text{spt } \partial R \subset \text{Hull}(\text{spt } \partial T)$. The remaining properties of P, S, R listed in the statement of the proposition easily follow from Proposition A.3 and the properties of h listed above. \square

When $k = m$, we can use similar techniques to approximate T by $P \in \mathcal{P}_m(X)$.

Proposition A.5. *Given $m \geq 1$ and $D, c \geq 1$ there is a positive constant C with the following property. Let X be a complete c -quasiconvex metric space admitting an (m, D, ε) -triangulation and $T \in \mathbf{I}_m(X)$. Then there exist $P, R \in \mathbf{I}_m(X)$ with $P \in \mathcal{P}_m(X)$ such that $T = P + R$ as well as*

$$\mathbf{M}(P) \leq C \mathbf{M}(T), \quad \mathbf{M}(\partial P) \leq C \mathbf{M}(\partial T), \quad \mathbf{M}(R) \leq \varepsilon C \mathbf{M}(\partial T)$$

and $\text{spt } R, \text{spt } \partial P \subset \text{Hull}(\text{spt } \partial T)$, and $\text{spt } P \subset \text{Hull}(\text{spt } T)$. Moreover,

$$\|P\|(\sigma) \leq C \|T\|(\sigma), \quad \|\partial P\|(\tau) \leq C \|\partial T\|(\text{st } \tau), \quad \|R\|(\sigma) \leq \varepsilon C \|\partial T\|(\text{int } \sigma)$$

for every $\sigma \in \mathcal{F}_m$ and every $\tau \in \mathcal{F}_{m-1}$.

Proof. We may suppose that $\varepsilon = 1$. After possibly changing the metric on X in a bilipschitz way we may assume by Corollary 4.3 that X is isometric to Σ equipped with the length metric. Let $q: X \rightarrow \Sigma$ be as above and let $\sigma \in \mathcal{F}_m$. Then $q|_\sigma$ is an isometry and we may view it as a map from σ to Δ^m . There exists a function $\theta_\sigma \in L^1(\sigma, \mathbb{Z})$ such that $T \llcorner \text{int } \sigma = \llbracket \theta_\sigma \rrbracket$ and $\|\theta_\sigma\|_1 \leq K_1 \|T\|(\sigma)$ for some $K_1 = K_1(m)$, where $\llbracket \theta_\sigma \rrbracket$ is the m -current in X given by integrating θ_σ along σ .

By the discussion preceding the proof of Proposition A.4, there exists $K_2 = K_2(m)$ such that for every $\sigma \in \mathcal{F}_m$ there are a point $a_\sigma \in \text{int } \sigma$ such that

$$\int_{\text{int } \sigma} d(x, a_\sigma)^{1-m} d\|\partial T\|(x) \leq K_2 \|\partial T\|(\text{int } \sigma) \tag{A.6}$$

and an integer z_σ such that $|z_\sigma| \leq K_2 \|T\|(\sigma)$ and

$$\lim_{r \rightarrow 0} \frac{1}{r^m} \int_{U(a_\sigma, r)} |\theta_\sigma - z_\sigma| d\mathcal{H}^m = 0. \tag{A.7}$$

In particular, $z_\sigma = 0$ for all but finitely many $\sigma \in \mathcal{F}_m$. We claim that

$$P = \sum_{\sigma \in \mathcal{F}_m} z_\sigma \llbracket \sigma \rrbracket$$

and $R = T - P$ satisfy the proposition. By construction, for every $\sigma \in \mathcal{F}_m$, we have $\|P\|(\sigma) \leq C \|T\|(\sigma)$, so $\text{spt } P \subset \text{Hull}(\text{spt } T)$ and $\mathbf{M}(P) \leq C \mathbf{M}(T)$ for some $C = C(m)$.

Let u and h be as above. We bound ∂P and R by applying Proposition A.3 to ∂T . For every admissible $r > 0$, let

$$(\partial T)_r := h_{1\#}(\partial T \llcorner \{u > r\}), \quad S_r := h_{\#}(\llbracket [0, 1] \rrbracket \times (\partial T \llcorner \{u > r\})).$$

As above, there is a $K_3 = K_3(m)$ such that for every $\tau \in \mathcal{F}_m$,

$$\int_{\text{int } \tau} u(x)^{1-m} d\|\partial T\|(x) \leq K_3 \|\partial T\|(\text{int } \tau).$$

Therefore, by Proposition A.3 applied to ∂T , there exist admissible $r_i \searrow 0$, $Q \in \mathbf{I}_{m-1}(X)$, and $S \in \mathbf{I}_m(X)$ with $\text{spt } Q \subset X^{(m-1)}$ and such that $(\partial T)_{r_i} \rightarrow Q$, $S_{r_i} \rightarrow S$, $\partial S = Q - \partial T$, and

$$\|S\|(\text{int } \sigma) \leq C \|\partial T\|(\text{int } \sigma) \quad \|Q\|(\text{int } \tau) \leq C \|\partial T\|(\text{st } \tau) \tag{A.8}$$

for every $\sigma \in \mathcal{F}_m$ and every $\tau \in \mathcal{F}_{m-1}$. We let $R = S$ and claim that $S = P - T$; it follows that $Q = \partial P$, and (A.8) implies the desired bounds on $\|R\|$ and $\|\partial P\|$.

For sufficiently small $r > 0$, let $g_r: X \rightarrow X$ be the continuous map such that

$$q(g_r(x)) = (1 - t_r(x))q(a_\sigma) + t_r(x)q_{q(a_\sigma)}(q(x))$$

for all $x \in \sigma \setminus \{a_\sigma\}$, where $t_r(x) = \min\{1, r^{-1}u(x)\}$. Let moreover $h^r: [0, 1] \times X \rightarrow X$ be the straight-line homotopy from id_X to g_r , i.e., $q(h^r(t, x)) = (1 - t)q(x) + tq(g_r(x))$. Then for $\sigma \in \mathcal{F}_m$, we have $g_r(\sigma \cap \{u \geq r\}) = h_1(\sigma \cap \{u \geq r\}) \subset \partial\sigma$ and $g_r(\sigma \cap \{u \leq r\}) = \sigma$. Furthermore, it follows as in the proof of Proposition A.4 that there exists $K_4 = K_4(m)$ such that $\text{lip } g_r(x) \leq K_4 r^{-1}$ and $\text{lip } h_t^r(x) \leq K_4 r^{-1}$ as well as $\text{lip } h_x^r(t) \leq K_4$, where we have used the notation $h_t^r(x) = h^r(t, x) = h_x^r(t)$ as above. In particular, g_r and h^r are Lipschitz.

The currents $U_i = g_{r_i\#}(T)$ and $V_i = h_{\#}^r(\llbracket [0, 1] \rrbracket \times \partial T)$ satisfy

$$U_i = T + V_i$$

because $W_i = h_{\#}^{r_i}(\llbracket 0, 1 \rrbracket \times T)$ is an $(m + 1)$ -chain in X , hence $W_i = 0$ and therefore

$$0 = \partial W_i = h_{1\#}^{r_i}(T) - h_{0\#}^{r_i}(T) - h_{\#}^{r_i}(\llbracket 0, 1 \rrbracket \times \partial T) = U_i - T - V_i.$$

Since $h^r(t, x) = h(t, x)$ whenever $u(x) > r$, we have $V_i - S_{r_i} = h_{\#}^{r_i}(\llbracket 0, 1 \rrbracket \times (\partial T \llcorner \{u \leq r\}))$. Therefore, letting $K_5 := mK_4^m$,

$$\begin{aligned} \mathbf{M}(V_i - S_{r_i}) &\leq K_5 r_i^{1-m} \mathbf{M}(\partial T \llcorner \{u \leq r_i\}) \\ &\leq K_5 \sum_{\sigma \in \mathcal{F}_m \cap \{u \leq r_i\}} \int d(x, a_\sigma)^{1-m} d\|\partial T\|(x) \leq K_5 K_2 \mathbf{M}(\partial T), \end{aligned}$$

where we use (A.6) in the last inequality. As $i \rightarrow \infty$, each term in the sum goes to zero, so by dominated convergence, $\mathbf{M}(S_{r_i} - V_i) \rightarrow 0$. Thus $V_i \rightarrow S$ in mass and (U_i) converges in mass too. Finally, since g_r is injective on $U(a_\sigma, r)$, we have for each $\sigma \in \mathcal{F}_m$ that $U_i \llcorner \text{int } \sigma = \llbracket \theta_\sigma \circ g_{r_i}^{-1} \llcorner \text{int } \sigma \rrbracket$. By (A.7), for every $\sigma \in \mathcal{F}_m$ we have that $\|\theta_\sigma \circ g_{r_i}^{-1} \llcorner \text{int } \sigma - z_\sigma\|_1 \rightarrow 0$ as $i \rightarrow \infty$. Since (U_i) converges in mass this implies that $U_i \rightarrow P$ in mass. This shows that $S = P - T$, as desired. \square

Remark A.6. It is not difficult to see that if X is an (n, D, ε) -triangulated metric space and $T \in \mathbf{I}_k(X)$ for some $k > n$ then $T = 0$. Indeed, for every simplex σ in X we have $\|T\|(\sigma) = 0$ because the Hausdorff dimension of σ is strictly smaller than k . Now, since $\text{spt } T$ is separable it follows from Lemma 2.1 that $\text{Hull}(\text{spt } T)$ is a countable simplicial complex and hence $\mathbf{M}(T) = \|T\|(\text{Hull}(\text{spt } T)) = 0$. Thus, $T = 0$, as desired.

A.3. Proof of the deformation theorem

Let $p: \Sigma \rightarrow X$ be an (n, D, ε) -triangulation of X and $T \in \mathbf{I}_k(X)$ for some $1 \leq k \leq n$. It is not difficult to show that for each $i = 1, \dots, n - k$ the restriction $p|_{\Sigma^{(n-i)}}$ is an $(n - i, D, \varepsilon)$ -triangulation of $X^{(n-i)}$ and that $X^{(n-i)}$ is quasiconvex with constant only depending on c, n , and D .

In the following, C and C_i will denote a constant depending on D, c, n and the constant may change from one occurrence to another. Set $P_0 := T$. By Proposition A.4, applied to $X^{(n-i+1)}$ and P_{i-1} for $i = 1, \dots, n - k$, we have $P_{i-1} = P_i + R_i + \partial S_i$ for some $P_i, R_i \in \mathbf{I}_k(X^{(n-i+1)})$, $S_i \in \mathbf{I}_{k+1}(X^{(n-i+1)})$ such that $\text{spt } P_i \subset X^{(n-i)}$ and

$$\begin{aligned} \mathbf{M}(P_i) &\leq C_i \mathbf{M}(T), & \mathbf{M}(\partial P_i) &\leq C_i \mathbf{M}(\partial T), \\ \mathbf{M}(S_i) &\leq \varepsilon C_i \mathbf{M}(T), & \mathbf{M}(R_i) &\leq \varepsilon C_i \mathbf{M}(\partial T). \end{aligned} \tag{A.9}$$

Moreover, $\text{spt } P_i, \text{spt } S_i \subset \text{Hull}(\text{spt } T)$ and $\text{spt } \partial P_i, \text{spt } R_i \subset \text{Hull}(\text{spt } \partial T)$ and the following property holds. For each simplex $\sigma \in \mathcal{F}$ we have

$$\begin{aligned} \|P_i\|(\text{int } \sigma) &\leq C\|P_{i-1}\|(\text{st } \sigma), & \|\partial P_i\|(\text{int } \sigma) &\leq C\|\partial P_{i-1}\|(\text{st } \sigma), \\ \|S_i\|(\text{int } \sigma) &\leq \varepsilon C\|P_{i-1}\|(\text{st } \sigma), & \|R_i\|(\text{int } \sigma) &\leq \varepsilon C\|\partial P_{i-1}\|(\text{st } \sigma). \end{aligned} \tag{A.10}$$

By Proposition A.5, we may write $P_{n-k} = P_{n-k+1} + R_{n-k+1}$ with $P_{n-k+1} \in \mathcal{P}_k(X)$ and $R_{n-k+1} \in \mathbf{I}_k(X)$ with $\text{spt } R_{n-k+1} \subset X^{(k)}$. These likewise satisfy $\text{spt } P_{n-k+1} \subset \text{Hull}(\text{spt } T)$ and $\text{spt } \partial P_{n-k+1}, \text{spt } R_{n-k+1} \subset \text{Hull}(\text{spt } \partial T)$, and their masses satisfy (A.9) and (A.10). Let $P := P_{n-k+1}, R := R_1 + \dots + R_{n-k+1}$, and $S := S_1 + \dots + S_{n-k}$. Then $T = P + R + \partial S$ and (A.1) as well as $\text{spt } P, \text{spt } S \subset \text{Hull}(\text{spt } T)$ and $\text{spt } \partial P, \text{spt } R \subset \text{Hull}(\text{spt } \partial T)$ are satisfied. Finally, we show the local mass estimates in (A.2). For any $i \geq 1$ and any $\sigma \in \mathcal{F}$,

$$\|P_i\|(\text{int } \sigma) \leq C\|P_{i-1}\|(\text{st } \sigma) = C \sum_{\sigma \subset \sigma_1} \|P_{i-1}\|(\text{int } \sigma_1),$$

where $\sigma_1 \in \mathcal{F}$. By induction,

$$\|P_i\|(\text{int } \sigma) \leq C^i \sum_{\sigma \subset \sigma_1 \subset \dots \subset \sigma_i} \|T\|(\text{int } \sigma_i),$$

where the sum is taken over all chains of simplices $\sigma_i \in \mathcal{F}$. Let N be the number of chains of simplices $\sigma_0 \subset \sigma_1 \subset \dots \subset \sigma_{n-k} \subset \Delta^n$. Then for any $\sigma \in \mathcal{F}$,

$$\|P\|(\text{int } \sigma) \leq C^{n-k+1} N \|T\|(\text{st } \sigma).$$

The same argument with P_i replaced by ∂P_i gives

$$\|\partial P\|(\text{int } \sigma) \leq C^{n-k+1} N \|\partial T\|(\text{st } \sigma).$$

The estimates of $\|S\|$ and $\|R\|$ then follow from (A.10).

References

[1] A. Abrams, N. Brady, P. Dani, M. Duchin, R. Young, Pushing fillings in right-angled Artin groups, *J. Lond. Math. Soc.* (2) 87 (3) (2013) 663–688.
 [2] F.J. Almgren Jr., The homotopy groups of the integral cycle groups, *Topology* 1 (1962) 257–299.
 [3] L. Ambrosio, B. Kirchheim, Currents in metric spaces, *Acta Math.* 185 (1) (2000) 1–80.
 [4] P. Assouad, Sur la distance de Nagata, *C. R. Acad. Sci. Paris Sér. I Math.* 294 (1) (1982) 31–34.
 [5] G. Bell, A. Dranishnikov, Asymptotic dimension, *Topol. Appl.* 155 (12) (2008) 1265–1296.
 [6] B.H. Bowditch, Bilipschitz triangulations of riemannian manifolds, Warwick, preprint, 2020.
 [7] N. Brodskiy, J. Dydak, J. Higes, A. Mitra, Assouad-Nagata dimension via Lipschitz extensions, *Isr. J. Math.* 171 (2009) 405–423.
 [8] N. Brodskiy, J. Dydak, U. Lang, Assouad-Nagata dimension of wreath products of groups, *Can. Math. Bull.* 57 (2) (2014) 245–253.
 [9] N. Brodskiy, J. Dydak, M. Levin, A. Mitra, A Hurewicz theorem for the Assouad-Nagata dimension, *J. Lond. Math. Soc.* (2) 77 (3) (2008) 741–756.
 [10] J. Burillo, J. Taback, Equivalence of geometric and combinatorial Dehn functions, *N.Y. J. Math.* 8 (2002) 169–179.

- [11] K.-U. Bux, K. Wortman, Finiteness properties of arithmetic groups over function fields, *Invent. Math.* 167 (2) (2007) 355–378.
- [12] D.B. Cohen, Lipschitz 1-connectedness for some solvable Lie groups, *Can. J. Math.* 71 (3) (2019) 533–555.
- [13] T. De Pauw, Approximation by polyhedral G chains in Banach spaces, *Z. Anal. Anwend.* 33 (3) (2014) 311–334.
- [14] J. Dydak, J. Higes, Asymptotic cones and Assouad-Nagata dimension, *Proc. Am. Math. Soc.* 136 (6) (2008) 2225–2233.
- [15] D.B.A. Epstein, J.W. Cannon, D.F. Holt, S.V.F. Levy, M.S. Paterson, W.P. Thurston, *Word Processing in Groups*, Jones and Bartlett Publishers, Boston, MA, 1992.
- [16] H. Federer, *Geometric Measure Theory*, Die Grundlehren der mathematischen Wissenschaften, vol. 153, Springer-Verlag New York Inc., New York, 1969.
- [17] H. Federer, W.H. Fleming, Normal and integral currents, *Ann. Math. (2)* 72 (1960) 458–520.
- [18] T. Goldhirsch, Lipschitz chain approximation of metric integral currents, *Anal. Geom. Metric Spaces* 10 (2022) 40–49.
- [19] T. Goldhirsch, U. Lang, Characterizations of higher rank hyperbolicity, arXiv preprint, arXiv:2112.15565, 2021.
- [20] M. Gromov, Filling Riemannian manifolds, *J. Differ. Geom.* 18 (1) (1983) 1–147.
- [21] M. Gromov, Asymptotic invariants of infinite groups, in: *Geometric Group Theory*, vol. 2, Sussex, 1991, in: *London Math. Soc. Lecture Note Ser.*, vol. 182, Cambridge Univ. Press, Cambridge, 1993, pp. 1–295.
- [22] A. Hatcher, *Algebraic Topology*, Cambridge University Press, Cambridge, 2015.
- [23] J. Huang, B. Kleiner, S. Stadler, Morse quasiflats. II, arXiv:2003.08912, 2020.
- [24] J. Huang, B. Kleiner, S. Stadler, Morse quasiflats. I, *J. Reine Angew. Math.* 784 (2022) 53–129.
- [25] J.R. Isbell, Six theorems about injective metric spaces, *Comment. Math. Helv.* 39 (1964) 65–76.
- [26] S. Keith, A differentiable structure for metric measure spaces, *Adv. Math.* 183 (2) (2004) 271–315.
- [27] B. Kleiner, U. Lang, Higher rank hyperbolicity, *Invent. Math.* 221 (2) (2020) 597–664.
- [28] U. Lang, Local currents in metric spaces, *J. Geom. Anal.* 21 (3) (2011) 683–742.
- [29] U. Lang, T. Schlichenmaier, Nagata dimension, quasisymmetric embeddings, and Lipschitz extensions, *Int. Math. Res. Not.* 58 (2005) 3625–3655.
- [30] E. Le Donne, T. Rajala, Assouad dimension, Nagata dimension, and uniformly close metric tangents, *Indiana Univ. Math. J.* 64 (1) (2015) 21–54.
- [31] E. Leuzinger, R. Young, The distortion dimension of \mathbb{Q} -rank 1 lattices, *Geom. Dedic.* 187 (2017) 69–87.
- [32] E. Leuzinger, R. Young, Filling functions of arithmetic groups, *Ann. Math. (2)* 193 (3) (2021) 733–792.
- [33] G. Link, Higher order Dehn functions for horospheres in products of Hadamard spaces, *Adv. Geom.* 19 (1) (2019) 15–20.
- [34] V. Magnani, Contact equations, Lipschitz extensions and isoperimetric inequalities, *Calc. Var. Partial Differ. Equ.* 39 (1–2) (2010) 233–271.
- [35] D. Meier, S. Wenger, Quasiconformal almost parametrizations of metric surfaces, arXiv preprint, arXiv:2106.01256, 2021.
- [36] J. Nagata, Note on dimension theory for metric spaces, *Fundam. Math.* 45 (1958) 143–181.
- [37] K. Rajala, S. Wenger, An upper gradient approach to weakly differentiable cochains, *J. Math. Pures Appl.* (9) 100 (6) (2013) 868–906.
- [38] T. Schlichenmaier, A quasisymmetrically invariant notion of dimension and absolute Lipschitz retracts, PhD thesis, ETH Zürich, 2005, Diss. Nr. 16216.
- [39] L. Simon, *Lectures on Geometric Measure Theory*, Proceedings of the Centre for Mathematical Analysis, vol. 3, Australian National University, Centre for Mathematical Analysis, Canberra, 1983.
- [40] E.H. Spanier, *Algebraic Topology*, Springer-Verlag, New York, 1995, Corrected reprint of the 1966 original.
- [41] B. Warhurst, Jet spaces as nonrigid Carnot groups, *J. Lie Theory* 15 (1) (2005) 341–356.
- [42] S. Wenger, Isoperimetric inequalities of Euclidean type in metric spaces, *Geom. Funct. Anal.* 15 (2) (2005) 534–554.
- [43] S. Wenger, Flat convergence for integral currents in metric spaces, *Calc. Var. Partial Differ. Equ.* 28 (2) (2007) 139–160.
- [44] S. Wenger, Gromov hyperbolic spaces and the sharp isoperimetric constant, *Invent. Math.* 171 (1) (2008) 227–255.

- [45] S. Wenger, Nilpotent groups without exactly polynomial Dehn function, *J. Topol.* 4 (1) (2011) 141–160.
- [46] S. Wenger, R. Young, Lipschitz extensions into jet space Carnot groups, *Math. Res. Lett.* 17 (6) (2010) 1137–1149.
- [47] B. White, The deformation theorem for flat chains, *Acta Math.* 183 (2) (1999) 255–271.
- [48] R. Young, Filling inequalities for nilpotent groups through approximations, *Groups Geom. Dyn.* 7 (4) (2013) 977–1011.
- [49] R. Young, Lipschitz connectivity and filling invariants in solvable groups and buildings, *Geom. Topol.* 18 (4) (2014) 2375–2417.
- [50] R. Young, Erratum: Lipschitz connectivity and filling invariants in solvable groups and buildings [MR3268779], *Geom. Topol.* 20 (4) (2016) 2433–2438.
- [51] R. Züst, Currents in snowflaked metric spaces, PhD thesis, ETH Zürich, 2011, Diss. Nr. 19966.