

## Nicolai maps with four-fermion interactions

---

Lorenzo Casarin,<sup>a,b</sup> Olaf Lechtenfeld<sup>a</sup> and Maximilian Rupprecht<sup>a</sup>

<sup>a</sup>*Institut für Theoretische Physik and Riemann Center for Geometry and Physics,  
Leibniz Universität Hannover,  
Appelstrasse 2, 30167 Hannover, Germany*

<sup>b</sup>*Max-Planck-Institut für Gravitationsphysik (Albert-Einstein-Institut),  
Am Mühlenberg 1, 14476 Potsdam, Germany*

*E-mail:* [lorenzo.casarin@itp.uni-hannover.de](mailto:lorenzo.casarin@itp.uni-hannover.de),  
[lechtenf@itp.uni-hannover.de](mailto:lechtenf@itp.uni-hannover.de),  
[maximilian.rupprecht@itp.uni-hannover.de](mailto:maximilian.rupprecht@itp.uni-hannover.de)

**ABSTRACT:** Nicolai maps offer an alternative description of supersymmetric theories via nonlinear and nonlocal transformations characterized by the so-called ‘free-action’ and ‘determinant-matching’ conditions. The latter expresses the equality of the Jacobian determinant of the transformation with the one obtained by integrating out the fermions, which so far have been considered only to quadratic terms. We argue that such a restriction is not substantial, as Nicolai maps can be constructed for arbitrary nonlinear sigma models, which feature four-fermion interactions. The fermionic effective one-loop action then gets generalized to higher loops and the perturbative tree expansion of such Nicolai maps receives quantum corrections in the form of fermion loop decorations. The ‘free-action condition’ continues to hold for the classical map, but the ‘determinant-matching condition’ is extended to an infinite hierarchy in fermion loop order. After general considerations for sigma models in four dimensions, we specialize to the case of  $\mathbb{C}P^N$  symmetric spaces and construct the associated Nicolai map. These sigma models admit a formulation with only quadratic fermions via an auxiliary vector field, which does not simplify our analysis.

**KEYWORDS:** Sigma Models, Supersymmetric Gauge Theory, Effective Field Theories, Space-Time Symmetries

ARXIV EPRINT: [2310.19946](https://arxiv.org/abs/2310.19946)

---

**Contents**

<b>1</b>	<b>Introduction and summary</b>	<b>1</b>
<b>2</b>	<b>Supersymmetric nonlinear sigma model</b>	<b>4</b>
<b>3</b>	<b>Supersymmetric <math>\mathbb{CP}^N</math> model</b>	<b>6</b>
<b>4</b>	<b>Adding an auxiliary vector field</b>	<b>11</b>

---

**1 Introduction and summary**

The Nicolai map  $T$  [1–3] is a (generically nonlocal and nonlinear) field transformation that relates supersymmetric theories at different values of their parameters, say coupling constants  $g$ . In particular, it allows one to compute the expectation value of any operator  $Y$  built from the bosonic fields  $\phi$  at coupling  $g$  by evaluating a free-field ( $g=0$ ) correlator of the inversely transformed operator  $T^{-1}Y$ ,<sup>1</sup>

$$\langle Y[\phi] \rangle_g = \langle (T_g^{-1}Y)[\phi] \rangle_0 = \langle Y[T_g^{-1}\phi] \rangle_0. \tag{1.1}$$

Here, we indicated the value of the coupling as a subscript on the correlator and also on the symbol of the map  $T_g : \phi \mapsto \phi'[g, \phi] = T_g\phi$ . The second equality expresses the distributivity of the map, i.e.  $T(\phi_1\phi_2) = (T\phi_1)(T\phi_2)$ . In the expectation values of (1.1) it is understood that all anticommuting degrees of freedom  $\psi$  have been integrated out.<sup>2</sup> This means that the Nicolai map operates in a nonlocal bosonic theory with an action

$$S_g[\phi] = S_g^b[\phi] + \hbar S_g^f[\phi], \tag{1.2}$$

where the local part  $S_g^b$  is the bosonic piece of the original supersymmetric action  $S_{\text{SUSY}}[\phi, \psi]$  and  $\exp\{iS_g^f\}$  arises from the path integral over the anticommuting fields in the partition function, both at coupling  $g$ . In particular,  $S_g^f$  is in general nonlocal and suppressed by  $\hbar$ . So far in the literature only theories with quadratic fermions have been considered, for which the integration over fermions can be formally carried out. As a result, changing path-integral variables  $Y \mapsto T_g Y$  on the right-hand side of (1.1) and sorting powers of  $\hbar$ , one recovers the original defining properties of the Nicolai map,

$$S_0^b[T_g\phi] = S_g^b[\phi] \quad \text{and} \quad S_0^f - i \operatorname{tr} \ln \frac{\delta T_g\phi}{\delta\phi} = S_g^f[\phi], \tag{1.3}$$

which are called the ‘free-action’ and ‘determinant-matching’ property, respectively. We note that  $S_0^f = S_0^f[T_g\phi]$  is a constant since this functional does not depend on the bosonic

---

<sup>1</sup>The vanishing of the vacuum energy in supersymmetric theories properly normalizes  $\langle 1 \rangle_g = 1$ .

<sup>2</sup>Auxiliary fields  $F$  may be kept as part of  $\phi$  or eliminated, see below.

fields. The name ‘determinant-matching’ was indeed coined because in theories with a flat target space integrating the quadratic fermions out produces a fermionic (or Matthews-Salam-Seiler) determinant, which has to be matched by the Jacobian determinant  $\det \frac{\delta T_g \phi}{\delta \phi}$ . Powers of  $\hbar$  are fully explicit because the fermionic determinant sums all fermionic loops in the bosonic background which is a one-loop exact operation. As a consequence, in the language of the ‘Nicolai rules’ to construct the map, only tree diagrams appear, and for this reason it is sometimes considered a classical construction.

In this paper we will show that the assumption of quadratic fermions is not necessary; Nicolai maps can be constructed for supersymmetric actions with higher-order fermion terms as well. The price to pay is quantum corrections to the Nicolai map and a more general dependence on  $\hbar$ , which upsets the conditions (1.3). In other words, the Nicolai map is a formal power series not only in  $g$  but also in  $\hbar$ ,

$$T_g \phi = T_g^{(0)} \phi + \sum_{r=1}^{\infty} \hbar^r T_g^{(r)} \phi, \tag{1.4}$$

where  $r$  counts the number of fermion loops in the diagrammatic representation. With higher-order fermionic terms in the action, the path integral over the anticommuting fields will produce an effective nonlocal action  $\sum_r \hbar^r S_g^{(r)}[\phi]$  extending the previous one-loop result  $\hbar S_g^f[\phi]$  to all orders in  $\hbar$ . Revising the argument leading from (1.1) to (1.3) we then find the combined identity

$$S_0^b[T_g \phi] + \hbar S_0^f - i \hbar \operatorname{tr} \ln \frac{\delta T_g \phi}{\delta \phi} = S_g^b[\phi] + \sum_r \hbar^r S_g^{(r)}[\phi]. \tag{1.5}$$

Inserting (1.4) into (1.5) and comparing powers in  $\hbar$  one obtains an infinite hierarchy of ‘Nicolai-map conditions’. The leading two represent the tree-level and one-loop contributions and read

$$S_0^b[T_g^{(0)} \phi] = S_g^b[\phi] \quad \text{and} \quad S_0^b[T_g \phi]|_{O(\hbar)} + S_0^f - i \operatorname{tr} \ln \frac{\delta T_g^{(0)} \phi}{\delta \phi} = S_g^{(1)}[\phi]. \tag{1.6}$$

Clearly, the ‘free-action condition’ is still valid for the classical part of the Nicolai map, but the ‘determinant-matching condition’ receives a free-action contribution from the one-loop correction to the map, and there are further conditions, each one balancing expressions of a fixed loop order.

For any off-shell supersymmetric theory, there exists a formalism and a universal formula which provides a formal power series expansion (in  $g$ ) of the map and its inverse [4–11].<sup>3</sup> The key player is the ‘coupling flow operator’

$$R_g[\phi] = \int dx (\partial_g T_g^{-1} \circ T_g) \phi(x) \frac{\delta}{\delta \phi(x)}, \tag{1.7}$$

where ‘ $x$ ’ stands for all coordinates the fields depend on. This functional differential operator governs the infinitesimal Nicolai map,

$$\partial_g \langle Y[\phi] \rangle_g = \langle (\partial_g + R_g[\phi]) Y[\phi] \rangle_g. \tag{1.8}$$

---

<sup>3</sup>Under certain conditions the construction works also in the absence of off-shell supersymmetry, e.g. for super Yang-Mills theory in dimensions 6 and 10 in the Landau gauge [8, 12–14].

The first step towards its construction is the observation that, in off-shell supersymmetric chiral theories,

$$S_{\text{SUSY}}[\phi, \psi] = \delta_\alpha \Delta_\alpha[\phi, \psi] \quad \Rightarrow \quad \partial_g S_{\text{SUSY}}[\phi, \psi] = \delta_\alpha \partial_g \Delta_\alpha[\phi, \psi], \quad (1.9)$$

where  $\alpha$  is a spinor index (we are being schematic here on the notation of spinors) and  $\Delta_\alpha$  is a certain anticommuting local functional. Super Yang-Mills theories are more complicated because  $[\partial_g, \delta_\alpha] \neq 0$ , and we exclude them in the following. Employing (1.9) and the supersymmetric Ward identity we integrate out the anticommuting variables to read off the coupling flow operator

$$R_g[\phi] = \frac{i}{\hbar} \partial_g \underbrace{\Delta_\alpha[\phi]} \delta_\alpha = \frac{i}{\hbar} \int dx \partial_g \underbrace{\Delta_\alpha[\phi]} \delta_\alpha \phi(x) \frac{\delta}{\delta \phi(x)}, \quad (1.10)$$

where the contraction indicates a fermionic expectation value to be taken.

Exponentiating the action of  $\partial_g + R_g$  generates the (finite-flow) inverse Nicolai map. Alternatively, a  $g$ -ordered exponential of  $-\int_0^g dg' R_{g'}$  directly yields the Nicolai map. In any case, the  $R_g$  action has to be iterated,  $R_{g_1} R_{g_2} \cdots R_{g_s} \phi$ , which grafts full fermionic  $2k$ -point functions onto previously generated diagrams. For the Wess-Zumino model (quadratic in the fermions) this produces fermionic tree diagrams, dressed with bosonic ‘leaves’. Still, fermion loops remain absent. For nonlinear sigma models, however, the graphical expansion of the Nicolai map will feature fermionic trees with all kinds of fermion loops embedded. Thus it can no longer be considered a classical map. Nevertheless, at any power of the coupling  $g$  a finite number of diagrams contributes to the Nicolai map, and we may still employ the universal formula to write it down. In the following, this will be demonstrated for four-dimensional supersymmetric sigma models, first in general and second for supersymmetric  $\mathbb{CP}^N$  models. The latter, being maximally symmetric, admit a formulation with only quadratic fermions via the introduction of an auxiliary vector field. We do not fully explore its implications here, as this falls outside the scope of the present paper, which presents a more general construction.

For the scope of this paper, relating the perturbative Nicolai map to the standard perturbative (Feynman) expansion solely concerns the generation of diagrams. These are in general divergent and in most applications require regularization. We assume that it has been done in an appropriate way, e.g. via dimensional regularization or with a UV cutoff.<sup>4</sup> These aspects, and more generally the interplay between the Nicolai map and regularization or renormalization, deserve further study.

There are several ways in which the work presented here can be further expanded or generalized. The implications of the auxiliary vector in the  $\mathbb{CP}^N$  model deserve further study.<sup>5</sup> It would be interesting to work out explicitly additional orders of the Nicolai map for  $\mathbb{CP}^N$  models presented here and study in detail the higher-loop identities (1.5).

---

<sup>4</sup>In particular, the perturbative non-renormalizability of the nonlinear sigma models described here is not of concern.

<sup>5</sup>For example, it might be possible to improve the construction presented here by choosing a different normalization for the fields, which is crucial in the construction of the map in the super Yang-Mills case [9]. Work in this direction is ongoing.

Conceivably, then, the formalism presented here can be naturally extended to gauge theories. Finally, a more ambitious goal is the application of the Nicolai map to supergravity, which has been excluded so far owing to the four-fermion contributions, with the potential application of shedding light on its UV behaviour.

The rest of the paper is organized as follows. Section 2 collects some general expressions for supersymmetric nonlinear sigma models and its coupling flow operator. In section 3 we specialize to  $\mathbb{C}P^N$  and construct the associated Nicolai map in perturbation theory. A possibility of eliminating the four-fermion interactions via auxiliary (Hubbard-Stratonovich) scalars or vectors is discussed in section 4. We then comment on the superfield origin of an auxiliary vector in  $\mathbb{C}P^N$  models and its relevance for the construction of the perturbative Nicolai map.

## 2 Supersymmetric nonlinear sigma model

The prototypical supersymmetric field theory with higher-than-quadratic fermion terms in the action is the supersymmetric nonlinear sigma model in (3+1)-dimensional Minkowski space [15], which is characterized by a Kähler potential  $K(\Phi_a^\dagger, \Phi^a)$  and a superpotential  $W(\Phi^a)$  for a collection

$$\{\Phi^a\} = \{\Phi^0, \Phi^A\} = \{\Phi^0, \Phi^1, \Phi^2, \dots, \Phi^N\} \tag{2.1}$$

of  $N+1$  chiral superfields, i.e.  $a, b, c, \dots = 0, 1, \dots, N$  and  $A, B, C, \dots = 1, \dots, N$ . Complex conjugation raises or lowers target-space indices. Adopting the Wess-Bagger notation [16] their component expansion (in  $x$  coordinates) reads

$$\Phi^a = \phi^a + i\theta\sigma^m\bar{\theta}\partial_m\phi^a + \frac{1}{4}\theta^2\bar{\theta}^2\Box\phi^a + \sqrt{2}\theta\psi^a - \frac{i}{\sqrt{2}}\theta^2\partial_m\psi^a\sigma^m\bar{\theta} + \theta^2 F^a =: \phi^a + \Xi^a, \tag{2.2}$$

where Weyl spinor indices  $\alpha, \dot{\alpha} = 1, 2$  are suppressed. We note that  $\Xi^3 = 0$ . For the purpose of this paper the superpotential is not essential (although it can easily be added), and thus we omit any  $F$ -term in the action and restrict ourselves to the Kähler potential in the supersymmetric D-term action

$$S_{\text{SUSY}} = \int d^4x \mathcal{L} \quad \text{with} \quad \mathcal{L} = \int d^2\theta d^2\bar{\theta} K(\Phi_a^\dagger, \Phi^a). \tag{2.3}$$

We further define the functional  $\Delta$  given by

$$\Delta = \int d^4x \mathcal{M} \quad \text{with} \quad \mathcal{M} = \int d\theta d^2\bar{\theta} K(\Phi_a^\dagger, \Phi^a), \tag{2.4}$$

so that

$$\mathcal{L} = -\frac{1}{4}\left(\frac{1+\kappa}{2}\delta^\alpha\mathcal{M}_\alpha + \frac{1-\kappa}{2}\bar{\delta}_{\dot{\alpha}}\bar{\mathcal{M}}^{\dot{\alpha}}\right) + \text{total derivative}, \tag{2.5}$$

with a free parameter  $\kappa$  controlling the relative weight of the chiral against antichiral contribution.<sup>6</sup>

---

<sup>6</sup>This freedom is R symmetry: a D-term can be reached in two ways, by applying either  $\delta^\alpha$  to  $\mathcal{M}_\alpha$  or  $\bar{\delta}_{\dot{\alpha}}$  to  $\bar{\mathcal{M}}^{\dot{\alpha}}$ .

We introduce the notation

$$K := K(\phi_a^*, \phi^a) \quad \text{and} \quad K_{ab\dots}^{cd\dots} := \frac{\partial}{\partial \phi^a} \frac{\partial}{\partial \phi^b} \cdots \frac{\partial}{\partial \phi_c^*} \frac{\partial}{\partial \phi_d^*} \cdots K. \quad (2.6)$$

Expanding

$$\begin{aligned} K &= K + K_a \Xi^a + K^a \bar{\Xi}_a + K_b^a \Xi^b \bar{\Xi}_a + \frac{1}{2} K_{ab} \Xi^a \Xi^b + \frac{1}{2} K^{ab} \bar{\Xi}_a \bar{\Xi}_b \\ &\quad + \frac{1}{2} K_{bc}^a \Xi^b \Xi^c \bar{\Xi}_a + \frac{1}{2} K_a^{bc} \Xi^a \bar{\Xi}_b \bar{\Xi}_c + \frac{1}{4} K_{cd}^{ab} \Xi^a \Xi^b \bar{\Xi}_c \bar{\Xi}_d \\ &= \dots + \bar{\theta}^2 \theta \mathcal{M} + \theta^2 \bar{\theta} \bar{\mathcal{M}} + \theta^2 \bar{\theta}^2 \mathcal{L} \end{aligned} \quad (2.7)$$

and employing the identity

$$\square K = K^{AB} \partial_m \phi_A^* \partial^m \phi_B^* + K_{AB} \partial_m \phi^A \partial^m \phi^B + 2K_B^A \partial_m \phi_A^* \partial^m \phi^B + K^A \square \phi_A^* + K_A \square \phi^A, \quad (2.8)$$

we read off the component lagrangian

$$\begin{aligned} \mathcal{L} &= K_a^b \left[ -\partial^m \phi^a \partial_m \phi_b^* - \frac{i}{2} \psi^a \sigma^m \partial_m \bar{\psi}_b + \frac{i}{2} \partial_m \psi^a \sigma^m \bar{\psi}_b + F^a F_b^* \right] \\ &\quad + \frac{1}{2} K_{ab}^c \left[ i \psi^b \sigma^m \bar{\psi}_c \partial_m \phi^a - \psi^a \psi^b F_c^* \right] + \frac{1}{2} K_c^{ab} \left[ i \bar{\psi}_b \bar{\sigma}^m \psi^c \partial_m \phi_a - \bar{\psi}_a \bar{\psi}_b F^c \right] \\ &\quad + \frac{1}{4} K_{cd}^{ab} \bar{\psi}_a \bar{\psi}_b \psi^c \psi^d \\ &= K_a^b \left[ -\partial^m \phi^a \partial_m \phi_b^* - \frac{i}{2} \psi^a \sigma^m \nabla_m \bar{\psi}_b + \frac{i}{2} \nabla_m \psi^a \sigma^m \bar{\psi}_b + F^a F_b^* \right] \\ &\quad - \frac{1}{2} K_{ab}^c \psi^a \psi^b F_c^* - \frac{1}{2} K_c^{ab} \bar{\psi}_a \bar{\psi}_b F^c + \frac{1}{4} K_{cd}^{ab} \bar{\psi}_a \bar{\psi}_b \psi^c \psi^d \end{aligned} \quad (2.9)$$

with target-space covariant derivative on spinors defined as

$$K_a^b \nabla_m \psi^a = K_a^b \partial_m \psi^a + K_{cd}^b \partial_m \phi^c \psi^d, \quad (2.10)$$

and we also obtain the penultimate component

$$\mathcal{M} = \sqrt{2} K_a^b [\psi^a F_b^* - i \sigma^m \bar{\psi}_b \partial_m \phi^a] + \frac{1}{\sqrt{2}} K_a^{bc} \psi^a \bar{\psi}_b \bar{\psi}_c, \quad (2.11)$$

which are both manifestly invariant under Kähler transformations  $K \mapsto K + \Lambda + \Lambda^\dagger$ .

In order to simplify the coupling-flow operator, it is advisable to also integrate out the auxiliary  $F^a$  by inserting its classical value

$$K_a^b F_b^* = \frac{1}{2} K_a^{cd} \bar{\psi}_c \bar{\psi}_d \quad \Rightarrow \quad F^a = \frac{1}{2} (K^{-1})_b^a K_{cd}^b \psi^c \psi^d \quad (2.12)$$

back into  $\mathcal{L}$  and  $\mathcal{M}$ , which produces

$$\mathcal{L} = K_a^b \left[ -\partial^m \phi^a \partial_m \phi_b^* - \frac{i}{2} \psi^a \sigma^m \nabla_m \bar{\psi}_b + \frac{i}{2} \nabla_m \psi^a \sigma^m \bar{\psi}_b \right] + \frac{1}{4} R_{cd}^{ab} \bar{\psi}_a \bar{\psi}_b \psi^c \psi^d \quad (2.13)$$

with the Riemann tensor

$$R_{cd}^{ab} = K_{cd}^{ab} - K_{cd}^r (K^{-1})_r^s K_s^{ab} \quad (2.14)$$

as well as

$$\mathcal{M} = -i\sqrt{2} K_a^b \sigma^m \bar{\psi}_b \partial_m \phi^a + \sqrt{2} K_a^{bc} \psi^a \bar{\psi}_b \bar{\psi}_c. \tag{2.15}$$

We note that the invertibility of the Kähler metric is not needed in the final expression for the coupling-flow operator. Still, as we shall see, if redundant target coordinates are employed we must constrain them (‘fix a gauge’) and also introduce a coupling  $g$  into the Kähler potential  $K$  in order to obtain a perturbative expansion for the Nicolai map.

Suppose now that  $S_{\text{SUSY}}$  features terms of order  $2k$  in the  $\psi$  fields, with  $k = 1, 2, \dots$ . Then,  $\partial_g \mathcal{M}_\alpha$  has terms of the form  $\psi^{2k-1}$ . Since  $\delta_\alpha$  acting on  $\phi$  is linear in  $\psi$ , the contraction in

$$R_g = -\frac{i}{4\hbar} \int dy \int dx \left\{ \frac{1+\kappa}{2} \partial_g \mathcal{M}_\alpha(y) \delta^\alpha(x) + \frac{1-\kappa}{2} \partial_g \bar{\mathcal{M}}^{\dot{\alpha}}(y) \bar{\delta}_{\dot{\alpha}}(x) \right\} \tag{2.16}$$

signifies a fermionic  $2k$ -point function in the  $\phi$  background, with  $2k-1$  legs fused. For a flat target space, we look at Wess-Zumino chiral models, and fermions appear only quadratically in the action. Thus,  $k=1$  implies only a full fermion propagator in (1.10),

$$\bar{\psi}_b^{\dot{\beta}}(x') \psi_\alpha^a(x) \equiv \hbar (G_2^{(g)})_{b\alpha}^{a\dot{\beta}}(x', x), \tag{2.17}$$

which contains all Feynman diagrams connected by a single fermion line with external  $\phi$  legs. Expanding the full fermion propagator in powers of  $g$ , one obtains chains of free fermion propagators, with vertices encoding the coupling to the bosonic background. No fermion loops arise. In case of a curved target space, instead, we face a nonlinear sigma model, whence  $S_{\text{SUSY}}$  has terms quadratic ( $k=1$ ) as well as quartic ( $k=2$ ) in the fermions. In this case, a  $\psi^3$  contribution in  $\mathcal{M}_\alpha$  produces a correlator of a composite  $\psi^3$  with another  $\psi$  in the bosonic background, which then occurs in (1.10),

$$\psi_\beta^b(x') \bar{\psi}_c^{\dot{\gamma}}(x') \bar{\psi}_d^{\dot{\delta}}(x') \psi_\alpha^a(x) \equiv \hbar^2 (G_4^{(g)})_{cd\beta\alpha}^{ba\dot{\gamma}\dot{\delta}}(x', x), \tag{2.18}$$

which contains all Feynman diagrams connecting a triple fermion vertex with another fermion. We note that both  $G_2^{(g)}$  and  $G_4^{(g)}$  include diagrams with fermion loops generated by the four-fermion coupling. Expanding in powers of  $g$  and taking into account the  $\psi^4$  interaction in  $S_{\text{SUSY}}$ , we encounter fermion loop diagrams in the graphical expansion of the coupling flow operator and hence of the Nicolai map.

### 3 Supersymmetric $\mathbb{CP}^N$ model

Let us become more concrete and specialize to a maximally symmetric Kähler target, namely the complex projective space  $\mathbb{CP}^N \simeq \frac{\text{SU}(N+1)}{\text{SU}(N) \times \text{U}(1)}$ . It is embedded into  $\mathbb{C}^{N+1} \ni \phi^a$  by the identification  $\phi^a \sim \lambda \phi^a$  with  $\lambda \in \mathbb{C}^*$ , which yields the Kähler potential

$$K(\phi_a^*, \phi^a) = \frac{\mu^2}{g} \log \left[ \frac{g}{\mu^2} \phi^* \phi \right] \quad \text{with} \quad \phi^* \phi \equiv \phi_a^* \phi^a, \tag{3.1}$$

where  $g$  is a dimensionless coupling, and  $\mu$  is a mass parameter accounting for the dimensionality of  $\phi$ .<sup>7</sup> The superfield extension identifies  $\Phi^a \sim \Lambda \Phi^a$  with a complex chiral superfield  $\Lambda$ . For later use, we introduce the abbreviations

$$f_g^{-1} := \frac{g}{\mu^2} \phi^* \phi \quad \text{and} \quad \Pi_a^b := \delta_a^b - \frac{\phi^b \phi_a^*}{\phi^* \phi}, \quad (3.2)$$

in terms of which the derivatives of  $K$  are as follows,

$$K_a^b = f_g \Pi_a^b, \quad K_a^{bc} = -2f_g^2 \frac{g}{\mu^2} \phi^{(b} \Pi_a^{c)}, \quad K_{ab}^{cd} = 4f_g^3 \frac{g^2}{\mu^4} \phi^{(c} \Pi_{(a}^{d)} \phi_{b)}^* - 2f_g^2 \frac{g}{\mu^2} \Pi_{(a}^{(c} \Pi_{b)}^{d)}. \quad (3.3)$$

We remark that  $\Pi_a^b \phi^a = 0 = \phi_b^* \Pi_a^b$  implements the projection transversal to the ‘radial direction’ of the identification  $\phi^a \sim \lambda \phi^a$ .

As it stands, this Kähler potential is singular in the  $g \rightarrow 0$  limit. In order to set up a perturbation theory around flat  $\mathbb{C}^N$ , we need to select a point on  $\mathbb{C}P^N$  and pick coordinates  $\phi^A \in \mathbb{C}^N$  centered around it. An appropriate superfield choice fixing the above redundancy is

$$\Phi^0 = \sqrt{\frac{\mu^2}{g}} \Leftrightarrow \{\phi^0, \psi^0, F^0\} = \left\{ \sqrt{\frac{\mu^2}{g}}, 0, 0 \right\}, \quad (3.4)$$

which leaves us with  $N$  proper (super)coordinates

$$\Phi^A = \phi^A + i\theta \sigma^m \bar{\theta} \partial_m \phi^A + \frac{1}{4} \theta^2 \bar{\theta}^2 \square \phi^A + \sqrt{2} \theta \psi^A - \frac{i}{\sqrt{2}} \theta^2 \partial_m \psi^A \sigma^m \bar{\theta} + \theta^2 F^A \quad (3.5)$$

and a well-behaved Kähler potential

$$K(\phi_A^*, \phi^A) = \frac{\mu^2}{g} \log \left[ 1 + \frac{g}{\mu^2} \hat{\phi}^* \hat{\phi} \right] = \hat{\phi}^* \hat{\phi} - \frac{1}{2} \frac{g}{\mu^2} (\hat{\phi}^* \hat{\phi})^2 + O(g^2) \quad \text{with} \quad \hat{\phi}^* \hat{\phi} \equiv \phi_A^* \phi^A. \quad (3.6)$$

Observing that (3.4) effectively reduces the summation ranges in (2.9) and (2.11) to  $1, \dots, N$ , i.e.  $a, b, \dots \rightarrow A, B, \dots$ , we only need to insert

$$K_A^B = f_g \Pi_A^B, \quad K_A^{BC} = -2f_g^2 \frac{g}{\mu^2} \phi^{(B} \Pi_A^{C)}, \quad K_{AB}^{CD} = 4f_g^3 \frac{g^2}{\mu^4} \phi^{(C} \Pi_{(A}^{D)} \phi_{B)}^* - 2f_g^2 \frac{g}{\mu^2} \Pi_{(A}^{(C} \Pi_{B)}^{D)} \quad (3.7)$$

with  $f_g^{-1} = 1 + \frac{g}{\mu^2} \hat{\phi}^* \hat{\phi}$  and

$$\Pi_A^B = \delta_A^B - f_g \frac{g}{\mu^2} \phi^B \phi_A^* \quad \Rightarrow \quad \Pi_A^B \phi^A = f_g \phi^B \quad \text{and} \quad \phi_B^* \Pi_A^B = f_g \phi_A^*. \quad (3.8)$$

The elimination of the auxiliary  $F$  via (2.12) commutes with the coordinate choice for  $\mathcal{M}$ ,

$$\begin{aligned} \mathcal{M} &= -i\sqrt{2} K_A^B \not{\partial} \phi^A \bar{\psi}_B + \sqrt{2} K_A^{BC} \psi^A \bar{\psi}_B \bar{\psi}_C \\ &= -i\sqrt{2} f_g \Pi_A^B \not{\partial} \phi^A \bar{\psi}_B - 2\sqrt{2} f_g^2 \frac{g}{\mu^2} \phi^C \Pi_A^B \psi^A \bar{\psi}_B \bar{\psi}_C, \end{aligned} \quad (3.9)$$

<sup>7</sup>There really is only one (dimensionful) parameter  $M^2 = \mu^2/g$ ; we introduce the dimensionless coupling  $g$  only for later convenience. Also,  $\mu^2$  need not be positive.



but not so for the lagrangian (2.9) because (2.14) requires invertibility of the Kähler metric  $K_a^b$ , which is degenerate in the redundant coordinates. However, after choosing (3.4) we simply obtain

$$K_A^B F_B^* = \frac{1}{2} K_A^{CD} \bar{\psi}_C \bar{\psi}_D \quad \Rightarrow \quad F^A = \frac{1}{2} (K^{-1})_B^A K_{CD}^B \psi^C \psi^D \quad \text{with} \quad (K^{-1})_B^A K_C^B = \delta_C^A \quad (3.10)$$

with an inverse  $K^{-1}$  on  $\mathbb{CP}^N$  and hence a four-fermion interaction

$$\frac{1}{4} R_{CD}^{AB} \bar{\psi}_A \bar{\psi}_B \psi^C \psi^D \quad \text{with} \quad R_{CD}^{AB} = K_{CD}^{AB} - K_{CD}^R (K^{-1})_R^S K_S^{AB}. \quad (3.11)$$

The  $\mathbb{CP}^N$  Riemann tensor computes to (cf. (3.7))

$$R_{CD}^{AB} = -2f_g^2 \frac{g}{\mu^2} \Pi_{(C}^{(A} \Pi_{D)}^{B)} = -\frac{g^2}{\mu^4} (K_C^A K_D^B + K_D^A K_C^B). \quad (3.12)$$

Working out the details one arrives at

$$\begin{aligned} \mathcal{L} = & f_g \Pi_A^B \left[ -\partial_m \phi^A \partial^m \phi_B^* - \frac{i}{2} (\psi^A \not{\partial} \bar{\psi}_B + \bar{\psi}_B \not{\partial} \psi^A) + \frac{i}{2} f_g \frac{g}{\mu^2} (\psi^A \phi^C \not{\partial} \phi_B^* \bar{\psi}_C + \bar{\psi}_B \phi_C^* \not{\partial} \phi^A \psi^C) \right] \\ & + \frac{i}{2} f_g^2 \frac{g}{\mu^2} \Pi_A^B \psi^A (\phi^C \not{\partial} \phi_C^* - \phi_C^* \not{\partial} \phi^C) \bar{\psi}_B + \frac{1}{4} f_g^2 \frac{g}{\mu^2} [\psi^C \sigma_m \Pi_C^A \bar{\psi}_A] [\psi^D \sigma^m \Pi_D^B \bar{\psi}_B], \end{aligned} \quad (3.13)$$

where we have employed the Fierz identity

$$[\bar{\psi}_A \bar{\psi}_B] [\psi^C \psi^D] = -\frac{1}{2} [\psi^C \sigma_m \bar{\psi}_A] [\psi^D \sigma^m \bar{\psi}_B]. \quad (3.14)$$

We now have all the ingredients for constructing the Nicolai map. Remembering that

$$\delta_\alpha \phi^A = \sqrt{2} \psi_\alpha^A, \quad \delta_\alpha \phi_A^* = 0, \quad , \quad \bar{\delta}^{\dot{\alpha}} \phi^A = 0, \quad \bar{\delta}^{\dot{\alpha}} \phi_A^* = \sqrt{2} \bar{\psi}_A^{\dot{\alpha}} \quad (3.15)$$

the coupling-flow operator takes the form

$$\begin{aligned} R_g = & -\frac{i}{4\hbar} \sqrt{2} \int dy \int dx \left\{ \frac{1+\kappa}{2} \partial_g \underline{\mathcal{M}_\alpha(y)} \psi^{A\alpha}(x) \frac{\delta}{\delta \phi^A(x)} + \frac{1-\kappa}{2} \partial_g \underline{\bar{\mathcal{M}}^{\dot{\alpha}}(y)} \bar{\psi}_{A\dot{\alpha}}(x) \frac{\delta}{\delta \phi_A^*(x)} \right\} \\ = & -\frac{1}{2\hbar} \int dy \int dx \frac{1+\kappa}{2} \left\{ \partial_g (f_g \Pi_A^B) \not{\partial}_{\alpha\dot{\alpha}} \phi^A(y) \underline{\bar{\psi}_B^{\dot{\alpha}}(y)} \psi^{D\alpha}(x) \frac{\delta}{\delta \phi^D(x)} \right. \\ & \left. - 2i \partial_g \left( f_g^2 \frac{g}{\mu^2} \Pi_A^B \right) \phi^C(y) \underline{\psi_\alpha^A(y) \bar{\psi}_{B\dot{\alpha}}(y) \bar{\psi}_C^{\dot{\alpha}}(y)} \psi^{D\alpha}(x) \frac{\delta}{\delta \phi^D(x)} \right\} \\ & - \frac{1}{2\hbar} \int dy \int dx \frac{1-\kappa}{2} \left\{ \partial_g (f_g \Pi_A^B) \bar{\not{\partial}}^{\dot{\alpha}\alpha} \phi_B^*(y) \underline{\psi_\alpha^A(y) \bar{\psi}_{D\dot{\alpha}}(x)} \frac{\delta}{\delta \phi_D^*(x)} \right. \\ & \left. - 2i \partial_g \left( f_g^2 \frac{g}{\mu^2} \Pi_A^B \right) \phi_C^*(y) \underline{\bar{\psi}_B^{\dot{\alpha}}(y) \psi_\alpha^A(y) \psi_\alpha^C(y)} \bar{\psi}_{D\dot{\alpha}}(x) \frac{\delta}{\delta \phi_D^*(x)} \right\} \end{aligned} \quad (3.16)$$

with full fermionic correlators indicated by the contractions, see (2.17) and (2.18).

Let us take a look at first order in the coupling  $g$ ,

$$(T_g \phi)^A(x) = \phi^A(x) - g (R_g \phi)|_{g=0}^A(x) + O(g^2). \quad (3.17)$$

We compute

$$\begin{aligned}
 (R_g \phi)|_{g=0}^A(x) &= \frac{1+\kappa}{4\hbar\mu^2} \int dy \left\{ \phi_B^* (\phi^B \not{\partial}_{\alpha\dot{\alpha}} \phi^C + \phi^C \not{\partial}_{\alpha\dot{\alpha}} \phi^B) \underbrace{\bar{\psi}_C^{\dot{\alpha}}(y)} \psi^{A\alpha}(x) \right. \\
 &\quad \left. + 2i \phi^C \underbrace{\psi_\alpha^B \bar{\psi}_{B\dot{\alpha}} \bar{\psi}_C^{\dot{\alpha}}(y)} \psi^{A\alpha}(x) \right\}, \\
 (R_g \phi^*)|_A^{g=0}(x) &= \frac{1-\kappa}{4\hbar\mu^2} \int dy \left\{ \phi^B (\phi_B^* \bar{\not{\partial}}^{\dot{\alpha}\alpha} \phi_C^* + \phi_C^* \bar{\not{\partial}}^{\dot{\alpha}\alpha} \phi_B^*) \underbrace{\psi_\alpha^C(y)} \bar{\psi}_{A\dot{\alpha}}(x) \right. \\
 &\quad \left. + 2i \phi_C^* \underbrace{\bar{\psi}_B^{\dot{\alpha}} \psi^{B\alpha} \psi_\alpha^C(y)} \bar{\psi}_{A\dot{\alpha}}(x) \right\},
 \end{aligned} \tag{3.18}$$

where now the contractions are free-field ones,

$$\begin{aligned}
 \sigma_{\alpha\dot{\alpha}}^m \underbrace{\bar{\psi}_C^{\dot{\alpha}}(y)} \psi^{A\alpha}(x) &= -2\hbar \delta_C^A \partial^m \square^{-1}(y-x) = \bar{\sigma}^{m\dot{\alpha}\alpha} \underbrace{\psi_\alpha^A(y)} \bar{\psi}_{B\dot{\alpha}}(x), \\
 \underbrace{\psi_\alpha^B \bar{\psi}_{B\dot{\alpha}} \bar{\psi}_C^{\dot{\alpha}}(y)} \psi^{A\alpha}(x) &= \hbar^2 (N+1) \delta_C^A \partial_m \square^{-1}(y-z) \partial^m \square^{-1}(y-x)|_{z=y},
 \end{aligned} \tag{3.19}$$

and arrive at the classical map

$$\begin{aligned}
 (T_g^{(0)} \phi)^A(x) &= \phi^A(x) + \frac{1+\kappa}{2} \frac{g}{\mu^2} \int dy \phi_B^* \partial_m (\phi^B \phi^A)(y) \partial^m \square^{-1}(y-x) + O(g^2), \\
 (T_g^{(0)} \phi^*)_A(x) &= \phi_A^*(x) + \frac{1-\kappa}{2} \frac{g}{\mu^2} \int dy \phi^B \partial_m (\phi_B^* \phi_A^*)(y) \partial^m \square^{-1}(y-x) + O(g^2),
 \end{aligned} \tag{3.20}$$

while the four-fermion contraction yields the leading (or one-loop) quantum contribution

$$\begin{aligned}
 (T_g^{(1)} \phi)^A(x) &= -i \frac{g}{\mu^2} \frac{1+\kappa}{2} (N+1) \int dy \phi^A(y) \partial_m \square^{-1}(y-z) \partial^m \square^{-1}(y-x)|_{z=y} \\
 &\quad + O(g^2), \\
 (T_g^{(1)} \phi^*)_A(x) &= -i \frac{g}{\mu^2} \frac{1-\kappa}{2} (N+1) \int dy \phi_A^*(y) \partial_m \square^{-1}(y-z) \partial^m \square^{-1}(y-x)|_{z=y} \\
 &\quad + O(g^2),
 \end{aligned} \tag{3.21}$$

The generalized free-action condition in (1.6) means that

$$S_0^b[T_g^{(0)} \phi] = - \int \partial_m (T_g \phi)^A \partial^m (T_g \phi^*)_A \stackrel{!}{=} - \int f_g \Pi_A^B \partial_m \phi^A \partial^m \phi_B^* = S_g^b[\phi], \tag{3.22}$$

which is met to first order in  $g$  because both sides are equal to

$$- \int \left\{ \partial_m \phi^A \partial^m \phi_A^* - \frac{g}{\mu^2} \phi_B^* (\phi^B \partial_m \phi^A + \phi^A \partial_m \phi^B) \partial^m \phi_A^* + O(g^2) \right\} \tag{3.23}$$

and  $\kappa$  cancels out. For the one-loop matching in (1.6) we obtain the  $O(g)$  contributions

$$\begin{aligned}
 S_0^b[T_g \phi]|_{O(\hbar)} &= i \frac{g}{\mu^2} (N+1) \int \left\{ \frac{1}{2} \phi^A \phi_A^* \delta(0) + \frac{\kappa}{2} (\phi^A \square \phi_A^* - \phi_A^* \square \phi^A) \square^{-1}(0) \right\}, \\
 -i \operatorname{tr} \ln \frac{\delta T_g^{(0)} \phi}{\delta \phi} &= i \frac{g}{\mu^2} (N+1) \int \left\{ \frac{1}{2} \phi^A \phi_A^* \delta(0) - \frac{\kappa}{2} (\phi^A \square \phi_A^* - \phi_A^* \square \phi^A) \square^{-1}(0) \right\}, \\
 S_g^{(1)}[\phi] &= i \frac{g}{\mu^2} (N+1) \int \left\{ \phi^A \phi_A^* \delta(0) \right\},
 \end{aligned} \tag{3.24}$$

which verifies the condition again with the expected cancellation of  $\kappa$ . In (3.24) we use the regularization-dependent quantities  $\delta(0) \equiv \delta^{(4)}(x-x)$  and  $\square^{-1}(0) \equiv \square^{-1}(x-x)$ , which contribute to a mass shift and wavefunction renormalization, respectively.<sup>8</sup> We kept them in this form to maintain the discussion as general as possible and to illustrate the matching at first order.

By choosing  $\kappa=+1$  or  $\kappa=-1$  we have the freedom to shift the Nicolai map entirely to  $\phi^A$  or  $\phi_A^*$  alone, respectively. A graphical representation of the Nicolai map to order  $g^2$  looks as follows,

$$\begin{aligned}
 T_g \phi = & \bullet \text{---} \text{wavy} + g \bullet \text{---} \text{wavy} + g^2 \left( \bullet \text{---} \text{wavy} + \bullet \text{---} \text{wavy} \right) + \dots \\
 & + \hbar g \bullet \text{---} \text{loop} + \hbar g^2 \left( \bullet \text{---} \text{loop} + \bullet \text{---} \text{loop} + \bullet \text{---} \text{loop} + \bullet \text{---} \text{loop} \right) + \dots \\
 & + \hbar^2 g^2 \left( \bullet \text{---} \text{loop} + \bullet \text{---} \text{loop} + \bullet \text{---} \text{loop} \right) + \dots \\
 & + O(g^3). \tag{3.25}
 \end{aligned}$$

Here, the thick dot at the left end of each diagram stands for the argument  $x$  of the map, other vertex positions are integrated over. Solid lines are free fermion propagators  $\not{\partial}\square^{-1}$  or  $\not{\partial}\square^{-1}$ , and wavy lines represent bosonic field insertions  $\phi$  or  $\phi^*$ . One of the bosonic legs emanating from each vertex not sourcing a loop carries a derivative (not shown). For the full ‘Nicolai rules’, one of course needs to add target-space indices, spinor traces, and weight factors. All diagrams shown above already appear in the first application of  $R_g$  on  $\phi$ . We see that in the  $\hbar$  expansion of the map an  $r$ -loop contribution arises first at order  $g^r$ , so that at each given order in perturbation theory only a finite number of diagrams contribute, as expected.

---

<sup>8</sup>In dimensional regularization both contributions vanish in a massless theory such as the one under consideration. With a UV cutoff they are quartically and quadratically divergent, respectively.

## 4 Adding an auxiliary vector field

In some field theories with four-fermion interactions one can ‘resolve’ the latter through a coupling with an auxiliary field  $A$ : the Hubbard-Stratonovich transformation. Schematically, one adds to an action with a  $(\bar{\psi}\psi)^2$  term an auxiliary-field coupling  $(\bar{\psi}\psi - A)^2$ , schematically

$$\bar{\psi} i \partial \psi + \frac{1}{4} g (\bar{\psi}\psi)^2 \quad \longrightarrow \quad \bar{\psi} i \partial \psi + \frac{1}{2} g A \bar{\psi}\psi - \frac{1}{4} g A^2 = \bar{\psi} i \left( \partial - \frac{i}{2} g A \right) \psi - \frac{1}{4} g A^2. \quad (4.1)$$

The four-fermion term has been cancelled, but eliminating  $A$  brings it back. Hence, the only price is an additional auxiliary field (or several of them). Filling in the indices in our schematic argument and allowing for the fierzing (3.14), we see that the transformation requires our four-fermion term (3.11) to be ‘factorizable’, i.e.

$$R_{CD}^{AB} = \lambda_s (\bar{m}_i)^{AB} (m^i)_{CD} + \lambda_v (\ell_I)_C^A (\ell^I)_D^B \quad (4.2)$$

with some coefficients  $\lambda_s$  and  $\lambda_v$  and indices  $i$  and  $I$  counting several such terms. For target geometries with this property we can remove the four-fermion interaction by introducing a bunch of complex scalar and real vector auxiliary fields

$$A^i = (m^i)_{CD} \psi^C \psi^D \quad \text{and} \quad A_m^I = (\ell^I)_D^B \psi^D \sigma_m \bar{\psi}_B, \quad (4.3)$$

respectively.

This is actually the case for all hermitian symmetric spaces [17, 18].<sup>9</sup> Therefore, these geometries allow for an auxiliary-field reformulation. The complex projective spaces  $\mathbb{C}P^N$  treated above are the maximally symmetric compact examples, and indeed (3.12) shows that a single real vector auxiliary  $A_m$  suffices. Actually, the Hubbard-Stratonovich trick can be slightly generalized by shifting  $A$  by an arbitrary function of bosonic fields. We make use of this option and choose

$$A_m = \frac{1}{\mu^2} f_g [i \phi^C \partial_m \phi_C^* - i \phi_C^* \partial_m \phi^C + \psi^A \Pi_A^B \sigma_m \bar{\psi}_B], \quad (4.4)$$

where the form of the (arbitrary) bosonic contribution will be justified later on. With some algebra this yields the enhanced lagrangian

$$\begin{aligned} \tilde{\mathcal{L}} = & -f_g \Pi_A^B \partial_m \phi^A \partial^m \phi_B^* + \frac{1}{4} f_g^2 \frac{g}{\mu^2} (\phi^A \partial_m \phi_A^* - \phi_A^* \partial_m \phi^A)^2 \\ & + \frac{i}{2} f_g g (\phi^A \partial_m \phi_A^* - \phi_A^* \partial_m \phi^A) A^m - \frac{1}{4} g \mu^2 A_m A^m \\ & - \frac{i}{2} f_g \Pi_A^B (\psi^A \not{\partial} \bar{\psi}_B + \bar{\psi}_B \not{\partial} \psi^A) - \frac{1}{2} f_g g \Pi_A^B \psi^A \not{A} \bar{\psi}_B \\ & + \frac{i}{2} f_g^2 \frac{g}{\mu^2} \Pi_A^B (\psi^A \phi^C \not{\partial} \phi_B^* \bar{\psi}_C + \bar{\psi}_B \phi_C^* \not{\partial} \phi^A \psi^C). \end{aligned} \quad (4.5)$$

There exists an instructive superfield formulation [17, 18],

$$\tilde{\mathcal{S}}_{\text{SUSY}} = \int d^4x \tilde{\mathcal{L}} \quad \text{with} \quad \tilde{\mathcal{L}} = \int d^2\theta d^2\bar{\theta} \{ e^{gV} \Phi_a^\dagger \Phi^a - \mu^2 V \}, \quad (4.6)$$

<sup>9</sup>See [19] for a more general review on nonlinear realization and hidden local symmetries.

and corresponding expressions for  $\tilde{\Delta}$  and  $\tilde{\mathcal{M}}$ , where the auxiliary real vector superfield  $V$  with components  $(C, L, A_m, \lambda, \chi, D)$  expands in  $x$  coordinates as

$$V = -C - i\theta\chi + i\bar{\theta}\bar{\chi} - \frac{i}{2}\theta^2 L + \frac{i}{2}\bar{\theta}^2 L^* + \theta\sigma^m\bar{\theta}A_m - i\theta^2\bar{\theta}\left[\bar{\lambda} + \frac{i}{2}\bar{\theta}\bar{\chi}\right] + i\bar{\theta}^2\theta\left[\lambda + \frac{i}{2}\theta\bar{\chi}\right] - \frac{1}{2}\theta^2\bar{\theta}^2\left[D + \frac{1}{2}\square C\right]. \quad (4.7)$$

The action (4.6) enjoys a complexified local U(1) invariance under

$$V \mapsto V + i\Lambda - i\Lambda^\dagger \quad \text{and} \quad \Phi^a \mapsto e^{-ig\Lambda} \Phi^a \quad (4.8)$$

with a chiral superfield parameter  $\Lambda$ . The coordinate choice (3.4) is not compatible with the Wess-Zumino gauge, and it completely breaks the gauge symmetry. Indeed, eliminating the vector superfield by its algebraic equation of motion,

$$V = -\frac{1}{g} \log \left[ \frac{g}{\mu^2} \Phi_a^\dagger \Phi^a \right] = -\frac{1}{\mu^2} K[\Phi^a, \Phi_a^\dagger], \quad (4.9)$$

brings back the original action (2.3) with the  $\mathbb{CP}^N$  Kähler potential (3.1) (up to a constant). We can interpret the action (4.6) and the gauge transformation (4.8) as a way of performing the  $\mathbb{CP}^N$  identification of scalar superfields under complex chiral parameters: the supergauge transformation realizes in a supersymmetric fashion a U(1) local transformation, thus extending the standard gauge invariance to a  $U(1)_{\mathbb{C}}$  symmetry including conformal rescaling.

Let us go now to the component level and eliminate auxiliary fields  $F, L, \chi, \lambda, C$  and  $D$  from the action by their equations of motion, e.g.

$$F^a = -\psi^a \frac{\psi^b \phi_b^*}{\phi^* \phi}, \quad \chi = -\frac{i\sqrt{2}}{g} \frac{\psi^a \phi_a^*}{\phi^* \phi}, \quad g e^{-gC} \phi^* \phi = \mu^2. \quad (4.10)$$

After lengthy but straightforward computations we arrive at

$$\begin{aligned} \tilde{\mathcal{L}} = f_g \Pi_a^b \left\{ -D_m \phi^a D^m \phi_b^* - \frac{i}{2} [\psi^a \not{D} \bar{\psi}_b + \bar{\psi}_b \not{D} \psi^a] + \frac{i}{2} \frac{1}{\phi^* \phi} [\psi^a \phi^c \not{D} \phi_b^* \bar{\psi}_c + \bar{\psi}^b \phi_c^* \not{D} \phi^a \psi^c] \right\} \\ + \frac{1}{4} f_g^2 \frac{g}{\mu^2} (\phi^a D_m \phi_a^* - \phi_a^* D_m \phi^a)^2 \end{aligned} \quad (4.11)$$

where  $D_m = \partial_m - \frac{i}{2} g A_m$  is a U(1)-covariant derivative. We note that  $\Pi_a^b D_m \phi^a = \Pi_a^b \partial_m \phi^a$ . For the penultimate components we find

$$\tilde{\mathcal{M}} = -i\sqrt{2} f_g \Pi_a^b \not{D} \phi^a \bar{\psi}_b + i\sqrt{2} f_g^2 \frac{g}{\mu^2} (\phi^b \not{D} \phi_b^* - \phi_b^* \not{D} \phi^b) \phi^a \bar{\psi}_a. \quad (4.12)$$

Both  $\tilde{\mathcal{L}}$  and  $\tilde{\mathcal{M}}$  are manifestly gauge invariant. We are left with the fields  $\phi^a, \psi^a$  and  $A_m$ . Eliminating the latter via its equation of motion

$$\mu^2 A_m = f_g [i\phi^c \partial_m \phi_c^* - i\phi_c^* \partial_m \phi^c + \Pi_a^b \psi^a \sigma_m \bar{\psi}_b] \Leftrightarrow i\phi^c D_m \phi_c^* - i\phi_c^* D_m \phi^c = -\Pi_a^b \psi^a \sigma_m \bar{\psi}_b \quad (4.13)$$

reverts to the four-fermion interaction, so we keep  $A_m$  in the lagrangian.

Instead, we now employ the local supersymmetric  $U(1)_{\mathbb{C}}$  invariance to fix one of the chiral superfields,  $\Phi^0 = \sqrt{\frac{\mu^2}{g}}$ , which explicitly connects the auxiliary-superfield formulation with our coordinate choice (3.4) for the nonlinear sigma model. It is not hard to see that this gauge fixing comes with a trivial Faddeev-Popov determinant. The gauge-fixed lagrangian then indeed agrees with (4.5). Using the identities  $\Pi_A^B \phi^A = f_g \phi^B$  and  $\frac{g}{\mu^2} \phi^A \phi_A^* = f_g^{-1} - 1$  as well as

$$\begin{aligned} \phi^c D \phi_c^* - \phi_c^* D \phi^c &= \phi^C \partial \phi_C^* - \phi_C^* \partial \phi^C + i f_g^{-1} \mu^2 A, \\ \Pi_a^b D \phi^a \bar{\psi}_b &= \Pi_A^B D \phi^A \bar{\psi}_B + \frac{i}{2} f_g g A \phi^A \bar{\psi}_A, \end{aligned} \quad (4.14)$$

we also obtain

$$\tilde{\mathcal{M}} = -i\sqrt{2} f_g \Pi_A^B \not{\partial} \phi^A \bar{\psi}_B + i\sqrt{2} f_g^2 \frac{g}{\mu^2} (\phi^C \not{\partial} \phi_C^* - \phi_C^* \not{\partial} \phi^C) \phi^A \bar{\psi}_A - \sqrt{2} f_g g A \phi^A \bar{\psi}_A. \quad (4.15)$$

Furthermore, after this gauge-fixing the value of  $A_m$  in (4.13) is exactly (4.4).

Again, we can set up the Nicolai map. Employing

$$\delta_\alpha Y = \sqrt{2} \psi_\alpha \frac{\delta Y}{\delta \phi} + \delta^\alpha A_m \frac{\delta Y}{\delta A_m}, \quad (4.16)$$

with a rather complicated expression for  $\delta^\alpha A_m$ , which we shall not reproduce here, and

$$\frac{\mu^2}{\sqrt{2}} \partial_g \tilde{\mathcal{M}}_\alpha = i f_g^2 \phi^B \phi_B^* \not{\partial}_{\alpha\dot{\alpha}} \phi^A \bar{\psi}_A^{\dot{\alpha}} + i f_g^3 (1 - \frac{g}{\mu^2} \phi^C \phi_C^*) \phi^B \not{\partial}_{\alpha\dot{\alpha}} \phi_B^* \phi^A \bar{\psi}_A^{\dot{\alpha}} - \mu^2 f_g^2 A_{\alpha\dot{\alpha}} \phi^A \bar{\psi}_A^{\dot{\alpha}}, \quad (4.17)$$

we may compose the coupling flow operator

$$\tilde{R}_g = -\frac{i}{4} \frac{1+\kappa}{2} \int dy \partial_g \tilde{\mathcal{M}}(y)_\alpha \delta^\alpha - \frac{i}{4} \frac{1-\kappa}{2} \int dy \partial_g \tilde{\mathcal{M}}(y)^{\dot{\alpha}} \bar{\delta}_{\dot{\alpha}}, \quad (4.18)$$

with the contractions now being defined in a  $(\phi, A)$  background.

It appears that we can arrive at a purely classical Nicolai map, in the sense described in the Introduction. After all, the fermions appear just quadratically in the lagrangian (4.11). To leading order in the coupling  $g$ , it looks as follows,

$$\begin{aligned} (T_g \phi)^A(x) &= \phi^A(x) + \frac{1+\kappa}{2} \frac{g}{\mu^2} \int dy [\phi^B (\phi_B^* \partial_m \phi^A - \phi^A \partial_m \phi_B^*) + i \mu^2 A_m \phi^A](y) \partial^m \square^{-1}(y-x) \\ &+ O(g^2), \end{aligned} \quad (4.19)$$

and similarly for  $(T_g A)_m$ . Checking the free-action condition

$$\begin{aligned} \int \partial_m (T_g \phi)^A \partial^m (T_g \phi^*)_A &\stackrel{!}{=} \int \left\{ f_g \Pi_A^B \partial_m \phi^A \partial^m \phi_B^* - \frac{1}{4} f_g^2 \frac{g}{\mu^2} (\phi^C \partial_m \phi_C^* - \phi_C^* \partial_m \phi^C)^2 \right. \\ &\left. - \frac{i}{2} f_g g (\phi^C \partial_m \phi_C^* - \phi_C^* \partial_m \phi^C) A^m + \frac{1}{4} g \mu^2 A_m A^m \right\} \end{aligned} \quad (4.20)$$

we obtain the first-order requirement

$$\begin{aligned}
 & \int \left\{ (\phi^* \phi) (\partial \phi^* \cdot \partial \phi) + \frac{1+\kappa}{2} (\phi \partial \phi^*) \cdot (\phi \partial \phi^*) + \frac{1-\kappa}{2} (\phi^* \partial \phi) \cdot (\phi^* \partial \phi) \right. \\
 & \quad \left. + \frac{1+\kappa}{2} i \mu^2 A \cdot (\phi \partial \phi^*) - \frac{1-\kappa}{2} i \mu^2 A \cdot (\phi^* \partial \phi) \right\} \\
 & \stackrel{!}{=} \int \left\{ (\phi^* \phi) (\partial \phi^* \cdot \partial \phi) + (\phi^* \partial \phi) \cdot (\phi \partial \phi^*) \right. \\
 & \quad \left. - \frac{1}{4} [\mu^2 A - i(\phi \partial \phi^* - \phi^* \partial \phi)] \cdot [\mu^2 A - i(\phi \partial \phi^* - \phi^* \partial \phi)] \right\}, \tag{4.21}
 \end{aligned}$$

where the dots indicate Lorentz contractions, and the round brackets enclose target coordinate contractions. We observe that the left-hand side is linear in  $A$  while the right-hand side is quadratic. Matching both sides (and cancelling  $\kappa$ ) requires putting

$$\mu^2 A_m \stackrel{!}{=} i(\phi^C \partial_m \phi_C^* - \phi_C^* \partial_m \phi^C) + O(g) + O(\hbar), \tag{4.22}$$

which agrees with the equation of motion (4.13) after integrating out the fermions! The crux of the mismatch, however, lies in the absence of  $A$  in the free action, which renders the  $g \rightarrow 0$  limit singular for the auxiliary field. In other words, the propagator for  $A$  is proportional to  $\frac{1}{g}$ , which upsets the perturbative expansion in  $g$  (not in powers of the fields) of correlation functions, with or without the Nicolai map. The remedy is to integrate out the auxiliary vector  $A$  and work with an effective theory of  $\phi$  alone. This, however, revives the four-fermion interaction: the classical solution (4.13) shows that  $A$  yields a fermion loop, and the ultralocal propagator  $\langle A(x) A(y) \rangle \sim \delta(x-y)$  glues two such loops together, effectively reproducing the four-fermion interaction. It seems that the auxiliary-field reformulation of the supersymmetric nonlinear sigma models does not simplify the Nicolai map in the end. However, the situation is somewhat reminiscent of the super Yang-Mills case [9], where one needs to use specific field redefinitions to define the map. A similar construction might apply to this case too and deserves further study.

Let us conclude with a comment on the  $N=1$  case. We have the accidental isomorphism  $\mathbb{CP}^1 \simeq S^2$ , the real 2-sphere, which is maximally symmetric in the real sense and whose Riemann tensor admits therefore the standard decomposition  $\text{Riem} = gg - gg$  in real coordinates. In this case indices  $A, B, \dots$  have only one value and  $\Pi \equiv f_g$ . Correspondingly, the four-fermion term of the lagrangian (3.13) takes the form

$$\mathcal{L}_{\psi^4} = -\frac{1}{2} f_g^3 \frac{g}{\mu^2} \psi \psi \bar{\psi} \bar{\psi}, \tag{4.23}$$

which can be resolved by means of a standard Hubbard-Stratonovich auxiliary complex scalar  $A \sim \psi \psi$ . Of course, the clash with the perturbative expansion in  $g$  described in (4.20)–(4.21) applies also in this case.

## Acknowledgments

We thank Hermann Nicolai for discussions.

**Open Access.** This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

## References

- [1] H. Nicolai, *On a new characterization of scalar supersymmetric theories*, *Phys. Lett. B* **89** (1980) 341 [[INSPIRE](#)].
- [2] H. Nicolai, *Supersymmetry and functional integration measures*, *Nucl. Phys. B* **176** (1980) 419 [[INSPIRE](#)].
- [3] H. Nicolai, *Supersymmetric functional integration measures*, in the proceedings of the 1984 *NATO ASI on supersymmetry*, Bonn, Germany, 20–31 August 1984, K. Dietz et al. eds., Plenum Press (1984), p. 393 [[INSPIRE](#)].
- [4] R. Flume and O. Lechtenfeld, *On the stochastic structure of globally supersymmetric field theories*, *Phys. Lett. B* **135** (1984) 91 [[INSPIRE](#)].
- [5] K. Dietz and O. Lechtenfeld, *Nicolai maps and stochastic observables from a coupling constant flow*, *Nucl. Phys. B* **255** (1985) 149 [[INSPIRE](#)].
- [6] O. Lechtenfeld, *Construction of the Nicolai mapping in supersymmetric field theories*, Ph.D. thesis, Bonn University, Bonn, Germany (1984) [[INSPIRE](#)].
- [7] O. Lechtenfeld, *Stochastic variables in ten-dimensions?*, *Nucl. Phys. B* **274** (1986) 633 [[INSPIRE](#)].
- [8] S. Ananth et al., *Perturbative linearization of supersymmetric Yang-Mills theory*, *JHEP* **10** (2020) 199 [[arXiv:2005.12324](#)] [[INSPIRE](#)].
- [9] O. Lechtenfeld and M. Rupperecht, *Universal form of the Nicolai map*, *Phys. Rev. D* **104** (2021) L021701 [[arXiv:2104.00012](#)] [[INSPIRE](#)].
- [10] O. Lechtenfeld and M. Rupperecht, *Is the Nicolai map unique?*, *JHEP* **09** (2022) 139 [[arXiv:2207.09471](#)] [[INSPIRE](#)].
- [11] O. Lechtenfeld, *The Nicolai-map approach to supersymmetry*, in the proceedings of the 12<sup>th</sup> *International Symposium on Quantum Theory and Symmetries*, (2023) [[arXiv:2309.00481](#)] [[INSPIRE](#)].
- [12] H. Malcha and H. Nicolai, *Perturbative linearization of super-Yang-Mills theories in general gauges*, *JHEP* **06** (2021) 001 [[arXiv:2104.06017](#)] [[INSPIRE](#)].
- [13] O. Lechtenfeld and M. Rupperecht, *Construction method for the Nicolai map in supersymmetric Yang-Mills theories*, *Phys. Lett. B* **819** (2021) 136413 [[arXiv:2104.09654](#)] [[INSPIRE](#)].
- [14] O. Lechtenfeld and M. Rupperecht, *An improved Nicolai map for super Yang-Mills theory*, *Phys. Lett. B* **838** (2023) 137681 [[arXiv:2211.07660](#)] [[INSPIRE](#)].
- [15] M. Bertolini, *Lectures on supersymmetry*, <https://people.sissa.it/~bertmat/susycourse.pdf>, 21 November 2023.
- [16] J. Wess and J. Bagger, *Supersymmetry and supergravity*, Princeton University Press, Princeton, NJ, U.S.A. (1992).



- [17] K. Higashijima and M. Nitta, *Supersymmetric nonlinear sigma models as gauge theories*, *Prog. Theor. Phys.* **103** (2000) 635 [[hep-th/9911139](#)] [[INSPIRE](#)].
- [18] K. Higashijima and M. Nitta, *Quantum equivalence of auxiliary field methods in supersymmetric theories*, *Prog. Theor. Phys.* **103** (2000) 833 [[hep-th/9911225](#)] [[INSPIRE](#)].
- [19] M. Bando, T. Kugo and K. Yamawaki, *Nonlinear realization and hidden local symmetries*, *Phys. Rept.* **164** (1988) 217 [[INSPIRE](#)].