



HAL
open science

Early release of the expanded atlas of the sky in continuous gravitational waves

Vladimir Dergachev, Maria Alessandra Papa

► **To cite this version:**

Vladimir Dergachev, Maria Alessandra Papa. Early release of the expanded atlas of the sky in continuous gravitational waves. 2023. hal-04314137

HAL Id: hal-04314137

<https://hal.science/hal-04314137>

Preprint submitted on 29 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Copyright

Early release of the expanded atlas of the sky in continuous gravitational waves

Vladimir Dergachev^{1,2,a} and Maria Alessandra Papa^{1,2,b}

¹Max Planck Institute for Gravitational Physics (Albert Einstein Institute), Callinstraße 38, 30167 Hannover, Germany

²Leibniz Universität Hannover, D-30167 Hannover, Germany

We present the early release of the atlas of continuous gravitational waves covering frequencies from 20 Hz to 1500 Hz and spindowns from -5×10^{-10} to 5×10^{-10} Hz/s. Compared to the previous atlas release we have greatly expanded the parameter space, and we now also provide polarization-specific data - both for signal-to-noise ratios and for the upper limits. Continuous wave searches are computationally difficult and take a long time to complete. The atlas enables new searches to be performed using modest computing power. To allow new searches to start sooner, we are releasing this data early, before our followup stages have completed.

I. INTRODUCTION

This paper presents the early release of the expanded atlas of continuous gravitational waves. The atlas contains upper limits and signal-to-noise ratios (SNR) as a function of signal frequency and polarization, and source sky position. Compared to the first atlas [1], we have greatly expanded the searched parameter space (Fig. 1) and provide polarization-specific information, while maintaining comparable sensitivity.

The atlas is constructed from the results of the first two stages of the Falcon search [1–4]. Subsequent stages do not improve the sensitivity of the analysis, but are used to identify potential signals. While we wait for these followups to complete, we are releasing the atlas data so that other scientists can use it earlier.

The data in the atlas can be used to find signals from neutron stars with equatorial deformations of $\approx 100 \mu\text{m}$ up to 200 pc away. Such sources could well exist in our immediate galactic neighbourhood.

The atlas contains seven billion records. It can be analyzed on a small personal computer thanks to the MVL library [5, 6]. The atlas data, example code, and documentation can be found online [7]. We also recommend to consult [1] and [8].

In the following sections, we present a brief overview of the key scientific aspects of the atlas.

II. CONTINUOUS WAVE SIGNALS

Falcon performs loosely coherent searches for bundles of signals with frequency evolution $f(t)$ of the IT2 type [3] plus a small higher order perturbation:

$$f(t) = f_0 + (t-t_0)f_1 + (t-t_0)^2 f_2/2 + \sum_{n=3}^{\infty} (t-t_0)^n \frac{f_n}{n!}, \quad (1)$$

This search covers signals with f_0 from 20 to 1500 Hz, $|f_1| \leq 5 \times 10^{-10}$ Hz/s, $|f_2| \leq 4 \times 10^{-20}$ Hz/s² and higher




FIG. 1. This plot shows the expected gravitational wave frequency and frequency-derivative of isolated neutron stars from the ATNF catalog [33] (circles). The curves show the frequency/frequency-derivative combinations for a gravitar (a neutron star losing energy solely due to gravitational wave emission) for different values of the ellipticity. Following [34] we observe that known pulsars have spin-down values that lie at or above the curve $\epsilon = 10^{-9}$. For gravitars this would imply that the minimum ellipticity is around 10^{-9} . The region above the grey trace and enclosed by the box of green straight lines shows the frequency/frequency-derivative combinations of optimally oriented gravitars at 100 pc, that our search could detect. The frequency-derivative of a detectable gravitar is proportional to the square of the distance, so if the distance increases by a factor of 10, correspondingly the grey trace moves up by a factor of 100.

order terms:

$$|f_{n \geq 3}| \leq \frac{2.9 \times 10^{-6} \text{ Hz} \cdot n!}{(1.5 \times 10^7 \text{ s})^n}. \quad (2)$$

The reference time (GPS epoch) is $t_0 = 1246070000$ (2019 Jul 2 02:33:02 UTC).

^a vladimir.dergachev@aei.mpg.de

^b maria.alessandra.papa@aei.mpg.de

It is expected that a rotating neutron star with an equatorial quadrupolar deformation will emit continuous gravitational waves. The amplitude of these waves grows as the first power of the deformation parameter ϵ (“ellipticity”) and with the second power of the frequency.

Theory predicts that the neutron star crust can support quadrupolar deformations of 10^{-6} and perhaps even larger for neutron stars with exotic composition [9–11], however, signals like this have not yet been detected [12–17].

Other potential sources are boson condensates formed around rapidly spinning blackholes. This source is highly hypothetical and depends on the existence of a particle with just the right mass to produce gravitational waves that can be registered with gravitational wave detectors [19–23]. Signals from boson condensates are expected to have $|\dot{f}| \leq 10^{-11}$ Hz/s during most of their lifetime [20].

The spindown range for this search was chosen to cover rotating neutron stars with ellipticities of 10^{-7} up to 1500 Hz plus an extra margin to allow for increased spindown due to magnetic fields or kinematic effects.

Our search uses data of the LIGO H1 and L1 interferometers, from the publicly released O3a set [24]. We limit our search to the LIGO detectors because no other detector comes within a factor of 2 of their sensitivity. The frequency band 20-36 Hz is used to veto short-duration power excess stemming from disturbances. This can affect upper limits in this band, but only for exceedingly loud signals that are excluded by prior searches.

We have reserved the O3b data [25] for validation of any potential outliers that may persist following our ongoing follow-up investigations. This approach is commonly used for searches conducted across an extensive parameter space. Given the substantial number of trials involved, noise alone can yield outliers, and these can be reliably discarded through verification using a separate dataset, as illustrated in [14, 26]. It is important to emphasize that this procedure is applicable because we assume the presence of the same physical signal in both datasets, a fundamental characteristic of continuous wave signals.

III. GRAVITATIONAL WAVE POLARIZATION DATA

A rotating source such as an equatorially deformed neutron star will emit circularly polarized waves along the rotation axis and linearly polarized waves perpendicular to the rotation axis.

Gravitational wave detectors presently in operation are not omnidirectional and at any point in time there are locations on the sky where a detector is completely insensitive to some linearly polarized signals. Because continuous wave searches aggregate data over months, the motion of the detector around the Earth and around the Sun will average the detector response and allow to detect an arbitrary source.

Depending on the orientation of the source to the detector line of sight, the detector will see either a single linearly polarized wave, or a mix of two. Correspondingly the amount of power received by the detector can vary by a factor of two. For this reason the sensitivity to continuous gravitational waves varies a lot depending on polarization as can be seen in figure 2, with circularly polarized upper limits more than a factor of two lower than linearly polarized ones.

In the past we only reported the largest upper limit of all polarizations (worst case) and the circular polarization upper limit. This is far from optimal if one is interested in an elliptically polarized signal - the circularly polarized limit is too small, and the worst case upper limit is unnecessarily large.

In this atlas release, we were able to include *functional upper limit* data [8]. That data consists of 14 numbers that are provided for each atlas frequency bin and each sky location. The 14 numbers are input parameters to a function of ι and ψ , and for every pair (ι, ψ) the function establishes a 95% confidence upper limit.




FIG. 2. Gravitational wave intrinsic amplitude h_0 upper limits at 95% confidence as a function of signal frequency. The upper limits are a measure of the sensitivity of the search. We also plot latest LIGO/Virgo and Einstein@Home all-sky population average results [14, 16].

IV. RESULTS

The Falcon pipeline calculates upper limits on the intrinsic amplitude h_0 of continuous gravitational waves across signal frequency and position. Our atlas includes upper limits for circularly polarized gravitational waves




FIG. 3. Reach of the search for stars with ellipticity of 10^{-8} . The search is also sensitive to sources with ellipticities of 10^{-7} with a distance from Earth that is 10 times higher. The X axis is the gravitational wave frequency, which is twice the pulsar rotation frequency for emission due to an equatorial ellipticity. R-modes and other emission mechanisms give rise to emission at different frequencies [35]. The top curve (purple) shows the reach for a population of circularly polarized signals; The bottom curve (blue) holds for linearly polarized signals.

originating from optimally aligned sources, and worst-case upper limits derived by maximizing over polarization. These upper limits are applicable to sources in the least favorable orientation.

Newly included are the functional upper limits. As this is the first implementation, there are imperfections that result in overly conservative figures. As discussed in [8], the functional upper limits can be as much as 5% larger than the original upper limits for noise-dominated data, and as much as 30% larger for high SNR signals. Example code showing how to use functional upper limits is included in the atlas release.

The new functional upper limits have been tested with software and hardware simulated signals (“injections”). Figure 4 shows established functional upper limits using injection-specific ι and ψ versus the actual injection strength. Table I shows the ratio of functional upper limits to injection amplitudes evaluated for ι and ψ corresponding to each hardware injection [27].

The atlas also includes the measured SNR as a function of signal frequency and source sky-position. Specifically, the atlas contains sky-resolved data for every 45 mHz signal-frequency band. The sky resolution increases proportionally to frequency, and at 1500 Hz at any location in the sky there is a grid point within a 0.17° radius. For each sky pixel and frequency bin, we also provide the




FIG. 4. Polarization-specific upper limits versus injected strain for a set of more than 31000 software injections. The upper limits are computed using ι and ψ of the injected signals.

highest SNR value and the corresponding frequency, as well as ι and ψ where the SNR peak has occurred.

The upper limits are summarized in Figures 2 and 3, by taking the maximum over the sky in 45 mHz frequency bands. This is the traditional way to present continuous wave upper limit results.

The amplitude of gravitational waves received at the detector depends on the distance d to the source, the source ellipticity, and the source frequency:

$$h_0 = \frac{4\pi^2 G I_{zz} f_0^2 \epsilon}{c^4 d}, \quad (3)$$

where $I_{zz} = 10^{38} \text{ kg m}^2$ is the moment of inertia of the star with respect to the principal axis aligned with the rotation axis.

Equation 3 suggests a convenient figure of merit to judge the sensitivity of the search, a sort of horizon distance; figure 3 shows the distance to which our upper limits exclude sources with ellipticity of 10^{-8} as a function of frequency, for different assumed polarizations.

This search is sensitive to sources with ellipticity of 10^{-8} up to 200 pc away - an improvement on our previous results [1] due to expanded reach at high frequencies.

The table I shows the atlas results associated to hardware injected signals, obtained by script `spatial_index_example2.R`. The upper limits are polarization-specific and were computed using the ι and ψ values of each injection.

The frequency corresponding to the SNR peak reflects the injection frequency only when the injection is loud enough and when its parameters are within the searched

area. Most injections within the searched parameter space are recovered within 0.1 mHz of the signal frequency. The only exception is hardware injection ip11 located in a highly contaminated low frequency band.

Signals outside the searched parameter space are not necessarily lost. For example, despite injection ip7 being on the edge of detectability, with SNR just shy of 12 and having a large spindown, it is clearly seen in Figure 5, with the peak slightly offset from the signal location. The established upper limit is however smaller than the injected amplitude, indicating, as expected, a loss of sensitivity due to parameter mismatch.

The findings in [16] span a much wider spin-down range than our search. They effectively rule out neutron stars with ellipticities larger than 10^{-6} within a few kiloparsecs, prompting a search for lower ellipticity sources. We accomplish this with a spindown range that covers ellipticities as high as 10^{-7} , with higher sensitivity than [16].

The atlas is made possible due to our efficient analytical upper limit procedure [28] that computes results separately for each point in the parameter space. In contrast, many searches [12, 14, 16, 21, 29] use time-consuming Monte-Carlo simulations to establish upper limits over the entire sky, and hence are unsuitable for the construction of an atlas.

All-sky surveys aimed at detecting continuous gravitational waves are extremely demanding in terms of computational resources. The search reported here required over 281 million CPU-hours (where CPU refers to a single execution thread). Our atlas data provides an opportunity for others to leverage this significant investment in several ways:

- **Followup of Astronomical Observations:** The data in our atlas enables followup of astronomical observations related to pulsations. Observations from a specific celestial location can be compared against high Signal-to-Noise Ratio (SNR) occurrences at the same location and consistent frequencies.
- **Efficient Frequency Scanning:** The atlas allows for rapid scanning across the entire frequency range at specific sky positions of interest. For example, this can be applied to the study of a supernova remnant, as demonstrated in the appendix, to search for significant results.
- **Ellipticity Constraints:** With an estimate of the distance to a gravitational wave source, it becomes straightforward to compute upper limits on its ellipticity.
- **Predictive Modeling:** Our atlas data can be directly incorporated into models of neutron star populations, facilitating predictions regarding their detectability. This same approach can be applied to primordial black hole populations emitting continuous gravitational waves through orbital energy

loss [30] or tens of solar mass black hole populations generating continuous waves through super-radiance [19].

- **Statistical Significance:** The fine granularity of our atlas provides a rich dataset. In the event of coincidences with other observations, this granularity greatly aids in estimating the chance probability of any discovery.

Label	f Hz	\dot{f} Hz/s	Binary	SNR	UL/ h_0 %	Δf mHz	In
ip0	265.57505	-4.15e-12	No	28.5	122.5	-0.1	Yes
ip1	848.93498	-3e-10	No	393.0	119.9	-0.1	Yes
ip2	575.16351	-1.37e-13	No	39.3	138.5	0.0	Yes
ip3	108.85716	-1.46e-17	No	23.7	141.6	0.1	Yes
ip4	1390.60583	-2.54e-08	No	7.6	21.3	-7.7	No
ip5	52.80832	-4.03e-18	No	155.9	130.2	0.0	Yes
ip6	145.39178	-6.73e-09	No	8.4	25.0	-11.2	No
ip7	1220.42586	-1.12e-09	No	7.3	68.1	3.6	No
ip8	190.03185	-8.65e-09	No	8.9	83.8	-2.9	No
ip9	763.84732	-1.45e-17	No	39.1	135.1	0.1	Yes
ip10	26.33210	-8.5e-11	No	63.9	124.9	0.0	Yes
ip11	31.42470	-5.07e-13	No	93.2	400.9	-12.1	Yes
ip12	37.75581	-6.25e-09	No	14.0	156.5	4.0	No
ip16	234.56700	0	Yes	8.3	29.6	42.7	No
ip17	890.12300	0	Yes	8.1	103.6	23.6	No

TABLE I. This table shows parameters of the hardware-injected continuous wave signals and atlas data for their locations and frequencies. The upper limits for the injections are polarization specific and were computed using ι and ψ of each injection. We show all the hardware injections within 20-1500 Hz range, including those outside of our search space, as indicated by the ‘‘In’’ column. We use the reference time (GPS epoch) $t_0 = 1246070000$ (2019 Jul 2 02:33:02 UTC).

ACKNOWLEDGMENTS

The authors thank the scientists, engineers and technicians of LIGO, whose hard work and dedication produced the data that made this search possible.

The search was performed on the ATLAS cluster at AEI Hannover. We thank Bruce Allen, Carsten Aulbert and Henning Fehrmann for their support.

This research has made use of data or software obtained from the Gravitational Wave Open Science Center (gw-openscience.org), a service of LIGO Laboratory, the LIGO Scientific Collaboration, the Virgo Collaboration, and KAGRA.

Appendix: Using the Falcon atlas to search supernova remnants

It is easy to use our atlas to examine arbitrary locations on the sky. An example R script




FIG. 5. Summary of atlas data from the bins between 1220.2-1220.4 Hz. The top panels show the highest SNR (left) and upper limit values (right) across the frequency band, for each pixel of the sky map, using equatorial coordinates. The red lines denote the galactic plane. The black triangle shows the location of hardware injection. The blue band of smaller SNRs near the ecliptic equator is due to large correlations between waveforms of sources in that region. The blue regions in the upper limit plot are due to the lower-SNR values in the ecliptic plane, and also occur near the ecliptic poles that are favored by the antenna pattern of the detectors. The bottom panels show the same data as a function of frequency and with the maximum taken over the sky. We mark the frequency of the band where the hardware injection was performed. The data and code used to produce this plot are available [7].

`spatial_index_example3.R` included with this release [7] shows how to produce upper limits and SNR values for any given sky location. By simply setting right ascension and declination to the coordinates of the Vela Jr or G189.1+3.0 supernova remnants, we obtain the data shown in Figures 6 and 7, respectively. Compared to plots shown in [1], our data covers a much larger frequency and spindown range.

We also plot the latest LIGO/Virgo/KAGRA results [31, 32]. Our spindown range is smaller than that searched by [31, 32], but it covers observed spindowns

of known pulsars. Furthermore, our results are rigorous 95% confidence level upper limits, rather than sensitivity estimates, and we have no excluded frequency bands.

These plots illustrate that an atlas user can establish rigorous upper limits that improve on the best dedicated directed searches.

This data can be used as is to place limits on the gravitational radiation coming from direction of these supernova remnants as we have done, or it can be the starting point for a deeper search.

-
- [1] Vladimir Dergachev and Maria Alessandra Papa, Frequency-Resolved Atlas of the Sky in Continuous Gravitational Waves, *Phys. Rev. X* **13**, 021020 (2023)
- [2] V. Dergachev, M. A. Papa, Results from the first all-sky search for continuous gravitational waves from small-ellipticity sources, *Phys. Rev. Lett.* **125**, no.17, 171101 (2020)
- [3] V. Dergachev, M. A. Papa, Results from high-frequency all-sky search for continuous gravitational waves from small-ellipticity sources, *Phys. Rev. D* **103**, 063019 (2021)
- [4] V. Dergachev, M. A. Papa, The search for continuous gravitational waves from small-ellipticity sources at low frequencies, *Phys. Rev. D* **104**, 043003 (2021)
- [5] Mappable Vector Library, <https://github.com/volodya31415/libMVL>
- [6] R package for Mappable Vector Library <https://cran.r-project.org/package=RMVL>
- [7] See EPAPS Document No. [number will be inserted by publisher] for numerical values of upper limits. Also at <https://www.aei.mpg.de/continuouswaves/03aFalcon20-1500> including the full atlas.
- [8] Vladimir Dergachev, Functional upper limits, arXiv:2311.09911 (2023)
- [9] N.K. Johnson-McDaniel and B.J. Owen, Maximum elastic deformations of relativistic stars, *Phys. Rev. D* **88**, 044004 (2013)
- [10] F. Gittins, N. Andersson and D. I. Jones, Modelling neutron star mountains, *Mon. Not. Roy. Astron. Soc.* **500**, no.4, 5570-5582 (2020)
- [11] F. Gittins and N. Andersson, Modelling neutron star mountains in relativity, *Mon. Not. Roy. Astron. Soc.* **507**, no.1, 116-128 (2021)
- [12] K. Riles, Searches for Continuous-Wave Gravitational Radiation, *Living Reviews in Relativity* **26**, 3 (2023)




FIG. 6. This plot shows upper limits similar to those shown in Figure 2, but for the location of Vela Jr. Latest LIGO/Virgo/KAGRA results [31] are shown for comparison. The triangles mark saturated bands for which the Weave results are invalid.




FIG. 7. This plot shows upper limits similar to those shown in Figure 2, but for the location of G189.1+3.0. Latest LIGO/Virgo/KAGRA results [32] are shown for comparison. The LVK upper limit curve was computed as minimum of Hanford, Livingston and Virgo data.

- [13] G. Pagliaro, M. A. Papa, J. Ming, J. Lian, D. Tsuna, C. Maraston and D. Thomas, Continuous Gravitational Waves from Galactic Neutron Stars: Demography, Detectability, and Prospects, *Astrophys. J.* **952**, no.2, 123 (2023)
- [14] B. Steltner, M. A. Papa, H. B. Eggenstein, R. Prix, M. Bensch, B. Allen and B. Machenschalk, Deep Einstein@Home All-sky Search for Continuous Gravitational Waves in LIGO O3 Public Data, *Astrophys. J.* **952**, no.1, 55 (2023)
- [15] R. Abbott *et al.* (LIGO Scientific Collaboration, Virgo Collaboration and KAGRA Collaboration), All-sky search for continuous gravitational waves from isolated neutron stars in the early O3 LIGO data, *Phys. Rev. D* **104**, 082004 (2021)
- [16] R. Abbott *et al.* (LIGO Scientific Collaboration, Virgo Collaboration and KAGRA Collaboration), All-sky search for continuous gravitational waves from isolated neutron stars using Advanced LIGO and Advanced Virgo O3 data, *Phys. Rev. D* **106**, 102008 (2022)
- [17] R. Abbott *et al.* [LIGO Scientific, VIRGO and KAGRA], Searches for Gravitational Waves from Known Pulsars at Two Harmonics in the Second and Third LIGO-Virgo Observing Runs, *Astrophys. J.* **935**, no.1, 1 (2022)
- [18] A. Tripathhee, K. Riles, Probing More Deeply in an All-Sky Search for Continuous Gravitational Waves in the LIGO O3 Data Set, arXiv:2311.04985
- [19] A. Arvanitaki and S. Dubovsky, Exploring the string axiverse with precision black hole physics, *Phys. Rev. D* **83** 044026 (2011)
- [20] A. Arvanitaki, M. Baryakhtar, X. Huang, Discovering the QCD Axion with Black Holes and Gravitational Waves, *Phys. Rev. D* **91**, 084011 (2015)
- [21] R. Abbott *et al.* (LIGO Scientific, VIRGO and KAGRA collaborations), All-sky search for gravitational wave emission from scalar boson clouds around spinning black holes in LIGO O3 data, *Phys. Rev. D* **105**, 102001 (2022)
- [22] S. J. Zhu, M. Baryakhtar, M. A. Papa, D. Tsuna, N. Kawanaka and H. B. Eggenstein, Characterizing the continuous gravitational-wave signal from boson clouds around Galactic isolated black holes, *Phys. Rev. D* **102**, no.6, 063020 (2020)
- [23] A. Arvanitaki, S. Dimopoulos, S. Dubovsky, N. Kaloper and J. March-Russell, String axiverse, *Phys. Rev. D* **81**, 123530 (2010)
- [24] The O3a Data Release <https://doi.org/10.7935/nfnt-hm34>
- [25] The O3b Data Release <https://doi.org/10.7935/pr1e-j706>
- [26] M. A. Papa, J. Ming, E. V. Gotthelf, B. Allen, R. Prix, V. Dergachev, H. B. Eggenstein, A. Singh and S. J. Zhu, Search for Continuous Gravitational Waves from the Central Compact Objects in Supernova Remnants Cassiopeia A, Vela Jr., and G347.3-0.5, *Astrophys. J.* **897**, no.1, 22 (2020)
- [27] C. Bower, D. Barker, J.C. Batch, J. Betzwieser, R.P. Fisher, E. Goetz, *et al.*, Validating gravitational-wave detections: The Advanced LIGO hardware injection system, *Phys. Rev. D* **95**, 062002 (2017)
- [28] V. Dergachev, A Novel Universal Statistic for Computing Upper Limits in Ill-behaved Background, *Phys. Rev. D* **87**, 062001 (2013).
- [29] P. B. Covas, M. A. Papa, R. Prix and B. J. Owen, Con-

- straints on r-modes and Mountains on Millisecond Neutron Stars in Binary Systems, *Astrophys. J. Lett.* **929**, no.2, L19 (2022)
- [30] A.L. Miller, S. Clesse, F. De Lillo, G. Bruno, A. Depasse, A. Tanasijczuk, *Physics of the Dark Universe* 32 100836 (2021)
- [31] R. Abbott *et al.*(LIGO Scientific Collaboration and Virgo), Search of the early O3 LIGO data for continuous gravitational waves from the Cassiopeia A and Vela Jr. supernova remnants, *Phys. Rev. D* **105**, 082005 (2022)
- [32] R. Abbott *et al.*(LIGO Scientific Collaboration, Virgo Collaboration and KAGRA Collaboration), Searches for Continuous Gravitational Waves from Young Supernova Remnants in the Early Third Observing Run of Advanced LIGO and Virgo, *ApJ* 921 80 (2021)
- [33] R.N. Manchester, G.B. Hobbs, A. Teoh, M. Hobbs, The Australia Telescope National Facility Pulsar Catalogue, *Astron. J.*, 129, 1993-2006 (2005)
- [34] G. Woan, M. D. Pitkin, B. Haskell, D. I. Jones, and P. D. Lasky, Evidence for a Minimum Ellipticity in Millisecond Pulsars, *ApJL* **863** L40 (2018)
- [35] B.J. Owen, How to adapt broad-band gravitational-wave searches for r-modes, *Phys. Rev. D* **82** 104002 (2010)