



Research paper

Clinical high risk state of major depressive episodes: Assessment of prodromal phase, its occurrence, duration and symptom patterns by the instrument the DEpression Early Prediction-INventory (DEEP-IN)

Eva Meisenzahl^{a,*}, Natalia Wege^{a,1}, Veronika Stegmüller^a, Gerd Schulte-Körne^b, Ellen Greimel^b, Udo Dannlowski^c, Tim Hahn^c, Georg Romer^d, Marcel Romanos^e, Lorenz Deserno^{e,f,g}, Cosima Klingele^b, Christian Theisen^a, Carolin Kieckhäfer^a, Andreas Forstner^h, Stefan Ruhrmannⁱ, Frauke Schultze-Lutter^a

^a Department of Psychiatry and Psychotherapy, Medical Faculty, Heinrich-Heine University, LVR Düsseldorf, Düsseldorf, Germany

^b Department of Child and Adolescent Psychiatry, Psychosomatics and Psychotherapy, University Hospital, LMU Munich, Munich, Germany

^c Institute for Translational Psychiatry, University of Muenster, Muenster, Germany

^d Department of Child Adolescence Psychiatry and Psychotherapy, University of Muenster, Muenster, Germany

^e Centre of Mental Health, Department of Child and Adolescent Psychiatry, Psychosomatics and Psychotherapy, University of Wuerzburg, Wuerzburg, Germany

^f Max Planck Institute for Human Cognitive and Brain Sciences, Leipzig, Germany

^g Neuroimaging Center, Technical University of Dresden, Dresden, Germany

^h Institute of Human Genetics, University of Bonn, School of Medicine & University Hospital Bonn, Bonn, Germany

ⁱ Department of Psychiatry and Psychotherapy, Faculty of Medicine and University Hospital, University of Cologne, Cologne, Germany



ARTICLE INFO

Keywords:

Clinical-high-risk state of depression

Early recognition

Indicated prevention

DEEP-IN depression-early-prediction-Inventory

Prodromal phase of depression

ABSTRACT

Background: To decrease the incidence of major depressive episodes, indicated prevention that targets clinical high-risk individuals with first detectable signs that forecast mental disorder is a highly relevant topic of preventive psychiatry. Still little is known about the prodrome of MDE. The aim of the current study was to identify the occurrence of a clinical high-risk state of depression, its duration and symptom constellation.

Methods: Seventy-three patients with a diagnosed affective disorder in partial remission were assessed with our newly developed semi-structured extensive clinical instrument, the DEpression Early Prediction-INventory (DEEP-IN). Within DEEP-IN the course of prodromal symptoms was explored by using a life-chart method.

Results: The significant majority of patients (93.2 %) reported a prodromal phase. The mean duration was 7.9 months (SD = 12.5). Within the group with an identified prodromal phase, psychopathological (95.6 %) as well as somatic symptoms (88.2 %) were reported. Somatic symptoms showed a moderate-to-strong effect of sex with higher prevalence in females than in males (97.6 % vs 73.1 %; $V = 0.370$).

Limitations: This feasibility study had only a small sample size.

Conclusions: The majority of patients with affective disorders reported a clinical prodromal phase with both psychopathological and somatic symptoms that developed months before the onset of the depressive episode. The development of structured instruments for the assessment of depressive risk states is a promising approach for indicated prevention of depression in the future.

1. Introduction

Major depressive episodes (MDE) are a common and serious health condition, with 12-month prevalence rates of unipolar depression of 7.7

(6.9–8.6) and of bipolar disorders of 1.5 % (1.1–2.0) as demonstrated by Jacobi et al. (2014). Each year, over 700,000 people with depression die by suicide. In order to reduce the incidence of MDE and the associated high burden, early detection and prevention are of high priority for

* Corresponding author at: Klinik und Poliklinik für Psychiatrie und Psychotherapie, LVR-Klinikum Düsseldorf, Kliniken der Heinrich-Heine-Universität Düsseldorf, Bergische Landstraße 2, 40629 Düsseldorf, Germany.

E-mail address: eva.meisenzahl@lvr.de (E. Meisenzahl).

¹ Equally contributed.

<https://doi.org/10.1016/j.jad.2023.12.084>

Received 6 June 2023; Received in revised form 20 December 2023; Accepted 28 December 2023

Available online 4 January 2024

0165-0327/© 2024 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

global health (WHO, 2021).

Peak of the first onset of affective disorders was reported to be around 20.5 years of age (Solmi et al., 2022). Despite functional impairment occurring in the early stages of the illness, MDE is often diagnosed with a significant delay (Sheehan, 2004; Cheung et al., 2017). This delay is largely caused by the lack of knowledge about the early diagnosis of the disease (Davidson and Meltzer-Brody, 1999; Cepoiu et al., 2008). Therefore, it is crucial to understand the prodrome in order to identify clear indications for early intervention in a preventive psychiatry. Currently, there is a lack of reliable instruments that comprehensively assess various dimensions associated with the prodrome of depression, including clinical and multimodal markers.

Significant progress has been made in predicting and preventing psychosis, incl. affective psychoses, over the last decades (Worthington and Cannon, 2021). The approach of assessing clinical high-risk (CHR) states of psychosis can serve as a model for indicated prevention in psychiatry. The concept of indicated prevention involves targeting individuals who exhibit minimal but identifiable signs or symptoms of a mental, emotional, or behavioural disorder, as well as biological markers indicating a predisposition to such a disorder, even if they do not meet diagnostic criteria at the time of intervention (O'Connell et al., 2009). Important impulse for indicated prevention of mental disorders was earlier given by conceptual advancements in prevention research (Bell, 1992; Yung and McGorry, 1996) and a paradigm shift from a deterministic prevention approach in somatic medicine oriented on well-defined cause–effect-relations towards a probabilistic approach based on risk factors (Gordon Jr, 1983). Extending the concept of primary prevention, this new approach distinguishes between indicated, selective, and universal prevention, with the primary goal of reducing the occurrence of a disorder (Ruhmann et al., 2010). This approach has gradually gained recognition and acceptance in both scientific and clinical settings since the 1990s. As an adaption to mental disorders, the definition of indicated prevention was broadened to enable the development of criteria that can include clinically significant signs and early symptoms of pathological mental changes, as long as the clinical picture does not meet diagnostic criteria for the manifest disorder (Mrazek and Haggerty, 1994).

As a first necessary step before the initiation of prospective studies with psychosis high-risk-individuals and inclusion of CHR states of psychosis into the research criteria of The Diagnostic and Statistical Manual of Mental Disorders (5th ed.; DSM–5; American Psychiatric Association, 2013) and European guidelines (National Collaborating Centre for Mental Health (UK), 2014; Deutsche Gesellschaft für Psychiatrie und Psychotherapie, Psychosomatik und Nervenheilkunde, 2019), extensive retrospective research has been conducted on the prodrome and risk states of psychosis (Ebel et al., 1989; Häfner et al., 1992; Yung and McGorry, 1996; McGorry, 2002; Pantelis et al., 2003; Schultze-Lutter et al., 2007a, 2007b; Schultze-Lutter et al., 2008; Schultze-Lutter et al., 2009; Schultze-Lutter et al., 2010; Schmidt et al., 2014; Fusar-Poli et al., 2018;). Similar to psychosis, depression is supposed to arise from a complex interplay of social, psychological, and biological factors (Remes et al., 2021). While predominantly psychosis research faced challenges in operationalizing the prodromal or clinical high-risk state of the condition in the past (David, 2004; Huber, 1979; Larsen et al., 2001), the complexity and heterogeneity of depression has been used as an argument against such operationalization. Nevertheless, recent studies on early detection of psychosis and the development of CHR of psychosis criteria with the support of advanced precision medicine techniques have shown that this complexity may in opposite offer opportunities to enhance individual-level prediction (Schultze-Lutter et al., 2015; Meisenzahl et al., 2020; Koutsouleris et al., 2021a, 2021b).

In the field of MDE, the concept of early recognition, in particular the development of appropriate clinical tools and early treatment still remains poorly addressed (Benasi et al., 2021). However, the identification of individuals at risk for MDE with appropriate clinical tools underlines the opportunity for future indicated prevention strategies in

the field of affective disorders.

The still limited studies on the prodromal stage of MDE already point to a prodromal phase and a pronounced prodromal symptomatology before the onset of the overt MDE. The investigations of possible prodromal phases and their characteristics have been carried out with different methods, and it is a scientific field that is constantly evolving. The approaches of the individual research groups differ, whereby most researchers understandably pursued a retrospective approach. This retrospective approach consisted of examining patient records and/or conducting structured or semi-structured interviews, in which symptoms were queried. The period of time the interviewers investigated for a prodromal phase was chosen differently. The patient groups were composed in terms of their disease phases and frequently included patients with both unipolar and bipolar disorders (Fava and Tossani, 2007, Benasi et al., 2021).

Regarding the definition of the prodromal phase in several previous studies, it was defined as a mild form of psychiatric symptoms of MDE, or assessed by unspecific symptoms – that, potentially depending on its stage (Otto et al., 2022), more or less differ from typical depressive symptoms of manifest depression. Molnar et al. (1988) considered the prodromal phase an early marker of MDE by referring to the **time interval between the onset of first signs and symptoms of mental distress, and the onset of the characteristic manifestations of the fully developed illness**. Pede et al. (2017) and Iacoviello et al. (2010) described the prodromal phase as the **period between the occurrence of at least one symptom, which must be consistently present until the acute phase**, and manifest depression. Similarly, Sahoo et al. (2012) asked patients and relatives about any symptoms antedating the onset of the full-blown episode **fixed by six months** or less, which were regarded as prodromal symptoms of the illness episode. The different definitions of the prodromal phase are presented by Benasi et al. (2021) in his recent review indicating a distinct prodromal phase before the onset of depression. In summary, regardless of the time period chosen, a prodromal phase was identified in the vast majority of patients in the studies to date (e.g. Fava et al., 1990; Sahoo et al., 2012; Pede et al., 2017).

The **duration of the prodromal phase** of MED varied greatly between the studies, from less than a month to several years (Benasi et al., 2021). Pede et al. (2017) found a mean prodromal phase duration of 115 days (range: 20–300 days). Sahoo et al. (2012) reported a prodromal phase of 42.7 days (range: 1–150 days). One prospective study presented the duration of the prodromal phase of 44.83 days (Iacoviello et al., 2010). These differences are partly due to the above different definitions time spans and, thus, observation periods of the prodromal phase.

As regards the prodromal symptoms of MDE, the vast majority of studies have focused on psychopathological signs (Benasi et al., 2021). In their excellent review, Benasi et al. (2021) grouped the prodromal symptoms described in the studies into (1) cognitive (2) emotional (3) physical and (4) psychomotor signs. Of these, anxiety, tension, irritability and a remarkable number of physical symptoms, such as reduced energy, fatigue, sleep disturbances and somatic complaints, were the most commonly reported signs (Benasi et al., 2021). In the overall view of the studies to date, also physical complaints as early signs of MDE were already described in the first studies in the sixties (Hopkinson, 1965; Widmer and Cadoret, 1978; Cadoret et al., 1980; Wilson et al., 1983; Fava et al., 1990; Young et al., 1991; Eaton et al., 1997; Perlis et al., 1997; Iacoviello et al., 2013; Pede et al., 2017). Interestingly, a first study (Snippe et al., 2023) using Ecological Momentary Assessment showed a significant increase in repetitive negative thinking (worry, negative thoughts about the self) as the most sensitive early sign of recurrence.

In parallel to the clinical examination of the prodrome, overarching framework concepts on the development and course of psychiatric disorders are evolving. Especially a transdiagnostic staging model was introduced in the first consensus statement from an International

Working Group on Transdiagnostic Clinical Staging in Youth Mental Health by Shah et al. (2020), based on the transdiagnostic concept developed by McGorry et al. (2006). Referring to this concept, Hetrick et al. (2008) described an earlier stage for affective disorders characterised by “mild, non-specific” or “subthreshold symptoms of anxiety or depression” with a decline in the Global Assessment of Functioning (GAF) score (<70) and neurocognitive changes associated with affective disorders as indicative of the prodromal phase, while Otto et al. (2022) suggested an earlier (stage 1a) and later (stage 1b) stage of progressing symptom severity. Fava and Tossani (2007) adapted the concept of a staging model specifically for depressive disorders, conducting an overview of studies on “early stage” of depression. The “first stage” of depression defined by Fava was a prodromal phase that showed mainly symptoms like anxiety, irritability, loss of interest and sleep disturbances (Fava and Tossani, 2007).

In the study presented here, first we used a retrospective approach with a new semi-structured interview, the DEpression Early Prediction-INventory (DEEP-IN) questionnaire, in order to interview stabilised patients with affective disorders for a possible prodromal phase of MDE, its duration and its symptoms. Secondly, we aimed to examine any potential impact of sex on the duration and presentation of symptoms in the prodromal phase. Thirdly, we investigated whether specific patterns of prodromal symptoms were associated with specific sociodemographic and disease-related factors.

2. Methods

2.1. Study sample

Patients were consecutively recruited from the in- and outpatient units of the Department of Psychiatry and Psychotherapy at the Ludwig-Maximilian University in Munich according to the inclusion and exclusion criteria by the trained clinician (E.M., V.S.). Patients between 18 and 65 years-of-age and with a past or current ICD-10 diagnosis of unipolar depressive disorder (F.32) or recurrent depressive disorder (F33) or bipolar disorder (F31), who were in partial remission (BDI II <20) and had the ability to provide written informed consent, were invited to participate (Köllner and Schauenburg, 2012). Exclusion criteria were insufficient German language skills, comorbid diagnosis of an organic mental disorders (ICD-10: F0.xx), comorbid diagnosis of schizophrenia and delusional disorders (ICD-10: F2.xx), comorbid diagnosis of pervasive developmental disorders (ICD-10: F84), acute suicidality, medical history of craniocerebral trauma or unconsciousness ≥ 5 min and medical history of organic or neurological disease with impairment of brain function. Data of 73 patients were available for analyses. All patients were included in the study, after written informed consent. The study was approved by the Ethics Committee of the Ludwig-Maximilian University (EK524-15).

2.2. Assessments

The semi-structured clinical interview DEpression Early Prediction-INventory (DEEP-IN), supported by the life-chart method (Honig et al., 2001; Supplementary Fig. 1), was used to explore occurrence and frequency of the prodromal phase, its duration, signs and symptoms, and clinical course of the affective disorder. The design of DEEP-IN follows that of the Interview for the Retrospective Assessment of Onset of Schizophrenia (IRAOS) developed by Häfner et al. (1992). The comprehensive questionnaire was used to collect detailed information on the prodromal development of psychosis, its onset and early course. The duration of the interview was 90–120 min. The interview was divided into different sections, described below.

The first section of DEEP-IN contained sociodemographic information, including sex, age, marital status, education, occupation, income, nationality/migrant status, and native language.

The second section registers the complete medical history (number

and begin of MDE, use of the health care system etc.). Additionally for the validation of the diagnosis and the course of the disease episodes, the patient’s medical records and doctors’ letters were examined. Moreover, current medication as well as information on physical illnesses and compliance were recorded.

In the third section, the investigation of possible prodromal symptoms of depression was conducted supported by the life-chart method according to Lyketsos et al. (1994) that was found to perform well in depressive patients (Honig et al., 2001). In a first step, trained clinicians (E.M., V.S.) recorded the number of previous MDEs, their onset and duration. Patients were asked to represent their course of disease graphically in a time-lifeline (onset: month/year; end: month/year; Supplementary Fig. 1). The onset of the prodromal phase was defined as the time at which patients noticed first somatic or psychopathological symptoms as subjectively perceived initial impairments in their well-being. The end of the prodromal phase was defined as the time at which the diagnosis of the manifest affective disorder was made by a physician. To improve the validity of recall, the best remembered MDE by patients was identified as the ‘index episode’. In patients with bipolar disorder, the selected ‘index episode’ was required to proceed the first (hypo) manic episode.

In the fourth section, the index episode was used to identify prodromal symptoms in patients. This was done by asking patients to describe the index period in more detail, using a second graphical time-lifeline. This graph allowed to capture the temporal span before the index episode. The exploration focussed on the first individually remembered changes of wellbeing, days, weeks, months or even years before the index MED.

At the beginning of the interview, the patient was asked to speak freely instead of initially presenting a list of symptoms. After the detailed clinical and in-depth exploration by the clinician, the symptoms were documented.

2.3. Statistical analysis

First, descriptive statistics of study variables were provided, and comparisons of the course of disease according to sociodemographic variables were performed. Second, means and standard deviation (SD) of the duration of prodromal phase in months by sex, course of disease and reported index episode were calculated. Third, a frequency analysis was performed for the evaluation of the prevalence of prodromal phase and prodromal symptoms. To this aim, psychopathological and somatic symptoms were dichotomized into presence and absence of symptom.

For group comparisons, *t*-tests for normally distributed interval data with Cohen’s *d* as effect size measure, Mann-Whitney *U* test for ordinal or non-normally distributed continuous data with Rosenthal’s *r* as effect size measure, and χ^2 tests for categorical variables with Cramer’s *V* as effect size measure were calculated. Considering the small sample size, effect sizes were prioritised in the interpretation of comparative analyses because they are regarded as less effected by sample size (Sullivan and Feinn, 2012). Furthermore, because this feasibility study was only used to investigate possible fruitful targets for a subsequent larger study, because of the critiques on the *P*-value (e.g., Amrhein et al., 2019) and because of our sample size-related emphasis on effect sizes, we did not adjust for multiple testing. The statistical Package for the Social Sciences (IBM SPSS Statistics, version 28.0) was used for these data analyses.

A latent class analysis (LCA) was performed using package *pLCA* for R (Linzer and Lewis, 2011) to determine possible subgroups of reported prodromal symptoms. LCA is used to identify qualitatively different subgroups within populations (Nylund et al., 1976), and is particularly appropriate for binary data, such as the presence or absence of symptoms. Following recent criteria for the final model selection (Weller et al., 2020), the fit of various models with 2–6 classes was compared using two statistical information criteria (ICs) - Bayesian information criteria (BIC) and Akaike information criterion (AIC), in which lower values suggest better model fit (see Supplementary Table 2).

Furthermore, theoretical interpretability was considered in the choice of the best solution (Muthén and Muthén, 2000; Nylund et al., 1976).

Finally, using R statistics, a multivariate logistic regression analysis was performed to investigate the association of demographic and disease-related factors with LCA classes of symptoms. The following variables were included in the analysis simultaneously: sex, age, educational level, marital status, living alone, children, net income, duration of prodromal phase, and index episode.

3. Results

3.1. Characteristics of the study sample

Thirty-three patients (45.2 %) were inpatients and 40 patients (54.8 %) were outpatients. Twenty patients (27.4 %) reported their first MDE (ICD10 F32.x), 51 patients (69.9 %) had recurrent MDEs (ICD10 F33.x) and two patients (2.74 %) had a bipolar disorder with a current MDE (ICD 10 F31.x).

Table 1 shows the main characteristics of the study sample by disease course (first episode vs recurrent). The mean age was 39.3 years (SD = 10.45; range 20–63 years), median was 38 years (25th and 75th percentiles: 31 and 47), women were more frequent than men (63.0 % vs 37.0) (Table 1). The majority of the patients were German native speakers, single, or separated or divorced (52.0 %), childless (68.8 %), had at least twelve years of schooling (74.0 %), were currently employed (57.5 %), and had a net income of 850 to 2500 Euros per month. About half of the patients (49.3 %) lived alone. 42.2 % of all treated patients received monotherapy with antidepressants (SNRI, SSRI, NDRI). 55.6 % of patients received antidepressive therapy with an additional augmentation strategy by a second antidepressant (SNRI, SSRI, NDRI, NaSSA), or antipsychotics (Quetiapine, Olanzapine, Aripipazole) or mood stabiliser (lamotrigine, valproate, lithium). Comorbid somatic diseases were reported by 23.3 % of all in the study included patients, 2.7 % patients reported two somatic illnesses. Reported were illnesses as: thyroid diseases, chronic inflammatory diseases, allergy or asthma, musculoskeletal illnesses, rheumatic disease, diabetes, and heart valve insufficiency.

3.2. Occurrence and frequency of a prodromal phase

A prodromal phase was reported by 93.2 % of patients (n = 68). This included 90.0 % of patients with a first manifestation (18 of 20 patients) and 94.3 % of patients with a recurrent depressive or bipolar disorder (50 of 53 patients). With regard to sex, 96.3 % of males (26 of 27 male patients) and 91.3 % of females (42 of 46 female patients) presented a prodromal phase.

In the group with recurrent depression, 11.3 % had chosen the first depressive episode as the index episode, while 30.2 % reported the second episode. Most patients (58.5 %) best recalled a later depressive episode and had chosen this as the index episode.

3.3. Duration of the prodromal phase

The mean total duration of the prodromal phase of the index episode was 7.2 months (SD 12.5). There was only an indication of a small difference in duration between first-episode patients compared to recurrent disorders, with a longer duration prior to first episodes. The patient's choice of the index phase (first or later episode), however, did not indicate an effect (Table 2). Sex had the most prominent, i.e., a small-to-moderate effect on the reported duration, with longer duration of the prodromal phase in males (10.75 months vs 6.11 months with Rosenthal's $r = 0.205$), in particular in prodromes of a recurrent depressive episode (9.40 months vs 5.23 months with Rosenthal's $r = 0.212$) (Table 2). Furthermore, age was unrelated to the duration of the prodrome (Kendall's tau = -0.087).

Table 1

Descriptive characteristic of the study sample (in total and by course of affective disorder, N = 73 (%)).

Variables	Total (N = 73)	First episode (N = 20 (27.4 %))	Recurrent * (N = 53 (72.6 %))	$\chi^2(df)/F; p$
Age, mean (SD)	39.38 (10.19)	36.85 (9.32)	40.36 (10.85)	T = -1.424; p = 0.248
Sex, n (%)				
– Male	27 (37 %)	7 (25.9 %)	20 (74.1 %)	$\chi^2(1) = 0.047; p = 0.829$
– Female	46 (63 %)	13 (28.3 %)	33 (71.7 %)	
First Language				
- German	64 (87.7 %)	16 (80.0 %)	48 (90.6 %)	
- Other European	9 (12.3 %)	4 (20 %)	5 (9.4 %)	$\chi^2(1) = 1.500; p = 0.201$
Highest school education				
– Not completed school education	3 (4.1 %)	–	3 (100 %)	$\chi^2(2) = 2.012; p = 0.366$
– Lower secondary education	16 (21.9 %)	6 (37.5 %)	10 (62.5 %)	
– Higher secondary education	54 (74 %)	14 (25.9 %)	40 (74.1 %)	
Marital status				
– Married	18 (24.6 %)	7 (38.9 %)	11 (61.1 %)	$\chi^2(3) = 5.522; p = 0.137$
– Steady partnership	17 (23.3 %)	7 (41.2 %)	10 (58.8 %)	
– Single	29 (39.7 %)	5 (17.2 %)	24 (82.8 %)	
– Divorced	9 (12.3 %)	1 (11.1 %)	8 (88.9 %)	
Children				
– Yes	20 (27.4 %)	6 (30.0 %)	14 (70.0 %)	$\chi^2(1) = 0.388; p = 0.533$
– No	44 (60.3 %)	10 (22.7 %)	34 (77.3 %)	
– n.n.	9 (12.3 %)			
Living conditions				
– Living alone	26 (35.6 %)	7 (26.9 %)	19 (73.1 %)	$\chi^2(2) = 0.040; p = 0.980$
– Living with a family member	37 (50.7 %)	10 (27.0 %)	27 (73.0 %)	
– Living with other flat member	10 (13.7 %)	3 (30.0 %)	7 (70.0 %)	
Employment status				
– Employed	42 (57.5 %)	14 (33.3 %)	28 (66.7 %)	$\chi^2(7) = 5.298; p = 0.624$
– Unemployed	13 (17.8 %)	3 (15.0 %)	10 (76.9 %)	
– Student	12 (16.4 %)	3 (25 %)	9 (75.0 %)	
– Retired	6 (8.2 %)	–	6 (100 %)	
Net income in Euro				
– ≤450	8 (11.8 %)	2 (25 %)	6 (75.0 %)	$\chi^2(5) = 0.408; p = 0.790$
– 451–850	6 (8.8 %)	1 (16.7 %)	5 (83.3 %)	
– 851–1500	14 (20.6 %)	6 (42.9 %)	8 (57.1 %)	
– 1501–2500	18 (26.5 %)	4 (22.2 %)	14 (77.8 %)	
– 2501–3500	16 (23.5 %)	4 (25.0 %)	12 (75.0 %)	
– > 3500	6 (8.8 %)	2 (27.9 %)	4 (66.7 %)	
– N.n.	5 (6.8 %)			
Explored index episode				
– 1	26 (35.6 %)	20 (100 %)	6 (11.3 %)	$\chi^2(7) = 49.797; p \leq 0.001$
– 2	16 (21.9 %)	–	16 (30.2 %)	

(continued on next page)

Table 1 (continued)

Variables	Total (N = 73)	First episode N = 20 (27.4 %)	Recurrent * N = 53 (72.6 %)	$\chi^2(df)/F; p$
- 3	14 (19.2 %)	-	14 (26.4 %)	
- 4	8 (11.0 %)	-	8 (15.1 %)	
- 5	4 (5.5 %)	-	4 (7.5 %)	
- 6	3 (4.1 %)	-	3 (5.7 %)	
- 8	2 (2.7 %)	-	2 (3.8 %)	

* Includes 2 patients with a bipolar disorder with a current depressive episode (ICD 10: F31.x).

Table 2

Comparison of the duration of the prodromal phase of the index episode in months between sex, course of disorder and chosen index episode, N = 68.

	Duration (Mean \pm SD)	None-parametric statistics; effect size (r) ⁻
Sex		U = 433.00;
- Male	10.75 \pm 15.47	z = -1.451; p = 0.147; r = 0.205*
- Female	6.11 \pm 10.05	
Course of disorder		U = 390.500;
- First episode	9.58 \pm 17.26	z = -0.842; p = 0.400;
- Recurrent disorder	7.22 \pm 10.23	r = 0.102
Chosen index episode		U = 522.000;
- First episode	8.79 \pm 16.06	z = -0.078; p = 0.938;
- Recurrent episode	7.38 \pm 10.22	r = 0.009.
First episode		U = 34.500;
- Male	15.17 \pm 27.63	z = -0.142; p = 0.892;
- Female	7.58 \pm 10.53	r = 0.033
Recurrent episode		U = 226.000;
- Male	9.40 \pm 10.30	z = -1.498; p = 0.134;
- Female	5.53 \pm 9.97	r = 0.212*

U - Mann-Whitney U test; Rosenthal's r : $r \geq 0.1$ (small effect), $r = 0.3$ (moderate effect) and $r = 0.5$ and above (large effect).

* Indicates at least small to moderate effect.

3.4. Characteristics of prodromal symptoms

The prevalence of reported somatic and psychopathological symptoms in total and by sex is presented in Table 3. The most frequent prodromal symptoms were sleep disturbances, followed by fears and worries, sadness, depressed mood, exhaustion, tiredness and appetite and weight changes. Of the 68 patients reporting a prodromal phase, 60 (88.2 %) described somatic symptoms and 65 (95.6 %) psychopathological symptoms. Sex had the strongest, albeit moderate effect on somatic symptoms, with female patients being significantly more likely to report somatic symptoms. The overall frequency of psychopathological symptoms did not differ between the sexes. On item level, a small effect of sex for headache and a nearly moderate effect for emotional lability was found; both showed consistently higher frequencies in females. Regarding the type of symptom occurring first in the prodrome of the index phase, 37 patients (54.4 %) described a simultaneous occurrence of psychopathological and somatic prodromal symptoms. Only 26 patients (33.8 %) reported solely psychopathological symptoms, and eight patients (11.8 %) reported solely somatic prodromal symptoms as the first sign. A difference in the occurrence of psychopathological or somatic prodromal symptoms of only small effect size was detected in relation to the index phase (Chi-square = 1.317, $df = 2$, $p = 0.518$, Cramer's $V = 0.139$).

3.5. Latent class analysis of prodromal symptoms

All 27 prodromal symptoms listed in Table 3 were included in the

LCA. The two-class solution with the best model fit (AIC = 1426.798; BIC = 544.432) was selected (Fig. 1). Class 1 (n = 26; 38.2 %) is characterised by higher likelihood of somatic complaints with fatigue and inflammation, low energy and anhedonia. Moreover, social withdrawal and the feeling of being overwhelmed were frequently reported (Fig. 1, Supplementary Table 1). Class 2 (n = 42; 61.8 %) was characterised by higher likelihood of psychopathological complaints including rumination, tension, emotional lability and restlessness. Sleep disturbances, poor appetite, headache and musculoskeletal complaints, fears and poor concentration did not differ significantly between the two classes (Fig. 1, Supplementary Table 1).

Comparisons of sociodemographic and disease-related factors between the two LCA classes revealed that patients in Class 2 (n = 26; 38.23 %) were more likely to be male, living alone and having a recurrent disorder with small and small-to-moderate effects (Table 4).

The logistic regression analysis revealed a significant model (AIC = 103.96; McFadden = 0.050; Nagelkerke = 0.087). It indicated that living alone and financial difficulties were the most likely predictors of membership to the classes of prodromal symptoms, with patients with lower income and living alone being more likely members of class 2; yet, results were only at statistical trend level ($p < 0.1$) (Table 5). Duration of the prodromal phase and course of disease were unlikely predictors of class membership (Table 5).

4. Discussion

The aim of the current study was to identify the occurrence and frequency of an initial prodromal phase of MDE, its duration and symptom constellation with the DEEP-INventory. Further, we explored sex differences in the prodromal phase presentation, and the presence of specific symptom patterns or classes.

The significant majority of patients (93.2 %) reported a prodromal phase with mean duration of 7.9 months (SD = 12.5). Both, psychopathological as well as somatic symptoms were reported. The most common prodromal symptoms were sleep disturbances. This was followed by fears and worries, sadness, depressed mood, exhaustion, tiredness and appetite as well as weight changes. There was a moderate sex effect in the duration of the prodromal phase with male patients showing a longer prodromal phase compared to female patients. Another small-to-moderate sex effect showed in somatic symptoms that were more prevalent in female than in male patients. Furthermore, two different prodromal patterns were detected: Class 1 was characterised by higher likelihood of somatic complaints with fatigue and inflammation, low energy, anhedonia, social withdrawal and feeling of being overwhelmed. Class 2 was characterised by higher likelihood of psychopathological complaints like rumination, tension, emotional lability and restlessness.

4.1. Occurrence, frequency and duration of the prodrome of major depressive episodes

Our findings of a high prevalence of a prodromal phase in our sample is within the prevalence range reported from previous studies (Benasi et al., 2021). However, the range of frequencies is relatively wide from 26 % to 100 %, and certainly related to the various chosen time periods and definitions of the prodromal phase. The lowest prevalence rates of prodromal phases of 26 % and 29.5 % were found in a study by Hopkinson (1963) and in the retrospective evaluation of possible prodromes in depression in a long-term population-based study from Eaton et al. (1997). Hopkinson (1963) described the prodromal phase using only medical records; while Eaton et al. (1997) defined the prodromal phase as a precursor period with mild depressive symptoms. Similar our findings are the results of Fava et al. (1990) and Pede et al. (2017) with an occurrence of 100 %, who used a comprehensive clinical interview of remitted patients with major depression. Similarly, the study of Sahoo et al. (2012) investigated the prodromal phase by a

Table 3

Somatic (in white) and psychopathological symptoms (in grey) of prodromal phase (N = 68), in total and by sex n (%).

Symptom, n (%)	Total N (%)	Males	Females	χ^2 (df); p; V
Sleep disturbances	30 (44.1%)	11 (42.3%)	19 (45.2%)	$\chi^2(1)=0.056$; $p=0.813$; $V=0.029$
Fears and worries	20 (29.4 %)	10 (38.5%)	10 (23.8%)	$\chi^2(1)=1.661$; $p=0.198$; $V=0.156$
Sadness, depressed mood	19 (27.9 %)	9 (36.45%)	10 (23.8%)	$\chi^2(1)=0.931$; $p=0.335$; $V=0.117$
Exhaustion, tiredness	17 (25.0 %)	7 (26.9%)	10 (23.8%)	$\chi^2(1)=0.083$; $p=0.773$; $V=0.035$
Appetite and weight changes	15 (22.1 %)	4 (15.4%)	11 (26.2%)	$\chi^2(1)=1.091$; $p=0.296$; $V=0.127$
Lack of energy	14 (20.6 %)	5 (19.2%)	9 (21.4%)	$\chi^2(1)=0.047$; $p=0.828$; $V=0.026$
Irritability, tension	13 (19.1 %)	3 (11.5%)	10 (23.8%)	$\chi^2(1)=1.564$; $p=0.211$; $V=0.197$
Rumination	13 (19.1 %)	6 (23.1%)	7 (16.7%)	$\chi^2(1)=0.427$; $p=0.514$; $V=0.079$
Emotional lability	12 (17.6 %)	1 (3.8%)	11 (26.2%)	$\chi^2(1)=5.517$; $p=0.019$; $V=0.285'$
Somatic misperception	11 (16.2 %)	2 (7.7%)	9 (21.4%)	$\chi^2(1)=2.235$; $p=0.135$; $V=0.181$
Social withdraw	11 (16.2%)	4 (15.4%)	7 (16.7%)	$\chi^2(1)=0.019$; $p=0.889$; $V=0.017$
Headache	10 (14.7 %)	1 (3.8%)	9 (21.4%)	$\chi^2(1)=3.958$; $p=0.047$; $V=0.241'$
Concentration impairment	10 (14.7 %)	5 (19.2%)	5 (11.9%)	$\chi^2(1)=0.687$; $p=0.407$; $V=0.101$
Gastrointestinal complaints	9 (13.2 %)	2 (7.7%)	7 (16.7%)	$\chi^2(1)=1.126$; $p=0.289$; $V=0.129$
Cardiovascular complaints	8 (11.8 %)	1 (3.8%)	7 (16.7%)	$\chi^2(1)=2.543$; $p=0.111$; $V=0.193$
Nervousness, restlessness	7 (10.3 %)	2 (7.7%)	5 (11.9%)	$\chi^2(1)=0.309$; $p=0.579$; $V=0.067$
Being overwhelmed	7 (10.3 %)	4 (15.4%)	3 (7.1%)	$\chi^2(1)=1.181$; $p=0.277$; $V=0.132$
Anhedonia	7 (10.3 %)	3 (11.5%)	4 (9.5%)	$\chi^2(1)=0.071$; $p=0.790$; $V=0.032$
Musculoskeletal complaints	6 (8.8 %)	2 (7.7%)	4 (9.5%)	$\chi^2(1)=0.067$; $p=0.796$; $V=0.031$
Inflammation	5 (7.4 %)	1 (3.8%)	4 (9.5%)	$\chi^2(1)=0.760$; $p=0.383$; $V=0.106$
Increased feelings of anger	5 (7.4 %)	1 (3.8%)	4 (9.5%)	$\chi^2(1)=0.760$; $p=0.383$; $V=0.106$
Obsessive/overvalued thoughts	4 (5.9 %)	0	4 (9.5%)	$\chi^2(1)=2.631$; $p=0.105$; $V=0.197$
Low self-esteem/self-blame	4 (5.9%)	1 (3.8%)	3 (7.1%)	$\chi^2(1)=0.315$; $p=0.574$; $V=0.068$
Respiratory complaints	2 (2.9 %)	1 (3.8%)	1 (2.4%)	$\chi^2(1)=0.121$; $p=0.728$; $V=0.042$
Libido problems	2 (2.9 %)	1 (3.8%)	1 (2.4%)	$\chi^2(1)=0.121$; $p=0.728$; $V=0.042$
Jaw problems/teeth grinding	1 (1.5 %)	0	1 (2.4%)	$\chi^2(1)=0.628$; $p=0.428$; $V=0.096$
Psychopathological symptoms	65 (95.6%)	25 (96.5%)	40 (95.2%)	$\chi^2(1)=0.032$; $p=0.858$; $V=0.022$
Somatic symptoms	60 (88.2%)	19 (73.1%)	41 (97.6%)	$\chi^2(1)=9.318$; $p=0.002$; $V=0.370^*$
First prodromal symptom				
Somatic symptom	8 (11.8%)	5 (19.2%)	3 (7.1%)	$\chi^2(2)=9.576$; $p=0.008$
Psychopathological symptom	23 (33.8%)	13 (50.0%)	10 (23.8%)	$V=0.375^*$
Both somatic and psychopathological symptoms	37 (54.4%)	8 (30.8%)	29 (69.0%)	

Significance level < 0.005; 'significance level < 0.05; Interpretation of effect size: V = 0.1' – small, V = 0.3 – medium, V = 0.5-large.




Fig. 1. Prodromal symptoms - probability estimates of the two-class solution.

retrospective interview of prodromal symptoms and reported a prodrome in 93 % of the patients.

In the current study a mean duration of the prodromal phase was of 7.2 months with a range up to several years (5.9 years). We would like to note that up to now only a few studies have systematically investigated the duration of the prodromal phase and that is a methodological challenge. To this aim, we had chosen the well-established life-chart method (Häfner et al., 1992; Honig et al., 2001) that assesses the individual duration retrospectively from the depressive index episode to the time point when the patient describes substantial mental or physical changes. The existing range of the duration of the prodromal phase of depression in literature is likely due to inclusion of different diagnostic groups and differences in the definition of the prodromal phase (Benasi et al., 2021).

4.2. Clinical picture of the prodrome of major depressive episodes

Both, psychopathological and somatic symptoms were reported by patients with an identified prodromal phase. Sleep disturbances were most frequent, followed by fears and worries.

Our psychopathological prodromal profile seems in line with the findings of another study (Benasi et al., 2021). In particular, the frequent report of fears and worries in our study is in concordance with earlier reported generalized anxiety symptoms as a commonly reported prodrome (Benasi et al., 2021). For example, Fava et al. (1990) identified generalized anxiety and irritability as the most common prodromal symptoms, and Pede et al. (2017) suggested that not classical depressive symptoms but other symptoms, such as irritability and insomnia, were core components of the prodrome.

Importantly, the majority of our patients (88 %) spontaneously reported somatic prodromal symptoms. This finding underlines the

importance of the exploration of somatic symptoms in the prodromal phase because of their possible prognostic value of transition into the manifest MDE. Somatic symptoms were already proposed to be coexistent with the well-known affective, behavioural, and cognitive symptoms of depression (Kapfhammer, 2006) and may be proxy indicators for underlying pathophysiological processes (Altemus et al., 2014). Earlier, investigation of somatic symptoms before depression onset were mostly based on medical records (Cadoret et al., 1980; Wilson et al., 1983), focused on particular one or several somatic symptoms (Wilson et al., 1983; Perlis et al., 1997) or investigated the special condition of postpartum depression (Chaudron et al., 2001; Okun et al., 2009; Okun et al., 2011). The relevance of physical complaints was pointed out by a review of Kapfhammer (2006) who stated that, from a primary care perspective, this unmet diagnostic need is deplorable, as the main mode of presenting a depression is by reporting somatic symptoms. This somatic presentation, however, significantly contributes to low rates of recognition in primary care (Kapfhammer, 2006). Yet, the vast majority of the work in the area of the prodromal phase of depression has focused mostly on psychopathological signs with little attention to the exploration of heterogenic somatic early signs (Benasi et al., 2021). As our analysis supports that physical complaints can be important prodromal signs, physical complaints should be considered in scientific studies of the prodromal phase of depression in future.

4.3. Sex differences of the prodromal phase of depression

Our analyses revealed some potentially meaningful sex effects of moderate and small-to-moderate size. Firstly, male patients had a significantly longer prodromal phase, a result that should be urgently investigated in larger studies. There are no studies on this so far, so we will have to wait for further studies to carry out this comparative

Table 4
Descriptive statistic, two classes of prodromal symptoms.*

Variables	Class 1 (n = 42)	Class 2 (n = 26)	Statistics; effect size
Age in years (mean; SD)	38.83 ± 10.32	40.23 ± 9.88	W = 492.5; p = 0.503
Sex			
– Male	26 14 (53.8 %)	12 (46.2 %)	$\chi^2(1) = 0.641$, p = 0.423; V = 0.128
– Female	42 28 (66.7 %)	14 (33.3 %)	
Years of school education			
– ≥ Higher secondary education	51 32 (62.75 %)	19 (37.25 %)	$\chi^2(1) = 0.083$, p = 0.773; V = 0.035
– <Low secondary education	17 10 (58.82 %)	7 (41.18 %)	
Marital status			
– Married / in relationship	32 21 (65.63 %)	11 (34.38 %)	$\chi^2(1) = 0.135$, p = 0.713; V = 0.075
– Single/Divorced	36 21 (65.63 %)	15 (44.67 %)	
Children			
– No	49 29 (59.18 %)	20 (40.82 %)	$\chi^2(1) = 0.181$, p = 0.671; V = 0.085
– Yes	19 13 (68.42 %)	6 (31.58 %)	
Living conditions			
– Living alone	23 11 (47.83 %)	12 (51.17 %)	$\chi^2(1) = 2.037$, p = 0.153; V = 0.205
– Living with a family/ Shared apartment	45 31 (68.89 %)	14 (31.11 %)	
Employment status			
– Employed	50 30 (60.0 %)	20 (40.0 %)	$\chi^2(1) = 0.047$, p = 0.829; V = 0.060
– Unemployed	18 12 (66.67 %)	6 (33.33 %)	
Net income in categories (range 1 to 6**; mean ± SD)	63 3.84 [1.53]	3.6 [1.38]	W = 528; p = 0.461
Duration of prodromal phase	68 8.02 [13.29]	7.65 [11.35]	W = 575.5; p = 0.710
Index episode			
- First	24 17 (70.83 %)	7 (29.17 %)	$\chi^2(1) = 0.766$, p = 0.381; V = 0.138
- Recurrent	44 25 (56.82 %)	19 (43.18 %)	
Course of disorder			
- First episode	18 13 (72.22 %)	5 (27.78 %)	$\chi^2(1) = 0.611$, p = 0.434; V = 0.129
- Recurrent disorder	50 29 (58 %)	21 (42 %)	

* Includes 2 patients with a bipolar disorder with a current depressive episode (ICD 10: F31.x);

** Net income categories: 1-<=450€; 2-451-850€; 3-851-1500€; 4-1501-2500€; 5-2501-3500€; 6-> 3500 €.

analysis. It is remarkable that male patients notice and name precisely the significant changes. The male patients thus have the same awareness for intrapsychic and physical processes as female patients and experiences this prodromal state over a significantly longer period of time.

Secondly, both groups showed predominantly somatic and psychopathological symptoms, but females reported significantly more physical signs in the prodromal phase, which is an interesting result. The fact that depression is diagnosed less often in male was partly attributed to the fact that they seek medical help less often than females (Moller-Leimkuhler, 2002). Conversely, it could be speculated that the higher incidence of somatic symptoms in women might bring them into contact with doctors more often, which might enhance an earlier diagnosis.

4.4. Latent class analysis of prodromal symptoms

The result of this preliminary study remarkably shows two symptomatically different prodromal patterns. Class 1 pattern was characterised by higher likelihood of somatic complaints with fatigue and inflammation, low energy, anhedonia, social withdrawal and feeling of

Table 5
Multivariate regression analysis for two LTA classes of prodromal symptoms.

	OR [95%CI]	Estimate	Std. error	z-value	p
Age	0.97 [0.90–1.05]	−0.03	0.04	−0.72	0.471
Female (ref. male)	2.50 [0.64–10.45]	0.92	0.70	1.30	0.193
No partner (ref. partnership/ married)	1.99 [0.52–8.31]	0.69	0.70	0.98	0.325
Low education (ref. higher school education)	0.91 [0.21–3.97]	−0.09	0.73	−0.12	0.904
Unemployed (ref. employed)	1.01 [0.24–4.52]	0.019	0.73	0.02	0.987
Living alone (ref. living with partner/ family)	3.41 [0.91–14.25]	1.23	0.69	1.78	0.076*
Children (ref. no children)	1.27 [0.26–6.74]	0.24	0.81	0.29	0.768
Low net income (ref. higher income)	1.54 [0.97–2.60]	0.43	0.25	1.76	0.078*
Recurrent depression (ref. first episode)	0.90 [0.08–11.95]	−0.10	1.24	−0.08	0.932
Duration of prodromal phase	1.03 [0.98–1.10]	0.03	0.03	1.15	0.252
Index episode not the first episode (ref. first episode)	0.50 [0.04–4.65]	−0.69	1.16	−0.59	0.552

being overwhelmed while pattern of Class 2 was characterised by higher likelihood of psychopathological complaints including rumination, tension, emotional lability and restlessness.

The classification of depressive disorders is controversial (Joyce, 2008; Gaebel et al., 2020). In the current diagnostic system, the model is uniterian and the earlier binary model was dropped primarily because no clear demarcation between two types of depression (melancholic and with neurotic/reactive) could be established (Kessing, 2007). Longitudinal studies are needed to reveal, which symptom may be a precursor of possible subtypes of depression. From another perspective, the investigation of syndromal clusters already in the prodromal phase may help to identify possible later clinical subtypes. Interestingly, the predominantly physical complaints detected in our feasibility study seems to form a separate prodromal ‘entity’ in this feasibility study, and it remains to be studied, if they are indeed more related to the hypothesized early stage (1a) of the prodrome of MDE rather than to the later, more progressed stage (1b) that is assumed to consist of more depression-like symptoms and the emergence of functional decline (Otto et al., 2022).

4.5. Strength and limitations

To our knowledge, this is the first study evaluating the prodromal phase for depression and its duration by using a semi-structured comprehensive clinical interview, DEEP-IN, supported by the life-chart method and including comprehensive exploration of somatic and psychopathological symptoms of the prodromal phase and evaluating sex differences.

As a limitation, these findings of this feasibility study are based on a small study sample. Further studies with larger case numbers are needed. In addition, retrospective elicitation of remembered prodromal symptoms always have a risk of recall bias. To reduce effects of memory problems, patients could choose the episode they remembered best as the index episode of a possible prodromal phase. This approach has been widely accepted as an important starting point in research on the prodrome of psychosis and has led to important results (Häfner, 1998, Huber, 1979, Janzarik, 1987; Young et al., 1991).

Another limitation might be the inclusion of patients with bipolar disorders, as these might slightly differ in their prodrome of MDEs

(Benasi et al., 2021). Yet, bipolar disorders frequently start with MDE (McIntyre et al., 2020), and prevention of the progression of a risk state for MDE might as well prevent both uni- and bipolar affective disorders.

Further, the missing adjustment for confounding variables might be perceived as potentially introducing bias; yet, the clearly heterogeneous nature of the sample with respect to the course of the disorder and sociodemographic variables makes systematic bias seem unlikely.

Finally, for the retrospective nature of our study, we cannot completely rule out that the assessed prodromal periods partly included the very early phase of the manifest MDE, as an exact dating of the onset of the manifest MDE was not always possible. However, as the diagnosis of MDE can only be made in retrospect, i.e., after the persistence of diagnostically relevant symptoms for two weeks, a certain overlap of the prodrome or risk state, and the manifest episode within this period seems unavoidable.

4.6. Conclusions and outlook

Our results show a high prevalence of a prodromal phase of MDE with a duration of several months that, alike in psychoses, offers the opportunity to intervene early in terms of an indicated prevention. In line with psychosis research, we suggest the term ‘**clinical high-risk of depression (CHR-D)**’ for future use in prospective studies. For the first time, it was described that the duration of the prodromal phase differs between male and female patients, with a longer prodromal phase in males and, in addition, more physical prodromal symptoms in female patients. Both results should be urgently investigated in larger studies as they may affect preventive measures. Finally, future studies should examine syndromal stratifications of the prodrome detected in this investigation to capture possible subtypes of depression and/or different risk stages. The final goal is the development of valid, reliable as well as economical clinical and multimodal instruments (clinical interviews and self-rating questionnaires) for the early detection of MDE in various settings with special conceptual focus on the lifespan, including the transition phase of early adulthood as well as older age. This developmental perspective is already well advanced in the field of psychoses (Schulze-Lutter et al., 2015, Koutsouleris et al., 2021a, b) and encourages the transfer to the field of affective disorders.

Role of funding

This research received no grant from any funding agency, commercial or not-for-profit sectors.

CRediT authorship contribution statement

Eva Meisenzahl: Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Project administration, Resources, Software, Supervision, Writing – original draft, Writing – review & editing. **Natalia Wege:** Data curation, Formal analysis, Methodology, Validation, Visualization, Writing – original draft, Writing – review & editing. **Veronika Stegmüller:** Conceptualization, Data curation, Formal analysis, Methodology, Validation, Writing – original draft, Writing – review & editing. **Gerd Schulte-Körne:** Conceptualization, Methodology, Resources, Supervision, Writing – original draft, Writing – review & editing. **Ellen Greimel:** Conceptualization, Formal analysis, Methodology, Validation, Writing – original draft, Writing – review & editing. **Udo Dannlowski:** Formal analysis, Investigation, Methodology, Writing – original draft, Writing – review & editing. **Tim Hahn:** Formal analysis, Methodology, Validation, Writing – original draft, Writing – review & editing. **Georg Romer:** Methodology, Validation, Writing – original draft, Writing – review & editing. **Marcel Romanos:** Methodology, Validation, Writing – original draft, Writing – review & editing. **Lorenz Deserno:** Methodology, Validation, Writing – original draft, Writing – review & editing. **Cosima Klingele:** Formal analysis, Methodology, Validation, Writing – original draft, Writing – review & editing.

Christian Theisen: Methodology, Writing – review & editing. **Carolin Kieckhäfer:** Methodology, Validation, Writing – original draft, Writing – review & editing. **Andreas Forstner:** Conceptualization, Methodology, Writing – Original Draft, Visualization – Preparation, Writing – Review & Editing. **Stefan Ruhrmann:** Methodology, Supervision, Writing – original draft, Writing – review & editing. **Frauke Schulze-Lutter:** Conceptualization, Investigation, Methodology, Supervision, Validation, Writing – original draft, Writing – review & editing.

Declaration of competing interest

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: Eva Meisenzahl reports financial support was provided by Heinrich Heine University Düsseldorf. Eva Meisenzahl reports a relationship with Heinrich Heine University Düsseldorf that includes: employment. Board member of the journal of affective disorder.

Acknowledgments

We would like to thank the staff and students of the Clinic and Polyclinic for Psychiatry, Psychosomatics and Psychotherapy of the University Hospital of LMU Munich for their participation and collegial support in conducting this study.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.jad.2023.12.084>.

References

- Altemus, M., Sarvaiya, N., Neill Epperson, C., 2014. Sex differences in anxiety and depression clinical perspectives. *Front. Neuroendocrinol.* 35 (3), 320–330. <https://doi.org/10.1016/j.yfrne.2014.05.004>.
- American Psychiatric Association, 2013. *Diagnostic and Statistical Manual of Mental Disorders*, 5th ed. <https://doi.org/10.1176/appi.books.9780890425596>.
- Amrhein, V., Greenland, S., McShane, B., 2019. Retire statistical significance, scientists rise up against statistical significance. *Nature* 567, 305–307.
- Bell, R.Q., 1992. Multiple-risk cohorts and segmenting risk as solutions to the problem of false positives in risk for the major psychoses. *Psychiatry* 55 (4), 370–381. <https://doi.org/10.1080/00332747.1992.11024610>.
- Benasi, G., Fava, G.A., Guidi, J., 2021. Prodromal symptoms in depression: a systematic review. *Psychother. Psychosom.* 90 (6), 365–372. <https://doi.org/10.1159/000517953>.
- Cadoret, R. J., Widmer, R. B., & Troughton, E. P. (1980). Somatic complaints – harbinger of depression in primary care. *J. Affect. Disord.*, 2(1), 61–70. doi:[https://doi.org/10.1016/0165-0327\(80\)90022-1](https://doi.org/10.1016/0165-0327(80)90022-1).
- Cepoiu, M., McCusker, J., Cole, M.G., Sewitch, M., Belzile, E., Ciampi, A., 2008. Recognition of depression by non-psychiatric physicians—a systematic literature review and meta-analysis. *J. Gen. Intern. Med.* 23, 25–36.
- Chaudron, L.H., Klein, M.H., Remington, P., Palta, M., Allen, C., Essex, M.J., 2001. Predictors, prodromes and incidence of postpartum depression. *J. Psychosom. Obstet. Gynaecol.* 22 (2), 103–112. <https://doi.org/10.3109/01674820109049960>.
- Cheung, R., O'Donnell, S., Madi, N., Goldner, E., 2017. Factors associated with delayed diagnosis of mood and/or anxiety disorders. *Facteurs associés au diagnostic tardif d'un trouble de l'humeur et/ou d'anxiété*. *Health Promot. Chronic Dis. Prev. Can. : Res. Policy Pract.* 37 (5), 137–148. <https://doi.org/10.24095/hpcdp.37.5.02>.
- David, A.S., 2004. Is early intervention a waste of valuable resources? In: McDonald, C., Schultz, K., Murray, R., Wright, P. (Eds.), *Schizophrenia: Challenging the Orthodox*. Taylor & Francis, London, New York, NY.
- Davidson, J.R., Meltzer-Brody, S.E., 1999. The underrecognition and undertreatment of depression: what is the breadth and depth of the problem? *J. Clin. Psychiatry* 60 (Suppl. 7), 4–11.
- Deutsche Gesellschaft für Psychiatrie und Psychotherapie, Psychosomatik und Nervenheilkunde, 2019. S3 Leitlinie Schizophrenie. <https://www.awmf.org/leitlinien/detail/ll/038-009.html> (accessed 20.01.2023).
- Eaton, W.W., Anthony, J.C., Gallo, J., Cai, G., Tien, A., Romanoski, A., Lyketsos, C., Chen, L.S., 1997. Natural history of diagnostic interview schedule/DSM-IV major depression. The Baltimore epidemiologic catchment area follow-up. *Arch. Gen. Psychiatry* 54 (11), 993–999. <https://doi.org/10.1001/archpsyc.1997.01830230023003>.
- Ebel, H., Gross, G., Klosterkötter, J., Huber, G., 1989. Basic symptoms in schizophrenic and affective psychoses. *Psychopathology* 22 (4), 224–232. <https://doi.org/10.1159/000284602>.

- Fava, G.A., Tossani, E., 2007. Prodromal stage of major depression. *Early Interv. Psychiatry* 1 (1), 9–18. <https://doi.org/10.1111/j.1751-7893.2007.00005.x>.
- Fava, G.A., Grandi, S., Canestrari, R., Molnar, G., 1990. Prodromal symptoms in primary major depressive disorder. *J. Affect. Disord.* 19 (2), 149–152. [https://doi.org/10.1016/0165-0327\(90\)90020-9](https://doi.org/10.1016/0165-0327(90)90020-9).
- Fusar-Poli, P., De Micheli, A., Cappucciati, M., Rutigliano, G., Davies, C., Ramella-Cravaro, V., Oliver, D., Bonoldi, I., Rocchetti, M., Gavaghan, L., Patel, R., McGuire, P., 2018. Diagnostic and prognostic significance of DSM-5 attenuated psychosis syndrome in Services for Individuals at ultra high risk for psychosis. *Schizophr. Bull.* 44 (2), 264–275. <https://doi.org/10.1093/schbul/sbx055>.
- Gaebel, W., Stricker, J., Kerst, A., 2020. Changes from ICD-10 to ICD-11 and future directions in psychiatric classification. *Dialogues Clin. Neurosci.* 22 (1), 7–15.
- Gordon Jr., R.S., 1983. An operational classification of disease prevention. *Public Health Rep.* 98 (2), 107–109.
- Häfner, H., 1998. Onset and course of the first schizophrenic episode. *Kaohsiung J. Med. Sci.* 14 (7), 413–431.
- Häfner, H., Riecher-Rössler, A., Hambrecht, M., Maurer, K., Meissner, S., Schmidtke, A., Fätkenheuer, B., Löffler, W., van der Heiden, W., 1992. IRAOS: an instrument for the assessment of onset and early course of schizophrenia. *Schizophr. Res.* 6 (3), 209–223. [https://doi.org/10.1016/0920-9964\(92\)90004-o](https://doi.org/10.1016/0920-9964(92)90004-o).
- Hetrick, S.E., Parker, A.G., Hickie, I.B., Purcell, R., Yung, A.R., McGorry, P.D., 2008. Early identification and intervention in prodromal disorders: towards a clinical staging model. *Psychother. Psychosom.* 77 (5), 263–270. <https://doi.org/10.1159/000140085>.
- Honig, A., Hendriks, C.H., Akkerhuis, G.W., Nolen, W.A., 2001. Usefulness of the retrospective Life-Chart method manual in outpatients with a mood disorder: a feasibility study. *Patient Educ. Couns.* 43 (1), 43–48. [https://doi.org/10.1016/s0738-3991\(00\)00144-0](https://doi.org/10.1016/s0738-3991(00)00144-0).
- Hopkinson, G., 1963. The onset of affective illness. *Psychiatr. Neurol. (Basel)* 146 (3), 133–140.
- Hopkinson, G., 1965. The prodromal phase of the depressive psychosis. *Psychiatr. Neurol.* 149, 1–6. <https://doi.org/10.1159/000128798>.
- Huber, G., 1979. Neuere Ansätze zur Überwindung des Mythos von den sog. Geisteskrankheiten [Newer concepts to overcome the myth of so-called mental illnesses (author's transl)]. *Fortschr. Neurol. Psychiatr. Grenzgeb.* 47 (9), 449–465.
- Iacoviello, B.M., Alloy, L.B., Abramson, L.Y., Choi, J.Y., 2010. The early course of depression: a longitudinal investigation of prodromal symptoms and their relation to the symptomatic course of depressive episodes. *J. Abnorm. Psychol.* 119 (3), 459–467. <https://doi.org/10.1037/a0020114>.
- Iacoviello, B.M., Alloy, L.B., Abramson, L.Y., Choi, J.Y., Morgan, J.E., 2013. Patterns of symptom onset and remission in episodes of hopelessness depression. *Depress. Anxiety* 30 (6), 564–573. <https://doi.org/10.1002/da.22085>.
- Jacobi, F., Höfler, M., Strehle, J., Mack, S., Gerschler, A., Scholl, L., Busch, M.A., Maske, U., Hapke, U., Gaebel, W., Maier, W., Wagner, M., Zielasek, J., Wittchen, H. U., 2014. Psychische Störungen in der Allgemeinbevölkerung: Studie zur Gesundheit Erwachsener in Deutschland und ihr Zusatzmodul Psychische Gesundheit (DEGS1-MH) [Mental disorders in the general population: Study on the health of adults in Germany and the additional module mental health (DEGS1-MH)]. *Nervenarzt* 85 (1), 77–87. <https://doi.org/10.1007/s00115-013-3961-y>.
- Janzari, W., 1987. The concept of schizophrenia: history and problems. In: Häfner, H., Gattaz, W.F., Janzari, W. (Eds.), *Search for the Courses of Schizophrenia*. Springer, Berlin Heidelberg New York, pp. 11–18.
- Joyce, P.R., 2008. Classification of mood disorders in DSM-V and DSM-VI. *Aust. N. Z. J. Psychiatry* 42 (10), 851–862.
- Kapfhammer, H.P., 2006. Somatic symptoms in depression. *Dialogues Clin. Neurosci.* 8 (2), 227–239. <https://doi.org/10.31887/DCNS.2006.8.2/hpkapfhammer>.
- Kessing, L.V., 2007. Epidemiology of subtypes of depression. *Acta Psychiatr. Scand. Suppl.* 433, 85–89. <https://doi.org/10.1111/j.1600-0447.2007.00966.x>.
- Köllner, V., Schauenburg, H., 2012. *Psychotherapie Im Dialog – Diagnostik Und Evaluation*. Georg Thieme Verlag, p. 38. ISBN 978-3-13-170041-4.
- Koutsouleris, N., Dwyer, D.B., Degenhardt, F., Maj, C., Urquijo-Castro, M.F., Sanfelici, R., Popovic, D., Oeztuerk, O., Haas, S.S., Weiske, J., Ruef, A., Kambeitz-Ilanovic, L., Antonucci, L.A., Neufang, S., Schmidt-Kraepelin, C., Ruhrmann, S., Penzel, N., Kambeitz, J., Haidl, T.K., Rosen, M., Meisenzahl, E., 2021a. Multimodal machine learning workflows for prediction of psychosis in patients with clinical high-risk syndromes and recent-onset depression. *JAMA Psychiatry* 78 (2), 195–209. <https://doi.org/10.1001/jamapsychiatry.2020.3604>.
- Koutsouleris, N., Worthington, M., Dwyer, D.B., Kambeitz-Ilanovic, L., Sanfelici, R., Fusar-Poli, P., Rosen, M., Ruhrmann, S., Anticevic, A., Addington, J., Perkins, D.O., Bearden, C.E., Cornblatt, B.A., Cadenhead, K.S., Mathalon, D.H., McGlashan, T., Seidman, L., Tsuang, M., Walker, E.F., Woods, S.W., Cannon, T.D., 2021b. Toward generalizable and transdiagnostic tools for psychosis prediction: an independent validation and improvement of the NAPLS-2 risk calculator in the multisite PRONIA cohort. *Biol. Psychiatry* 90 (9), 632–642. <https://doi.org/10.1016/j.biopsych.2021.06.023>.
- Larsen, T.K., Friis, S., Haahr, U., Joa, I., Johannessen, J.O., Melle, I., Opjordsmoen, S., Simonsen, E., Vaglum, P., 2001. Early detection and intervention in first-episode schizophrenia: a critical review. *Acta Psychiatr. Scand.* 103 (5), 323–334. <https://doi.org/10.1034/j.1600-0447.2001.00131.x>.
- Linzer, D.A., Lewis, J.B., 2011. poLCA: an R package for polytomous variable latent class analysis. *J. Stat. Softw.* 42 (10), 1–29.
- Lyketso, C.G., Nestadt, G., Cwi, J., Heithoff, K., et al., 1994. The life chart interview: a standardized method to describe the course of psychopathology. *Int. J. Methods Psychiatr. Res.* 4 (3), 143–155.
- McGorry, P.D., 2002. The recognition and optimal management of early psychosis: an evidence-based reform. *World Psychiatry* 1 (2), 76–83.
- McGorry, P.D., Hickie, I.B., Yung, A.R., Pantelis, C., Jackson, H.J., 2006. Clinical staging of psychiatric disorders: a heuristic framework for choosing earlier, safer and more effective interventions. *Aust. N. Z. J. Psychiatry* 40 (8), 616–622. <https://doi.org/10.1080/j.1440-1614.2006.01860.x>.
- McIntyre, R.S., Berk, M., Brietzke, E., Goldstein, B.I., López-Jaramillo, C., Kessing, L.V., Malhi, G.S., Nierenberg, A.A., Rosenblat, J.D., Majeed, A., Vieta, E., Vinberg, M., Young, A.H., Mansur, R.B., 2020. Bipolar disorders. *Lancet* 396 (10265), 1841–1856. [https://doi.org/10.1016/S0140-6736\(20\)31544-0](https://doi.org/10.1016/S0140-6736(20)31544-0).
- Meisenzahl, E., Walger, P., Schmidt, S.J., Koutsouleris, N., Schultze-Lutter, F., 2020. Früherkennung und Prävention von Schizophrenie und anderen Psychosen [Early recognition and prevention of schizophrenia and other psychoses]. *Nervenarzt* 91 (1), 10–17. <https://doi.org/10.1007/s00115-019-00836-5>.
- Moller-Leimkuhler, A.M., 2002. Barriers to help-seeking by men: a review of sociocultural and clinical literature with particular reference to depression. *J. Affect. Disord.* 71 (1–3), 1–9. [https://doi.org/10.1016/s0165-0327\(01\)00379-2](https://doi.org/10.1016/s0165-0327(01)00379-2).
- Molnar, G., Feeney, M.G., Fava, G.A., 1988. Duration and symptoms of bipolar prodromes. *Am. J. Psychiatry* 145 (12), 1576–1578. <https://doi.org/10.1176/ajp.145.12.1576>.
- Mrazek, P.J., Haggerty, R.J., 1994. New directions in definitions. In: Mrazek, P.J., Haggerty, R.J. (Eds.), *Reducing Risks for Mental Disorders: Frontiers for Preventive Intervention Research*. National Academy Press, Washington, pp. 19–29.
- Muthén, B., Muthén, L.K., 2000. Integrating person-centered and variable-centered analyses: growth mixture modeling with latent trajectory classes. *Alcohol. Clin. Exp. Res.* 24 (6), 882–891.
- National Collaborating Centre for Mental Health (UK), 2014. *Psychosis and Schizophrenia in Adults: Treatment and Management*. National Institute for Health and Care Excellence (UK). 157–263.
- Nylund, K., Bellmore, A., Nishina, A., Winokur, G., 1976. Duration of illness prior to hospitalization (onset) in the affective disorders. *Neuropsychobiology* 2 (2–3), 87–93. <https://doi.org/10.1159/000117535>.
- O'Connell, M.E., Boat, T., Warner, K.E. (Eds.), 2009. *National Research Council (US) and Institute of Medicine (US) Committee on the Prevention of Mental Disorders and Substance Abuse Among Children, Youth, and Young Adults: Research Advances and Promising Interventions*. Preventing Mental, Emotional, and Behavioral Disorders Among Young People: Progress and Possibilities. National Academies Press (US), Washington (DC). PMID: 20662125.
- Okun, M.L., Hanusa, B.H., Hall, M., Wisner, K.L., 2009. Sleep complaints in late pregnancy and the recurrence of postpartum depression. *Behav. Sleep Med.* 7 (2), 106–117. <https://doi.org/10.1080/15402000902762394>.
- Okun, M.L., Luther, J., Prather, A.A., Perel, J.M., Wisniewski, S., Wisner, K.L., 2011. Changes in sleep quality, but not hormones predict time to postpartum depression recurrence. *J. Affect. Disord.* 130 (3), 378–384. <https://doi.org/10.1016/j.jad.2010.07.015>.
- Otto, M.W., Birk, J.L., Fitzgerald, H.E., Chauvin, G.V., Gold, A.K., Carl, J.R., 2022. Stage models for major depression: cognitive behavior therapy, mechanistic treatment targets, and the prevention of stage transition. *Clin. Psychol. Rev.* 95, 102172. <https://doi.org/10.1016/j.cpr.2022.102172>.
- Pantelis, C., Velakoulis, D., McGorry, P.D., Wood, S.J., Suckling, J., Phillips, L.J., Yung, A.R., Bullmore, E.T., Brewer, W., Soulsby, B., Desmond, P., McGuire, P.K., 2003. Neuroanatomical abnormalities before and after onset of psychosis: a cross-sectional and longitudinal MRI comparison. *Lancet (London, England)* 361 (9354), 281–288. [https://doi.org/10.1016/S0140-6736\(03\)12323-9](https://doi.org/10.1016/S0140-6736(03)12323-9).
- Pede, V.B., Jaiswal, S.V., Sawant, V.A., 2017. Study of prodromal and residual symptoms of depression. *Ind. Psychiatry J.* 26 (2), 121–127. <https://doi.org/10.4103/ipj.ipj.19.18>.
- Perlis, M.L., Giles, D.E., Buysse, D.J., Tu, X., Kupfer, D.J., 1997. Self-reported sleep disturbance as a prodromal symptom in recurrent depression. *J. Affect. Disord.* 42 (2–3), 209–212. [https://doi.org/10.1016/s0165-0327\(96\)01411-5](https://doi.org/10.1016/s0165-0327(96)01411-5).
- Remes, O., Mendes, J.F., Templeton, P., 2021. Biological, psychological, and social determinants of depression: a review of recent literature. *Brain Sci.* 11 (12), 1633. <https://doi.org/10.3390/brainsci11121633>.
- Ruhrmann, S., Schultze-Lutter, F., & Klosterkötter, J. (2010). Probably at-risk, but certainly ill—advocating the introduction of a psychosis spectrum disorder in DSM-V. *Schizophr. Res.*, 120(1–3), 23–37. doi:<https://doi.org/10.1016/j.schres.2010.03.015>.
- Sahoo, M.K., Chakrabarti, S., Kulhara, P., 2012. Detection of prodromal symptoms of relapse in mania and unipolar depression by relatives and patients. *Indian J. Med. Res.* 135 (2), 177–183.
- Schmidt, A., Smieskova, R., Simon, A., Allen, P., Fusar-Poli, P., McGuire, P.K., Bendfeldt, K., Aston, J., Lang, U.E., Walter, M., Radue, E.W., Riecher-Rössler, A., Borgwardt, S.J., 2014. Abnormal effective connectivity and psychopathological symptoms in the psychosis high-risk state. *J. Psychiatry Neurosci.* 39 (4), 239–248. <https://doi.org/10.1503/jpn.130102>.
- Schultze-Lutter, F., Ruhrmann, S., Hoyer, C., Klosterkötter, J., Leweke, F.M., 2007a. The initial prodrome of schizophrenia: different duration, different underlying deficits? *Compr. Psychiatry* 48 (5), 479–488. <https://doi.org/10.1016/j.comppsy.2007.04.001>.
- Schultze-Lutter, F., Ruhrmann, S., Picker, H., von Reventlow, H.G., Brockhaus-Dumke, A., Klosterkötter, J., 2007b. Basic symptoms in early psychotic and depressive disorders. *Br. J. Psychiatry Suppl.* 51, s31–s37. <https://doi.org/10.1192/bjp.191.51.s31>.
- Schultze-Lutter, F., Picker, H., Ruhrmann, S., Klosterkötter, J., 2008. Das Kölner Früh-Erkennungs- & Therapie-Zentrum für psychische Krisen (FETZ): evaluation der inanspruchnahme [The Cologne Early Recognition and Intervention Center for mental crises (FETZ). Evaluation of service use]. *Med Klin (Munich)* 103 (2), 81–89. <https://doi.org/10.1007/s00063-008-1012-4>. German.

- Schultze-Lutter, F., Ruhrmann, S., Klosterkötter, J., 2009. Early detection of psychosis - establishing a service for persons at risk. *Eur. Psychiatry* 24 (1), 1–10. <https://doi.org/10.1016/j.eurpsy.2008.08.004>.
- Schultze-Lutter, F., Ruhrmann, S., Berning, J., Maier, W., Klosterkötter, J., 2010. Basic symptoms and ultrahigh risk criteria: symptom development in the initial prodromal state. *Schizophr. Bull.* 36 (1), 182–191. <https://doi.org/10.1093/schbul/sbn072>.
- Schultze-Lutter, F., Michel, C., Schmidt, S.J., Schimmelmann, B.G., Maric, N.P., Salokangas, R.K., Riecher-Rössler, A., van der Gaag, M., Nordentoft, M., Raballo, A., Meneghelli, A., Marshall, M., Morrison, A., Ruhrmann, S., Klosterkötter, J., 2015. EPA guidance on the early detection of clinical high risk states of psychoses. *Eur. Psychiatry* 30 (3), 405–416. <https://doi.org/10.1016/j.eurpsy.2015.01.010>.
- Shah, J.L., Scott, J., McGorry, P.D., Cross, S.P.M., Keshavan, M.S., Nelson, B., Wood, S.J., Marwaha, S., Yung, A.R., Scott, E.M., Öngür, D., Conus, P., Henry, C., Hickie, I.B., International Working Group on Transdiagnostic Clinical Staging in Youth Mental Health, 2020. Transdiagnostic clinical staging in youth mental health: a first international consensus statement. *World Psychiatry* 19 (2), 233–242. <https://doi.org/10.1002/wps.20745>.
- Sheehan, D.V., 2004. Depression: underdiagnosed, undertreated, underappreciated. *Manag. Care Pa.*, 13(6 Suppl Depression), 6–8.
- Snippe, E., Smit, A.C., Kuppens, P., Burger, H., Ceulemans, E., 2023. Recurrence of depression can be foreseen by monitoring mental states with statistical process control. *J. Psychopathol. Clin. Sci.* 132 (2), 145–155. <https://doi.org/10.1037/abn0000812>.
- Solmi, M., Radua, J., Olivola, M., Croce, E., Soardo, L., Salazar de Pablo, G., Il Shin, J., Kirkbride, J.B., Jones, P., Kim, J.H., Kim, J.Y., Carvalho, A.F., Seeman, M.V., Correll, C.U., Fusar-Poli, P., 2022. Age at onset of mental disorders worldwide: large-scale meta-analysis of 192 epidemiological studies. *Mol. Psychiatry* 27 (1), 281–295. <https://doi.org/10.1038/s41380-021-01161-7>.
- Sullivan, G.M., Feinn, R., 2012. Using effect size-or why the P value is not enough. *J. Grad. Med. Educ.* 4 (3), 279–282. <https://doi.org/10.4300/JGME-D-12-00156.1>.
- Weller, B.E., Bowen, N.K., Faubert, S.J., 2020. Latent class analysis: A guide to best practice. *J. Black Psychol.* 46 (4), 287–311. <https://doi.org/10.1177/0095798420930932>.
- Widmer, R.B., Cadoret, R.J., 1978. Depression in primary care: changes in pattern of patient visits and complaints during a developing depression. *J. Fam. Pract.* 7 (2), 293–302.
- Wilson, D.R., Widmer, R.B., Cadoret, R.J., Judiesch, K., 1983. Somatic symptoms. A major feature of depression in a family practice. *J. Affect. Disord.* 5 (3), 199–207. [https://doi.org/10.1016/0165-0327\(83\)90042-3](https://doi.org/10.1016/0165-0327(83)90042-3).
- World Health Organization, 13 September 2021. <https://www.who.int/news-room/fact-sheets/detail/depression> (accessed 22 December 2022).
- Worthington, M.A., Cannon, T.D., 2021. Prediction and prevention in the clinical high-risk for psychosis paradigm: a review of the current status and recommendations for future directions of inquiry. *Front. Psychol.* 12, 770774 <https://doi.org/10.3389/fpsy.2021.770774>.
- Young, M.A., Watel, L.G., Lahmeyer, H.W., Eastman, C.I., 1991. The temporal onset of individual symptoms in winter depression: differentiating underlying mechanisms. *J. Affect. Disord.* 22 (4), 191–197. [https://doi.org/10.1016/0165-0327\(91\)90065-z](https://doi.org/10.1016/0165-0327(91)90065-z).
- Yung, A.R., McGorry, P.D., 1996. The initial prodrome in psychosis: descriptive and qualitative aspects. *Aust. N. Z. J. Psychiatry* 30 (5), 587–599. <https://doi.org/10.3109/00048679609062654>.