

A phenomenological gravitational waveform model of binary black holes incorporating horizon fluxes

Samanwaya Mukherjee^{1,*}, Sayak Datta^{2,3,†}, Khun Sang Phukon⁴ and Sukanta Bose⁵

¹*Inter-University Centre for Astronomy and Astrophysics, Post Bag 4, Ganeshkhind, Pune - 411007, India*

²*Max-Planck-Institut für Gravitationsphysik (Albert-Einstein-Institut), D-30167 Hannover, Germany*

³*Leibniz Universität Hannover, D-30167 Hannover, Germany*

⁴*School of Physics and Astronomy and Institute for Gravitational Wave Astronomy, University of Birmingham, Edgbaston, Birmingham, B15 9TT, United Kingdom*

⁵*Department of Physics and Astronomy, Washington State University, 1245 Webster, Pullman, Washington 99164-2814, U.S.A.*

Subjected to the tidal field of its companion, each component of a coalescing binary suffers a slow change in its mass (tidal heating) and spin (tidal torquing) during the inspiral and merger. For black holes, these changes are associated with their absorption of energy and angular momentum fluxes. This effect modifies the inspiral rate of the binary, and consequently, the phase and amplitude of its gravitational waveform. Numerical relativity waveforms contain these effects inherently, whereas analytical approximants for the early inspiral phase have to include them manually in the energy balance equation. In this work, we construct a frequency-domain gravitational waveform model that incorporates the effects of tidal heating of black holes. This is achieved by recalibrating the inspiral phase of the waveform model `IMRPhenomD` to incorporate the phase corrections for tidal heating. We also include corrections to the amplitude, but add them directly to the inspiral amplitude model of `IMRPhenomD`. We show that the new model is faithful, with less than 1% mismatch, against a set of hybrid waveforms, except for one outlier that barely breaches this limit. The recalibrated model shows mismatches of up to $\sim 16\%$ with `IMRPhenomD` for high mass ratios and spins. Amplitude corrections become less significant for higher mass ratios, whereas the phase corrections leave more impact – suggesting that the former is practically irrelevant for gravitational wave data analysis in Advanced LIGO (aLIGO, Virgo and KAGRA). Comparing with a set of 219 numerical relativity waveforms, we find that the median of mismatches decreases by $\sim 4\%$ in aLIGO zero-detuned high power noise curve, and by $\sim 2\%$ with a flat noise curve. This implies a modest but notable improvement in waveform accuracy.

I. INTRODUCTION

The launch of gravitational wave (GW) astronomy has had a stellar start, with the detection of over 90 compact binary coalescences (CBCs) so far [1–3]. The fourth (O4) observation run of the ground-based GW detectors LIGO [4], Virgo [5] and KAGRA [6] are expected to detect many more CBCs, which enables us to subject General Relativity (GR) to unprecedented tests. Such tests demand high precision CBC waveforms. In the field of GW data analysis, waveform models serve as templates against which the real data are compared and estimates of the source parameters are made by the matched filtering technique [7] and Bayesian inference [8–10]. Since a sizable subset of CBCs are high-mass binaries, which have a good fraction of their signal power arriving from the late-inspiral and merger parts, these tests benefit from employing inspiral-merger-ringdown (IMR) waveforms. One needs to solve the full Einstein equations numerically to extract gravitational waveforms predicted by GR to meet the accuracy standard imperative for describing the merger-ringdown parts of a CBC. However, such simulations of numerical relativity (NR) are

of extreme computational cost, impeding the usage of this scheme for creating long waveforms that span the entire frequency range of GW detectors for intermediate to stellar-mass binaries. On the other hand, post-Newtonian (PN) approximation (see, e.g., Ref. [11] and the references therein) or the effective-one-body (EOB) framework [12–14] describes the inspiral part in analytical forms, but fail to be reliably accurate in the late inspiral to the merger regime.

Efforts to construct accurate IMR models have been based on the EOB formalism with calibrations to NR data (SEOBNR [13, 15, 16] and TEOBResumS [17–19] models), phenomenological waveform models built by combining PN and NR waveforms [20–25], and more recently, surrogate models [26–28]. The computational cost of EOB models, however, is burdened by the need to tackle the orbital dynamics through solving a complex system of ordinary differential equations. Phenomenological models, in comparison, are equipped with closed-form expressions for the phase and amplitude of the waveforms for a given set of binary parameters, and are much faster to evaluate. The latter models contain PN-inspired phase and amplitude behaviors augmented with higher-order terms calibrated against a set of “hybrid waveforms”. The hybrids are constructed by stitching either PN (e.g., in `IMRPhenomC` [21]) or EOB (e.g. `IMRPhenomD` [24, 25], `IMRPhenomXAS` [23] for aligned-spin cases) waveforms with NR ones at a suitable frequency,

* samanwaya@iucaa.in

† sayak.datta@aei.mpg.de

to allow smooth transitions of phase and amplitude from the inspiral to the merger-ringdown regime. Utmost care needs to be borne when producing complete IMR waveforms by combining these parts, to limit introducing waveform systematics, which can reduce the effectiveness of tests of GR. The increasing modeling errors and computational cost of gravitational waveforms with increasing mass ratio and spins [29] have inspired the advent of reduced-order models (ROMs) and surrogate models. By fitting interpolated decomposed waveform data pieces over the binary parameter space, surrogate models can significantly accelerate NR (e.g. [28, 30, 31]) or EOB waveforms (e.g. [32–34]), while maintaining high accuracy within their parameter space of validity.

In phenomenological models for binary black holes (BBHs), the inspiral waveform is usually constructed with the “point-particle” approximation, which also acts as a baseline for creating generic waveforms with tidal effects arising from the finite size of the components, e.g. for binary neutron stars (BNSs) [35, 36]. However, even for black holes (BHs) there exists a finite-size effect that needs to be consistently incorporated within a complete BBH waveform model, namely, the absorption of energy and angular momentum by their horizons. Due to their causal structure, BHs in GR are perfect absorbers that behave as dissipative systems [37, 38]. A significant feature of a BH is its horizon, which is a null surface and a “one-way membrane” that does not allow energy to escape outward. In a binary, a BH experiences tidal interactions from its companion, which cause changes in its mass, angular momentum, and horizon area. This phenomenon is called tidal heating (TH) [39–43]. This effect is weak but important in many respects. Recent studies [44, 45] have found that the presence of TH can be measurable, especially in the future era of GW detectors with orders of magnitude higher signal-to-noise ratios (SNRs) compared to the current ones. Combined with the fact that TH is much more significant for BHs than horizonless compact objects, this makes TH a viable discriminator for horizons to identify BHs against theoretically possible exotic compact objects (ECOs) [46], which can mimic BHs in their GW signals [47, 48].

In NR simulations, TH arises naturally due to the presence of the BH apparent horizons. While evolving the binary, BH singularities are expunged from the computational domain by excision techniques or by the “moving puncture” method. The properties of the apparent horizons – the masses, spins, and the horizon areas – continue to get impacted by the energy and angular momentum fluxes throughout the binary evolution up to the merger. Changes in BH masses and spins impact the inspiral rate of the binary, leaving its imprint in the phase of GW signals. Complete IMR waveforms created with NR data for late inspiral and merger, then, would have incomplete information about BBH systems if they are joined with inspiral approximants devoid of the imprints of horizon fluxes. Ignoring this effect may also lead to ambiguities in the tests of GR from GW observations of CBCs, where

small deviations from GR predictions are probed under the strong gravity conditions of a CBC. Lesser systematics will facilitate a more accurate evaluation of GR’s validity in such cases.

In the EOB formalism, progress has been made to include the effects of horizon absorption for nonspinning binaries [49–51] and for aligned spins [52]. These works use numerical evaluation of horizon fluxes using a frequency-domain perturbative approach along stable and unstable orbits in the test-mass limit. This test-mass knowledge is then hybridized with lower-order analytical information that is valid for comparable masses. These calculations were implemented in the construction of `TEOBResumS` [17], an NR-informed EOB waveform model which includes the horizon-absorption effects in a PN-inspired resummed form. Alvi [41] has calculated, under PN approximation, leading-order expressions of mass and spin evolution of BBH systems. Alvi’s expressions are explicitly valid for any mass ratio. A resummed version of these expressions was used by Damour and Nagar [52] to describe them under the EOB framework.

In Phenomenological models, horizon absorption was taken into account in the flux calculations of `IMRPhenomC` [21], up to 2.5PN order. However, later models with more accurate calibrations spanning larger regions of the parameter space (e.g. `IMRPhenomD` or `IMRPhenomXAS`) do not account for this effect explicitly in their inspiral parts.

In this work, we construct a phenomenological BBH waveform model with a consistent inclusion of TH effects, and study its contribution to an improvement in waveform systematics. A complete BBH model, which includes the phase and amplitude modifications for TH from the early inspiral to the merger, not only provides a comprehensive foundation but also helps to lay the groundwork for a more general waveform model with the *horizon parameters*, which can act as discriminators for BHs from ECOs or NSs [44, 45].

We first describe some preliminaries and notations of gravitational waveforms in Sec. II. Then in Sec. III A, we build an inspiral model containing explicit corrections due to TH at 2.5PN, 3.5PN, and 4PN orders in their phase and amplitude. The point-particle baseline for the inspiral waveform is based on an aligned-spin EOB approximant. In Sec. III B, we describe the publicly available NR waveforms of the SXS catalog [53] used for merger and ringdown phases. In Sec. IV, we describe the construction of new hybrid waveforms by stitching together the TH-corrected inspiral and merger-ringdown parts. Following `IMRPhenomD` (henceforth referred to as `PhenomD`), we model the hybrids in the frequency domain in Sec. V and calibrate the inspiral part with augmented PN expressions, but against the new set of hybrid waveforms. Section VI discusses the faithfulness of the new model, and in Sec. VII, we compare it with `PhenomD` and NR data within the parameter range of calibration. We conclude in Sec. VIII with discussions and future directions of this work.

Throughout the article, we will use geometric units by considering the gravitational constant (G) and the speed of light in vacuum (c) set to unity, except while calculating physical quantities.

II. WAVEFORM CONVENTIONS AND NOTATION

We consider $l = |m| = 2$ spherical harmonic modes of gravitational waves from coalescing binary black holes. In the time domain, the complex GW strain can be written as,

$$h_{2,2}(\Theta, t) = A(\Theta, t)e^{-i\phi(\Theta, t)}, \quad (1)$$

where Θ is the parameter vector containing the parameters of the binary. The intrinsic parameters are the dimensionless spin vectors χ_1, χ_2 and the two masses m_1 and m_2 . Since we are concerned with non-precessing binaries only, the spin vectors have nonzero components only along (or opposite to) the direction of the orbital angular momentum. In terms of the normalized orbital angular momentum \hat{L} and the dimensionful spin vectors \mathbf{S}_i ($i = 1, 2$ for the two objects), the components of dimensionless spins can be expressed as

$$\chi_i = \frac{\mathbf{S}_i \cdot \hat{L}}{m_i^2}. \quad (2)$$

We define the mass ratio as $q = m_1/m_2 \geq 1$, total mass $M = m_1 + m_2$, and the symmetric mass ratio $\eta = m_1 m_2 / M^2$.

The complex strain in Eqn. (1) can be constructed from the plus and cross polarization states as

$$h(\Theta, t) = h_+(\Theta, t) - ih_\times(\Theta, t). \quad (3)$$

The Fourier transform of the complex strain is defined as

$$\tilde{h}(f) = \int_{-\infty}^{\infty} h(t)e^{-i2\pi ft} dt, \quad (4)$$

where f denotes frequency. For the non-eccentric waveforms that we use, f can be written as a monotonically increasing function of time t until the merger.

In gravitational waveform modeling, the dimensionless frequency Mf plays an important role, since the total mass M acts as a scaling factor. For better readability, we define $F \equiv Mf$.

III. INPUT WAVEFORMS

A. Inspiral waveforms

1. Effective-one-body description of a compact binary coalescence

In the effective-one-body approach to the two-body problem in GR, originally developed by Buonanno and

Damour [12], the dynamics of two compact objects of masses m_1 and m_2 and spins χ_1 and χ_2 is mapped onto the dynamics of an effective particle of mass $\mu = m_1 m_2 / (m_1 + m_2)$ and spin χ_* moving in the background of a deformed Kerr geometry with mass $M = m_1 + m_2$ and spin χ_{Kerr} . The spin mapping $\{\chi_1, \chi_2\} \rightarrow \chi_*$ and the deformation of the Kerr background, parametrized by the symmetric mass ratio η , implies that the test-particle dynamics reproduces the PN-expanded dynamics of the original two-body system. Free parameters are introduced into the models that represent unknown, higher-order PN terms, or additional physical effects like corrections due to eccentricity. Such free parameters are calibrated to NR simulations. With the EOB system specified, its conservative dynamics can be described by an EOB Hamiltonian [15, 16], while the non-conservative dynamics is contained in a parametrized radiation-reaction term that is inserted in the equations of motion. This term sums over the outgoing GW modes and is calibrated to reproduce NR simulations. The combination of these two pieces describes the binary inspiral through to the merger, at which point a ringdown waveform is attached to the inspiral-merger waveform. The ringdown waveform is constructed as a linear superposition of the dominant quasinormal modes (QNMs) of the Kerr BH formed at merger [52, 54], with the amplitude and phase of each QNM mode determined by the process of stitching the ringdown part with the inspiral-merger parts.

For the purpose of creating hybrid waveforms in our work, we choose SEOBv2 [16], an aligned-spin EOB approximant, as the point-particle baseline. More recent versions of the SEOBv2 family of waveforms are available [13] at present, but we create our model based on the former one for a direct comparison with PhenomD. We also mention here that for PhenomD, NR calibrations of SEOBv2 were removed before creating hybrid waveforms, referred to as SEOBv2. In this work, however, we use the original calibrated version of the EOB model since in the hybrids, stitched at a frequency much lower than the innermost stable circular orbit (ISCO), we expect little to no NR information from the inspiral approximant to be present.

2. Corrections for tidal heating of black holes

The early inspiral part of a CBC can be described by the analytical formalism of post-Newtonian (PN) expansions [11], under the approximation that the source is slowly moving and weakly gravitating. In this formalism, the evolution of the orbital phase $\Psi(t)$ of a compact binary is computed as a perturbative expansion in a small parameter, typically taken to be the characteristic velocity $v = (\pi M f)^{1/3}$. This analytical procedure demands $v \ll 1$, which makes it useful in the early inspiral phase of a CBC.

An electrically neutral spinning black hole in GR –

the Kerr black hole (KBH) – is stationary when it is isolated. On the other hand, when a KBH is a member of a binary, it feels its companion’s tidal field, which acts as a non-axisymmetric perturbation [39]. This perturbation causes changes in the mass, spin, and horizon area of the KBH over time [41]. Since the KBH experiences the tidal field of its orbiting companion, it absorbs (emits) energy from (into) the orbit. The absorption part is present in non-spinning BHs as well. Additionally, for a KBH, a spin frequency higher than the angular frequency of the tidal field slows down due to tidal interactions, which in turn makes the KBH lose its rotational energy. The slowing down of a rotating BH due to the gravitational dissipation produced by exterior mass is analogous to the slowing down of a rotating planet by viscous dissipation due to tides raised by an exterior moon that increases its internal thermal content - a phenomenon known as tidal heating. Due to this similarity, the energy and angular momentum flux in BBHs is also termed tidal heating [40].

During the early stages of the binary evolution, the change in the orbital frequency due to emission of GWs is much smaller than the orbital frequency itself. In this *adiabatic inspiral* regime, loss of binding energy $E(v)$ of the two-body system with time equals the GW flux emitted to future null infinity ($\mathcal{F}^\infty(v)$) plus the energy flux absorbed by the two BH horizons ($\mathcal{F}^H(v)$). So the energy balance condition becomes

$$-\frac{dE(v)}{dt} = \mathcal{F}^\infty(v) + \mathcal{F}^H(v). \quad (5)$$

Evolution of the orbital phase ϕ and the characteristic velocity v , obtained from this equation, read

$$\frac{d\phi}{dt} = \frac{v^3}{M}, \quad \frac{dv}{dt} = -\frac{\mathcal{F}(v)}{E'(v)}, \quad (6)$$

where $\mathcal{F}(v) = \mathcal{F}^\infty(v) + \mathcal{F}^H(v)$. These equations yield a solution for the phase $\Psi(f)$ of the frequency-domain waveform $\tilde{h}(f) = \tilde{A}(f)e^{-i\Psi(f)}$ [55]:

$$\Psi(f) = 2(t_c/M)v^3 - 2\phi_c - \pi/4 - \frac{2}{M} \int (v^3 - \bar{v}^3) \frac{E'(\bar{v})}{\mathcal{F}(\bar{v})} d\bar{v}, \quad (7)$$

where $E'(v) = dE(v)/dv$.

The frequency-domain amplitude $\tilde{A}(f)$ can be written as¹ [56]

$$\tilde{A}(f) = \mathcal{C} \frac{2\eta M^{3/2}}{d_L} v \sqrt{\frac{\pi}{3}} \left[-\frac{E'(v)}{\mathcal{F}(v)} \right]^{1/2}, \quad (8)$$

where \mathcal{C} is a constant that depends on the inclination of the orbital plane with the line of sight, and d_L denotes the distance to the binary.

Splitting \mathcal{F} into \mathcal{F}^∞ and \mathcal{F}^H in Eq. (7) enables one to write the frequency-domain phase in the form

$$\Psi(f) = \Psi_{\text{PP}}(f) + \Psi_{\text{TH}}(f), \quad (9)$$

with $\Psi_{\text{PP}}(f)$ being the GW phase under point particle (PP) approximation that corresponds to \mathcal{F}^∞ , and $\Psi_{\text{TH}}(f)$ being the phase correction due to TH. The absorbed flux can be expressed as the sum of the rates of mass increment [41] for the two BHs:

$$\begin{aligned} \mathcal{F}^H(v) &= \frac{32}{5} \eta^2 \frac{v^{15}}{4} \sum_{i=1}^2 \left(\frac{m_i}{M} \right)^3 (1 + 3\chi_i^2) \\ &\times \left\{ -(\hat{L} \cdot \hat{S}_i) \chi_i + 2 \left[1 + \sqrt{1 - \chi_i^2} \right] \frac{m_i v^3}{M} \right\}, \end{aligned} \quad (10)$$

where the indices $i = 1, 2$ denote the two BHs, and \hat{S}_i and \hat{L} are the unit vectors along the directions of the i th object’s spin and the orbital angular momentum, respectively.

Using this expression, one can calculate the phase contribution of tidal heating in gravitational waveforms of BBHs to be [44, 45]

$$\begin{aligned} \Psi_{\text{TH}} &= \frac{3}{128\eta} \left[-\frac{10}{3} H_{\text{eff5}}^{(\text{BBH})} \log(v) \right. \\ &- \frac{5}{168} v^2 H_{\text{eff5}}^{(\text{BBH})} (952\eta + 995) \\ &\left. + \frac{5}{3} v^3 \log(v) (-4H_{\text{eff8}}^{(\text{BBH})} + H_{\text{eff5}}^{(\text{BBH})} \Psi_{\text{SO}}) \right], \end{aligned} \quad (11)$$

where the quantities $H_{\text{eff5}}^{(\text{BBH})}$ and $H_{\text{eff8}}^{(\text{BBH})}$ are defined as

$$H_{\text{eff5}}^{(\text{BBH})} \equiv \sum_{i=1}^2 \left(\frac{m_i}{M} \right)^3 (\hat{L} \cdot \hat{S}_i) \chi_i (3\chi_i^2 + 1), \quad (12a)$$

$$\begin{aligned} H_{\text{eff8}}^{(\text{BBH})} &\equiv 4\pi H_{\text{eff5}}^{(\text{BBH})} + \sum_{i=1}^2 \left(\frac{m_i}{M} \right)^4 (3\chi_i^2 + 1) \\ &\times \left(\sqrt{1 - \chi_i^2} + 1 \right). \end{aligned} \quad (12b)$$

Ψ_{SO} is a “spin-orbit” term, given by

$$\begin{aligned} \Psi_{\text{SO}} &= \frac{73}{3(1+q)^2} \left\{ q^2 (\hat{L} \cdot \hat{S}_1) \chi_1 + (\hat{L} \cdot \hat{S}_2) \chi_2 \right\} \\ &+ \frac{15q}{(1+q)^2} \left\{ (\hat{L} \cdot \hat{S}_1) \chi_1 + (\hat{L} \cdot \hat{S}_2) \chi_2 \right\}. \end{aligned} \quad (13)$$

We find the PN expansion of the total amplitude by expanding $[-E'(v)/\mathcal{F}(v)]^{1/2}$ in Eq. (8) in powers of v up to v^7 , and separate the contribution due to TH:

$$\tilde{A}(f) = \tilde{A}_{\text{PP}}(f) + \tilde{A}_{\text{TH}}(f). \quad (14)$$

¹ Note that we define $E(v)$ as the total binding energy. Ref [56] treats $E(v)$ as the specific binding energy (binding energy per unit mass), so the power of M in our treatment is different from theirs in this particular equation.

$\tilde{A}_{\text{PP}}(f)$ is mentioned in Eq. (5.7) of Ref. [56], and we get the expression for $\tilde{A}_{\text{TH}}(f)$ as

$$\begin{aligned} \tilde{A}_{\text{TH}}(f) = & \mathcal{C} \frac{M^{5/6}}{d_L \pi^{2/3}} \sqrt{\frac{5\eta}{24}} f^{-7/6} \left[\frac{1}{8} H_{\text{eff5}}^{(\text{BBH})} v^5 \right. \\ & + \left(\frac{1079}{1792} + \frac{103}{192} \eta \right) H_{\text{eff5}}^{(\text{BBH})} v^7 \\ & + \frac{1}{32} \left\{ \tilde{A}_{\text{SO}} H_{\text{eff5}}^{(\text{BBH})} \right. \\ & \left. \left. - 8 \left(H_{\text{eff8}}^{(\text{BBH})} - \pi H_{\text{eff5}}^{(\text{BBH})} \right) \right\} v^8 \right], \end{aligned} \quad (15)$$

where \tilde{A}_{SO} is the spin-orbit term

$$\begin{aligned} \tilde{A}_{\text{SO}} = & \frac{179}{6(1+q)^2} \left\{ q^2 (\hat{L} \cdot \hat{S}_1) \chi_1 + (\hat{L} \cdot \hat{S}_2) \chi_2 \right\} \\ & + \frac{35q}{2(1+q)^2} \left\{ (\hat{L} \cdot \hat{S}_1) \chi_1 + (\hat{L} \cdot \hat{S}_2) \chi_2 \right\}. \end{aligned} \quad (16)$$

B. Numerical relativity waveforms for merger-ringdown

As a BBH progresses towards the merger phase, the BHs come so close to each other that the system becomes too compact for the weak-gravity condition to hold. In addition, their orbital velocities culminate to values comparable to c , breaching the slow-motion approximation. The analytical PN results, with their highest order of expansion available currently, become increasingly inaccurate to describe the system evolving under such extreme conditions. Starting from this *late inspiral* phase to the merger and ringdown phase where the two BHs merge and the final remnant BH settles to equilibrium, one needs to solve the full Einstein equations numerically for extracting GWs to conform to the accuracy standards of the waveforms required for searches of GW signals or their parameter estimation (PE). This is the field of NR, which has seen exciting breakthroughs [57–59] in the early 2000s, and great strides towards simulating merging black holes with different masses and spins [60–63].

We use NR data from the publicly available SXS catalog [53], computed using the Spectral Einstein Code (SpEC) [64, 65]. SpEC uses the excision techniques to remove the BH singularity for extracting gravitational waves from binary black hole systems. Unlike the PN framework where the fluxes of energy and angular momentum absorbed by the BHs have to be considered explicitly in the energy balance equation, numerical simulations of full Einstein equations capture this effect inherently. Scheel *et al.* [66] have demonstrated the slow change in the BH masses and spins through the binaries' temporal evolution in NR simulations with SpEC. They track the apparent horizons as a function of time, and at frequent time intervals, they measure both the surface area and the spin of the horizons. The spin computation has been carried out using the approximate Killing vector formalism [67]. The mass of the black hole is then

computed using Christodoulou's formula [68],

$$M^2 = M_{\text{irr}}^2 + \frac{S^2}{4M_{\text{irr}}^2}, \quad (17)$$

where M_{irr} is the irreducible mass of the Kerr BH,

$$M_{\text{irr}}^2 = \frac{1}{16\pi} \int_{\mathcal{H}} dA. \quad (18)$$

Here S is the (dimensionful) spin of the KBH and the integration covers the total horizon area A .

During the evolution, each SXS NR waveform is extracted at a series of times on a set of concentric co-ordinate spheres surrounding the binary, decomposed in modes of spin-weighted spherical harmonic functions. Then the waveforms are extrapolated to future null infinity \mathcal{I}^+ . The dataset of each SXS waveform contains several gravitational waveform modes (l, m) , and the orders of extrapolation $N = 2, 3, 4$. A higher order of extrapolation is preferred when accurate waveforms are needed in the inspiral, and lower order extrapolation is preferred for more accuracy at the merger-ringdown phase [53]. Since we use these waveforms for constructing hybrids, we need more accuracy for the merger-ringdown part, due to the fact that some part of the inspiral has to be replaced by the PN waveforms in the hybrids. We use 20 aligned-spin non-eccentric BBH waveforms from SXS with $1 \leq q \leq 8$ and $-0.95 \leq \chi_1, \chi_2 \leq 0.85$ for calibrating our model. We choose (2, 2) modes, and the extrapolation order $N = 2$ for our purpose.

IV. HYBRID CONSTRUCTION

For constructing a hybrid, the NR waveform has to be 'stitched' with the analytical inspiral waveform in a frequency region where both the waveforms have sufficient accuracy, which also demands that the morphology of these waveforms do not deviate significantly from one another. For PN waveforms, one conventionally chooses ISCO of the binary to be the endmost point of validity. The corresponding GW frequency f_{ISCO} is considered to be the highest frequency for PN expansions. In our work, we ensured that the frequency at the stitching region is below this threshold for all the hybrids - which also ensures that the NR calibrations in SEOBNRv2 are minimally present before the stitching starts.

While stitching at a frequency as low as possible (considering ISCO) should enable one to incorporate the maximum number of NR cycles into the hybrid, there are two more factors that affect the choice of the stitching region:

- The initial NR data do not perfectly describe two black holes in quasi-equilibrium. At the start of each simulation, the geometry relaxes to equilibrium on the dynamical time-scale of the individual BHs, changing the mass and spin of each BH by a fractional amount of order 10^{-5} , and emitting a

#	Simulation label	q	χ_1	χ_2	χ_{PN}	N_{orb}	F_{stitch}
1	SXS:BBH:0156	1	-0.95	-0.95	-0.79	13	0.0057
2	SXS:BBH:0151	1	-0.6	-0.6	-0.5	15	0.0054
3	SXS:BBH:0001	1	0	0	0	28	0.0041
4	SXS:BBH:0152	1	0.6	0.6	0.5	23	0.0052
5	SXS:BBH:0153	1	0.85	0.85	0.7	25	0.0052
6	SXS:BBH:0234	2	-0.85	-0.85	-0.72	28	0.0037
7	SXS:BBH:0238	2	-0.5	-0.5	-0.43	32	0.0037
8	SXS:BBH:0169	2	0	0	0	16	0.0062
9	SXS:BBH:0253	2	0.5	0.5	0.43	29	0.0046
10	SXS:BBH:2131	2	0.85	0.85	0.72	25	0.0056
11	SXS:BBH:1936	4	-0.8	-0.8	-0.71	17	0.0057
12	SXS:BBH:1418	4	-0.4	-0.5	-0.37	67	0.005
13	SXS:BBH:0167	4	0	0	0	16	0.0071
14	SXS:BBH:1417	4	0.4	0.5	0.37	80	0.0057
15	SXS:BBH:1907	4	0	0.8	0.12	21	0.0062
16	SXS:BBH:1423	8	-0.6	-0.75	-0.57	18	0.0066
17	SXS:BBH:0064	8	-0.5	0	-0.43	19	0.0066
18	SXS:BBH:0063	8	0	0	0	26	0.0065
19	SXS:BBH:0065	8	0.5	0	0.43	34	0.0063
20	SXS:BBH:1426	8	0.48	0.75	0.47	26	0.0078

TABLE I. Hybrid waveforms used to calibrate the model. The first column lists the simulation IDs of the corresponding NR data in the SXS catalog. The last column reports the dimensionless frequency at the midpoint of the stitching region.




FIG. 1. Hybrid waveforms for four different configurations in the parameter space of q, χ_1, χ_2 . Inspirational waveforms are generated by adding the phase and amplitude corrections due to TH to the `SEOBNRv2` model, shown in blue. NR waveforms are shown in orange, and the hybrid waveforms are shown in black dashed-dotted lines. The x and y axes denote time (in units of total mass) and the real part of time-domain strain, $h_+(t)$ (in units of M/D_L), respectively. The blue shaded areas denote the stitching regions.

spurious pulse of gravitational radiation (often referred to as ‘junk radiation’).

- A robust hybrid should depend weakly on the small changes to the stitching region. A monochromatic signal would be completely degenerate under a shift in the coalescence time t_c and the coalescence phase ϕ_c ; this degeneracy is broken by the increase in fre-

quency with time. The start and end of the stitching interval should reflect enough change in the frequency to break this degeneracy.

The first issue is addressed in our work by choosing the stitching region beyond the specified *relaxation time* for each NR simulation in the SXS catalog, the time interval (in units of M) by which the junk radiation dies

out. To address the second requirement, MacDonald *et al.* [69] have recommended the stitching interval to satisfy $\delta\omega/\omega_m \geq 0.1$, where ω_m is the GW frequency at the midpoint of the stitching interval, and $\delta\omega$ is the change in frequency over the interval. This choice ensures that the residual oscillations in t_c with changing ω_m are below $1M$. In our work, we perform the stitching over 4 GW cycles, and we place the stitching region in a way to satisfy this condition. In the literature, the construction of hybrids has been performed both in time [20, 23, 24] and frequency domains [21, 70]. In our work, we choose the former, primarily because the NR waveforms are available in the time domain, and typically they have too few numbers of GW cycles for a reliable Fourier transform.

In time domain, any two non-precessing binary waveforms $h^A(\Theta, t)$ and $h^B(\Theta, t)$, with the same set of intrinsic parameters Θ , only differ by a relative time shift (Δt) and an overall constant phase difference (ϕ_0):

$$h^A(\Theta, t) = e^{i\phi_0} h^B(\Theta, t + \Delta t).$$

In the region where we expect the PN and NR waveforms to be morphologically similar, we can write

$$h^{\text{PN}}(\Theta, t) = e^{i\phi_0} h^{\text{NR}}(\Theta, t + \Delta t). \quad (19)$$

However, in reality, $h^{\text{PN}}(\Theta, t)$ and $h^{\text{NR}}(\Theta, t)$ differ from each other morphologically due to waveform systematics, so Eq. (19) does not hold in general [69]. To generate a hybrid waveform, then, one needs to find a suitable region where PN and NR waveforms are approximately the same and minimize the square of the difference between the left-hand and right-hand sides in Eq. (19), by varying Δt and ϕ_0 . Since the GW frequency of a CBC increases with time, one can alternatively write the angular GW frequency as a function of time ($\omega(t)$) and minimize the quantity

$$\delta = \int_{t_1}^{t_2} \left(\omega^{\text{NR}}(t) - \omega^{\text{PN}}(t + \Delta t) \right)^2 dt \quad (20)$$

with respect to Δt . Here t_1 and t_2 denote the start and end of the stitching region.

We list the steps taken to construct the hybrids as follows:

1. SEOBNRv2 is a time-domain waveform, and the phase and amplitude corrections for TH are calculated in the frequency domain. So, we have to first convert the time-domain data to the Fourier domain to apply the corrections. Since the time-domain data is finite, we use a tapered window function to minimize Gibbs artifacts, called the Planck-taper window [69]:

$$w(x) = \begin{cases} 0, & x \leq x_1 \\ [e^{y(x)} + 1]^{-1}, & x_1 < x < x_2 \\ 1, & x_2 \leq x \leq x_3 \\ [e^{z(x)} + 1]^{-1}, & x_3 < x < x_4 \\ 0, & x_4 \leq x, \end{cases} \quad (21)$$

where $y(x) = (x_2 - x_1)/(x - x_1) + (x_2 - x_1)/(x - x_2)$, and $z(x) = (x_3 - x_4)/(x - x_3) + (x_3 - x_4)/(x - x_4)$.

The frequency-domain waveform, after correcting for TH, is then converted back to the time domain for hybridization.

2. NR data from the SXS catalog sample the GW strain $h^{\text{NR}}(t)$ in non-uniform timesteps, with a higher sampling rate in regions of higher GW frequency. We resample the data with the coarsest sampling rate by using cubic interpolation.
3. Having obtained $h^{\text{PN}}(t)$ and $h^{\text{NR}}(t)$, we minimize the quantity δ defined in Eq. (20) with respect to Δt . Phase alignments are done at the temporal midpoint of the stitching region, $t = (t_1 + t_2)/2$.
4. The hybrid waveform is then constructed as

$$h^{\text{Hyb}}(t) = \mathcal{F}(t)h^{\text{PN}}(t) + [1 - \mathcal{F}(t)]e^{i\phi'_0}h^{\text{NR}}(t + \Delta t'), \quad (22)$$

where ϕ'_0 and $\Delta t'$ are the phase offset at the midpoint of the stitching region and the value of Δt that minimizes δ , respectively. $\mathcal{F}(t)$ is a blending function defined by

$$\mathcal{F}(t) = \begin{cases} 1, & t < t_1 \\ \frac{1}{2} \left(1 + \cos \frac{\pi(t-t_1)}{t_2-t_1} \right), & t_1 \leq t \leq t_2 \\ 0, & t_2 < t. \end{cases} \quad (23)$$

In the time domain hybrids, parts of the waveforms at $t < t_1$ are purely post-Newtonian, and for $t > t_2$ the waveforms contain purely NR data. In between, the blending function $\mathcal{F}(t)$ smoothly stitches the two waveforms together after proper alignment in phase and time.

Figure 1 shows four of the total twenty hybrids made for calibrating the waveform model, at the extreme points of the calibration parameter space. The blue shaded regions correspond to the stitching intervals, $t_1 \leq t \leq t_2$. In Table I, we list the NR data used to create the set of hybrids for calibration. We report the simulation labels of aligned-spin NR data from SXS, dimensionless spin magnitudes for the two component BHs, the effective spin parameter used for modeling the inspiral (discussed in Sec. VC), the number of orbits present in the NR data, and the value of dimensionless frequency Mf at the midpoint of the stitching regions.

V. MODELING THE HYBRIDS IN THE FREQUENCY DOMAIN

We perform phenomenological parametrization of the hybrid waveforms in the frequency domain. Template waveforms in the frequency domain are of particular preference since the parametrization is easier, and also a search using frequency-domain templates is computationally inexpensive compared to one using time-domain templates. Our parametrization follows the construction of **PhenomD** [24, 25] for modeling the phase and amplitude of the frequency-domain hybrids.

PhenomD used uncalibrated **SEOBNRv2** as the inspiral approximant, while 19 NR waveforms extracted by **SpEC** and **BAM** [71] codes were used for hybrid construction and calibration. The model was constructed in the frequency domain by dividing the entire waveform into three parts – inspiral, intermediate, and merger-ringdown. This modular nature allows one to use a different inspiral model while keeping the merger and ringdown intact. In that model, inspiral is defined to be the region below $F = 0.018$. Merger and ringdown phases are separated by identifying the ringdown frequencies of different hybrids that are used to calibrate the model. In **PhenomD**, the end of inspiral ($F = 0.018$) was chosen in such a way that beyond that frequency, each of the hybrids contained purely NR data. The merger-ringdown model of **PhenomD**, then, is a model of purely NR data, which we do not need to recalibrate in this work. In fact, as discussed in Ref. [24], the modular nature of **PhenomD** allows independent modeling of the inspiral, intermediate, and merger-ringdown parts separately, without even needing any hybrid waveform. The hybrid dataset, however, serves as a benchmark to compare the final model with and provide long time-domain waveforms for better frequency-domain modeling. Additionally, the hybrids are created with a smooth transition of phase and amplitude by the stitching process, producing accurate values of the pseudo-PN parameters capturing higher-order terms in frequency.

A. Inspiral phase model

The inspiral approximant used in the hybrid data is **SEOBNRv2** with corrections in phase and amplitude due to tidal heating, as discussed earlier. Since the final model is a phenomenological one, it requires ready-to-use analytical forms for the phase and amplitude with extra coefficients calibrated to the hybrid data. We write the frequency-domain phase of a waveform in the inspiral as

$$\begin{aligned} \Psi_{\text{INS}}(F; \Theta) &= \Psi_{\text{TF2}}(F; \Theta) + \Psi_{\text{TH}}(F; \Theta) \\ &+ \frac{1}{\eta} \left[\sigma_0 + \sigma_1 F + \frac{3}{4} \sigma_2 F^{4/3} \right. \\ &\quad \left. + \frac{3}{5} \sigma_3 F^{5/3} + \frac{1}{2} \sigma_4 F^2 \right], \end{aligned} \quad (24)$$

where $\Psi_{\text{TF2}}(F; \Theta)$ and $\Psi_{\text{TH}}(F; \Theta)$ are 3.5PN phase of **TaylorF2** (described in Appendix A) and the TH correction (Eq. (11)), respectively. σ_i ($i = 0 - 4$) are 5 phenomenological pseudo-PN parameters. The analytical ansatz in Eq. (24) is the same as in **PhenomD** inspiral model. Since we want to leverage the modular nature of **PhenomD** to concatenate its merger-ringdown model with our own inspiral model, the end of the inspiral phase in the current work needs to remain unchanged, which is $F = 0.018$. We also note here that one can, in principle, directly add the correction $\Psi_{\text{TH}}(F; \Theta)$ to the **PhenomD** inspiral phase (as is done for the amplitude, discussed in Sec. VB), without any recalibration. However, since we intend to avoid adding the correction to the NR information present in the last few cycles of inspiral, recalibrating the phase allows a smooth transition from PN to NR. To find the parameter values corresponding to each hybrid waveform, we fit the first derivative of the phase with respect to F :

$$\begin{aligned} \Psi'_{\text{INS}}(F; \Theta) &= \Psi'_{\text{TF2}}(F; \Theta) + \Psi'_{\text{TH}}(F; \Theta) \\ &+ \frac{1}{\eta} \left[\sigma_1 + \sigma_2 F^{1/3} \right. \\ &\quad \left. + \sigma_3 F^{2/3} + \sigma_4 F \right], \end{aligned} \quad (25)$$

where $\Psi' = \partial\Psi/\partial F$. σ_0 is determined by imposing $C^{(1)}$ continuity in phase at the boundary between inspiral and merger. The fits are performed over the frequency range $F \in [0.0035, 0.019]$, ending at a slightly higher frequency, to reduce boundary effects at the interface and find robust fits for σ_i . For all the 20 hybrids used for calibration, the stitching regions are placed within this frequency range, which warrants that there is purely NR data beyond $F = 0.018$.

B. Inspiral and intermediate amplitude model

To model the amplitude within the frequency range of the inspiral, we build the inspiral amplitude model by adding the correction due to TH, given by Eq. (15), to the **PhenomD** amplitude:

$$\tilde{A}_{\text{INS}}(F) = \tilde{A}_{\text{D}}(F) + \tilde{A}_{\text{TH}}(F). \quad (26)$$

We do not recalibrate the amplitude pseudo-PN parameters of **PhenomD**, since the inspiral amplitude model ends (at $F = 0.014$) before the phase model does (at $F = 0.018$), and the amount of NR information is less than that of phase. Moreover, we find that the effect of the amplitude correction is significantly less than the phase correction, diminishing the necessity of a recalibration.

At the interface of two frequency intervals modeled separately, $C^{(1)}$ continuity on the amplitude cannot be imposed in a straightforward manner. For the frequency-domain phase, one has the freedom to tune the coalescence time t_c and the coalescence phase ϕ_c to impose continuity of the phase and its derivative across an interface.




FIG. 2. Fits of σ_i calculated by the `NonLinearModelFit` module of Mathematica. The surfaces correspond to the 2D fits with η and χ_{PN} as described in Eq. (30).

For amplitude, however, there is no such freedom. If the amplitude is modeled separately in different frequency regimes, a different strategy has to be followed. To ensure that the amplitude and its derivative are continuous throughout the three phases, one needs a frequency interval to fit a polynomial that satisfies these conditions at the beginning and the end. This ‘intermediate’ interval is defined to be $F \in [0.014, F_{\text{peak}}]$, where $F_{\text{peak}} \equiv M f_{\text{peak}}$ is the frequency corresponding to the peak amplitude. For $F < 0.014$, the amplitude is given by Eq. (26). In the intermediate region, the amplitude is approximated as a polynomial in f :

$$\tilde{A}_{\text{int}} = A_0 [\delta_0 + \delta_1 f + \delta_2 f^2 + \delta_3 f^3 + \delta_4 f^4]. \quad (27)$$

Here A_0 includes the leading order $f^{-7/6}$ behaviour. Evaluation of δ_i follows the steps of `PhenomD`, described in Sec. V(C) of Ref [25]. We briefly summarize it here:

- Equation (27) has five parameters, requiring five independent equations for unique solutions. Two of them come from the $C^{(1)}$ continuity of \tilde{A}_{int} with $\tilde{A}_{\text{INS}}(f)$ at the beginning ($F = 0.014$), and two from $C^{(1)}$ continuity with `PhenomD` amplitude at F_{peak} . The fifth equation appears from an additional condition that the polynomial coincides with `PhenomD` amplitude at the mid-frequency (chosen as a collocation point), $F = (0.014 + F_{\text{peak}})/2$.
- Solving the aforementioned set of equations, one finds the form of \tilde{A}_{int} which smoothly connects the intermediate region with inspiral and merger.
- It is worth mentioning that while choosing more collocation points would improve the model accuracy, it would also encumber the model with a larger set of equations to solve. Keeping in mind that amplitude errors are less consequential than phase errors, we refrain from considering more collocation points.

In Appendix B, we describe the evaluation of the peak frequency f_{peak} and the ringdown frequency f_{RD} as functions of the BH masses and spins.

C. Correspondence between the phenomenological and physical parameters

Our model has 4 phenomenological parameters corresponding to the inspiral phase. These parameters play an intermediate role in generating gravitational waveforms of aligned-spin compact binaries. To generate waveforms for arbitrary values of the masses and spins, one needs to build a correspondence between the set of physical parameters $\{\eta, \chi_1, \chi_2\}$ and the set of phenomenological parameters $\{\sigma_i\}$. The total mass M of the binary works as a trivial scaling factor.

In PN expansions, the leading order contribution of spin in the phase of the waveform appears as a function of the combination

$$\chi_{\text{PN}} = \chi_{\text{eff}} - \frac{38\eta}{113}(\chi_1 + \chi_2), \quad (28)$$

where

$$\chi_{\text{eff}} = \frac{m_1 \chi_1 + m_2 \chi_2}{M}. \quad (29)$$

In this model, χ_{PN} is used as a single spin parameter to generate the phenomenological parameters. This ‘effective spin approximation’ works well for aligned-spin binaries, especially in the inspiral regime [25].

To establish the correspondence between physical and phenomenological parameters, we fit σ_i obtained from different hybrids with a polynomial of η (up to second order) and the effective spin parameter χ_{PN} (up to third order):

$$\begin{aligned} \sigma_i = & \lambda_{00}^i + \lambda_{10}^i \eta + (\chi_{\text{PN}} - 1)(\lambda_{01}^i + \lambda_{11}^i \eta + \lambda_{21}^i \eta^2) \\ & + (\chi_{\text{PN}} - 1)^2(\lambda_{02}^i + \lambda_{12}^i \eta + \lambda_{22}^i \eta^2) \\ & + (\chi_{\text{PN}} - 1)^3(\lambda_{03}^i + \lambda_{13}^i \eta + \lambda_{23}^i \eta^2). \end{aligned} \quad (30)$$




FIG. 3. Phase, phase derivative, and phase difference between the hybrids (same as in Fig. 1), individual fits obtained from Eq. (25), and the final model prediction. *Top panels*: Inspiral phase $\Psi(Mf)$. *Middle panels*: $-\Psi'(Mf) = -\partial\Psi/\partial(Mf)$ plotted against Mf . *Bottom panels*: $\Delta\Psi_{\text{Fit}} = (\Psi_{\text{Hybrid}} - \Psi_{\text{Fit}})$, and $\Delta\Psi_{\text{Model}} = (\Psi_{\text{Hybrid}} - \Psi_{\text{Model}})$. All the plots are shown within the inspiral frequency range, $0.0035 \leq Mf \leq 0.018$.




FIG. 4. Rescaled amplitude of the full IMR waveforms as functions of Mf . The blue shaded regions denote the intermediate frequency range, $0.014 \leq Mf \leq F_{\text{peak}}$.

For each σ_i , this polynomial generates a set $\{\lambda_{jk}^i\}$ to build the correspondence between physical and phenomenological parameter sets. Figure 2 shows the surfaces of σ_i as functions of $\{\eta, \chi_{\text{PN}}\}$ with the ansatz in Eq. (30). These best-fit surfaces generate a constant set of $\{\lambda_{jk}^i\}$ which define the inspiral model, along with the TaylorF2 phase and the TH correction. In Appendix C, we list the coefficient values corresponding to each σ_i .

A schematic flowchart for generating the inspiral phase model starting from the intrinsic binary parameters can be described as:

$$\{\eta, \chi_1, \chi_2\} \xrightarrow[\text{spin}]{\text{Effective}} \{\eta, \chi_{\text{PN}}\} \xrightarrow{\lambda^i} \{\sigma_i\} \rightarrow \text{Inspiral model.}$$

D. Full IMR model

For the phase, we have separately modeled the frequency region $0.0035 \leq F \leq 0.018$, and we have the phase model of PhenomD thereafter. For the amplitude, we have a corrected model in the range $F \leq 0.014$, and a polynomial function for $0.014 \leq F \leq F_{\text{peak}}$. This gives us one interface for the phase and two interfaces for the amplitude to impose $C^{(1)}$ continuity. For the amplitude, however, $C^{(1)}$ continuity is ensured by the intermediate polynomial itself. For the phase, we can vary t_c and ϕ_c of the inspiral phase. After imposing $C^{(1)}$ continuity, we can generate the complete IMR phase and amplitude by defining a step function

$$\theta(f - f_0) = \begin{cases} -1, & f < f_0 \\ +1, & f \geq f_0. \end{cases} \quad (31)$$

Using this function, one can define

$$\theta^\pm(f; f_0) = \frac{1}{2}[1 \pm \theta(f - f_0)], \quad (32)$$

so that the IMR phase can be written as

$$\Phi_{\text{IMR}}(F) = \Phi_{\text{INS}}(F)\theta^-(F; F_2) + \theta^+(F; F_2)\Phi_{\text{D}}(F), \quad (33)$$

where $F_2 = 0.018$, and $\Phi_D(F)$ corresponds to the phase model of `PhenomD`. $\Phi_{\text{INS}}(F)$ is given by Eq. (24).

The IMR amplitude model follows a similar treatment, which can be expressed as

$$\begin{aligned} \tilde{A}_{\text{IMR}}(F) = & \tilde{A}_{\text{INS}}(F)\theta^-(F; F_3) \\ & + \theta^+(F; F_3)\tilde{A}_{\text{int}}(F)\theta^-(F; F_4) + \theta^+(F; F_4)\tilde{A}_D(F), \end{aligned} \quad (34)$$

where $F_3 = 0.014$, and $F_4 = F_{\text{peak}}$. $\tilde{A}_{\text{INS}}(F)$ is the inspiral amplitude in Eq. (26), $\tilde{A}_{\text{int}}(F)$ is the intermediate amplitude defined in Eq. (27), and $\tilde{A}_D(F)$ is the `PhenomD` amplitude.

Finally, the plus and cross polarization states of the complete frequency-domain waveform read,

$$\begin{aligned} \tilde{h}_+(f) = & \tilde{A}_{\text{IMR}}(f) \left(\frac{1 + \cos^2 \iota}{2} \right) \\ & \times \exp\{-i(\Phi_{\text{IMR}}(f) - \phi_0 - 2\pi f t_0)\}, \end{aligned} \quad (35)$$

and

$$\begin{aligned} \tilde{h}_\times(f) = & -i\tilde{A}_{\text{IMR}}(f) \cos \iota \\ & \times \exp\{-i(\Phi_{\text{IMR}}(f) - \phi_0 - 2\pi f t_0)\}. \end{aligned} \quad (36)$$

Here ι is the angle of inclination of the binary plane to the line of sight, ϕ_0 and t_0 carry the overall phase and timeshift freedom of the complete waveform. We call the final waveform model `IMRPhenomD_Horizon`, and abbreviate it to `PhenomD_Horizon`.

In Fig. 3, we show the inspiral phase $\Psi(Mf)$, its derivative with respect to Mf , and the difference between the phase of the hybrid data and the model, for the same configurations as in Fig. 1. We show both the individual fits from the ansatz in Eq. (24) and the final model prediction obtained from the best fit values of λ_{jk}^i from Eq. (30). In Fig. 4, we show the inspiral and intermediate amplitude, rescaled by the leading order factor $A_0 = \frac{1}{\pi^{2/3}} \sqrt{5\eta/24} f^{-7/6}$. The intermediate regions are shown in a blue shaded colour. We also show the amplitude of `PhenomD` on the same plots.

VI. FAITHFULNESS OF THE MODEL

A phenomenological model, calibrated with a set of “target hybrids”, should be compared with a larger set of “test hybrids” to assess its accuracy. The test hybrids should contain both the target hybrids and a new set of hybrids interspersed within the parameter space of calibration. To study the model’s behavior beyond the calibration range, test hybrids can be extended further in the parameter space. How well the final model can predict these test hybrids is denoted by its *effectualness* and *faithfulness* [20, 72]. A model is effectual if it is accurate enough to predict a GW signal from the detector noise, while to be faithful it also has to have enough accuracy to estimate the binary parameters. A faithful model is also effectual, but the converse may not be true. Given

#	Simulation label	q	χ_1	χ_2
1	SXS:BBH:0159	1	-0.9	-0.9
2	SXS:BBH:0154	1	-0.8	-0.8
3	SXS:BBH:0148	1	-0.44	-0.44
4	SXS:BBH:0150	1	0.2	0.2
5	SXS:BBH:0170	1	0.44	0.44
6	SXS:BBH:0155	1	0.8	0.8
7	SXS:BBH:0160	1	0.9	0.9
8	SXS:BBH:0157	1	0.95	0.95
9	SXS:BBH:0014	1.5	-0.5	0
10	SXS:BBH:0008	1.5	0	0
11	SXS:BBH:0013	1.5	0.5	0
12	SXS:BBH:0046	3	-0.5	-0.5
13	SXS:BBH:0036	3	-0.5	0
14	SXS:BBH:0168	3	0	0
15	SXS:BBH:0031	3	0.5	0
16	SXS:BBH:0047	3	0.5	0.5
17	SXS:BBH:0056	5	0	0
18	SXS:BBH:0181	6	0	0
19	SXS:BBH:1424	6.465	-0.66	-0.8
20	SXS:BBH:0298	7	0	0

TABLE II. List of the extra hybrids created for validating the model.




FIG. 5. Mismatches (%) between `IMRPhenomD_Horizon` and the hybrid waveforms (Table I and Table II) in aLIGO ZDHP noise PSD with a lower cutoff of 10 Hz, as a function of the total mass.

any two signals $h(t)$ and $g(t)$, buried in the noise $n(t)$, one can define a noise-weighted inner product

$$\langle h|g \rangle = 2 \int_0^\infty \frac{\tilde{h}(f)\tilde{g}^*(f) + \tilde{h}^*(f)\tilde{g}(f)}{S_n(f)} df, \quad (37)$$

where $\tilde{h}(f)$ is the Fourier transform of $h(t)$ defined by

$$\tilde{h}(f) = \int_{-\infty}^\infty h(t)e^{-2\pi ift} dt, \quad (38)$$

and $S_n(f)$ is the power spectral density (PSD) of the noise.

Equipped with this definition of an inner product between two waveforms, one can define the *match* between them by maximizing the inner product with respect to the constant phase and timeshift freedom between the waveforms $\tilde{h}(f)$ and $\tilde{g}(f)$, normalized by their individual norms:

$$\mathcal{M} = \min_{t_c, \phi_c} \frac{\langle h(\lambda) | g(\lambda') \rangle}{\sqrt{\langle h(\lambda) | h(\lambda) \rangle \langle g(\lambda') | g(\lambda') \rangle}}. \quad (39)$$

Here λ and λ' denote the intrinsic parameters. The *mismatch*, defined as $1 - \mathcal{M}$, quantifies how much the waveform $\tilde{h}(f)$ morphologically differs from $\tilde{g}(f)$. The faithfulness of a model is quantified by demanding that the mismatches between the model and the validation dataset (test hybrids) do not exceed a certain threshold. For **PhenomD**, the threshold was 1%.

Since the total mass of the binary works as a scaling factor for the gravitational waveforms, it defines how many GW cycles are contained within the sensitive frequency band of a detector. The lower the total mass, the more time the binary spends in its inspiral phase, resulting in a longer waveform given a lower cutoff for the frequency. In Advanced LIGO, complete IMR models become important above a critical value $\sim 12M_\odot$ [73] of the binary mass. To assess our model's faithfulness, we create a set of test hybrids that are long enough to cover the sensitive frequency band of Advanced LIGO for a total binary mass as low as $10M_\odot$. At 10 Hz, the dimensionless frequency corresponding to a $10M_\odot$ binary is $Mf \approx 0.0005$.

In Table II we list the extra test hybrids created for assessing the model accuracy. Figure 5 shows the mismatches between the entire set of test hybrids (Table I and Table II) against the BBH waveform model presented in this paper. We have used the Advanced LIGO zero-detuned high power [74] (ZDHP) noise curve with a lower frequency cutoff of 10 Hz, and upper cutoff of $Mf = 0.3$. The mismatches are plotted against the total binary mass. We find that the model accuracy is at par with the accuracy standard of **PhenomD**, with almost all the hybrids showing mismatches well below 1%, the majority of them having mismatches around 0.1% within the mass range $10M_\odot \leq M \leq 100M_\odot$. The worst mismatch we find is for the $\{q = 4, \chi_1 = 0, \chi_2 = 0.8\}$ configuration, which crosses the 1% level above $\sim 50M_\odot$, and reaches a maximum of 1.21% at $70M_\odot$.

VII. MODEL COMPARISON

Figure 6 shows the mismatches between **PhenomD_Horizon** and **PhenomD**, keeping $\chi_1 = \chi_2$. We explore the mismatches within the parameter space $q \in [1, 8]$ and $\chi_1, \chi_2 \in [-0.9, 0.9]$, for six values of the binary mass, $M/M_\odot \in [10, 20, 30, 40, 50, 60]$. In Fig. 6(a), we show the mismatches including only the phase

correction - i.e. using the recalibrated inspiral phase, but keeping the IMR amplitude the same as **PhenomD**. Since the phase correction due to TH written in Eq. (11) is $\propto 1/\eta$ at the leading order, higher mismatches are found for higher mass ratios, for a fixed value of the spins. With increasing spins, mismatches rise due to the increased significance of TH. The mismatch reaches a maximum of $\sim 16\%$ for $30M_\odot$ binaries with $q \gtrsim 7$ and $\chi_1 = \chi_2 \gtrsim 0.8$, and a minimum of $\sim 10^{-5}\%$ for heavier binaries. For $M \leq 30M_\odot$, we see substantial regions of the parameter space with mismatches between 1-10% for spin magnitudes of $|\chi_1| = |\chi_2| \gtrsim 0.5$. For $M > 30M_\odot$, the 1% level is crossed only for $q \gtrsim 4$.

Figure 6(b) shows mismatches when only the amplitude correction is added to the inspiral and the intermediate region is re-evaluated accordingly, but the phase is kept the same as **PhenomD**. The mismatches, in this case, are orders of magnitude less than the phase-corrected model, and never exceed 0.001% within this parameter space. This result also vindicates our choice of not recalibrating the amplitude model, as discussed in Sec VB. We also note from Eq. (15) that at the leading order, $\tilde{A}_{\text{TH}} \propto \sqrt{\eta}$, which implies that the TH corrections are more significant for more symmetric masses. This is in contrast to the behaviour of the phase correction, and renders the amplitude correction ineffective for high mass-ratio binaries with notable TH contribution in the phase. Nevertheless, \tilde{A}_{TH} still rises with increasing spins.

In Fig. 6(c), we use the complete **IMRPhenomD_Horizon** model, by including both the phase and amplitude corrections. The contours of mismatches $\geq 0.1\%$ are almost the same as in Fig. 6(a), but amplitude corrections elevate them above 0.01% throughout other regions of the parameter space. However, mismatches $< 0.1\%$ are below the accuracy level of the model, for which we considered a 1% tolerance level of modeling errors (Fig. 5). This suggests that including the amplitude correction in the TH corrected model has negligible effect compared to the phase correction in GW data analysis.

To demonstrate the improvement in accuracy that our model introduces over **PhenomD**, we use 219 NR waveforms from the SXS catalog and compare them against **PhenomD** and **PhenomD_Horizon** separately by computing mismatches. In Fig. 7, we show the distributions of the mismatches, by using the Advanced LIGO ZDHP noise curve. We plot histograms of $\log_{10}(1 - \mathcal{M})$, where $(1 - \mathcal{M})$ is the average mismatch in the binary mass range $12M_\odot \leq M \leq 100M_\odot$. The dashed lines show the medians of the two histogram plots, where a slight overall improvement is seen. The fractional shift in the medians (denoted by an overbar), is

$$\frac{\overline{(1 - \mathcal{M})_{\text{PhenomD}}} - \overline{(1 - \mathcal{M})_{\text{PhenomD}_\text{Horizon}}}}{\overline{(1 - \mathcal{M})_{\text{PhenomD}}}} \approx 0.04, \quad (40)$$

showing an improvement of $\sim 4\%$. Figure 8 shows similar histograms with a flat noise curve. In this case, also we see a shift in the median value towards lower mismatches,




FIG. 6. Mismatches (%) between `PhenomD_Horizon` and `PhenomD`. (a) Mismatches with only the phase correction but no amplitude correction in `PhenomD_Horizon`. (b) Mismatches with only the amplitude correction without any phase correction. (c) Mismatches with corrections in both the phase and amplitude. The plots show that the dominant contributor to these mismatches is the phase correction.




FIG. 7. Mismatches of 219 non-precessing non-eccentric NR waveforms from SXS with `PhenomD_Horizon` and `PhenomD`, in aLIGO ZDHP noise curve, averaged in the binary mass range $12 - 100M_{\odot}$. The dashed lines show the median values of the distributions.

with an improvement of $\sim 2\%$.

VIII. DISCUSSION AND FUTURE WORK

We have presented a phenomenological gravitational waveform model for binary black holes that carries the signatures of tidal heating of the holes explicitly throughout the binary evolution, starting from the early inspiral phase and ending at their merger. In the frequency domain, it contains corrections due to this effect both in the phase and amplitude. The entire IMR model for the phase is divided into two parts. The inspiral phase model, defined by the frequency range $0.0035 \leq Mf \leq 0.018$, is a recalibrated version of `IMRPhenomD` [24, 25] to account




FIG. 8. Same as in Fig. 7, but with a flat noise curve.

for the modifications due to tidal heating. The merger-ringdown parts, defined as $Mf \geq 0.018$ in terms of the dimensionless frequency, is the same as `PhenomD` since this region contains purely NR data, which inherently incorporates the dephasing due to tidal heating. The amplitude corrections are added by dividing the model into three parts. The inspiral model, with $0.0035 \leq Mf \leq 0.014$, is constructed by directly adding the correction term to the amplitude model of `PhenomD` in this region. The intermediate region, $0.014 \leq Mf \leq Mf_{\text{peak}}$, f_{peak} being the frequency corresponding to the peak amplitude, is approximated as a polynomial in f and evaluated by imposing $C^{(1)}$ continuity at the two ends and a collocation point at their mid-frequency. The amplitude model is identical to `PhenomD` for $f > f_{\text{peak}}$. The final model is calibrated within the range in the mass ratio $q \in [1, 8]$ and the effective spin parameter $\chi_{\text{PN}} \in [-0.79, 0.72]$.

We conducted a study of faithfulness, by calculating mismatches between the model and a validation dataset

of 40 hybrid waveforms constructed with tidal heating included. We demonstrated that modeling errors do not lead to mismatches higher than 1% (barring a single outlier), with most of the mismatch values lying around 0.1%. This standard of faithfulness is comparable to **PhenomD**, which also considered a maximum tolerance level of 1% mismatch.

How the new model differs from its parent model **PhenomD** was assessed by calculating mismatches between them within the calibration range of the intrinsic parameter space, for binary masses ranging from $10M_\odot$ to $60M_\odot$. Phase corrections in the inspiral result in mismatches between these two models as high as $\sim 16\%$, whereas amplitude corrections alone are unable to produce mismatches above $\sim 0.001\%$. With increasing mass asymmetry, the impact of the phase correction increases, but the amplitude correction becomes less effective. Combined with the fact that the former has more significance in GW searches and PE, this reciprocal nature of these two correction terms leaves the latter mostly superfluous for practical purposes.

We compared the model with a set of 219 NR waveforms from the SXS catalog [53], and show the mismatch distribution along with the same for **PhenomD**, averaged over the binary mass range $12M_\odot \leq M \leq 100M_\odot$. With the Advanced LIGO ZDHP noise curve, we see an improvement of $\sim 4\%$ in the medians of the two distributions. With a flat noise curve, the improvement is relatively modest, about 2%. These overall improvements point towards a mild yet non-negligible increase in waveform accuracy.

Tidal heating of black holes, after all, is a weak effect in the context of comparable-mass binaries. Nevertheless, an absence of its signatures from the complete waveform models can bias the estimated parameters of a binary. Its absence can also mimic a deviation from GR predictions, raising complications in tests of GR. We leave these studies to future work. Additionally, tidal heating has the important property of varying significantly according to the nature of the component objects, which makes it a potential discriminator for BHs from other objects. However, to utilize the full power of TH as a BH identifier, one needs to model the energy and

angular momentum fluxes down the BH horizons also in the late inspiral regime described by NR, where the tidal fields are the strongest. The feasibility and effectiveness of modeling these quantities can be explored in future studies, where the model presented in this work can be used directly.

The current work, in its entirety, is based on the frequency-domain waveform model **IMRPhenomD**. For this reason, we perform all the comparisons of the model presented here with the said approximant only. More recent and state-of-the-art phenomenological waveform models are available currently, namely, the **IMRPhenomX** [23, 75] family of waveforms. These waveforms are calibrated with a much larger set of hybrid waveforms, and use superior techniques like hierarchical modeling [76] to reduce modeling errors, and also to generate more accurate waveforms with unequal spins. These models boast 1 to 2 orders of magnitude better faithfulness than **PhenomD**, and a dramatic shift in the mismatch distribution against NR waveforms towards lower values [23]. Including the effects of TH in these models by recalibrating the inspiral is beyond the scope of this work, which uses only publicly available NR data from the SXS catalog. We leave them as future possibilities.

ACKNOWLEDGMENTS

SM thanks the University Grants Commission (UGC), India, for providing financial support as a senior research fellowship. The high performance computing facility **Pegasus** at IUCAA was used for this work. The authors thank Vaishak Prasad, Apratim Ganguly, Anuj Mishra, Mark Scheel, Nathan Johnson-McDaniel, P. Ajith, and Shrobona Ghosh for useful comments and suggestions. The Python software packages **PyCBC** [77], **Scipy** [78], **Numpy** [79], **Matplotlib** [80] were used for computational works, analyses, and plots. **Mathematica** [81] was used for data fitting purposes. SB acknowledges the hospitality of IUCAA with gratitude and support from the NSF under Grant PHY-2309352.

Appendix A: Post-Newtonian inspiral phase

We write the 3.5PN frequency-domain phase (without TH) as

$$\Psi_{\text{TF2}}(f) = 2\pi f t_c - \phi_c - \frac{\pi}{4} + \frac{3}{128\eta v^5} \sum_{n=0}^7 \psi_{(n)} v^n, \quad (\text{A1})$$

where $v = (\pi M f)^{1/3}$.

The coefficients $\{\psi_{(n)}\}$ ($n=0-7$) are given by [82–84],

$$\psi_{(0)} = 1, \quad (\text{A2})$$

$$\psi_{(1)} = 0, \quad (\text{A3})$$

$$\psi_{(2)} = \frac{3715}{756} + \frac{55}{9}\eta, \quad (\text{A4})$$

$$\psi_{(3)} = -16\pi + \frac{113}{3}\delta\chi_a + \left(\frac{113}{3} - \frac{76}{3}\eta\right)\chi_s, \quad (\text{A5})$$

$$\psi_{(4)} = \frac{15293365}{508032} + \frac{27145}{504}\eta + \frac{3085}{72}\eta^2 - \frac{405}{4}\delta\chi_a\chi_s + \left(-\frac{405}{8} + 200\eta\right)\chi_a^2 + \left(-\frac{405}{8} + \frac{5}{2}\eta\right)\chi_s^2, \quad (\text{A6})$$

$$\psi_{(5)} = \left[\frac{38645\pi}{756} - \frac{65\pi}{9}\eta - \left(\frac{732985}{2268} - \frac{24260}{81}\eta - \frac{340}{9}\eta^2\right)\chi_s - \left(\frac{732985}{2268} + \frac{140}{9}\eta\right)\delta\chi_a\right](1 + 3\ln v), \quad (\text{A7})$$

$$\begin{aligned} \psi_{(6)} = & \frac{11583231236531}{4694215680} - \frac{640\pi^2}{3} - \frac{6848}{21}\gamma_E + \eta \left(-\frac{15737765635}{3048192} + \frac{2255\pi^2}{12}\right) + \frac{76055}{1728}\eta^2 - \frac{127825}{1296}\eta^3 - \frac{6848}{21}\ln(4v) \\ & + \pi \left\{ \frac{2270}{3}\delta\chi_a + \left(\frac{2270}{3} - 520\eta\right)\chi_s \right\} + \left(\frac{75515}{144} - \frac{8225}{18}\eta\right)\delta\chi_a\chi_s + \left(\frac{75515}{288} - \frac{263245}{252}\eta - 480\eta^2\right)\chi_a^2 \\ & + \left(\frac{75515}{288} - \frac{232415}{504}\eta + \frac{1255}{9}\eta^2\right)\chi_s^2, \end{aligned} \quad (\text{A8})$$

$$\begin{aligned} \psi_{(7)} = & \frac{77096675\pi}{254016} + \frac{378515\pi}{1512}\eta - \frac{74045\pi}{756}\eta^2 + \left\{ -\frac{25150083775}{3048192} + \frac{10566655595}{762048}\eta - \frac{1042165}{3024}\eta^2 + \frac{5345}{36}\eta^3 \right. \\ & \left. + \left(\frac{14585}{8} - 7270\eta + 80\eta^2\right)\chi_a^2 \right\}\chi_s + \left\{ \left(\frac{-25150083775}{3048192} + \frac{26804935}{6048}\eta - \frac{1985}{48}\eta^2\right)\chi_a \right. \\ & \left. + \left(\frac{14585}{24} - 2380\eta\right)\chi_a^3 + \left(\frac{14585}{8} - \frac{215}{2}\eta\right)\chi_a\chi_s^2 \right\}\delta + \left(\frac{14585}{24} - \frac{475}{6}\eta + \frac{100}{3}\eta^2\right)\chi_s^3. \end{aligned} \quad (\text{A9})$$

Here $\delta = \sqrt{1-4\eta}$, $\chi_s = (\chi_1 + \chi_2)/2$, $\chi_a = (\chi_1 - \chi_2)/2$. γ_E is the Euler's constant, $\gamma_E = 0.5772156\dots$.

Appendix B: Ringdown and peak frequency

The quasi-normal mode (QNM) frequency of a BH can be expressed in terms of its oscillatory (real) and damping (imaginary) parts as,

$$f_{\text{QNM}} = f_{\text{RD}} - if_{\text{damp}}. \quad (\text{B1})$$

f_{RD} can be obtained from the fitting formula [85]

$$f_{\text{RD}} = \frac{1}{2\pi M_f} [f_1 + f_2(1 - \chi_f)^{f_3}], \quad (\text{B2})$$

where M_f and χ_f are the mass and spin of the merger remnant BH. Evaluation of these quantities is described in Appendix C of Ref [86]. For the dominant harmonic mode $l = m = 2$, the coefficients are given by, $f_1 = 1.5251$, $f_2 = -1.1568$, $f_3 = 0.1292$.

f_{damp} can be written as

$$f_{\text{damp}} = \frac{f_{\text{RD}}}{2Q}, \quad (\text{B3})$$

where the quality factor Q can be expressed by the fitting formula

$$Q = q_1 + q_2(1 - \chi_f)^{q_3}. \quad (\text{B4})$$

For the $l = m = 2$ mode, $q_1 = 0.7$, $q_2 = 1.4187$, $q_3 = -0.499$.

The frequency corresponding to the peak amplitude in the Fourier domain, f_{peak} , can then be obtained from the phenomenological relation

$$f_{\text{peak}} = \left| f_{\text{RD}} + \frac{f_{\text{damp}}\gamma_3(\sqrt{1-\gamma_2^2}-1)}{\gamma_2} \right|, \quad (\text{B5})$$

γ_2 and γ_3 being two phenomenological parameters that can be evaluated for given values of $\{\eta, \chi_1, \chi_2\}$ from the ansatz in Eq. (30). The corresponding coefficients are given in Appendix C of Ref. [25].

Appendix C: Coefficients for the pseudo-PN parameters

	σ_1	σ_2	σ_3	σ_4
λ_{00}	1650.156	14056.472	-86832.697	146221.791
λ_{10}	-3531.389	-75378.11	412796.518	-667838.551
λ_{01}	-2661.071	137530.39	-623322.321	920614.957
λ_{11}	15385.008	-944130.978	4.1416×10^6	-5.959×10^6
λ_{21}	-219.668	1.7166×10^6	-7.1273×10^6	9.8223×10^6
λ_{02}	-6845.194	169601.655	-737033.591	1.0493×10^6
λ_{12}	38637.011	-1.2802×10^6	5.4343×10^6	-7.5726×10^6
λ_{22}	-45127.308	2.648×10^6	-1.09142×10^7	1.4893×10^7
λ_{03}	-2076.335	50611.846	-213200.291	288184.3
λ_{13}	17433.08	-468141.986	1.9402×10^6	-2.641×10^6
λ_{23}	-31696.712	1.10645×10^6	-4.5663×10^6	6.2489×10^6

TABLE III. Best-fit values of λ_{jk}^i parameters in Eq. (30) for the pseudo-PN parameters $\{\sigma_i\}$ in the inspiral phase.

-
- [1] R. Abbott *et al.* (LIGO Scientific, VIRGO, KAGRA), GWTC-3: Compact Binary Coalescences Observed by LIGO and Virgo During the Second Part of the Third Observing Run, (2021), [arXiv:2111.03606 \[gr-qc\]](#).
- [2] A. H. Nitz, S. Kumar, Y.-F. Wang, S. Kasta, S. Wu, M. Schäfer, R. Dhurkunde, and C. D. Capano, 4-OGC: Catalog of Gravitational Waves from Compact Binary Mergers, *Astrophys. J.* **946**, 59 (2023), [arXiv:2112.06878 \[astro-ph.HE\]](#).
- [3] S. Olsen, T. Venumadhav, J. Mushkin, J. Roulet, B. Zackay, and M. Zaldarriaga, New binary black hole mergers in the LIGO-Virgo O3a data, *Phys. Rev. D* **106**, 043009 (2022), [arXiv:2201.02252 \[astro-ph.HE\]](#).
- [4] J. Aasi *et al.* (LIGO Scientific), Advanced LIGO, *Class. Quant. Grav.* **32**, 074001 (2015), [arXiv:1411.4547 \[gr-qc\]](#).
- [5] F. Acernese, M. Agathos, K. Agatsuma, D. Aisa, N. Allemandou, A. Allocca, J. Amarni, P. Astone, G. Balestri, G. Ballardin, *et al.*, Advanced Virgo: a second-generation interferometric gravitational wave detector, *Classical and Quantum Gravity* **32**, 024001 (2015), [arXiv:1408.3978 \[gr-qc\]](#).
- [6] T. Akutsu *et al.* (KAGRA), KAGRA: 2.5 Generation Interferometric Gravitational Wave Detector, *Nature Astron.* **3**, 35 (2019), [arXiv:1811.08079 \[gr-qc\]](#).
- [7] B. J. Owen and B. S. Sathyaprakash, Matched filtering of gravitational waves from inspiraling compact binaries: Computational cost and template placement, *Phys. Rev. D* **60**, 022002 (1999), [arXiv:gr-qc/9808076](#).
- [8] J. Veitch *et al.*, Parameter estimation for compact binaries with ground-based gravitational-wave observations using the LALInference software library, *Phys. Rev. D* **91**, 042003 (2015), [arXiv:1409.7215 \[gr-qc\]](#).
- [9] G. Ashton *et al.*, BILBY: A user-friendly Bayesian inference library for gravitational-wave astronomy, *Astrophys. J. Suppl.* **241**, 27 (2019), [arXiv:1811.02042 \[astro-ph.IM\]](#).
- [10] I. M. Romero-Shaw *et al.*, Bayesian inference for compact binary coalescences with bilby: validation and application to the first LIGO-Virgo gravitational-wave transient catalogue, *Mon. Not. Roy. Astron. Soc.* **499**, 3295 (2020), [arXiv:2006.00714 \[astro-ph.IM\]](#).
- [11] L. Blanchet, Gravitational Radiation from Post-Newtonian Sources and Inspiralling Compact Binaries, *Living Rev. Rel.* **17**, 2 (2014), [arXiv:1310.1528 \[gr-qc\]](#).
- [12] A. Buonanno and T. Damour, Effective one-body approach to general relativistic two-body dynamics, *Phys. Rev. D* **59**, 084006 (1999), [arXiv:gr-qc/9811091](#).
- [13] A. Bohé *et al.*, Improved effective-one-body model of spinning, nonprecessing binary black holes for the era of gravitational-wave astrophysics with advanced detectors, *Phys. Rev. D* **95**, 044028 (2017), [arXiv:1611.03703 \[gr-qc\]](#).
- [14] A. Nagar and P. Retteno, Efficient effective one body time-domain gravitational waveforms, *Phys. Rev. D* **99**, 021501 (2019), [arXiv:1805.03891 \[gr-qc\]](#).
- [15] A. Taracchini, Y. Pan, A. Buonanno, E. Barausse, M. Boyle, T. Chu, G. Lovelace, H. P. Pfeiffer, and

- M. A. Scheel, Prototype effective-one-body model for nonprecessing spinning inspiral-merger-ringdown waveforms, *Phys. Rev. D* **86**, 024011 (2012), [arXiv:1202.0790 \[gr-qc\]](#).
- [16] A. Taracchini *et al.*, Effective-one-body model for black-hole binaries with generic mass ratios and spins, *Phys. Rev. D* **89**, 061502 (2014), [arXiv:1311.2544 \[gr-qc\]](#).
- [17] A. Nagar *et al.*, Time-domain effective-one-body gravitational waveforms for coalescing compact binaries with nonprecessing spins, tides and self-spin effects, *Phys. Rev. D* **98**, 104052 (2018), [arXiv:1806.01772 \[gr-qc\]](#).
- [18] R. Gamba, S. Akçay, S. Bernuzzi, and J. Williams, Effective-one-body waveforms for precessing coalescing compact binaries with post-Newtonian twist, *Phys. Rev. D* **106**, 024020 (2022), [arXiv:2111.03675 \[gr-qc\]](#).
- [19] A. Nagar, G. Riemenschneider, G. Pratten, P. Rettengo, and F. Messina, Multipolar effective one body waveform model for spin-aligned black hole binaries, *Phys. Rev. D* **102**, 024077 (2020), [arXiv:2001.09082 \[gr-qc\]](#).
- [20] P. Ajith *et al.*, A Template bank for gravitational waveforms from coalescing binary black holes. I. Non-spinning binaries, *Phys. Rev. D* **77**, 104017 (2008), [Erratum: *Phys.Rev.D* 79, 129901 (2009)], [arXiv:0710.2335 \[gr-qc\]](#).
- [21] L. Santamaria *et al.*, Matching post-Newtonian and numerical relativity waveforms: systematic errors and a new phenomenological model for non-precessing black hole binaries, *Phys. Rev. D* **82**, 064016 (2010), [arXiv:1005.3306 \[gr-qc\]](#).
- [22] M. Hannam, P. Schmidt, A. Bohé, L. Haegel, S. Husa, F. Ohme, G. Pratten, and M. Pürrer, Simple Model of Complete Precessing Black-Hole-Binary Gravitational Waveforms, *Phys. Rev. Lett.* **113**, 151101 (2014), [arXiv:1308.3271 \[gr-qc\]](#).
- [23] G. Pratten, S. Husa, C. Garcia-Quiros, M. Colleoni, A. Ramos-Buades, H. Estelles, and R. Jaume, Setting the cornerstone for a family of models for gravitational waves from compact binaries: The dominant harmonic for nonprecessing quasicircular black holes, *Phys. Rev. D* **102**, 064001 (2020), [arXiv:2001.11412 \[gr-qc\]](#).
- [24] S. Husa, S. Khan, M. Hannam, M. Pürrer, F. Ohme, X. Jiménez Forteza, and A. Bohé, Frequency-domain gravitational waves from nonprecessing black-hole binaries. I. New numerical waveforms and anatomy of the signal, *Phys. Rev. D* **93**, 044006 (2016), [arXiv:1508.07250 \[gr-qc\]](#).
- [25] S. Khan, S. Husa, M. Hannam, F. Ohme, M. Pürrer, X. Jiménez Forteza, and A. Bohé, Frequency-domain gravitational waves from nonprecessing black-hole binaries. II. A phenomenological model for the advanced detector era, *Phys. Rev. D* **93**, 044007 (2016), [arXiv:1508.07253 \[gr-qc\]](#).
- [26] M. Tiglio and A. Villanueva, Reduced order and surrogate models for gravitational waves, *Living Rev. Rel.* **25**, 2 (2022), [arXiv:2101.11608 \[gr-qc\]](#).
- [27] S. E. Field, C. R. Galley, J. S. Hesthaven, J. Kaye, and M. Tiglio, Fast prediction and evaluation of gravitational waveforms using surrogate models, *Phys. Rev. X* **4**, 031006 (2014), [arXiv:1308.3565 \[gr-qc\]](#).
- [28] J. Blackman, S. E. Field, M. A. Scheel, C. R. Galley, C. D. Ott, M. Boyle, L. E. Kidder, H. P. Pfeiffer, and B. Szilágyi, Numerical relativity waveform surrogate model for generically precessing binary black hole mergers, *Phys. Rev. D* **96**, 024058 (2017), [arXiv:1705.07089 \[gr-qc\]](#).
- [29] P. Kumar, T. Chu, H. Fong, H. P. Pfeiffer, M. Boyle, D. A. Hemberger, L. E. Kidder, M. A. Scheel, and B. Szilágyi, Accuracy of binary black hole waveform models for aligned-spin binaries, *Phys. Rev. D* **93**, 104050 (2016), [arXiv:1601.05396 \[gr-qc\]](#).
- [30] J. Blackman, S. E. Field, C. R. Galley, B. Szilágyi, M. A. Scheel, M. Tiglio, and D. A. Hemberger, Fast and Accurate Prediction of Numerical Relativity Waveforms from Binary Black Hole Coalescences Using Surrogate Models, *Phys. Rev. Lett.* **115**, 121102 (2015), [arXiv:1502.07758 \[gr-qc\]](#).
- [31] V. Varma, S. E. Field, M. A. Scheel, J. Blackman, L. E. Kidder, and H. P. Pfeiffer, Surrogate model of hybridized numerical relativity binary black hole waveforms, *Phys. Rev. D* **99**, 064045 (2019), [arXiv:1812.07865 \[gr-qc\]](#).
- [32] M. Pürrer, Frequency domain reduced order model of aligned-spin effective-one-body waveforms with generic mass-ratios and spins, *Phys. Rev. D* **93**, 064041 (2016), [arXiv:1512.02248 \[gr-qc\]](#).
- [33] B. D. Lackey, S. Bernuzzi, C. R. Galley, J. Meidam, and C. Van Den Broeck, Effective-one-body waveforms for binary neutron stars using surrogate models, *Phys. Rev. D* **95**, 104036 (2017), [arXiv:1610.04742 \[gr-qc\]](#).
- [34] B. D. Lackey, M. Pürrer, A. Taracchini, and S. Marsat, Surrogate model for an aligned-spin effective one body waveform model of binary neutron star inspirals using Gaussian process regression, *Phys. Rev. D* **100**, 024002 (2019), [arXiv:1812.08643 \[gr-qc\]](#).
- [35] K. Barkett *et al.*, Gravitational waveforms for neutron star binaries from binary black hole simulations, *Phys. Rev. D* **93**, 044064 (2016), [arXiv:1509.05782 \[gr-qc\]](#).
- [36] T. Dietrich *et al.*, Matter imprints in waveform models for neutron star binaries: Tidal and self-spin effects, *Phys. Rev. D* **99**, 024029 (2019), [arXiv:1804.02235 \[gr-qc\]](#).
- [37] E. Poisson, Tidal interaction of black holes and Newtonian viscous bodies, *Phys. Rev. D* **80**, 064029 (2009), [arXiv:0907.0874 \[gr-qc\]](#).
- [38] V. Cardoso and P. Pani, Tidal acceleration of black holes and superradiance, *Class. Quant. Grav.* **30**, 045011 (2013), [arXiv:1205.3184 \[gr-qc\]](#).
- [39] J. B. Hartle, Tidal Friction in Slowly Rotating Black Holes, *Phys. Rev.* **D8**, 1010 (1973).
- [40] E. Poisson, Absorption of mass and angular momentum by a black hole: Time-domain formalisms for gravitational perturbations, and the small-hole / slow-motion approximation, *Phys. Rev.* **D70**, 084044 (2004), [arXiv:gr-qc/0407050 \[gr-qc\]](#).
- [41] K. Alvi, Energy and angular momentum flow into a black hole in a binary, *Phys. Rev.* **D64**, 104020 (2001), [arXiv:gr-qc/0107080 \[gr-qc\]](#).
- [42] K. Chatziioannou, E. Poisson, and N. Yunes, Improved next-to-leading order tidal heating and torquing of a Kerr black hole, *Phys. Rev. D* **94**, 084043 (2016), [arXiv:1608.02899 \[gr-qc\]](#).
- [43] S. Datta, Horizon fluxes of binary black holes in eccentric orbits, (2023), [arXiv:2305.03771 \[gr-qc\]](#).
- [44] S. Datta, K. S. Phukon, and S. Bose, Recognizing black holes in gravitational-wave observations: Challenges in telling apart impostors in mass-gap binaries, *Phys. Rev. D* **104**, 084006 (2021), [arXiv:2004.05974 \[gr-qc\]](#).
- [45] S. Mukherjee, S. Datta, S. Tiwari, K. S. Phukon, and S. Bose, Toward establishing the presence or absence of horizons in coalescing binaries of compact objects by using their gravitational wave signals, *Phys. Rev. D* **106**,

- 104032 (2022), [arXiv:2202.08661 \[gr-qc\]](#).
- [46] V. Cardoso, S. Hopper, C. F. B. Macedo, C. Palenzuela, and P. Pani, Gravitational-wave signatures of exotic compact objects and of quantum corrections at the horizon scale, *Phys. Rev. D* **94**, 084031 (2016), [arXiv:1608.08637 \[gr-qc\]](#).
- [47] S. Datta, R. Brito, S. Bose, P. Pani, and S. A. Hughes, Tidal heating as a discriminator for horizons in extreme mass ratio inspirals, *Phys. Rev. D* **101**, 044004 (2020), [arXiv:1910.07841 \[gr-qc\]](#).
- [48] S. Datta and S. Bose, Probing the nature of central objects in extreme-mass-ratio inspirals with gravitational waves, *Phys. Rev. D* **99**, 084001 (2019), [arXiv:1902.01723 \[gr-qc\]](#).
- [49] A. Nagar and S. Akcay, Horizon-absorbed energy flux in circularized, nonspinning black-hole binaries and its effective-one-body representation, *Phys. Rev. D* **85**, 044025 (2012), [arXiv:1112.2840 \[gr-qc\]](#).
- [50] S. Bernuzzi, A. Nagar, and A. Zenginoglu, Horizon-absorption effects in coalescing black-hole binaries: An effective-one-body study of the non-spinning case, *Phys. Rev. D* **86**, 104038 (2012), [arXiv:1207.0769 \[gr-qc\]](#).
- [51] T. Damour, A. Nagar, and S. Bernuzzi, Improved effective-one-body description of coalescing nonspinning black-hole binaries and its numerical-relativity completion, *Phys. Rev. D* **87**, 084035 (2013), [arXiv:1212.4357 \[gr-qc\]](#).
- [52] T. Damour and A. Nagar, New effective-one-body description of coalescing nonprecessing spinning black-hole binaries, *Phys. Rev. D* **90**, 044018 (2014), [arXiv:1406.6913 \[gr-qc\]](#).
- [53] M. Boyle *et al.*, The SXS Collaboration catalog of binary black hole simulations, *Class. Quant. Grav.* **36**, 195006 (2019), [arXiv:1904.04831 \[gr-qc\]](#).
- [54] A. Buonanno, Y. Pan, H. P. Pfeiffer, M. A. Scheel, L. T. Buchman, and L. E. Kidder, Effective-one-body waveforms calibrated to numerical relativity simulations: Coalescence of non-spinning, equal-mass black holes, *Phys. Rev. D* **79**, 124028 (2009), [arXiv:0902.0790 \[gr-qc\]](#).
- [55] W. Tichy, E. E. Flanagan, and E. Poisson, Can the post-Newtonian gravitational wave form of an inspiraling binary be improved by solving the energy balance equation numerically?, *Phys. Rev. D* **61**, 104015 (2000), [arXiv:gr-qc/9912075](#).
- [56] P. Ajith, Addressing the spin question in gravitational-wave searches: Waveform templates for inspiralling compact binaries with nonprecessing spins, *Phys. Rev. D* **84**, 084037 (2011), [arXiv:1107.1267 \[gr-qc\]](#).
- [57] F. Pretorius, Evolution of binary black hole spacetimes, *Phys. Rev. Lett.* **95**, 121101 (2005), [arXiv:gr-qc/0507014](#).
- [58] M. Campanelli, C. O. Lousto, P. Marronetti, and Y. Zlochower, Accurate evolutions of orbiting black-hole binaries without excision, *Phys. Rev. Lett.* **96**, 111101 (2006), [arXiv:gr-qc/0511048](#).
- [59] J. G. Baker, J. Centrella, D.-I. Choi, M. Koppitz, and J. van Meter, Gravitational wave extraction from an inspiraling configuration of merging black holes, *Phys. Rev. Lett.* **96**, 111102 (2006), [arXiv:gr-qc/0511103](#).
- [60] H. P. Pfeiffer, Numerical simulations of compact object binaries, *Class. Quant. Grav.* **29**, 124004 (2012), [arXiv:1203.5166 \[gr-qc\]](#).
- [61] J. Centrella, J. G. Baker, B. J. Kelly, and J. R. van Meter, Black-hole binaries, gravitational waves, and numerical relativity, *Rev. Mod. Phys.* **82**, 3069 (2010), [arXiv:1010.5260 \[gr-qc\]](#).
- [62] A. Le Tiec, The Overlap of Numerical Relativity, Perturbation Theory and Post-Newtonian Theory in the Binary Black Hole Problem, *Int. J. Mod. Phys. D* **23**, 1430022 (2014), [arXiv:1408.5505 \[gr-qc\]](#).
- [63] M. Hannam, Modelling gravitational waves from precessing black-hole binaries: Progress, challenges and prospects, *Gen. Rel. Grav.* **46**, 1767 (2014), [arXiv:1312.3641 \[gr-qc\]](#).
- [64] M. A. Scheel, M. Boyle, T. Chu, L. E. Kidder, K. D. Matthews, and H. P. Pfeiffer, High-accuracy waveforms for binary black hole inspiral, merger, and ringdown, *Phys. Rev. D* **79**, 024003 (2009), [arXiv:0810.1767 \[gr-qc\]](#).
- [65] S. Ossokine, L. E. Kidder, and H. P. Pfeiffer, Precession-tracking coordinates for simulations of compact-object-binaries, *Phys. Rev. D* **88**, 084031 (2013), [arXiv:1304.3067 \[gr-qc\]](#).
- [66] M. A. Scheel, M. Giesler, D. A. Hemberger, G. Lovelace, K. Kuper, M. Boyle, B. Szilágyi, and L. E. Kidder, Improved methods for simulating nearly extremal binary black holes, *Class. Quant. Grav.* **32**, 105009 (2015), [arXiv:1412.1803 \[gr-qc\]](#).
- [67] G. B. Cook and B. F. Whiting, Approximate Killing Vectors on S^{**2} , *Phys. Rev. D* **76**, 041501 (2007), [arXiv:0706.0199 \[gr-qc\]](#).
- [68] D. Christodoulou and R. Ruffini, Reversible transformations of a charged black hole, *Phys. Rev. D* **4**, 3552 (1971).
- [69] I. MacDonald, S. Nissanke, H. P. Pfeiffer, and H. P. Pfeiffer, Suitability of post-Newtonian/numerical-relativity hybrid waveforms for gravitational wave detectors, *Class. Quant. Grav.* **28**, 134002 (2011), [arXiv:1102.5128 \[gr-qc\]](#).
- [70] T. Damour, A. Nagar, and M. Trias, Accuracy and effectualness of closed-form, frequency-domain waveforms for non-spinning black hole binaries, *Phys. Rev. D* **83**, 024006 (2011), [arXiv:1009.5998 \[gr-qc\]](#).
- [71] B. Bruegmann, J. A. Gonzalez, M. Hannam, S. Husa, U. Sperhake, and W. Tichy, Calibration of Moving Puncture Simulations, *Phys. Rev. D* **77**, 024027 (2008), [arXiv:gr-qc/0610128](#).
- [72] L. Lindblom, B. J. Owen, and D. A. Brown, Model Waveform Accuracy Standards for Gravitational Wave Data Analysis, *Phys. Rev. D* **78**, 124020 (2008), [arXiv:0809.3844 \[gr-qc\]](#).
- [73] A. Buonanno, B. Iyer, E. Ochsner, Y. Pan, and B. S. Sathyaprakash, Comparison of post-Newtonian templates for compact binary inspiral signals in gravitational-wave detectors, *Phys. Rev. D* **80**, 084043 (2009), [arXiv:0907.0700 \[gr-qc\]](#).
- [74] B. P. Abbott *et al.* (KAGRA, LIGO Scientific, Virgo, VIRGO), Prospects for observing and localizing gravitational-wave transients with Advanced LIGO, Advanced Virgo and KAGRA, *Living Rev. Rel.* **21**, 3 (2018), [arXiv:1304.0670 \[gr-qc\]](#).
- [75] G. Pratten *et al.*, Computationally efficient models for the dominant and subdominant harmonic modes of precessing binary black holes, *Phys. Rev. D* **103**, 104056 (2021), [arXiv:2004.06503 \[gr-qc\]](#).
- [76] X. Jiménez-Forteza, D. Keitel, S. Husa, M. Hannam, S. Khan, and M. Pürrer, Hierarchical data-driven approach to fitting numerical relativity data for nonprecessing binary black holes with an application to final spin and radiated energy, *Phys. Rev. D* **95**, 064024 (2017), [arXiv:1611.00332 \[gr-qc\]](#).

- [77] A. Nitz, I. Harry, D. Brown, C. M. Biwer, J. Willis, T. D. Canton, C. Capano, T. Dent, L. Pekowsky, S. De, M. Cabero, A. R. Williamson, B. Machenschalk, D. Macleod, G. S. C. Davies, P. Kumar, F. Pannarale, S. Reyes, dfinstad, S. Kumar, M. Tápai, L. Singer, S. Khan, S. Fairhurst, veronica villa, K. Chandra, S. Wu, A. Nielsen, S. Singh, and T. Massinger, [gwastro/pycbc: v2.0.6 release of pycbc](#) (2023).
- [78] P. Virtanen *et al.*, SciPy 1.0: Fundamental Algorithms for Scientific Computing in Python, *Nature Meth.* **17**, 261 (2020), [arXiv:1907.10121 \[cs.MS\]](#).
- [79] C. R. Harris *et al.*, Array programming with NumPy, *Nature* **585**, 357 (2020), [arXiv:2006.10256 \[cs.MS\]](#).
- [80] J. D. Hunter, Matplotlib: A 2D Graphics Environment, *Computing in Science & Engineering* **9**, 90 (2007).
- [81] W. R. Inc., *Mathematica, Version 13.3* (2023), Champaign, IL.
- [82] K. G. Arun, A. Buonanno, G. Faye, and E. Ochsner, Higher-order spin effects in the amplitude and phase of gravitational waveforms emitted by inspiraling compact binaries: Ready-to-use gravitational waveforms, *Phys. Rev. D* **79**, 104023 (2009), [Erratum: *Phys.Rev.D* 84, 049901 (2011)], [arXiv:0810.5336 \[gr-qc\]](#).
- [83] C. K. Mishra, A. Kela, K. G. Arun, and G. Faye, Ready-to-use post-Newtonian gravitational waveforms for binary black holes with nonprecessing spins: An update, *Phys. Rev. D* **93**, 084054 (2016), [arXiv:1601.05588 \[gr-qc\]](#).
- [84] M. Wade, J. D. E. Creighton, E. Ochsner, and A. B. Nielsen, Advanced LIGO's ability to detect apparent violations of the cosmic censorship conjecture and the no-hair theorem through compact binary coalescence detections, *Phys. Rev. D* **88**, 083002 (2013), [arXiv:1306.3901 \[gr-qc\]](#).
- [85] H. Nakano, T. Narikawa, K.-i. Oohara, K. Sakai, H.-a. Shinkai, H. Takahashi, T. Tanaka, N. Uchikata, S. Yamamoto, and T. S. Yamamoto, Comparison of various methods to extract ringdown frequency from gravitational wave data, *Phys. Rev. D* **99**, 124032 (2019), [arXiv:1811.06443 \[gr-qc\]](#).
- [86] M. Favata, C. Kim, K. G. Arun, J. Kim, and H. W. Lee, Constraining the orbital eccentricity of inspiralling compact binary systems with Advanced LIGO, *Phys. Rev. D* **105**, 023003 (2022), [arXiv:2108.05861 \[gr-qc\]](#).