



Testosterone is positively associated with coronary artery calcium in a low cardiovascular disease risk population

Benjamin C. Trumble^{1,*}, Jacob Negrey¹, Stephanie V. Koebele¹, Randall C. Thompson², L. Samuel Wann³, Adel H. Allam⁴, Bret Beheim⁵, M. Linda Sutherland⁶, James D. Sutherland⁶, Daniel Eid Rodriguez⁷, David E. Michalik^{8,9}, Chris J. Rowan¹⁰, Guido P. Lombardi¹¹, Angela R. Garcia¹, Daniel K. Cummings¹², Edmond Seabright¹³, Sarah Alami¹³, Thomas S. Kraft¹⁴, Paul Hooper¹², Kenneth Buetow¹, Andrei Irimia¹⁵, Margaret Gatz¹⁵, Jonathan Stieglitz^{16,#}, Michael D. Gurven^{17,#}, Hillard Kaplan^{12,#} and Gregory S. Thomas^{#,6,18}, Tsimane Health and Life History Project Teams[†]

¹Arizona State University, School of Human Evolution and Social Change, Center for Evolution and Medicine, Institute of Human Origins, Tempe, AZ, USA; ²Saint Luke's Mid America Heart Institute, Department of Cardiology, Kansas City, MO, USA; ³University of New Mexico, School of Medicine, Albuquerque, NM, USA; ⁴Al Azhar University, School of Medicine, Cairo, Egypt; ⁵Max Planck Institute for Evolutionary Anthropology, Department of Human Behavior, Ecology and Culture, Leipzig, Germany; ⁶MemorialCare Health System, Fountain Valley, CA, USA; ⁷Universidad de San Simón, Department of Medicine, Cochabamba, Bolivia; ⁸University of California Irvine, School of Medicine, Irvine, CA, USA; ⁹Miller Women's and Children's Hospital Long Beach, CA, USA; ¹⁰University of Nevada, School of Medicine, NV, USA; ¹¹Universidad Peruana Cayetano Heredia, Laboratorio de Paleopatología, Lima, Peru; ¹²Chapman University, Economic Science Institute, Orange, CA, USA; ¹³Mohammed VI Polytechnic University, School of Collective Intelligence, Ben Guerir, Morocco; ¹⁴University of Utah, Anthropology Department, Salt Lake City, UT, USA; ¹⁵University of Southern California, Psychology Department, Los Angeles, CA, USA; ¹⁶Toulouse School of Economics, Institute for Advanced Study Toulouse, Toulouse, France; ¹⁷University of California Santa Barbara, Department of Anthropology, Santa Barbara, CA, USA; ¹⁸University of California Irvine, Division of Cardiology, Orange, CA, USA
#Co-senior.

[†]For the most up-to-date group information, see <https://tsimane.anth.ucsb.edu/>.

*Corresponding author. School of Human Evolution and Social Change, Arizona State University, Center for Evolution and Medicine, Institute of Human Origins, 427 E Tyler Mall, LSC 218, Tempe AZ 85281, USA. Tel: +206-708-9012.

E-mail: trumble@asu.edu

Received 27 March 2023; revised version accepted 11 October 2023.

ABSTRACT

Background: In industrialized populations, low male testosterone is associated with higher rates of cardiovascular mortality. However, coronary risk factors like obesity impact both testosterone and cardiovascular outcomes. Here, we assess the role of endogenous testosterone on coronary artery calcium in an active subsistence population with relatively low testosterone levels, low cardiovascular risk and low coronary artery calcium scores.

Methodology: In this cross-sectional community-based study, 719 Tsimane forager-horticulturalists in the Bolivian Amazon aged 40+ years underwent computed tomography (49.8% male, mean age 57.6 years).

Results: Coronary artery calcium levels were low; 84.5% had no coronary artery calcium. Zero-inflated negative binomial models found testosterone was positively associated with coronary artery calcium for the full sample (Incidence Rate Ratio [IRR] = 1.477, 95% Confidence Interval [CI] 1.001–2.170, $P = 0.031$), and in a male-only subset (IRR = 1.532, 95% CI 0.993–2.360, $P = 0.053$). Testosterone was also positively associated with clinically relevant coronary atherosclerosis (calcium >100 Agatston units) in the full sample (Odds Ratio [OR] = 1.984, 95% CI 1.202–3.275, $P = 0.007$) and when limited to male-only sample (OR = 2.032, 95% CI 1.118–4.816, $P = 0.024$). Individuals with coronary artery calcium >100 had 20% higher levels of testosterone than those with calcium <100 ($t = -3.201$, $P = 0.007$).

Conclusions and Implications: Among Tsimane, testosterone is positively associated with coronary artery calcium despite generally low normal testosterone levels, minimal atherosclerosis and rare cardiovascular disease (CVD) events. Associations between low testosterone and CVD events in industrialized populations are likely confounded by obesity and other lifestyle factors.

LAY SUMMARY In sedentary urban male participants, low testosterone is associated with coronary heart disease. However, obesity can confound this association, resulting in lower testosterone and more heart disease. Among the physically active Tsimane of Bolivia, heart disease and obesity are rare, and male participants with higher testosterone had more atherosclerosis of their coronary arteries.

KEYWORDS: testosterone; cardiovascular disease; evolutionary mismatch; coronary artery calcium

INTRODUCTION

Testosterone plays an important role in male reproductive trade-offs across a number of species including humans. Testosterone is associated with energetically expensive muscle tissue [1–3], and during energetic stressors like illness or injury [4–6], high energetic expenditure [7], or reduced caloric intake [8, 9], testosterone declines rapidly within hours to days. Because of these energetic trade-offs, male participants in industrialized populations tend to have higher circulating testosterone than male age-matched subsistence populations with high parasite and pathogen loads [10–12]. However, in industrialized populations, increases in body fat with age result in the aromatization of testosterone to estradiol and declines in circulating testosterone with age [13]. As such, low endogenous male testosterone in industrialized populations is often associated with a higher risk of morbidity and all-cause mortality, including higher rates of cardiovascular disease (CVD) [14–21].

While low testosterone has been associated with cardiovascular morbidity [14, 15, 19–21], a causal relationship between low testosterone and atherosclerotic CVD has not been established [22] and remains controversial [18, 19]. Notably, most CVD data are drawn from industrialized populations that exhibit species-atypical (and perhaps maladaptive) lifestyles characterized by excessive sedentary inactivity [23, 24], diets high in saturated fats and in simple sugars, low in fiber [25, 26], with high rates of chronic disease [27]. The cumulative physiological disparities distinguishing industrialized from subsistence populations are reflected in testosterone levels: industrialized populations exhibit significantly higher testosterone levels in early adulthood

than do members of small-scale societies [10–12]. Yet, despite elevated levels of testosterone across much of adulthood, industrialized populations exhibit higher rates of CVD and associated risk factors than do small-scale societies [28, 29]. Given divergent patterns of testosterone secretion and CVD prevalence across populations, observed associations between testosterone and atherosclerotic CVD in industrialized populations may reflect a species-atypical pattern that blurs or obscures any mechanistic relationships between testosterone and coronary artery disease (CAD) progression.

Confounding factors such as obesity that are prevalent in industrialized societies make it difficult to isolate the impact of testosterone on CVD, as obesity is associated with both increased risk of CVD and low testosterone [13, 30]. In the Multiethnic Study of Atherosclerosis (MESA), lower male testosterone was associated with higher rates of hypertension, diabetes, higher body mass index and higher rates of atherosclerosis as measured by coronary artery calcium (CAC) [31]; see Fig. 1. Males in industrial populations with many chronic diseases from cancer, hyperuricemia, T2DM, high inflammation and CVD have lower circulating testosterone [14, 15, 18–20, 22, 32–36]. As these studies are observational in nature, it is not possible to assess causality, but it should be noted that there is also the potential for reverse causality; inflammatory processes that are involved in CVD and many chronic diseases can downregulate testosterone production [5, 6]. Adipose tissue is a significant source of inflammation in industrialized populations [37], sometimes referred to as sterile inflammation (as opposed to infectious-based inflammation [38]), and thus could present a multifactorial impact decreasing testosterone production, as well as increasing CVD risk [39].




Figure 1. Conceptual figure highlighting differences in testosterone physiology and the associations between testosterone and CVD risk in industrial and subsistence populations.

While observational studies suggest that lower testosterone is associated with CVD, testosterone supplementation showed little benefit [40] or even increased CAC following treatment [41]. Commonly reported associations between testosterone and CVD in relatively sedentary obese industrialized populations may be additionally confounded by lifestyle factors like diet or physical activity, where obesity is typically correlated with both low testosterone and CVD [13] (Fig. 1). While coronary atherosclerosis begins as a non-calcified plaque in the arterial wall, as a plaque matures calcium hydroxyapatite is deposited in the plaque and its presence in a coronary artery is diagnostic of atherosclerosis [42, 43]. Measurement of CAC by chest computed tomography scanning is often used to detect pre-clinical CAD and is a tool to assess the burden of coronary atherosclerosis [44]. Furthermore, CAC is correlated with coronary atherosclerotic burden, and is predictive of future cardiovascular events and all-cause mortality [45, 46].

Here, we test for a testosterone- CAD association in a physically active population with few CAD risk factors, but relatively low testosterone, compared to age-matched US males [47]. The Tsimane, a remote population of forager-horticulturalists in the Bolivian Amazon [48], show levels of testosterone that are generally in the low-to-normal range by US standards across their lifespan, with testosterone 37.3% lower than age-matched US males [47]. Unlike documented patterns in the USA and other high-income countries, male Tsimane displays relatively minimal age-related declines in testosterone [49],

perhaps due to low body fat deposition with age [50], which leads to lower testosterone in US males [18]. Body fat can also influence many immune processes, including inflammation, which can lead to lower testosterone; thus, we also examine associations between cytokines, body fat and testosterone. The Tsimane practice a physically intensive, traditional subsistence lifestyle based on small-scale horticulture, fishing, hunting and gathering [48], and are one of the most heart-healthy populations ever examined clinically, with minimal obesity and hypertension, low blood lipids, negligible atrial fibrillation and minimal CAC [51, 52].

METHODS

Coronary artery calcium

Tsimane adults ($n = 719$, 49.8% male) aged 40–94 years underwent thoracic non-contrast CT scanning using a 16-detector row, multi-slice scanner (GE Brightspeed, Milwaukee, WI). A licensed radiology technician acquired electrocardiogram-gated CT scans. All scans were supervised and reviewed by HORUS team cardiologists and radiologists. Calcium in the coronary arteries was scored using Siemens (Erlangen, Germany) Syngo.via calcium scoring semi-automatic software, which was then reviewed by a technician blind to participant data. Candidate lesions were reviewed and scored [53] using protocols similar to those employed in large population samples [54].

Since 2010, the Tsimane Health and Life History Project (THLHP) has conducted continuous epidemiological surveillance on >85% of the Tsimane population. Among adults over 40 years of age, clinical exams, biomarkers, demographic information and other interviews are collected annually [48]. Body fat is measured via bioelectric impedance (Tanita BC-1500), and age is verified through a multi-step process involving known dated events, birth cohort-rankings, demographic histories and census records [48]. For this study, 719 individuals were sampled from 1214 who met the inclusion criteria (40+ years of age and self-identifying as Tsimane). Age stratification of the sample ensures even decile representation, requiring sampling of all available individuals ≥ 60 years and a random subset of individuals 40–59 years [51]. Individuals over age 60, not sampled had either recently migrated following major flooding ($n = 49$), were presently engaged in horticultural labor or hunting ($n = 15$), or refused to participate ($n = 2$). Only one individual refused to participate because of poor health or inability to travel (recovering from a hernia surgery). To address potential bias, those who had computed tomography scans (CTs) were compared to those without CTs; there were no significant differences in sex ($P = 0.634$), systolic ($P = 0.301$) or diastolic ($P = 0.301$) blood pressure, or body fat ($P = 0.942$), indicating a representative sample of adults over 40 years of age.

Blood lipids and testosterone and immune markers

Fasting morning blood was drawn shortly after waking (6–8 am), and serum was stored in liquid nitrogen until transferred to the laboratory. Lipids were measured (Stat Fax 1908, Awareness Technology, Palm City, FL, USA) in the THLHP laboratory in San Borja, Beni, Bolivia, and testosterone and cytokines were measured in singlicate via enzyme immunoassay in the University of California Santa Barbara Biodemography lab [49]. Within (intra plate) and between (inter) plate coefficients of variation were 5.9% and 14.9% for the high controls, and 6.3% and 11.7% for the low controls, which were run in duplicate on all plates. Cytokines (interleukin [IL]-1B, IL-2, IL-4, IL-5, IL-6, IL-10, IL-13, granulocyte-macrophage colony-stimulating factor and interferon [IFN]-gamma INFG) were measured via Luminex multiplex (EMD Millipore, Darmstadt, Germany). Following the manufacturer's recommendation, calibrators were prepared with a serum matrix, and specimens below the limits of detection were assigned the lower limit of detection. Quality control specimens were within the expected ranges provided by the manufacturer, see Ref. [51].

Statistical approach

To evaluate the association between testosterone and CAC scores while adjusting for known covariates of Tsimane CAC

[51], we performed a multivariate zero-inflated negative binomial regression [55] to account for the large fraction of 'zero' CAC scores and over-dispersion of data. The model performs a logistic regression to test for factors that predict excess CAC absence, and a simultaneous negative binomial regression to examine factors associated with CAC scores (including zeros and positive scores). CAC >100 is considered moderate, clinically relevant atherosclerosis [51, 54]; additional logistic regressions examined the association between CAC >100 and testosterone with relevant control variables, but see limitations. Testosterone was log transformed for normality; other variables had normal distributions. Models were selected by Akaike's Information Criteria, starting from the base model from previously published work on Tsimane CAC [51]. Males and females differ in testosterone level and variation as well as CAC, resulting in significant heteroscedasticity (unequal variance across sex), necessitating separate analyses for males and females. It should be noted that testosterone had more explanatory value in explaining differences in CAC score than biological sex. US studies focus only on males so it is not possible to examine sex differences in the effects of testosterone across populations.

Institutional review board approval

Informed consent was collected at three levels: by the individual, by the community and by the Tsimane Gran Consejo (Tsimane governing body). The procedures were approved by the institutional review boards at the University of California, Santa Barbara and University of New Mexico, and Universidad San Simon Mayor, Cochabamba Bolivia.

Role of the funding source

This study was funded by the National Institute of Health, National Institute on Aging (NIH/NIA) 1R01-AG054442 and R01-AG024119. The funding organization played no role in the study design, methods, or analyses.

RESULTS

Risk factors

The Tsimane are a relatively lean population with low atherosclerotic risk factors, and in this sample of 719 adults, there are low levels of CAC through the eighth decade of life (Table 1). Despite no statin use, 0% of male participants, and only 0.9% of female participants had elevated total cholesterol (>240 mg/dL) while 13.8% of male and 15.6% of female participants had LDL greater than 130 mg/dL. HDL is relatively low in this population with 53.2% of male and 49.4% of female participants below 40


Table 1. Biometrics and CVD risk factors by sex for $n = 719$ Tsimane

Biomarker (mean)	Male	Male SD	Female	Female SD	P value
Coronary Artery Calcium score (AU)	11.3	43.1	6.9	57.1	0.239
Age, years	57.2	10.3	58.0	10.7	0.302
BMI, kg/m ²	24.1	2.9	24.0	4.0	0.777
Body fat, %	18.2	6.2	26.0	7.9	<0.001
Systolic BP, mmHg	117.3	12.4	115.0	12.7	0.015
Diastolic BP, mmHg	74.9	10.2	72.3	9.9	<0.001
Total testosterone, ng/dL	779.9	1034.1	297.8	370.6	<0.001
IL-5, pg/mL	2.5	2.5	3.0	6.2	0.145
IL-10, pg/mL	4.8	5.6	4.2	6.8	0.251
Cholesterol, mg/dL	148.7	31.8	152.4	28.9	0.105
LDL, mg/dL	96.0	31.0	101.8	30.5	0.016
HDL, mg/dL	40.2	8.1	38.8	7.4	0.017
Triglycerides, mg/dL	106.0	43.3	98.6	47.2	0.031
Glucose, mg/dL	79.1	11.8	78.9	10.3	0.900
% High LDL (>130 mg/dL)	13.8%		15.6%		0.484
% High cholesterol (>240 mg/dL)	0.0%		0.9%		0.083
% High triglycerides (>200 mg/dL)	3.3%		4.3%		0.752
% Low HDL (<40 mg/dL)	52.1%		57.6%		0.170
% High glucose (>125 mg/dL)	0.6%		0.0%		0.158
% Obese BMI (>30 kg/m ²)	3.3%		8.8%		0.002
% Hypertensive (Systolic >140 or Diastolic >90 mmHg)	6.2%		4.7%		0.484

AU, Agatston units; BMI, body mass index; BP, blood pressure; HDL, high-density lipoprotein; IL, interleukin; LDL, low-density lipoprotein; SD, standard deviation. Bold indicates statistically significant at $P < 0.05$.

mg/dL. Obesity is unusual with only 3.3% of male and 8.8% of female Tsimane presenting with BMI >30 kg/m². Previous studies report that testosterone is significantly lower (37.3% lower) among the Tsimane compared to aged-matched US males [47], and has a shallower rate of decline with age [47, 49].

Low levels of coronary artery calcium

As previously reported [51], the Tsimane have the lowest known levels of CAC for any population to date (Table 1). Only 15.5% of Tsimane presented with any CAC, while comparative samples of MESA data show more than 86% of Americans in the same age range have CAC. Similarly, only 2.7% of Tsimane had CAC scores >100 Agatston units (AU), while more than 50% of MESA participants had calcium scores >100 AU [56].

Testosterone and CAC

While CAC levels are extremely low in this population, higher testosterone was associated with higher CAC values for male and female participants who had CAC (IRR = 1.474, $P = 0.049$, Table 2; Fig. 2a). There was no interaction between sex and testosterone

on overall CAC extent, though males were more likely to have non-zero CAC. Male sex, older age, HDL and triglycerides all positively predicted CAC presence, while higher testosterone, age, body fat and lower IL-10 were associated with continuous CAC levels above zero (Table 2). When the model was limited to males only, testosterone did not significantly associate with the probability of presenting with CAC, but there was a positive trend between testosterone levels and the extent of CAC for those who had CAC (IRR = 1.532, $P = 0.054$, Table 2). Age, HDL and triglycerides all positively predicted CAC presence in male Tsimane, while age and IL-10 were associated with CAC extent (Table 2). A female-only subset found no significant effect of testosterone on CAC.

Testosterone and clinically relevant atherosclerosis

In addition to its association with subclinical CAC, testosterone was associated with a higher likelihood of having CAC >100 for both the full sample (OR = 1.984, $P = 0.007$, Pseudo $R^2 = 0.156$, Table 3), and when the sample was limited to males only (OR = 2.320, $P = 0.024$, Pseudo $R^2 = 0.1932$, Table 3). Males and females with CAC >100 had significantly higher testosterone than those

Table 2. Associations between testosterone and CAC

A. Zero-inflated negative binomial model predicting Tsimane CAC scores by log testosterone for both sexes together and individually

	IRR	P value	95% Conf. Int	IRR	P value	95% Conf. Int	IRR	P value	95% Conf. Int			
Predictor of CAC score, both sexes			Male only			Female only						
Log testosterone (pg/mL)	1.474	0.049	1.001	2.170	1.532	0.054	0.993	2.360	0.938	0.902	0.338	2.604
Age (years)	1.070	<0.001	1.032	1.109	1.071	0.002	1.026	1.117	1.047	0.127	0.987	1.110
Body fat (%)	1.060	0.017	1.011	1.112	1.015	0.685	0.945	1.090	1.121	0.020	1.018	1.234
IL-5 (pg/mL)	0.848	0.167	0.670	1.072	1.113	0.538	0.792	1.564	0.624	0.005	0.448	0.870
IL-10 (pg/mL)	0.924	0.042	0.856	0.997	0.897	0.010	0.826	0.974	0.950	0.637	0.768	1.175
Intercept	0.010	0.043	0.000	0.869	0.010	0.109	0.000	2.774	0.398	0.843	0.000	3521.9
Predictor of CAC absence (zero Inflation)			Male			Female						
Male	-1.243	<0.001	-1.841	-0.645								
Age (years)	-0.071	<0.001	-0.098	-0.045	-0.083	<0.001	-0.120	-0.045	-0.060	0.002	-0.098	-0.022
HDL (mg/dL)	-0.029	0.105	-0.063	0.006	-0.043	0.049	-0.085	0.000	-0.0001	0.996	-0.061	0.601
Triglycerides (mg/dL)	-0.009	0.007	-0.016	-0.003	-0.009	0.077	-0.019	0.001	-0.009	0.056	-0.018	0.0002
Intercept	8.434	0.000	5.987	10.880	8.493	<0.001	5.378	11.608	6.514	0.001	2.824	10.203

Zero-inflated negative binomial model predicting Tsimane CAC scores by log testosterone for both sexes.

AU, Agatston units; CAC, coronary artery calcium; HDL, high-density lipoprotein; IL, interleukin.




Figure 2. (A) Male CAC by age and tertile of testosterone. (B) Associations between testosterone and body fat differ for male Tsimane above and below median body fat. (C) Tsimane males have low body fat compared to age-matched US males (NHANES). Only 3.3% of US males fall below the median Tsimane body fat, and thus nearly all US males fall on the portion of the regression line where higher body fat is associated with low testosterone.

without high CAC (median testosterone with CAC >100 = 417 ng/dL, median testosterone with CAC <100 = 341 ng/dL; $t = -3.201, P = 0.0007$).

Testosterone and CAD risk factors

Both male and female Tsimane generally showed low levels of classical CVD risk factors (Table 1). However, testosterone had a

non-linear association with one CAD risk factor-body fat. Controlling for age, Tsimane male body fat was positively (though not significantly) associated with testosterone for those below the median body fat (17.6%), and negatively associated with testosterone for those above median body fat ($\beta = -0.05$, $P = 0.002$), Fig. 2B. Age matched representative data from US males (NHANES) show significantly higher male body fat; the median US male has a body fat of 28.9%, with only 3.3% of US males falling below the Tsimane median (Fig. 2C). In female Tsimane, testosterone was not associated with body fat or any other traditional risk factors.

Inflammation, testosterone and body fat

In a combined model with both sexes, those with higher log IL-2 had significantly lower levels of log testosterone ($\beta = -0.13$, $P = 0.002$) controlling for age, sex and body fat, following Bonferroni correction (see Supplementary Table S1). Body fat was not significantly associated with any cytokines in the full model.

In the male-only subset, those with higher log IL-2 and log IL-5 had lower log testosterone, but these results were not statistically significant following Bonferroni correction, (see Supplementary Table S2). Higher levels of log IL-1b were associated with higher body fat controlling for age and testosterone, but these results were not statistically significant following Bonferroni correction.

Comparison to clinical cutoffs

Various clinical cutoffs for 'low' male testosterone or clinical hypogonadism have been used in the past in industrialized

populations [57]. MESA studies used cutoffs of 300 ng/dL and 275 ng/dL, finding that US males below either of these cutoffs showed increased CAC [31]. For male Tsimane, having testosterone lower than these cutoffs was not associated with increased CAC (14.2% of male Tsimane in this sample had testosterone <300 ng/dL). On the contrary, participants below either of the MESA cutoffs had less CAC than those above the cutoffs. Similarly, male Tsimane below the clinical cutoff of hypogonadism of 230 ng/dL also had lower CAC scores than those with higher CAC [31]. Table 4 shows CAC scores by tercile for Tsimane testosterone.

DISCUSSION

The Tsimane population offers a unique test of the association between endogenous testosterone and coronary calcium with low levels of confounding CVD risk factors like obesity seen in industrialized populations. While no single population is a perfect exemplar of the myriad of environments in which humans evolved, sedentary urban environments are evolutionarily novel and not a good representation of what healthy aging was like prior to the last several hundred years. Most medical research is conducted in these sedentary environments, and thus our understanding of associations between cardiovascular disease and testosterone may be skewed. By working with populations that still have a physically active subsistence life style, we can better understand how human physiology is associated with healthy aging, without the confounding of obesity and other related



Table 3. Logistic regression models testing association between log testosterone and clinically relevant atherosclerosis (CAC >100 AU)

CAC >100 AU both sexes	CAC >100 AU male only			CAC >100 AU female only					
	OR	P value	95% Conf. Int	OR	P value	95% Conf. Int	OR	P value	95% Conf. Int
Log testosterone (pg/mL)	1.984	0.007	1.202 3.275	2.320	0.024	1.118 4.816	1.212	0.700	0.456 3.220
Age (years)	1.010	<0.001	1.056 1.138	1.114	<0.001	1.060 1.171	1.076	0.022	1.011 1.145



Table 4. Testosterone in males and CAC score by testosterone tercile (low, middle, high)

	Low tercile <i>T n</i> = 114	Middle tercile <i>T n</i> = 114	High tercile <i>T n</i> = 115
Median testosterone ng/dL	322.7	513.0	931.4
Median age	60	52	52.5
Mean CAC	7.5	13.7	3.6
% CAC > 0 AU	27.4%	13.4%	21.4%
% CAC > 100 AU	4.4%	2.6%	5.3%

AU, Agatston units; CAC, coronary artery calcium.

factors. Though studies from industrial urban centers like MESA report higher coronary calcium in male participants with low testosterone [31], we find that male Tsimane with higher testosterone have *more* coronary calcium, while there is no relationship between testosterone and coronary calcium in female Tsimane participants (note: MESA investigators did not describe associations between testosterone in female participants and CAC) [7]. While high levels of CAC indicative of clinically relevant atherosclerosis are roughly 10-fold lower among Tsimane than in the USA [51], Tsimane with higher circulating testosterone were 2.5 times more likely to exhibit CAC scores over 100 than Tsimane with low testosterone.

Despite having one-third lower levels of testosterone than age-matched US males, even Tsimane with the lowest levels of testosterone were not at risk of atherosclerosis. In fact, not only does Tsimane have low levels of testosterone and low CVD risk, but Tsimane with higher levels of testosterone actually present with higher levels of atherosclerosis, and have a higher risk of clinically relevant CAC scores. These results call into question whether low testosterone plays a causal role in atherosclerosis, or whether the association between low testosterone and CVD is indirect. While male participants in industrial populations with low testosterone often are obese or have other cardiovascular risk factors such as diabetes [18, 30, 31], low testosterone was not associated with traditional cardiovascular risk factors in the Tsimane. The complex and often inconsistent results available in the testosterone and cardiovascular morbidity literature may stem from heterogeneity in underlying levels of CVD risk factors. This study is unique in that the Tsimane show exceptionally low levels of CVD risk factors, making it an ideal population in which to measure the effects of testosterone on CAD without confounding other risk factors.

Aromatization of testosterone by adipose tissue has been linked to the lower circulating levels of testosterone in obese male participants in industrialized populations [13], and obesity can alter the gut microbiome in ways that impact steroid hormone concentrations as well [58]. However, males in industrialized populations have significantly higher body fat (median US male body fat is 48.6% higher than median Tsimane body fat). We speculate that if US males fell within the range of body fat observed in Tsimane males, we would not see a purely detrimental impact of body fat on testosterone (Fig. 2B and C). Indeed, among male Olympians, androgen levels were significantly lower in athletes considered 'lean' (body fat 11.7% SD \pm 3.4%) versus 'nonlean' (body fat 16.4% SD \pm 5.8%) [59]. Individuals in a depleted energetic condition are likely to have downregulated testosterone and low body fat. Those in better energetic condition can maintain higher testosterone and higher body fat, but once body fat levels get above a threshold, testosterone will begin to decrease as it is aromatized into estradiol. Among the

Tsimane, high body fat is negatively associated with testosterone while there is no association between testosterone and below median levels of body fat. Nearly all US males have high body fat by Tsimane standards, which may help explain why testosterone is consistently negatively associated with body fat in industrialized populations (Fig. 2B and C). The relatively low levels of testosterone among Tsimane may instead be due to their high pathogen burden and low food security as a result of energetic tradeoffs [48, 60]. It is physiologically expensive to maintain high levels of testosterone, and males experiencing illness [5], injury [6] or sustained caloric deficiency show rapid, substantive declines in testosterone [7]. Similarly, low levels of testosterone have been reported among other subsistence populations facing high pathogen burden [11]. Notably, due to relatively low testosterone, Tsimane have relatively small prostate volumes compared to other populations with higher testosterone, resulting in some of the lowest population levels of benign prostatic hyperplasia [49]. Yet, similar to what is reported in the USA, we find a positive association between testosterone and cognitive function among the Tsimane [61]. We suspect that in these populations and in similar contexts, lower testosterone may also be associated with lower, not greater, atherosclerotic burden.

We did not find strong associations between inflammation and testosterone in this sample. Of nine cytokines measured, only IL-2 was significantly associated with testosterone, with higher IL-2 predicting lower levels of testosterone. This finding fits with the experimental literature where IL-2 infusions decrease testosterone production in human males in industrialized populations [62]. While higher levels of inflammation are often associated with low testosterone in industrialized populations, we do not find the same associations here, nor do we see that higher adiposity is associated with higher inflammation [37]. The lack of association between inflammation and body fat may be because of differences in the type of inflammation; while individuals in industrialized populations experience sterile inflammation from obesity, smoking and pollution, the Tsimane are experiencing high inflammation from parasites and pathogens [37, 38, 60].

To date, the causal role of testosterone in cardiovascular health has not been well established [63]; and it should be noted that we are only reporting relationships in this study and not causal associations. Studies have reported associations between low testosterone and CVD risk factors, cardiovascular disease rates (as measured by CAC), percutaneous coronary intervention-related major adverse cardiac events, and all-cause mortality [14, 15, 18–21, 31]. There are several biologically plausible mechanisms by which exogenous testosterone could be cardioprotective either directly or by reducing CVD risk factors [22, 64]; exogenous testosterone increases exercise tolerance in males with stable angina [64], increases coronary blood flow [65], and cardiac output [65, 66], and is associated with decreased reperfusion injury

[66]. However, testosterone therapy may come at a health risk; it is associated with an increase in myocardial infarction [67, 68], increased risk of thrombosis [69], and higher rates of stroke in most [70], though not all, studies [71]. Testosterone administration is associated with an increase in total plaque volume but does not appear to impact CAC deposition over a one-to-three-year period [40, 41]. Thus, testosterone administration may have both harmful effects with an increased rate of plaque formation and thrombosis risk as well as positive effects on arterial elasticity and cardiac perfusion [22, 68, 72, 73].

This creates an apparent contradiction where low endogenous testosterone is associated with higher rates of CAD [14, 15, 18–21, 31], but exogenous testosterone can also have deleterious cardiovascular effects [40, 41, 67–70]. This may be explained by the underlying cause of low endogenous testosterone in males with CVD: males with higher body fat have lower testosterone due to aromatization to estradiol [13]. Thus, obesity and its related disease states, not circulating testosterone, are likely responsible for poor CVD outcomes in males with low testosterone. In the USA, males taking supplemental testosterone can maintain both high body fat and high testosterone, an unnatural and potentially risk-inducing combination if both testosterone and obesity independently contribute to CVD.

These contradictory impacts of testosterone on cardiovascular risk could explain why the testosterone supplementation literature shows some positive impacts of testosterone [22, 64], despite overall negative impacts of testosterone supplementation on cardiovascular morbidity and mortality [68, 73]. There has been a large increase in the number of prescriptions for testosterone supplementation over the last decade: nearly a 10-fold increase in the USA and 100-fold increase worldwide [63, 74]. While existing data may not yet be sufficient to assess the effects of testosterone on CVD, new data from the large numbers of US males now taking exogenous testosterone may illuminate additional pathways and mechanisms.

LIMITATIONS

Most medical research is conducted in sedentary urban populations, and while this study adds a very underrepresented type of population, there are limits to the generalizability of these results. While the Tsimane have relatively low CVD risk factors, and while that allows us to examine associations between testosterone and CVD without certain confounders, the high parasite and pathogen burden [60], and differential genetic structure of the population [75] may limit cross-population comparisons. A relatively small proportion of this sample had CAC scores over 100 AU (2.7% of the total sample, $n = 20$ individuals, including 14 male and 6 female participants). Thus, these results (Table 3) should be interpreted with caution. Due to limitations in sample

volume and cost, testosterone and cytokines were run in single, though assay standards and controls and run in duplicate to assess assay functionality. This could potentially add additional random variation to the sample.

We used a single testosterone measurement, which may not be indicative of an individual's entire lifetime exposure to testosterone, as testosterone can fluctuate rapidly under multiple types of stressors [5–8]. That said, testosterone is relatively stable with age in this population [76, 77], and specimens collected 2 years apart ($n = 135$) had an intraclass correlation (ICC) of 0.45, indicating relatively fair stability across time points with approximately half of the variability due to differences between individuals. Our use of single-point measures of testosterone is comparable to what is reported in other studies in industrialized populations [14–16, 19, 20, 31]. We measured total testosterone, but not sex hormone binding globulin or albumin, and thus cannot assess how much of that testosterone was biologically available or bound to carrier proteins. Enzyme immunoassays designed to measure just free testosterone are not considered accurate [78], and the Endocrine Society position statement recommends serum total testosterone measurements for both males and females [78]. Total testosterone has been measured in other testosterone-CVD studies in industrialized populations [14–16, 19, 20, 31].

CONCLUSIONS

While prior studies have shown that low levels of serum testosterone are associated with increased rates of coronary artery calcification, we found the opposite effect in a population with very low rates of atherosclerosis risk factors and low rates of CAC. Our findings suggest that the previously described association between CAC and low testosterone is indirect rather than causal.

SUPPLEMENTARY DATA

Supplementary data is available at *EMPH* online.

ACKNOWLEDGMENTS

We thank the HORUS and THLHP teams and our collaborators at the Hospital Presidente German Busch of Trinidad, Bolivia, including Edila Arteaga and Editha Alpire and CT core lab leader James K. Min and Kimberly Elmore of the Dalio Institute of Cardiovascular Imaging, Weill Cornell Medical College; JSacknowledges support from the Agence Nationale de la Recherche—Labex Institute of Advanced Study in Toulouse. National Institute on Aging within the National Institute of Health (R01AG054442) National Science Foundation (1748282). National Institute on

Aging within the National Institute of Health (R01AG054442) National Science Foundation (1748282).

AUTHOR CONTRIBUTIONS

Benjamin Trumble (Conceptualization [Lead], Data curation [Lead], Formal analysis [Lead], Funding acquisition [Equal], Investigation [Equal], Methodology [Equal], Project administration [Equal], Resources [Equal], Software [Equal], Supervision [Lead], Validation [Equal], Visualization [Lead], Writing—original draft [Lead], Writing—review & editing [Lead]), Jacob Negrey (Conceptualization [Equal], Writing—review & editing [Equal]), Stephanie Koebele (Conceptualization [Equal], Investigation [Equal], Writing—review & editing [Equal]), randall Thompson (Conceptualization [Equal], Funding acquisition [Equal], Investigation [Equal], Methodology [Equal], Writing—review & editing [Equal]), L. Samuel Wann (Conceptualization [Equal], Funding acquisition [Equal], Investigation [Equal], Methodology [Equal], Writing—review & editing [Equal]), Adel Allam (Conceptualization [Equal], Investigation [Equal], Methodology [Equal], Writing—review & editing [Equal]), Bret Beheim (Conceptualization [Equal], Data curation [Equal], Investigation [Equal], Methodology [Equal], Project administration [Equal], Writing—review & editing [Equal]), M. Linda Sutherland (Data curation [Equal], Investigation [Equal], Methodology [Equal], Project administration [Equal], Supervision [Equal], Validation [Equal], Visualization [Equal], Writing—review & editing [Equal]), James D. Sutherland (Data curation [Equal], Investigation [Equal], Methodology [Equal], Project administration [Equal], Supervision [Equal], Validation [Equal], Visualization [Equal], Writing—review & editing [Equal]), Daniel Eid Rodriguez (Conceptualization [Equal], Funding acquisition [Equal], Investigation [Equal], Methodology [Equal], Project administration [Equal], Supervision [Equal], Writing—review & editing [Equal]), David Michalik (Investigation [Equal], Methodology [Equal], Writing—review & editing [Equal]), Chris J. Rowan (Investigation [Equal], Methodology [Equal], Writing—review & editing [Equal]), Guido Lombardi (Investigation [Equal], Methodology [Equal], Project administration [Equal], Writing—review & editing [Equal]), Angela Garcia (Conceptualization [Equal], Funding acquisition [Equal], Investigation [Equal], Methodology [Equal], Project administration [Equal], Writing—review & editing [Equal]), Daniel Cummings (Conceptualization [Equal], Data curation [Equal], Investigation [Equal], Methodology [Equal], Project administration [Equal], Writing—review & editing [Equal]), Edmond Seabright (Data curation [Equal], Investigation [Equal], Methodology [Equal], Project administration [Equal], Writing—review & editing [Equal]), Sarah Alami (Data curation [Equal], Investigation [Equal], Methodology [Equal], Project administration [Equal], Writing—review & editing [Equal]), Thomas Kraft (Data curation [Equal], Investigation

[Equal], Methodology [Equal], Writing—review & editing [Equal]), Paul Hooper (Conceptualization [Equal], Data curation [Equal], Investigation [Equal], Methodology [Equal], Project administration [Equal], Writing—review & editing [Equal]), Kenneth Buetow (Conceptualization [Equal], Funding acquisition [Equal], Project administration [Equal], Writing—review & editing [Equal]), Andrei Irimia (Conceptualization [Equal], Funding acquisition [Equal], Investigation [Equal], Methodology [Equal], Validation [Equal], Visualization [Equal], Writing—review & editing [Equal]), Margaret Gatz (Conceptualization [Equal], Funding acquisition [Equal], Investigation [Equal], Methodology [Equal], Project administration [Equal], Writing—review & editing [Equal]), Jonathan Stieglitz (Conceptualization [Equal], Funding acquisition [Equal], Investigation [Equal], Methodology [Equal], Project administration [Equal], Writing—review & editing [Equal]), Michael Gurven (Conceptualization [Equal], Funding acquisition [Equal], Investigation [Equal], Methodology [Equal], Project administration [Equal], Writing—review & editing [Equal]), Hillard Kaplan (Conceptualization [Equal], Data curation [Equal], Funding acquisition [Equal], Investigation [Equal], Methodology [Equal], Project administration [Lead], Writing—review & editing [Equal]), and Gregory Thomas (Conceptualization [Equal], Funding acquisition [Equal], Investigation [Equal], Methodology [Equal], Project administration [Equal], Writing—review & editing [Equal])

CONFLICT OF INTERESTS

No conflicts of interest to disclose.

DATA AVAILABILITY

Individual-level data are stored in the Tsimane Health and Life History Project (THLHP) Data Repository and are available through restricted access for ethical reasons. THLHP's highest priority is the safeguarding of human subjects and minimization of risk to study participants. The THLHP adheres to the 'CARE Principles for Indigenous Data Governance' (Collective Benefit, Authority to Control, Responsibility, and Ethics), which assure that the Tsimane (i) have sovereignty over how data are shared, (ii) are the primary gatekeepers determining ethical use, (iii) are actively engaged in the data generation and (iv) derive benefit from data generated and shared for use whenever possible. The THLHP is also committed to the 'FAIR Guiding Principles for scientific data management and stewardship' (Findable, Accessible, Interoperable, Reusable). Requests for individual-level data should take the form of an application that details the exact uses of the data and the research questions to be addressed, procedures that will be employed for data security and individual privacy, potential benefits to the study communities, and

procedures for assessing and minimizing stigmatizing interpretations of the research results (see the following webpage for links to the data sharing policy and data request forms: <https://tsimane.anth.ucsb.edu/data.html>). Requests for individual-level data will require institutional IRB approval (even if exempt) and will be reviewed by an Advisory Council composed of Tsimane community leaders, community members, Bolivian scientists, and the THLHP leadership. The study authors and the THLHP leadership are committed to open science and are available to assist interested investigators in preparing data access requests.

REFERENCES

- Bribiescas RG. An evolutionary and life history perspective on human male reproductive senescence. *Ann N Y Acad Sci* 2010;**1204**:54–64. DOI: [10.1111/j.1749-6632.2010.05524.x](https://doi.org/10.1111/j.1749-6632.2010.05524.x).
- Falquetto H, Júnior JL, Silvério MN *et al.* Can conditions of skeletal muscle loss be improved by combining exercise with anabolic–androgenic steroids? A systematic review and meta-analysis of testosterone-based interventions. *Rev Endocr Metab Disord* 2021;**22**:161–78.
- Trumble BC, Pontzer H, Stieglitz J *et al.* Energetic costs of testosterone in two subsistence populations. *Am J Hum Biol* 2023;**35**:e23949.
- Muehlenbein MP, Hirschtick JL, Bonner JZ *et al.* Toward quantifying the usage costs of human immunity: altered metabolic rates and hormone levels during acute immune activation in men. *Am J Hum Biol* 2010;**22**:546–56. DOI: [10.1002/ajhb.21045](https://doi.org/10.1002/ajhb.21045).
- Spratt DI, Cox P, Orav J *et al.* Reproductive axis suppression in acute illness is related to disease severity. *J Clin Endocrinol Metab* 1993;**76**:1548–54. DOI: [10.1210/jcem.76.6.8501163](https://doi.org/10.1210/jcem.76.6.8501163).
- Spratt DI, Kramer RS, Morton JR *et al.* Characterization of a prospective human model for study of the reproductive hormone responses to major illness. *Am J Physiol Endocrinol Metab* 2008;**295**:E63–9. DOI: [10.1152/ajpendo.00472.2007](https://doi.org/10.1152/ajpendo.00472.2007).
- Nindl BC, Barnes BR, Alemany JA *et al.*; Physiological consequences of U.S. Army Ranger training. *Med Sci Sports Exerc* 2007;**39**:1380–7.
- Trumble BC, Brindle E, Kupsik M *et al.* Responsiveness of the reproductive axis to a single missed evening meal in young adult males. *Am J Hum Biol* 2010;**22**:775–81.
- Cameron JL, Weltzin TE, McConaha C *et al.* Slowing of pulsatile luteinizing hormone secretion in men after forty-eight hours of fasting. *J Clin Endocrinol Metab* 1991;**73**:35–41. DOI: [10.1210/jcem-73-1-35](https://doi.org/10.1210/jcem-73-1-35).
- Trumble BC, Stieglitz J, Rodriguez DE *et al.* Challenging the inevitability of prostate enlargement: low levels of benign prostatic hyperplasia among Tsimane Forager-Horticulturalists. *J Gerontol A Biol Sci Med Sci* 2015;**70**:1262–8.
- Ellison PT, Bribiescas RG, Bentley GR *et al.* Population variation in age-related decline in male salivary testosterone. *Hum Reprod* 2002;**17**:3251–3. DOI: [10.1093/humrep/17.12.3251](https://doi.org/10.1093/humrep/17.12.3251).
- Jasienska G, Bribiescas RG, Furberg A-S *et al.* Human reproduction and health: an evolutionary perspective. *Lancet (London, England)* 2017;**390**:510–20.
- Gates MA, Mekary RA, Chiu GR *et al.* Sex steroid hormone levels and body composition in men. *J Clin Endocrinol Metab* 2013;**98**:2442–50. DOI: [10.1210/jc.2012-2582](https://doi.org/10.1210/jc.2012-2582).
- Ohlsson C, Barrett-Connor E, Bhasin S *et al.* High serum testosterone is associated with reduced risk of cardiovascular events in elderly men. The MrOS (Osteoporotic Fractures in Men) study in Sweden. *J Am Coll Cardiol* 2011;**58**:1674–81. DOI: [10.1016/j.jacc.2011.07.019](https://doi.org/10.1016/j.jacc.2011.07.019).
- Khaw K-T, Dowsett M, Folkard E *et al.* Endogenous testosterone and mortality due to all causes, cardiovascular disease, and cancer in men: European prospective investigation into cancer in Norfolk (EPIC-Norfolk) Prospective Population Study. *Circulation* 2007;**116**:2694–701. DOI: [10.1161/circulationaha.107.719005](https://doi.org/10.1161/circulationaha.107.719005).
- Laughlin GA, Barrett-Connor E, Bergstrom J. Low serum testosterone and mortality in older men. *J Clin Endocrinol Metab* 2008;**93**:68–75. DOI: [10.1210/jc.2007-1792](https://doi.org/10.1210/jc.2007-1792).
- Maggio M, Lauretani F, Ceda GP *et al.* Relationship between low levels of anabolic hormones and 6-year mortality in older men: the aging in the Chianti Area (InCHIANTI) study. *Arch Intern Med* 2007;**167**:2249–54. DOI: [10.1001/archinte.167.20.2249](https://doi.org/10.1001/archinte.167.20.2249).
- Muehlenbein MP, Gassen J, Shattuck EC, *et al.* Lower testosterone levels are associated with higher risk of death in men. *Evol Med Public Health* 2022;**11**:30–40. DOI: [10.1093/emph/eoac044](https://doi.org/10.1093/emph/eoac044).
- Gettler LT, Sarma MS, Gengo RG *et al.* Adiposity, CVD risk factors, and testosterone: variation by partnering status and residence with children in US men. *Evol Med Public Health* 2017;**2017**:67–80. DOI: [10.1093/emph/eox005](https://doi.org/10.1093/emph/eox005).
- Lopez DS, Taha S, Gutierrez S *et al.* Association of total and free testosterone with cardiovascular disease in a nationally representative sample of white, black, and Mexican American men. *Int J Impot Res* 2022:1–9. DOI: [10.1038/s41443-022-00660-7](https://doi.org/10.1038/s41443-022-00660-7).
- Chiang C-H, Hung W-T, Liu E-S *et al.* The influence of testosterone on the risk of cardiovascular events after percutaneous coronary intervention. *Front Cardiovasc Med* 2022;**9**:998056. DOI: [10.3389/fcvm.2022.998056](https://doi.org/10.3389/fcvm.2022.998056).
- Kloner RA, Carson C, Dobs A *et al.* Testosterone and cardiovascular disease. *J Am Coll Cardiol* 2016;**67**:545–57.
- Gurven M, Jaeggi AV, Kaplan H, *et al.* Physical activity and modernization among Bolivian Amerindians. *PLoS One* 2013;**8**:e55679. DOI: [10.1371/journal.pone.0055679](https://doi.org/10.1371/journal.pone.0055679).
- Raichlen DA, Pontzer H, Zderic TW *et al.* Sitting, squatting, and the evolutionary biology of human inactivity. *Proc Natl Acad Sci USA* 2020;**117**:7115–21. DOI: [10.1073/pnas.1911868117](https://doi.org/10.1073/pnas.1911868117).
- Cordain L, Eaton SB, Sebastian A *et al.* Origins and evolution of the Western diet: health implications for the 21st century. *Am J Clin Nutr* 2005;**81**:341–54. DOI: [10.1093/ajcn.81.2.341](https://doi.org/10.1093/ajcn.81.2.341).
- Kraft TS, Stieglitz J, Trumble BC *et al.* Nutrition transition in 2 lowland Bolivian subsistence populations. *Am J Clin Nutr* 2018;**108**:1183–95. DOI: [10.1093/ajcn/nqy250](https://doi.org/10.1093/ajcn/nqy250).
- Trumble BC, Schneider-Crease I. Chronic diseases of aging in an evolutionary context. *Evol Med Public Health* 2020;**2020**:84–5. DOI: [10.1093/emph/eoaa013](https://doi.org/10.1093/emph/eoaa013).
- Kaplan H, Thompson RC, Trumble BC *et al.* Coronary atherosclerosis in indigenous South American Tsimane: a cross-sectional cohort study. *Lancet (London, England)* 2017;**389**:1730–9. DOI: [10.1016/S0140-6736\(17\)30752-3](https://doi.org/10.1016/S0140-6736(17)30752-3).
- Raichlen DA, Pontzer H, Harris JA *et al.* Physical activity patterns and biomarkers of cardiovascular disease risk in hunter-gatherers. *Am J Hum Biol* 2017;**29**:e22919. DOI: [10.1002/ajhb.22919](https://doi.org/10.1002/ajhb.22919).
- Wang C, Jackson G, Jones TH *et al.* Low testosterone associated with obesity and the metabolic syndrome contributes to sexual dysfunction

- and cardiovascular disease risk in men with type 2 diabetes. *Diabetes Care* 2011;**34**:1669–75. DOI: [10.2337/dc10-2339](https://doi.org/10.2337/dc10-2339).
31. Khazai B, Golden SH, Colangelo LA *et al.* Association of endogenous testosterone with subclinical atherosclerosis in men: the multi-ethnic study of atherosclerosis. *Clin Endocrinol (Oxf)* 2016;**84**:700–7. DOI: [10.1111/cen.12997](https://doi.org/10.1111/cen.12997).
 32. Svartberg J, Midtby M, Bonna KH *et al.* The associations of age, lifestyle factors and chronic disease with testosterone in men: the Tromsø Study. *Eur J Endocrinol* 2003;**149**:145–52. DOI: [10.1530/eje.0.1490145](https://doi.org/10.1530/eje.0.1490145).
 33. Grossmann M. Low testosterone in men with type 2 diabetes: significance and treatment. *J Clin Endocrinol Metab* 2011;**96**:2341–53. DOI: [10.1210/jc.2011-0118](https://doi.org/10.1210/jc.2011-0118).
 34. Rastrelli G, Corona G, Maggi M. Both comorbidity burden and low testosterone can explain symptoms and signs of testosterone deficiency in men consulting for sexual dysfunction. *Asian J Androl* 2020;**22**:265–73. DOI: [10.4103/aja.aja_61_19](https://doi.org/10.4103/aja.aja_61_19).
 35. Erenpreiss J, Fodina V, Pozarska R *et al.* Prevalence of testosterone deficiency among aging men with and without morbidities. *Aging Male* 2020;**23**:901–5. DOI: [10.1080/13685538.2019.1621832](https://doi.org/10.1080/13685538.2019.1621832).
 36. Tsai M-K, Hung K-C, Liao C-C *et al.* The Association between serum testosterone and hyperuricemia in males. *J Clin Med* 2022;**11**:2743.
 37. Calder PC, Ahluwalia N, Brouns F *et al.* Dietary factors and low-grade inflammation in relation to overweight and obesity. *Br J Nutr* 2011;**106**:S1–S78. DOI: [10.1017/S0007114511005460](https://doi.org/10.1017/S0007114511005460).
 38. Garcia AR, Finch C, Gatz M *et al.* APOE4 is associated with elevated blood lipids and lower levels of innate immune biomarkers in a tropical Amerindian subsistence population. *eLife* 2021;**10**:e68231. DOI: [10.7554/eLife.68231](https://doi.org/10.7554/eLife.68231).
 39. Cordain L, Eaton SB, Sebastian A *et al.* Origins and evolution of the Western diet: health implications for the 21st century1,2. *Am J Clin Nutr* 2005;**81**:341–54. DOI: [10.1093/ajcn.81.2.341](https://doi.org/10.1093/ajcn.81.2.341).
 40. Basaria S, Harman S, Travison TG *et al.* Effects of testosterone administration for 3 years on subclinical atherosclerosis progression in older men with low or low-normal testosterone levels: a randomized clinical trial. *JAMA* 2015;**314**:570–81. DOI: [10.1001/jama.2015.8881](https://doi.org/10.1001/jama.2015.8881).
 41. Budoff MJ, Ellenberg SS, Lewis CE *et al.* Testosterone treatment and coronary artery plaque volume in older men with low testosterone. *JAMA* 2017;**317**:708–16. DOI: [10.1001/jama.2016.21043](https://doi.org/10.1001/jama.2016.21043).
 42. Agatston AS, Janowitz WR, Hildner FJ *et al.* Quantification of coronary artery calcium using ultrafast computed tomography. *J Am Coll Cardiol* 1990;**15**:827–32.
 43. Stary HC, Chandler AB, Dinsmore RE *et al.* A definition of advanced types of atherosclerotic lesions and a histological classification of atherosclerosis: a report from the Committee on Vascular Lesions of the Council on Arteriosclerosis, American Heart Association. *Circulation* 1995;**92**:1355–74.
 44. Rumberger JA, Sheedy PF III, Breen JF *et al.* Coronary calcium, as determined by electron beam computed tomography, and coronary disease on arteriogram: effect of patient's sex on diagnosis. *Circulation* 1995;**91**:1363–7.
 45. Detrano R, Guerci AD, Carr JJ *et al.* Coronary calcium as a predictor of coronary events in four racial or ethnic groups. *N Engl J Med* 2008;**358**:1336–45.
 46. Nakanishi R, Li D, Blaha MJ *et al.* All-cause mortality by age and gender based on coronary artery calcium scores. *Eur Heart J* 2016;**17**:1305–14.
 47. Trumble BC, Cummings D, von Rueden C *et al.* Physical competition increases testosterone among Amazonian forager-horticulturalists: a test of the 'challenge hypothesis'. *Proc R Soc B Biol Sci* 2012;**279**:2907–12. DOI: [10.1098/rspb.2012.0455](https://doi.org/10.1098/rspb.2012.0455).
 48. Gurven M, Stieglitz J, Trumble BC *et al.* The Tsimane Health and Life History Project: integrating anthropology and biomedicine. *Evol Anthropol* 2017;**26**:54–73.
 49. Trumble BC, Stieglitz J, Rodriguez DE *et al.* Challenging the inevitability of prostate enlargement: low levels of benign prostatic hyperplasia among Tsimane forager-horticulturalists. *J Gerontol A Biol Sci Med Sci* 2015;**70**:1262–8. DOI: [10.1093/geronol/glv051](https://doi.org/10.1093/geronol/glv051).
 50. Kaplan H, Gatz M, Mack WJ *et al.* Brain volume, energy balance and cardiovascular health in two non-industrial South American populations. *Proc Natl Acad Sci USA* 2023;**120**:e2205448120.
 51. Kaplan H, Thompson R, Trumble BC *et al.* Indigenous South American Tsimane demonstrate the lowest levels of coronary atherosclerosis. *Lancet* 2017;**S0140**:30752–3.
 52. Rowan CJ, Eskander MA, Seabright E *et al.* Very low prevalence and incidence of atrial fibrillation among Bolivian forager-farmers. *Ann Global Health* 2021;**87**:18.
 53. Agatston AS, Janowitz WR, Hildner FJ *et al.* Quantification of coronary artery calcium using ultrafast computed tomography. *J Am Coll Cardiol* 1990;**15**:827–32. DOI: [10.1016/0735-1097\(90\)90282-t](https://doi.org/10.1016/0735-1097(90)90282-t).
 54. Thompson RCM, Ai MK, O'Keefe JH, Jr *et al.* Clinical utility of coronary calcium scoring after nonischemic myocardial perfusion imaging. *J Nucl Cardiol* 2005;**12**:392–400.
 55. Gudmundsson EF, Gudnason V, Sigurdsson S *et al.* Coronary artery calcium distributions in older persons in the AGES-Reykjavik study. *Eur J Epidemiol* 2012;**27**:673–87. DOI: [10.1007/s10654-012-9730-6](https://doi.org/10.1007/s10654-012-9730-6).
 56. Tota-Maharaj R, Blaha MJ, Blankstein R *et al.* Association of coronary artery calcium and coronary heart disease events in young and elderly participants in the multi-ethnic study of atherosclerosis: a secondary analysis of a prospective, population-based cohort. *Mayo Clin Proc* 2014;**89**:1350–9. DOI: [10.1016/j.mayocp.2014.05.017](https://doi.org/10.1016/j.mayocp.2014.05.017).
 57. Jasuja R, Pencina KM, Spencer DJ *et al.* Reference intervals for free testosterone in adult men measured using a standardized equilibrium dialysis procedure. *Andrology* 2023;**11**:125–33.
 58. Mayneris-Perxachs J, Arnoriaga-Rodríguez M, Luque-Córdoba D *et al.* Gut microbiota steroid sexual dimorphism and its impact on gonadal steroids: influences of obesity and menopausal status. *Microbiome* 2020;**8**:1–15.
 59. Hagmar M, Berglund B, Brismar K *et al.* Body composition and endocrine profile of male olympic athletes striving for leanness. *Clin J Sport Med* 2013;**23**:197–201.
 60. Blackwell AD, Trumble BC, Maldonado Suarez I *et al.* Immune function in Amazonian horticulturalists. *Ann Hum Biol* 2016;**43**:382–96. DOI: [10.1080/03014460.2016.1189963](https://doi.org/10.1080/03014460.2016.1189963).
 61. Trumble BC, Stieglitz J, Thompson ME *et al.* Testosterone and male cognitive performance in Tsimane forager-horticulturalists. *Am J Hum Biol* 2015;**27**:582–6.
 62. Veldhuis J, Yang R, Roelfsema F *et al.* Proinflammatory cytokine infusion attenuates LH's feedforward on testosterone secretion: modulation by age. *J Clin Endocrinol Metab* 2016;**101**:539–49. DOI: [10.1210/jc.2015-3611](https://doi.org/10.1210/jc.2015-3611).
 63. Handelsman DJ. Testosterone and male aging: faltering hope for rejuvenation. *JAMA* 2017;**317**:699–701. DOI: [10.1001/jama.2017.0129](https://doi.org/10.1001/jama.2017.0129).
 64. Corona G, Rastrelli G, Vignozzi L *et al.* Testosterone, cardiovascular disease and the metabolic syndrome. *Best Pract Res Clin Endocrinol Metab* 2011;**25**:337–53. DOI: [10.1016/j.beem.2010.07.002](https://doi.org/10.1016/j.beem.2010.07.002).

65. Deenadayalu V, Puttabyatappa Y, Liu AT *et al.* Testosterone-induced relaxation of coronary arteries: activation of BKCa channels via the cGMP-dependent protein kinase. *Am J Physiol Heart Circ Physiol* 2011;**302**:H115–23.
66. Pongkan W, Chattipakorn SC, Chattipakorn N. Roles of testosterone replacement in cardiac ischemia–reperfusion injury. *J Cardiovasc Pharmacol Ther* 2015;**21**:27–43. DOI: [10.1177/1074248415587977](https://doi.org/10.1177/1074248415587977).
67. Finkle WD, Greenland S, Ridgeway GK *et al.* Increased risk of non-fatal myocardial infarction following testosterone therapy prescription in men. *PLoS One* 2014;**9**:e85805. DOI: [10.1371/journal.pone.0085805](https://doi.org/10.1371/journal.pone.0085805).
68. Basaria S, Coviello AD, Travison TG *et al.* Adverse events associated with testosterone administration. *N Engl J Med* 2010;**363**:109–22. DOI: [10.1056/NEJMoa1000485](https://doi.org/10.1056/NEJMoa1000485).
69. Martinez C, Suissa S, Rietbrock S *et al.* Testosterone treatment and risk of venous thromboembolism: population based case-control study. *BMJ (Clinical research ed.)* 2016;**355**:i5968. DOI: [10.1136/bmj.i5968](https://doi.org/10.1136/bmj.i5968).
70. Vigen R, O'Donnell CI, Barón AE *et al.* Association of testosterone therapy with mortality, myocardial infarction, and stroke in men with low testosterone levels. *JAMA* 2013;**310**:1829–36. DOI: [10.1001/jama.2013.280386](https://doi.org/10.1001/jama.2013.280386).
71. Snyder PJ, Bhasin S, Cunningham GR *et al.*; Testosterone Trials Investigators. Effects of testosterone treatment in older men. *N Engl J Med* 2016;**374**:611–24. DOI: [10.1056/NEJMoa1506119](https://doi.org/10.1056/NEJMoa1506119).
72. Albert SG, Morley JE. Testosterone therapy, association with age, initiation and mode of therapy with cardiovascular events: a systematic review. *Clin Endocrinol (Oxf)* 2016;**85**:436–43.
73. Xu L, Freeman G, Cowling BJ *et al.* Testosterone therapy and cardiovascular events among men: a systematic review and meta-analysis of placebo-controlled randomized trials. *BMC Med* 2013;**11**:108. DOI: [10.1186/1741-7015-11-108](https://doi.org/10.1186/1741-7015-11-108).
74. Handelsman DJ. Global trends in testosterone prescribing, 2000–2011: expanding the spectrum of prescription drug misuse. *Med J Aust* 2013;**199**:548–51. DOI: [10.5694/mja13.10111](https://doi.org/10.5694/mja13.10111).
75. Lea AJ, Garcia A, Arevalo J *et al.* Natural selection of immune and metabolic genes associated with health in two lowland Bolivian populations. *Proc Natl Acad Sci USA* 2023;**120**:e2207544120.
76. Trumble BC, Cummings D, von Rueden C *et al.* Physical competition increases testosterone among Amazonian forager-horticulturalists: a test of the 'challenge hypothesis'. *Proc Biol Sci* 2012;**279**:2907–12.
77. Trumble BC, Cummings DK, O'Connor KA *et al.* Age-independent increases in male salivary testosterone during horticultural activity among Tsimane forager-farmers. *Evolut Human Behav* 2013;**34**:350–7.
78. Rosner W, Auchus RJ, Azziz R *et al.* Utility, limitations, and pitfalls in measuring testosterone: an endocrine society position statement. *J Clin Endocrinol Metab* 2007;**92**:405–13. DOI: [10.1210/jc.2006-1864](https://doi.org/10.1210/jc.2006-1864).