



Degenerating products of flag varieties and applications to the Breuil–Mézard conjecture

Robin Bartlett¹

Accepted: 10 November 2023 / Published online: 19 January 2024
© Crown 2024

Abstract

We consider closed subschemes in the affine grassmannian obtained by degenerating e -fold products of flag varieties, embedded via a tuple of dominant cocharacters. For $G = \mathrm{GL}_2$, and cocharacters small relative to the characteristic, we relate the cycles of these degenerations to the representation theory of G . We then show that these degenerations smoothly model the geometry of (the special fibre of) low weight crystalline subspaces inside the Emerton–Gee stack classifying p -adic representations of the Galois group of a finite extension of \mathbb{Q}_p . As an application we prove new cases of the Breuil–Mézard conjecture in dimension two.

Mathematics Subject Classification Primary 11F80; Secondary 14M15

Contents

1	Introduction	2
	Overview	2
	Main result	3
	Method	4
	Part 1: local models in the affine grassmannian	4
	Part 2: from local models to moduli of crystalline Galois representations	6
	Part 3: upper and lower multiplicity bounds	8
2	Notation	10
3	Cycles	11
4	Local models	13
5	Cohomology	17
6	Multiplicity bounds	18
7	Topological descriptions	21
8	Duality	27
9	Breuil–Kisin modules	28
10	Crystalline Breuil–Kisin modules	31
11	Naive Galois actions	33

✉ Robin Bartlett
robin.bartlett@glasgow.ac.uk

¹ University of Glasgow, Glasgow, UK

12 Comparison with local models 36
 13 Lower bounds 40
 14 Main result 43
 15 Miscellany 46
 References 47

1 Introduction

Overview

Let K be a finite extension of \mathbb{Q}_p with residue field k and let \mathcal{X}_d denote the Emerton–Gee stack classifying d -dimensional p -adic representations of G_K . Inside \mathcal{X}_d there are closed substacks $\mathcal{X}_d^{\mu, \tau}$ classifying potentially crystalline representations of type (μ, τ) , for μ and τ respectively Hodge and inertial types. When μ is regular (i.e. consists of distinct integers) these closed substacks have maximal dimension and the Breuil–Mézard conjecture [4, 5, 7] predicts the existence of top dimensional cycles \mathcal{C}_λ in the special fibre $\overline{\mathcal{X}}_d$ such that

$$[\overline{\mathcal{X}}_d^{\mu, \tau}] = \sum_{\lambda} m(\lambda, \mu, \tau) \mathcal{C}_\lambda \tag{1.1}$$

where

- λ runs over irreducible $\overline{\mathbb{F}}_p$ -representations of $\mathrm{GL}_d(k)$.
- $m(\lambda, \mu, \tau)$ denotes the multiplicity with which λ appears in an explicit \mathbb{F} -representation $V(\mu, \tau)$ of $\mathrm{GL}_d(k)$ attached to μ and τ .

(there is also a version of the conjecture for substacks of potentially semistable representations; the conjecture has the same shape but with altered $V(\mu, \tau)$). These identities have been verified in only a small number of cases:

- (1) When $K = \mathbb{Q}_p$ and $d = 2$, using the p -adic Langlands correspondence. See [14, 18, 23, 25, 27].
- (2) When $d = 2$ and $\mu = (1, 0)$, as consequence of certain modularity lifting theorems. See [10].
- (3) When K is unramified over \mathbb{Q}_p , d is arbitrary, and both p and τ are *generic* relative to μ . See [21]. Again modularity lifting technique play an important role.

In this paper we construct Breuil–Mézard identities in a fourth setting: we are interested in the two dimensional case where $\tau = 1$ (i.e. we consider only crystalline rather than potentially crystalline representations) and μ is bounded so that the representation theory of $\mathrm{GL}_2(k)$ in the conjecture behaves as it does in characteristic zero. We do this by constructing analogous identities involving certain degenerations of products of flag varieties embedded in the affine grassmannian, and then relating the geometry of these degenerations to the geometry of the $\overline{\mathcal{X}}_d^{\lambda, 1}$.

Main result

First we describe a bound on the Hodge types considered above, which is natural in the sense that the $GL_d(k)$ -representation theory appearing in the conjecture changes markedly once the bound is passed. Recall that a Hodge type μ consists of a d -tuple of integers

$$\mu_\kappa = (\mu_{\kappa,1} \geq \dots \geq \mu_{\kappa,d})$$

for each embedding $\kappa : K \rightarrow \overline{\mathbb{Q}}_p$. If one assumes that

$$\sum_{\kappa|k=\kappa_0} \mu_{\kappa,1} - \mu_{\kappa,d} \leq e + p - 1 \tag{1.2}$$

for each embedding $\kappa_0 : k \hookrightarrow \overline{\mathbb{F}}_p$ then:

- $V(\mu, 1)$ is a tensor product over the embeddings κ of representations of highest weight μ_κ and the Jordan–Holder factors of this tensor product are computed in characteristic p just as they are in characteristic zero, by Littlewood–Richardson coefficients.
- Each Jordan–Holder factor λ of $V(\mu, 1)$ can be written as $V(\tilde{\lambda}, 1)$ for some Hodge type $\tilde{\lambda}$ uniquely determined up to an ordering of the embeddings κ .

In particular, the cycles \mathcal{C}_λ appearing in (1.1) for these small μ are uniquely determined by the conjectured identity for $\mu = \tilde{\lambda}$; one has $[\overline{\mathcal{X}}_d^{\tilde{\lambda}}] = \mathcal{C}_\lambda$. Thus, the following theorem establishes new cases of the conjecture:

Theorem 1.3 *Assume that $d = 2$, $p > 2$, μ is regular, and that*

$$\sum_{\kappa|k=\kappa_0} \mu_{\kappa,1} - \mu_{\kappa,d} \leq p$$

for each embedding $\kappa_0 : k \rightarrow \overline{\mathbb{F}}_p$. Then

$$[\overline{\mathcal{X}}_2^{\mu,1}] = \sum_{\lambda} m(\lambda, \mu, 1) [\overline{\mathcal{X}}_2^{\tilde{\lambda},1}] \tag{1.4}$$

There are some comments to make before we discuss what goes into the proof. Firstly, the theorem has two clear limitations: the assumption that $d = 2$ and the fact that the bound on μ is stronger than that in (1.2) (we have no expectation whatever that the methods in this paper apply beyond (1.2)).

As we explain in more detail below, the proof of the theorem has two key inputs. The first involves relating the $\overline{\mathcal{X}}_d^{\mu,1}$ with certain local models we define inside the affine grassmannian. This can be done without any restriction on d but our current argument requires the stronger bound on μ . The second key input is a lower bound

on the Breuil–Mézard identities which has been established when $d = 2$ using global techniques from [10] (this is also where the assumption $p > 2$ appears).

Finally, taking $\tilde{\lambda} = \mu$ shows that the cycle $[\overline{X}_2^\lambda]$ is independent of the choice of “lift” $\tilde{\lambda}$ of λ . We also show that each of these cycles consists of a single irreducible component.

Method

The proof of the theorem divides into three parts:

Part 1: local models in the affine grassmannian

The starting point of the proof is the construction of certain projective schemes whose special fibres give upper bounds on the multiplicities appearing in Theorem 1.3. To explain their construction we fix a sufficiently large extension E of \mathbb{Q}_p , with ring of integers \mathcal{O} and residue field \mathbb{F} , and consider a mixed characteristic version of the affine grassmannian $\text{Gr}_{\mathcal{O}}$ over \mathcal{O} whose special and generic fibres are given by

$$\text{Gr}_{\mathcal{O}} \otimes_{\mathcal{O}} \mathbb{F} \cong \prod_{\kappa_0: k \rightarrow \mathbb{F}} \text{Gr} \otimes_{\mathcal{O}_K, \kappa_0} k, \quad \text{Gr}_{\mathcal{O}} \otimes_{\mathcal{O}} E \cong \prod_{\kappa: K \rightarrow E} \text{Gr} \otimes_{\mathcal{O}_K, \kappa} E$$

Here κ_0 and κ are embeddings and Gr is the affine grassmannian over \mathcal{O}_K whose A points, for A a p -adically complete \mathcal{O}_K -algebra, classify rank d -projective $A[[u]]$ -modules satisfying

$$(u - \pi)^a A[[u]]^d \subset \mathcal{E} \subset (u - \pi)^{-a} A[[u]]^d$$

for some $a \in \mathbb{Z}_{\geq 0}$ and $\pi \in K$ a fixed choice of uniformiser. For each dominant cocharacter λ of $G = \text{GL}_d$ there is a closed immersion of the flag variety $G/P_\lambda \rightarrow \text{Gr}$ ($P_\lambda \subset G$ being the parabolic corresponding to λ). This allows us to define, for any Hodge type $\mu = (\mu_\kappa)$, an \mathcal{O} -flat closed subscheme M_μ in $\text{Gr}_{\mathcal{O}}$ by taking the closure in $\text{Gr}_{\mathcal{O}}$ of

$$\prod_{\kappa} (G/P_{\mu_\kappa} \otimes_{\mathcal{O}_K, \kappa} E) \hookrightarrow \prod_{\kappa} (\text{Gr} \otimes_{\mathcal{O}_K, \kappa} E) = \text{Gr}_{\mathcal{O}} \otimes_{\mathcal{O}} E$$

The following summarises the key results we prove regarding these M_μ ’s

Proposition 1.5 *Assume that μ is regular.*

- (1) *If μ satisfies (1.2) then there exist $n(\lambda, \mu) \in \mathbb{Z}$ such that in the group of $\sum_{\kappa} \dim G/P_{\mu_\kappa}$ -dimensional cycles*

$$[M_\mu \otimes_{\mathcal{O}} \mathbb{F}] = \sum_{\lambda} n(\lambda, \mu) [M_{\tilde{\lambda}} \otimes_{\mathcal{O}} \mathbb{F}]$$

with the sum running those irreducible $GL_d(k)$ -representations for which the Hodge type $\tilde{\lambda}$ also satisfies (1.2).

- (2) If $d = 2$ then the $M_{\tilde{\lambda}} \otimes_{\mathcal{O}} \mathbb{F}$ appearing in (1) are irreducible, generically reduced, and produce pairwise distinct cycles. In particular, $n(\lambda, \mu) \geq 0$ in this case.
- (3) If, for every λ , one has $n(\lambda, \mu) \geq m(\lambda, \mu, 1)$ where $m(\lambda, \mu, 1)$ denotes the multiplicity from the Breuil–Mézard conjecture, then $n(\lambda, \mu) = m(\lambda, \mu, 1)$.

The first part is proved by constructing an explicit closed locus $Gr_{\mathcal{O}}^{\nabla} \subset Gr_{\mathcal{O}}$ defined in terms of a differential operator ∇ (this is a variant of locus considered in [21]). A direct computation shows that if we bound the height according to (1.2) then the resulting closed subscheme of $Gr_{\mathcal{O}}^{\nabla} \otimes_{\mathcal{O}} \mathbb{F}$ consists of irreducible components of dimension $\leq \dim M_{\mu}$. Furthermore, those components with maximal dimension are labelled by the λ 's appearing in (1). One can also show that $M_{\mu} \otimes_{\mathcal{O}} \mathbb{F}$ is contained in this closed subscheme. From these observations we are able to prove (1).

Remark 1.6 Unfortunately, this explicit moduli interpretation is only a good topological approximation of $M_{\mu} \otimes_{\mathcal{O}} \mathbb{F}$; typically the components appear with much too high multiplicity.

Part (2) is proved by constructing an explicit resolution of $X \rightarrow M_{\tilde{\lambda}}$ with X smooth and which is an isomorphism on the generic fibre. Unfortunately, we do not know how to construct such resolutions when $d > 2$ (or whether they are likely to exist).

For part (3) we consider the restriction of the determinant line bundle on $Gr_{\mathcal{O}}$ to M_{μ} . Since the generic fibre of M_{μ} is a product of flag varieties it is easy to compute that for

$$H^0(M_{\mu} \otimes_{\mathcal{O}} E, \mathcal{L}_{\det}) = \bigotimes_{\kappa} H^0(\mu_{\kappa}) \otimes_{K,\kappa} E \tag{1.7}$$

where $H^0(\mu_{\kappa})$ denotes the algebraic representation of G over K of highest weight μ_{κ} . We point out that this tensor product differs from the $V(\mu, 1)$ appearing in the Breuil–Mézard conjecture in that $V(\mu, 1)$ is obtained as the reduction modulo p of such a tensor product, but in which μ_{κ} is replaced by $\mu_{\kappa} - \rho$ for $\rho = (d - 1, d - 2, \dots, 1, 0)$. Nevertheless, these multiplicities are approximately the same, in the sense that if, in the Grothendieck group of E -representations, one has

$$\left[\bigotimes_{\kappa} H^0(\mu_{\kappa} - \rho) \otimes_{K,\kappa} E \right] = \sum_{\lambda} m(\lambda, \mu) \left[\bigotimes_{\kappa} H^0(\tilde{\lambda}_{\kappa} - \rho) \otimes_{K,\kappa} E \right]$$

then, for $n > 0$,

$$\dim \left(\bigotimes_{\kappa} H^0(n\mu_{\kappa}) \otimes_{K,\kappa} E \right) - \sum_{\lambda} m(\lambda, \mu) \dim \left(\bigotimes_{\kappa} H^0(n\tilde{\lambda}_{\kappa}) \otimes_{K,\kappa} E \right)$$

equals the value at n of a polynomial of degree $< \dim M_{\mu}$. Since the representations $\bigotimes_{\kappa} H^0(n\mu_{\kappa}) \otimes_{\mathcal{O},\kappa} E$ can be obtained by replacing \mathcal{L}_{\det} with $\mathcal{L}_{\det}^{\otimes n}$ in (1.7), the identity of cycles in part (1) implies that

$$\dim \left(\bigotimes_{\kappa} H^0(n\mu_{\kappa}) \otimes_{K,\kappa} E \right) - \sum_{\lambda} n(\lambda, \mu) \dim \left(\bigotimes_{\kappa} H^0(n\tilde{\lambda}_{\kappa}) \otimes_{K,\kappa} E \right)$$

is also the value of a polynomial in n of degree $< \dim M_{\mu}$, at least for $n \gg 0$. Taking the difference shows that

$$\sum_{\lambda} (n(\lambda, \mu) - m(\lambda, \mu)) \dim \left(\bigotimes_{\kappa} H^0(n\tilde{\lambda}_{\kappa}) \otimes_{K,\kappa} E \right) \tag{1.8}$$

is polynomial in n of degree $< \dim M_{\mu}$ for $n \gg 0$. For μ satisfying (1.2) the multiplicities $m(\lambda, \mu)$ computed in characteristic zero coincide with the $m(\lambda, \mu, 1)$ computed in characteristic p . Thus, the assumption in (3) is that $n(\lambda, \mu) - m(\lambda, \mu) \geq 0$. Each term in (1.8) is a polynomial in n of degree $\dim M_{\mu}$ and positive leading term. Therefore we must have $n(\lambda, \mu) = m(\lambda, \mu)$.

Part 2: from local models to moduli of crystalline Galois representations

The second step is to relate the M_{μ} 's to the geometry of \mathcal{X}_d . The basic strategy is to study the geometry of \mathcal{X}_d via a resolution

$$Y_d \rightarrow \mathcal{X}_d$$

with Y_d a stack whose A -points classify Breuil–Kisin modules with A -coefficients (i.e. projective $(W(k) \otimes_{\mathbb{Z}_p} A)[[u]]$ -modules equipped with a semilinear endomorphism φ). A local version of this construction was first made in [19] (with \mathcal{X}_d replaced by Spec of a deformation ring) and its globalisation to stacks first appeared in [24], before being built upon in [7].

In our case, we take Y_d as the stack classifying pairs (\mathfrak{M}, σ) with \mathfrak{M} a rank d Breuil–Kisin module and σ a φ -equivariant action of G_K on $\mathfrak{M} \otimes_{W(k)[[u]]} A_{\text{inf}}$ satisfying a “crystalline” condition (which means that $\sigma - 1$ is sufficiently divisible). Inside Y_d there are \mathbb{Z}_p -flat closed substacks Y_d^{μ} whose \mathcal{O} -valued points correspond to Breuil–Kisin modules associated to crystalline representations of Hodge type μ whenever \mathcal{O} is the ring of integers in a finite extension of \mathbb{Q}_p . Then $\mathcal{X}_d^{\mu,1}$ is, by definition, the scheme theoretic image of the morphism $Y_d^{\mu} \rightarrow \mathcal{X}_d$.

To relate Y_d to the affine grassmannian we use the following diagram:

$$Y_d \xleftarrow{\Gamma} \tilde{Y}_d \xrightarrow{\Psi} \text{Gr}_{\mathcal{O}} \tag{1.9}$$

Here \tilde{Y}_d classifies Breuil–Kisin modules in Y_d together with a choice of basis (to stay in the world of finite type stacks this basis is taken modulo u^N for $N \gg 0$) and, using this choice of basis, the morphism Ψ takes a Breuil–Kisin module \mathfrak{M} to the relative position of \mathfrak{M} and its image of Frobenius. The morphism Γ forgets this choice of basis. The key result we prove is then

- Proposition 1.10** (1) *If μ satisfies the bound from Theorem 1.3 then the restriction of $\tilde{Y}_d \rightarrow \text{Gr}_{\mathcal{O}}$ to $\tilde{Y}_d^\mu \otimes_{\mathcal{O}} \mathbb{F}$ (for \tilde{Y}_d^μ the preimage of Y_d^μ in \tilde{Y}_d) factors through $M_{-w_0\mu} \otimes_{\mathcal{O}} \mathbb{F}$ for $w_0 \in W$ the longest element.*
 (2) *For such μ , the morphism $\tilde{Y}_d \rightarrow \text{Gr}_{\mathcal{O}}$ is smooth over $M_\mu \otimes_{\mathcal{O}} \mathbb{F}$ with irreducible fibres of dimension equal the relative dimension of $\tilde{Y}_d \rightarrow Y_d$.*

To prove (1) it suffices to show this factorisation for \bar{A} -points for every finite \mathbb{F} -algebra \bar{A} . For simplicity, we sketch the argument only in the case where $\bar{A} = \mathbb{F}$. The general case requires only minor technical changes. We also assume $k = \mathbb{F}_p$ as this greatly simplifies the notation. If $e = 1$ then $M_\mu = G/P_\mu$ is just a single flag variety and the claimed factorisation comes down to showing that for any $\bar{\mathfrak{M}} \in Y_d^{\mu,1}(\bar{A})$ and any basis $\bar{\beta}$ the module $\bar{\mathfrak{M}}$ is generated by

$$\varphi(\bar{\beta})g \begin{pmatrix} u^{-\mu_1} & & & \\ & \ddots & & \\ & & \ddots & \\ & & & u^{-\mu_d} \end{pmatrix}$$

for some $g \in \text{GL}_d(\bar{A})$. This follows from results in [11] where it is shown, for any lift of $\bar{\mathfrak{M}}$ to $\mathfrak{M} \in Y_d^\mu(A)$, with A the ring of integers in a finite extension of E , and any basis β that $(u - \pi)^p \mathfrak{M}$ is generated by

$$\varphi(\beta)g \left[\begin{pmatrix} (u - \pi)^{p-\mu_1} & & & \\ & \ddots & & \\ & & \ddots & \\ & & & (u - \pi)^{p-\mu_d} \end{pmatrix} + X_{\text{err}} \right]$$

for a matrix X_{err} divisible by a power of $\pi^{p-\mu_1+\mu_d+1}$ and $g \in \text{GL}_d(A)$. Here $\pi \in K$ is a fixed uniformiser. This result does not directly extend to the case $e > 1$. However, a variant of the method is able to show that, for each embedding $\kappa : K \rightarrow E$, the module $\mathfrak{M}^\varphi \cap (u - \kappa(\pi))^p \mathfrak{M}$ can be generated by

$$\varphi(\beta)g_\kappa \left[\begin{pmatrix} (u - \kappa(\pi))^{p-\mu_{\kappa,1}} & & & \\ & \ddots & & \\ & & \ddots & \\ & & & (u - \kappa(\pi))^{p-\mu_{\kappa,d}} \end{pmatrix} + X_{\text{err},\kappa} \right]$$

for some $g_\kappa \in \text{GL}_d(A)$ and $X_{\text{err},\kappa}$ a matrix divisible by $\pi^{p-\mu_{\kappa,1}+\mu_{\kappa,d}+1}$. This was done in [12] (actually they only consider the case $d = 2$ but it is straightforward to extend their arguments to higher dimensions). If the $X_{\text{err},\kappa}$'s are divisible by a high enough power of π then it follows that

$$\prod_{\kappa} (u - \kappa(\pi))^p \mathfrak{M} = \bigcap_{\kappa} (\mathfrak{M}^\varphi \cap (u - \kappa(\pi))^p \mathfrak{M})$$

is congruent modulo π to the intersection of the submodules generated by

$$\varphi(\beta)_{g_\kappa} \begin{pmatrix} (u - \kappa(\pi))^{p-\mu_{\kappa,1}} & & & \\ & \ddots & & \\ & & \ddots & \\ & & & (u - \kappa(\pi))^{p-\mu_{\kappa,d}} \end{pmatrix}$$

This sufficient divisibility is ensured by the bound on μ from Theorem 1.3 and this congruence is precisely what it means of $\overline{\mathfrak{M}}$ to be mapped onto an element of $M_{-w_0\mu} \otimes_{\mathcal{O}} \mathbb{F}$ by Ψ .

For the proof of (2) we factor the morphism Ψ as $\tilde{Y}_d \rightarrow \tilde{Z}_d \rightarrow \text{Gr}_{\mathcal{O}}$ where \tilde{Z}_d denotes the moduli stack of Breuil–Kisin modules (without a crystalline Galois action) and $\tilde{Y}_d \rightarrow \tilde{Z}_d$ forgets the Galois action. An easy calculation shows that over the special fibre $\tilde{Z}_d \rightarrow \text{Gr}_{\mathcal{O}}$ is smooth with irreducible fibres of dimension equal the relative dimension of $\tilde{Y}_d \rightarrow Y_d$. Part (2) therefore reduces to understanding when $\tilde{Y}_d \rightarrow \tilde{Z}_d$ is an isomorphism. To address this we note that for any Breuil–Kisin module \mathfrak{M} with basis β we can define a naive Galois action $\sigma_{\text{naive},\beta}$ on \mathfrak{M} by semilinearly extending the trivial G_K -action on $\varphi(\beta)$. Usually $\sigma_{\text{naive},\beta}$ will not be φ -equivariant or crystalline. However, we show that if $\sigma_{\text{naive},\beta} - 1$ is suitably divisible and if \mathfrak{M} satisfies height conditions imposed by (1.2) (actually a very slight strengthening of this bound is required to avoid certain “Steinberg” situations) then

$$\lim_{n \rightarrow \infty} \varphi^n \circ \sigma_{\text{naive},\beta} \circ \varphi^{-n}$$

converges to a unique φ -equivariant crystalline G_K -action. It turns out that the locus of \tilde{Z}_d on which $\sigma_{\text{naive},\beta} - 1$ is sufficiently divisible is closed, and obtained as the preimage of a closed subscheme in $\text{Gr}_{\mathcal{O}}$. Part (2) is then proved by showing that $M_{\mu} \otimes_{\mathcal{O}} \mathbb{F}$ is contained in this closed subscheme.

Part 3: upper and lower multiplicity bounds

The final ingredient which goes into the proof of Theorem 1.3 is a lower bound on the multiplicities appearing in the Breuil–Mézard conjecture. This is the most critical place where we require $d = 2$. It is also where we use that $p > 2$. Under these assumptions it is shown in [7, 8.6] (using global automorphy lifting techniques from [10]) that one always has

$$[\overline{\mathcal{X}}_2^{\mu,\tau}] \geq \sum_{\lambda} m(\lambda, \mu, \tau) \mathcal{C}_{\lambda}$$

This holds without any assumption on μ or τ . Combining [10] with the potential diagonalisability established in [1] one also obtains that $\mathcal{C}_{\lambda} = [\overline{\mathcal{X}}_2^{\lambda,1}]$ so long as λ is not Steinberg (for $d = 2$ this means λ is not a twist of $\bigotimes_{\kappa_0:k \rightarrow \mathbb{F}} \text{Sym}^{p-1} \mathbb{F}^2$). The bounds on μ ensure Steinberg λ do not appear in Theorem 1.3 (except if K/\mathbb{Q}_p is unramified, but in this case the theorem is trivial). Therefore, for μ as in Theorem 1.3,

we have

$$[\bar{\mathcal{X}}_2^{\mu,1}] \geq \sum_{\lambda} m(\lambda, \mu, 1)[\bar{\mathcal{X}}_2^{\tilde{\lambda},1}]$$

To finish the proof we have to show that the results from parts 1. and 2. can be combined to give equality.

First we consider the identity $[M_{\mu} \otimes_{\mathcal{O}} \mathbb{F}] = \sum_{\lambda} n(\lambda, \mu)[M_{\tilde{\lambda}} \otimes_{\mathcal{O}} \mathbb{F}]$ from Proposition 1.5. Applying an involution of $\text{Gr}_{\mathcal{O}}$ which sends a lattice onto its dual allows us to replace μ and each $\tilde{\lambda}$ in this identity with $-w_0\mu$ and $-w_0\tilde{\lambda}$. Thus

$$[M_{-w_0\mu} \otimes_{\mathcal{O}} \mathbb{F}] = \sum_{\lambda} n(\lambda, \mu)[M_{-w_0\tilde{\lambda}} \otimes_{\mathcal{O}} \mathbb{F}]$$

Part (2) of Proposition 1.10 ensures $\tilde{Y}_2 \rightarrow \text{Gr}_{\mathcal{O}}$ is smooth over the closed subschemes appearing in this identity of cycles. This allows us to pull the identity back to \tilde{Y}_d to obtain

$$[Y_2^{\mu,\text{flag}}] = \sum_{\lambda} n(\lambda, \mu)[Y_2^{\tilde{\lambda},\text{flag}}]$$

where $Y_2^{\mu,\text{flag}}$ equals the preimage of $M_{-w_0\mu} \otimes_{\mathcal{O}} \mathbb{F}$ under this map. Part (1) of Proposition 1.10 (together with a dimension comparison) implies $[\tilde{Y}_d^{\mu} \otimes_{\mathcal{O}} \mathbb{F}] \leq [Y_2^{\mu,\text{flag}}]$. Using part (2) of Proposition 1.5 we are even able to deduce this is an equality when $\mu = \tilde{\lambda}$. Therefore

$$[\tilde{Y}_2^{\mu} \otimes_{\mathcal{O}} \mathbb{F}] \leq \sum_{\lambda} n(\lambda, \mu)[\tilde{Y}_2^{\tilde{\lambda}} \otimes_{\mathcal{O}} \mathbb{F}]$$

Since $\tilde{Y}_d \rightarrow Y_d$ is smooth and surjective it follows that also

$$[Y_2^{\mu} \otimes_{\mathcal{O}} \mathbb{F}] \leq \sum_{\lambda} n(\lambda, \mu)[Y_2^{\tilde{\lambda}} \otimes_{\mathcal{O}} \mathbb{F}]$$

Pushing this identity forward along the proper morphism $Y_2 \rightarrow \mathcal{X}_2$ gives an inequality $[\bar{\mathcal{X}}_2^{\mu}] \leq \sum_{\lambda} n(\lambda, \mu)[\bar{\mathcal{X}}_2^{\tilde{\lambda}}]$. Combining this with the lower bound we obtain $n(\lambda, \mu) \geq m(\lambda, \mu, 1)$. By part (3) of Proposition 1.5 this must be an equality, which proves Theorem 1.3. Actually, in the paper we follow the same argument, but for the final step we prefer to work with deformation rings rather than $\bar{\mathcal{X}}_2$. This allows us to avoid dealing with stacks. As explained in [7, 8.3], Theorem 1.3 is implied by its analogue in the setting of deformation rings.

2 Notation

2.1 We fix a finite extension K of \mathbb{Q}_p with residue field k of degree f over \mathbb{F}_p and ramification degree e . Let C denote the completed algebraic closure of K , with ring of integers \mathcal{O}_C , and fix a compatible system π^{1/p^∞} of p -th power roots of a fixed choice of uniformiser $\pi \in K$ in \mathcal{O}_C . Set $K_\infty = K(\pi^{1/p^\infty})$. Write $E(u) \in W(k)[u]$ for the minimal polynomial of π . Thus $E(u)$ is Eisenstein of degree equal to the ramification degree e of K over \mathbb{Q}_p .

We also fix another finite extension E of \mathbb{Q}_p with ring of integers \mathcal{O} and residue field \mathbb{F} . We assume that E contains a Galois closure of K . We typically use κ and κ_0 respectively to denote embeddings $K \rightarrow E$ and $k \rightarrow \mathbb{F}$. For each κ_0 we fix an embedding $\tilde{\kappa}_0 : K \rightarrow E$ with $\tilde{\kappa}_0|_k = \kappa_0$.

2.2 For any \mathbb{Z}_p -algebra A we write $\mathfrak{S}_A = (W(k) \otimes_{\mathbb{Z}_p} A)[[u]]$. This comes equipped with the A -linear endomorphism φ which on $W(k)$ acts as the lift of the p -th power map on k and sends $u \mapsto u^p$. We also consider

$$A_{\text{inf},A} = \varprojlim_a \varprojlim_i (W(\mathcal{O}_{C^b})/p^a \otimes_{\mathbb{Z}_p} A)/u^i$$

where $\mathcal{O}_{C^b} = \varprojlim_{x \mapsto x^p} \mathcal{O}_C/p$ and $u = [(\pi, \pi^{1/p}, \pi^{1/p^2}, \dots)] \in W(\mathcal{O}_{C^b})$. We view $A_{\text{inf},A}$ as an \mathfrak{S}_A -algebra via u . Note that the lift of Frobenius on $W(\mathcal{O}_{C^b})$ induces a Frobenius φ on $A_{\text{inf},A}$ which is compatible with that on \mathfrak{S}_A . The natural G_K -action on \mathcal{O}_C also induces a continuous (for the (u, p) -adic topology) G_K -action on $A_{\text{inf},A}$ commuting with φ . Write

$$W(C^b)_A = \varprojlim_a A_{\text{inf},A}[\frac{1}{u}]/p^a$$

If A is topologically of finite type (i.e. $A \otimes_{\mathbb{Z}_p} \mathbb{F}_p$ is of finite type) then $\mathfrak{S}_A \rightarrow A_{\text{inf},A}$ is faithfully flat (in particular injective) [7, 2.2.13].

We also fix a compatible system $(1, \epsilon_1, \epsilon_2, \dots)$ of p -th power roots of unity in \mathcal{O}_C which we view as an element $\epsilon \in \mathcal{O}_{C^b}$. We write $\mu = [\epsilon] - 1 \in A_{\text{inf},A}$.

Lemma 2.3 *Let $\Theta : A_{\text{inf}} \rightarrow \mathcal{O}_C$ denote the surjection given by $\sum p^i x_i \mapsto \sum p^i x_i^{(0)}$ with $x_i = [(x_i^{(j)})_{j \geq 1}]$ and extend this to a surjection $A_{\text{inf},\mathcal{O}} \rightarrow \mathcal{O} \otimes_{\mathbb{Z}_p} \mathcal{O}_C$. Then*

$$\mu A_{\text{inf},\mathcal{O}} = \{x \in A_{\text{inf},\mathcal{O}} \mid \Theta(\varphi^n(x)) = 0 \text{ for all } n \geq 0\}$$

Proof By choosing a \mathbb{Z}_p -basis of \mathcal{O} this follows immediately from the assertion that $\mu A_{\text{inf}} = \{x \in A_{\text{inf}} \mid \Theta(\varphi^n(x)) = 0 \text{ for all } n \geq 0\}$ which is [8, 5.1.3]. \square

2.4 We frequently consider modules as in 2.2 defined over $\mathcal{O} \otimes_{\mathbb{Z}_p} W(k)$ for an \mathcal{O} -algebra A . Using the isomorphism

$$\mathcal{O} \otimes_{\mathbb{Z}_p} W(k) \xrightarrow{\sim} \prod_{\kappa_0: k \rightarrow \mathbb{F}} \mathcal{O} \cong \prod_{\kappa_0} \mathcal{O} \otimes_{W(k), \kappa_0} W(k)$$

given by $a \otimes b \mapsto (a\kappa_0(b))_{\kappa_0}$ (here we write κ_0 to its extension to an embedding $W(k) \rightarrow \mathcal{O}$) we see that any such module M can be expressed as a product

$$M = \prod_{\kappa_0} M_{\kappa_0}$$

where M_{κ_0} can be identified with the submodule of M on which the two actions of $W(k)$ given by $(1 \otimes a)m$ and $a \mapsto (\kappa_0(a) \otimes 1)m$ coincide. Similarly, there is an isomorphism

$$E \otimes_{\mathbb{Z}_p} \mathcal{O}_K \xrightarrow{\sim} \prod_{\kappa:K \rightarrow E} E \cong \prod_{\kappa} E \otimes_{\mathcal{O}_{K,\kappa}} \mathcal{O}_K \tag{2.5}$$

given by $a \otimes b \mapsto (\kappa(b)a)_{\kappa}$ which allows us to write an $E \otimes_{\mathbb{Z}_p} \mathcal{O}_K$ -module M as

$$M = \prod_{\kappa} M_{\kappa}$$

where again M_{κ} can be identified with the submodule consisting of $m \in M$ with $(1 \otimes a)m = (\kappa(a) \otimes 1)m$ for all $a \in \mathcal{O}_K$. We warn the reader that the idempotents in (2.5) will not be contained in $\mathcal{O} \otimes_{\mathbb{Z}_p} \mathcal{O}_K$ whenever K/\mathbb{Q}_p is ramified and so the product decomposition $M = \prod_{\kappa} M_{\kappa}$ is not valid integrally, i.e. when M is an $\mathcal{O} \otimes_{\mathbb{Z}_p} \mathcal{O}_K$ -module.

2.6 Applying the previous discussion to $(A \otimes_{\mathbb{Z}_p} W(k))[u]$ allows us to write

$$(A \otimes_{\mathbb{Z}_p} W(k))[u] = \prod_{\kappa_0} A[u]$$

Using this identification we define $E_{\kappa}(u) \in (A \otimes_{\mathbb{Z}_p} W(k))[u]$ for every embedding $\kappa : K \rightarrow E$ as the element corresponding to

$$(1, \dots, 1, u - \pi_{\kappa}, 1, \dots, 1) \in \prod_{\kappa_0} A[u]$$

where $\pi_{\kappa} := \kappa(\pi)$ and the $u - \pi_{\kappa}$ appears in the $\kappa|_k$ -th factor in the product. Notice that $E(u) = \prod_{\kappa} E_{\kappa}(u)$ inside $(A \otimes_{\mathbb{Z}_p} W(k))[u]$.

3 Cycles

3.1 For a Noetherian scheme X let $Z_m(X)$ denote the free abelian group generated by integral closed subschemes $Z \subset X$ of dimension m . If \mathcal{F} is a coherent sheaf on X with support of dimension $\leq m$ then we define

$$[\mathcal{F}] = \sum_Z \text{length}_{\mathcal{O}_{X,\xi}}(\mathcal{F}_{\xi})[Z] \in Z_m(X)$$

for $\xi \in Z$ the generic point. If $i : Y \rightarrow X$ is a closed immersion write $[Y] = [i_*\mathcal{O}_Y]$. Any flat morphism $f : X \rightarrow Y$ of relative dimension d produces a homomorphism $f^* : Z_m(Y) \rightarrow Z_{m+d}(X)$ with $f^*[\mathcal{F}] = [f^*\mathcal{F}]$. See [26, 02RE]. If instead f is proper then there is a pushforward homomorphism $f_* : Z_m(X) \rightarrow Z_m(Y)$ with $f_*[\mathcal{F}] = [f_*\mathcal{F}]$. See [26, 02R6].

Lemma 3.2 *Let X be a projective scheme over k equipped with an ample line bundle \mathcal{L} . Suppose that Y, Y_1, \dots, Y_s are m -dimensional closed subschemes in X and that*

$$[Y] = \sum n_i [Y_i] \in Z_m(X)$$

Then

$$\dim H^0(Y, \mathcal{L}^{\otimes n}) - \sum n_i \dim H^0(Y_i, \mathcal{L}^{\otimes n})$$

is, for large n , the value at n of a polynomial of degree $< m$.

Proof This follows from [26, 0BEN] and the fact that, since \mathcal{L} is ample, the higher cohomologies of $\mathcal{L}^{\otimes n}$ vanish for $n \gg 0$ [26, 0B5U]. \square

Lemma 3.3 *Suppose that $f : X \rightarrow Y$ is a proper morphism between equidimensional flat \mathcal{O} -schemes which becomes an isomorphism after applying $\otimes_{\mathcal{O}} E$. Suppose $Z_X \subset X, Z_Y \subset Y$ are \mathcal{O} -flat top dimensional closed subschemes for which f restricts to an isomorphism*

$$f : Z_X \otimes_{\mathcal{O}} E \xrightarrow{\sim} Z_Y \otimes_{\mathcal{O}} E$$

Then $f_*[Z_X \otimes_{\mathcal{O}} \mathbb{F}] = [Z_Y \otimes_{\mathcal{O}} \mathbb{F}]$.

Proof Let $A_m(X)$ denote the quotient of $Z_m(X)$ by rational equivalence. Since X is \mathcal{O} -flat there is a specialisation homomorphism

$$\sigma_X : A_m(X \otimes_{\mathcal{O}} E) \rightarrow A_m(X \otimes_{\mathcal{O}} \mathbb{F})$$

with $\sigma_X([Z_X \otimes_{\mathcal{O}} E]) = [Z_X \otimes_{\mathcal{O}} \mathbb{F}]$ whenever $Z_X \subset X$ is a closed \mathcal{O} -flat subscheme of relative dimension m . Furthermore σ_X commutes with proper pushforward. All this is explained in [9, 20.3].

If $m = \dim X$ then $A_m(X) = Z_m(X)$ by [9, 1.3.2]. Therefore, the two stated properties of the specialisation map give

$$\begin{aligned} f_*[Z_X \otimes_{\mathcal{O}} \mathbb{F}] &= f_*\sigma_X([Z_X \otimes_{\mathcal{O}} E]) = \sigma_Y(f_*[Z_X \otimes_{\mathcal{O}} E]) \\ &= \sigma_Y([Z_Y \otimes_{\mathcal{O}} E]) = [Z_Y \otimes_{\mathcal{O}} \mathbb{F}] \end{aligned}$$

in $Z_{\dim Y \otimes_{\mathcal{O}} \mathbb{F}}(Y \otimes_{\mathcal{O}} \mathbb{F})$. \square

4 Local models

4.1 We begin by defining an ind-scheme Gr over \mathcal{O}_K whose A -points classify rank d -projective $A[u]$ -modules satisfying

$$(u - \pi)^a A[u]^d \subset \mathcal{E} \subset \prod (u - \pi)^{-a} A[u]^d \tag{4.2}$$

for some $a \geq 0$. For each $\kappa_0 : k \rightarrow \mathbb{F}$, which we extend to an embedding $W(k) \rightarrow \mathcal{O}$, we also define Gr_{κ_0} as the ind-scheme over \mathcal{O} whose A -points classify rank d -projective $A[u]$ -modules satisfying

$$\kappa_0(E(u))^a A[u]^d \subset \mathcal{E} \subset \prod \kappa_0(E(u))^{-a} A[u]^d$$

for some $a \geq 0$.

Note that, for each $\kappa : K \rightarrow E$, we can view an A -valued point of $\text{Gr} \otimes_{\mathcal{O}_K, \kappa} \mathcal{O}$ as a rank d projective $A[u]$ -module satisfying $(u - \kappa(\pi))^a A[u]^d \subset \mathcal{E} \subset (u - \kappa(\pi))^{-a} A[u]^d$ for some $a \geq 0$. Therefore, if $\kappa|_k = \kappa_0$ then there is a natural closed embedding

$$\text{Gr} \otimes_{\mathcal{O}_K, \kappa} \mathcal{O} \rightarrow \text{Gr}_{\kappa_0}$$

Remark 4.3 Recall that an $A[u]$ -submodule as in (4.2) is $A[u]$ -projective of rank d if and only if $(u - \pi)^{-a} A[u]^d / \mathcal{E}$ is A -projective. In particular, this illustrates the ind-representability of the functor; the locus of \mathcal{E} as in (4.2) identifies with a closed subscheme of the usual grassmannian classifying submodules of $(u - \pi)^{-a} A[u]^d / (u - \pi)^a A[u]^d$.

4.4 Write $X(T)$ for the group of characters of GL_d relative to T , the diagonal torus, and identify $X(T) = \mathbb{Z}^d$ as usual. We say an element $\mu = (\mu_1, \dots, \mu_d) \in X(T)$ is dominant if $\mu_i \geq \mu_{i+1}$ and for any such dominant μ we write $\mathcal{E}_\mu \in \text{Gr}_K$ for the \mathcal{O}_K -point generated by

$$(e_1, \dots, e_d) \begin{pmatrix} (u - \pi)^{\mu_1} & & & \\ & \ddots & & \\ & & \ddots & \\ & & & (u - \pi)^{\mu_d} \end{pmatrix}$$

for (e_1, \dots, e_d) the standard basis in $\mathcal{O}_K[u]^d$. There is an obvious action of $G = \text{GL}_d$ on Gr and, since the stabiliser of \mathcal{E}_μ under this action is a parabolic subgroup $P_\mu \subset G$, the orbit map induces a proper monomorphism

$$G/P_\mu \rightarrow \text{Gr}$$

i.e. a closed immersion. Notice that if $\mu_i \geq \mu_{i+1}$ then P_μ is the opposite Borel of lower triangular matrices.

An alternative description of $G/P_\mu \rightarrow \text{Gr}$ can be given after interpreting A -points of G/P_μ as filtrations

$$\dots \subset \text{Fil}^{n+1} \subset \text{Fil}^n \subset \text{Fil}^{n-1} \subset \dots$$

of type μ on A^d (which means the n -th graded piece is A -projective of rank equal to the multiplicity of $-n$ in μ). Then on A -points the closed immersion $G/P_\lambda \rightarrow \text{Gr}$ is given by

$$\text{Fil}^\bullet \mapsto \sum_{i \geq \mu_d} (u - \pi)^i A[u] \text{Fil}^{-i}$$

where we view A^d as a submodule of $A[u]^d$ in the obvious way.

Lemma 4.5 *If A is a p -adically complete \mathcal{O} -algebra then Gr identifies with the set of rank d projective $A[[u]]$ -modules satisfying*

$$(u - \pi)^a A[[u]] \subset \mathcal{E} \subset (u - \pi)^{-a} A[[u]]$$

for some $a \geq 0$. Similarly, for each Gr_{κ_0} .

Proof This follows from the Beauville–Laszlo gluing lemma [3]. □

Lemma 4.6 *For each $\kappa_0 : k \rightarrow \mathbb{F}$ there is an isomorphism*

$$\text{Gr}_{\kappa_0} \otimes_{\mathcal{O}} E \rightarrow \prod_{\kappa|_k = \kappa_0} (\text{Gr} \otimes_{\mathcal{O}_{K,\kappa}} E)$$

with inverse given by $(\mathcal{E}_\kappa) \mapsto \bigcap_{\kappa|_k = \kappa_0} \mathcal{E}_\kappa$.

Proof Let $U \subset \mathbb{A}_A^1$ denote the open obtained by inverting $\kappa_0(E(u))$ and write $U_\kappa \subset \mathbb{A}_A^1$ for the open obtained by inverting $(u - \kappa'(\pi))$ for each $\kappa' \neq \kappa$ with $\kappa'|_k = \kappa_0$. Then $U = \bigcap U_\kappa$ and if A is an E -algebra then the U_κ form an open cover of \mathbb{A}_A^1 .

Note that an A -valued point of Gr_{κ_0} is the same thing as a rank d vector bundle on \mathbb{A}_A^1 which is trivial over U while an A -valued point of $\text{Gr} \otimes_{\mathcal{O}_{K,\kappa}} \mathcal{O}$ is likewise vector bundle trivial over $\bigcup_{\kappa' \neq \kappa} U_{\kappa'}$.

The map in the lemma can therefore be expressed as $\mathcal{E} \mapsto (\mathcal{E}_\kappa)$ where \mathcal{E}_κ is the vector bundle obtained by glueing $\mathcal{E}|_{U_\kappa}$ with the trivial bundle on $\bigcup_{\kappa' \neq \kappa} U_{\kappa'}$. The inverse of this map sends (\mathcal{E}_κ) onto the vector bundle obtained by glueing the $\mathcal{E}_\kappa|_{U_\kappa}$. Concretely, this glueing corresponds to taking the intersection of each of the \mathcal{E}_κ 's which gives the lemma. □

4.7 We define one last ind-scheme $\text{Gr}_{\mathcal{O}}$ whose A -points now classify rank d projective $(A \otimes_{\mathbb{Z}_p} W(k))[u]$ -modules satisfying

$$E(u)^a (A \otimes_{\mathbb{Z}_p} W(k))[u]^d \subset \mathcal{E} \subset E(u)^{-a} (A \otimes_{\mathbb{Z}_p} W(k))[u]^d$$

for some $a \geq 0$. From 2.4 we see that

$$\text{Gr}_{\mathcal{O}} \cong \prod_{\kappa_0} \text{Gr}_{\kappa_0}$$

Lemma 4.6 implies that the generic fibre of $\text{Gr}_{\mathcal{O}}$ identifies with $\prod_{\kappa} (\text{Gr} \otimes_{\mathcal{O}_{K,\kappa}} E)$ with the product running over all embeddings $\kappa : K \rightarrow E$. Note also that the analogue of Lemma 4.5 applies to $\text{Gr}_{\mathcal{O}}$ and identifies its points valued in p -adically complete \mathcal{O} -algebras A with rank d projective \mathfrak{S}_A -modules satisfying

$$E(u)^a \mathfrak{S}_A^d \subset \mathcal{E} \subset E(u)^{-a} \mathfrak{S}_A^d$$

Definition 4.8 Let $\mu = (\mu_{\kappa})$ be a Hodge type, i.e. a collection of dominant $\mu_{\kappa} \in X(T)$ indexed by embeddings $\kappa : K \rightarrow E$. Then we define M_{μ} as the closure in $\text{Gr}_{\mathcal{O}}$ of

$$\prod_{\kappa} (G/P_{\mu_{\kappa}} \otimes_{\mathcal{O}_{K,\kappa}} E) \hookrightarrow \prod_{\kappa} (\text{Gr} \otimes_{\mathcal{O}_{K,\kappa}} E) \cong \text{Gr}_{\mathcal{O}} \otimes_{\mathcal{O}} E$$

Lemma 4.9 Let μ be a Hodge type and suppose $n_{\kappa} \geq 0$ so that $\mu_{\kappa,d} \geq -n_{\kappa}$ for every κ .

- (1) Let A be an E -algebra. Then $\mathcal{E} \in \text{Gr}_{\mathcal{O}}(A)$ is contained in M_{μ} if and only if there are filtrations $\text{Fil}_{\kappa}^{\bullet}$ on A^d of type μ_{κ} so that

$$\left(\prod_{\kappa} E_{\kappa}(u)^{n_{\kappa}} \right) \mathcal{E} = \bigcap_{\kappa} \left(\sum_{i \geq \mu_{\kappa,d} + n_{\kappa}} E_{\kappa}(u)^{i+n_{\kappa}} (A \otimes_{\mathbb{Z}_p} W(k))[u] \text{Fil}_{\kappa}^{-i} \right)$$

(recall the elements $E_{\kappa}(u)$ from 2.4).

- (2) Let A be a Noetherian \mathcal{O} -algebra and suppose there are A -submodules

$$\dots \subset \text{Fil}_{\kappa}^{i+1} \subset \text{Fil}_{\kappa}^i \subset \text{Fil}_{\kappa}^{i-1} \subset \dots \subset A^d$$

for each κ such that $\text{Fil}_{\kappa}^i / \text{Fil}_{\kappa}^{i+1}$ becomes $A[\frac{1}{p}]$ -projective of constant rank after inverting p . If $\mathcal{E} \in \text{Gr}(A)$, when viewed as an \mathfrak{S}_A -module, can be expressed as

$$\left(\prod_{\kappa} E_{\kappa}(u)^{n_{\kappa}} \right) \mathcal{E} = \bigcap_{\kappa} \left(\sum_{i \geq \mu_{\kappa,d} + n_{\kappa}} E_{\kappa}(u)^{i+n_{\kappa}} \mathfrak{S}_A \text{Fil}_{\kappa}^{-i} \right)$$

then $\mathcal{E} \in M_{\mu}(A)$ for μ_{κ} the type of $\text{Fil}_{\kappa}[\frac{1}{p}]^{\bullet}$.

- (3) If A is the ring of integers in a finite extension of E then every A -valued point of M_{μ} is as in (2).

Proof Note that multiplication by $(\prod_{\kappa} E_{\kappa}(u)^{n_{\kappa}})$ identifies M_{μ} with $M_{\mu'}$ for $\mu'_{\kappa} = \mu_{\kappa} + (n_{\kappa})$. Thus we can assume $n_{\kappa} = 0$ throughout.

For (1) we first decompose $\mathcal{E} = \prod_{\kappa_0} \mathcal{E}_{\kappa_0} \in \prod_{\kappa_0} \text{Gr}_{\kappa_0}$ according to the action of $W(k)$. Then Lemma 4.6 and the description of $G/P_{\mu_\kappa} \hookrightarrow \text{Gr}$ from 4.4 implies $\mathcal{E} \in M_\mu$ if and only if, for each κ_0 ,

$$\mathcal{E}_{\kappa_0} = \bigcap_{\kappa_\kappa | \kappa = \kappa_0} \left(\sum_{i \geq \lambda_d} (u - \kappa(\pi))^i A[u] \text{Fil}_\kappa^{-i} \right)$$

for filtrations $\text{Fil}_\kappa^\bullet$ on A^d of type μ_κ . Since $\lambda_d \geq 0$ we have $\mathcal{E}_{\kappa_0} \subset A[u]^d$ for each κ_0 and so

$$\mathcal{E} = \bigcap_{\kappa_0} \left(A[u]^d \times \dots \times A[u]^d \times \underbrace{\mathcal{E}_{\kappa_0}}_{\kappa_0\text{-th position}} \times A[u]^d \times \dots \times A[u]^d \right)$$

Thus, to prove (1) we just need to identify the κ_0 -th term inside this intersection with $\bigcap_{\kappa_\kappa | \kappa = \kappa_0} \left(\sum_{i \geq \mu_{\kappa,d}} E_\kappa(u)^i (A \otimes_{\mathbb{Z}_p} W(k))[u] \text{Fil}_\kappa^{-i} \right)$. This is clear since $E_\kappa(u)$ corresponds to $(1, \dots, 1, (u - \kappa(\pi)), 1, \dots, 1)$ under the identification $(\mathcal{O} \otimes_{\mathbb{Z}_p} W(k))[u] \cong \prod_{\kappa_0} A[u]$.

For part (2) we use that A is Noetherian to ensure $(A \otimes_{\mathbb{Z}_p} W(k))[u] \rightarrow \mathfrak{S}_A$ is flat. Thus $\otimes_{(A \otimes_{\mathbb{Z}_p} W(k))[u]} \mathfrak{S}_A$ commutes with finite intersections and so

$$\mathcal{E} = \left(\bigcap_{\kappa} \left(\sum_{i \geq \mu_{\kappa,d}} E_\kappa(u)^{i+n} (A \otimes_{\mathbb{Z}_p} W(k))[u] \text{Fil}_\kappa^{-i} \right) \right) \otimes_{(A \otimes_{\mathbb{Z}_p} W(k))[u]} \mathfrak{S}_A$$

As a consequence of (1) it follows that $\mathcal{E}[\frac{1}{p}] \in M_\mu$ and so $\mathcal{E} \in M_\mu$ also.

Part (3) relies on the fact that, for A as in the proposition, being A -projective is equivalent to being p -torsion free and finitely generated. Applying (1) to $\mathcal{E}[\frac{1}{p}]$ produces filtrations on $A[\frac{1}{p}]^d$ for each κ . If Fil_κ^i denotes the intersection of this filtration with A^d then the graded pieces are p -torsionfree. This is equivalent to asking that each $\mathfrak{S}_A^d / \left(\sum_{i \geq \mu_{\kappa,d}} E_{ij}(u)^{i+n} \mathfrak{S}_A \text{Fil}_\kappa^{-i} \right)$ is p -torsionfree. Therefore $\mathfrak{S}_A^d / \bigcap_{\kappa} \left(\sum_{i \geq \mu_{\kappa,d}} E_{ij}(u)^{i+n} \mathfrak{S}_A \text{Fil}_\kappa^{-i} \right)$ is also p -torsionfree, and so

$$\mathcal{E}' = \bigcap_{\kappa} \left(\sum_{i \geq \mu_{\kappa,d}} E_{ij}(u)^{i+n} \mathfrak{S}_A \text{Fil}_\kappa^{-i} \right)$$

is \mathfrak{S}_A -projective and $\mathcal{E}' \in \text{Gr}_{\mathcal{O}}$. Since $\mathcal{E}'[\frac{1}{p}] = \mathcal{E}[\frac{1}{p}]$ the valuative criterion for properness implies $\mathcal{E}' = \mathcal{E}$. □

5 Cohomology

5.1 Recall $G = \text{GL}_d$ viewed as an algebraic group over \mathcal{O}_K . Let $\lambda \in X(T)$ be dominant and set

$$\dots \subset \text{Fil}^{n+1} \subset \text{Fil}^n \subset \text{Fil}^{n-1} \subset \dots \subset \mathcal{O}_{G/P_\lambda}^d$$

equal to the universal filtration on G/P_λ of type λ . Then

$$\mathcal{L}(\lambda) := \bigotimes_n \det(\text{Fil}^{-n} / \text{Fil}^{-n+1})^{\otimes n}$$

is a G -equivariant line bundle on G/P_λ and $H^0(G/P_\lambda, \mathcal{L}(\lambda)^{\otimes n})$ can be viewed as an algebraic representation of G on a flat \mathcal{O}_K -module.

Lemma 5.2 *If $\lambda = (\lambda_1, \dots, \lambda_d) \in X(T)$ is such that $\lambda_i > \lambda_{i+1}$ and $n \geq 1$ then*

$$H^0(n\lambda) := H^0(G/P_\lambda, \mathcal{L}(\lambda)^{\otimes n}) \otimes_{\mathcal{O}_K} K$$

is the unique irreducible algebraic representation of G over K with highest weight $n\lambda$.

Proof It follows from [15, I.5.12] that

$$H^0(G/P_\lambda, \mathcal{L}(\lambda)^{\otimes n}) \otimes_{\mathcal{O}_K} K = \text{Ind}_{P_\lambda}^G(\lambda_*(n))$$

where $\lambda_*(n)$ is the character through which T acts on the fibre of $\mathcal{L}(\lambda)$ over the class of 1 in G/P_λ . Since $\lambda_i > \lambda_{i+1}$ we have $P_\lambda = B^-$ the opposite Borel consisting of lower triangular matrices and the class of 1 in G/B^- is the standard filtration of type λ , i.e. the filtration with

$$\text{Fil}^{-\lambda_i} / \text{Fil}^{-\lambda_i+1} = Ae_i \oplus \dots \oplus Ae_d / Ae_{i+1} \oplus \dots \oplus Ae_d$$

As the fibre of $\mathcal{L}(\lambda)$ over this standard filtration is

$$\bigotimes_{i=1}^d \det(\text{Fil}^{-\lambda_i} / \text{Fil}^{-\lambda_i+1})^{\otimes n\lambda_i}$$

we see that T acts on this fibre via the character $\text{diag}(a_1, \dots, a_d) \mapsto \prod a_i^{n\lambda_i}$. The representation $\text{Ind}_{B^-}^G(\lambda_*(n))$ therefore has highest weight $n\lambda$ by [15, II.2.2] and is irreducible in characteristic zero by [15, II.5.6]. □

5.3 On Gr there is an ample G -equivariant line bundle \mathcal{L}_{\det} whose fibre over any A -valued point $\mathcal{E} \in \text{Gr}(A)$ with $\mathcal{E} \subset (u - \pi)^{-a} \mathfrak{S}_A^d$ for $a \geq 0$ is given by

$$\det(\mathcal{E}_0/\mathcal{E}) \otimes \det(\mathcal{E}_0/A[u]^d)^{-1}$$

for any $\mathcal{E}_0 \in \text{Gr}$ with \mathcal{E} , $A[u]^d \subset \mathcal{E}_0$. Note this is G -equivariantly independent of \mathcal{E}_0 . We also write \mathcal{L}_{\det} for the G -equivariant ample line bundle constructed analogously on $\text{Gr}_{\mathcal{O}}$.

Lemma 5.4 *The restriction of \mathcal{L}_{\det} to G/P_λ inside Gr identifies G -equivariantly with $\mathcal{L}(\lambda)$.*

Proof Suppose $\mathcal{E} \in \text{Gr}_K$ corresponds to an A -valued point in the image of $G/P_\lambda \rightarrow \text{Gr}_K$. Then $\mathcal{E} = \sum_{i \geq \lambda_d} (u - \pi)^i A[u] \text{Fil}^{-i}$ for a filtration Fil^\bullet of type λ on A^d and so, as A -modules, $(u - \pi)^{\lambda_d} A[u]/\mathcal{E} = \bigoplus_{i \geq \lambda_d} A^d / \text{Fil}^{-i}$. Thus,

$$\det((u - \pi)^{\lambda_d} A[u]/\mathcal{E}) = \bigotimes_{i \geq \lambda_d} \det(\text{Fil}^{-i} / \text{Fil}^{-i+1})^{\otimes(i-\lambda_d)}$$

and so the fibre of \mathcal{L}_{\det} over \mathcal{E} equals

$$\begin{cases} \bigotimes_{i \geq \lambda_d} \det(\text{Fil}^{-i} / \text{Fil}^{-i+1})^{\otimes(i-\lambda_d)} \otimes \det((A[u]^d / (u - \pi)^{\lambda_d} A[u]^d)) & \text{if } \lambda_d \geq 0 \\ \bigotimes_{i \geq \lambda_d} \det(\text{Fil}^{-i} / \text{Fil}^{-i+1})^{\otimes(i-\lambda_d)} \otimes \det(((u - \pi)^{\lambda_d} A[u]^d / A[u]^d))^{-1} & \text{if } \lambda_d \leq 0 \end{cases}$$

In either case, the second factor in these tensor products can be identified with $\bigotimes_{i \geq \lambda_d} \det(\text{Fil}^{-i} / \text{Fil}^{-i+1})^{\lambda_d}$, which finishes the proof. \square

Corollary 5.5 *For any $n > 0$, there is an identification*

$$H^0(M_\mu \otimes_{\mathcal{O}} E, \mathcal{L}_{\det}^{\otimes n}) = \bigotimes_K (H^0(n\mu_\kappa) \otimes_{K,\kappa} E)$$

of G -representations.

Proof Let $p_\kappa : M_\mu \otimes_{\mathcal{O}} E \rightarrow G/P_{\mu_\kappa} \otimes_{\mathcal{O}_{K,\kappa}} E$ be the κ -th projection. Then the restriction of \mathcal{L}_{\det} on $\text{Gr}_{\mathcal{O}}$ to $M_\mu \otimes_{\mathcal{O}} E$ coincides with $\bigotimes_\kappa p_\kappa^*(\mathcal{L}_{\det} \otimes_{\mathcal{O}_{K,\kappa}} E)$ where \mathcal{L}_{\det} here denotes the restriction to G/P_{μ_κ} of the determinant line bundle on Gr . The Kunnetth formula [26, 0BED] gives

$$H^0(M_\mu \otimes_{\mathcal{O}} E, \mathcal{L}_{\det}^{\otimes n}) = \bigotimes_\kappa H^0(G/P_{\mu_\kappa}, \mathcal{L}_{\det}^{\otimes n}) \otimes_{\mathcal{O}_{K,\kappa}} E$$

as G -representations. Therefore, we just have to show $H^0(G/P_{\mu_\kappa}, \mathcal{L}_{\det}^{\otimes n}) \otimes_{\mathcal{O}_K} K = H^0(n\mu_\kappa)$ as G -representations, and this follows from Lemma 5.4. \square

6 Multiplicity bounds

6.1 The formal character of an algebraic representation V of G on a finite dimensional vector space is defined as

$$\text{ch}(V) = \sum_\lambda V_\lambda e(\lambda) \in \mathbb{Z}[X(T)]$$

where V_λ is the λ -weight space of V and $e(\lambda)$ denotes λ viewed as an element of the group ring $\mathbb{Z}[X(T)]$. This induces an isomorphism between the Grothendieck group of such representations and $\mathbb{Z}[X(T)]^W$ where W denotes the Weyl group of G [15, II.5.7].

6.2 For $\lambda \in X(T)$ dominant recall the G -representation $H^0(\lambda)$ over K from 5.1. Weyl’s character formula gives

$$\text{ch}(H^0(\lambda)) = \frac{A(\lambda + \rho)}{A(\rho)}, \quad \rho := (d - 1, d - 2, \dots, 1, 0) \in X(T)$$

where $A(\lambda) := \sum_{w \in W} \det(w)e(w\lambda)$ [15, II.5.10]. If we write $\dim : \mathbb{Z}[X(T)] \rightarrow \mathbb{Z}$ for the map $\sum_\lambda a_\lambda e(\lambda) \mapsto \sum a_\lambda$ then one also has

$$\dim \text{ch}(H^0(\lambda)) = \dim H^0(\lambda) = \prod_{i > j} \frac{\lambda_j - \lambda_i + i - j}{i - j}$$

Though here $H^0(\lambda)$ is defined over a field of characteristic zero, all the above goes through with $H^0(\lambda)$ replaced by the representation over a field of characteristic p of highest weight λ . What differs in characteristic p is that this highest weight representation may not be irreducible.

Lemma 6.3 *Let $\mu_1, \dots, \mu_e \in X(T)$ with $\mu_i - \rho$ dominant for each i and suppose that*

$$\text{ch} \left(\bigotimes_{i=1}^e H^0(\mu_i - \rho) \right) = \sum_{\lambda \in X(T)} m(\lambda, \mu) \text{ch} H^0(\lambda)$$

for $m(\lambda, \mu) \in \mathbb{Z}$. Then

$$\dim \left(\bigotimes_{i=1}^e H^0(n\mu_i) \right) - \sum_{\lambda \in X(T)} m(\lambda, \mu) \dim H^0(n(\lambda + \rho)) \text{ch} \left(H^0(n\rho)^{\otimes(e-1)} \right)$$

is a polynomial in n of degree $< \sum_i \dim G/P_{\mu_i}$.

Proof Using Weyl’s character formula from 6.2 and multiplying by $A(\rho)^e$ gives $\prod_i A(\mu_i) = \sum_\lambda m(\lambda, \mu) A(\lambda + \rho) A(\rho)^{e-1}$. Taking the image of this identity under the endomorphism of $\mathbb{Z}[X(T)]$ induced by multiplication by n on $X(T)$ gives

$$\prod_i A(n\mu_i) = \sum_\lambda m(\lambda, \mu) A(n(\lambda + \rho)) A(n\rho)^{e-1}$$

(because the formation of A commutes with this endomorphism). Dividing by $A(\rho)^e$ then gives

$$\prod_i \text{ch}(H^0(n\mu_i - \rho)) = \sum_\lambda m(\lambda, \mu) \text{ch}(H^0(n(\lambda + \rho) - \rho)) \left(\text{ch}(H^0(n\rho - \rho)) \right)^{e-1}$$

The lemma therefore follows by considering dimensions and observing that the difference $\dim H^0(n\lambda) - \dim H^0(n\lambda - \rho)$ is a polynomial in n of degree $< \dim G/P_\lambda$ for any dominant $\lambda \in X(T)$ (use the last equation from 6.2). \square

Remark 6.4 Since K has characteristic zero each $H^0(\lambda)$ is irreducible. Moreover, every irreducible G -representation is isomorphic to one such $H^0(\lambda)$. The observation from 6.2 that ch induces an identification between $\mathbb{Z}[X(T)]$ and the Grothendieck group of G -representations shows that the integers $m(\lambda, \mu)$ in the previous lemma are ≥ 0 and are uniquely determined.

Notation 6.5 Recall the fixed elements $\tilde{\kappa}_0 : K \rightarrow E$ lifting the κ_0 from 2.1. Then, for any tuple $\lambda = (\lambda_{\kappa_0})_{\kappa_0:k \rightarrow \mathbb{F}}$ of dominant $\lambda_{\kappa_0} \in X(T)$ write $\tilde{\lambda} = (\tilde{\lambda}_\kappa)$ for the Hodge type defined by

$$\tilde{\lambda}_\kappa = \begin{cases} \lambda_{\kappa_0} + \rho & \text{if } \kappa = \tilde{\kappa}_0 \\ \rho & \text{otherwise} \end{cases}$$

Proposition 6.6 *Let μ be a Hodge type with $\mu_\kappa - \rho$ dominant for every κ and suppose that*

$$[M_\mu \otimes_{\mathcal{O}} \mathbb{F}] = \sum n(\lambda, \mu) [M_{\tilde{\lambda}} \otimes_{\mathcal{O}} \mathbb{F}]$$

for integers

$$n(\lambda, \mu) \geq \prod_{\kappa_0} m(\lambda_{\kappa_0}, (\mu_\kappa)_{\kappa|_{\kappa=\kappa_0}})$$

Then this inequality is an equality for each λ .

Proof We apply Lemma 3.2 to the line bundle \mathcal{L}_{\det} . This gives that

$$\dim H^0(\overline{M}_\mu, \mathcal{L}_{\det}^{\otimes n}) - \sum_{\lambda} n(\lambda, \mu) \dim H^0(\overline{M}_{\tilde{\lambda}}, \mathcal{L}_{\det}^{\otimes n})$$

is, for large n , equal the value at n of a polynomial of degree $< \sum_{\kappa} \dim G/P_{\mu_\kappa}$. Since M_μ is \mathcal{O} -flat the same is true of the coherent sheaf $\mathcal{L}_{\det}^{\otimes n}$. It therefore follows that the Euler characteristic of $\mathcal{L}_{\det}^{\otimes n}$ on \overline{M}_μ equals the Euler characteristic of $\mathcal{L}_{\det}^{\otimes n}$ on $M_\mu \otimes_{\mathcal{O}} E$ (see for example [20, 5.3.28]). Since \mathcal{L}_{\det} is ample we conclude that

$$\dim_{\mathbb{F}} H^0(\overline{M}_\mu, \mathcal{L}_{\det}^{\otimes n}) = \dim_E H^0(M_\mu \otimes_{\mathcal{O}} E, \mathcal{L}_{\det}^{\otimes n})$$

for sufficiently large n . Therefore, Corollary 5.5 shows that

$$\prod_{\kappa} \dim(H^0(n\mu_\kappa)) - \sum_{\lambda=(\lambda_{\kappa_0})} n(\lambda, \mu) \prod_{\kappa_0} (\dim(H^0(n(\lambda_{\kappa_0} + \rho))) \dim(H^0(n\rho))^{e-1})$$

is also, for large n , equal the value at n of a polynomial of degree $< \sum_{\kappa} \dim G/P_{\mu_{\kappa}}$. Set $m_{\lambda} = \prod_{\kappa_0} m(\lambda_{\kappa_0}, (\mu_{\kappa})_{\kappa|_{\kappa=\kappa_0}})$. Lemma 6.3 gives that

$$\prod_{\kappa} \dim(H^0(n\mu_{\kappa})) - \sum_{\lambda=(\lambda_{\kappa_0})} m_{\lambda} \prod_{\kappa_0} \left(\dim(H^0(n(\lambda_{\kappa_0} + \rho))) \dim(H^0(n\rho))^{e-1} \right)$$

is a polynomial of degree $< \sum_{\kappa} \dim G/P_{\mu_{\kappa}}$ in n . We conclude that the dimension of

$$\sum_{\lambda=(\lambda_{\kappa_0})} (n(\lambda, \mu) - m(\lambda, \mu)) \prod_{\kappa_0} \left(\dim(H^0(n(\lambda_{\kappa_0} + \rho))) \dim(H^0(n\rho))^{e-1} \right)$$

is also polynomial of degree $< \sum_{\kappa} \dim G/P_{\mu_{\kappa}}$ in n for $n \gg 0$. Since $n(\lambda, \mu) - m(\lambda, \mu) \geq 0$, each term in the above sum is a polynomial in n of degree $\sum_{\kappa} \dim G/P_{\mu_{\kappa}}$ with non-negative leading term. We must therefore have $n(\lambda, \mu) = m(\lambda, \mu)$ for each λ . □

7 Topological descriptions

7.1 Recall that for $\lambda, \lambda' \in X(T)$ we write

$$\lambda' \leq \lambda$$

if $\lambda_d + \dots + \lambda_i \leq \lambda'_d + \dots + \lambda'_i$ for each i with equality when $i = 1$.

Proposition 7.2 *Let μ be a Hodge type with $\mu_{\kappa} - \rho$ dominant for each κ . Assume that*

$$\sum_{\kappa|_{k=\kappa_0}} \mu_{\kappa,1} - \mu_{\kappa,d} \leq e + p - 1$$

for each $\kappa_0 : k \rightarrow \mathbb{F}$. Then:

(1) *There are integers $n(\lambda, \mu) \in \mathbb{Z}$ such that*

$$[M_{\mu} \otimes_{\mathcal{O}} \mathbb{F}] = \sum_{\lambda} n(\lambda, \mu) [M_{\tilde{\lambda}} \otimes_{\mathcal{O}} \mathbb{F}]$$

with the sum running over tuples $\lambda = (\lambda_{\kappa_0})$ with each $\lambda_{\kappa_0} \in X(T)$ dominant and satisfying $\lambda_{\kappa_0} \leq \sum_{\kappa|_{k=\kappa_0}} (\mu_{\kappa} - \rho)$ and $\tilde{\lambda}$ defined as in Notation 6.5.

(2) *If $d = 2$ then each $M_{\tilde{\lambda}} \otimes_{\mathcal{O}} \mathbb{F}$ appearing in this sum is irreducible, generically reduced, and pairwise distinct.*

In particular, when $d = 2$ part (2) implies that the $n(\lambda, \mu)$ in part (1) are ≥ 0 .

7.3 To prove the proposition we will approximate $M_{\mu} \otimes_{\mathcal{O}} \mathbb{F}$ via explicit moduli conditions. In fact we give two such moduli interpretations, based on the following two operators:

- For any \mathcal{O} -algebra A set

$$\nabla := u \frac{d}{du} : \mathfrak{S}_A \left[\frac{1}{E(u)} \right] \rightarrow \mathfrak{S}_A \left[\frac{1}{E(u)} \right]$$

We also write ∇ for the coordinate-wise extension to $\mathfrak{S}_A[\frac{1}{E(u)}]^d$.

- If A is a p -adically complete \mathcal{O} -algebra of topologically finite type then for each $\sigma \in G_K$ we can also define

$$\nabla_\sigma := \frac{\sigma - \text{Id}}{\mu} : A_{\text{inf},A} \left[\frac{1}{\mu} \right] \rightarrow A_{\text{inf},A} \left[\frac{1}{\mu} \right]$$

Note that this is well defined because $\sigma(\mu)$ and μ generate the same ideal in A_{inf} (as follows from e.g. Lemma 2.3). Again write ∇_σ also for the coordinate-wise extension to $A_{\text{inf},A}[\frac{1}{\mu}]^d$.

The advantage of ∇ is that it is easier to compute with. The advantage of ∇_σ is that it is more directly related to Galois representations.

Lemma 7.4 *There exist closed subfunctors $\text{Gr}_{\mathcal{O}}^{\nabla_\sigma}, \text{Gr}_{\mathcal{O}}^\nabla \subset \text{Gr}_{\mathcal{O}}$ such that*

- (1) $\mathcal{E} \in \text{Gr}^\nabla(A)$ if and only if

$$E(u)\nabla(\mathcal{E}) \subset u\mathcal{E}$$

as submodules of $\mathfrak{S}_A[\frac{1}{E(u)}]^d$.

- (2) Whenever A is a p -adically complete \mathcal{O} -algebra of topologically finite type then $\mathcal{E} \in \text{Gr}^{\nabla_\sigma}(A)$ if and only if

$$E(u)\nabla_\sigma(\mathcal{E}) \subset u\mathcal{E} \otimes_{\mathfrak{S}_A} A_{\text{inf},A}$$

as submodules of $A_{\text{inf},A}[\frac{1}{\mu}]^d$ for every $\sigma \in G_K$.

Remark 7.5 Since $E(u)$ and $\frac{\mu}{\varphi^{-1}(\mu)}$ generate the same ideal in A_{inf} the condition defining $\text{Gr}_{\mathcal{O}}^{\nabla_\sigma}$ can also be expressed as

$$(\sigma - 1)(\mathcal{E}) \subset u\varphi^{-1}(\mu)\mathcal{E} \otimes_{\mathfrak{S}_A} A_{\text{inf},A}$$

This description may be more familiar from the point of view of crystalline Breuil–Kisin modules.

Proof That $\text{Gr}_{\mathcal{O}}^\nabla$ is a closed subfunctor is clear, so we focus on $\text{Gr}_{\mathcal{O}}^{\nabla_\sigma}$. First, suppose that A is any Noetherian \mathcal{O}/p^n -algebra, for some $n \geq 1$, and that $\mathcal{E} \in \text{Gr}_{\mathcal{O}}(A)$. An application of [7, B.29] shows that the condition

$$E(u)\nabla_\sigma(\mathcal{E}) \subset u\mathcal{E} \otimes_{\mathfrak{S}_A} A_{\text{inf},A} \tag{7.6}$$

is closed in Spec A . More precisely, we specialise the general setting of [7, B.11] to $R = W(k)/p^n$, $C^+ = A_{\text{inf}}/p^n$, and u equal to $u\varphi^{-1}(\mu) \in C^+$. Then, for each $m \in \mathcal{E}$ and each $\sigma \in G_K$, apply [7, B.29] with $M = \mathcal{E} \otimes_{\mathfrak{S}_A} u\varphi^{-1}(\mu)A_{\text{inf},A}$ and $x = (\sigma - 1)(m) \in M$. This implies that the condition

$$(\sigma - 1)(m) \in u\varphi^{-1}(\mu)\mathcal{E} \otimes_{\mathfrak{S}_A} A_{\text{inf},A}$$

is closed in Spec A . Taking the intersection of the resulting ideals as m runs over all elements of \mathcal{E} and considering Remark 7.5 shows (7.6) is closed in Spec A .

Since $\text{Gr}_{\mathcal{O}}$ is of ind-finite type over Spec \mathcal{O} the previous paragraph shows that for each $n \geq 1$ there exists closed ind-subchemes $\text{Gr}_{\mathcal{O},n}^{\nabla\sigma} \subset \text{Gr}_{\mathcal{O}} \otimes_{\mathcal{O}} \mathcal{O}/p^n$ which are compatible with the closed immersions $\text{Gr}_{\mathcal{O}} \otimes_{\mathcal{O}} \mathcal{O}/p^n \hookrightarrow \text{Gr}_{\mathcal{O}} \otimes_{\mathcal{O}} \mathcal{O}/p^{n-1}$. Since $\text{Gr}_{\mathcal{O},1}^{\nabla\sigma}$ is proper over \mathcal{O}/p (since $\text{Gr}_{\mathcal{O}}$ is itself ind-projective over \mathcal{O}) it follows from [26, Tag 0899] that there exists a closed ind-subscheme $\text{Gr}_{\mathcal{O}}^{\nabla\sigma}$ of $\text{Gr}_{\mathcal{O}}$ with $\text{Gr}_{\mathcal{O},n}^{\nabla\sigma} = \text{Gr}_{\mathcal{O}}^{\nabla\sigma} \otimes_{\mathcal{O}} \mathcal{O}/p^n$ for each n . This closed ind-subscheme has the desired property because if $\mathcal{E} \in \text{Gr}_{\mathcal{O}}(A)$ for A a p -adically complete \mathcal{O} -algebra of topologically finite type then $\mathcal{E} \in \text{Gr}_{\mathcal{O}}^{\nabla\sigma}(A)$ if and only if $\mathcal{E}_n = \mathcal{E} \otimes_{\mathcal{O}} \mathcal{O}/p^n \in \text{Gr}_{\mathcal{O},n}^{\nabla\sigma}(A/p^n)$ for all $n \geq 1$, i.e. if and only if $E(u)\nabla_{\sigma}(\mathcal{E}_n) \subset u\mathcal{E}_n \otimes_{\mathfrak{S}_A} A_{\text{inf},A}$ for all $n \geq 1$. Since A is p -adically complete and \mathcal{E} is \mathfrak{S}_A -projective one has $\mathcal{E} = \varprojlim \mathcal{E}_n$ and so this is equivalent to asking that $E(u)\nabla_{\sigma}(\mathcal{E}) \subset u\mathcal{E}$, as required. \square

Proposition 7.7 *For every Hodge type μ one has $M_{\mu} \subset \text{Gr}_{\mathcal{O}}^{\nabla\sigma}$ and $M_{\mu} \subset \text{Gr}_{\mathcal{O}}^{\nabla}$.*

Proof Since $\text{Gr}_{\mathcal{O}}^{\nabla}$ is closed it suffices to show $\mathcal{E} \in M_{\mu}(A)$ is contained in $\text{Gr}_{\mathcal{O}}^{\nabla}$ for any E -algebra A . Lemma 4.9 allows us to write \mathcal{E} as an intersection of $\mathcal{E}_{\kappa} = \frac{1}{E(u)^n} \sum E_{\kappa}(u)^{i+n} (A \otimes_{\mathbb{Z}_p} W(k))[u] \text{Fil}_{\kappa}^{-i}$ for some $n \geq 0$. It is therefore enough to show that $E(u)\nabla(\mathcal{E}_{\kappa}) \subset u\mathcal{E}_{\kappa}$ and this follows since

$$\nabla(\mathcal{E}_{\kappa}) = \sum_i \nabla \left(\frac{E_{\kappa}(u)^{i+n}}{E(u)^n} \right) (A \otimes_{\mathbb{Z}_p} W(k))[u] \text{Fil}_{\kappa}^{-i}$$

and $E(u)\nabla\left(\frac{E_{\kappa}(u)^{i+n}}{E(u)^n}\right) \in u\frac{E_{\kappa}(u)^{i+n}}{E(u)^n} (A \otimes_{\mathbb{Z}_p} W(k))[u]$.

There is a slight difficulty in giving an identical argument to show $M_{\mu} \subset \text{Gr}_{\mathcal{O}}^{\nabla\sigma}$ because the moduli description for $\text{Gr}_{\mathcal{O}}^{\nabla\sigma}$ does not apply when A is an E -algebra. To address this we first note that the generic fibre of M_{μ} is reduced so to show $M_{\mu} \otimes_{\mathcal{O}} E \subset \text{Gr}_{\mathcal{O}}^{\nabla\sigma}$ it suffices to show this on A -points whenever A is a finite extension of E . By the valuative criterion for properness, any such A -valued point is induced from a point valued in the ring of integers of A . Thus we are reduced to showing $M_{\mu}(A) \subset \text{Gr}_{\mathcal{O}}^{\nabla\sigma}(A)$ whenever A is the ring of integers in a finite extension of E . Using part (3) of Lemma 4.9 this comes down to proving that

$$E(u)\nabla_{\sigma}(\mathcal{E}_{\kappa}) \subset u\mathcal{E}_{\kappa} \otimes_{\mathfrak{S}_A} A_{\text{inf},A}$$

for $\mathcal{E}_\kappa = \frac{1}{E(u)^n} \sum E_\kappa(u)^{i+n} \mathfrak{S}_A \text{Fil}_\mathfrak{g}^{-i}$. This would follow from the claim that

$$E(u) \nabla_\sigma \left(\frac{E_\kappa(u)^{i+n}}{E(u)^n} \right) \in u \frac{E_\kappa(u)^{i+n}}{E(u)^n} A_{\text{inf}, \mathcal{O}}$$

To prove the claim first note that $\sigma(E_\kappa(u)) - E_\kappa(u) = \sigma(u) - u \in u\mu A_{\text{inf}}$. Similarly $\sigma(E(u)) - E(u) \in u\mu A_{\text{inf}}$. Writing

$$\nabla_\sigma(E_\kappa(u)^i) = \nabla_\sigma(E_\kappa(u)^{i-1})\sigma(E_\kappa(u)) + E_\kappa(u)\nabla_\sigma(E_\kappa(u))$$

and arguing by induction on i then gives that $\nabla_\sigma(E_\kappa(u)^i) \in uE_\kappa(u)^{i-1}A_{\text{inf}, \mathcal{O}}$. Similarly $\nabla_\sigma(E(u)^i) \in uE(u)^{i-1}A_{\text{inf}}$. Since we can write

$$E(u)^n(\sigma - 1)\left(\frac{E_\kappa(u)^{i+n}}{E(u)^n}\right) = (\sigma - 1)(E_\kappa(u)^{i+n}) - \frac{\sigma(E_\kappa(u)^{i+n})}{\sigma(E(u)^n)}(\sigma - 1)(E(u)^n)$$

the claim follows. □

Remark 7.8 After possibly replacing the compatible system of primitive p -th power roots of unity ϵ we can choose $\sigma \in G_K$ so that $\sigma(u)/u = [\epsilon]$. Then

$$\nabla_\sigma(u^i) = u^i \left(\frac{(\frac{\sigma(u)}{u})^i - 1}{[\epsilon] - 1} \right) = u^i \left(\frac{[\epsilon]^i - 1}{[\epsilon] - 1} \right) = u^i(1 + [\epsilon] + \dots + [\epsilon]^{i-1})$$

Thus $\nabla_\sigma = u\nabla_q$ where ∇_q is the q -derivation for $q = [\epsilon]$. In particular $\nabla_\sigma \equiv u \frac{d}{du} = \nabla$ modulo $[\epsilon] - 1$. This illustrates the close relationship between the Gr_σ^∇ and the locus $\text{Gr}_\mathcal{O}^\nabla$.

7.9 For the rest of this section we focus on $\text{Gr}_\mathcal{O}^\nabla \otimes_{\mathcal{O}} \mathbb{F}$. Note that since ∇ is $W(k)$ -linear we have

$$\text{Gr}_\mathcal{O}^\nabla = \prod_{\kappa_0} \text{Gr}_{\kappa_0}^\nabla$$

where $\text{Gr}_{\kappa_0}^\nabla$ is defined similarly. Let us write $\overline{\text{Gr}} = \text{Gr}_{\kappa_0} \otimes_{\mathcal{O}} \mathbb{F}$ (note this is independent of κ_0) and $\overline{\text{Gr}}^\nabla = \text{Gr}_{\kappa_0}^\nabla \otimes_{\mathcal{O}} \mathbb{F}$. The description from Lemma 4.5 shows that the group scheme

$$LG^+ : A \mapsto \text{GL}_d(A[[u]])$$

acts on $\overline{\text{Gr}}$. For $\lambda \in X(T)$ dominant we set $\overline{\text{Gr}}_\lambda$ equal to the LG^+ -orbit of $\mathcal{E}_\lambda \in \text{Gr}$ (recall \mathcal{E}_λ is defined in 4.4) and we set $\overline{\text{Gr}}_{\leq \lambda}$ equal to its reduced closure. Then

$$\overline{\text{Gr}}_{\leq \lambda} = \bigcup_{\lambda' \leq \lambda} \overline{\text{Gr}}_{\lambda'}$$

Lemma 7.10 *Suppose $\lambda \in X(T)$ is dominant with*

$$\lambda_1 - \lambda_d \leq e + p - 1$$

Set \mathcal{C}_λ equal to the closure of $\overline{\text{Gr}}_\lambda \cap \overline{\text{Gr}}^\nabla$ in $\overline{\text{Gr}}$. Then \mathcal{C}_λ is reduced and irreducible of dimension

$$\sum_{\kappa_0} \sum_{i < j} \max\{\lambda_i - \lambda_j, e\}$$

Proof We begin by giving an open cover of $\overline{\text{Gr}}_\lambda$: let $\mathcal{U}_\lambda \subset L^+G$ denote the subfunctor whose A -points consist of unipotent upper triangular matrices

$$\begin{pmatrix} 1 & & a_{ij} \\ & \ddots & \\ & & 1 \end{pmatrix} \in L^+G(A)$$

where for each $i > j$, $a_{ij} \in A[u]$ has degree $< \lambda_j - \lambda_i$. Consider the morphism $\mathcal{U}_\lambda \rightarrow \overline{\text{Gr}}_\lambda$ sending $g \mapsto g\mathcal{E}_\lambda$. Recall that $g\mathcal{E}_\lambda \mapsto g_0\mathcal{E}_\lambda$, for $g_0 = g$ modulo u , defines a morphism $\overline{\text{Gr}}_\lambda \rightarrow G/P_\lambda$. Since the parabolic P_λ is contained in the Borel of lower triangular matrices $B^- \subset G$ we can compose this map with $G/P_\lambda \rightarrow G/B^-$. Then the morphism $\mathcal{U} \rightarrow \overline{\text{Gr}}_\lambda$ identifies \mathcal{U}_λ with the preimage under this composite of the open $U \subset G/B^-$ consisting of upper triangular unipotent matrices. In particular, $\mathcal{U}_\lambda \rightarrow \overline{\text{Gr}}_\lambda$ is an open immersion and $\overline{\text{Gr}}_\lambda = \bigcup_w w\mathcal{U}_\lambda$ with w running over the permutation matrices in G (as follows by considering the open cover $G/B^- = \bigcup wU$).

Since $\nabla(w) = 0$ we have $w\mathcal{U}_\lambda \cap \overline{\text{Gr}}^\nabla = w(\mathcal{U}_\lambda \cap \overline{\text{Gr}}^\nabla)$. Therefore the lemma reduces to showing $\mathcal{U}_\lambda \cap \overline{\text{Gr}}^\nabla$ is an affine space of the claimed dimension. Observe that $g \in \mathcal{U}_\lambda \cap \overline{\text{Gr}}^\nabla$ if and only if

$$u^{e-1}g^{-1}\nabla(g) \in \begin{pmatrix} u^{\lambda_1} & & \\ & \ddots & \\ & & u^{\lambda_d} \end{pmatrix} \text{Mat}(A[u]) \begin{pmatrix} u^{-\lambda_1} & & \\ & \ddots & \\ & & u^{-\lambda_d} \end{pmatrix} \tag{7.11}$$

If we write $g^{-1} = (b_{ij})_{ij}$ then, using that $b_{jj} = 1$, $b_{lj} = 0$ for $l < j$, and $\nabla(a_{ii}) = 0$, we see that (7.11) is equivalent to asking that

$$\nabla(a_{ij}) + \sum_{j < l < i} \nabla(a_{il})b_{lj} \in u^{\lambda_j - \lambda_i - e + 1} A[u] \tag{7.12}$$

for every $i > j$. By assumption $\lambda_j - \lambda_i - e + 1 \leq p$ and so $\sum_{j < l < i} \nabla(a_{il})b_{lj}$ modulo $u^{\lambda_j - \lambda_i - e + 1}$ admits an antiderivative; in other words, there exists a unique $X \in uA[u]$ of degree $< \lambda_j - \lambda_i - e + 1$ with

$$\nabla(X) \equiv - \sum_{j < l < i} \nabla(a_{il})b_{lj} \quad \text{modulo } u^{\lambda_j - \lambda_i - e + 1}$$

Since a_{ij} has degree $< \lambda_j - \lambda_i$ it follows that

$$a_{ij} = \begin{cases} X + a_{ij}^{(0)} + u^{\lambda_j - \lambda_i - e + 1} a_{ij}^{(1)} + \dots + u^{\lambda_j - \lambda_i - 1} a_{ij}^{(e-1)} & \text{if } \lambda_j - \lambda_i \geq e \\ X + a_{ij}^{(0)} + u a_{ij}^{(1)} + \dots + u^{\lambda_j - \lambda_i - 1} a_{ij}^{\lambda_i - \lambda_j - 1} & \text{if } \lambda_j - \lambda_i < e \end{cases}$$

for some $a_{ij}^{(l)} \in A$. Note that, for $i > j$, the ij -th entry of $gg^{-1} = 1$ is

$$0 = \sum_{l=0}^d a_{il} b_{lj} = b_{ij} + a_{ij} + \sum_{j < l < i} a_{il} b_{lj}$$

This shows, by an inductive argument, that b_{ij} is a function of a_{lk} for $l < k$ with $k - l \leq i - j$. Therefore the element $X \in uA[u]$ considered above depends on a_{lk} with $k - l < i - j$. As a consequence the morphism

$$U_\lambda \cap \overline{\text{Gr}}^\nabla \rightarrow \prod_{ij} \mathbb{A}_{\mathbb{F}}^{\min\{e, \lambda_j - \lambda_i\}}$$

given by $(a_{ij}) \mapsto (a_{ij}^{(l)})$ has a well-defined inverse which finishes the proof. □

Proof of Proposition First observe that under the identification $\text{Gr}_{\mathcal{O}} \otimes_{\mathcal{O}} \mathbb{F} \cong \prod_{\kappa_0} \overline{\text{Gr}}$ we have $(M_\mu \otimes_{\mathcal{O}} \mathbb{F})_{\text{red}} \hookrightarrow \prod_{\kappa_0} \overline{\text{Gr}}_{\leq \sum_{\kappa|k=\kappa_0} \mu_\kappa}$. Thus $(M_\mu \otimes_{\mathcal{O}} \mathbb{F})_{\text{red}}$ is contained in $\bigcup_{\lambda=(\lambda_{\kappa_0})} \prod \overline{\text{Gr}}_\lambda$ where the product runs over $\lambda = (\lambda_{\kappa_0})$ with $\lambda_{\kappa_0} \leq \sum_{\kappa|k=\kappa_0} \mu_\kappa$. Since $M_\mu \subset \text{Gr}_{\mathcal{O}}^\nabla$ and each $\mu_\kappa - \rho$ is dominant the dimension calculations from Lemma 7.10 imply that

$$(M_\mu \otimes_{\mathcal{O}} \mathbb{F})_{\text{red}} \subset \bigcup_{\lambda=(\lambda_{\kappa_0})} \mathcal{C}_{\lambda+e\rho}$$

where the union now runs over $\lambda = (\lambda_{\kappa_0})$ with $\lambda_{\kappa_0} + e\rho \leq \sum_{\kappa|k=\kappa_0} \mu_\kappa$ and where we write $\mathcal{C}_{\lambda+e\rho} = \prod_{\kappa_0} \mathcal{C}_{\lambda_{\kappa_0}+e\rho}$. Thus, one can write

$$[M_\mu \otimes_{\mathcal{O}} \mathbb{F}] = \sum_{\lambda} n(\lambda, \mu) [\mathcal{C}_{\lambda+e\rho}]$$

as cycles, for integers $n(\lambda, \mu) \geq 0$. Furthermore, since $\overline{\text{Gr}}_{\sum_{\kappa|k=\kappa_0} \mu_\kappa} \subset \overline{\text{Gr}}_{\leq \sum_{\kappa|k=\kappa_0} \mu_\kappa}$ is open it follows that $\mathcal{C}_{\lambda+e\rho} \cap (M_\mu \otimes_{\mathcal{O}} \mathbb{F})$ is open in $(M_\mu \otimes_{\mathcal{O}} \mathbb{F})$ for $\lambda = (\lambda_{\kappa_0})$ with $\lambda_{\kappa_0} = \sum_{\kappa|k=\kappa_0} (\mu_\kappa - \rho)$. Since this intersection is clearly non-empty it follows that $n(\lambda, \mu) = 1$ for this particular λ .

This shows that

$$[M_{\tilde{\lambda}} \otimes_{\mathcal{O}} \mathbb{F}] = [\mathcal{C}_{\lambda+e\rho}]$$

can be expressed as a $\mathbb{Z}_{\geq 0}$ -linear combination of $[\mathcal{C}_{\lambda'+e\rho}]$'s for $\lambda' = (\lambda'_{\kappa_0})$ with $\lambda'_{\kappa_0} \leq \lambda_{\kappa_0}$. Arguing by induction we conclude that we can always write $[\mathcal{C}_{\lambda+e\rho}] = \sum_{\lambda'} n_{\lambda'} [M_{\tilde{\lambda}'} \otimes_{\mathcal{O}} \mathbb{F}]$ for $\lambda' = (\lambda'_{\kappa_0})$ satisfying $\lambda'_{\kappa_0} \leq \lambda_{\kappa_0}$ and some $n_{\lambda'} \in \mathbb{Z}$. This proves the first part of the proposition.

The second part follows from the first provided we can show $M_{\tilde{\lambda}}$ is irreducible whenever $\lambda_{\kappa_0,1} - \lambda_{\kappa_0,d} \leq p - 1$. To establish this irreducibility we require $d = 2$. Choose an indexing $\kappa_{0,1}, \dots, \kappa_{0,e}$ of those κ with $\kappa|_k = \kappa_0$ so that $\kappa_{0,1} = \tilde{\kappa}_0$. Then construct a scheme X which classifies tuples $(\mathcal{E}_e \subset \dots \subset \mathcal{E}_1)$ with $\mathcal{E}_1 \in \prod_{\kappa_0} (G/P_{\kappa_{0,1}} \otimes_{\mathcal{O}_{K,\kappa_{0,1}}} \mathcal{O})$ and

$$\left(\prod_{\kappa_0} E_{\kappa_{0,i}}(u) \right) \mathcal{E}_i \subset \mathcal{E}_{i+1} \subset \mathcal{E}_i$$

with $\mathcal{E}_i/\mathcal{E}_{i+1}$ of rank one over $(A \otimes_{\mathbb{Z}_p} W(k))$ for each i . Then the map $(\mathcal{E}_i) \mapsto \mathcal{E}_e$ produces a proper morphism $X \rightarrow \text{Gr}_{\mathcal{O}}$ which on the generic fibre identifies $X \otimes_{\mathcal{O}} E$ with $M_{\tilde{\lambda}} \otimes_{\mathcal{O}} E$. In particular, this shows that $X \otimes_{\mathcal{O}} \mathbb{F} \rightarrow M_{\tilde{\lambda}} \otimes_{\mathcal{O}} \mathbb{F}$ is surjective. On the other hand, X is a successive extension of (products of) grassmannians over a (product of) flag varieties. Thus X is \mathcal{O} -smooth, and so $X \otimes_{\mathcal{O}} \mathbb{F}$ is irreducible. We conclude the same is true of $M_{\tilde{\lambda}} \otimes_{\mathcal{O}} \mathbb{F}$.

Finally, note that the $M_{\tilde{\lambda}} \otimes_{\mathcal{O}} \mathbb{F}$ are pairwise distinct because if $M_{\tilde{\lambda}} \otimes_{\mathcal{O}} \mathbb{F} = M_{\tilde{\lambda}'} \otimes_{\mathcal{O}} \mathbb{F}$ then $\mathcal{C}_{\lambda'_{\kappa_0}+e\rho} = \mathcal{C}_{\lambda_{\kappa_0}+e\rho}$ for each κ_0 . But this implies $\mathcal{E}_{\lambda_{\kappa_0}+e\rho} \in \overline{\text{Gr}}_{\leq \lambda'_{\kappa_0}+e\rho}$ and so $\lambda_{\kappa_0} \leq \lambda'_{\kappa_0}$. By symmetry $\lambda_{\kappa_0} = \lambda'_{\kappa_0}$. \square

8 Duality

In this section we introduce an involution of $\text{Gr}_{\mathcal{O}}$ which is useful when dealing with certain normalisation issues which arise when passing between the affine grassmannian and moduli of Breuil–Kisin modules.

8.1 If \mathcal{E} corresponds to an A -valued point of $\text{Gr}_{\mathcal{O}}$ then

$$\mathcal{E}^* := \text{Hom}_{(A \otimes_{\mathbb{Z}_p} W(k))[u]}(\mathcal{E}, (A \otimes_{\mathbb{Z}_p} W(k))[u])$$

is again $(A \otimes_{\mathbb{Z}_p} W(k))[u]$ -projective and, under the natural identification $(A \otimes_{\mathbb{Z}_p} W(k))[u]^{d,*} \cong (A \otimes_{\mathbb{Z}_p} W(k))[u]^d$, we can view \mathcal{E}^* as an A -valued point of $\text{Gr}_{\mathcal{O}}$. Since $\mathcal{E}^{**} = \mathcal{E}$ the endomorphism of $\text{Gr}_{\mathcal{O}}$ induced by

$$\mathcal{E} \mapsto \mathcal{E}^*$$

is an automorphism.

Lemma 8.2 *The above automorphism identifies M_{λ} with $M_{-w_0\lambda}$ where $w_0 \in W$ denotes the longest element.*

In other words, $-w_0\mu = (-w_0\mu_{\kappa})$ where $-w_0\mu_{\kappa} = (-\mu_{\kappa,d}, \dots, -\mu_{\kappa,1}) \in X(T)$.

Proof It suffices to prove this on the generic fibre. Thus, one is reduced to prove that for any $\lambda \in X(T)$, $G/P_\lambda \subset \text{Gr}$ is identified with $G/P_{-w_0\lambda}$ by the version of $\mathcal{E} \mapsto \mathcal{E}^*$ on Gr . But this follows easily from the fact that if \mathcal{E} is generated by $(e_1, \dots, e_d)X$ then \mathcal{E}^* is generated by $(e_1, \dots, e_d)(X^{-1})^t$ for $(X^{-1})^t$ the conjugate transpose. In particular, the G -orbit of any \mathcal{E} is mapped onto the G -orbit of \mathcal{E}^* . Since $\mathcal{E}_\lambda^* = w_0\mathcal{E}_{-w_0\lambda}$ the lemma follows. \square

9 Breuil–Kisin modules

9.1 Let A be a p -adically complete \mathcal{O} -algebra. Then a *Breuil–Kisin module* \mathfrak{M} over A is a finite projective \mathfrak{S}_A -module equipped with an \mathfrak{S}_A -linear homomorphism

$$\varphi_{\mathfrak{M}} = \varphi : \mathfrak{M} \otimes_{\varphi, \mathfrak{S}_A} \mathfrak{S}_A \rightarrow \mathfrak{M}$$

whose cokernel is killed by a power of $E(u)$. We say \mathfrak{M} has height $\leq h$ if the cokernel is killed by $E(u)^h$. We write \mathfrak{M}^φ for the image of $\varphi_{\mathfrak{M}}$ and $\varphi(\mathfrak{M})$ for the image of the composite $\mathfrak{M} \xrightarrow{m \mapsto m \otimes 1} \mathfrak{M} \otimes_{\varphi, \mathfrak{S}_A} \mathfrak{S}_A \rightarrow \mathfrak{S}$. Thus $\varphi(\mathfrak{M})$ is an $\varphi(\mathfrak{S}_A)$ -submodule of \mathfrak{M}^φ which generates \mathfrak{M}^φ over \mathfrak{S}_A .

Definition 9.2 For any p -adically complete \mathcal{O} -algebra A write

- $Z_d^{\leq h}(A)$ for the groupoid of rank d Breuil–Kisin modules over A with height $\leq h$. Morphisms are \mathfrak{S}_A -linear isomorphisms compatible with the Frobenius.
- $\tilde{Z}_d^{\leq h}(A)$ for the groupoid of pairs (\mathfrak{M}, β) where $\mathfrak{M} \in Z_d^{\leq h}(A)$ and $\beta = (\beta_1, \dots, \beta_d)$ is an \mathfrak{S}_A -basis of \mathfrak{M} . Morphisms are \mathfrak{S}_A -linear isomorphisms compatible with the Frobenius and identifying the bases.

With pull-backs defined by base-change these categories form fpqc stacks $Z^{\leq h}, \tilde{Z}_d^{\leq h}$ over $\text{Spf } \mathcal{O}$.

9.3 We consider the following diagram:

$$\begin{array}{ccc} & \tilde{Z}_d^{\leq h} & \\ & \swarrow \Gamma & \searrow \Psi \\ Z_d^{\leq h} & & \text{Gr}_{\mathcal{O}} \end{array} \tag{9.4}$$

where Γ forgets the choice of basis and, with $\text{Gr}_{\mathcal{O}}$ viewed as a formal scheme over $\text{Spf } \mathcal{O}$, the morphism Ψ sends (\mathfrak{M}, β) onto $\mathcal{E} \in \text{Gr}_{\mathcal{O}}$ obtained via

$$\mathcal{E} := \mathfrak{M} \hookrightarrow \frac{1}{E(u)^h} \mathfrak{M}^\varphi \xrightarrow{\varphi(\beta)} \frac{1}{E(u)^h} \mathfrak{S}_A^d$$

(here $\varphi(\beta)$ is interpreted as an identification $\mathfrak{M}^\varphi \cong \mathfrak{S}_A^d$). Concretely, if C is the matrix with entries in \mathfrak{S}_A for which $\varphi(\beta) = \beta C$ then, since \mathfrak{M} is generated by $\varphi(\beta)C^{-1}$, we have $\Psi(\mathfrak{M}, \beta) = \mathcal{E}$ for $\mathcal{E} \subset \mathfrak{S}_A[\frac{1}{E(u)}]^d$ the submodule generated by C^{-1} .

Writing $L^+ \text{GL}_d$ for the group scheme given by $A \mapsto \text{GL}_d(\mathfrak{S}_A)$ we see that Γ is an $L^+ \text{GL}_d$ -torsor for the action on $\tilde{Z}_d^{\leq h}$ given by $g \cdot (\mathfrak{M}, \beta) = (\mathfrak{M}, \beta g)$. A second action of $L^+ \text{GL}_d$ on $\tilde{Z}_d^{\leq h}$ can be given by


$$g * (\mathfrak{M}, \beta) = (\mathfrak{M}_g, \beta)$$

where $\mathfrak{M}_g = \mathfrak{M}$ as an \mathfrak{S}_A -module, with Frobenius given by $\varphi_g(\beta) = \beta g C$ for C the matrix determined by $\varphi(\beta) = \beta C$. It is easy to see that this action makes Ψ into an $L^+ \text{GL}_d$ -torsor over its image in $\text{Gr}_{\mathcal{O}}$.

9.5 An alternative viewpoint on 9.3 is as follows. Let $L^{\leq h} \text{GL}_d$ denote the group scheme over \mathcal{O} given by

$$A \mapsto \{C \in \text{GL}_d(\mathfrak{S}_A[\frac{1}{E(u)}]) \mid E(u)^h C^{-1}, C \in \text{Mat}(\mathfrak{S}_A)\}$$

Then the morphism $\tilde{Z}_d^{\leq h} \rightarrow L^{\leq h} \text{GL}_d$ given by $(\mathfrak{M}, \beta) \mapsto C$, for C defined by $\varphi(\beta) = \beta C$, is an isomorphism (here we view $L^{\leq h} \text{GL}_d$ as a formal scheme over $\text{Spf } \mathcal{O}$). Under this isomorphism the action $g \cdot (\mathfrak{M}, \beta)$ corresponds to the (right) action of $L^+ \text{GL}_d$ by φ -conjugation: $C \mapsto g^{-1} C \varphi(g)$. The action $g * (\mathfrak{M}, \beta)$ corresponds to the (left) multiplication action: $C \mapsto g C$. Thus (9.4) identifies with the diagram



Here $L^{\leq h} \text{GL}_d /_{\varphi} L^+ \text{GL}_d$ indicates the quotient by Frobenius conjugation

The issue with the construction in 9.3 is that $\tilde{Z}_d^{\leq h} \cong L^{\leq h} \text{GL}_d$ is not of finite type over $\text{Spf } \mathcal{O}$. To address this we instead consider:


Definition 9.6 For $N \geq 1$ let $\tilde{Z}_d^{\leq h, N}(A)$ denote the category of pairs $(\mathfrak{M}, \beta) \in \tilde{Z}_d^{\leq h}(A)$ whose morphisms are φ -equivariant morphisms of \mathfrak{S}_A -modules which identify the bases modulo u^N . Equivalently, $\tilde{Z}_d^{\leq h, N}$ is the quotient of $\tilde{Z}_d^{\leq h}$ under the action of the group scheme

$$U_N : A \mapsto 1 + u^N \text{Mat}(\mathfrak{S}_A)$$

Thus $\tilde{Z}_d^{\leq h} \cong [L^{\leq h} \text{GL}_d / U_N]$ (the quotient being by Frobenius conjugation).

The following is a essentially [24, 2.1].

Proposition 9.7 Fix $n \geq 1$. Then, for N sufficiently large, (9.4) induces a diagram



in which Γ_N is a \mathcal{G}_N -torsor for \mathcal{G}_N the group scheme defined by $A \mapsto \mathrm{GL}_d(\mathfrak{S}_A/u^N)$ and Ψ_N is smooth of relative dimension $\dim_{\mathcal{O}} \mathcal{G}_N$ with irreducible fibres. More precisely, the action $g * (\mathfrak{M}, \beta)$ from 9.3 induces an action of \mathcal{G}_N on $\tilde{Z}_d^{\leq h, N} \otimes_{\mathcal{O}} \mathcal{O}/\pi^n$ making Ψ_N into a \mathcal{G}_N -torsor over its image in $\mathrm{Gr}_{\mathcal{O}} \otimes_{\mathcal{O}} \mathcal{O}/\pi^n$.

We will see in the proof that N is required large enough that $E(u)^h$ divides $u^{(p-1)N-1}$ in $\mathfrak{S}_{\mathcal{O}/\pi^n}$ (this can be made explicit using e.g. [6, 5.2.6]).

Proof The crucial observation is that if $E(u)^h$ divides $u^{(p-1)N-1}$ in $\mathfrak{S}_{\mathcal{O}/\pi^n}$ then, as stacks over $\mathrm{Spec} \mathcal{O}/\pi^n$, the identity map $L^{\leq h} \mathrm{GL}_d \rightarrow L^{\leq h} \mathrm{GL}_d$ induces an isomorphism

$$\left[L^{\leq h} \mathrm{GL}_d /_{\varphi} U_N \right] \cong [U_N \setminus L^{\leq h} \mathrm{GL}_d]$$

(the quotient on the left being by Frobenius conjugation and that on the right being by left multiplication). This immediately shows that the morphism Ψ induces a morphism Ψ_N as claimed, and that Ψ_N is a $\mathcal{G}_N = L^+ \mathrm{GL}_d / U_N$ -torsor over the closed \mathcal{O}/π^n -subscheme $[L^{\leq h} \mathrm{GL}_d / L^+ \mathrm{GL}_d] \hookrightarrow \mathrm{Gr}_{\mathcal{O}}$.

Concretely, the claimed isomorphism follows from the two assertions: If $C \in L^{\leq h} \mathrm{GL}_d(A)$ then

- For each $g_0 \in U_N(A)$ we have $g_0^{-1} C \varphi(g_0) = gC$ for a unique $g \in U_N(A)$.
- For each $g \in U_N(A)$ there exists a unique $g_0 \in U_N(A)$ with $g_0^{-1} C \varphi(g_0) = gC$.

Note these are exactly the same assertions as (1) and (2) in [24, 2.2]. The first point is easy because $g_0 \in U_N(A)$ implies

$$g = g_0^{-1} C \varphi(g_0) C^{-1} = g_0^{-1} \left(1 + u^{pN} C_1 C^{-1} \right) = 1 + u^N C_2 + u^{pN} C_3 C^{-1}$$

Therefore, the fact that $E(u)^h C^{-1} \in \mathrm{Mat}(\mathfrak{S}_A)$ and that $E(u)^h$ divides u^{pN} ensures $g \in U_N(A)$. For the second point notice that, if $J_n = (gC)\varphi(gC) \dots \varphi^n(gC)$ and $I_n = C\varphi(C) \dots \varphi^{n-1}(C)$, then any such g_0 satisfies

$$g_0 = I_n \varphi^n(g_0) J_n^{-1} = I_n J_n^{-1} + I_n (\varphi^n(g_0) - 1) J_n^{-1}$$

for every $n \geq 1$. We claim that for any $g_1 \in U_N(A)$ we have $I_n (\varphi^n(g_1) - 1) J_n^{-1} \in u^N \mathrm{Mat}(\mathfrak{S}_A)$ and that this element converges u -adically to zero as $n \rightarrow \infty$. This ensures that $I_n J_n^{-1}$ converges in $U_N(A)$ as $n \rightarrow \infty$ and that this limit is the unique g_0 satisfying $g_0^{-1} C \varphi(g_0) = gC$. Since $g_1 \equiv 1$ modulo u^N we can write $\varphi^n(g_1) - 1 = u^{N+(p^n-1)N} g'_1$. Therefore the claim will follow if $u^{(p^n-1)N} J_{n-1}^{-1} \in \mathrm{Mat}(\mathfrak{S}_A)$ converges u -adically to zero. Since $\varphi^i(E(u)^h)$ divides $u^{((p-1)N-1)p^i}$ and

$$E(u)^h \varphi(E(u)^h) \dots \varphi^{n-1}(E(u)^h) I_{n-1}^{-1} \in \mathrm{Mat}(\mathfrak{S}_A)$$

the claim follows from the observation that $(p^n - 1)N - ((p - 1)N - 1)(1 + \dots + p^{n-1}) = 1 + \dots + p^{n-1}$ is ≥ 0 and $\rightarrow \infty$. □

Corollary 9.8 $\tilde{Z}_d^{\leq h, N} \times_{\mathcal{O}} \mathcal{O}/\pi^n$ is a finite type \mathcal{O} -scheme for $N \gg 0$ and $Z_d^{\leq h}$ is a p -adic formal algebraic stack (in the sense of [7, A7]) of finite type over $\mathrm{Spf} \mathcal{O}$.

Proof The first part follows since we’ve just seen that $\tilde{Z}_d^{\leq h, N} \times_{\mathcal{O}} \mathcal{O}/\pi^n$ is a torsor for a finite type group scheme over a finite type \mathcal{O}/π^n -scheme. The second part follows from the first and the definition of a p -adic formal algebraic stack. \square

10 Crystalline Breuil–Kisin modules

10.1 If A is a p -adically complete \mathcal{O} -algebra which is of topologically finite type then a *crystalline Breuil–Kisin module over A* is a pair (\mathfrak{M}, σ) with \mathfrak{M} a Breuil–Kisin module over A and σ a continuous φ -equivariant $A_{\mathrm{inf}, A}$ -semilinear action of G_K on $\mathfrak{M} \otimes_{\mathfrak{S}_A} A_{\mathrm{inf}, A}$ satisfying

$$(\sigma - 1)(m) \in \mathfrak{M} \otimes_{\mathfrak{S}_A} [\pi^b] \varphi^{-1}(\mu) A_{\mathrm{inf}, A}, \quad (\sigma_{\infty} - 1)(m) = 0$$

for every $m \in \mathfrak{M}$ and every $\sigma \in G_K, \sigma_{\infty} \in G_{K_{\infty}}$.

Definition 10.2 Write $Y_d^{\leq h}(A)$ for the groupoid consisting of rank d crystalline Breuil–Kisin modules over A with height $\leq h$.

10.3 One can attach a Hodge type to crystalline Breuil–Kisin modules in $Y_d^{\leq h}(A)$ whenever A is \mathcal{O} -flat. For $n \in \mathbb{Z}$ define $\mathrm{Fil}^n(\mathfrak{M}^{\varphi}) = \mathfrak{M}^{\varphi} \cap E(u)^n \mathfrak{M}$ and equip the finite projective $\mathcal{O}_K \otimes_{\mathbb{Z}_p} A$ -module $\mathfrak{M}^{\varphi}/E(u)$ with the filtration whose n -th filtered piece is the image of $\mathrm{Fil}^n(\mathfrak{M}^{\varphi})$. It is shown in [7, 4.7.2] that the graded pieces of this filtration become $(A \otimes_{\mathbb{Z}_p} W(k)[\frac{1}{p}])$ -projective after inverting p . This allows us to say that (\mathfrak{M}, σ) has Hodge type $\mu = (\mu_{\kappa})$ if the part of $\mathrm{gr}^n \mathfrak{M}^{\varphi}[\frac{1}{p}]/E(u)$ on which K acts via κ has E -dimension equal the multiplicity of n in μ_{κ} .

Remark 10.4 In other words, \mathfrak{M} has Hodge type μ if the part of $\mathfrak{M}^{\varphi}[\frac{1}{p}]/E(u)$ on which K acts via κ is a filtration of type

$$-w_0 \mu_{\kappa} = (-\mu_{\kappa, d}, \dots, -\mu_{\kappa, 1})$$

in the sense of 4.4.

Theorem 10.5 If $(\mathfrak{M}, \sigma) \in Y_d^{\leq h}(A)$ with A a finite flat \mathcal{O} -algebra then

$$V = (\mathfrak{M} \otimes_{\mathfrak{S}_A} W(C^b)_A)^{\varphi=1}$$

equipped with the G_K -action induced by σ is a crystalline representation of G_K on a finite projective A -module. Furthermore, the Hodge type of (\mathfrak{M}, σ) coincides with that attached to V via the filtered module $D_{\mathrm{crys}}(V)_K := (V \otimes_{\mathbb{Z}_p} B_{\mathrm{dR}})^{G_K}$ with n -th filtered piece given by $(V \otimes_{\mathbb{Z}_p} t^n B_{\mathrm{dR}}^+)^{G_K}$.

Proof The theorem as stated is taken from [2, 2.1.12], but the result originates from a combination of ideas appearing in [11, 16, 22]. \square

Remark 10.6 These conventions mean that the Hodge type of the cyclotomic character is -1 .

Proposition 10.7 *There exists a limit preserving p -adic algebraic formal stack $Y_d^{\leq h}$ of topologically finite type over \mathcal{O} whose groupoid of A -valued points, for any p -adically complete \mathcal{O} -algebra topologically of finite type, is canonically equivalent to $Y_d^{\leq h}(A)$.*

For each Hodge type μ there exists a unique \mathcal{O} -flat closed substack Y_d^μ of $Y_d^{\leq h}$ with the property that the full subcategory $Y_d^\mu(A)$ of $Y_d^{\leq h}(A)$ consists of all crystalline Breuil–Kisin modules with height $\leq h$ and Hodge type μ whenever A is finite flat \mathcal{O} -algebra.

Proof The first part follows from [7, §4.5]. There, algebraic stacks $\mathcal{C}_{\pi^b, s, d, h}^a$ over $\text{Spec } \mathcal{O}/\pi^a$ are constructed [7, 4.5.8] with $\pi^b = (\pi, \pi^{1/p}, \dots)$ and s some sufficiently large integer. In the proof of [7, 4.5.15] it is explained how $Y_\mu^{\leq h} \times_{\mathcal{O}} \mathcal{O}/\pi^a$ can be realised as a closed substack of $\mathcal{C}_{\pi^b, s, d, h}^a$. The second part is [7, 4.8.2]. \square

We conclude with a useful lemma giving a description of the points of Y_d^μ valued in a finite local \mathbb{F} -algebra. For this we make the following construction:

Construction 10.8 Let \mathbb{F}'/\mathbb{F} be a finite extension. Write $R_{\bar{\rho}}$ for the framed \mathcal{O} -deformation ring corresponding to some $\bar{\rho} : G_K \rightarrow \text{GL}_d(\mathbb{F}')$. In [2, 2.2.11] a projective $R_{\bar{\rho}}$ -scheme $\mathcal{L}_{\bar{\rho}}^{\leq h}$ is constructed with A' -points, for A' any p -adically complete \mathcal{O} -algebra, classifying pairs (\mathfrak{M}, ρ) with ρ a framed deformation of $\bar{\rho}$ to A' and $\mathfrak{M} \in Z_{\bar{\rho}}^{\leq h}(A')$ satisfying

$$\mathfrak{M} \otimes_{\mathfrak{S}_A} W(C^b)_A = \rho \otimes_A W(C^b)_A$$

so that φ (induced semilinearly from that on \mathfrak{M}) is the identity on $\bar{\rho}$ and so that the G_K -action (induced semilinearly from that on ρ) satisfies

$$(\sigma - 1)(m) \in \mathfrak{M} \otimes_{\mathfrak{S}_A} [\pi^b] \varphi^{-1}(\mu) A_{\text{inf}, A}, \quad (\sigma_\infty - 1)(m) = 0$$

for every $\sigma \in G_K, \sigma_\infty \in G_{K_\infty}$ and $m \in \mathfrak{M}$. In [2, 2.2.15] it is also shown that the morphism $\mathcal{L}_{\bar{\rho}}^{\leq h} \rightarrow \text{Spec } R_{\bar{\rho}}^\square$ becomes a closed immersion after inverting p , and, if $\mu_\kappa \subset [0, h]$ for all κ , that the closed subscheme $\mathcal{L}_{\bar{\rho}}^{\leq h}[\frac{1}{p}] \hookrightarrow \text{Spec } R_{\bar{\rho}}^\square[\frac{1}{p}]$ contains $\text{Spec } R_{\bar{\rho}}^\mu[\frac{1}{p}]$, where $R_{\bar{\rho}}^\mu$ is the reduced \mathcal{O} -flat quotient of $R_{\bar{\rho}}$ constructed in [17, 3.3.8].

If $\widehat{\mathcal{L}}_{\bar{\rho}}^{\leq h}$ denotes the $\mathfrak{m}_{R_{\bar{\rho}}}$ -adic completion of $\mathcal{L}_{\bar{\rho}}^{\leq h}$ then we see there is a morphism

$$\widehat{\mathcal{L}}_{\bar{\rho}}^{\leq h} \rightarrow Y_d^{\leq h}$$

given by $(\mathfrak{M}, \rho) \mapsto (\mathfrak{M}, \sigma)$. Notice $\widehat{\mathcal{L}}_{\bar{\rho}}^{\leq h}$ can equivalently be constructed as the fibre product $Y_d^{\leq h} \times_{\mathcal{X}_{K, d}} \text{Spf } R_{\bar{\rho}}$ where $\text{Spf } R_{\bar{\rho}} \rightarrow \mathcal{X}_{K, d}$ is the morphism defined in [7, 3.6.3].

Lemma 10.9 *Assume that $\mu_\kappa \subset [0, h]$ for each κ and suppose that $(\overline{\mathfrak{M}}, \overline{\sigma})$ corresponds to an \overline{A} -valued point of Y_d^μ for \overline{A} some finite local \mathbb{F} -algebra. Then there exists a local finite flat \mathcal{O} -algebra A with $\overline{A} = A \otimes_{\mathcal{O}} \mathbb{F}$ and an A -valued point (\mathfrak{M}, σ) of Y_d^μ with special fibre $(\overline{\mathfrak{M}}, \overline{\sigma})$.*

Proof To begin let $R_{\overline{\rho}}$ be a framed deformation ring as in Construction 10.8 and set $\widehat{\mathcal{L}}_{\overline{\rho}}^\mu = Y_d^\mu \times_{\mathcal{X}_{K,d}} \text{Spf } R_{\overline{\rho}}$. Since $\mu_\kappa \subset [0, h]$ for each κ we can view $\widehat{\mathcal{L}}_{\overline{\rho}}^\mu$ as the preimage of Y_d^μ under the morphism $\widehat{\mathcal{L}}_{\overline{\rho}}^{\leq h} \rightarrow Y_d^{\leq h}$ from Construction 10.8. Notice this is a special case of the formal scheme denoted \widehat{C} in the proof of [7, 4.8.10]. In loc. cit it is shown that the scheme theoretic image of the morphism $\widehat{\mathcal{L}}_{\overline{\rho}}^\mu \rightarrow \text{Spf } R_{\overline{\rho}}$ is precisely $\text{Spf } R_{\overline{\rho}}^\mu$ (as in Construction 10.8, $R_{\overline{\rho}}^\mu$ denotes the reduced \mathcal{O} -flat quotient of $R_{\overline{\rho}}$ classifying crystalline representations of Hodge type μ). Using [7, 4.5.26] we see that $\widehat{\mathcal{L}}_{\overline{\rho}}^\mu$ arises as the $m_{R_{\overline{\rho}}}$ -adic completion of a projective $R_{\overline{\rho}}$ -scheme $\mathcal{L}_{\overline{\rho}}^\mu$. This scheme embeds as a closed subscheme in $\mathcal{L}_{\overline{\rho}}^{\leq h}$ and, since $\mathcal{L}_{\overline{\rho}}^{\leq h}[\frac{1}{p}] \rightarrow \text{Spec } R_{\overline{\rho}}$ is a closed immersion, we have $\mathcal{L}_{\overline{\rho}}^\mu[\frac{1}{p}] = \text{Spec } R_{\overline{\rho}}^\mu[\frac{1}{p}]$.

By [7, A.31] we know that $\text{Spf } R_{\overline{\rho}} \rightarrow \mathcal{X}_{K,d}^\mu$ is flat, and so the same is true of $\widehat{\mathcal{L}}_{\overline{\rho}}^\mu \rightarrow Y_d^\mu$. Since $Y_d^\mu \rightarrow \text{Spf } \mathcal{O}$ is flat we conclude that $\widehat{\mathcal{L}}_{\overline{\rho}}^\mu$ is \mathcal{O} -flat also. Therefore, reducedness of $R_{\overline{\rho}}^\mu$ implies $\mathcal{L}_{\overline{\rho}}^\mu$ is reduced too.

Now apply the above construction with $\overline{\rho} = (\overline{\mathfrak{M}} \otimes_{\mathfrak{S}_{\overline{A}}} W(C^b)_{\overline{A}})^{\varphi=1} \otimes_{\overline{A}} \mathbb{F}'$ where \mathbb{F}' denotes the residue field of \overline{A} . Then $(\overline{\mathfrak{M}}, \overline{\sigma})$ form the lemma induces an \overline{A} -valued point of $\mathcal{L}_{\overline{\rho}}^\mu$. Since $\mathcal{L}_{\overline{\rho}}^\mu$ is reduced, \mathcal{O} -flat, and of finite type over $R_{\overline{\rho}}^\mu$ we can apply [2, 4.1.2] to the local ring of $\mathcal{L}_{\overline{\rho}}^\mu$ at this points to produce a finite flat \mathcal{O} -algebra A with $A \otimes_{\mathcal{O}} \mathbb{F} = \overline{A}$ and an A -valued point (\mathfrak{M}, ρ) of $\mathcal{L}_{\overline{\rho}}^\mu$ pulling back to our \overline{A} -valued point. Write σ for the G_K -action on $\mathfrak{M} \otimes_{\mathfrak{S}_A} A_{\text{inf},A}$ induced by ρ . Then (\mathfrak{M}, σ) is a crystalline Breuil–Kisin module with Hodge type μ and so we obtain an A -valued point in $Y_d^\mu(A)$ as desired. \square

11 Naive Galois actions

In this section we consider the morphism $Y_d^{\leq h} \rightarrow Z_d^{\leq h}$ which forgets the Galois action. More precisely, we consider its base-change $Y_d^{\leq h} \times_{Z_d^{\leq h}} \widetilde{Z}_d^{\leq h,N} \rightarrow \widetilde{Z}_d^{\leq h,N}$ for $N \gg 0$, and show this is an isomorphism over certain closed subschemes in the special fibre of $\widetilde{Z}_d^{\leq h,N}$.

Construction 11.1 The aim is to establish conditions which allow the following “naive” crystalline G_K -action on $(\mathfrak{M}, \beta) \in \widetilde{Z}_d^{\leq h,N}(A)$ to be perturbed into one which is φ -equivariant. Let $\sigma_{\text{naive},\beta}$ denote the continuous $A_{\text{inf},A}$ -semilinear action of G_K on $\mathfrak{M} \otimes_{\mathfrak{S}_A} A_{\text{inf},A}$ obtained from the coordinate-wise action on $A_{\text{inf},A}^d$ via the identification

$$\mathfrak{M}^\varphi \otimes_{\mathfrak{S}_A} A_{\text{inf},A} \cong A_{\text{inf},A}^d$$

induced by $\varphi(\beta)$. Thus, $\sigma_{\text{naive},\beta}$ is uniquely determined as the semilinear G_K -action fixing $\varphi(\beta)$.

11.2 Let us fix integers $0 \leq r_\kappa \leq h$ for each κ . Then we consider the closed subfunctor $\text{Gr}_{\mathcal{O}}^{\nabla_{\sigma},r} \subset \text{Gr}_{\mathcal{O}}^{\nabla_{\sigma}}$ defined by requiring that

$$\mathfrak{S}_A^d \subset \mathcal{E} \subset \prod_{\kappa} E_{\kappa}(u)^{-r_{\kappa}} \mathfrak{S}_A^d$$

Recall $\text{Gr}_{\mathcal{O}}^{\nabla_{\sigma}}$ was defined in Lemma 7.4. We then choose N sufficiently large so that Proposition 9.7 produces the morphism $\Psi_N : \tilde{Z}_d^{\leq h,N} \otimes_{\mathcal{O}} \mathbb{F} \rightarrow \text{Gr}_{\mathcal{O}} \otimes_{\mathcal{O}} \mathbb{F}$ and define

$$\tilde{Z}_{d,\mathbb{F}}^{\nabla_{\sigma},r} \hookrightarrow \tilde{Z}_d^{\leq h,N} \otimes_{\mathcal{O}} \mathbb{F}$$

as the preimage of $\text{Gr}_{\mathcal{O}}^{\nabla_{\sigma},r} \otimes_{\mathcal{O}} \mathbb{F}$ under the morphism Ψ_N . Notice we suppress the dependence of $\tilde{Z}_{d,\mathbb{F}}^{\nabla_{\sigma},r}$ on N from the notation. The A -points of $\tilde{Z}_d^{\nabla_{\sigma},r}$ for A any finite type \mathbb{F} -algebra are precisely those $(\mathfrak{M}, \beta) \in \tilde{Z}_{d,\mathbb{F}}^{\leq h,N}(A)$ satisfying

(1) For every $\sigma \in G_K$

$$(\sigma_{\text{naive},\beta} - 1)(\mathfrak{M}) \subset \mathfrak{M} \otimes_{\mathfrak{S}_A} [\pi^b] \varphi^{-1}(\mu) A_{\text{inf},A,i}$$

See Remark 7.5.

(2) For each $i = 1, \dots, f$

$$\prod_j E_{\kappa}(u)^{r_{\kappa}} \mathfrak{M}_i \subset \mathfrak{M}_i^{\varphi} \subset \mathfrak{M}_i$$

Proposition 11.3 Assume that $\sum_{\kappa|k=\kappa_0} r_{\kappa} \leq e + p - 1$ for each κ_0 with at least one inequality strict. Then

$$Y_d^{\leq h} \times_{Z_d^{\leq h}} \tilde{Z}_{d,\mathbb{F}}^{\nabla_{\sigma},r} \rightarrow \tilde{Z}_{d,\mathbb{F}}^{\nabla_{\sigma},r}$$

is an isomorphism.

Proof By Lemma 15.1 is enough to show that this morphism induces equivalences on groupoids of \bar{A} -valued points for any local finite type \mathbb{F} -algebra \bar{A} . Equivalently, we must show that for any $(\mathfrak{M}, \beta) \in \tilde{Z}_{d,\mathbb{F}}^{\nabla_{\sigma},r}(\bar{A})$ there exists a unique action σ of G_K making (\mathfrak{M}, σ) into an object of $Y_d^{\leq h}$. Existence implies essential surjectivity on \bar{A} -valued points and full-faithfulness follows from the uniqueness.

Write $\text{Hom}(\mathfrak{M}, \mathfrak{M})$ for the \mathfrak{S}_A -module of \mathfrak{S}_A -linear endomorphisms of \mathfrak{M} and equip $\text{Hom}(\mathfrak{M}, \mathfrak{M})$ with the Frobenius φ_{Hom} given by $h \mapsto \varphi \circ h \circ \varphi^{-1}$. If we identify $\text{Hom}(\mathfrak{M}, \mathfrak{M})$ with matrices in \mathfrak{S}_A using the basis β then φ_{Hom} acts by

$$M \mapsto C\varphi(M)C^{-1}$$

where φ acts entry wise on the matrix M and C is such that $\varphi(\beta) = \beta C$. The following claim shows that, after extending scalars to $A_{\text{inf},A}$, this operator is topologically nilpotent on matrices with entries divisible by $[\pi^b]\varphi^{-1}(\mu)$.

Claim *Set $\mathcal{H} := \text{Hom}(\mathfrak{M}, \mathfrak{M}) \otimes [\pi^b]\varphi^{-1}(\mu)A_{\text{inf},\bar{A}}$. Then \mathcal{H} is φ_{Hom} -stable and φ_{Hom} is topologically nilpotent on \mathcal{H} .*

Proof of claim Recall that $\mathfrak{M}_{\kappa_0} \subset \mathfrak{M}$ is the submodule on which $W(k)$ acts via κ_0 and φ on \mathfrak{M} restricts to a semilinear map $\mathfrak{M}_{\kappa_0} \rightarrow \mathfrak{M}_{\kappa_0 \circ \varphi^{-1}}[\frac{1}{E(u)}]$. Our assumption on \mathfrak{M} implies $(\prod_{\kappa|k=\kappa_0} E_{\kappa}(u)^{r_{\kappa}})\mathfrak{M}_{\kappa_0} \subset \mathfrak{M}_{\kappa_0}^{\varphi}$ and therefore

$$\varphi_{\text{Hom}}(\text{Hom}(\mathfrak{M}, \mathfrak{M})_{\kappa_0}) \subset \left(\prod_{\kappa|k=\kappa_0} E_{\kappa}(u)^{-r_{\kappa}} \right) \text{Hom}(\mathfrak{M}, \mathfrak{M})_{\kappa_0 \circ \varphi^{-1}}$$

Since $\frac{[\pi^b]^p \mu}{[\pi^b]\varphi^{-1}(\mu)} A_{\text{inf},\bar{A}} = [\pi^b]^{p-1} E(u) A_{\text{inf},\bar{A}}$ we also have

$$\varphi_{\text{Hom}}(\mathcal{H}_{\kappa_0}) \subset [\pi^b] E(u) \left(\prod_{\kappa|k=\kappa_0} E_{\kappa}(u)^{-r_{\kappa}} \right) \mathcal{H}_{\kappa_0 \circ \varphi^{-1}} = u^{e+p-1-\sum_j r_{ij}} \mathcal{H}_{i-1}$$

(the last equality uses that \bar{A} is an \mathbb{F} -algebra) and so, as $\sum_{\kappa|k=\kappa_0} r_{\kappa} \leq e + p - 1$, it follows that \mathcal{H} is φ_{Hom} -stable. Since the inequality is strict at least once we have $\varphi_{\text{Hom}}(\mathcal{H}_{\kappa}) \subset u \mathcal{H}_{\kappa_0 \circ \varphi^{-1}}$ for at least one κ_0 . In particular φ_{Hom} is topologically nilpotent. \square

Set $\mathfrak{M}_{\text{inf}} = \mathfrak{M} \otimes_{\mathfrak{S}_A} A_{\text{inf},A}$. Note that for each $\sigma \in G_K$ the endomorphism $\sigma_{\text{naive},\beta}$ is σ -semilinear and so defines an element of

$$\text{Hom}(\mathfrak{M}_{\text{inf}}, \mathfrak{M}_{\text{inf}}^{\sigma})$$

where $\mathfrak{M}_{\text{inf}}^{\sigma} := \mathfrak{M}_{\text{inf}} \otimes_{A_{\text{inf},A},\sigma} A_{\text{inf},A}$. Any such semilinear map is determined by where β is sent; thus we obtain an (additive) identification $\text{Hom}(\mathfrak{M}_{\text{inf}}, \mathfrak{M}_{\text{inf}}^{\sigma}) = \text{Hom}(\mathfrak{M}, \mathfrak{M}) \otimes_{\mathfrak{S}_A} A_{\text{inf},A}$ which identifies $\text{Hom}(\mathfrak{M}_{\text{inf}}, [\pi^b]\varphi^{-1}(\mu)\mathfrak{M}_{\text{inf}}^{\sigma})$ and \mathcal{H} . Via this identification we view φ_{Hom} as a Frobenius on $\text{Hom}(\mathfrak{M}_{\text{inf}}, \mathfrak{M}_{\text{inf}}^{\sigma})$. We claim that

$$\varphi_{\text{Hom}}^n(\sigma_{\text{naive},\beta}) \in \text{Hom}(\mathfrak{M}_{\text{inf}}, \mathfrak{M}_{\text{inf}}^{\sigma})$$

for all $n \geq 0$ and that this sequence converges to a σ -semilinear endomorphism which we simply write as σ . By construction this endomorphism is φ -equivariant. To see this claim note that by assumption $\sigma_{\text{naive},\beta} - 1^{\sigma} \in \text{Hom}(\mathfrak{M}_{\text{inf}}, [\pi^b]\varphi^{-1}(\mu)\mathfrak{M}_{\text{inf}}^{\sigma})$ where 1^{σ} the σ -semilinear extension of the map $\beta \mapsto \beta$. Since we can write

$$\begin{aligned} & \lim_{n \rightarrow \infty} \varphi_{\text{Hom}}^n(\sigma_{\text{naive},\beta}) \\ &= \sigma_{\text{naive},\beta} + \sum_{n \geq 1} \left(\varphi_{\text{Hom}}^n(\sigma_{\text{naive},\beta} - 1^{\sigma}) - \varphi_{\text{Hom}}^{n-1}(\sigma_{\text{naive},\beta} - 1^{\sigma}) \right) \end{aligned}$$

the claimed convergence follows from the topological nilpotence of the operator φ_{Hom} on $\text{Hom}(\mathfrak{M}_{\text{inf}}, [\pi^b]\varphi^{-1}(\mu)\mathfrak{M}_{\text{inf}}^\sigma) = \mathcal{H}$. This formula also shows that $\sigma(x) - x \in \mathfrak{M} \otimes_{\mathfrak{S}_A} [\pi^b]\varphi^{-1}(\mu)A_{\text{inf}, A}$ for each $x \in \mathfrak{M}$. Since $\sigma_{\text{naive}, \beta}$ defines a G_K -action so to does σ . Similarly, continuity of $\sigma_{\text{naive}, \beta}$ implies continuity of σ .

Finally, to see uniqueness suppose that σ' was another such G_K -action. Then for each $\sigma \in G_K$ one has

$$\sigma - \sigma' \in \text{Hom}(\mathfrak{M}_{\text{inf}}, [\pi^b]\varphi^{-1}(\mu)\mathfrak{M}_{\text{inf}}^\sigma) = \mathcal{H}$$

Since $\sigma - \sigma'$ is φ_{Hom} -fixed the topological nilpotence of φ_{Hom} on \mathcal{H} implies $\sigma - \sigma' = 0$. □

12 Comparison with local models

Set $\bar{Y}_d^\mu = Y_d^\mu \otimes_{\mathcal{O}} \mathbb{F}$ and choose N sufficiently large that Proposition 9.7 produces the morphism $\Psi_N : \tilde{Z}_d^{\leq h, N} \otimes_{\mathcal{O}} \mathbb{F} \rightarrow \text{Gr}_{\mathcal{O}} \otimes_{\mathcal{O}} \mathbb{F}$.

Theorem 12.1 *Assume that $\mu_\kappa \subset [0, r_\kappa]$ for integers $r_\kappa \leq h$ satisfying*

$$\sum_{\kappa|k=\kappa_0} r_\kappa \leq \frac{p-1}{v_{\kappa_0}} + 1, \quad v_{\kappa_0} = \max_{\kappa|k=\kappa_0} \{v_\pi(\pi_\kappa - \pi_{\kappa'})\}$$

for all $\kappa_0 : k \rightarrow \mathbb{F}$. Then the composite

$$\bar{Y}_d^\mu \times_{Z_d^{\leq h}} \tilde{Z}_d^{\leq h, N} \rightarrow \tilde{Z}_d^{\leq h, N} \otimes_{\mathcal{O}} \mathbb{F} \xrightarrow{\Psi_N} \text{Gr}_{\mathcal{O}} \otimes_{\mathcal{O}} \mathbb{F}$$

factors through $\bar{M}_{-w_0\mu}$ (recall this notation from Lemma 8.2)

Remark 12.2 If K is tamely ramified, i.e. if e is not divisible by p , then $\pi_\kappa - \pi_{\kappa'}$ generates the same ideal of \mathcal{O}_K as π whenever $\kappa' \neq \kappa$. Therefore $v_{\kappa_0} = 0$ in this case. To see this consider the π -adic valuation of $\frac{d}{du}\kappa_0(E(u))|_{u=\pi_\kappa} = \prod_{\kappa \neq \kappa', \kappa'|k=\kappa_0} (\pi_\kappa - \pi_{\kappa'})$.

The following proposition is the key technical result which goes into the proof of the theorem. It is a reworking of techniques originally developed in [11, 12].

Proposition 12.3 *Let A be a finite flat \mathcal{O} -algebra and suppose $(\mathfrak{M}, \sigma, \beta) \in Y_d^\mu(A)$. Define*

$$M_\kappa = \mathfrak{M}^\varphi / E_\kappa(u)$$

Use the $\varphi(\mathfrak{S}_A)$ -basis $\varphi(\beta)$ to define a section s of $\varphi(\mathfrak{M}) \rightarrow \mathfrak{M}^\varphi \rightarrow M_\kappa$. Then there exists a filtration $\text{Fil}_\kappa^\bullet$ on M_κ by A -submodules with p -torsionfree graded pieces such that

$$\sum_{n=0}^{r_\kappa} E_\kappa(u)^{r_\kappa-n} \mathfrak{S}_A \text{Fil}_\kappa^n + \mathfrak{M}_{\text{err}, \kappa}^\varphi = \mathfrak{M}^\varphi \cap E_\kappa(u)^{r_\kappa} \mathfrak{M} + \mathfrak{M}_{\text{err}, \kappa}^\varphi$$

when Fil_κ^n is viewed as a submodule of \mathfrak{M}^φ via s and $\mathfrak{M}_{\text{eff},\kappa}^\varphi := \sum_{l=1}^{p-1} \pi^{p-l} E_\kappa(u)^l \mathfrak{M}^\varphi$.

Note that, by construction, the image of the section s generates \mathfrak{M}^φ over \mathfrak{S}_A .

Proof First we define the filtration $\text{Fil}_\kappa^\bullet$. Recall that we equipped $M := \mathfrak{M}^\varphi / E(u)$ with the filtration whose n -th piece is the image of $\mathfrak{M}^\varphi \cap E(u)^n \mathfrak{M}$. Then $D_K := M[\frac{1}{p}]$ is a filtered $A \otimes_{\mathbb{Z}_p} K$ -module and can be written as $\prod_\kappa D_{K,\kappa}$ with each $D_{K,\kappa}$ a filtered $A[\frac{1}{p}]$ -module. The composite $\mathfrak{M}^\varphi \rightarrow D_K \rightarrow D_{K,\kappa}$ is obtained by base-change along the map $\mathfrak{S} \rightarrow A[\frac{1}{p}]$ given by $u \mapsto \pi_\kappa$. Therefore, its kernel is $E_\kappa(u) \mathfrak{M}^\varphi$. This means M_κ can be viewed as a submodule of $D_{K,\kappa}$ and $M_\kappa[\frac{1}{p}] = D_{K,\kappa}$. Define

$$\text{Fil}_\kappa^n = \text{Fil}^n(D_{K,\kappa}) \cap M_\kappa$$

The filtered pieces of $D_{K,\kappa}$ are \mathbb{Q}_p -vector spaces so the graded pieces of $\text{Fil}_\kappa^\bullet$ are p -torsionfree. This also shows that $\text{Fil}_\kappa^n[\frac{1}{p}] = \text{Fil}^n(D_{K,\kappa})$ which proves Corollary 12.4 below.

Next we use:

Claim For $\bar{x} \in \text{Fil}_\kappa^n$ with $n \leq p$ there exists $x_1, \dots, x_{p-1} \in \mathfrak{M}^\varphi$ such that

$$s(\bar{x}) + E_\kappa(u) \pi^{p-1} x_1 + \dots + E_\kappa(u)^{p-1} \pi x_{p-1} \in \mathfrak{M}^\varphi \cap E_\kappa(u)^n \mathfrak{M}$$

Proof of Claim This follows from results in [1, §5]. To apply these first note that in *loc. cit.* the embeddings $K \rightarrow E$ are indexed by integers $1 \leq i \leq f$ and $1 \leq j \leq e$ so that $\kappa_{ij}|_k$ depends only on i . This labelling can be chosen so that κ from the proposition equals κ_{i1} for some i . In [1, 5.2.5] it is shown that for any $x \in \varphi(\mathfrak{M})$ there exist $x_1, \dots, x_{p-1} \in \mathfrak{M}^\varphi$ so that

$$x^{(n)} - x + E_1(u) \pi^{p-1} x_1 + \dots + E_1(u)^{p-1} \pi x_{p-1} \in E_1(u)^p \mathfrak{M}^\varphi \otimes_{\mathfrak{S}} S[\frac{1}{p}]$$

where

- S is the ring defined in [1, §5.1],
- $x^{(i)}$ is defined as in [1, §5.2],
- $E_1(u) = \prod_{i=1}^f E_{i1}(u) \in \mathfrak{S}_{\mathcal{O}}$ for $E_{ij}(u) := E_{\kappa_{ij}}(u)$.

We apply this to $x = s(\bar{x})$. Then the image of x in $D_{K,\kappa}$ is contained in $\text{Fil}^n(D_{K,\kappa})$ and so [1, 5.2.2] implies $x^{(n)}$ is contained in a submodule of $\mathfrak{M}^\varphi \otimes_{\mathfrak{S}} S[\frac{1}{p}]$ denoted $\text{Fil}^{(n,0,\dots,0)}$. In [1, 5.1.3] it is shown that $\text{Fil}^{(n,0,\dots,0)} \cap \mathfrak{M}^\varphi = \mathfrak{M}^\varphi \cap E_1(u)^n \mathfrak{M}$. Therefore

$$s(\bar{x}) + E_1(u) \pi^{p-1} x_1 + \dots + E_1(u)^{p-1} \pi x_{p-1} \in \mathfrak{M}^\varphi \cap E_1(u)^n \mathfrak{M} \subset \mathfrak{M}^\varphi \cap E_\kappa(u)^n \mathfrak{M}$$

(the inclusion following because $E_\kappa(u)$ divides $E_1(u)$). Under the identification $\mathfrak{M}^\varphi = \prod_{\kappa_0} \mathfrak{M}_{\kappa_0}^\varphi$ the κ_0 -th part of $\mathfrak{M}^\varphi \cap E_\kappa(u)^n \mathfrak{M}$ is just $\mathfrak{M}_{\kappa_0}^\varphi$ for $\kappa_0 \neq \kappa|_k$. Therefore, in the above identity we can replace each $E_1(u)$ with $E_\kappa(u)$, and the claim follows. \square

The claim shows that

$$\underbrace{\sum_{n=0}^m E_\kappa(u)^{m-n} \mathfrak{S}_A \text{Fil}_\kappa^n + \mathfrak{M}_{\text{err},\kappa}^\varphi}_{:=Y_m} \subset \mathfrak{M}^\varphi \cap E_\kappa(u)^m \mathfrak{M} + \mathfrak{M}_{\text{err},\kappa}^\varphi$$

for any $0 \leq m \leq r_\kappa$ and we want to prove the opposite inclusion for $0 \leq m \leq r_\kappa$ by induction on m . When $m = 0$ this is clear since both sides equal \mathfrak{M}^φ (recall that the section s was chosen so that $s(M_\kappa)$ generates \mathfrak{M}^φ over \mathfrak{S}_A). For $m > 0$ note that the image of $\mathfrak{M}^\varphi \cap E_\kappa(u)^m \mathfrak{M}$ in M_κ is contained in Fil_κ^m , while Fil_κ^m equals the image of Y_m . The above inclusion therefore shows these images are equal. As a consequence, if $x \in \mathfrak{M}^\varphi \cap E_\kappa(u)^m \mathfrak{M}$ then there exists $x' \in Y_m$ so that

$$\begin{aligned} x - x' &\in E_\kappa(u)\mathfrak{M}^\varphi \cap (\mathfrak{M}^\varphi \cap E_\kappa(u)^m \mathfrak{M} + \mathfrak{M}_{\text{err},\kappa}^\varphi) \\ &= (E_\kappa(u)\mathfrak{M}^\varphi \cap E_\kappa(u)^m \mathfrak{M}) + \mathfrak{M}_{\text{err},\kappa}^\varphi \\ &= E_\kappa(u) (\mathfrak{M}^\varphi \cap E_\kappa(u)^{m-1} \mathfrak{M}) + \mathfrak{M}_{\text{err},\kappa}^\varphi \end{aligned}$$

The second equality uses that $\mathfrak{M}_{\text{err},\kappa}^\varphi \subset E_\kappa(u)\mathfrak{M}^\varphi$. The inductive hypothesis therefore gives that $x - x' \in E_\kappa(u)Y_{m-1} + \mathfrak{M}_{\text{err},\kappa}^\varphi$. Since $E_\kappa(u)Y_{m-1} \subset Y_m$ it follows that $x \in Y_m + \mathfrak{M}_{\text{err},\kappa}^\varphi$ as desired. \square

Corollary 12.4 *The graded pieces of $\text{Fil}_\kappa^\bullet$ become $A[\frac{1}{p}]$ -projective after inverting p and $\text{Fil}_\kappa^\bullet[\frac{1}{p}]$ has type $-w_0\mu_\kappa = (-\mu_{\kappa,d} \geq \dots \geq -\mu_{\kappa,1})$.*

Proof of Theorem 12.1 By Corollary 15.2 it suffices to prove the factorisation on the level of \bar{A} -valued points for \bar{A} any finite local \mathbb{F} -algebra. Let $(\bar{\mathfrak{M}}, \bar{\sigma}, \bar{\beta})$ be such a point. Applying Lemma 10.9 we obtain a local finite flat \mathcal{O} -algebra A with a map $A \rightarrow \bar{A}$ and $(\mathfrak{M}, \sigma) \in Y_d^\mu(A)$ lifting $(\bar{\mathfrak{M}}, \bar{\sigma})$. Additionally, choose an \mathfrak{S}_A -basis β lifting $\bar{\beta}$. We will then be done if we can show that the special fibre of $(\mathfrak{M}, \sigma, \beta)$ is mapped into $\bar{M}_{-w_0\mu}$ by Ψ_N . We can assume that $\bar{A} = A \otimes_{\mathcal{O}} \mathbb{F}$.

Applying Proposition 12.3 for each κ we obtain filtrations $\text{Fil}_\kappa^\bullet$. Define \mathcal{E} by

$$\left(\prod_{\kappa} E_\kappa(u)^{r_\kappa} \right) \mathcal{E} = \bigcap_{\kappa} \left(\sum_{n=0}^{r_\kappa} E_\kappa(u)^{r_\kappa-n} \mathfrak{S}_A \text{Fil}_\kappa^n \right)$$

As in Proposition 12.3 the Fil_κ^n 's are viewed as submodules of \mathfrak{M}^φ using the basis $\varphi(\beta)$. Corollary 12.4 together with part (2) of Lemma 4.9 (taking $n_\kappa = r_\kappa$) shows that $\mathcal{E}[\frac{1}{p}]$ defines an $A[\frac{1}{p}]$ point of $M_{-w_0\mu}$ under the identification $\mathfrak{M}^\varphi = \mathfrak{S}_A^d$ induced by $\varphi(\beta)$. Note, however, that it is not a priori clear \mathcal{E} defines an A -valued point. We will be done if we can show this is the case, and if we can show that $\mathcal{E} \otimes_{\mathcal{O}} \mathbb{F}$ coincides with the image of $(\bar{\mathfrak{M}}, \bar{\sigma}, \bar{\beta})$ under Ψ_N . In other words, if \mathcal{E} defines an A -valued point and if $\mathcal{E} \otimes_{\mathcal{O}} \mathbb{F} = \mathfrak{M} \otimes_{\mathcal{O}} \mathbb{F}$.

We begin with the second assertion. Take $z \in (\prod_{\kappa} E_{\kappa}(u)^{r_{\kappa}}) \mathfrak{M}$. Then $z \in \mathfrak{M}^{\varphi} \cap E_{\kappa}(u)^{r_{\kappa}} \mathfrak{M}$ for each κ and so Proposition 12.3 ensures the existence of $m_{\kappa} \in \mathfrak{M}_{\text{err},\kappa}^{\varphi}$ such that

$$z - m_{\kappa} \in \left(\sum_{n=0}^{r_{\kappa}} E_{\kappa}(u)^{r_{\kappa}-n} \mathfrak{S}_A \text{Fil}_{\kappa}^n \right)$$

We claim there then exists $m \in \pi \mathfrak{M}^{\varphi}$ such that

$$m \equiv m_{\kappa} \pmod{E_{\kappa}(u)^{r_{\kappa}} \mathfrak{M}^{\varphi}}$$

for each κ . Since \mathfrak{M}^{φ} is \mathfrak{S}_A -free this claim follows from Lemma 12.5 below. This is where we use the bound on the r_{κ} . Since

$$E_{\kappa}(u)^{r_{\kappa}} \mathfrak{M}^{\varphi} \subset \left(\sum_{n=0}^{r_{\kappa}} E_{\kappa}(u)^{r_{\kappa}-n} \mathfrak{S}_A \text{Fil}_{\kappa}^n \right)$$

for each κ (due to the filtration Fil_{κ}^n being concentrated in degrees $[0, r_{\kappa}]$ we have $\text{Fil}_{\kappa}^0 = M_{\kappa}$) it follows that $z - m \in (\prod_{\kappa} E_{\kappa}(u)^{r_{\kappa}}) \mathcal{E}$. Since $m \in \pi \mathfrak{M}^{\varphi}$ the image of z in $\mathfrak{M}^{\varphi} \otimes_{\mathcal{O}} \mathbb{F}$ is contained in the image of $(\prod_{\kappa} E_{\kappa}(u)^{r_{\kappa}}) \mathcal{E}$. A symmetrical argument shows also that if $z \in (\prod_{\kappa} E_{\kappa}(u)^{r_{\kappa}}) \mathcal{E}$ then its image in $\mathfrak{M}^{\varphi} \otimes_{\mathcal{O}} \mathbb{F}$ is contained in the image of $(\prod_{\kappa} E_{\kappa}(u)^{r_{\kappa}}) \mathfrak{M}$.

Next we show that \mathcal{E} is \mathfrak{S}_A -projective. This is equivalent to $\mathfrak{M}^{\varphi} / (\prod_{\kappa} E_{\kappa}(u)^{r_{\kappa}}) \mathcal{E}$ being A -projective. From the definitions we see that $\mathfrak{M}^{\varphi} / (\prod_{\kappa} E_{\kappa}(u)^{r_{\kappa}}) \mathcal{E}$ is p -torsionfree. This means that $(\prod_{\kappa} E_{\kappa}(u)^{r_{\kappa}}) \mathcal{E} \otimes_{\mathcal{O}} \mathbb{F}$ equals its image in $\mathfrak{M}^{\varphi} \otimes_{\mathcal{O}} \mathbb{F}$. It also means that, by [26, 00ML], A -projectivity of $\mathfrak{M}^{\varphi} / (\prod_{\kappa} E_{\kappa}(u)^{r_{\kappa}}) \mathcal{E}$ follows from $A \otimes_{\mathcal{O}} \mathbb{F}$ -projectivity of $\mathfrak{M}^{\varphi} \otimes_{\mathcal{O}} \mathbb{F} / (\prod_{\kappa} E_{\kappa}(u)^{r_{\kappa}}) \mathcal{E} \otimes_{\mathcal{O}} \mathbb{F}$. But we saw in the previous paragraph that $(\prod_{\kappa} E_{\kappa}(u)^{r_{\kappa}}) \mathcal{E} \otimes_{\mathcal{O}} \mathbb{F} = (\prod_{\kappa} E_{\kappa}(u)^{r_{\kappa}}) \mathfrak{M} \otimes_{\mathcal{O}} \mathbb{F}$. Since $\mathfrak{M}^{\varphi} / (\prod_{\kappa} E_{\kappa}(u)^{r_{\kappa}}) \mathfrak{M}$ is A -projective the claimed \mathfrak{S}_A -projectivity of \mathcal{E} follows. This establishes the two required conditions mentioned in the second paragraph, and therefore finishes the proof. □

Lemma 12.5 *Let A be a finite flat \mathcal{O} -algebra and suppose*

$$m_{\kappa} = \sum_{l=1}^p E_{\kappa}(u)^{p-l} \pi^l m_{\kappa,l}$$

are given with $m_{\kappa,l} \in A$. Then there exists $m \in \pi \mathfrak{S}_A$ with $m \equiv m_{\kappa}$ modulo $E_{\kappa}(u)^{r_{\kappa}}$ for each κ .

Proof Firstly, by choosing an \mathcal{O} -basis of A we can reduce to the case $A = \mathcal{O}$. Secondly, we can fix κ and assume that $m_{\kappa'} = 0$ for all $\kappa' \neq \kappa$. Using the identification $\mathfrak{S}_{\mathcal{O}} = \prod_{\kappa_0} \mathcal{O}[[u]]$ we are left proving that if $m \in \mathcal{O}[[u]]$ can be written as

$$m = \sum_{l=1}^p (u - \pi_{\kappa})^{p-l} \pi^l m_l, \quad m_l \in \mathcal{O}$$

then there exists $M \in \pi\mathcal{O}[[u]]$ with

- M divisible by $(u - \pi_{\kappa'})^{r_{\kappa'}}$ for every $\kappa' \neq \kappa$ with $\kappa'|_k = \kappa|_k$.
- $M \equiv m$ modulo $(u - \pi_{\kappa})^{r_{\kappa}}$.

We will construct M explicitly. For κ' with $\kappa'|_k = \kappa|_k$ and $\kappa' \neq \kappa$ set

$$X_{\kappa'} := (-1)^{r_{\kappa}} \sum_{n=0}^{r_{\kappa}-1} \binom{r_{\kappa'} - 1 + j}{r_{\kappa'} - 1} \frac{(u - \pi_{\kappa})^n}{(\pi_{\kappa} - \pi_{\kappa'})^{n+r_{\kappa}'}}$$

Using the formal identity $\frac{1}{(1-y)^r} = \sum_{n=0}^{\infty} \binom{r-1+i}{r-1} y^i$ with $r = r_{\kappa'}$ and $y = \frac{u-\pi_{\kappa}}{\pi_{\kappa'}-\pi_{\kappa}}$ shows that

$$X_{\kappa'}(u - \pi_{\kappa'})^{r_{\kappa'}} \equiv 1 \text{ modulo } (u - \pi_{\kappa})^{r_{\kappa}}$$

Define $N \in E[u]$ to be the polynomial of degree $< r_{\kappa}$ when viewed as a polynomial in $(u - \pi_{\kappa})$, obtained by truncating $m \prod_{\kappa' \neq \kappa} X_{\kappa'}$. Then $N \equiv m \prod_{\kappa' \neq \kappa} X_{\kappa'}$ modulo $(u - \pi_{\kappa})^{r_{\kappa}}$ and so

$$M := N \prod_{\kappa' \neq \kappa} (u - \pi_{\kappa'})^{r_{\kappa}'}$$

satisfies the two bullet points above. To finish it suffices to show that N , and hence M also, is contained in $\pi\mathcal{O}[[u]]$.

For this view N as a polynomial in $(u - \pi_{\kappa})$. By assumption the coefficient of $(u - \pi_{\kappa})^n$ in m has valuation $\geq p - n$. On the other hand, the coefficient of $(u - \pi_{\kappa})$ in $X_{\kappa'}$ has valuation $\geq -(n + r_{\kappa}')v$ for $v := v_{\kappa|_k}$. Since $v \geq 1$ we have $p - n \geq p - n v$ and as such the coefficient of $(u - \pi_{\kappa})^n$ in $m \prod_{\kappa' \neq \kappa} X_j$ has valuation

$$\geq p - \left(\sum_{\kappa' \neq \kappa} r_{\kappa'} + n \right) v$$

We will be done if $p - (\sum_{\kappa' \neq \kappa} r_{\kappa'} + n)v \geq 1$ for all $n = 0, \dots, r_{\kappa} - 1$, i.e. if $p - (\sum_{\kappa'|_k = \kappa|_k} r_{\kappa'} - 1)v \geq 1$. This is equivalent to asking that $\sum_{\kappa'|_k = \kappa|_k} r_{\kappa'} \leq \frac{p-1}{v} + 1$ so we are done. \square

13 Lower bounds

In this section we recall from [7, 10] the lower bound on the cycles appearing in the Breuil–Mézard conjecture attained when $d = 2$. We do this in the context of cycles in deformation rings. We also give a minor improvement using the potential diagonalisability established in [1].

13.1 First, recall that isomorphism classes of absolutely irreducible representations of $GL_d(k)$ on \mathbb{F} -vector spaces are in bijection with tuples $\lambda = (\lambda_{\kappa_0})$ indexed by embeddings $\kappa_0 : k \rightarrow \mathbb{F}$ with each λ_{κ_0} a tuple of integers satisfying

$$\lambda_{\kappa_0,1} - \lambda_{\kappa_0,d} \leq p - 1$$

This bijection sends (λ_{κ_0}) onto the $GL_d(k)$ -representation obtained by evaluating the algebraic representation of $G = GL_d$

$$\bigotimes_{\kappa_0} (L(\lambda_{\kappa_0}) \otimes_{k,\kappa_0} \mathbb{F})$$

on k -points [13]. Here $L(\lambda_{\kappa_0}) \subset H^0(G/P_{\lambda_{\kappa_0}}, \mathcal{L}(\lambda_{\kappa_0})) \otimes_{\mathcal{O}_K} k$ denotes the unique irreducible algebraic G -submodule.

13.2 We also recall from the introduction the \mathbb{F} -representation $V(\mu, \tau)$ of $GL_d(k)$ attached to any pair (μ, τ) with μ a Hodge type with each $\mu_{\kappa} - \rho$ dominant and τ an inertial type. Taking $\tau = 1$ we obtain $V(\mu, 1)$ by evaluating the algebraic representation

$$\bigotimes_{\kappa} (H^0(G/P_{\mu_{\kappa}-\rho}, \mathcal{L}(\mu_{\kappa} - \rho)) \otimes_{\mathcal{O}_{K,\kappa}} \mathbb{F})$$

(here we write κ also for its composite with the surjection $\mathcal{O} \rightarrow \mathbb{F}$) on k -points. Since the exact definition of $V(\mu, \tau)$ will not be needed for $\tau \neq 1$ we refer to [7, 8.2] for the general construction.

Lemma 13.3 *Suppose $d = 2$ and μ is a Hodge type with $\mu_{\kappa} \subset [0, r_{\kappa}]$ for $r_{\kappa} \geq 0$ satisfying*

$$\sum_{\kappa|k=\kappa_0} r_{\kappa} \leq e + p - 1$$

Then the multiplicity of λ in $V(\mu, 1)$, for λ an absolutely irreducible \mathbb{F} -representation of $GL_d(k)$ corresponding to (λ_{κ_0}) under 13.1, equals the product

$$\prod_{\kappa_0} m(\lambda_{\kappa_0}, (\mu_{\kappa})_{\kappa|k=\kappa_0})$$

from Proposition 6.6.

Proof Recall from Lemma 6.3 that each $m(\lambda_{\kappa_0}, (\mu_{\kappa})_{\kappa|k=\kappa_0})$ can be interpreted as the multiplicity of $H^0(\lambda_{\kappa_0})$ in $\bigotimes_{\kappa|k=\kappa_0} H^0(\mu_{\kappa} - \rho)$ where $H^0(-)$ denotes the generic fibre of $H^0(G/P_{-}, \mathcal{L}(-))$. This coincides with the corresponding multiplicities on the special fibre, i.e. the multiplicity of $H^0(G/P_{\lambda_{\kappa_0}}, \mathcal{L}(\lambda_{\kappa_0})) \otimes_{\mathcal{O}_K} k$ in

$$\bigotimes_{\kappa|k=\kappa_0} (H^0(G/P_{\lambda_{\mu_{\kappa}-\rho}}, \mathcal{L}(\mu_{\kappa} - \rho)) \otimes_{\mathcal{O}_K} k)$$

The product of these multiplicities therefore equals the multiplicity of

$$\bigotimes_{\kappa_0} H^0(G/P_{\lambda_{\kappa_0}}, \mathcal{L}(\lambda_{\kappa_0})) \otimes_{\mathcal{O}_{K, \kappa_0}} \mathbb{F}$$

inside

$$\bigotimes_{\kappa} \left(H^0(G/P_{\lambda_{\mu_{\kappa} - \rho}}, \mathcal{L}(\mu_{\kappa} - \rho)) \otimes_{\mathcal{O}_{K, \kappa}} \mathbb{F} \right)$$

The lemma will therefore follow if each $H^0(G/P_{\lambda_{\kappa_0}}, \mathcal{L}(\lambda_{\kappa_0})) \otimes_{\mathcal{O}_K} k$ is simple, i.e. equals $L(\lambda_{\kappa_0})$. Since $d = 2$ this can be seen from the explicit description given in, for example, [15, II.2.16]. \square

Notation 13.4 If λ denotes an isomorphism class of absolutely irreducible \mathbb{F} -representation of $GL_d(k)$ then we write $\tilde{\lambda}$ for the Hodge type obtained as in 6.5 for (λ_{κ_0}) the tuple corresponding to λ in 13.1.

13.5 In the next proposition we fix a continuous homomorphism $\bar{\rho} : G_K \rightarrow GL_d(\mathbb{F})$ and, as in the proof of Lemma 10.9, we write $R_{\bar{\rho}}$ for the \mathcal{O} -framed deformation ring of $\bar{\rho}$. If (μ, τ) is a pair consisting of a Hodge type μ and an inertial type τ then we also write $R_{\bar{\rho}}^{\mu, \tau}$ for the unique reduced \mathcal{O} -flat quotient of $R_{\bar{\rho}}$ whose points valued in a finite extension of E are correspond to potentially crystalline representations of type (μ, τ) .

We also say that an absolutely irreducible representation \mathbb{F} -representation λ of $GL_2(k)$ is non-Steinberg if λ corresponds to a tuple (λ_{κ_0}) with $\lambda_{\kappa_0,1} - \lambda_{\kappa_0,2} \neq p - 1$ for at least one κ_0 .

Proposition 13.6 Assume $p > 2$, $d = 2$, and that $\mu_{\kappa} - \rho$ is dominant for each κ . Then

- (1) There are cycles $\mathcal{C}_{\bar{\rho}, \lambda}$ in $\text{Spec } R_{\bar{\rho}}$, indexed by isomorphism classes of absolutely irreducible \mathbb{F} -representations λ of $GL_d(k)$, such that, for any pair (μ, τ) , one has an inequality

$$[\text{Spec } R_{\bar{\rho}}^{\mu, \tau} \otimes_{\mathcal{O}} \mathbb{F}] \geq \sum_{\lambda} m(\lambda, \mu, \tau) \mathcal{C}_{\bar{\rho}, \lambda}$$

for $m(\lambda, \mu, \tau)$ the multiplicity of λ in $V(\mu, \tau)$ (where $V(\mu, \tau)$ is the $GL_d(k)$ -representation attached to μ, τ).

- (2) If λ is non-Steinberg then $\mathcal{C}_{\bar{\rho}, \lambda} = [\text{Spec } R_{\bar{\rho}}^{\tilde{\lambda}, 1} \otimes_{\mathcal{O}} \mathbb{F}]$.

This is the single point where the assumption $p > 2$ arises.

Proof In [7, 8.6.6] it is shown that, if $\mathcal{X}_2^{\mu, \tau}$ denotes the closed algebraic substack of the Emerton–Gee stack \mathcal{X}_2 whose A -points (for A a finite flat \mathcal{O} -algebra) correspond to potentially crystalline G_K -representations of type (μ, τ) , then there are top dimensional cycles $\mathcal{C}_{\lambda} \subset \overline{\mathcal{X}}_2$ such that

$$[\overline{\mathcal{X}}_2^{\mu, \tau}] \geq \sum_{\lambda} m(\lambda, \mu, \tau) \mathcal{C}_{\lambda}$$

The cycles C_λ are also described explicitly in [7, 8.6.2]. In particular, if λ is non-Steinberg then C_λ is the irreducible component of $\overline{\mathcal{X}}_2$ labelled by λ as in [7, 5.5.11]. As explained in [7, 8.3] the above inequality implies part (1) of the proposition by pulling back along the formally smooth morphism $\mathrm{Spf} R_{\overline{\rho}} \otimes_{\mathbb{O}} \mathbb{F} \rightarrow \overline{\mathcal{X}}_2$.

For part (2), we observe that the irreducibility of $M_{\tilde{\lambda}} \otimes_{\mathbb{O}} \mathbb{F}$ implies, for non-Steinberg λ , irreducibility of $Y_2^{\tilde{\lambda}} \otimes_{\mathbb{O}} \mathbb{F}$ (see the paragraph before (14.3)). By definition $\mathcal{X}_2^{\tilde{\lambda},1}$ is the scheme theoretic image of Y_2^{μ} and so the topological space of $\overline{\mathcal{X}}_2^{\tilde{\lambda},1}$ is the image of that of $Y_2^{\tilde{\lambda}} \otimes_{\mathbb{O}} \mathbb{F}$. Therefore $\overline{\mathcal{X}}_2^{\tilde{\lambda},1}$ is irreducible and $[\overline{\mathcal{X}}_2^{\tilde{\lambda},1}] = m_\lambda C_\lambda$ for some $m_\lambda \in \mathbb{Z}_{\geq 0}$. We have to show $m_\lambda = 1$.

If $\overline{\rho}$ is a smooth point of C_λ then pulling back this identity along $\mathrm{Spf} R_{\overline{\rho}} \otimes_{\mathbb{O}} \mathbb{F} \rightarrow \overline{\mathcal{X}}_2$ and taking Hilbert–Samuel multiplicities gives $m_\lambda = e(R_{\overline{\rho}}^{\tilde{\lambda},1} \otimes_{\mathbb{O}} \mathbb{F})$ (smoothness of C_λ at $\overline{\rho}$ implying the Hilbert–Samuel multiplicity of C_λ at this $\overline{\rho}$ is 1). We therefore have to show $e(R_{\overline{\rho}}^{\tilde{\lambda},1} \otimes_{\mathbb{O}} \mathbb{F}) = 1$. As explained in [7, 6.5.1], the component of $\overline{\mathcal{X}}_2$ representing C_λ contains an open substack consisting of representations which are maximally non-split of niveau 1. We can therefore assume the smooth point $\overline{\rho}$ chosen above is of this form. Since λ is non-Steinberg we ensure $\overline{\rho}$ is not of the form

$$\psi \otimes \begin{pmatrix} 1 & * \\ 0 & \chi_{\mathrm{cyc}}^{-1} \end{pmatrix}$$

for ψ an unramified character and χ_{cyc} the mod p cyclotomic character. This means $\overline{\rho}$ is cyclotomic-free in the sense of [1, 1.1.1] and so [1, 1.1.2] ensures every crystalline lift of $\overline{\rho}$ with Hodge type $\tilde{\lambda}$ is potentially diagonalisable. Combined with [10, 3.5.5] this implies $e(R_{\overline{\rho}}^{\tilde{\lambda},1} \otimes_{\mathbb{O}} \mathbb{F}) = \mu_\lambda(\overline{\rho})$ where $\mu_\lambda(\overline{\rho})$ equals the Hilbert–Samuel multiplicity of $C_{\lambda,\overline{\rho}}$. Thus $e(R_{\overline{\rho}}^{\tilde{\lambda},1} \otimes_{\mathbb{O}} \mathbb{F}) = 1$ as desired. □

14 Main result

We can now prove our main result.

Theorem 14.1 *Assume $p > 2$ and $d = 2$. Let μ be a Hodge type with each $\mu_\kappa - \rho$ dominant and*

$$\sum_{\kappa \mid k = \kappa_0} (\mu_{\kappa,1} - \mu_{\kappa,2} - 1) \leq p$$

for each $\kappa_0 : k \rightarrow \mathbb{F}$. Then

$$[\mathrm{Spec} R_{\overline{\rho}}^{\mu,1} \otimes_{\mathbb{O}} \mathbb{F}] = \sum_{\lambda} m(\lambda, \mu, 1) [\mathrm{Spec} R_{\overline{\rho}}^{\tilde{\lambda},1} \otimes_{\mathbb{O}} \mathbb{F}]$$

for $m(\lambda, \mu, 1)$ the multiplicity of λ in $V(\mu, 1)$.

Proof First, by a simple twisting argument, we can assume $\mu_{\kappa,2} = 0$ for each κ . Since $\mu_{\kappa,1} \geq 1$ the bound $\sum_{\kappa|k=\kappa_0} \mu_{\kappa,1} \leq p$ implies that either $e < p$ or $e = p$ and $\mu_{\kappa,1} = 1$ for each κ . In the latter case $\mu = \tilde{\lambda}$ for λ the trivial representation and in this case there is nothing to prove. Thus, we can assume $e < p$. This means K is tamely ramified over \mathbb{Q}_p and so Remark 12.2 indicates that Theorem 12.1 applies. We can also assume that $e > 1$, since if $e = 1$ then again $\mu = \tilde{\lambda}$ and the theorem is again trivial. This means that $\sum_{\kappa|k=\kappa_0} \mu_{\kappa,1} < p + e - 1$ for each κ and so Proposition 11.3 also applies, with $r_{\kappa} = \mu_{\kappa,1}$.

Proposition 7.2 gives an identity of cycles

$$[M_{\mu} \otimes_{\mathcal{O}} \mathbb{F}] = \sum_{\lambda} n(\lambda, \mu) [M_{\tilde{\lambda}} \otimes_{\mathcal{O}} \mathbb{F}]$$

for integers $n(\lambda, \mu) \geq 0$ and λ running over tuples (λ_{κ_0}) with $\lambda_{\kappa_0} \leq \sum_{\kappa|k=\kappa_0} (\mu_{\kappa} - \rho)$. Each such λ_{κ_0} then satisfies $\lambda_{\kappa_0,1} - \lambda_{\kappa_0,2} \leq p - 1$ so we can also view the sum as running over absolutely irreducible \mathbb{F} -representations of $GL_2(k)$ by 13.1. Applying the automorphism from Sect. 8 gives

$$[M_{-w_0\mu} \otimes_{\mathcal{O}} \mathbb{F}] = \sum_{\lambda} n(\lambda, \mu) [M_{-w_0\tilde{\lambda}} \otimes_{\mathcal{O}} \mathbb{F}]$$

Proposition 7.7 allows us to view this identity of cycles as occurring within the closed subscheme $Gr_{\mathcal{O}}^{\nabla\sigma,r} \otimes_{\mathcal{O}} \mathbb{F}$ from 11.2 for $r = (r_{\kappa})$. We want to consider its preimage under the composite

$$Y_2^{\leq h} \times_{Z_2^{\leq h}} (\tilde{Z}_2^{\nabla\sigma,r} \otimes_{\mathcal{O}} \mathbb{F}) \rightarrow \tilde{Z}_2^{\nabla\sigma,r} \otimes_{\mathcal{O}} \mathbb{F} \xrightarrow{\Psi_N} Gr_{\mathcal{O}}^{\nabla\sigma,r} \otimes_{\mathcal{O}} \mathbb{F} \tag{14.2}$$

(here the auxiliary integer N is chosen sufficiently large so that Proposition 9.7 applies). To do this we need to show the composite is flat. As the first map is an isomorphism by Proposition 11.3, and \mathcal{G}_N is a smooth and irreducible group scheme, this composite is smooth with irreducible fibres. Smooth morphisms are flat so the pull-back of cycles is well defined and we obtain

$$[Y_2^{\mu, \text{flag}}] = \sum_{\lambda} n(\lambda, \mu) [Y_2^{\tilde{\lambda}, \text{flag}}]$$

where $Y_2^{\mu, \text{flag}}$ denotes the preimage of $M_{-w_0\mu} \otimes_{\mathcal{O}} \mathbb{F}$. These are identities of $\dim \mathcal{G}_N + \sum_{\kappa} \dim G/P_{\mu_{\kappa}}$ -dimensional cycles. Theorem 12.1 shows that

$$\bar{Y}_2^{\mu} \times_{Z_2^{\leq h}} \tilde{Z}_2^{\leq h, N} \hookrightarrow Y_2^{\mu, \text{flag}},$$

from which we conclude that $[\bar{Y}_2^{\mu} \times_{Z_2^{\leq h}} \tilde{Z}_2^{\leq h, N}] \leq [Y_2^{\mu, \text{flag}}]$ as cycles. We also point out that since each $M_{\tilde{\lambda}} \otimes_{\mathcal{O}} \mathbb{F}$ is irreducible and generically reduced (see Proposition 7.2) the same is true of $M_{-w_0\tilde{\lambda}} \otimes_{\mathcal{O}} \mathbb{F}$. The same is then also true of $Y_2^{\tilde{\lambda}, \text{flag}}$ since (14.2) is smooth

with irreducible fibres. In particular, this implies the inequality $[\bar{Y}_2^\mu \times_{Z_2^{\leq h}} \tilde{Z}_2^{\leq h, N}] \leq [Y_2^{\mu, \text{flag}}]$ is an equality when $\mu = \tilde{\lambda}$. As a consequence

$$[\bar{Y}_2^\mu \times_{Z_2^{\leq h}} \tilde{Z}_2^{\leq h, N}] \leq \sum_{\lambda} n(\lambda, \mu) [\bar{Y}_2^\lambda \times_{Z_2^{\leq h}} \tilde{Z}_2^{\leq h, N}] \tag{14.3}$$

as $\dim \mathcal{G}_N + \sum_{\kappa} \dim G/P_{\mu_{\kappa}}$ -dimensional cycles inside the scheme $Y_2^{\leq h} \times_{Z_2^{\leq h}} (\tilde{Z}_2^{\nabla_{\sigma, r}} \otimes_{\mathcal{O}} \mathbb{F})$.

The next goal is to descend this identity to an inequality of cycles in $\text{Spec } R_{\bar{p}} \otimes_{\mathcal{O}} \mathbb{F}$. For this we recall the projective $R_{\bar{p}}$ -scheme $\mathcal{L}_{\bar{p}}^{\leq h}$ introduced in Construction 10.8. The morphism $\mathcal{L}_{\bar{p}}^{\leq h} \otimes_{\mathcal{O}} \mathbb{F} \rightarrow Y_2^{\leq h} \otimes_{\mathcal{O}} \mathbb{F}$ is clearly formally smooth, with relative dimension $d^2 = 4$ arising from the framing variables in $R_{\bar{p}}$. Pulling back the previous inequality along

$$(\mathcal{L}_{\bar{p}}^{\leq h} \otimes_{\mathcal{O}} \mathbb{F}) \times_{Z_2^{\leq h}} \tilde{Z}_2^{\leq h, N} \rightarrow (Y_2^{\leq h} \otimes_{\mathcal{O}} \mathbb{F}) \times_{Z_2^{\leq h}} \tilde{Z}_2^{\leq h, N}$$

(being a formally smooth morphism between Noetherian schemes, this map is flat and so the pull-back is defined) gives an inequality

$$[\bar{\mathcal{L}}_{\bar{p}}^\mu \times_{Z_2^{\leq h}} \tilde{Z}_2^{\leq h, N}] \leq \sum_{\lambda} n(\lambda, \mu) [\bar{\mathcal{L}}_{\bar{p}}^{\tilde{\lambda}} \times_{Z_2^{\leq h}} \tilde{Z}_2^{\leq h, N}]$$

where $\bar{\mathcal{L}}_{\bar{p}}^\mu = \mathcal{L}_{\bar{p}}^\mu \otimes_{\mathcal{O}} \mathbb{F}$ for $\mathcal{L}_{\bar{p}}^\mu$ the preimage of Y_2^μ in $\mathcal{L}_{\bar{p}}^{\leq h}$ (constructed just as in the proof of Lemma 10.9). This is an identity of $d^2 + \dim \mathcal{G}_N + \sum_{\kappa} \dim G/P_{\mu_{\kappa}}$ -dimensional cycles. Since the morphism $\mathcal{L}_{\bar{p}}^{\leq h} \times_{Z_2^{\leq h}} (\tilde{Z}_2^{\leq h, N} \otimes_{\mathcal{O}} \mathbb{F}) \rightarrow \mathcal{L}_{\bar{p}}^{\leq h}$ is a \mathcal{G}_N -torsor (in particular smooth, surjective, and of relative dimension $\dim \mathcal{G}_N$) it follows that

$$[\bar{\mathcal{L}}_{\bar{p}}^\mu] \leq \sum_{\lambda} n(\lambda, \mu) [\bar{\mathcal{L}}_{\bar{p}}^{\tilde{\lambda}}]$$

as $d^2 + \sum_{\kappa} \dim G/P_{\mu_{\kappa}}$ -dimensional cycles inside $\mathcal{L}_{\bar{p}}^{\leq h}$. Recall that the projective morphism $\Theta : \mathcal{L}_{\bar{p}}^{\leq h} \rightarrow \text{Spec } R_{\bar{p}}$ becomes a closed immersion after inverting p and this closed immersion identifies $\mathcal{L}_{\bar{p}}^\mu[\frac{1}{p}] = \text{Spec } R_{\bar{p}}^{\mu, 1}[\frac{1}{p}]$. This was discussed in the proof of Lemma 10.9. Since the $R_{\bar{p}}^{\mu, 1}$ are \mathcal{O} -flat, an application of Lemma 3.3 shows that

$$\Theta_*[\bar{\mathcal{L}}_{\bar{p}}^\mu] = [\text{Spec } R_{\bar{p}}^{\mu, 1} \otimes_{\mathcal{O}} \mathbb{F}]$$

Therefore, pushing forward the previous inequality of cycles gives

$$[\text{Spec } R_{\bar{p}}^{\mu, 1} \otimes_{\mathcal{O}} \mathbb{F}] \leq \sum_{\lambda} n(\lambda, \mu) [\text{Spec } R_{\bar{p}}^{\tilde{\lambda}, 1} \otimes_{\mathcal{O}} \mathbb{F}]$$

now as $d^2 + \sum_{\kappa} \dim G/P_{\kappa}$ -dimensional cycles inside $\text{Spec } R_{\bar{p}} \otimes_{\mathcal{O}} \mathbb{F}$. Proposition 13.6 then gives that $n(\lambda, \mu) \geq m(\lambda, \mu, 1)$. Combining Lemma 13.3 and Proposition 6.6 shows that this must be an equality. The theorem follows. \square

15 Miscellany

Let \mathcal{X} and \mathcal{Y} be algebraic stacks of finite type over a field k and let $f : \mathcal{X} \rightarrow \mathcal{Y}$ be a morphism of stacks.

Lemma 15.1 *If, for A any local finite k -algebra, the induced functor $\mathcal{X}(A) \rightarrow \mathcal{Y}(A)$:*

- (1) *is fully faithful then f is a monomorphism (which by our definition implies being representable by algebraic spaces and separated).*
- (2) *is an equivalence then $\mathcal{X} \rightarrow \mathcal{Y}$ is an isomorphism.*

Proof First we prove (2) under the additional assumption that f is representable by algebraic spaces. Then, by choosing a smooth surjection $U \rightarrow \mathcal{Y}$ with U an algebraic space, we can assume that \mathcal{X} and \mathcal{Y} are algebraic spaces. With this reduction the argument given in [21, 7.2.4] goes through with schemes replaced by algebraic spaces. Indeed, by [26, 0APP] this morphism is smooth and quasi-finite, and hence étale. By [26, 05W1] the diagonal $\mathcal{X} \rightarrow \mathcal{X} \times_{\mathcal{Y}} \mathcal{X}$ is an open immersion. Since it is surjective on finite type points it is an isomorphism and so $\mathcal{X} \rightarrow \mathcal{Y}$ is a monomorphism. By [26, 05W5] it is an open immersion, and so an isomorphism, again by surjectivity on finite type points.

Now we prove (1). By [26, 04XS] the diagonal $\Delta_f : \mathcal{X} \rightarrow \mathcal{X} \times_{\mathcal{Y}} \mathcal{X}$ is representable by algebraic spaces. Full faithfulness of f on A -valued points implies that Δ_f is an equivalence on such points. Therefore the first paragraph implies Δ_f is an isomorphism. From [26, 04ZZ] we obtain (1).

To finish the proof of (2) note that by (1) we have f representable by algebraic spaces. \square

Corollary 15.2 *Suppose \mathcal{Z} is a closed substack of \mathcal{Y} and that for every morphism $\text{Spec } A \rightarrow \mathcal{X}$, with A any local finite k -algebra, the composite $\text{Spec } A \rightarrow \mathcal{X} \rightarrow \mathcal{Y}$ factors through \mathcal{Z} . Then $\mathcal{X} \rightarrow \mathcal{Y}$ factors through \mathcal{Z} .*

Proof This follows since by Lemma 15.1 the map $\mathcal{X} \rightarrow \mathcal{X} \times_{\mathcal{Y}} \mathcal{Z}$ is an isomorphism. \square

Acknowledgements I would like to thank the referee for many helpful comments and corrections. I would also like to thank the Max Planck Institute for Mathematics Bonn, the University of Münster, and the University of Glasgow, where parts of this work was done. In particular, the project was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation)–Project ID 427320536-SFB1442, as well as under Germany’s Excellence Strategy EXC2044390685587, Mathematics Münster: Dynamics–Geometry–Structure.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included

in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Bartlett, R.: Potential diagonalisability of pseudo-Barsotti–Tate representations (2019). Preprint. [arXiv:2001.08660](https://arxiv.org/abs/2001.08660)
2. Bartlett, R.: On the irreducible components of some crystalline deformation rings. *Forum Math. Sigma* **8**, e22 (2020)
3. Beauville, A., Laszlo, Y.: Un lemme de descente. *C. R. Acad. Sci. Paris Sér. I Math.* **320**, 335–340 (1995)
4. Breuil, C., Mézard, A.: Multiplicités modulaires et représentations de $GL_2(\mathbf{Z}_p)$ et de $\text{Gal}(\overline{\mathbf{Q}}_p/\mathbf{Q}_p)$ en $l = p$. *Duke Math. J.* **115**(2), 205–310, With an appendix by Guy Henniart (2002)
5. Emerton, M., Gee, T.: A geometric perspective on the Breuil–Mézard conjecture. *J. Inst. Math. Jussieu* **13**(1), 183–223 (2014)
6. Emerton, M., Gee, T.: ‘Scheme-theoretic images’ of morphisms of stacks. *Algebr. Geom.* **8**(1), 1–132 (2021)
7. Emerton, M., Gee, T.: Moduli Stacks of Étale (ϕ, Γ) -modules and the Existence of Crystalline Lifts. *Annals of Mathematics Studies*, vol. 215. Princeton University Press, Princeton (2023)
8. Fontaine, J.-M.: Le corps des périodes p -adiques. *Astérisque* **223**, 59–111, With an appendix by Pierre Colmez, *Périodes p -adiques (Bures-sur-Yvette, 1988)* (1994)
9. Fulton, W.: Intersection theory, *Second, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*, vol. 2. Springer, Berlin (1998)
10. Gee, T., Kisin, M.: The Breuil–Mézard conjecture for potentially Barsotti–Tate representations. *Forum Math. Pi* **2**, e1, 56 (2014)
11. Gee, T., Liu, T., Savitt, D.: The Buzzard–Diamond–Jarvis conjecture for unitary groups. *J. Am. Math. Soc.* **27**(2), 389–435 (2014)
12. Gee, T., Liu, T., Savitt, D.: The weight part of Serre’s conjecture for $GL(2)$. *Forum Math. Pi* **3**, e2, 52 (2015)
13. Herzig, F.: The weight in a Serre-type conjecture for tame n -dimensional Galois representations. *Duke Math. J.* **149**(1), 37–116 (2009)
14. Hu, Y., Tan, F.: The Breuil–Mézard conjecture for non-scalar split residual representations. *Ann. Sci. Éc. Norm. Supér. (4)* **48**(6), 1383–1421 (2015)
15. Jantzen, J.C.: *Representations of Algebraic Groups, Second, Mathematical Surveys and Monographs*, vol. 107. American Mathematical Society, Providence (2003)
16. Kisin, M.: Crystalline Representations and F -crystals, *Algebraic Geometry and Number Theory, Progress in Mathematics*, vol. 253, pp. 459–496. Birkhäuser, Boston (2006)
17. Kisin, M.: Potentially semi-stable deformation rings. *J. Am. Math. Soc.* **21**(2), 513–546 (2008)
18. Kisin, M.: The Fontaine–Mazur conjecture for GL_2 . *J. Am. Math. Soc.* **22**(3), 641–690 (2009)
19. Kisin, M.: Moduli of finite flat group schemes, and modularity. *Ann. Math. (2)* **170**(3), 1085–1180 (2009)
20. Liu, Q.: *Algebraic Geometry and Arithmetic Curves, Oxford Graduate Texts in Mathematics*, vol. 6. Oxford University Press, Oxford. Translated from the French by Reinie Ern e, Oxford Science Publications (2002)
21. Le, D., Hung, B.V.L., Levin, B., Morra, S.: Local models for Galois deformation rings and applications (2020). [arXiv:2007.05398](https://arxiv.org/abs/2007.05398)
22. Ozeki, Y.: Full faithfulness theorem for torsion crystalline representations. *N. Y. J. Math.* **20**, 1043–1061 (2014)
23. Paškūnas, V.: On the Breuil–Mézard conjecture. *Duke Math. J.* **164**(2), 297–359 (2015)
24. Pappas, G., Rapoport, M.: Φ -modules and coefficient spaces. *Mosc. Math. J.* **9**(3), 625–663 (2009)
25. Sander, F.: Hilbert–Samuel multiplicities of certain deformation rings. *Math. Res. Lett.* **21**(3), 605–615 (2014)

26. The Stacks Project Authors, *stacks project*. <http://stacks.math.columbia.edu> (2017)
27. Tung, S.-N.: On the modularity of 2-adic potentially semi-stable deformation rings. *Math. Z.* **298**(1–2), 107–159 (2021)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.