



Cosmological Einstein- λ -Perfect-Fluid Solutions with *Asymptotic Dust* or *Asymptotic Radiation* Equations of State

Helmut Friedrich

Abstract. This article introduces the notions of *asymptotic dust* and *asymptotic radiation* equations of state. With these non-linear generalizations of the well known *dust* or (incoherent) *radiation* equations of state the perfect-fluid equations lose any conformal covariance or privilege. We analyse the conformal field equations induced with these equations of state. It is shown that the Einstein- λ -perfect-fluid equations with an asymptotic radiation equation of state allow for large sets of Cauchy data that develop into solutions which admit smooth conformal boundaries in the future and smooth extensions beyond. In the case of asymptotic dust equations of state sharp results on the future asymptotic behaviour are not available yet.

1. Introduction

This article extends our investigation of the long term behaviour of solutions to Einstein's equation

$$R_{\mu\nu}[\hat{g}] - \frac{1}{2}R[\hat{g}]\hat{g}_{\mu\nu} + \lambda\hat{g}_{\mu\nu} = \hat{T}_{\mu\nu}, \quad (1.1)$$

with positive cosmological constant λ and energy-momentum tensor $\hat{T}_{\mu\nu}$ for various matter models. The ones considered in the present paper are given by perfect fluids with certain non-linear equations of state. Several authors studied the future asymptotic behaviour of solutions to the Einstein- λ -perfect fluid equations with a linear equation of state $\hat{p} = w(\hat{\rho}) = w_*\hat{\rho}$, $w_* = \text{const.}$ [5, 9, 12, 13, 15, 16], often assuming that $0 < w_* < 1/3$. More general equations of state have been considered in the articles [11, 14]. In the present article

the asymptotic behaviour is analysed from a particular point of view. It is motivated by the following observation.

De Sitter space is a geodesically complete, spatially compact, conformally flat solution of FLRW type to the Einstein- λ -vacuum equations with cosmological constant $\lambda > 0$. It admits smooth conformal boundaries at future and past time-like infinity. It generalizes as follows. Smooth Cauchy data for the same equations on a Cauchy hypersurface S of de Sitter space that are sufficiently close (with respect to suitable Sobolev norms) to the de Sitter data develop into solutions that are also time-like and null geodesically complete and admit smooth conformal boundaries. The *conformal Einstein equations*, which have been used for this purpose, determine in fact smooth solutions that extend beyond these boundaries and define on ‘the other sides’ again vacuum solutions. Perturbations of the conformal curvature, only restricted by the smallness condition on S , travel unimpeded across the boundaries [1, 2].

The question of interest in this article is: For which matter models can be established similar results on the future asymptotic behaviour of solutions to the Einstein- λ -matter equations? For a discussion of the answers obtained so far the reader is referred to the article [6]. The latter can also be read as an extended non-technical introduction to the present article. It explains our interest in the question above and also may be helpful for readers not acquainted with the conformal methods used in the following.

The two simplest cases with a positive answer are the FLRW-type solutions to the Einstein- λ -perfect-fluid equations with the *pure (incoherent) radiation* equation of state $\hat{p} = w(\hat{\rho}) = \frac{1}{3} \hat{\rho}$ and the *pure dust* equation of state $\hat{p} = w(\hat{\rho}) = 0$ (the unusual word *pure* has been added here to avoid confusions with notions introduced below).

These cases have been generalized by showing that these FLRW solutions are future stable in the class of general solutions to the Einstein- λ -perfect-fluid equations with the respective equation of state. The perturbed solutions are not only time-like geodesically future complete but also admit smooth future conformal boundaries and extensions beyond [5, 12]. These cases are, however, still somewhat special. The perfect fluid equations with a pure radiation equation of state considered in [12] are conformally invariant with vanishing trace of the energy momentum tensor. The perfect fluid equations with a pure dust equation of state are not conformally invariant and have an energy momentum tensor with non-vanishing trace. They are, however, *conformally privileged* (see [6]) by being related to a conformally invariant structure which helps establish the result of [5].

It may be reasonable to base models of the universe on solutions to the Einstein- λ -perfect-fluid equations where the future development is determined by one of the equations of state above. It appears unlikely, however, that it makes sense to cover the whole stretch from the Big Bang to future time-like infinity by one and the same linear equation of state. One can expect to need a transition where

$$\hat{p} = w^{**}(\hat{\rho}) \hat{\rho},$$

with some function $w^{**}(\hat{\rho})$ of which we will assume that $0 \leq w^{**}(\hat{\rho}) \leq 1/3$. At late time $w^{**}(\hat{\rho})$ should then acquire the value $w^{**}(\hat{\rho}) = 0$ if we want to model the end by a pure dust equation of state and $w^{**}(\hat{\rho}) = 0$ in the case of pure radiation.

There is nothing, however, which fixes a natural notion of ‘late time’. The only meaningful requirement would be that these values are assumed in the limit when the space-time approaches future time-like infinity. But then the equation of state would still need to recognize where and when this limit will be achieved.

In the conformal analysis of the two FLRW models mentioned above the physical density $\hat{\rho}$ and the conformal density ρ satisfy a relation of the form

$$\hat{\rho} = \Omega^e \rho, \quad (1.2)$$

with $e = 3$ in the case of pure dust and $e = 4$ in the case of pure radiation. These values are chosen because they imply that $\rho = \rho_* = \text{const.} > 0$ (assuming that $\hat{\rho} > 0$ on some initial slice). Since then $\hat{\rho} \rightarrow 0$ at future time-like infinity where $\Omega \rightarrow 0$, the behaviour of $\hat{\rho}$ can be understood as an indicator for the approach to future time-like infinity. In generalizing the situation we shall keep (1.2), hoping it to serve as an indicator function of the far future where $\Omega \rightarrow 0$. The relation $\rho = \rho_* = \text{const.}$ may not be preserved under the generalization considered below and we will have to control that ρ remains positive and bounded in the limit $\Omega \rightarrow 0$. The pure dust and pure radiation equations of state are now generalized as follows.

An *asymptotic dust equation of state* is given by a function of the form

$$\hat{p} = w(\hat{\rho}) = (\hat{\rho}^k w^*(\hat{\rho})) \hat{\rho} \quad \text{with some } k \in \mathbb{N}, \quad (1.3)$$

combined with (1.2) where $e = 3$. It implies $w'(\hat{\rho}) = (1+k)\hat{\rho}^k w^*(\hat{\rho}) + \hat{\rho}^{1+k} (w^*)'(\hat{\rho})$, where here and in the following the notation $' = \partial/\partial\hat{\rho}$ is used.

An *asymptotic radiation equation of state* is given by a function of the form

$$\hat{p} = w(\hat{\rho}) = \left(\frac{1}{3} - \hat{\rho}^k w^*(\hat{\rho}) \right) \hat{\rho} \quad \text{with some } k \in \mathbb{N}, \quad (1.4)$$

combined with (1.2) where $e = 4$. It implies $w'(\hat{\rho}) = \frac{1}{3} - (1+k)\hat{\rho}^k w^*(\hat{\rho}) - \hat{\rho}^{1+k} (w^*)'(\hat{\rho})$. In both cases $w^*(\hat{\rho})$ is a smooth function of $\hat{\rho}$ that satisfies

$$0 < w^*(\hat{\rho}) < c = \text{const.}$$

To ensure that $0 \leq w^{**}(\hat{\rho}) \leq 1/3$ as required above, we would need to impose more detailed conditions on $w^*(\hat{\rho})$ but for the analysis of the effect of the new equations of state in the far future where Ω becomes small the conditions above are sufficient. The limits $w^*(\hat{\rho}) \rightarrow 0$ give back the pure dust and the pure radiation equations of state.

The factor $\hat{\rho}^k$ has been inserted in the definitions as a simple means to control in terms of k the speed at which the pure dust or the pure radiation situations is approximated as $\hat{\rho} \rightarrow 0$. Eventually the conformal field equations may force us to impose less crude conditions on $w^*(\hat{\rho})$ or to introduce more subtle definitions of asymptotic equations of state.

Definition (1.3) implies

$$\hat{T} = \hat{g}^{\mu\nu} \hat{T}_{\mu\nu} = 3w(\hat{\rho}) - \hat{\rho} = -\Omega^3 \rho + 3\Omega^{3+3k} \rho^{1+k} w^*(\Omega^3 \rho),$$

while definition (1.4) gives

$$\hat{T} = -3\Omega^{4+4k} \rho^{1+k} w^*(\Omega^4 \rho).$$

In both case $\hat{T} \neq 0$ if $\rho > 0$ and $\hat{T} \rightarrow 0$ as $\Omega \rightarrow 0$ if ρ remains bounded in this limit.

Because the principal parts of the matter equations are affected by the equations of state above any conformal covariance or privilege is lost.

The Cauchy problem local in time for Einstein- λ -perfect fluids with an asymptotic dust or radiation equations of state as above poses no problems. This follows from the results of [3, 8] where only weak conditions on the equation of state are assumed.

First conditions on the admissible values of k are found if the equations of state above are considered in the conformal analogues of the FLRW Friedmann and the energy conservation equation. In the case of asymptotic dust the system reads

$$(\dot{\Omega})^2 = \frac{\lambda}{3} - \frac{R[\hat{h}]}{6} \Omega^2 + \Omega^3 \frac{\rho}{3}, \quad \dot{\rho} = \Omega^{3k-1} \rho^{1+k} w^*(\Omega^3 \rho) \dot{\Omega},$$

where $R[\hat{h}] = \text{const.} \geq 0$. It can be integrated across $\Omega = 0$ with ρ bounded and positive if $3k - 1 \in \mathbb{N}_0$. In the case of asymptotic radiation the system reads

$$(\dot{\Omega})^2 = \frac{\lambda}{3} - \frac{R[\hat{h}]}{6} \Omega^2 + \Omega^4 \frac{\rho}{3}, \quad \dot{\rho} = -\Omega^{4k-1} \rho^{1+k} w^*(\Omega^4 \rho) \dot{\Omega}.$$

It can be integrated across $\Omega = 0$ with ρ bounded and positive if $4k - 1 \in \mathbb{N}_0$.

In the following sections we derive the conformal Einstein- λ -perfect fluid equations, introduce a gauge that involves in particular an orthonormal frame e_j , $j = 0, 1, 2, 3$, with $e_0 = U$, and discuss the regularity as $\Omega \rightarrow 0$ of the equations in this gauge with any of the two asymptotic equations of state assumed.

In the case of an asymptotic dust equation of state there arise problems. The use of this equation of state affects the principal part of the fluid equations in a way which does not allow us to apply the methods which were successful in the case of the pure dust equation of state [5]. This case, which I consider as particularly interesting (see [6]), is left open.

In the case of an asymptotic radiation equation of state regularity of the equations can be ensured by a suitable assumption and there can be derived a symmetric hyperbolic reduced system that is well defined irrespective of the sign of Ω . This allows us to give the following answer to the question posed in the beginning.

Consider the reduced system of the conformal Einstein- λ -perfect-fluid equations in the gauge discussed below with an asymptotic radiation equation of

state where $k \geq 1$. On a compact 3-dimensional manifold \mathcal{J}^+ let be given smooth Cauchy data for the reduced equations with $\Omega = 0$, $U^k = \delta^k_0$ time-like future directed, and $U^k \nabla_k \Omega < 0$ that satisfy the constraints induced by the conformal field equations and the special properties implied on a space-like hypersurface on which $\Omega = 0$. These data determine a smooth solution to the reduced equations with $\Omega < 0$ in the future of \mathcal{J}^+ and $\Omega > 0$ in the past of \mathcal{J}^+ . Where $\Omega > 0$ the solution defines a unique solution to the system of the Einstein- λ -perfect-fluid equations with an asymptotic radiation equation of state that is time-like geodesically future complete and for which \mathcal{J}^+ represents the conformal boundary at the infinite time-like future.

Let S be a Cauchy hypersurface for this solution in the past of \mathcal{J}^+ and denote by Δ the Cauchy data induced by the solution on S . Any Cauchy data Δ' on S for the same system which are sufficiently close to Δ develop into a solution which is also time-like geodesically future complete, admits a smooth conformal boundary in its future, and a smooth conformal extension beyond.

At the end of this article are discussed observations about solutions for which U is orthogonal to the hypersurface \mathcal{J}^+ and consequently hypersurface orthogonal on physical solution in the past of \mathcal{J}^+ .

2. The Conformal Einstein- λ -Euler System

We consider the Einstein- λ -perfect fluid system with cosmological constant λ that consists of the Einstein equations (1.1) with cosmological constant $\lambda > 0$ and an energy momentum tensor of a simple ideal fluid given by

$$\hat{T}_{\mu\nu} = (\hat{\rho} + \hat{p}) \hat{U}_\mu \hat{U}_\nu + \hat{p} \hat{g}_{\mu\nu}.$$

The unknowns are a Lorentz metric $\hat{g}_{\mu\nu}$ on a four-dimensional manifold \hat{M} , a future directed time-like fluid flow vector field \hat{U}^μ satisfying $\hat{U}_\mu \hat{U}^\mu = -1$, the total energy density $\hat{\rho}$ and the pressure \hat{p} as measured by co-moving observers. The system is completed by an equation of state

$$\hat{p} = w(\hat{\rho}) \geq 0. \quad (2.1)$$

The matter equations, equivalent to $\hat{\nabla}^\mu \hat{T}_{\mu\nu} = 0$, are given by

$$\hat{U}^\mu \hat{\nabla}_\mu \hat{\rho} + (\hat{\rho} + \hat{p}) \hat{\nabla}_\mu \hat{U}^\mu = 0, \quad (2.2)$$

$$(\hat{\rho} + \hat{p}) \hat{U}^\mu \hat{\nabla}_\mu \hat{U}_\nu + \{\hat{U}^\mu \hat{U}_\nu + \hat{g}^\mu{}_\nu\} \hat{\nabla}_\mu \hat{p} = 0. \quad (2.3)$$

We shall assume that $\hat{\rho} > 0$ on an initial Cauchy slice.

Let \hat{e}_a , $a = 1, 2, 3$, be vector fields that satisfy $\hat{g}(\hat{U}, \hat{e}_a) = 0$, $\hat{g}(\hat{e}_a, \hat{e}_b) = \delta_{ab}$ and $F = F(\hat{\rho})$ a function that satisfies

$$F' = -(\hat{\rho} + w)^{-1} w', \quad (2.4)$$

where we use, as in the following chapters, the notation $' = \frac{\partial}{\partial \hat{\rho}}$. With (2.3) follows

$$\hat{f}_a \equiv \hat{U}^\mu \hat{e}^\nu{}_a \hat{\nabla}_\mu \hat{U}_\nu = \hat{e}_a(F). \quad (2.5)$$

Equations (2.2), (2.3) imply

$$w' \hat{\nabla}_\nu \hat{\rho} = (\hat{\rho} + w) \left(w' \hat{U}_\nu \hat{\nabla}_\rho \hat{U}^\rho - \hat{U}^\rho \hat{\nabla}_\rho \hat{U}_\nu \right). \quad (2.6)$$

Taking a derivative and observing that $\hat{\rho} + w > 0$ we get the equation

$$\begin{aligned} w'' \hat{\nabla}_\mu \hat{\rho} \hat{\nabla}_\nu \hat{\rho} + w' \hat{\nabla}_\mu \hat{\nabla}_\nu \hat{\rho} &= w' \frac{1+w'}{\hat{\rho}+w} \hat{\nabla}_\mu \hat{\rho} \hat{\nabla}_\nu \hat{\rho} \\ &+ (\hat{\rho} + w) \left(w'' \hat{\nabla}_\mu \hat{\rho} \hat{U}_\nu \hat{\nabla}_\rho \hat{U}^\rho + w' \hat{\nabla}_\mu \hat{U}_\nu \hat{\nabla}_\rho \hat{U}^\rho + w' \hat{U}_\nu \hat{\nabla}_\mu \hat{\nabla}_\rho \hat{U}^\rho \right) \\ &- (\hat{\rho} + w) \left(\hat{\nabla}_\mu \hat{U}^\rho \hat{\nabla}_\rho \hat{U}_\nu + \hat{U}^\rho \hat{\nabla}_\mu \hat{\nabla}_\rho \hat{U}_\nu \right). \end{aligned}$$

Where $w' \neq 0$ anti-symmetrization of this relation implies the integrability condition

$$\hat{U}^\rho \hat{\nabla}_\rho \hat{\nabla}_{[\mu} \hat{U}_{\nu]} - \hat{\nabla}_\rho \hat{U}_{[\mu} \hat{\nabla}_{\nu]} \hat{U}^\rho = \quad (2.7)$$

$$w' \hat{\nabla}_{[\mu} \hat{U}_{\nu]} \hat{\nabla}_\rho \hat{U}^\rho - w' \hat{U}_{[\mu} \hat{\nabla}_{\nu]} \hat{\nabla}_\rho \hat{U}^\rho - w'' \frac{\hat{\rho}+w}{w'} \hat{\nabla}_\lambda \hat{U}^\lambda \hat{U}^\rho \hat{\nabla}_\rho \hat{U}_{[\mu} \hat{U}_{\nu]}.$$

Conformal transformations of the curvature fields.

The conformal Weyl tensor $C^\mu{}_{\nu\lambda\eta}$ and the Schouten tensor $L_{\mu\nu} = 1/2 \left(R_{\mu\nu} - 1/6 R g_{\mu\nu} \right)$ of a given metric satisfy the contracted Bianchi identity

$$\nabla_\mu C^\mu{}_{\nu\lambda\eta} = 2 \nabla_{[\lambda} L_{\eta]\nu}.$$

Under the rescaling

$$\hat{g}_{\mu\nu} \rightarrow g_{\mu\nu} = \Omega^2 \hat{g}_{\mu\nu}, \quad \Omega > 0,$$

the conformal Weyl tensor is invariant, $C^\mu{}_{\nu\lambda\eta} = \hat{C}^\mu{}_{\nu\lambda\eta}$, the Ricci tensor transform as

$$\hat{R}_{\mu\nu} \rightarrow R_{\mu\nu} = \hat{R}_{\mu\nu} - 2 \Omega^{-1} \nabla_\mu \nabla_\nu \Omega - g^{\lambda\delta} \left(\Omega^{-1} \nabla_\lambda \nabla_\delta \Omega - 3 \Omega^{-2} \nabla_\lambda \Omega \nabla_\delta \Omega \right) g_{\mu\nu},$$

and the Schouten tensor as

$$\hat{L}_{\mu\nu} \rightarrow L_{\mu\nu} = \hat{L}_{\mu\nu} - \Omega^{-1} \nabla_\mu \nabla_\nu \Omega + 1/2 \Omega^{-2} g^{\lambda\delta} \nabla_\lambda \Omega \nabla_\delta \Omega g_{\mu\nu}. \quad (2.8)$$

The conformal Weyl tensor satisfies $\nabla_\mu \left(\Omega^{-1} C^\mu{}_{\nu\lambda\eta} \right) = \Omega^{-1} \hat{\nabla}_\mu \hat{C}^\mu{}_{\nu\lambda\eta}$ where ∇ and $\hat{\nabla}$ denote the Levi-Civita connections of $g_{\mu\nu}$ and $\hat{g}_{\mu\nu}$. The identity above thus implies for the *rescaled conformal Weyl tensor* $W^\mu{}_{\nu\lambda\eta} = \Omega^{-1} C^\mu{}_{\nu\lambda\eta}$ the relation

$$\nabla_\mu W^\mu{}_{\nu\lambda\eta} = 2 \Omega^{-1} \hat{\nabla}_{[\lambda} \hat{L}_{\eta]\nu}, \quad (2.9)$$

whence

$$2 \nabla_{[\lambda} L_{\eta]\nu} - \nabla_\mu \Omega W^\mu{}_{\nu\lambda\eta} = 2 \hat{\nabla}_{[\lambda} \hat{L}_{\eta]\nu}. \quad (2.10)$$

Conformal transformation of the matter fields.

We combine the conformal rescaling of the metric with the rescalings

$$U^\mu = \Omega^{-1} \hat{U}^\mu \quad U_\mu = \Omega \hat{U}_\mu, \quad \rho = \Omega^{-e} \hat{\rho}, \quad \text{with } e = \text{const.} > 0.$$

In the following it will always be understood that

$$w = w(\hat{\rho}) = w(\Omega^e \rho), \quad w' = w'(\hat{\rho}) = w'(\Omega^e \rho),$$

and similarly with the function w^* introduced later. Equations (1.1) imply

$$\hat{T} = \hat{g}^{\mu\nu} \hat{T}_{\mu\nu} = 3w - \hat{\rho}, \quad \hat{R} = 4\lambda - \hat{T} = 4\lambda + \hat{\rho} - 3w,$$

and $\hat{L}_{\mu\nu} = \hat{L}_{\mu\nu}^* + \frac{1}{6} \lambda \hat{g}_{\mu\nu}$ with

$$\hat{L}_{\mu\nu}^* = \frac{1}{2} (\hat{\rho} + w) \hat{U}_\mu \hat{U}_\nu + \frac{1}{6} \hat{\rho} \hat{g}_{\mu\nu} = \Omega^{-2} \left(\frac{1}{2} (\hat{\rho} + w) U_\mu U_\nu + \frac{1}{6} \hat{\rho} g_{\mu\nu} \right). \quad (2.11)$$

It follows that $\hat{\nabla}_\lambda \hat{L}_{\eta\nu} = \hat{\nabla}_\lambda \hat{L}_{\eta\nu}^*$. The Ricci scalar of the conformal metric satisfies

$$6\Omega \nabla_\mu \nabla^\mu \Omega + \Omega^2 R - 12 \nabla_\mu \Omega \nabla^\mu \Omega = 4\lambda + \hat{\rho} - 3w.$$

Written in the form

$$6\Omega s - 3 \nabla_\mu \Omega \nabla^\mu \Omega = \lambda + \frac{1}{4} (\hat{\rho} - 3w) \quad \text{with} \quad s = \frac{1}{4} \nabla_\mu \nabla^\mu \Omega + \frac{1}{24} \Omega R, \quad (2.12)$$

it will be referred to as λ -equation. Equations (2.8), (2.11), (2.12) give

$$\nabla_\mu \nabla_\nu \Omega = -\Omega L_{\mu\nu} + s g_{\mu\nu} + \Omega^{-1} \frac{\hat{\rho} + w}{2} \left(U_\mu U_\nu + \frac{1}{4} g_{\mu\nu} \right) \quad (2.13)$$

which will be referred to as Ω -equation. Applying a derivative to the λ -equation (2.12) gives with (2.13) the s -equation

$$\begin{aligned} \nabla_\mu s + L_{\mu\nu} \nabla^\nu \Omega &= \Omega^{-2} \frac{\hat{\rho} + w}{2} \left(U_\mu U^\nu \nabla_\nu \Omega + \frac{1}{4} \nabla_\mu \Omega \right) \\ &+ \frac{1}{24} (1 - 3w') \Omega^{-1} \nabla_\mu \hat{\rho}. \end{aligned} \quad (2.14)$$

It follows

$$\hat{\nabla}_{[\lambda} \hat{L}_{\eta]\nu} = \nabla_{[\lambda} \hat{L}_{\eta]\nu}^* - \Omega^{-1} \left(\hat{L}_{\nu[\lambda}^* \nabla_{\eta]} \Omega + g_{\nu[\lambda} \hat{L}_{\eta]\rho}^* \nabla^\rho \Omega \right),$$

which gives with (2.11) and $\nabla_\lambda \hat{\rho} = \Omega^e (\nabla_\lambda \rho + e \rho \Omega^{-1} \nabla_\lambda \Omega)$ the relation

$$\begin{aligned} 2 \hat{\nabla}_{[\lambda} \hat{L}_{\eta]\nu} &= \Omega^{-3} (\hat{\rho} + w) (U_\nu U_{[\lambda} \nabla_{\eta]} \Omega - g_{\nu[\lambda} U_{\eta]} U_\rho \nabla^\rho \Omega) \\ &+ e \rho \Omega^{e-3} \left((1 + w') \nabla_{[\lambda} \Omega U_{\eta]} U_\nu + \frac{1}{3} \nabla_{[\lambda} \Omega g_{\eta]\nu} \right) \\ &+ \Omega^{-2} (\hat{\rho} + w) (\nabla_{[\lambda} U_{\eta]} U_\nu + U_{[\eta} \nabla_{\lambda]} U_\nu) \\ &+ \Omega^{e-2} \left((1 + w') \nabla_{[\lambda} \rho U_{\eta]} U_\nu + \frac{1}{3} \nabla_{[\lambda} \rho g_{\eta]\nu} \right). \end{aligned} \quad (2.15)$$

The conformal matter equations

Equations (2.2) and (2.3) transform with $\nabla_\mu \hat{\rho} = \Omega^e \{ \nabla_\mu \rho + e \rho \Omega^{-1} \nabla_\mu \Omega \}$ into

$$0 = U^\mu \nabla_\mu \rho + (\rho + \Omega^{-e} w) \nabla_\mu U^\mu - \Omega^{-1} (-e \rho + 3 \rho + 3 \Omega^{-e} w) U^\mu \nabla_\mu \Omega, \quad (2.16)$$

$$0 = (\rho + \Omega^{-e} w) U^\mu \nabla_\mu U_\nu + w' (g^\mu{}_\nu + U^\mu U_\nu) \nabla_\mu \rho - \Omega^{-1} (\rho + \Omega^{-e} w - e \rho w') (g^\mu{}_\nu + U^\mu U_\nu) \nabla_\mu \Omega. \quad (2.17)$$

In the case of pure dust, where $w = 0$ and $e = 3$, the equations read

$$0 = U^\mu \nabla_\mu \rho + \rho \nabla_\mu U^\mu, \quad 0 = \rho U^\mu \nabla_\mu U_\nu - \Omega^{-1} \rho (g^\mu{}_\nu + U^\mu U_\nu) \nabla_\mu \Omega.$$

The Ω^{-1} term in the second equation reflects the conformal non-covariance of the system. In the case of pure radiation with $w = \frac{1}{3} \hat{\rho}$, and $e = 4$ the conformal equations reduce to

$$U^\mu \nabla_\mu \rho + \frac{4}{3} \rho \nabla_\mu U^\mu = 0, \quad \frac{4}{3} \rho U^\mu \nabla_\mu U_\nu + \frac{1}{3} (g^\mu{}_\nu + U^\mu U_\nu) \nabla_\mu \rho = 0,$$

and have thus the same form as their physical versions.

Equation (2.7) transforms into

$$\begin{aligned} & U^\rho \nabla_\rho \nabla_{[\mu} U_{\nu]} + w' U_{[\mu} \nabla_{\nu]} \nabla_\rho U^\rho \\ & - \nabla_\pi U_{[\mu} \nabla_{\nu]} U^\pi - (1 - 3 w') U_{[\mu} L_{\nu]\rho} U^\rho - w' \nabla_{[\mu} U_{\nu]} \nabla_\pi U^\pi \\ & = +(1 - 3 w') \left\{ \Omega^{-1} \nabla_{[\mu} U_{\nu]} U^\rho \nabla_\rho \Omega - \Omega^{-1} U_{[\mu} \nabla_{\nu]} \right. \\ & \quad \left. U_\rho \nabla^\rho \Omega + \Omega^{-2} U_{[\mu} \nabla_{\nu]} \Omega U^\rho \nabla_\rho \Omega \right\} \\ & - w'' \frac{\hat{\rho} + w}{w'} \left\{ \nabla_\pi U^\pi U^\rho \nabla_\rho U_{[\mu} U_{\nu]} - 3 \Omega^{-1} U^\pi \nabla_\pi \Omega U^\rho \nabla_\rho U_{[\mu} U_{\nu]} \right. \\ & \quad \left. - \Omega^{-1} \nabla_\pi U^\pi \nabla_{[\mu} \Omega U_{\nu]} + 3 \Omega^{-2} U^\pi \nabla_\pi \Omega \nabla_{[\mu} \Omega U_{\nu]} \right\}. \end{aligned} \quad (2.18)$$

With $\hat{e}^\mu{}_a = \Omega e^\mu{}_a$ the relation (2.5) transforms into

$$e_a(F) = \Omega^{-1} \hat{f}_a = U^\mu \nabla_\mu U_\nu e^\nu{}_a - e_a(\log \Omega). \quad (2.19)$$

3. The Gauge and the Implied Equations

We express the equations in terms of an orthonormal frame field $e_k = e^\mu{}_k \partial_{x^\mu}$, $k = 0, 1, 2, 3$, so that $g_{jk} \equiv g(e_j, e_k) = \eta_{jk} = \text{diag}(-1, 1, 1, 1)$ and e_0 is a time-like vector field. The space-like frame vector fields are then given by the e_a , where $a, b, c = 1, 2, 3$ denote spatial indices to which the summation convention also applies. In the following all tensor fields considered in the previous sections will be expressed in terms of this frame field. The contravariant coordinate version of the metric is given by $g^{\mu\nu} = \eta^{jk} e^\mu{}_j e^\nu{}_k$.

The connection coefficients, defined by $\nabla_j e_k \equiv \nabla_{e_j} e_k = \Gamma_j^l{}^k e_l$, satisfy $\Gamma_{jlk} = -\Gamma_{jkl}$ with $\Gamma_{jlk} = \Gamma_j^i{}^k g_{il}$, because $\nabla_i g_{jk} = 0$. The covariant derivative of a tensor field $X^\mu{}_\nu$, given in the frame by $X^i{}_j$, takes the form $\nabla_k X^i{}_j = X^i{}_{j,\mu} e^\mu{}_k + \Gamma_k^i{}^l X^l{}_j - \Gamma_k^i{}^l X^i{}_j$. The frame and the connection coefficients satisfy the *first structural equations*

$$e^\mu{}_{i,\nu} e^\nu{}_j - e^\mu{}_{j,\nu} e^\nu{}_i = (\Gamma_j^k{}^i - \Gamma_i^k{}^j) e^\mu{}_k, \quad (3.1)$$

which ensures that the connection is torsion free, and the *second structural equations*

$$\begin{aligned} & \Gamma_l^i{}_{j,\mu} e^\mu{}_k - \Gamma_k^i{}_{j,\mu} e^\mu{}_l + 2\Gamma_{[k}^i{}^p \Gamma_{l]pj} - 2\Gamma_{[k}^p{}^l \Gamma_p^i{}^j] \\ & = \Omega W^i{}_{jkl} + 2\{g^i{}_{[k} L_{l]j} + L^i{}_{[k} g_{l]j}\}. \end{aligned} \quad (3.2)$$

To restrict the gauge freedom for the frame we set $e_0 = U$ so that $U = U^k e_k$ with $U^k = \delta^k_0$, choose at the points of a given smooth space-like hypersurface S transverse to the flow line of U vector fields e_a , $a = 1, 2, 3$, so that $g(e_j, e_k) = \eta_{jk}$, assume the orthonormal frame e_k to be extended by Fermi-transport in the direction of $e_0 = U$ so that $0 = \mathbb{F}_U e_k = \nabla_U e_k - g(e_k, \nabla_U U) U + g(e_k, U) \nabla_U U$, and assume the e_a to be chosen on S so that the resulting orthonormal frame field is smooth. In terms of this frame the Fermi transport law reduces to

$$\Gamma_0^a{}_b = 0. \quad (3.3)$$

To restrict the gauge freedom for the coordinates x^μ , we assume that $\tau \equiv x^0 = 0$ on S and the x^μ be dragged along with U so that

$$\langle e_0, dx^\mu \rangle = e^\mu{}_0 = U^\mu = \delta^\mu_0.$$

The remaining not necessarily vanishing connection coefficients are given by

$$f_a \equiv \Gamma_0^0{}_a = \delta_{ab} \Gamma_0^b{}_0, \quad \chi_{ab} \equiv \Gamma_a^0{}_b = \delta_{bc} \Gamma_a^c{}_0 \quad \text{and} \quad \Gamma_a^b{}_c.$$

It holds $\nabla_i U_k = \Gamma_i^0{}_k = \chi_{ab} \delta^a{}_i \delta^b{}_k + f_b \delta^0{}_i \delta^b{}_k$ and $U^i \nabla_i U_k = f_b \delta^b{}_k$, $\nabla_i U^i = \chi_a^a$.

The first structural equations supply the constraint

$$e^\mu{}_{a,\nu} e^\nu{}_b - e^\mu{}_{b,\nu} e^\nu{}_a = (\Gamma_b^c{}^a - \Gamma_a^c{}^b) e^\mu{}_c + (\chi_{ba} - \chi_{ab}) \delta^\mu_0, \quad (3.4)$$

and the evolution equations

$$e^\mu{}_{a,0} = -\chi_a^b e^\mu{}_b + f_a \delta^\mu_0. \quad (3.5)$$

The second structural equations imply for $\Gamma_a^b{}_c$ and χ_{ab} the constraints

$$\begin{aligned} & \Gamma_b^a{}_{c,\mu} e^\mu{}_d - \Gamma_d^a{}_{c,\mu} e^\mu{}_b + 2\Gamma_{[d}^a{}^i \Gamma_{b]ic} - 2\Gamma_{[d}^i{}^b \Gamma_i^a{}^c] \\ & = \Omega W^a{}_{cdb} + 2\{g^a{}_{[d} L_{b]c} + L^a{}_{[d} g_{b]c}\}, \end{aligned} \quad (3.6)$$

$$\begin{aligned} & \chi_{ab,\mu} e^\mu{}_c - \chi_{cb,\mu} e^\mu{}_a + 2\Gamma_{[c}^0{}^p \Gamma_{a]pb} - 2\Gamma_{[c}^p{}^a \Gamma_p^0{}^b] \\ & + 2L^0{}_{[c} g_{a]b}, \end{aligned} \quad (3.7)$$

and the evolution equations

$$\Gamma_a{}^b{}_{c,0} + f^b \chi_{ac} - \chi_a{}^b f_c + \chi_a{}^d \Gamma_d{}^b{}_{c} = \Omega W^b{}_{c0a} - g^b{}_a L_{0c} + L^b{}_0 g_{ac}, \quad (3.8)$$

$$\chi_{ac,0} - f_{c,\mu} e^\mu{}_a + f_b \Gamma_a{}^b{}_{c} - f_a f_b + \chi_a{}^b \chi_{bc} = \Omega W^0{}_{c0a} + L_{ac} + L^0{}_0 g_{ac}. \quad (3.9)$$

No equation for f_a is implied by the structural equations.

The fluid equations (2.16), (2.17) imply in our gauge the evolution equation

$$\nabla_0 \rho + (\rho + \Omega^{-e} w) \chi_a{}^a - \Omega^{-1} (-e \rho + 3 \rho + 3 \Omega^{-e} w) \nabla_0 \Omega = 0, \quad (3.10)$$

and the constraint

$$(\rho + \Omega^{-e} w) f_a + w' \nabla_a \rho - \Omega^{-1} (\rho + \Omega^{-e} w - e \rho w') \nabla_a \Omega = 0. \quad (3.11)$$

With $\hat{\rho} = \Omega^e \rho$ the latter is seen to be (2.19), which reads with $f_a = U^\mu \nabla_\mu U_\nu e^\nu{}_a$

$$f_a = e_a(F + \log \Omega). \quad (3.12)$$

Equation (2.18) is equivalent to the two equations

$$\begin{aligned} & e_0(f_a) - w' e_a(\chi_c{}^c) + \chi_{ac} f^c + (1 - 3 w') L_{a0} - w' \chi_c{}^c f_a \\ &= (1 - 3 w') (\Omega^{-1} \nabla_0 \Omega f_a + \Omega^{-1} \chi_{ab} \nabla^b \Omega - \Omega^{-2} \nabla_0 \Omega \nabla_a \Omega) \\ & - w'' \frac{\hat{\rho} + w}{w'} (\chi_c{}^c f_a - 3 \Omega^{-1} \nabla_0 \Omega f_a - \Omega^{-1} \chi_c{}^c \nabla_a \Omega + \Omega^{-2} \nabla_0 \Omega \nabla_a \Omega), \end{aligned} \quad (3.13)$$

and

$$e_0(\chi_{[ab]}) - \chi_{c[a} \chi_{b]}{}^c - w' \chi_c{}^c \chi_{[ab]} = (1 - 3 w') \Omega^{-1} \nabla_0 \Omega \chi_{[ab]}. \quad (3.14)$$

With $\chi_{c[a} \chi_{b]}{}^c = -\chi_{[ac]} \sigma_b{}^c + \chi_{[bc]} \sigma_a{}^c$, where $\sigma_{ab} = \chi_{(ab)}$, the latter can be read as a linear homogeneous ODE for $\chi_{[ab]}$.

The system of conformal field equations reads with $e = \text{const}$.

$$6 \Omega s - 3 \nabla_k \Omega \nabla^k \Omega - \lambda = \frac{1}{4} (\hat{\rho} - 3 w), \quad (3.15)$$

$$\nabla_k \nabla_j \Omega + \Omega L_{kj} - s g_{kj} = \Omega^{-1} \frac{\hat{\rho} + w}{2} \left(U_k U_j + \frac{1}{4} g_{kj} \right), \quad (3.16)$$

$$\begin{aligned} & \nabla_k s + \nabla^i \Omega L_{ik} \\ &= \Omega^{-2} \frac{\hat{\rho} + w}{2} \left(U_k U^i \nabla_i \Omega + \frac{1}{4} \nabla_k \Omega \right) + \frac{1}{24} (1 - 3 w') \Omega^{e-1} \\ & \quad \{ \nabla_\mu \rho + e \rho \Omega^{-1} \nabla_k \Omega \}, \end{aligned} \quad (3.17)$$

$$\nabla_i L_{jk} - \nabla_j L_{ik} - \nabla_l \Omega W^l{}_{kij} = 2 \hat{\nabla}_{[i} \hat{L}_{j]k}, \quad (3.18)$$

$$\nabla_l W^l{}_{kij} = 2 \Omega^{-1} \hat{\nabla}_{[i} \hat{L}_{j]k}, \quad (3.19)$$

with

$$\begin{aligned}
 2 \hat{\nabla}_{[i} \hat{L}_{j]k} &= \Omega^{-3} (\hat{\rho} + w) (U_k U_{[i} \nabla_{j]} \Omega - g_{k[i} U_{j]} U^l \nabla_l \Omega) \\
 &+ e \rho \Omega^{e-3} \left((1 + w') \nabla_{[i} \Omega U_{j]} U_k + \frac{1}{3} \nabla_{[i} \Omega g_{j]k} \right) \\
 &+ \Omega^{-2} (\hat{\rho} + w) (\nabla_{[i} U_{j]} U_k + U_{[j} \nabla_{i]} U_k) \\
 &+ \Omega^{e-2} \left((1 + w') \nabla_{[i} \rho U_{j]} U_k + \frac{1}{3} \nabla_{[i} \rho g_{j]k} \right). \quad (3.20)
 \end{aligned}$$

4. Regularity of the Equations

In the following an equation will be called ‘regular’ if no negative or non-integer powers of Ω occur in it. The structural equations are regular in this sense but the tensorial equations can be problematic. I will use from now on the notation

$$\nabla_k \Omega = \Sigma_k. \quad (4.1)$$

The asymptotic dust equation of state.

With the asymptotic dust equation of state and $e = 3$ there enter in various places of the equations functions such as

$$\begin{aligned}
 w &= (\Omega^3 \rho)^{k+1} w^*, & w' &= \Omega^{3k} \rho^k ((k+1) w^* + \Omega^3 \rho (w^*)'), \\
 \hat{\rho} + w &= \Omega^3 \rho (1 + \Omega^{3k} \rho^k w^*), \\
 1 - 3w' &= 1 - 3\Omega^{3k} \rho^k ((k+1) w^* + \Omega^3 \rho (w^*)').
 \end{aligned}$$

As $w^* \rightarrow 0$ they approach the values of the corresponding quantities of the pure dust equation of state. Therefore it is clear that equations which are singular in that case must also be singular in the present case.

In the present case the singular terms cannot be handled as in the pure dust case because equation (2.3) remains a partial differential equation and the Ω^{-1} terms cannot be compensated by suitable choices of k . Moreover, the expression

$$w'' \frac{\hat{\rho} + w}{w'} = (1 + \hat{\rho}^k w^*) \left(k + \hat{\rho} \frac{(2+k)(w^*)' + \hat{\rho}(w^*)''}{(1+k)w^* + \hat{\rho}(w^*)'} \right),$$

which comes with a singular factor in the equation

$$\begin{aligned}
 e_0(f_a) - w' e_a(\chi_c{}^c) + \chi_{ac} f^c + (1 - 3w') L_{a0} - w' \chi_c{}^c f_a \\
 = (1 - 3w') (\Omega^{-1} \Sigma_0 f_a + \Omega^{-1} \chi_{ab} \Sigma^b - \Omega^{-2} \Sigma_0 \Sigma_a) \\
 - w'' \frac{\hat{\rho} + w}{w'} (\chi_c{}^c f_a - 3\Omega^{-1} \Sigma_0 f_a - \Omega^{-1} \chi_c{}^c \Sigma_a + \Omega^{-2} \Sigma_0 \Sigma_a),
 \end{aligned}$$

is not even defined if $w' = 0$. Since there is no obvious way to handle it, this case will not be considered any further in this article.

The asymptotic radiation equation of state.

The asymptotic radiation equation of state with $e = 4$ gives rise to expressions like

$$\begin{aligned}\hat{\rho} + w &= \Omega^4 \rho \left(\frac{4}{3} - (\Omega^4 \rho)^k w^* \right), & w' &= \frac{1}{3} - (\Omega^4 \rho)^k ((k+1)w^* + \Omega^4 \rho (w^*)'), \\ \hat{\rho} - 3w &= 3\Omega^4 (\Omega^4 \rho)^{k+1} \rho^{k+1} w^* (\Omega^4 \rho), & 1 - 3w' &= 3(\Omega^4 \rho)^k ((k+1)w^* + \Omega^4 \rho (w^*)'), \\ w'' \frac{\hat{\rho} + w}{w'} &= -(\Omega^4 \rho)^k (4 - 3\hat{\rho}^k w^*) \frac{k(1+k)w^* + 2(1+k)\hat{\rho}(w^*)' + \hat{\rho}^2 (w^*)''}{1 - 3(1+k)\hat{\rho}^k w^* + 3\hat{\rho}^{1+k} (w^*)'}.\end{aligned}$$

The limits as $w^* \rightarrow 0$ are well defined and yield the corresponding functions in the pure radiation case. If $\rho > 0$ and bounded as $\Omega \rightarrow 0$ we can assume for small $|\Omega|$ that

$$w' = \frac{1}{3} - \Omega^4 \rho^k ((k+1)w^* + \Omega^4 \rho (w^*)') > 0. \quad (4.2)$$

The structural equation are regular with no condition on k . Equations (3.15), (3.16), (3.17) are immediately seen to be regular with $4k \in \mathbb{N}_0$. The fluid equation (3.10) reads

$$\nabla_0 \rho + \rho \left\{ \left(\frac{4}{3} - (\Omega^4 \rho)^k w^* \right) \chi_a^a + 3\Omega^4 k^{-1} \rho^k w^* \right\} \Sigma_0 = 0. \quad (4.3)$$

It is regular if $4k \in \mathbb{N}$. As long as the term in curly brackets is continuous, the function ρ will stay positive if it is positive on some Cauchy surface. With (4.2) the constraint (3.11) can be solved for $\nabla_a \rho$ in a neighbourhood of $\Omega = 0$

$$\nabla_a \rho = -\rho \frac{4 - 3(\Omega^4 \rho)^k w^*}{3w'} f_a - \Omega^4 k^{-1} \rho^{k+1} \frac{(4k+3)w^* + 4\Omega^4 \rho (w^*)'}{w'} \Sigma_a. \quad (4.4)$$

The equation is regular if $4k \in \mathbb{N}$. Inspection of the equations

$$\begin{aligned}e_0(f_a) - w' e_a(\chi_c^c) + \chi_{ac} f^c + (1 - 3w') L_{a0} - w' \chi_c^c f_a \\ = (1 - 3w') (\Omega^{-1} \Sigma_0 f_a + \Omega^{-1} \chi_{ab} \Sigma^b - \Omega^{-2} \Sigma_0 \Sigma_a) \\ - w'' \frac{\hat{\rho} + w}{w'} (\chi_c^c f_a - 3\Omega^{-1} \Sigma_0 f_a - \Omega^{-1} \chi_c^c \Sigma_a + \Omega^{-2} \Sigma_0 \Sigma_a),\end{aligned} \quad (4.5)$$

and

$$e_0(\chi_{[ab]}) - \chi_{c[a} \chi_{b]}^c - w' \chi_c^c \chi_{[ab]} = (1 - 3w') \Omega^{-1} \Sigma_0 \chi_{[ab]}, \quad (4.6)$$

shows that they are regular if $4k - 1 \in \mathbb{N}$. With the expressions above the equations

$$6\Omega s - 3\Sigma_k \Sigma^k - \lambda = \frac{1}{4} (\hat{\rho} - 3w), \quad (4.7)$$

$$\nabla_k \Sigma_j = -\Omega L_{kj} + s g_{kj} + \Omega^{-1} \frac{\hat{\rho} + w}{2} \left(U_k U_j + \frac{1}{4} g_{kj} \right), \quad (4.8)$$

$$\begin{aligned}\nabla_k s + \Sigma^i L_{ik} &= \Omega^{-2} \frac{\hat{\rho} + w}{2} \\ &\left(U_k U^i \Sigma_i + \frac{1}{4} \Sigma_k \right) + \frac{1}{24} (1 - 3w') \Omega^3 \{ \nabla_k \rho + 4\rho \Omega^{-1} \Sigma_k \},\end{aligned} \quad (4.9)$$

are seen to be regular if $4k \in \mathbb{N}_0$. We have finally

$$\nabla_i L_{jk} - \nabla_j L_{ik} - \Sigma_l W^l{}_{kij} = \Omega M_{ijk}, \quad (4.10)$$

$$\nabla_l W^l{}_{kij} = M_{ijk}, \quad (4.11)$$

where

$$\begin{aligned} M_{ijk} &= 2\Omega^{-1} \hat{\nabla}_{[i} \hat{L}_{j]k} = \Omega^{-4} (\hat{\rho} + w) (U_k U_{[i} \Sigma_{j]} - g_{k[i} U_{j]} \Sigma_0) \\ &+ 4\rho \left((1 + w') \Sigma_{[i} U_{j]} U_k + \frac{1}{3} \Sigma_{[i} g_{j]k} \right) \\ &+ \Omega^{-3} (\hat{\rho} + w) (\nabla_{[i} U_{j]} U_k + U_{[j} \nabla_{i]} U_k) \\ &+ \Omega \left((1 + w') \nabla_{[i} \rho U_{j]} U_k + \frac{1}{3} \nabla_{[i} \rho g_{j]k} \right). \end{aligned} \quad (4.12)$$

Its components, given by

$$M_{0b0} = \Omega^{-4} \frac{\hat{\rho} + w}{2} (\Sigma_b - \Omega f_b) - \frac{2 + 3w'}{6} (\Omega \nabla_b \rho + 4\rho \Sigma_b), \quad (4.13)$$

$$M_{ab0} = -\Omega^{-3} (\hat{\rho} + w) \chi_{[ab]} \quad (4.14)$$

$$M_{0bc} = \left\{ -\Omega^{-4} \frac{\hat{\rho} + w}{2} \Sigma_0 + \frac{1}{6} (\Omega \nabla_0 \rho + 4\rho \Sigma_0) \right\} g_{bc} \quad (4.15)$$

$$M_{abc} = \frac{1}{3} \left\{ \Omega \nabla_{[a} \rho g_{b]c} + 4\rho \Sigma_{[a} g_{b]c} \right\}, \quad (4.16)$$

are regular if $4k \in \mathbb{N}_0$.

5. Cauchy Problems

To solve the conformal equations with an asymptotic radiation equation of state we extract from the complete system a symmetric hyperbolic *reduced system* for the unknown

$$\mathbf{Z} = (\Omega, \Sigma_k, s, e^\mu{}_a, \Gamma_a{}^b{}_c, \rho, \chi_{ab}, f_a, L_{jk}, C^i{}_{jkl}),$$

where the Ricci scalar $R = R[g] = 6L_j{}^j$ of $g_{\mu\nu}$ plays the role of a conformal gauge source function that controls implicitly the evolution of the conformal factor. It can be prescribed as an arbitrary function of the coordinates. The present discussion is concerned with the transition from the ‘physical part of the solution’ to scri and beyond. In physical terms this involves a discussion of a domain of infinite temporal extent but in conformal terms it involves only a finite step in the conformal time τ . The specification of $R[g]$ is thus not particularly critical. *It is chosen to be constant.*

Symmetric hyperbolic reduced equations.

The reduced system is essentially built from the equations which contain derivatives in the direction of U . However, sometimes these equations are modified by using constraints, i.e. equations which do not contain derivatives in

the direction of U . For the first six components of \mathbf{Z} we get the system

$$\begin{aligned} e_0(\Omega) &= \Sigma_0, \\ e_0(\Sigma_0) - f^c \Sigma_c &= -\Omega L_{00} - s + \Omega^{-1} \frac{3(\hat{\rho} + w)}{8}, \\ e_0(\Sigma_a) - f_a \Sigma_0 &= -\Omega L_{0a}, \\ \nabla_0 s + \Sigma^i L_{i0} &= \frac{1}{8} \Omega^{-2} \{(1 - 3w')w - 3(\hat{\rho} + w)\} \Sigma_0 \\ &\quad - \frac{1}{24} (1 - 3w') (\Omega^3 \rho + \Omega^{-1} w) \chi_a^a, \end{aligned}$$

where (3.10) resp. (4.3) has been used to replace $\nabla_0 \rho$,

$$\begin{aligned} e^\mu{}_{a,0} &= -\chi_a^b e^\mu{}_b + f_a \delta^\mu{}_0, \\ \Gamma_a{}^b{}_{c,0} &= -f^b \chi_{ac} + \chi_a^b f_c - \chi_a^d \Gamma_d{}^b{}_c + \Omega W^b{}_{c0a} - g^b{}_a L_{0c} + L^b{}_0 g_{ac}, \\ \nabla_0 \rho + (\rho + \Omega^{-4} w) \chi_a^a + \Omega^{-1} (\rho - 3\Omega^{-4} w) \Sigma_0 &= 0, \end{aligned}$$

Adding a suitable contraction of (3.7) to equation (4.5) gives the equation

$$\begin{aligned} e_0(f_a) - w' e_c(\chi_a^c) &= -\chi_{ac} f^c - (1 - 3w') L_{a0} + w' \chi_c^c f_a \\ &\quad + (1 - 3w') (\Omega^{-1} \Sigma_0 f_a + \Omega^{-1} \chi_{ab} \Sigma^b - \Omega^{-2} \Sigma_0 \Sigma_a) \\ &\quad - w'' \frac{\hat{\rho} + w}{w'} (\chi_c^c f_a - 3\Omega^{-1} \Sigma_0 f_a - \Omega^{-1} \chi_c^c \Sigma_a + \Omega^{-2} \Sigma_0 \Sigma_a), \\ &\quad + w' (2\chi_{[c}{}^e \Gamma_{a]e}{}^c - 2f^c \chi_{ca} - 2\Gamma_{[c}{}^e{}_{a]} \chi_e^c - 2L^0{}_a). \end{aligned} \quad (5.1)$$

Adding (4.6) to (3.9) gives

$$\begin{aligned} e_0(\chi_{ba}) - e_a(f_b) &= -f_d \Gamma_a{}^d{}_b + f_a f_b - \chi_a^d \chi_{db} + \Omega W^0{}_{b0a} + L_{ab} + L^0{}_0 g_{ab}, \\ &\quad - 2(\chi_{c[a} \chi_b]{}^c + w' \chi_c^c \chi_{[ab]}) + (1 - 3w') \Omega^{-1} \Sigma_0 \chi_{[ab]}. \end{aligned} \quad (5.2)$$

If equations (5.1) and (5.2) are written as a system for f_a and χ_{bc} with principal part

$$\frac{1}{w'} e_0(f_a) - e_c(\chi_a^c) = \dots, \quad e_0(\chi_{ba}) - e_a(f_b) = \dots$$

they represent, with given right hand sides, a symmetric hyperbolic system.

To derive the reduced systems for the curvature fields we follow the methods of [5] and earlier articles. Equation (4.10) provides a system for the unknowns $L_{0a} = L_{a0}$ and $L_{ab} = L_{ba}$ that contains no derivatives of L_{00} . Where this quantity appears in the equation the contraction $-L_{00} + g^{ab} L_{ab} = L_j{}^j = R/6$ can be used to express it in terms of L_{ab} and the gauge source function $R = R[g] = \text{const}$. The system is given by

$$\nabla_0 L_{0a} - g^{bc} \nabla_b L_{ac} = \Omega (M_{0a0} - g^{bc} M_{bac}), \quad a = 1, 2, 3, \quad (5.3)$$

$$\nabla_0 L_{aa} - \nabla_a L_{0a} = \Sigma_l W^l{}_{a0a} + \Omega M_{0aa}, \quad a = 1, 2, 3, \quad (5.4)$$

$$2\nabla_0 L_{ab} - \nabla_a L_{0b} - \nabla_b L_{0a} = -2\Sigma_l W^l{}_{(ab)0} + 2\Omega M_{0(ab)}, \quad (5.5)$$

$$a, b = 1, 2, 3, \quad a < b.$$

To extract the desired system from (4.11) consider the fields

$$\begin{aligned} h^j{}_k &= g^j{}_k + U^j U_k, \quad l^j{}_k = g^j{}_k + 2U^j U_k, \quad \epsilon_{ijkl} = \epsilon_{[ijkl]}, \\ \epsilon_{jkl} &= U^i \epsilon_{ijkl}, \quad \text{with } \epsilon_{0123} = 1. \end{aligned}$$

The symmetric, trace-free U -electric part w_{jl} and the U -magnetic part w_{jl}^* of $W^i{}_{jkl}$,

$$w_{jl} = W_{ipkq} U^i h^p{}_j U^k h^q{}_l, \quad w_{jl}^* = 1/2 W_{ipmn} \epsilon^{mn}{}_{kq} U^i h^p{}_j U^k h^q{}_l,$$

allow us to represent the rescaled conformal Weyl tensor in form

$$W_{ijkl} = 2(l_{i[k} w_{l]j} - l_{j[k} w_{l]i} - U_{[k} w_{l]p}^* \epsilon^p{}_{ij} - U_{[i} w_{j]p}^* \epsilon^p{}_{kl}).$$

Suitable evolution equations for w_{jl} and w_{jl}^* is given by $\nabla^i W_{i(a|dc|} \epsilon_b)^{dc} = K_{dc(a} \epsilon_b)^{dc}$, $\nabla^i W_{i(a|0|b)} = K_{0(ab)}$ which read in detail

$$\begin{aligned} e_0(w_{ab}^*) - D_d w_{c(a} \epsilon_b)^{dc} &= -\chi_{ed} w_{cf}^* \epsilon_{(a}{}^{fe} \epsilon_b)^{dc} \\ &\quad + (\chi^e{}_{(a} w_{b)e}^* - \chi w_{ab}^*) + f_d w_{c(a} \epsilon_b)^{dc}, \end{aligned} \quad (5.6)$$

and

$$\begin{aligned} e_0(w_{ab}) + D_d w_{c(a} \epsilon_b)^{dc} &= -2\chi w_{ab} + 2\chi^c{}_{(a} w_{b)c} \\ &\quad + \chi_{(a}{}^c w_{b)c} - \chi^{cd} w_{cd} g_{ab} + 2w_{(a}{}^e \epsilon_{b)ed} f^d - M_{0(ab)}, \end{aligned} \quad (5.7)$$

with ‘spatial derivatives’ $D_a w_{bc}^* \equiv e_a(w_{bc}^*) - \Gamma_a{}^e{}_b w_{ec}^* - \Gamma_a{}^e{}_c w_{be}^* = \nabla_a w_{bc}^*$. The components of M_{jkl} on the right hand sides of equations (5.3) – (5.7) are given by

$$\begin{aligned} M_{0a0} - g^{bc} M_{bac} &= \Omega^{-4} \frac{\hat{\rho} + w}{2} (\Sigma_a - \Omega f_a) - \frac{w'}{2} (\Omega \nabla_a \rho + 4\rho \Sigma_a), \\ M_{0ab} &= - \left\{ \Omega^{-4} \frac{\hat{\rho} + w}{2} \Sigma_0 - \frac{1}{6} (\Omega \nabla_0 \rho + 4\rho \Sigma_0) \right\} g_{ab}, \quad M_{dc(a} \epsilon_b)^{dc} = 0, \end{aligned}$$

where here and in (5.3), (5.4), (5.5) equations (4.3) and (4.4) must be used to replace $\nabla_0 \rho$ and $\nabla_a \rho$.

Observing the symmetry of w_{ab} , w_{ab}^* (but ignoring their trace-freeness, which can later be recovered as a consequence of the initial data and the equations above), equations (5.6) and (5.7) can written as system for the unknowns w_{ab} and w_{ab}^* with $a \leq b$, $a, b = 1, 2, 3$. If the equations $w_{ab,0} = \dots$, $w_{ab,0}^* = \dots$ are then written with a factor 2 if $a < b$, the system is seen to be manifestly symmetric hyperbolic.

Together with the previous equations we have obtained now a quasi-linear system for the unknown \mathbf{Z} which can be written in the form $\mathbf{A}^\mu \partial_\mu \mathbf{Z} = \mathbf{B}$ with matrix-valued functions $\mathbf{B} = \mathbf{B}(\mathbf{Z})$ and $\mathbf{A}^\mu = \mathbf{A}^\mu(\mathbf{Z})$ that are symmetric, i.e. ${}^t \mathbf{A}^\mu = \mathbf{A}^\mu$, with \mathbf{A}^0 defining a positive definite bilinear form if the $e^0{}_a$ are not too large.

Cauchy data for the conformal equations.

Solutions that admit smooth conformal extensions at future time-like infinity can be constructed from data for the conformal field equations on a compact Cauchy hypersurface S in the ‘physical domain’, where $\Omega > 0$, or on the compact 3-manifold $S = \mathcal{J}^+ = \{\Omega = 0\}$ that represents future time-like infinity.

Unless the field U is assumed to be orthogonal to S , the data, which must satisfy the constraints induced by the conformal field equations on S , are in general expressed in terms of the unit normal to S and then transformed into a frame e_k with $e_0 = U$. This requires calculations involving the complete system of conformal field equations which are fairly tedious (see [7] where the presence of a boundary requires this) and give little insight. The calculation will be skipped here.

We will first construct *asymptotic end data* on \mathcal{J}^+ with the assumption that U is orthogonal to \mathcal{J}^+ . It is interesting to note that in the case of a pure dust equation of state one is forced into this requirement without forcing the field U to be hypersurface orthogonal in the physical domain (see [5]). In the present case this assumption implies a genuine restriction. The analysis of the asymptotic end data follows essentially the one given in the vacuum case [1] with some modifications if matter fields are involved [4, 5]

To analyse the constraints, i.e. the equations that do not involve derivatives in the direction of U , we assume that $k \geq 1$. The restrictions of the equations to \mathcal{J}^+ are then regular and the expressions containing the quantities w^* , $(w^*)'$ drop out. The unknowns are then determined as follows.

The requirement that U is orthogonal to \mathcal{J}^+ implies

$$e^0_a = 0, \quad \Sigma_a = 0, \quad \chi_{ab} \text{ is the second fundamental form induced on } \mathcal{J}^+.$$

The first structural equation (3.1) implies on \mathcal{J}^+ the first structural equation with respect to the frame e_a and metric h_{ab} induced on \mathcal{J}^+ . The $\Gamma_a^c_b$ are thus the connection coefficients of the covariant derivative operator D induced by the metric h_{ab} in the frame e_a . The second structural equation (3.2) reduces to the relation $R^a_{\ cdb}[h] = 2g^a_{\ [d}L_{b]c} + 2L^a_{\ [d}g_{b]c}$, equivalent to

$$L_{ab}[g] = l_{ab}[h] \quad \text{with} \quad l_{ab}[h] = R_{ab}[h] - \frac{1}{4}R[h]h_{ab}.$$

This gives $L_{00} = g^{ab}L_{ab} - L_j^j = R[h]/4 - R/6$ where $R = R[g]$ represents the conformal gauge source function. Equations (3.11), (3.15), (3.16), (3.17) imply

$$f_a = -\frac{1}{4}\nabla_a\rho, \quad \Sigma^0 = \nu \equiv \sqrt{\lambda/3} > 0, \quad \chi_{ab} = s/\nu g_{ab}, \quad \nabla_a s = -\nu L_{0a},$$

where it is assumed that U is future directed and Ω is decreasing in the direction of U . By a rescaling $g_{\mu\nu} \rightarrow \theta^2 g_{\mu\nu}$ and $\Omega \rightarrow \theta\Omega$, where $\theta > 0$ is smooth with prescribed value and suitable normal derivative on \mathcal{J}^+ , it can be achieved that

$$s = 0 \quad \text{whence} \quad \chi_{ab} = 0, \quad L_{0a} = 0 \quad \text{on } \mathcal{J}^+.$$

Equation (3.18), which reduces to

$$D_c l_{da} \epsilon_b{}^{cd} = \nu w_{ab}^*,$$

were $l_{ab} = l_{ab}[h]$ denotes here the Schouten tensor of h , relates the Cotton tensor of h , given on the left hand side, to the U -magnetic part of $W^i{}_{jkl}$. The constraints induced by (3.19), given by $\nabla_l W^l{}_{0ij} = M_{ij0}$, translate with $M_{ab0} = 0$ and $M_{0b0} = 0$ to the constraint

$$D^a w_{ab} = 0, \quad (5.8)$$

and to $D^a w_{ab}^* = 0$, which is not a constraint but the identity satisfied by the Cotton tensor.

If the data

$$e^\alpha{}_a, \quad \rho, \quad w_{ab},$$

are prescribed on \mathcal{J}^+ so that $h^{\alpha\beta} = \delta^{ab} e^\alpha{}_a e^\beta{}_b$ defines the contravariant version of a Riemannian metric, $\rho > 0$, and $w_{ab} = w_{ba}$, $w_c{}^c = 0$ with w_{ab} satisfying (5.8) with the covariant derivative operator D defined by h , all the unknowns subsumed by \mathbf{Z} can be determined in our gauge.

On the preservation of the constraints.

Smooth data on \mathcal{J}^+ as described above, determine a smooth solution to the symmetric hyperbolic reduced equations that cover a neighbourhood of $\mathcal{J}^+ = \{\tau = 0\}$ with $\Omega > 0$ in the past of \mathcal{J}^+ , where the parameter τ on the flow lines of U is negative, and $\Omega < 0$ in the future of \mathcal{J}^+ where $\tau > 0$ [10]. We can assume it to exist in a range $-\tau_* \leq \tau \leq \tau_*$ with some $\tau_* > 0$ and $\tau = 0$ on \mathcal{J}^+ .

Once this solution has been obtained, it remains to show that the solution to the reduced equations does in fact also satisfy the constraints and thus the complete system of conformal equations. Since the proof follows a standard recipe which has been worked out in detail in earlier articles (see [5], and in particular [3, 8] for a discussion of the Einstein-perfect-fluid equations with a general equation of state), we skip this step.

There remains, however, an open question. Since we assumed U to be orthogonal to \mathcal{J}^+ we should expect U to be hypersurface orthogonal, that is $\chi_{[ab]} = 0$ in the range $-\tau_* \leq \tau \leq \tau_*$. Because $\chi_{[ab]} = 0$ on \mathcal{J}^+ , one could think this to be a consequence of equation (4.6). Since this equation is neither a constraint nor satisfied as part of the reduced system, we use a different argument. The evolution of χ_{ab} is governed by equation (5.2), which implies

$$\begin{aligned} e_0(\chi_{[ba]}) &= D_{[a} f_{b]} + \chi_{[ac]} \sigma_b{}^c - \chi_{[bc]} \sigma_a{}^c - 2w' \chi_c{}^c \chi_{[ab]} \\ &\quad - 2(1 - 3w') \Omega^{-1} \Sigma_0 \chi_{[ab]}, \end{aligned}$$

with $\sigma_{ab} = \chi_{(ac)}$. We formally write here $D_a f_b = e_a(f_b) - f_d \Gamma_a{}^d{}_b$, not implying that D denotes a covariant derivative away from \mathcal{J}^+ . Among the constraints which we can assume to be satisfied there is the relation (3.11) or its more concise version (3.12), i.e.

$$f_a = e_a(f) \quad \text{with} \quad f = F + \log \Omega \quad \text{and} \quad F' = -(\hat{\rho} + w)^{-1} w'. \quad (5.9)$$

Since the connection ∇ is torsion free it follows

$$0 = \nabla_a \nabla_b f - \nabla_b \nabla_a f = 2(D_{[a} f_{b]} - \chi_{[ab]} e_0(f)),$$

where the function

$$e_0(f) = e_0(F + \log \Omega) = -\frac{w'}{\rho + \Omega^{-4} w} \nabla_0 \rho + \Omega^{-1} \frac{\rho + \Omega^{-4} w - 4\rho w'}{\rho + \Omega^{-4} w} \nabla_0 \Omega,$$

is regular if $k \geq 1$. The resulting ODE for $\chi_{[ab]}$ implies that $\chi_{[ab]} = 0$, equation (4.6) is satisfied, and U is hypersurface orthogonal in the range $-\tau_* \leq \tau \leq \tau_*$.

With this we discussed all the ingredients needed to obtain existence results. General results in symmetric hyperbolic systems [10] allow us to draw the following conclusions.

Let the reduced conformal Einstein- λ -perfect-fluid equations with asymptotic radiation equation of state where $k \geq 1$ be given in terms of the gauge discussed above so that $e_0 = U$. Assume the future directed flow vector U to be orthogonal to the compact 3-manifold $\mathcal{J}^+ = \{\Omega = 0\}$. Let on \mathcal{J}^+ be given a minimal end data set consisting of a smooth frame field $e_a = e^\alpha_a \partial_{x^\alpha}$, a positive function ρ , and a symmetric trace-free tensor field w_{ab} that satisfies $D^\alpha w_{ab} = 0$ where D is the covariant derivative operator associated with the metric h that satisfies $h(e_a, e_b) = \delta_{ab}$. Then:

- *As discussed above, a complete set of Cauchy data for the conformal field equations can be calculated from the minimal set if the gauge conditions, the constraints, and the special features of $\mathcal{J}^+ = \{\Omega = 0\}$ are taken into account.*
- *These data determine a smooth solution Ω, g, \dots , to the conformal field equation (unique up to extensions) so that $\Omega < 0$ in the future of \mathcal{J}^+ , $\Omega > 0$ in the past of \mathcal{J}^+ , and $\hat{g} = \Omega^{-2} g$, $\hat{U} = \Omega U$, $\hat{\rho} = \Omega^4 \rho$ satisfy where $\Omega \neq 0$ the Einstein- λ -perfect-fluid equations with asymptotic radiation equation of state. The flow field \hat{U} is hypersurface orthogonal.*
- *Let S be a Cauchy hypersurface in the ‘physical domain’ of this solution where $\Omega > 0$ and denote by Δ the Cauchy data induced by the solution on S . Let Δ' be Cauchy data on S for the same system of equations. If these data are sufficiently close to Δ they develop into a solution which is time-like and null geodesically future complete, admits a smooth conformal boundary \mathcal{J}'^+ representing its future time-like infinity and a smooth conformal extension beyond with $\rho > 0$.*
- *If the value $w^* = 0$ is admitted, the resulting set of solutions, which contain as special cases the FLRM-pure-radiation solutions, develop a smooth conformal boundary in the future.*

Given the data Δ' whence the corresponding flow field U' on S , the field equations allow us to determine the quantity $U'_{[i} \nabla'_j U'_{k]}$ on S . In the last statement it is not required that this quantity vanishes on S . Thus the flow field U' determined by the data Δ' will not necessarily be hypersurface orthogonal and U' need not be orthogonal to \mathcal{J}'^+ .

If Cauchy data are given on \mathcal{J}^+ with a flow vector field U that is not necessarily orthogonal to \mathcal{J}^+ the result stated at the end of the introduction is obtained.

Funding Open Access funding enabled and organized by Projekt DEAL.

Open Access. This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

References

- [1] Friedrich, H.: Existence and structure of past asymptotically simple solution of Einstein's field equations with positive cosmological constant. *J. Geom. Phys.* **3**, 101–117 (1986)
- [2] Friedrich, H.: On the existence of n-geodesically complete or future complete solutions of Einstein's field equations with smooth asymptotic structure. *Commun. Math. Phys.* **107**, 587–609 (1986)
- [3] Friedrich, H.: Evolution equations for gravitating ideal fluid bodies in general relativity. *Phys. Rev. D* **57**, 2317–2322 (1998)
- [4] Friedrich, H.: Smooth non-zero rest-mass evolution across time-like infinity. *Ann. Henri Poincaré* **16**, 2215–2238 (2015)
- [5] Friedrich, H.: Sharp asymptotics for Einstein- λ -dust flows. *Commun. Math. Phys.* (2016). <https://doi.org/10.1007/s00220-016-2716-6>
- [6] Friedrich, H.: Cosmological models with positive λ : hierarchies of future behaviour. *Phil. Trans. R. Soc. A* **382**, 20230044 (2023)
- [7] Friedrich, H., Nagy, G.: The initial boundary value problem for Einstein's vacuum field equations. *Comm. Math. Phys.* **201**, 619–655 (1999)
- [8] Friedrich, H., Rendall, A.: The Cauchy Problem for the Einstein Equations. In: Schmidt, B. (ed.) *Einstein's Field Equations and Their Physical Implications*. Springer, Lecture Notes in Physics, Berlin (2000)
- [9] Hadžić, H., Speck, J.: The global future stability of the FLRW solutions to the Dust–Einstein system with a positive cosmological constant. *J. Hyperbolic Differ. Equ.* **12**, 87 (2015)
- [10] Kato, T.: The Cauchy problem for quasi-linear symmetric hyperbolic systems. *Arch. Ration. Mech. Anal.* **58**, 181–205 (1975)

- [11] Liu, C., Wei, C.: Future stability of the FLRW space-time for a large class of perfect fluids. *Ann. Henri Poincaré* **22**, 715–770 (2021)
- [12] Lübbe, C., Valiente Kroon, J.A.: A conformal approach to the analysis of the non-linear stability of radiation cosmologies. *Ann. Phys.* **328**, 1–25 (2013)
- [13] Olyinyk, T.A.: Future stability of the FLRW fluid solutions in the presence of a positive cosmological constant. *Commun. Math. Phys.* **346**, 293–312 (2016)
- [14] Reula, O.: Exponential decay for small non-linear perturbations of expanding flat homogeneous cosmologies. *Phys. Rev. D* **60**, 083507 (1999)
- [15] Rodnianski, I., Speck, J.: The nonlinear future stability of the FLRW family of solutions to the irrotational Euler–Einstein system with a positive cosmological constant. *J. Eur. Math. Soc.* **15**, 2369–2462 (2013)
- [16] Speck, J.: The nonlinear future stability of the FLRW family of solutions to the Euler–Einstein system with a positive cosmological constant. *Sel. Math. New Ser.* **18**, 633–715 (2012)

Helmut Friedrich
Max-Planck-Institut für Gravitationsphysik
Am Mühlenberg 1
14476 Golm
Germany
e-mail: hcf@aei.mpg.de

Communicated by Mihalis Dafermos.

Received: November 3, 2023.

Accepted: January 7, 2024.