

Academy of Aphasia 2011

Lateralisation of the Arcuate Fasciculus Predicts Aphasia Recovery at 6 Months

Stephanie J. Forkel^{a,*}, Flavio Dell'Acqua^b, Lalit Kalra^c, Steven C. Williams^d, Marco Catani^e

^a *Natbrainlab, Department of Clinical Neurosciences and Neuroimaging, Institute of Psychiatry, King's College London, London, UK*

^b *Natbrainlab, Centre for Neuroimaging Sciences & NIHR Biomedical Research Centre, Institute of Psychiatry, King's College London, London, UK*

^c *Department of Medicine, Guy's, King's and St Thomas's School of Medicine, King's College London, UK*

^d *Department of Neuroimaging, Institute of Psychiatry, King's College London, London, UK*

^e *Natbrainlab, Department of Forensic and Neurodevelopmental Sciences, Institute of Psychiatry, King's College London, London, UK*

Background

Perisylvian language areas are connected through the arcuate fasciculus, a long association pathway that can be visualised using MR-based diffusion tensor imaging (DTI) tractography (Catani et al., 2005). Volumetric measurements of the arcuate fasciculus in healthy subjects show that direct connections between temporal and frontal areas are left-lateralised in most but not all subjects (Catani et al., 2007). In this study, we combined language assessments (Western Aphasia Battery-Revised (WAB-R); Kertesz, 2007) and DTI tractography in patients with post-stroke aphasia to identify anatomical predictors of recovery based on lateralisation patterns (Figure 1.A).

Methods

Nine right-handed subjects (6 males, 3 females) were assessed within 2 weeks of symptom onset using WAB-R and DTI. Follow-up assessment was conducted 6 months thereafter. We scored each patient on the WAB-R to obtain i) the severity of impairment (Aphasia Quotient (AQ) = 0-100, with 0 = very severe, 100 = normal language) and ii) taxonomical classification. Lateralisation of the arcuate was determined by virtual dissections of arcuate fibres in both hemispheres; patients were clustered into three groups accordingly (left-lateralised; left-lateralised/bilateral; bilateral). Statistical analysis was performed in IBM SPSS 19.

Results

At baseline, all subjects scored below cut-off (AQ = 93.8) for clinically significant aphasia. At follow-up, all subjects improved and two recovered. Tractography dissections of the arcuate fasciculus showed that three male patients were extremely left-lateralised with an $AQ_{(\text{median})} = 81$. Another three subjects (2F, 1M) were left-lateralised/bilateral with an $AQ_{(\text{median})} = 87.20$. The remaining subjects (1F, 2M) were bilateral with an $AQ_{(\text{median})} = 95.20$. The latter included the two recovered subjects. Severity of impairment highly correlated with

* Corresponding author.

E-mail address: stephanie.forkel@kcl.ac.uk.

laterality ($r = 0.896, p \leq 0.001$). Comparison of means of language performances across all groups revealed significant differences ($H(2) = 6.489, p < 0.05$) (Figure 1.B). Post hoc pair-wise comparison showed a significant difference between left-lateralised and bilateral subjects ($p < 0.05$).




Figure 1. **A** shows the three laterality groups of the arcuate fasciculus in healthy subjects (left-lateralised, left-lateralised/bilateral, bilateral) depicting the three segments of the arcuate fasciculus for both hemispheres. For both bilateral groups one can appreciate the long segment in the right hemisphere (arrow). **B** shows the 6 months follow-up language performance (AQ, Aphasia Quotient) in relation to arcuate laterality. Dotted line indicates the cut-off for normal language function (AQ=93.8).

Conclusion

Our preliminary results indicate a strong relationship between the degree of lateralisation of the arcuate fasciculus and the language performances at 6 months after stroke-induced aphasia. We showed that the group with the most bilateral representation of the arcuate fasciculus significantly outperforms the left-lateralised group. These results may indicate that the underlying white matter anatomy may be relevant to aphasia recovery post stroke.

References

Catani, M., Allin, M.P., Husain, M., Pugliese, L., Mesulam, M.M., Murray, R.M., Jones, D.K. (2007). Symmetries in human brain language pathways correlate with verbal recall. *Proceedings of the National Academy of Sciences of the USA*, 104 (43), 17163-8.

Catani, M., Jones DK, ffytche, D.H. (2005). Perisylvian language networks of the human brain. *Annals of Neurology*, 57, 8-16.

Kertesz, A. (2007). *Western Aphasia Battery-Revised*. San Antonio, TX: Harcourt Assessment, Inc.