



A Q-Operator for Open Spin Chains II: Boundary Factorization

Alec Cooper¹, Bart Vlaar^{1,2,3} , Robert Weston¹

¹ Department of Mathematics, Heriot-Watt University, Edinburgh EH14 4AS, Scotland, UK.

E-mail: a.cooper@hw.ac.uk

² Beijing Institute for Mathematical Sciences and Applications, 544 Hefangkou Village, Huaibei Town, Huairou District, Beijing 101408, China. E-mail: b.vlaar@bimsa.cn

³ Max Planck Institute for Mathematics, Vivatsgasse 7, 53111 Bonn, Germany. E-mail: r.a.weston@hw.ac.uk

Received: 28 January 2023 / Accepted: 19 February 2024

© The Author(s) 2024

Abstract: One of the features of Baxter’s Q-operators for many closed spin chain models is that all transfer matrices arise as products of two Q-operators with shifts in the spectral parameter. In the representation-theoretical approach to Q-operators, underlying this is a factorization formula for L-operators (solutions of the Yang–Baxter equation associated to particular infinite-dimensional representations). To extend such a formalism to open spin chains, one needs a factorization identity for solutions of the reflection equation (boundary Yang–Baxter equation) associated to these representations. In the case of quantum affine \mathfrak{sl}_2 and diagonal K-matrices, we derive such an identity using the recently formulated theory of universal K-matrices for quantum affine algebras.

Contents

1.	Introduction	
1.1	Background and overview	
1.2	Present work	
1.3	Outline	
2.	Quantum Affine \mathfrak{sl}_2 and its Universal R-Matrix	
2.1	General overview of finite-dimensional R-matrix theory	
2.2	Quantum affine \mathfrak{sl}_2	
2.3	Quantized Kac–Moody algebra	
2.4	Co-opposite Hopf algebra structure	
2.5	Weight modules	
2.6	Quasitriangularity	
2.7	Level-0 representations	
2.8	The action of \mathcal{R} on tensor products of level-0 modules	
2.9	Adjusting the grading	
3.	The Augmented q-Onsager Algebra, its Twist and its Universal K-Matrix	
3.1	The twist map ψ	

- 3.2 The augmented q-Onsager algebra
- 3.3 Universal K-matrix
- 3.4 The action of the universal K-matrix on level-0 representations
- 4. Borel Representations in Terms of the q-Oscillator Algebra
 - 4.1 The infinite-dimensional vector space W
 - 4.2 Diagonal operators from functions and an extended q-oscillator algebra
 - 4.3 Endomorphisms of $W \otimes W$
 - 4.4 The Borel representations
 - 4.5 The $U_q(\widehat{\mathfrak{b}^+})$ -intertwiner \mathcal{O}
 - 4.6 Formalism for $U_q(\widehat{\mathfrak{b}^-})$
- 5. L-Operators and R-Operators
 - 5.1 L-operators for $U_q(\widehat{\mathfrak{b}^+})$ -modules
 - 5.2 Actions of $\mathcal{R}(z)$ on tensor products of infinite-dimensional Borel representations
- 6. K-Operators
 - 6.1 Right K-operators
 - 6.2 Left K-operators
- 7. Fusion Intertwiners Revisited
- 8. Boundary Factorization Identity
- 9. Discussion
- A. Deformed Pochhammer Symbols and Exponentials
 - A.1 Deformed Pochhammer symbols
 - A.2 Deformed exponentials
 - A.3 Deformed exponentials as linear maps
- B. Explicit Expressions for R-Operators
 - B.1 Explicit expression for $R_{\nu\phi}(z)$
 - B.2 The automorphism χ and the q-oscillator subalgebra $\widetilde{\mathcal{A}}$
 - B.3 Explicit expression for $\mathcal{R}_{\mathcal{Q}\bar{\rho}}(z)$
- C. An Alternative Proof of the Main Theorem

1. Introduction

1.1. Background and overview. Baxter first introduced his Q-operator in [Ba72, Ba73] as an auxiliary tool in the derivation of Bethe Equations for the eigenvalues of the 8-vertex model transfer matrix. The key characters in the story are the transfer matrix $\mathcal{T}(z)$ and the Q-operator $\mathcal{Q}(z)$. A detailed description of the essential properties of $\mathcal{T}(z)$ and $\mathcal{Q}(z)$ can be found in [BLZ97] (also see [VW20] and references therein); the key relation that they satisfy that leads directly to the Bethe equations is of the form

$$\mathcal{T}(z)\mathcal{Q}(z) = \alpha_+(z)\mathcal{Q}(qz) + \alpha_-(z)\mathcal{Q}(q^{-1}z), \tag{1.1}$$

where $\alpha_{\pm}(z)$ are meromorphic functions and $q \in \mathbb{C}^{\times}$ is not a root of unity.

In the original papers of Baxter, the operator $\mathcal{Q}(z)$ was constructed by a brilliant but ad hoc argument; the representation-theoretic construction of $\mathcal{Q}(z)$ had to wait more than 20 years until the work of Bazhanov, Lukyanov and Zamolodchikov [BLZ96, BLZ97, BLZ99]. The main idea of the latter approach is to construct both $\mathcal{T}(z)$ and $\mathcal{Q}(z)$ as partial traces over different representations of the universal R-matrix \mathcal{R} of $U_q(\widehat{\mathfrak{sl}}_2)$. The operator $\mathcal{T}(z)$ is a twisted trace over a two-dimensional $U_q(\widehat{\mathfrak{sl}}_2)$ -representation Π_z , and $\mathcal{Q}(z)$ is a similarly twisted trace over an infinite-dimensional $U_q(\widehat{\mathfrak{b}^+})$ -representation ρ_z ,

where $\widehat{\mathfrak{b}}^+$ is the upper Borel subalgebra of $\widehat{\mathfrak{sl}}_2$ (the relevant representations are defined in Sect. 4.4 of the current paper). The relation (1.1) for closed spin chains then follows immediately by considering a short exact sequence (SES) of $U_q(\widehat{\mathfrak{b}}^+)$ -representations with $\Pi_z \otimes \rho_z$ as its ‘middle’ object (cf. [FR99, Lem. 2 (2)]). For an arbitrary untwisted affine Lie algebra $\widehat{\mathfrak{g}}$ with upper Borel subalgebra $\widehat{\mathfrak{b}}^+$, the level-0 representation theory of $U_q(\widehat{\mathfrak{b}}^+)$ was studied in [HJ12]; for the general connection with the theory of Baxter’s Q-operators see [FH15].

As well as this direct SES route to the equation, there is an alternative strategy which we refer to as the *factorization approach*; for closed chains see [BS90, De05, DKK06, De07, BJMST09, BLMS10]. In fact, this approach was the one taken by Bazhanov, Lukyanov and Zamolodchikov. The work that developed this formalism in language most similar to the current paper is [KT14].

In this approach, a second operator $\overline{\mathcal{Q}}(z)$ with similar properties to $\mathcal{Q}(z)$ is introduced as a trace of \mathcal{R} over another infinite-dimensional representation $\overline{\rho}_z$ of $U_q(\widehat{\mathfrak{b}}^+)$. The affinized version ν_z of the $U_q(\mathfrak{sl}_2)$ -Verma module is also considered as well as another infinite-dimensional filtered $U_q(\widehat{\mathfrak{b}}^+)$ -module ϕ_z ; these two representations depend on a complex parameter μ . The key connection between all representations is given by Theorem 4.4, which expresses the fact that particular pairwise tensor products are isomorphic as $U_q(\widehat{\mathfrak{b}}^+)$ -modules by means of an explicit intertwiner \mathcal{O} (defined in Sect. 4.5 of the current paper). At the level of the L-operators this implies

$$\mathcal{O}_{12} \mathcal{L}_\rho(q^{-\mu/2} z)_{13} \mathcal{L}_{\overline{\rho}}(q^{\mu/2} z)_{23} = \mathcal{L}_\nu(z)_{13} \mathcal{L}_\phi(z)_{23} \mathcal{O}_{12}, \tag{1.2}$$

(see Theorem 5.2 of the current paper), which is referred to as *factorization* of the Verma module L-operator $\mathcal{L}_\nu(z)$ in terms of the L-operators $\mathcal{L}_\rho(z)$ and $\mathcal{L}_{\overline{\rho}}(z)$ which are used to define $\mathcal{Q}(z)$, $\overline{\mathcal{Q}}(z)$ (the transfer matrix corresponding to the additional operator $\mathcal{L}_\phi(z)$ is trivial).

Defining $\mathcal{T}_\mu(z)$ to be the transfer matrix that is the trace over the μ -dependent representation ν_z of \mathcal{R} in the first space, Theorem 5.2 yields a relation of the following form:

$$\mathcal{T}_\mu(z) \propto \mathcal{Q}(zq^{-\mu/2}) \overline{\mathcal{Q}}(zq^{\mu/2}). \tag{1.3}$$

The SES associated with ν_z in the case μ is an integer then leads to the key relation (1.1).

1.2. Present work. In the current work we are interested in an analogue of (1.2) for open chains, setting out an approach to Q-operators which complements the SES approach of [VW20].

The problem of Q-operators for open XXX chains with diagonal boundaries was discussed in [BT18] and in [Ts21]. The XXX version of this problem was solved already in [FS15]. Earlier, Baxter TQ-relations with more general boundary conditions were found in [YNZ06] (XXX) and [YZ06] (XYZ) by spin- j transfer matrix asymptotics.

Our main result is the following analogue of Theorem 5.2, which we call the *boundary factorization identity*. Its existence answers in the positive a question raised in [BT18, Sec. 5]:

$$\mathcal{K}_\nu(z)_1 \mathcal{R}_{\nu\phi}(z^2) \mathcal{K}_\phi(z)_2 \mathcal{O} = \mathcal{O} \mathcal{K}_\rho(q^{-\mu/2} z)_1 \mathcal{R}_{\overline{\rho}}(z^2) \mathcal{K}_{\overline{\rho}}(q^{\mu/2} z)_2 \tag{1.4}$$

where z is a formal parameter (which can be specialized to generic complex numbers). The precise statement is given in Theorem 8.1. This formula involves the actions of the universal R-matrix of $U_q(\widehat{\mathfrak{sl}}_2)$ in tensor products of the various infinite-dimensional representations introduced. In addition, the various K-operators are diagonal solutions of reflection equations (boundary Yang–Baxter equations) [Ch84, Sk88]. They arise as actions of the universal K-matrix associated to the augmented q-Onsager algebra, a particular coideal subalgebra of $U_q(\widehat{\mathfrak{sl}}_2)$, which featured also in e.g. [BB13, RSV15, BT18, VW20]. More precisely, diagonal solutions of the reflection equation with a free parameter, considered by Sklyanin in his 2-boundary version of the algebraic Bethe ansatz in [Sk88], are intertwiners for this algebra.

Equation (1.4) has a natural diagrammatic formulation, see Sect. 8. In a subsequent paper the authors will explain how (1.4) yields relations analogous to (1.3) and hence (1.1) for open chains.

The proof of (1.4) and of the well-definedness of the various K-operators is an application of the universal K-matrix formalism developed in [AV22a, AV22b] which is built on the earlier works [BW18, BK19]. More precisely, it relies on an extension of the theory of K-matrices for finite-dimensional representations of quantum affine algebras in [AV22b] to level-0 representations of $U_q(\widehat{\mathfrak{b}}^+)$, which we discuss in Sect. 3. The key point is that, for the special case of the augmented q-Onsager algebra, there exists a universal element \mathcal{K} , centralizing the augmented q-Onsager algebra up to a twist, with three desirable properties.

- (i) The element \mathcal{K} lies in (a completion of) the Borel subalgebra $U_q(\widehat{\mathfrak{b}}^+)$, so that the resulting family of linear maps is itself compatible with $U_q(\widehat{\mathfrak{b}}^+)$ -intertwiners (which play an essential role in the algebraic theory of Baxter Q-operators).
- (ii) The coproduct of \mathcal{K} is of a particularly simple form, which is relevant for the proof of the boundary factorization identity.
- (iii) The linear operators accomplishing the action of \mathcal{K} in level-0 representations satisfy the untwisted reflection equation.

Thus we obtain the factorization identity (1.4) as a natural consequence of the representation theory of $U_q(\widehat{\mathfrak{sl}}_2)$. The main benefit of this universal approach is that laborious linear-algebraic computations are avoided; in particular, we not even need explicit expressions for the various factors. Nevertheless, we do provide these explicit expressions, as we expect them to be useful in further work in this direction. We also give an alternative computational proof of (1.4), to illustrate the power of the universal approach.

This is a ‘boundary counterpart’ to the level-0 theory of the universal R-matrix, which we also include for reference. We do this in Sect. 2, staying close to the original work by Drinfeld and Jimbo [Dr85, Dr86, Ji86a, Ji86b]. In particular, Theorem 2.4 states that the grading-shifted universal R-matrix has a well-defined action as a linear-operator-valued formal power series on any tensor product of level-0 representations of $U_q(\widehat{\mathfrak{b}}^+)$ and $U_q(\widehat{\mathfrak{b}}^-)$ (including finite-dimensional representations). Often this well-definedness is tacitly assumed, see e.g. [VW20, Sec. 2.3]. Alternatively, it follows from the Khoroshkin-Tolstoy factorization [KT92] of the universal R-matrix, see also [BGKNR10, BGKNR13, BGKNR14]; however we are unaware of such a factorization for the universal K-matrix.

1.3. Outline. In Sect. 2 we study the action of the universal R-matrix of quantum affine \mathfrak{sl}_2 on tensor products of level-0 representations of Borel subalgebras. Section 3 is a ‘boundary counterpart’ to Sect. 2, where we consider the augmented q-Onsager algebra.

We show that its *(semi-)standard* universal K-matrix, see [AV22a, AV22b], has a well-defined action on level-0 representations of $U_q(\widehat{\mathfrak{b}^+})$, see Theorem 3.6, and, with a simple correction, satisfies the above three desirable properties.

In Sect. 4 we discuss the relevant representations of $U_q(\widehat{\mathfrak{b}^+})$ in terms of (an extension of) the q-oscillator algebra, as well as the $U_q(\widehat{\mathfrak{b}^+})$ -intertwiner \mathcal{O} . Various solutions of Yang–Baxter equations are obtained in Sect. 5 as actions of the universal R-matrix in tensor products of Borel representations. Similarly, in Sect. 6 we introduce solutions of the reflection equation as actions of the universal K-matrix in Borel representations.

We revisit the SES approach to Baxter’s Q-operators for the open XXZ spin chain in light of the universal K-matrix formalism in Sect. 7. Next, in Sect. 8 we give a diagrammatic motivation of the boundary factorization identity (1.4) for the open XXZ spin chain, and provide a short proof using the level-0 theory developed in Sect. 3. Finally, in Sect. 9 we summarize the main results and point out future work.

Some supplementary material is given in appendices. Namely, Appendix A provides some background material on deformed Pochhammer symbols and exponentials. Moreover, Appendix B contains derivations of the explicit expressions of the two R-operators appearing in (1.4). In Appendix C we provide a computational alternative proof of the boundary factorization identity (1.4), relying on the explicit expressions of all involved factors. The key tool of this proof is provided by Lemma C.1, which consists in two product formulas involving deformed Pochhammer symbols and exponentials. We emphasize that the main text and its results do not rely on Appendices B and C.

2. Quantum Affine \mathfrak{sl}_2 and its Universal R-Matrix

In this section we study the action of the universal R-matrix of the quasitriangular Hopf algebra quantum affine \mathfrak{sl}_2 on tensor products of level-0 representations (including infinite-dimensional representations) of the Borel subalgebras. We give a basic survey of the algebras involved, the representations and the quasitriangular structure and show that the universal R-matrix has a well-defined action on tensor products of all level-0 representations of the Borel subalgebras.

2.1. General overview of finite-dimensional R-matrix theory. To formulate a quantum integrable system in terms of a transfer matrix built out of R-matrices, one needs finite-dimensional representations of a suitable quasitriangular Hopf algebra. To get trigonometric R-matrices, one can proceed as follows.

Let \mathfrak{g} be a finite-dimensional simple Lie algebra and note that the untwisted loop algebra $L\mathfrak{g} = \mathfrak{g} \otimes \mathbb{C}[t, t^{-1}]$ has a central extension $\widehat{\mathfrak{g}} = L\mathfrak{g} \oplus \mathbb{C}c$. In turn, this can be extended to $\widetilde{\mathfrak{g}} = \widehat{\mathfrak{g}} \oplus \mathbb{C}d$ where d satisfies $[d, \cdot] = t \frac{d}{dt}$. For a fixed Cartan subalgebra $\mathfrak{h} \subset \mathfrak{g}$ we define

$$\widehat{\mathfrak{h}} := \mathfrak{h} \oplus \mathbb{C}c, \quad \widetilde{\mathfrak{h}} := \widehat{\mathfrak{h}} \oplus \mathbb{C}d.$$

The Lie algebra $\widetilde{\mathfrak{g}}$ is a Kac–Moody algebra and hence has a non-degenerate bilinear form (\cdot, \cdot) , which restricts to a non-degenerate bilinear form on $\widetilde{\mathfrak{h}}$. See e.g. [Ka90] for more detail.

The universal enveloping algebras $U(\widehat{\mathfrak{g}})$ and $U(\widetilde{\mathfrak{g}})$ can be q-deformed, yielding non-cocommutative Hopf algebras (Drinfeld–Jimbo quantum groups) $U_q(\widehat{\mathfrak{g}})$ and $U_q(\widetilde{\mathfrak{g}})$, see e.g. [Dr85, Dr86, Ji86a, KT92, Lu94]. The nondegenerate bilinear form (\cdot, \cdot) lifts to $U_q(\widetilde{\mathfrak{g}})$

inducing a pairing between the q -deformed Borel subalgebras and hence a quasitriangular structure. On the other hand, the subalgebra $U_q(\widehat{\mathfrak{g}})$ has a rich finite-dimensional representation theory, see e.g. [CP94, CP95, Ch02, HJ12]. The grading-shifted universal R -matrix has a well-defined action on tensor products of finite-dimensional representations of $U_q(\widehat{\mathfrak{g}})$ as a formal power series, see e.g. [Dr86, FR92, KS95, EM03, He19]). We now discuss the natural extension of this theory to level-0 representations of Borel subalgebras, including various infinite-dimensional representations. We will restrict to the case $\mathfrak{g} = \mathfrak{sl}_2$ (but the theory generalizes to any quantum untwisted affine algebra).

2.2. *Quantum affine \mathfrak{sl}_2 .* Denoting the canonical Cartan generator of \mathfrak{sl}_2 by h_1 , $\widehat{\mathfrak{h}}$ is spanned by $h_0 = c - h_1$ and h_1 . The bilinear form on $\widehat{\mathfrak{h}}$ is defined by

$$(h_0, h_0) = (h_1, h_1) = -(h_0, h_1) = 2, \quad (h_0, d) = 1, \quad (h_1, d) = (d, d) = 0.$$

Fix $\epsilon \in \mathbb{C}$ such that $q = \exp(\epsilon)$ is not a root of unity. For all $\mu \in \mathbb{C}$ we will denote $\exp(\epsilon\mu)$ by q^μ . First, we define $U_q(\mathfrak{g})$ as the algebra generated over \mathbb{C} by e, f and invertible k subject to the relations

$$ke = q^2ek, \quad kf = q^{-2}fk, \quad [e, f] = \frac{k - k^{-1}}{q - q^{-1}}. \tag{2.1}$$

The following assignments determine a coproduct $\Delta : U_q(\mathfrak{g}) \rightarrow U_q(\mathfrak{g}) \otimes U_q(\mathfrak{g})$:

$$\Delta(e) = e \otimes 1 + k \otimes e, \quad \Delta(f) = f \otimes k^{-1} + 1 \otimes f, \quad \Delta(k^{\pm 1}) = k^{\pm 1} \otimes k^{\pm 1}. \tag{2.2}$$

It uniquely extends to a Hopf algebra structure on $U_q(\mathfrak{g})$. Now the main algebra of interest, $U_q(\widehat{\mathfrak{g}})$, arises as follows.

Definition 2.1 (*Quantum affine \mathfrak{sl}_2*). We denote by $U_q(\widehat{\mathfrak{g}})$ the Hopf algebra generated by two triples $\{e_i, f_i, k_i\}$ ($i \in \{0, 1\}$), such that:

(i) the following assignments for $i \in \{0, 1\}$ define Hopf algebra embeddings from $U_q(\mathfrak{g})$ to $U_q(\widehat{\mathfrak{g}})$:

$$e \mapsto e_i, \quad f \mapsto f_i, \quad k \mapsto k_i; \tag{2.3}$$

(ii) the following cross relations are satisfied:

$$k_i k_j = k_j k_i, \quad k_i e_j = q^{-2} e_j k_i, \quad k_i f_j = q^2 f_j k_i, \quad [e_i, f_j] = 0, \tag{2.4}$$

$$[e_i, [e_i, [e_i, e_j]_{q^2}]_1]_{q^{-2}} = [f_i, [f_i, [f_i, f_j]_{q^2}]_1]_{q^{-2}} = 0, \tag{2.5}$$

for $i \neq j$, where we have introduced the notation $[x, y]_p := xy - pyx$. ∅

Consider the affine Cartan subalgebra $\widehat{\mathfrak{h}} = \mathbb{C}h_0 \oplus \mathbb{C}h_1$. Note that its q -deformation $U_q(\widehat{\mathfrak{h}}) = \langle k_0^{\pm 1}, k_1^{\pm 1} \rangle$ is isomorphic to the group algebra of the affine co-root lattice

$$\widehat{Q}^\vee = \mathbb{Z}h_0 + \mathbb{Z}h_1 \subset \widehat{\mathfrak{h}}. \tag{2.6}$$

The nontrivial diagram automorphism Φ of the affine Dynkin diagram, i.e. the nontrivial permutation of the index set $\{0, 1\}$, lifts to a linear automorphism Φ of $\widehat{\mathfrak{h}}$ which preserves the lattice \widehat{Q}^\vee . Accordingly, it also lifts to an involutive Hopf algebra automorphism of $U_q(\widehat{\mathfrak{g}})$, also denoted Φ , via the assignments

$$\Phi(e_i) = e_{\Phi(i)}, \quad \Phi(f_i) = f_{\Phi(i)}, \quad \Phi(k_i^{\pm 1}) = k_{\Phi(i)}^{\mp 1} \quad \text{for } i \in \{0, 1\}. \tag{2.7}$$

2.3. Quantized Kac–Moody algebra. To define the quantized Kac–Moody algebra $U_q(\widehat{\mathfrak{g}})$, one chooses an extension \widetilde{Q}^\vee of \widehat{Q}^\vee (a lattice of rank 3 contained in \mathfrak{h}) preserved by Φ .

Remark 2.2. The standard extension of the affine co-root lattice $\mathbb{Z}h_0 + \mathbb{Z}h_1 + \mathbb{Z}d$ is not so convenient for us, mainly in view of the construction of the universal K-matrix in Sect. 3.3. Namely, extensions of Φ to $\widehat{\mathfrak{h}}$ which are compatible with the bilinear form on $\widehat{\mathfrak{h}}$ do not preserve this lattice, see also [Ko14, Sec. 2.6] and [AV22a, Sec. 3.14]. \square

The most convenient choice is to use the *principal grading* and set

$$d_{\text{pr}} := -\frac{1}{8}h_0 + \frac{3}{8}h_1 + 2d \in \mathfrak{h}, \tag{2.8}$$

so that

$$(d_{\text{pr}}, h_0) = (d_{\text{pr}}, h_1) = 1, \quad (d_{\text{pr}}, d_{\text{pr}}) = 0.$$

Now we set $\Phi(d_{\text{pr}}) = d_{\text{pr}}$ and obtain a linear automorphism Φ of $\widetilde{\mathfrak{h}}$ preserving the lattice

$$\widetilde{Q}^\vee := \mathbb{Z}h_0 + \mathbb{Z}h_1 + \mathbb{Z}d_{\text{pr}}.$$

The corresponding dual map on $\widetilde{\mathfrak{h}}^*$, also denoted by Φ , preserves the extended affine weight lattice

$$\widetilde{P} = \{\lambda \in \widetilde{\mathfrak{h}}^* \mid \lambda(\widetilde{Q}^\vee) \subseteq \mathbb{Z}\}. \tag{2.9}$$

Accordingly, we define $U_q(\widetilde{\mathfrak{g}})$ as the Hopf algebra obtained by extending $U_q(\widehat{\mathfrak{g}})$ by a group-like element¹ g satisfying

$$ge_i = qe_i g, \quad gf_i = q^{-1}f_i g, \quad gk_i = k_i g. \tag{2.10}$$

Hence, the assignment $\Phi(g) = g$ together with (2.7) defines an involutive Hopf algebra automorphism of $U_q(\widetilde{\mathfrak{g}})$.

2.4. Co-opposite Hopf algebra structure. For any \mathbb{C} -algebra A , denote by σ the algebra automorphism of $A \otimes A$ which sends $a \otimes a'$ to $a' \otimes a$ for all $a, a' \in A$. If $X \in A \otimes A$ we will also write X_{21} for $\sigma(X)$.

If A is a bialgebra with coproduct Δ , the *co-opposite bialgebra*, denoted A^{cop} , is the bialgebra with the same underlying algebra structure and counit as A but with Δ replaced by

$$\Delta^{\text{op}} := \sigma \circ \Delta \tag{2.11}$$

(if A is a Hopf algebra with invertible antipode S , then A^{cop} is also a Hopf algebra with antipode S^{-1}). The assignments

$$\omega(e_i) = f_i, \quad \omega(f_i) = e_i, \quad \omega(k_i^{\pm 1}) = k_i^{\mp 1} \quad \text{for } i \in \{0, 1\}, \quad \omega(g) = g^{-1} \tag{2.12}$$

define a bialgebra isomorphism from $U_q(\widetilde{\mathfrak{g}})$ to $U_q(\widetilde{\mathfrak{g}})^{\text{cop}}$ (in particular, $(\omega \otimes \omega) \circ \Delta = \Delta^{\text{op}} \circ \omega$) which commutes with Φ .

¹ It is equal to $\exp(\epsilon d_{\text{pr}})$ if we define $U_q(\widetilde{\mathfrak{g}})$ as a topological Hopf algebra over $\mathbb{C}[[\epsilon]]$.

2.5. *Weight modules.* We review some basic representation-theoretic notions for $U_q(\tilde{\mathfrak{g}})$ by means of which its universal R-matrix can be described. Consider the commutative subalgebra

$$U_q(\tilde{\mathfrak{h}}) = \langle k_0^{\pm 1}, k_1^{\pm 1}, g^{\pm 1} \rangle \subset U_q(\tilde{\mathfrak{g}}). \tag{2.13}$$

Call a $U_q(\tilde{\mathfrak{g}})$ -module M a $U_q(\tilde{\mathfrak{h}})$ -weight module if

$$M = \bigoplus_{\lambda \in \tilde{P}} M_\lambda, \quad M_\lambda = \{m \in M \mid k_i \cdot m = q^{\lambda(h_i)} m \text{ for } i \in \{0, 1\}, g \cdot m = q^{\lambda(d_{\text{pr}})} m\}.$$

Elements of M_λ are said to have weight λ . The adjoint action of $U_q(\tilde{\mathfrak{h}})$ (with its generators acting by conjugation) endows $U_q(\tilde{\mathfrak{g}})$ itself with a $U_q(\tilde{\mathfrak{h}})$ -weight module structure, with elements of $U_q(\tilde{\mathfrak{h}})$ of weight 0. More precisely, the weights of $U_q(\tilde{\mathfrak{g}})$ are given by the affine root lattice

$$\widehat{Q} := \mathbb{Z}\alpha_0 + \mathbb{Z}\alpha_1 \subset \tilde{P}$$

(e_i has weight α_i , f_i has weight $-\alpha_i$). The adjoint action of $U_q(\tilde{\mathfrak{h}})$ preserves the subalgebras

$$U_q(\widehat{\mathfrak{n}}^+) = \langle e_0, e_1 \rangle, \quad U_q(\widehat{\mathfrak{n}}^-) = \langle f_0, f_1 \rangle \tag{2.14}$$

and the corresponding weights are given by the monoids $\pm \widehat{Q}^+$ respectively, where $\widehat{Q}^+ := \mathbb{Z}_{\geq 0}\alpha_0 + \mathbb{Z}_{\geq 0}\alpha_1$.

2.6. *Quasitriangularity.* The universal R-matrix for $U_q(\tilde{\mathfrak{g}})$ is an element of a completion of $U_q(\tilde{\mathfrak{g}}) \otimes U_q(\tilde{\mathfrak{g}})$ satisfying

$$\mathcal{R}\Delta(u) = \Delta^{\text{op}}(u)\mathcal{R} \quad \text{for all } u \in U_q(\tilde{\mathfrak{g}}), \tag{2.15}$$

$$(\Delta \otimes \text{id})(\mathcal{R}) = \mathcal{R}_{13}\mathcal{R}_{23}, \quad (\text{id} \otimes \Delta)(\mathcal{R}) = \mathcal{R}_{13}\mathcal{R}_{12} \tag{2.16}$$

and hence

$$\mathcal{R}_{12}\mathcal{R}_{13}\mathcal{R}_{23} = \mathcal{R}_{23}\mathcal{R}_{13}\mathcal{R}_{12}. \tag{2.17}$$

Consider the quantum analogues of the Borel subalgebras, which are the Hopf subalgebras

$$U_q(\tilde{\mathfrak{b}}^\pm) = \langle U_q(\tilde{\mathfrak{h}}), U_q(\widehat{\mathfrak{n}}^\pm) \rangle.$$

The element \mathcal{R} arises as the canonical element of the bialgebra pairing between $U_q(\tilde{\mathfrak{b}}^+)$ and the algebra $U_q(\tilde{\mathfrak{b}}^-)^{\text{op}}$ (the bialgebra isomorphic as a coalgebra to $U_q(\tilde{\mathfrak{b}}^-)$ but with the opposite multiplication), see [Dr85, Lu94]. In particular, \mathcal{R} lies in a completion of $U_q(\tilde{\mathfrak{b}}^+) \otimes U_q(\tilde{\mathfrak{b}}^-)$. Further, invariance properties of the bialgebra pairing imply

$$(\omega \otimes \omega)(\mathcal{R}) = \mathcal{R}_{21}, \tag{2.18}$$

$$(\Phi \otimes \Phi)(\mathcal{R}) = \mathcal{R}. \tag{2.19}$$

Also, this pairing has a non-degenerate restriction to $U_q(\widehat{\mathfrak{n}^+})_\lambda \times U_q(\widehat{\mathfrak{n}^-})_{-\lambda}$ for all $\lambda \in \widehat{Q}^+$; denote the canonical element of this restricted pairing by Θ_λ . With our choice of the coproduct we have

$$\mathcal{R} = \Theta^{-1} \cdot \kappa^{-1}, \quad \Theta = \sum_{\lambda \in \widehat{Q}^+} \Theta_\lambda, \tag{2.20}$$

A priori, Θ acts naturally on $U_q(\widehat{\mathfrak{g}})$ -modules with a locally finite action of $U_q(\widehat{\mathfrak{n}^+})$ or $U_q(\widehat{\mathfrak{n}^-})$. We briefly explain one possible definition² of the element κ . The non-degenerate bilinear form (\cdot, \cdot) on $\widetilde{\mathfrak{h}}$ induces one on $\widetilde{\mathfrak{h}}^*$, which we denote by the same symbol. If M, M' are $U_q(\mathfrak{h})$ -weight modules we define a linear map $\kappa_M : M \otimes M' \rightarrow M \otimes M'$ by stipulating that it acts on $M_\lambda \otimes M'_{\lambda'}$ ($\lambda, \lambda' \in \widetilde{P}$) as multiplication by $q^{(\lambda, \lambda')}$. The family of these maps κ_M , where M runs through all $U_q(\widetilde{\mathfrak{h}})$ -weight modules, is compatible with $U_q(\widetilde{\mathfrak{h}})$ -intertwiners. Hence it gives rise to a well-defined weight-0 element κ of the corresponding completion of $U_q(\widehat{\mathfrak{g}}) \otimes U_q(\widehat{\mathfrak{g}})$ which we call here *weight completion*. Similarly, we will define weight-0 elements of the weight completion of $U_q(\widehat{\mathfrak{g}})$ itself using functions from \widetilde{P} to \mathbb{C} . See also [AV22a, Sec. 4.8] for more detail.

2.7. *Level-0 representations.* Consider the following subalgebras of $U_q(\widehat{\mathfrak{g}})$:

$$U_q(\widehat{\mathfrak{b}^\pm}) = \langle U_q(\widehat{\mathfrak{h}}), U_q(\widehat{\mathfrak{n}^\pm}) \rangle = U_q(\widetilde{\mathfrak{b}^\pm}) \cap U_q(\widehat{\mathfrak{g}}). \tag{2.21}$$

Then $U_q(\widehat{\mathfrak{b}^+})$ is isomorphic to the algebra with generators e_i, k_i ($i \in \{0, 1\}$) subject to those relations in Definition 2.1 which do not involve the f_i (the proof of e.g. [Ja96, Thm. 4.21] applies). We say that a $U_q(\widehat{\mathfrak{b}^+})$ -module V is *level-0* if it decomposes as

$$V = \bigoplus_{c \in \mathbb{C}^\times} V(c), \quad V(c) = \{v \in V \mid k_0 \cdot v = c^{-1}v, \quad k_1 \cdot v = cv\} \tag{2.22}$$

with each *weight subspace* $V(c)$ finite-dimensional. Note that the class of finitely generated level-0 modules is closed under tensor products. By the $U_q(\widehat{\mathfrak{g}})$ -relations we have $e_0 \cdot V(c) \subseteq V(q^{-2}c)$, $e_1 \cdot V(c) \subseteq V(q^2c)$. It is convenient to call the subset $\{c \in \mathbb{C}^\times \mid \dim(V(c)) \neq 0\}$ the *support* of V . If V is a finite-dimensional $U_q(\widehat{\mathfrak{g}})$ -module then it is level-0 with support contained in $\pm q^{\mathbb{Z}}$, see e.g. [CP95, Prop. 12.2.3].

Remark 2.3. The $U_q(\widehat{\mathfrak{g}})$ -action on a nontrivial finite-dimensional module does not extend to a $U_q(\widehat{\mathfrak{g}})$ -action. By [HJ12, Prop. 3.5], this is a special case of the following observation. If V is an irreducible level-0 $U_q(\widehat{\mathfrak{b}^+})$ -module with $\dim(V) > 1$, then the $U_q(\widehat{\mathfrak{b}^+})$ -action does not extend to a $U_q(\widetilde{\mathfrak{b}^+})$ -action. To see this, choose distinct $c, c' \in \mathbb{C}^\times$ in the support of V . By irreducibility, for any nonzero $v \in V(c), v' \in V(c')$ there exist $x, x' \in U_q(\widehat{\mathfrak{b}^+})$ such that $x \cdot v = v', x' \cdot v' = v$. Without loss of generality, we may assume both x and x' have no term in $U_q(\widehat{\mathfrak{h}})$. Hence $x'x$ is not a scalar. For any nonzero $v \in V(c)$, since the action of g preserves $V(c)$, applying g to $(x'x) \cdot v = v$ now yields a contradiction with (2.10). ∅

Analogous definitions and comments can be made for $U_q(\widehat{\mathfrak{b}^-})$ -modules.

² Note that in the topological Hopf algebra setting one simply has $\kappa = q^{c \otimes d + d \otimes c + h_1 \otimes h_1 / 2}$.

2.8. *The action of \mathcal{R} on tensor products of level-0 modules.* We wish to connect the quasitriangular structure of $U_q(\widehat{\mathfrak{g}})$ with the level-0 representation theory of $U_q(\widehat{\mathfrak{g}})$, i.e. let the universal R-matrix of $U_q(\widehat{\mathfrak{g}})$ act on tensor products of level-0 modules. To do this, we follow the ideas from [Dr86, Sec. 13] (also see [FR92, Sec. 4], [He19, Sec. 1]). If we write the action of k_1 on an arbitrary level-0 module V as $\exp(\epsilon H_V)$, then note that the factor κ naturally acts on tensor products $V \otimes V'$ of level-0 modules as $\exp(\epsilon H_V \otimes H_{V'}/2)$.

To let Θ act on such tensor products, we extend the field of scalars \mathbb{C} to the Laurent polynomial ring $\mathbb{C}[z, z^{-1}]$, where z is a formal parameter. The action of Θ is particularly well-behaved if we use the principal grading. That is, we define a Hopf algebra automorphism Σ_z of $U_q(\widehat{\mathfrak{g}})[z, z^{-1}]$ such that

$$\Sigma_z(e_i) = ze_i, \quad \Sigma_z(f_i) = z^{-1}f_i, \quad \Sigma_z|_{U_q(\widehat{\mathfrak{h}})} = \text{id}. \tag{2.23}$$

Straightforwardly one sees that

$$\omega \circ \Sigma_z = \Sigma_{z^{-1}} \circ \omega, \tag{2.24}$$

$$\Phi \circ \Sigma_z = \Sigma_z \circ \Phi. \tag{2.25}$$

Let the height function $\text{ht} : \widehat{Q} \rightarrow \mathbb{Z}$ be defined by $\text{ht}(m_0\alpha_0 + m_1\alpha_1) = m_0 + m_1$ for all $m_0, m_1 \in \mathbb{Z}$ and note that the number of elements of \widehat{Q}^+ of given height is finite. The key observation is that

$$(\Sigma_z \otimes \text{id})(\Theta) = (\text{id} \otimes \Sigma_{z^{-1}})(\Theta) = \sum_{r \geq 0} z^r \sum_{\lambda \in \widehat{Q}^+, \text{ht}(\lambda)=r} \Theta_\lambda, \tag{2.26}$$

is a formal power series in z whose coefficients are finite sums and hence lie in $U_q(\widehat{\mathfrak{n}^+}) \otimes U_q(\widehat{\mathfrak{n}^-})$. Hence $(\Sigma_z \otimes \text{id})(\Theta) = (\text{id} \otimes \Sigma_{z^{-1}})(\Theta)$ has a well-defined action as a linear-operator-valued formal power series on a tensor product of any $U_q(\widehat{\mathfrak{n}^+})$ -representation with any $U_q(\widehat{\mathfrak{n}^-})$ -representation. Consider now the *grading-shifted universal R-matrix*:

$$\mathcal{R}(z) := (\Sigma_z \otimes \text{id})(\mathcal{R}) = (\text{id} \otimes \Sigma_{z^{-1}})(\mathcal{R}). \tag{2.27}$$

Note that by applying $\Sigma_z \otimes \text{id}$ to (2.15) we deduce that $\mathcal{R}(z)$ commutes with $\Delta(k_1) = \Delta^{\text{op}}(k_1) = k_1 \otimes k_1$. We collect the results obtained thus far, writing

$$M[[z]] = M \otimes \mathbb{C}[[z]]$$

for any complex vector space M (in particular, any complex unital associative algebra).

Theorem 2.4. *Consider a pair of level-0 representations $\pi^\pm : U_q(\widehat{\mathfrak{b}^\pm}) \rightarrow \text{End}(V^\pm)$. Then³*

$$\mathcal{R}_{\pi^+\pi^-}(z) := (\pi^+ \otimes \pi^-)(\mathcal{R}(z)) \in \text{End}(V^+ \otimes V^-)[[z]] \tag{2.28}$$

is well-defined and commutes with $(\pi^+ \otimes \pi^-)(\Delta(k_1)) = \pi^+(k_1) \otimes \pi^-(k_1)$.

³ Note that in Sect. 5 we will use the notation $\mathcal{R}_{\pi^+\pi^-}(z)$ for a rescaled version of the action of the grading-shifted universal R-matrix.

From now on we will use the standard convention that if π is any level-0 representation then the corresponding grading-shifted representation is denoted by a subscript z :

$$\pi_z := \pi \circ \Sigma_z. \tag{2.29}$$

Hence we may write

$$\mathcal{R}_{\pi^+\pi^-}(z) = (\pi_z^+ \otimes \pi^-)(\mathcal{R}) = (\pi^+ \otimes \pi_{1/z}^-)(\mathcal{R}).$$

Consider two indeterminates z_1, z_2 . Applying, say, $\Sigma_{z_1} \otimes \text{id} \otimes \Sigma_{1/z_2}$, to (2.17), we obtain a $\mathbb{C}[[z_1, z_2]]$ -version of the universal Yang–Baxter equation which can be evaluated on suitable triple tensor products.

Proposition 2.5. *If $\pi^+ : U_q(\widehat{\mathfrak{b}}^+) \rightarrow \text{End}(V^+)$, $\pi : U_q(\widehat{\mathfrak{g}}) \rightarrow \text{End}(V)$ and $\pi^- : U_q(\widehat{\mathfrak{b}}^-) \rightarrow \text{End}(V^-)$ are level-0 representations, then we have the following identity of linear-operator-valued formal power series in two indeterminates:*

$$\mathcal{R}_{\pi^+\pi}(z_1)_{12} \mathcal{R}_{\pi^+\pi^-}(z_1 z_2)_{13} \mathcal{R}_{\pi\pi^-}(z_2)_{23} = \mathcal{R}_{\pi\pi^-}(z_2)_{23} \mathcal{R}_{\pi^+\pi^-}(z_1 z_2)_{13} \mathcal{R}_{\pi^+\pi}(z_1)_{12}. \tag{2.30}$$

Given a pair of level-0 representations $\pi^\pm : U_q(\widehat{\mathfrak{b}}^\pm) \rightarrow \text{End}(V^\pm)$ it is often convenient to have an explicit expression of $\mathcal{R}_{\pi^+\pi^-}(z)$ which does not rely on computing the coefficients of the series $\mathcal{R}(z)$. Essentially following Jimbo’s approach from [Ji86b], we may try to solve a linear equation for $\mathcal{R}_{\pi^+\pi^-}(z)$. To derive such a linear equation, it is convenient to assume that, say, π^+ extends to a representation of $U_q(\widehat{\mathfrak{g}})$. In this case⁴, one directly obtains the following result.

Proposition 2.6. *If π^+ is a level-0 $U_q(\widehat{\mathfrak{g}})$ -representation and π^- a level-0 $U_q(\widehat{\mathfrak{b}}^-)$ -representation, then for all $u \in U_q(\widehat{\mathfrak{b}}^-)$ we have*

$$\mathcal{R}_{\pi^+\pi^-}(z) \cdot (\pi_z^+ \otimes \pi^-)(\Delta(u)) = (\pi_z^+ \otimes \pi^-)(\Delta^{\text{op}}(u)) \cdot \mathcal{R}_{\pi^+\pi^-}(z). \tag{2.31}$$

Obviously there is a counterpart of Proposition 2.6 with the role of $U_q(\widehat{\mathfrak{b}}^-)$ replaced by $U_q(\widehat{\mathfrak{b}}^+)$.

Remark 2.7. If the solution space of the linear equation (2.31) is 1-dimensional, Proposition 2.6 implies that any solution must be a scalar multiple of $\mathcal{R}_{\pi^+\pi^-}(z)$ and hence satisfy the Yang–Baxter equation. This is well-known if both V^\pm extend to finite-dimensional $U_q(\widehat{\mathfrak{g}})$ -modules. In this case the existence of the universal R-matrix implies the existence of a solution of the intertwining condition (2.31) depending rationally on z . If π^+ and π^- are also both irreducible then it is known, see e.g. [KS95, Sec. 4.2] and [Ch02, Thm. 3], that $V^+(z) \otimes V^-$ is irreducible as a representation of $U_q(\widehat{\mathfrak{g}})(z)$ (extension of scalars to formal Laurent series); hence an application of Schur’s lemma yields the 1-dimensionality of the solution space of (2.31). In this case, the rational intertwiner is called *trigonometric R-matrix*. For more background and detail, see e.g. [He19] and [AV22b, Secs. 2.6 & 2.7].

In the absence of a linear relation such as (2.31), one can use the Yang–Baxter equation (2.30) to determine an explicit expression for one of $\mathcal{R}_{\pi^+\pi}(z)$, $\mathcal{R}_{\pi^+\pi^-}(z)$, or $\mathcal{R}_{\pi\pi^-}(z)$, provided the other two are known. ∅

⁴ One can of course apply $\pi_z^+ \otimes \pi^-$ to (2.15) for arbitrary $U_q(\widehat{\mathfrak{b}}^\pm)$ -representations π^\pm , yielding (2.31) for all $u \in U_q(\widehat{\mathfrak{g}})$ such that $\Delta(u)$ and $\Delta^{\text{op}}(u)$ both lie in $U_q(\widehat{\mathfrak{b}}^+) \otimes U_q(\widehat{\mathfrak{b}}^-)$. However, by applying counits this subalgebra is seen to be equal to $U_q(\widehat{\mathfrak{b}}^+) \cap U_q(\widehat{\mathfrak{b}}^-) = U_q(\widehat{\mathfrak{h}})$. Hence, one would just recover the second statement of Theorem 2.4.

2.9. Adjusting the grading. In this approach the use of the principal grading in Theorem 2.4 avoids further constraints on the representations (e.g. local finiteness conditions). For completeness we briefly explain how to extend the results of Sect. 2.8 to arbitrary grading. For nonnegative integers s_0, s_1 such that $s_0 + s_1$ is nonzero, define a more general Hopf algebra automorphism $\Sigma_z^{s_0, s_1}$ of $U_q(\widehat{\mathfrak{g}})[z, z^{-1}]$ by

$$\Sigma_z^{s_0, s_1}(e_i) = z^{s_i} e_i, \quad \Sigma_z^{s_0, s_1}(f_i) = z^{-s_i} f_i, \quad \Sigma_z^{s_0, s_1}|_{U_q(\widehat{\mathfrak{h}})} = \text{id} \quad (2.32)$$

(note that the choice $s_0 = 0, s_1 = 1$ is used in in [KT14, Eq. (2.11)]).

Rather than giving generalized versions of the main results above and of various statements in the remainder of this work, we make an observation which will allow the reader to generate these statements, as required. Recalling the decomposition (2.22) and the associated terminology, suppose the level-0 $U_q(\widehat{\mathfrak{b}^+})$ -module V is generated by a nonzero element of $V(c_0)$ for some $c_0 \in \mathbb{C}^\times$ (which includes all modules considered in this paper and all irreducible finite-dimensional $U_q(\widehat{\mathfrak{g}})$ -modules). Then the support of V , see Sect. 2.7, is contained in $q^{2\mathbb{Z}}c_0$. Now for any indeterminate y and any integer m , let y^{mD} denote the linear map on V which acts on $V(q^{-2m}c_0)[y, y^{-1}]$ as scalar multiplication by y^m .

Writing the corresponding representation as $\pi : U_q(\widehat{\mathfrak{b}^+}) \rightarrow \text{End}(V)$, the more general grading-shifted representation $\pi_z^{s_0, s_1} := \pi \circ \Sigma_z^{s_0, s_1}$ can be related to the representation shifted by the principal grading as follows. Adjoining to the ring $\mathbb{C}[z, z^{-1}]$ a square root Z of z , we have

$$\pi_z^{s_0, s_1} = \text{Ad}(Z^{(s_0 - s_1)D}) \circ \pi_{Z^{s_0 + s_1}}, \quad (2.33)$$

where on the right-hand side Ad stands for ‘conjugation by’. See [AV22b, Sec. 5.2] for essentially the same point in the context of irreducible finite-dimensional $U_q(\widehat{\mathfrak{g}})$ -representations.

3. The Augmented q-Onsager Algebra, its Twist and its Universal K-Matrix

In parallel with the previous section, we consider a particular subalgebra of $U_q(\widehat{\mathfrak{g}})$ and extend some recent results on universal K-matrices [AV22a, AV22b] in the context of (possibly infinite-dimensional) level-0 representations of Borel subalgebras of quantum affine \mathfrak{sl}_2 . For a related approach tailored to evaluation representations involving essentially the same subalgebra, see [BT18].

3.1. The twist map ψ . We consider the following algebra automorphism and coalgebra antiautomorphism of $U_q(\widehat{\mathfrak{g}})$:

$$\psi := \omega \circ \Phi. \quad (3.1)$$

From (2.18–2.19) and (2.24–2.25), respectively, we immediately deduce

$$(\psi \otimes \psi)(\mathcal{R}) = \mathcal{R}_{21}, \quad (3.2)$$

$$\psi \circ \Sigma_z = \Sigma_{z^{-1}} \circ \psi. \quad (3.3)$$

By the following result, P-symmetric R-matrices $(\mathcal{R}(z))_{21} = \mathcal{R}(z)$ naturally arise in tensor products of representations of the upper and lower Borel subalgebras on the same vector space, provided they are related through ψ and the principal grading is used in the definition of grading-shifted universal R-matrix $\mathcal{R}(z)$, see (2.27).

Lemma 3.1. *Consider two pairs of level-0 representations $\pi^\pm, \varrho^\pm : U_q(\widehat{\mathfrak{b}}^\pm) \rightarrow \text{End}(V)$ such that*

$$\varrho^\mp = \pi^\pm \circ \psi. \tag{3.4}$$

Then $\mathcal{R}_{\pi^+\pi^-}(z)_{21} = \mathcal{R}_{\varrho^+\varrho^-}(z)$.

Proof. Unpacking the definitions (2.28) and (2.27), we have

$$\mathcal{R}_{\pi^+\pi^-}(z)_{21} = \left(((\pi^+ \otimes \pi^-) \circ (\Sigma_z \otimes \text{id}))(\mathcal{R}) \right)_{21} = ((\pi^- \otimes \pi^+) \circ (\text{id} \otimes \Sigma_z))(\mathcal{R}_{21}).$$

Now using (3.2–3.3) we deduce

$$\mathcal{R}_{\pi^+\pi^-}(z)_{21} = ((\pi^- \otimes \pi^+) \circ (\psi \otimes \psi) \circ (\text{id} \otimes \Sigma_{z^{-1}}))(\mathcal{R}).$$

Applying (3.4) and using (2.28) and (2.27) once again, we obtain $\mathcal{R}_{\varrho^+\varrho^-}(z)$ as required. \square

3.2. The augmented q -Onsager algebra. The map ψ plays an important role in the theory of diagonal matrix solutions with a free parameter of the reflection equation in $U_q(\widehat{\mathfrak{g}})$ -modules. Namely, fix a parameter $\xi \in \mathbb{C}^\times$ and consider the following subalgebra of $U_q(\widehat{\mathfrak{g}})$, also called the *(embedded) augmented q -Onsager algebra*:

$$U_q(\mathfrak{k}) := \mathbb{C}\langle e_0 - q^{-1}\xi^{-1}k_0f_1, e_1 - q^{-1}\xi k_1f_0, k_0k_1^{-1}, k_0^{-1}k_1 \rangle. \tag{3.5}$$

This is a left coideal:

$$\Delta(U_q(\mathfrak{k})) \subseteq U_q(\widehat{\mathfrak{g}}) \otimes U_q(\mathfrak{k}). \tag{3.6}$$

The automorphism ψ is the trivial q -deformation of a Lie algebra automorphism of $\widehat{\mathfrak{g}}$, also denoted ψ , and $U_q(\mathfrak{k})$ is the (ξ -dependent) coideal q -deformation of the universal enveloping algebra of the fixed-point subalgebra $\mathfrak{k} = \widehat{\mathfrak{g}}^\psi$, in the style of [Ko14] but with opposite conventions.

Remark 3.2. See [VW20, Rmk. 2.3] for more background on this subalgebra. Note that the definition of $U_q(\mathfrak{k})$ in *loc. cit.* has a misprint: ξ should be replaced by ξ^{-1} . \square

To connect with the universal K -matrix formalism of [AV22a, AV22b], let \widetilde{S} be the bialgebra isomorphism⁵ from $U_q(\widetilde{\mathfrak{g}})$ to $U_q(\widetilde{\mathfrak{g}})^{\text{op, cop}}$ (also known as the *unitary antipode*) defined by the assignments

$$\widetilde{S}(e_i) = -qk_i^{-1}e_i, \quad \widetilde{S}(f_i) = -q^{-1}f_ik_i, \quad \widetilde{S}(k_i^{\pm 1}) = k_i^{\mp 1}, \quad \widetilde{S}(g^{\pm 1}) = g^{\mp 1}. \tag{3.7}$$

Note that $\widetilde{S}^2 = \text{id}$. Now consider⁶ the right coideal subalgebra

$$U_q(\mathfrak{k})' = \widetilde{S}(U_q(\mathfrak{k})) = \mathbb{C}\langle f_0 - q\xi^{-1}e_1k_0^{-1}, f_1 - q\xi e_0k_1^{-1}, k_0k_1^{-1}, k_0^{-1}k_1 \rangle$$

which is the subalgebra considered in [AV22a, Sec. 9.7], forming part of a more general family of right coideal subalgebras (quantum symmetric pair subalgebras) of quantum affine algebras as considered in [Ko14, AV22a, AV22b].

⁵ In particular, \widetilde{S} , like the antipode S itself, is an algebra antiautomorphism and a coalgebra antiautomorphism.

⁶ In general, each element or map in the right coideal setting of [Ko14, AV22a, AV22b] is denoted by a prime on the corresponding object in the current left coideal setting.

3.3. *Universal K-matrix.* By [AV22a, Thm. 8.5], $U_q(\widehat{\mathfrak{g}})$ is endowed with a so-called *standard* universal K-matrix, which is an invertible element in a completion of $U_q(\widehat{\mathfrak{b}}^+)$ satisfying a twisted $U_q(\mathfrak{k})$ -intertwining property and a twisted coproduct formula involving the universal R-matrix⁷

$$\mathcal{R}' = \mathcal{R}_{21}^{-1}. \tag{3.8}$$

There is an action of invertible elements of a completion of $U_q(\widehat{\mathfrak{g}})$, gauge-transforming the universal K-matrix and the twisting operator simultaneously, see [AV22b, Sec. 3.6]. For the case under consideration, there exists a gauge transformation (a ‘Cartan correction’, see [AV22a, Sec. 8.8]) that brings both the intertwining property and the coproduct formula for the universal K-matrix into a particularly nice form. Moreover, the gauge-transformed universal K-matrix still resides in a completion of $U_q(\widehat{\mathfrak{b}}^+)$ and, when shifted by the principal grading, acts as a linear-operator-valued formal power series for all level-0 $U_q(\widehat{\mathfrak{b}}^+)$ -modules.

To make this more precise, let $\Omega : \widetilde{P} \rightarrow \mathbb{C}^\times$ be any group homomorphism such that $\Omega(\alpha_0) = -\xi$ and $\Omega(\alpha_1) = -\xi^{-1}$. Now define a function $G' : \widetilde{P} \rightarrow \mathbb{C}^\times$ by

$$G'(\lambda) = \Omega(\lambda)q^{-(\Phi(\lambda), \lambda)/2}. \tag{3.9}$$

Note that this is not a group homomorphism. Define the corresponding linear operator acting on $U_q(\widehat{\mathfrak{h}})$ -weight modules as follows:

$$G' \cdot v = G'(\lambda)v, \quad v \in V_\lambda, \quad \lambda \in \widetilde{P}. \tag{3.10}$$

Analogously to our definition of the factor κ of the universal R-matrix, we thus obtain an invertible element G' of the weight completion of $U_q(\widehat{\mathfrak{g}})$. Finally, let $\delta = \alpha_0 + \alpha_1$ be the basic imaginary root of $\widehat{\mathfrak{g}}$. Then the following result is a special case of [AV22a, Sec. 9.7], with the coproduct formula a direct consequence of [AV22a, (8.21)].

Proposition 3.3. *There exists an invertible element*

$$\Upsilon' = \sum_{\lambda \in \mathbb{Z}_{\geq 0}\delta} \Upsilon'_\lambda, \quad \Upsilon'_\lambda \in U_q(\widehat{\mathfrak{n}}^+)_\lambda, \tag{3.11}$$

such that the invertible element

$$\mathcal{K}' := G' \cdot \Upsilon' \tag{3.12}$$

satisfies

$$\mathcal{K}' \cdot u = \psi(u) \cdot \mathcal{K}' \quad \text{for all } u \in U_q(\mathfrak{k})', \tag{3.13}$$

$$\Delta(\mathcal{K}') = (1 \otimes \mathcal{K}') \cdot (\psi \otimes \text{id})(\mathcal{R}') \cdot (\mathcal{K}' \otimes 1). \tag{3.14}$$

Remark 3.4. In general, a universal K-matrix \mathcal{K}' satisfying the simple 3-factor coproduct formula (3.14) is called *semi-standard*, see [AV22a, Sec. 8.10] and cf. [AV22b, Ex. 3.6.3 (2)]. It corresponds to a particular choice of a *twist pair* (ψ, J) where ψ is a *bialgebra isomorphism* from $U_q(\widehat{\mathfrak{g}})$ to $U_q(\widehat{\mathfrak{g}})^{\text{cop}}$ (essentially the composition of ω with a diagram automorphism determined by the coideal subalgebra) and J is the trivial Drinfeld twist

⁷ Note that our convention for the coproduct is as in [AV22a], but the ordering of the tensor product of the two Borel subalgebras is opposite. Hence the R-matrix in [AV22a], denoted here by \mathcal{R}' , is equal to \mathcal{R}_{21}^{-1} .

$1 \otimes 1$, see [AV22a, Sec. 2.4 and 2.5]. The semi-standard K-matrix is always available; what is rather special in the case of the augmented q-Onsager algebra is that the semi-standard K-matrix coincides with the standard K-matrix (and hence lies in a completion of $U_q(\widehat{\mathfrak{b}^+})$). ∅

Now we transform this formalism [AV22a] for the right coideal subalgebra $U_q(\mathfrak{k})'$, expressed in terms of the universal R-matrix \mathcal{R}' , to a formalism for the left coideal subalgebra $U_q(\mathfrak{k}) = \widetilde{S}(U_q(\mathfrak{k})')$, expressed in terms of the universal R-matrix \mathcal{R} as used in this paper. To do this, note that, when going from a $U_q(\widehat{\mathfrak{g}})$ -weight module to its dual, weights transform as $\lambda \mapsto -\lambda$. This defines the extension of S and \widetilde{S} to a map on the weight completion of $U_q(\widehat{\mathfrak{g}})$. Therefore $\widetilde{S}(\Omega) = \Omega^{-1}$ but the non-group-like factor of G' is fixed by \widetilde{S} . We define $G : \widetilde{P} \rightarrow \mathbb{C}^\times$ by

$$G(\lambda) := \Omega(\lambda)q^{(\Phi(\lambda), \lambda)/2} \tag{3.15}$$

so that $G = \widetilde{S}(G')^{-1}$. Also, we set

$$\Upsilon := \widetilde{S}(\Upsilon')^{-1} = \sum_{\lambda \in \mathbb{Z}_{\geq 0}\delta} \Upsilon_\lambda, \quad \Upsilon_\lambda \in \widetilde{S}(U_q(\widehat{\mathfrak{n}^+})_\lambda) \subset U_q(\widehat{\mathfrak{h}}) \cdot U_q(\widehat{\mathfrak{n}^+})_\lambda. \tag{3.16}$$

Proposition 3.5. *The element*

$$\mathcal{K} := \widetilde{S}(\mathcal{K}')^{-1} = G \cdot \Upsilon \tag{3.17}$$

satisfies

$$\mathcal{K} \cdot u = \psi(u) \cdot \mathcal{K} \quad \text{for all } u \in U_q(\mathfrak{k}), \tag{3.18}$$

$$\Delta(\mathcal{K}) = (\mathcal{K} \otimes 1) \cdot (\text{id} \otimes \psi)(\mathcal{R}) \cdot (1 \otimes \mathcal{K}). \tag{3.19}$$

Proof. This follows straightforwardly from Proposition 3.3. Namely, we apply \widetilde{S} to (3.13) and $(\widetilde{S} \otimes \widetilde{S}) \circ \sigma$ to (3.14), and use the fact that $\widetilde{S} \circ \psi = \psi \circ \widetilde{S}$ and $(\widetilde{S} \otimes \widetilde{S})(\mathcal{R}) = \mathcal{R}$. \square

Note that $U_q(\widehat{\mathfrak{b}^+})$ is a bialgebra and, as expected, the right-hand side of (3.19) lies in a completion of $U_q(\widehat{\mathfrak{b}^+}) \otimes U_q(\widehat{\mathfrak{b}^+})$, since ψ interchanges the two Borel subalgebras $U_q(\widehat{\mathfrak{b}^\pm})$. The reflection equation satisfied by the universal element \mathcal{K} is as follows:

$$\mathcal{R} \cdot (\mathcal{K} \otimes 1) \cdot (\text{id} \otimes \psi)(\mathcal{R}) \cdot (1 \otimes \mathcal{K}) = (1 \otimes \mathcal{K}) \cdot (\text{id} \otimes \psi)(\mathcal{R}) \cdot (\mathcal{K} \otimes 1) \cdot \mathcal{R}. \tag{3.20}$$

This is a consequence of the linear relation (2.15) for \mathcal{R} and the coproduct formula (3.19) for \mathcal{K} , alongside (3.2) and $\psi^2 = \text{id}$.

3.4. The action of the universal K-matrix on level-0 representations. To deduce that \mathcal{K} has a well-defined action on level-0 representations of, say, $U_q(\widehat{\mathfrak{b}^+})$, we can proceed in a similar way to the case of the R-matrix. This builds on the finite-dimensional theory for more general quantum symmetric pair subalgebras in [AV22b, Sec. 4].

First note that if π is a level-0 representation, π and the twisted representation $\pi \circ \psi$ coincide on $U_q(\widehat{\mathfrak{h}})$. Now let z once again be a formal variable. Note that by (3.15) the function G sends the basic imaginary root δ to 1. Hence the proof of [AV22b, Prop. 4.3.1 (3)] implies that the corresponding factor G of the universal K-matrix descends to level-0 modules. Furthermore, the argument that shows $\Sigma_z(\Theta)$ is a $U_q(\widehat{\mathfrak{n}^+}) \otimes U_q(\widehat{\mathfrak{n}^-})$ -valued

formal power series can be easily adapted to Υ ; it yields a formal power series with coefficients in $\widehat{S}(U_q(\widehat{\mathfrak{n}^+}) \subset U_q(\widehat{\mathfrak{b}^+})$:

$$\Sigma_z(\Upsilon) = \sum_{r \geq 0} z^r \sum_{\lambda \in \mathbb{Z}_{\geq 0} \delta, \text{ht}(\lambda)=r} \Upsilon_\lambda.$$

Now consider the grading-shifted universal K-matrix:

$$\mathcal{K}(z) = \Sigma_z(\mathcal{K}). \tag{3.21}$$

Noting that the form of Υ implies that \mathcal{K} commutes with k_1 , we arrive at the following main result, which is a boundary analogue of Theorem 2.4.

Theorem 3.6. *Consider a level-0 representation $\pi : U_q(\widehat{\mathfrak{b}^+}) \rightarrow \text{End}(V)$. Then⁸*

$$\mathcal{K}_\pi(z) := \pi(\mathcal{K}(z)) \in \text{End}(V)[[z]] \tag{3.22}$$

is well-defined and commutes with $\pi(k_1)$.

We will also need boundary counterparts of Propositions 2.5 and 2.6. Consider two indeterminates z_1, z_2 . Applying $\Sigma_{z_1} \otimes \Sigma_{z_2}$ to (3.20) and using (3.3), we obtain the following reflection equation for the grading-shifted universal operators:

$$\begin{aligned} \mathcal{R}(z_1/z_2) \cdot (\mathcal{K}(z_1) \otimes 1) \cdot (\text{id} \otimes \psi)(\mathcal{R}(z_1 z_2)) \cdot (1 \otimes \mathcal{K}(z_2)) &= \\ = (1 \otimes \mathcal{K}(z_2)) \cdot (\text{id} \otimes \psi)(\mathcal{R}(z_1 z_2)) \cdot (\mathcal{K}(z_1) \otimes 1) \cdot \mathcal{R}(z_1/z_2). \end{aligned} \tag{3.23}$$

Recalling that the universal R-matrix \mathcal{R} lies in a completion of $U_q(\widehat{\mathfrak{b}^+}) \otimes U_q(\widehat{\mathfrak{b}^-})$ and applying a tensor product of suitable representations to (3.23), one obtains the *right reflection equation* with multiplicative spectral parameters for P-symmetric R-matrices, as we now state precisely.

Proposition 3.7. *Consider level-0 representations $\pi^+ : U_q(\widehat{\mathfrak{b}^+}) \rightarrow \text{End}(V^+)$ and $\pi : U_q(\widehat{\mathfrak{g}}) \rightarrow \text{End}(V)$ such that $\pi \circ \psi = \pi$. Then*

$$\begin{aligned} \mathcal{R}_{\pi^+ \pi}(z_1/z_2)(\mathcal{K}_{\pi^+}(z_1) \otimes \text{Id}_V) \mathcal{R}_{\pi^+ \pi}(z_1 z_2)(\text{Id}_{V^+} \otimes \mathcal{K}_\pi(z_2)) &= \\ = (\text{Id}_{V^+} \otimes \mathcal{K}_\pi(z_2)) \mathcal{R}_{\pi^+ \pi}(z_1 z_2)(\mathcal{K}_{\pi^+}(z_1) \otimes \text{Id}_V) \mathcal{R}_{\pi^+ \pi}(z_1/z_2). \end{aligned} \tag{3.24}$$

The use of linear relations to find explicit solutions of reflection equations was proposed in [MN98, DG02, DM03]. As before, we assume that π extends to a $U_q(\widehat{\mathfrak{g}})$ -representation,⁹ in which case it restricts to a $U_q(\mathfrak{k})$ -representation and we obtain the following result as a consequence of (3.3).

Proposition 3.8. *If $\pi : U_q(\widehat{\mathfrak{g}}) \rightarrow \text{End}(V)$ is a level-0 representation such that $\pi \circ \psi = \pi$, then*

$$\mathcal{K}_\pi(z) \cdot \pi_z(u) = \pi_{1/z}(u) \cdot \mathcal{K}_\pi(z) \quad \text{for all } u \in U_q(\mathfrak{k}). \tag{3.25}$$

We close this section with some comments parallel to Remark 2.7.

⁸ In Sect. 6 we will use this notation for a rescaled version of the action of the grading-shifted universal K-matrix.

⁹ Analogous to the case of the R-matrix, we can observe that the intersection of $U_q(\mathfrak{k})$ and $U_q(\widehat{\mathfrak{b}^+})$ is contained in $U_q(\widehat{\mathfrak{h}})$. Therefore, applying a level-0 representation π to (3.18) just recovers the second part of Theorem 3.6.

Remark 3.9. If the solution space of (3.25) is 1-dimensional, Proposition 3.8 implies that any solution must be a scalar multiple of $\mathcal{K}(z)$ and hence automatically satisfy the reflection equation (3.24). In the case that $\pi : U_q(\widehat{\mathfrak{b}^+}) \rightarrow \text{End}(V)$ extends to a representation and V is finite-dimensional, there is an analogue to Remark 2.7. Namely, the solution space of (3.25) for irreducible representations is 1-dimensional and the existence of a solution of the intertwining condition (3.25) depending rationally on z leads to a *trigonometric K-matrix*. See [AV22b, Secs. 5 and 6] for more detail.

To explicitly determine $\mathcal{K}_{\pi^+}(z)$ in the cases where $\pi^+ : U_q(\widehat{\mathfrak{b}^+}) \rightarrow \text{End}(V)$ does not extend to a $U_q(\widehat{\mathfrak{g}})$ -representation, we will use the reflection equation (3.24), with the other K-matrix $\mathcal{K}_\pi(z)$ determined using Proposition 3.8. ∅

4. Borel Representations in Terms of the q-Oscillator Algebra

4.1. The infinite-dimensional vector space W. The countably-infinite-dimensional vector space plays a central role in the theory of Baxter’s Q-operators. We may define it as the free \mathbb{C} -module over a given set $\{w_j\}_{j \in \mathbb{Z}_{\geq 0}}$:

$$W = \bigoplus_{j \geq 0} \mathbb{C}w_j. \tag{4.1}$$

Given this distinguished basis, elements of $\text{End}(W)$ naturally identify with infinite-by-infinite matrices with the property that all but finitely many entries of each column are zero.

It is convenient to work with a particular subalgebra of $\text{End}(W)$ depending on the deformation parameter q . More precisely, consider the \mathbb{C} -linear maps a, a^\dagger on W defined by

$$a \cdot w_{j+1} = w_j, \quad a \cdot w_0 = 0, \quad a^\dagger \cdot w_j = (1 - q^{2(j+1)})w_{j+1} \tag{4.2}$$

for all $j \in \mathbb{Z}_{\geq 0}$. For the description of L-operators associated to $U_q(\widehat{\mathfrak{g}})$ acting on $W \otimes \mathbb{C}^2$ (particular solutions of the Yang–Baxter equation), it is convenient to consider a linear operator q^D which is a square root of $1 - a^\dagger a$, i.e. $q^D \cdot w_j = q^j w_j$ for $j \in \mathbb{Z}_{\geq 0}$. Note that q^D is invertible and we let q^{-D} denote its inverse.

Remark 4.1. Often the q-oscillator algebra is defined as an abstract algebra, generated by a, a^\dagger and $q^{\pm D}$ subject to certain relations, which naturally embeds into $\text{End}(W)$. This version of the q-oscillator algebra appeared in the guise of a topological algebra for instance in [BGKNR10, Sec. 2.3] and with slightly different conventions in [KT14]¹⁰. ∅

4.2. Diagonal operators from functions and an extended q-oscillator algebra. To accommodate the action of the universal R and K-matrices on certain level-0 modules, we will need an extension of the commutative subalgebra $\langle q^{\pm D} \rangle$ and work over the commutative ring $\mathbb{C}[[z]]$.

Denote by \mathcal{F} the commutative algebra of functions from $\mathbb{Z}_{\geq 0}$ to $\mathbb{C}[[z]]$. For any $f \in \mathcal{F}$ we define $f(D) \in \text{End}(W)[[z]]$ via

$$f(D) \cdot w_j = f(j)w_j. \tag{4.3}$$

¹⁰ The two vector spaces W_1 and W_2 introduced in [KT14, Sec. 2.3] are naturally isomorphic, so that the two algebras Osc_1 and Osc_2 can be identified with the same subalgebra of $\text{End}(W_1) \cong \text{End}(W_2)$.

Thus, we obtain an algebra embedding $\mathcal{F} \rightarrow \text{End}(W)[[z]]$. Now we combine this with the maps a, a^\dagger defined above (viewed as maps on $W \otimes \mathbb{C}[[z]]$, acting trivially on the second factor).

Definition 4.2. The (extended) q -oscillator algebra is the subalgebra $\mathcal{A} \subset \text{End}(W)[[z]]$ generated by a^\dagger, a and $\mathcal{F}(D)$. ∅

As can be verified on basis vectors, in \mathcal{A} one has the relations

$$\begin{aligned} aa^\dagger &= 1 - q^{2(D+1)}, & a^\dagger a &= 1 - q^{2D}, \\ af(D) &= f(D+1)a, & f(D)a^\dagger &= a^\dagger f(D+1). \end{aligned} \tag{4.4}$$

One straightforwardly verifies that the subalgebras $\mathcal{F}(D), \langle a^\dagger \rangle$ and $\langle a \rangle$ are self-centralizing. Note that the operator

$$\bar{a}^\dagger := -q^{-2D}a^\dagger \in \text{End}(W) \tag{4.5}$$

sends w_j to $(1 - q^{-2(j+1)})w_{j+1}$. Clearly, \mathcal{A} is also generated by \bar{a}^\dagger, a and $\mathcal{F}(D)$. The transformation $q \mapsto q^{-1}$ defines an algebra automorphism of \mathcal{A} , preserving the subalgebra $\mathcal{F}(D)$, fixing the generator a and interchanging the generators a^\dagger and \bar{a}^\dagger .

4.3. Endomorphisms of $W \otimes W$. The linear maps

$$a_1 := a \otimes \text{Id}_W, \quad a_1^\dagger := a^\dagger \otimes \text{Id}_W, \quad a_2 := \text{Id}_W \otimes a, \quad a_2^\dagger := \text{Id}_W \otimes a^\dagger$$

together with $\mathcal{F}(D_1) \cup \mathcal{F}(D_2)$ generate $\mathcal{A} \otimes \mathcal{A}$ over $\mathbb{C}[[z]]$. We will need a larger subalgebra of $\text{End}(W \otimes W)$: we will allow all functions of two nonnegative integers as well as formal power series in certain locally nilpotent endomorphisms.

Denote by $\mathcal{F}^{(2)}$ the commutative algebra of functions from $\mathbb{Z}_{\geq 0} \times \mathbb{Z}_{\geq 0}$ to $\mathbb{C}[[z]]$. For any $f \in \mathcal{F}^{(2)}$ we define $f(D_1, D_2) \in \text{End}(W \otimes W)[[z]]$ via

$$f(D_1, D_2) \cdot (w_j \otimes w_k) = f(j, k)w_j \otimes w_k, \tag{4.6}$$

yielding an algebra embedding $\mathcal{F}^{(2)} \rightarrow \text{End}(W \otimes W)[[z]]$. Now note that $a_1 a_2^\dagger$ and $a_1^\dagger a_2$ are locally nilpotent endomorphisms of $W \otimes W$. Hence, for any $g_{k,\ell}, h_{k,\ell} \in \mathcal{F}^{(2)}$ series of the form

$$\sum_{k,\ell \geq 0} (a_2^\dagger)^\ell g_{k,\ell}(D_1, D_2) a_1^k, \quad \sum_{k,\ell \geq 0} (a_1^\dagger)^k h_{k,\ell}(D_1, D_2) a_2^\ell \tag{4.7}$$

truncate when applied to any basis vector $w_j \otimes w_{j'}$. We obtain a class of well-defined elements of $\text{End}(W \otimes W)[[z]]$. We denote by $\mathcal{A}^{(2)}$ the $\mathbb{C}[[z]]$ -span of the operator-valued formal series (4.7), which is easily seen to be a subalgebra of $\text{End}(W \otimes W)[[z]]$.

4.4. *The Borel representations.* We introduce four level-0 representations of $U_q(\widehat{\mathfrak{b}^+})$. First of all, let $\mu \in \mathbb{C}$ be a free parameter. It is straightforward to check that the following assignments define a representation ν of $U_q(\widehat{\mathfrak{g}})$ on W :

$$\begin{aligned} \nu(e_0) = \nu(f_1) &= \frac{1}{1 - q^2} a^\dagger, & \nu(k_0) &= q^{-\mu+1+2D}, \\ \nu(e_1) = \nu(f_0) &= \frac{q^2}{1 - q^2} a(q^{-\mu} - q^{\mu-2D}), & \nu(k_1) &= q^{\mu-1-2D}. \end{aligned} \quad (4.8)$$

The module structure on W defined by ν is the evaluation Verma module: affinizations of finite-dimensional irreducible $U_q(\mathfrak{sl}_2)$ -modules arise as quotients if $\mu \in \mathbb{Z}_{>0}$ (also see [KT14, Sec. 2.2]).

We will in addition consider three $U_q(\widehat{\mathfrak{b}^+})$ -representations which do not extend to representations of $U_q(\widehat{\mathfrak{g}})$. A useful reducible representation $\phi : U_q(\widehat{\mathfrak{b}^+}) \rightarrow \text{End}(W)$ is given by

$$\phi(e_0) = 0, \quad \phi(e_1) = \frac{q}{1 - q^2} a, \quad \phi(k_0) = q^{\mu+1+2D}, \quad \phi(k_1) = q^{-\mu-1-2D} \quad (4.9)$$

which is closely related to the special evaluation homomorphism defined in [KT14, Eq. (4.6)]. The following representations $\varrho, \bar{\varrho} : U_q(\widehat{\mathfrak{b}^+}) \rightarrow \text{End}(W)$ play an essential role in the definition of Baxter Q-operators:

$$\begin{aligned} \varrho(e_0) &= \frac{1}{1 - q^2} a^\dagger, & \varrho(e_1) &= \frac{q^2}{1 - q^2} a, & \varrho(k_0) &= q^{2D}, & \varrho(k_1) &= q^{-2D}, \\ \bar{\varrho}(e_0) &= \frac{q^2}{1 - q^2} \bar{a}^\dagger, & \bar{\varrho}(e_1) &= \frac{1}{1 - q^2} a, & \bar{\varrho}(k_0) &= q^{2(D+1)}, & \bar{\varrho}(k_1) &= q^{-2(D+1)}. \end{aligned} \quad (4.10)$$

They correspond to the representations $L_{1,a}^\pm$ introduced in [HJ12, Def. 3.7] for suitable $a \in \mathbb{C}^\times$ (called *prefundamental* representations in [FH15] which considers their role in the construction of Q-operators for closed chains).

We will henceforth repeatedly denote grading-shifted representations by the notation (2.29). Note that the grading-shifted representation ν_z is an algebra homomorphism from $U_q(\widehat{\mathfrak{g}})$ to $\text{End}(W)[z, z^{-1}]$. Furthermore, the grading-shifted representations $\nu_z|_{U_q(\widehat{\mathfrak{b}^+})}$, $\phi_z, \varrho_z, \bar{\varrho}_z$ are algebra homomorphisms from $U_q(\widehat{\mathfrak{b}^+})$ to $\text{End}(W)[z] \subset \text{End}(W)[[z]]$. Finally, note that $\varrho_z, \bar{\varrho}_z$ correspond to the representations defined by [KT14, Eq. (3.5)].

Remark 4.3. The grading-shifted representation in [VW20, Eq. (2.9)] is related to ϱ_z by a factor of -1 in the actions of e_0 and e_1 : in other words it is equal to ϱ_{-z} . Since the Baxter Q-operators only depend on z^2 , see [VW20, Lem. 4.5], there are no serious discrepancies. The benefit of the current choice is its consistency across the relevant level-0 representations, with ν having the same sign convention as finite-dimensional representations such as Π , see Sect. 5. ∅

4.5. *The $U_q(\widehat{\mathfrak{b}^+})$ -intertwiner \mathcal{O} .* The tensor products $\varrho_{q^{-\mu/2z}} \otimes \bar{\varrho}_{q^{\mu/2z}}$ and $\nu_z \otimes \phi_z$ of shifted representations are closely related in the following sense: the two induced $U_q(\widehat{\mathfrak{b}^+})$ -actions on $W \otimes W$ are conjugate by an element in $\mathcal{A}^{(2)}$ which is independent of z . More precisely, consider the deformed exponential

$$e_{q^2}(x) = \sum_{k=0}^{\infty} \frac{x^k}{(q^2; q^2)_k}. \tag{4.11}$$

We refer to Appendix A for more detail. We now define the following element of $\text{GL}(W \otimes W)$:

$$\mathcal{O} = e_{q^2}(q^2 a_1 \bar{a}_2^\dagger)^{-1} q^{\mu(D_1 - D_2)/2}. \tag{4.12}$$

The following statement is [KT14, Eq. (4.4)] and connects to [FH15, Thm. 3.8]; for completeness we provide a proof in the present conventions.

Theorem 4.4. *The $U_q(\widehat{\mathfrak{b}^+})$ -representations $\varrho_{q^{-\mu/2z}} \otimes \bar{\varrho}_{q^{\mu/2z}}$ and $\nu_z \otimes \phi_z$ are intertwined by \mathcal{O} :*

$$\mathcal{O} (\varrho_{q^{-\mu/2z}} \otimes \bar{\varrho}_{q^{\mu/2z}})(\Delta(u)) = (\nu_z \otimes \phi_z)(\Delta(u)) \mathcal{O} \quad \text{for all } u \in U_q(\widehat{\mathfrak{b}^+}). \tag{4.13}$$

Proof. The relations (A.13–A.15) can be evaluated at $y = q^2$, yielding

$$\begin{aligned} q^{\mu(D_2 - D_1)/2} e_{q^2}(q^2 a_1 \bar{a}_2^\dagger) \bar{a}_2^\dagger &= (q^{-\mu/2} a_1^\dagger + q^{2(D_1+1)+\mu/2} \bar{a}_2^\dagger) q^{\mu(D_2 - D_1)/2} e_{q^2}(q^2 a_1 \bar{a}_2^\dagger), \\ q^{\mu(D_2 - D_1)/2} e_{q^2}(q^2 a_1 \bar{a}_2^\dagger) (a_1(q^{-2\mu} - q^{-2D_1}) + q^{-2(D_1+1)} a_2) &= \\ &= (a_1 q^{-3\mu/2} + q^{-\mu/2 - 2(D_1+1)} a_2) q^{\mu(D_2 - D_1)/2} e_{q^2}(q^2 a_1 \bar{a}_2^\dagger), \\ q^{\mu(D_2 - D_1)/2} e_{q^2}(q^2 a_1 \bar{a}_2^\dagger) q^{2(D_1+D_2+1)} &= q^{2(D_1+D_2+1)} q^{\mu(D_2 - D_1)/2} e_{q^2}(q^2 a_1 \bar{a}_2^\dagger), \\ q^{\mu(D_2 - D_1)/2} e_{q^2}(q^2 a_1 \bar{a}_2^\dagger) q^{-2(D_1+D_2+1)} &= q^{-2(D_1+D_2+1)} q^{\mu(D_2 - D_1)/2} e_{q^2}(q^2 a_1 \bar{a}_2^\dagger). \end{aligned}$$

These directly imply (4.13) for $u \in \{e_0, e_1, k_0, k_1\}$. □

4.6. *Formalism for $U_q(\widehat{\mathfrak{b}^-})$.* Recall from (3.1) the automorphism ψ which interchanges the two Borel subalgebras. Note that the representation $\nu : U_q(\widehat{\mathfrak{g}}) \rightarrow \text{End}(W)$ satisfies

$$\nu = \nu \circ \psi. \tag{4.14}$$

Hence, it is natural to define representations of $U_q(\widehat{\mathfrak{b}^-})$ corresponding to ϱ , $\bar{\varrho}$ and ϕ , as follows:

$$\varrho^- := \varrho \circ \psi, \quad \bar{\varrho}^- := \bar{\varrho} \circ \psi, \quad \phi^- := \phi \circ \psi. \tag{4.15}$$

Explicitly, we have

$$\begin{aligned} \varrho^-(f_0) &= \frac{q^2}{1 - q^2} a, & \varrho^-(f_1) &= \frac{1}{1 - q^2} a^\dagger, & \varrho^-(k_0) &= q^{2D}, & \varrho^-(k_1) &= q^{-2D}, \\ \bar{\varrho}^-(f_0) &= \frac{1}{1 - q^2} a, & \bar{\varrho}^-(f_1) &= \frac{q^2}{1 - q^2} \bar{a}^\dagger, & \bar{\varrho}^-(k_0) &= q^{2(D+1)}, & \bar{\varrho}^-(k_1) &= q^{-2(D+1)}, \\ \phi^-(f_0) &= \frac{q}{1 - q^2} a, & \phi^-(f_1) &= 0, & \phi^-(k_0) &= q^{\mu+1+2D}, & \phi^-(k_1) &= q^{-\mu-1-2D}. \end{aligned} \tag{4.16}$$

By (3.3), whereas the grading-shifted representations $\varrho_z, \bar{\varrho}_z, \phi_z$ take values in $\text{End}(W) \otimes \mathbb{C}[z]$, their negative counterparts $\varrho_z^-, \bar{\varrho}_z^-, \phi_z^-$ take values in $\text{End}(W) \otimes \mathbb{C}[z^{-1}]$.

Since ψ is a coalgebra antiautomorphism, using (3.3) we immediately deduce the following characterization of the tensorial opposite of the intertwiner \mathcal{O} .

Corollary 4.5. *The linear map*

$$\mathcal{O}_{21} = e_{q^2} (q^2 \bar{a}_1^\dagger a_2)^{-1} q^{\mu(D_2 - D_1)/2} \in \text{End}(W \otimes W). \tag{4.17}$$

intertwines the $U_q(\widehat{\mathfrak{b}^-})$ -representations $\bar{\varrho}_{q^{-\mu/2z}}^- \otimes \varrho_{q^{\mu/2z}}^-$ and $\phi_z^- \otimes \nu_z$, viz.

$$\mathcal{O}_{21} (\bar{\varrho}_{q^{-\mu/2z}}^- \otimes \varrho_{q^{\mu/2z}}^-)(\Delta(u)) = (\phi_z^- \otimes \nu_z)(\Delta(u)) \mathcal{O}_{21} \text{ for all } u \in U_q(\widehat{\mathfrak{b}^-}). \tag{4.18}$$

5. L-Operators and R-Operators

In order to define L-operators, we recall the standard 2-dimensional representation $\Pi : U_q(\widehat{\mathfrak{g}}) \rightarrow \text{End}(\mathbb{C}^2)$ determined by

$$\begin{aligned} \Pi(e_0) = \Pi(f_1) &= \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, & \Pi(k_0) &= \begin{pmatrix} q^{-1} & 0 \\ 0 & q \end{pmatrix}, \\ \Pi(e_1) = \Pi(f_0) &= \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, & \Pi(k_1) &= \begin{pmatrix} q & 0 \\ 0 & q^{-1} \end{pmatrix}. \end{aligned} \tag{5.1}$$

In analogy with (4.14), we have

$$\Pi = \Pi \circ \psi. \tag{5.2}$$

5.1. L-operators for $U_q(\widehat{\mathfrak{b}^+})$ -modules. We will now obtain explicit formulas for certain scalar multiples of the four different actions of the grading-shifted universal R-matrix on $W \otimes \mathbb{C}^2$. In these cases both Theorem 2.4 and Proposition 2.6 apply. It turns out that the relevant linear equations all have 1-dimensional solution spaces over $\mathbb{C}[[z]]$. The following linear operators are convenient scalar multiples.

$$\mathcal{L}_\varrho(z) = \begin{pmatrix} q^D & a^\dagger q^{-D-1} z \\ a q^{D+1} z & q^{-D} - q^{D+2} z^2 \end{pmatrix}, \tag{5.3}$$

$$\mathcal{L}_{\bar{\varrho}}(z) = \begin{pmatrix} q^{D+1} - q^{-D+1} z^2 & \bar{a}^\dagger q^{-D} z \\ a q^D z & q^{-D-1} \end{pmatrix}, \tag{5.4}$$

$$\mathcal{L}_\nu(z) = \begin{pmatrix} q^D - q^{-D+\mu} z^2 & a^\dagger q^{-D-2+\mu} z \\ a q (q^{D-\mu} - q^{-D+\mu}) z & q^{-D-1+\mu} - q^{D+1} z^2 \end{pmatrix}, \tag{5.5}$$

$$\mathcal{L}_\phi(z) = \begin{pmatrix} q^{D+1} & 0 \\ a q^{D+1} z & q^{-D-\mu} \end{pmatrix}. \tag{5.6}$$

Remark 5.1. We have abused notation by representing linear operators on $\text{End}(W \otimes \mathbb{C}^2)$ as 2×2 matrices with coefficients in $\text{End}(W)$ (as opposed to the conventional usage that realizes operators on $\text{End}(\mathbb{C}^2 \otimes W)$ in this way). ∅

The following result is [KT14, Cor. 4.2].

Theorem 5.2. *The above L-operators satisfy the following relation in $\text{End}(W \otimes W \otimes \mathbb{C}^2)[[z]]$:*

$$\mathcal{O}_{12} \mathcal{L}_\varrho(q^{-\mu/2}z)_{13} \mathcal{L}_{\bar{\varrho}}(q^{\mu/2}z)_{23} = \mathcal{L}_\nu(z)_{13} \mathcal{L}_\phi(z)_{23} \mathcal{O}_{12}. \tag{5.7}$$

Proof. From (2.16) one deduces

$$\begin{aligned} \mathcal{L}_\varrho(q^{-\mu/2}z)_{13} \mathcal{L}_{\bar{\varrho}}(q^{\mu/2}z)_{23} &\propto (\varrho_{q^{-\mu/2}z} \otimes \bar{\varrho}_{q^{\mu/2}z} \otimes \Pi)((\Delta \otimes \text{id})(\mathcal{R})), \\ \mathcal{L}_\nu(z)_{13} \mathcal{L}_\phi(z)_{23} &\propto (\nu_z \otimes \phi_z \otimes \Pi)((\Delta \otimes \text{id})(\mathcal{R})). \end{aligned}$$

Now Theorem 4.4 implies (5.7) up to a scalar. By applying both sides to $w_0 \otimes w_0 \otimes \binom{1}{0}$ one observes that the scalar is 1. □

Given the L-operators for the various $U_q(\widehat{\mathfrak{b}}^+)$ -representations, Lemma 3.1 provides us with L-operators for the corresponding $U_q(\widehat{\mathfrak{b}}^-)$ -representations: $\mathcal{L}_\pi^-(z) = \mathcal{L}_\pi(z)_{21}$ for $\pi \in \{\varrho, \bar{\varrho}, \nu, \phi\}$. These are scalar multiples of $\mathcal{R}_{\Pi\varrho^-}(z)$, $\mathcal{R}_{\Pi\bar{\varrho}^-}(z)$, $\mathcal{R}_{\Pi\nu}(z)$ and $\mathcal{R}_{\Pi\phi^-}(z)$, respectively. Theorem 5.2 immediately yields the following result:

Corollary 5.3. *The following relation in $\text{End}(\mathbb{C}^2 \otimes W \otimes W)[[z]]$ is satisfied:*

$$\mathcal{O}_{32} \mathcal{L}_\varrho^-(q^{-\mu/2}z)_{13} \mathcal{L}_{\bar{\varrho}}^-(q^{\mu/2}z)_{12} = \mathcal{L}_\nu^-(z)_{13} \mathcal{L}_\phi^-(z)_{12} \mathcal{O}_{32}. \tag{5.8}$$

5.2. Actions of $\mathcal{R}(z)$ on tensor products of infinite-dimensional Borel representations. By Theorem 2.4, the grading-shifted universal R-matrix has well-defined actions on the tensor product of the level-0 modules (ν, W) and (ϕ^-, W) and on the tensor product of the level-0 modules (ϱ, W) and $(\bar{\varrho}^-, W)$ as $\text{End}(W \otimes W)$ -valued formal power series. Note that, using the terminology of Sect. 2.7, $\mathbb{C}w_0 \otimes w_0 \subset W \otimes W$ is the subspace of weight q^{-2} and hence $w_0 \otimes w_0$ is an eigenvector of both actions of the universal R-matrix with invertible eigenvalues. It is convenient to use rescaled linear-operator-valued formal power series

$$\mathcal{R}_{\varrho\bar{\varrho}}(z), \mathcal{R}_{\nu\phi}(z) \in \text{End}(W \otimes W)[[z]], \tag{5.9}$$

uniquely defined by the condition that they fix $w_0 \otimes w_0$:

$$\begin{aligned} \mathcal{R}_{\varrho\bar{\varrho}}(z) &\propto (\varrho \otimes \bar{\varrho}^-)(\mathcal{R}(z)), & \mathcal{R}_{\varrho\bar{\varrho}}(z) \cdot (w_0 \otimes w_0) &= w_0 \otimes w_0, \\ \mathcal{R}_{\nu\phi}(z) &\propto (\nu \otimes \phi^-)(\mathcal{R}(z)), & \mathcal{R}_{\nu\phi}(z) \cdot (w_0 \otimes w_0) &= w_0 \otimes w_0. \end{aligned} \tag{5.10}$$

These power series will appear in the boundary factorization identity. In Appendix B we obtain explicit expressions for $\mathcal{R}_{\varrho\bar{\varrho}}(z)$ and $\mathcal{R}_{\nu\phi}(z)$, although we will not need these for the proof of the boundary factorization identity using the universal K-matrix formalism of Sect. 3.

6. K-Operators

In this section we consider solutions of reflection equations associated to the subalgebra $U_q(\mathfrak{k})$.

6.1. *Right K-operators.* By Theorem 3.6, applying any of the level-0 $U_q(\widehat{\mathfrak{b}}^+)$ -representations $\varrho, \bar{\varrho}, \nu, \phi$ to the grading-shifted universal K-matrix associated to $U_q(\mathfrak{k})$ we obtain $\text{End}(W)$ -valued formal power series, satisfying the reflection equation (3.7). Moreover, in terms of the terminology of Sect. 2.7, the weight subspaces of all four actions are all 1-dimensional and therefore w_0 is an eigenvector of each action with invertible eigenvalue. We will consider the scalar multiples of these linear operators which fix w_0 :

$$\mathcal{K}_\pi(z) \propto \pi(\mathcal{K}(z)), \quad \mathcal{K}_\pi(z) \cdot w_0 = w_0. \tag{6.1}$$

for $\pi \in \{\varrho, \bar{\varrho}, \nu, \phi\}$. It is convenient to obtain explicit expressions by applying Propositions 3.7 and 3.8. These could be found independently of the universal K-matrix formalism, either by solving the reflection equations directly in all cases or by following the approach outlined in [DG02, DM03, RV16] (this relies on the irreducibility of certain tensor products as $U_q(\mathfrak{k})((z))$ -modules; otherwise the reflection equation must be verified directly).

First of all, the linear operator

$$K_\Pi(z) = \begin{pmatrix} \xi z^2 - 1 & 0 \\ 0 & \xi - z^2 \end{pmatrix} \in \text{End}(\mathbb{C}^2)[[z]] \tag{6.2}$$

is, up to a scalar, the unique solution of the $U_q(\mathfrak{k})$ -intertwining condition

$$K_\Pi(z)\Pi_z(u) = \Pi_{1/z}(u)K_\Pi(z) \quad \text{for all } u \in U_q(\mathfrak{k}). \tag{6.3}$$

By Theorem 3.6, it is proportional to the action of the grading-shifted universal K-matrix in the representation (Π, \mathbb{C}^2) .

Recall that $\Pi \circ \psi = \Pi$. Hence, motivated by Proposition 3.7, we consider the right reflection equation for $\pi \in \{\varrho, \bar{\varrho}, \nu, \phi\}$:

$$\mathcal{L}_\pi\left(\frac{y}{z}\right)\mathcal{K}_\pi(y)\mathcal{L}_\pi(yz)K_\Pi(z) = K_\Pi(z)\mathcal{L}_\pi(yz)\mathcal{K}_\pi(y)\mathcal{L}_\pi\left(\frac{y}{z}\right) \in \text{End}(W \otimes \mathbb{C}^2)[[y/z, z]]. \tag{6.4}$$

Lemma 6.1. *We have*

$$\begin{aligned} \mathcal{K}_\varrho(z) &= (-q^{-D}\xi)^D (q^2\xi^{-1}z^2; q^2)_D, & \mathcal{K}_{\bar{\varrho}}(z) &= (qz^2)^{-D} (q^2\xi^{-1}z^{-2}; q^2)_D^{-1}, \\ \mathcal{K}_\nu(z) &= z^{-2D} \frac{(q^{2-\mu}\xi^{-1}z^2; q^2)_D}{(q^{2-\mu}\xi^{-1}z^{-2}; q^2)_D}, & \mathcal{K}_\phi(z) &= (-q^{-\mu-D-1}\xi)^D. \end{aligned} \tag{6.5}$$

Note that these expressions were already given in [BT18] in different conventions. For completeness we sketch a proof relying on the universal K-matrix formalism.

Proof of Lemma 6.1. For $\mathcal{K}_\nu(z)$, by a straightforward check, the intertwining condition

$$\mathcal{K}_\nu(z)v_z(u) = v_{1/z}(u)\mathcal{K}_\nu(z) \quad \text{for all } u \in U_q(\mathfrak{k}) \tag{6.6}$$

can be solved to find $\mathcal{K}_\nu(z)$, making use of Proposition 3.8. Since $\mathcal{K}(z)$ commutes with the action of k_1 it follows that $\mathcal{K}_\nu(z) = f(D)$ for some $f \in \mathcal{F}$. Now imposing (6.6) for the generators $e_0 - q^{-1}\xi^{-1}k_0f_1$ and $e_1 - q^{-1}\xi k_1f_0$ yields the recurrence relation

$$\frac{f(D+1)}{f(D)} = \frac{1 - q^{2(D+1)-\mu}\xi^{-1}z^2}{z^2 - q^{2(D+1)-\mu}\xi^{-1}}.$$

In particular, the linear relation (6.6) has a 1-dimensional solution space. Together with the constraint $f(0) = 1$ it yields the formula given in (6.5).

For $\pi \in \{\varrho, \bar{\varrho}, \phi\}$, it is convenient to consider the linear space

$$\text{RE}_\pi := \{\mathcal{K}_\pi(y) \in \mathcal{F}(D) \mid (6.4) \text{ is satisfied}\} \tag{6.7}$$

and use Proposition 3.7 to find the explicit expression, relying on the second part of Theorem 3.6 for the fact that $\mathcal{K}_\pi(y)$ lies in $\mathcal{F}(D)$. Indeed, the operator $K_\varrho(z)$ was obtained in [VW20, Sec. 2.4] as the unique element of the 1-dimensional linear space RE_ϱ which fixes w_0 . In an analogous way we obtain the result for $K_{\bar{\varrho}}(z)$.

Note that ϕ is a reducible representation. Indeed, the solution space of (6.4) with $\pi = \phi$ is infinite-dimensional: the general solution $\mathcal{K}_\phi(z)$ is of the form $(-q^{-\mu-D-1} \xi)^D p$ with p in the centralizer of a in \mathcal{A} , i.e. a polynomial in a with coefficients in $\mathbb{C}[[z]]$. Since $\mathcal{K}_\phi(z) \in \mathcal{F}(D)$, p is a scalar. The requirement that w_0 is fixed forces $p = 1$. \square

6.2. Left K -operators. We now obtain linear-operator-valued power series satisfying a reflection equation for the left boundary by using a well-established bijection, see [Sk88, Eq. (15)], between its solution set and the solution set of the right reflection equation. For fixed $\xi \in \mathbb{C}^\times$ we define

$$\begin{aligned} \tilde{K}_\Pi(z) &:= (1 - q^2 \tilde{\xi}^{-1} z^2)^{-1} (1 - q^2 \tilde{\xi} z^2)^{-1} (K_\Pi(qz)^{-1}|_{\xi \mapsto \tilde{\xi}^{-1}}) \\ &= \begin{pmatrix} q^2 \tilde{\xi} z^2 - 1 & 0 \\ 0 & \tilde{\xi} - q^2 z^2 \end{pmatrix}. \end{aligned} \tag{6.8}$$

Also, for $\pi \in \{\varrho, \bar{\varrho}, \nu, \phi\}$ we define

$$\tilde{\mathcal{K}}_\pi(z) := \mathcal{K}_\pi(qz)^{-1}|_{\xi \mapsto \tilde{\xi}^{-1}}. \tag{6.9}$$

Similarly, note that $\mathcal{L}_\pi(\gamma z)$ is invertible in $\text{End}(W \otimes \mathbb{C}^2)[[z]]$ for all $\gamma \in \mathbb{C}$. We define

$$\tilde{\mathcal{L}}_\pi(z) = \mathcal{L}_\pi(q^2 z)^{-1}. \tag{6.10}$$

Lemma 6.2. *For all $\pi \in \{\varrho, \bar{\varrho}, \nu, \phi\}$ the left reflection equation holds:*

$$\tilde{\mathcal{K}}_\pi(y) \tilde{\mathcal{L}}_\pi(yz) \tilde{K}_\Pi(z) \mathcal{L}_\pi\left(\frac{y}{z}\right) = \mathcal{L}_\pi\left(\frac{y}{z}\right) \tilde{K}_\Pi(z) \tilde{\mathcal{L}}_\pi(yz) \tilde{\mathcal{K}}_\pi(y) \in \text{End}(W \otimes \mathbb{C}^2)[[y/z, z]]. \tag{6.11}$$

Proof. The desired equation (6.11) can be rewritten as

$$\tilde{K}_\Pi(z)^{-1} \tilde{\mathcal{L}}_\pi(yz)^{-1} \tilde{\mathcal{K}}_\pi(y)^{-1} \mathcal{L}_\pi\left(\frac{y}{z}\right) = \mathcal{L}_\pi\left(\frac{y}{z}\right) \tilde{\mathcal{K}}_\pi(y)^{-1} \tilde{\mathcal{L}}_\pi(yz)^{-1} \tilde{K}_\Pi(z)^{-1}.$$

By (6.8–6.10), this is equivalent to the right-reflection equation (6.4) with $y \mapsto qy$, $z \mapsto qz$ and $\xi \mapsto \tilde{\xi}^{-1}$. \square

Using the explicit formulas (6.2) and (6.4) we obtain that the solutions of the left reflection equations (6.9) are the following $\text{End}(W)$ -valued formal power series in z :

$$\begin{aligned} \tilde{\mathcal{K}}_\varrho(z) &= (-q^D \tilde{\xi})^D (q^4 \tilde{\xi} z^2; q^2)_D^{-1}, & \tilde{\mathcal{K}}_{\bar{\varrho}}(z) &= (q^3 z^2)^D (\tilde{\xi} z^{-2}; q^2)_D, \\ \tilde{\mathcal{K}}_\nu(z) &= (qz)^{2D} \frac{(q^{-\mu} \tilde{\xi} z^{-2}; q^2)_D}{(q^{4-\mu} \tilde{\xi} z^2; q^2)_D}, & \tilde{\mathcal{K}}_\phi(z) &= (-q^{\mu+D+1} \tilde{\xi})^D. \end{aligned} \tag{6.12}$$

7. Fusion Intertwiners Revisited

In this short intermezzo we explain how the universal K-matrix formalism naturally leads to relations involving K-operators and $U_q(\widehat{\mathfrak{b}^+})$ -intertwiners, called *fusion intertwiners*, which play a key role in the short exact sequence approach to the Q-operator. These intertwiners were discussed in [VW20] and the relevant relations with K-matrices were shown by a linear-algebraic computation relying on the explicit expressions of the various constituent factors, see [VW20, Lem. 3.2]. In other words, the representation-theoretic origin of these relations was unclear, which we now remedy.

Level-0 representations of $U_q(\widehat{\mathfrak{b}^+})$ are amenable to scalar modifications of the action of $U_q(\mathfrak{h}) = \langle k_1^{\pm 1} \rangle$, see also [HJ12, Rmk. 2.5]. In particular, for $r \in \mathbb{C}^\times$, define a modified Borel representation ϱ as follows:

$$\varrho_r(e_i) = \varrho(e_i), \quad \varrho_r(k_0) = r\varrho(k_0), \quad \varrho_r(k_1) = r^{-1}\varrho(k_1) \tag{7.1}$$

and consider the grading-shifted representation $\varrho_{r,z} := (\varrho_r)_z$. There exist $U_q(\widehat{\mathfrak{b}^+})$ -intertwiners

$$\begin{aligned} \iota(r) &: (\varrho_{qr,qz}, W) \rightarrow (\varrho_{r,z} \otimes \Pi_z, W \otimes \mathbb{C}^2), \\ \tau(r) &: (\varrho_{r,z} \otimes \Pi_z, W \otimes \mathbb{C}^2) \rightarrow (\varrho_{q^{-1}r,q^{-1}z}, W), \end{aligned}$$

called *fusion intertwiners*, which take part in the following short exact sequence:

$$0 \longrightarrow (\varrho_{qr,qz}, W) \xrightarrow{\iota(r)} (W \otimes \mathbb{C}^2, \varrho_{r,z} \otimes \pi_z) \xrightarrow{\tau(r)} (\varrho_{q^{-1}r,q^{-1}z}, W) \longrightarrow 0 \tag{7.2}$$

Explicitly¹¹, we have

$$\iota(r) = \begin{pmatrix} q^{-D} a^\dagger \\ -q^{D+1} r \end{pmatrix}, \quad \tau(r) = (q^D, q^{-D} r^{-1} a^\dagger). \tag{7.3}$$

Analogously to Theorem 5.2, fusion relations for the L-operators $\mathcal{L}(r, z)$, defined as suitable scalar multiples of $(\varrho_{r,z} \otimes \Pi)(\mathcal{R})$, now follow from these intertwining properties and the coproduct formulas for \mathcal{R} (2.16), see [VW20, Eqns. (3.8) and (3.9)].

Recalling the universal object \mathcal{K} and Theorem 3.6, we define the corresponding K-operator $\mathcal{K}_\varrho(r, z)$ as the unique scalar multiple of $\varrho_{r,z}(\mathcal{K})$ which fixes w_0 (cf. [VW20, Prop. 2.5]). Then

$$(\varrho_{r,z} \otimes \Pi_z)(\Delta(\mathcal{K})) \quad \propto \quad \mathcal{K}_\varrho(r, z)_1 \mathcal{L}(r, z^2) \mathcal{K}_\Pi(z)_2 \tag{7.4}$$

as a consequence of (3.19). Since \mathcal{K} lies in a completion of $U_q(\widehat{\mathfrak{b}^+})$, the intertwining properties of $\iota(r)$ and $\tau(r)$ now directly yield the following fusion relation for the K-operator:

$$\begin{aligned} \mathcal{K}_\varrho(r, z)_1 \mathcal{L}(r, z^2) \mathcal{K}_\Pi(z)_2 \iota(r) &\quad \propto \quad \iota(r) \mathcal{K}_\varrho(qr, qz) \\ \tau(r) \mathcal{K}_\varrho(r, z)_1 \mathcal{L}(r, z^2) \mathcal{K}_\Pi(z)_2 &\quad \propto \quad \mathcal{K}_\varrho(q^{-1}r, q^{-1}z) \tau(r), \end{aligned}$$


with the scalar factors determined by applying the two sides of the equation to w_0 , say. We will be able to prove a boundary counterpart of the factorization identity (5.7) using similar ideas.

¹¹ The sign mismatch with [VW20, Eq. (3.1)] is explained in Remark 4.3.


We recover, with a much smaller computational burden, the key result [VW20, Lemma 3.2] (a similar relation for left K-operators can easily be deduced from this, as explained in the last sentence of [VW20, Proof of Lemma 3.2]). In the approach to Baxter’s Q-operator using short exact sequences, the fusion relations for L and K-operators induce fusion relations for 2-boundary monodromy operators, see [VW20, Lem. 4.2] from which Baxter’s relation (1.1) follows by taking traces, see [VW20, Sec. 5.2].

8. Boundary Factorization Identity

In motivating and presenting the key boundary relations, it is very useful to introduce a graphical representation of spaces and operators. Let us introduce the following pictures for the different representations from Sects. 4 and 5:



For any vector spaces V, V' , denote by \mathcal{P} the linear map from $V \otimes V'$ to $V' \otimes V$ such that $\mathcal{P}(v \otimes v') = v' \otimes v$ for all $v \in V, v' \in V'$. Also set $z = z_1/z_2$. We then have the following pictures for L-operators and R-operators:




We now make the following definitions¹²:

$$\tilde{\mathcal{R}}_{\varrho\bar{\varrho}}(z) := \mathcal{R}_{\varrho\bar{\varrho}}(q^2z)^{-1}, \quad \tilde{\mathcal{R}}_{\nu\phi}(z) := \mathcal{R}_{\nu\phi}(q^2z)^{-1}, \quad (8.1)$$


and represent these modified R-matrices by the following pictures:




The various right-boundary K-matrices are represented as follows:



The left-boundary K-matrices defined in Sect. 6.2 are represented by the natural analogues of these pictures. For example:




Making use of these pictures, we see that Theorem 5.2 and Corollary 5.3 are represented by



For the compatibility with the right boundary we claim that

¹² These are the modified forms of the R-matrices that appear in the corresponding left reflection equations, see [Sk88, Eq. (13)].



which corresponds to the following identity in $\mathcal{A}^{(2)}$:

$$\mathcal{K}_v(z)_1 \mathcal{R}_{v\phi}(z^2) \mathcal{K}_\phi(z)_2 \mathcal{O} = \mathcal{O} \mathcal{K}_\rho(q^{-\mu/2}z)_1 \mathcal{R}_{\rho\bar{\rho}}(z^2) \mathcal{K}_{\bar{\rho}}(q^{\mu/2}z)_2, \quad (8.2)$$

which we call the *right boundary factorization identity*. The diagrams above serve as a motivation for the identity, which we now prove using results from Sect. 3 (an alternative computational proof of Theorem 8.1 is given in Appendix C).

Theorem 8.1. *For all $\mu \in \mathbb{C}$, all $q \in \mathbb{C}^\times$ not a root of unity and all $\xi \in \mathbb{C}^\times$, relation (8.2) is satisfied.*

Proof. The proof is analogous to the proof of Theorem 5.2. We first note that


$$\begin{aligned} (\varrho_{q^{-\mu/2}z} \otimes \bar{\varrho}_{q^{\mu/2}z})(\text{id} \otimes \psi)(\mathcal{R}) &= (\varrho_{q^{-\mu/2}z} \otimes \bar{\varrho}_{q^{-\mu/2}z^{-1}})(\mathcal{R}) \quad \propto \mathcal{R}_{\rho\bar{\rho}}(z^2), \\ (v_z \otimes \phi_z)(\text{id} \otimes \psi)(\mathcal{R}) &= (v_z \otimes \phi_{z^{-1}})(\mathcal{R}) \quad \propto \mathcal{R}_{v\phi}(z^2). \end{aligned}$$

Noting the coproduct formula (3.19), we obtain

$$\begin{aligned} \mathcal{K}_\rho(q^{-\mu/2}z)_1 \mathcal{R}_{\rho\bar{\rho}}(z^2) \mathcal{K}_{\bar{\rho}}(q^{\mu/2}z)_2 &\propto (\varrho_{q^{-\mu/2}z} \otimes \bar{\varrho}_{q^{\mu/2}z})(\Delta(\mathcal{K})), \\ \mathcal{K}_v(z)_1 \mathcal{R}_{v\phi}(z^2) \mathcal{K}_\phi(z)_2 &\propto (v_z \otimes \phi_z)(\Delta(\mathcal{K})). \end{aligned}$$

Now Theorem 4.4 implies (8.2) up to a scalar. The fact that all factors fix $w_0 \otimes w_0$ shows that the scalar is 1. \square

Compatibility with the left boundary requires that



The identity in $\mathcal{A}^{(2)}$ corresponding to this is

$$\tilde{\mathcal{K}}_{\bar{q}}(q^{\mu/2}z, \tilde{\xi})_2 \tilde{\mathcal{R}}_{\bar{q}\bar{q}}(z^2) \tilde{\mathcal{K}}_{\bar{q}}(q^{-\mu/2}z, \tilde{\xi})_1 \mathcal{O}^{-1} = \mathcal{O}^{-1} \tilde{\mathcal{K}}_{\phi}(z, \tilde{\xi})_2 \tilde{\mathcal{R}}_{\nu\phi}(z^2) \tilde{\mathcal{K}}_{\nu}(z, \tilde{\xi})_1. \tag{8.3}$$

Theorem 8.2. *Relation (8.3) is satisfied.*

Proof. Given the definitions (6.12) and (8.1), this follows straightforwardly by inverting (8.2) and replacing $(z, \xi) \mapsto (qz, \tilde{\xi}^{-1})$. □

9. Discussion

The main result of this paper is Theorem 8.1 which can be viewed as a boundary analogue of Theorem 5.2. To establish this result, first we needed to show that all R and K-operators involved in equation (8.2) are well-defined actions of the universal elements \mathcal{R} and \mathcal{K} on the infinite-dimensional $U_q(\widehat{\mathfrak{b}}^+)$ -modules involved. The key fact that allows for this is that \mathcal{R} and \mathcal{K} live in completions of $U_q(\widehat{\mathfrak{b}}^+) \otimes U_q(\widehat{\mathfrak{b}}^-)$ and of $U_q(\widehat{\mathfrak{b}}^+)$, respectively. This is very familiar for \mathcal{R} but for \mathcal{K} relies on the recent works [AV22a, AV22b]. Introducing the $U_q(\widehat{\mathfrak{b}}^+)$ -intertwiner \mathcal{O} and the formula for $\Delta(\mathcal{K})$ given by (3.19), relation (8.2) follows immediately from the intertwining property of \mathcal{O} .

The open Q-operator $\mathcal{Q}(z)$ of [VW20] is the trace of a product of R and K-operators over the $U_q(\widehat{\mathfrak{b}}^+)$ -module (ϱ_z, W) and there is a similar construction of an open Q-operator $\bar{\mathcal{Q}}(z)$. In a future paper, the authors will present this construction and the use of Theorem 8.2 in deriving a boundary analogue of the factorization relation $\mathcal{T}_{\mu}(z) \propto \mathcal{Q}(zq^{-\mu/2})\bar{\mathcal{Q}}(zq^{\mu/2})$. They will also develop the analogous theory for different coideal subalgebras, in particular those for which non-diagonal solutions of the reflection equation are intertwiners. There is a quite subtle rational degeneration of the construction in the present paper. The first-named author will study this in a separate paper, giving an alternative approach to Q-operators for the open XXX spin chain, cf. [FS15].

Acknowledgments B.V. would like to thank A. Appel, P. Baseilhac and N. Reshetikhin for useful discussions. This research was supported in part by funding from EPSRC grant EP/R009465/1, from the Simons Foundation and the Centre de Recherches Mathématiques (CRM), through the Simons-CRM scholar-in-residence programme, and by the Galileo Galilei Institute (GGI) scientific programme on ‘Randomness, Integrability and Universality’. R.W. would like to acknowledge and thank CRM and the GGI for their hospitality and support. A.C. is grateful for support from EPSRC DTP award EP/M507866/1.

Data Availability Data sharing is not applicable to this article as no datasets were generated or analysed during the current study.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Publisher’s Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

A. Deformed Pochhammer Symbols and Exponentials

This appendix is independent from the main text, but provides identities which are used there. We review some basic theory of deformed Pochhammer symbols and exponentials (as formal power series) with a deformation parameter $p \in \mathbb{C}^\times$, which corresponds to q^2 in the main text.

A.1. Deformed Pochhammer symbols. Let x be a formal variable. For $n \in \mathbb{Z}$, the (finite) deformed Pochhammer symbol $(x; p)_n \in \mathbb{C}[[x]]$ is defined by

$$(x; p)_n := \begin{cases} \prod_{m=0}^{n-1} (1 - xp^m) & \text{if } n \geq 0, \\ \prod_{m=n}^{-1} (1 - xp^m)^{-1} & \text{if } n < 0 \end{cases} \tag{A.1}$$

(the definition for $n < 0$ is understood as a product of geometric series); since its constant coefficient is nonzero, it is invertible. For all $p \in \mathbb{C}^\times$ and $n \in \mathbb{Z}_{\geq 0}$ we have the following basic identity in $\mathbb{C}[[x]]$, see [GR90, (I.2), (I.3)]:

$$(x; p)_{-n} = (p^{-n}x; p)_n^{-1} = (x/p; p^{-1})_n^{-1} = (-x)^{-n} p^{n(n+1)/2} (p/x; p)_n^{-1}. \tag{A.2}$$

Assuming $|p| < 1$, the infinite deformed Pochhammer symbol

$$(x; p)_\infty := \prod_{m=0}^\infty (1 - xp^m) \tag{A.3}$$

is an invertible formal power series with well-defined coefficients in \mathbb{C} . The following identity holds in $\mathbb{C}[[x]]$, see [GR90, (I.5)]:

$$(x; p)_n = \frac{(x; p)_\infty}{(p^n x; p)_\infty}. \tag{A.4}$$

A.2. Deformed exponentials. From now on we assume that p is not a root of unity. In particular, $(p; p)_k \neq 0$ for all $k \in \mathbb{Z}_{\geq 0}$. The *deformed exponential* is the invertible formal power series

$$e_p(x) := {}_1\phi_0(0; -; p, x) = \sum_{k=0}^\infty \frac{x^k}{(p; p)_k}. \tag{A.5}$$

The ordinary exponential formal power series arises as the termwise limit $\lim_{p \rightarrow 1} e_p((1 - p)x) = e^x$. This series satisfies the functional relation

$$e_p(px) = (1 - x)e_p(x), \tag{A.6}$$

see [GR90, Sec. 1.3]. Since constants are the only formal power series which are invariant under $x \mapsto px$, an inspection of constant coefficients shows that (A.6) implies

$$e_p(x) = \frac{1}{(x; p)_\infty} \quad \text{if } |p| < 1. \tag{A.7}$$

Similarly we consider the invertible formal power series

$$E_p(x) := {}_0\phi_0(-; -; p, -x) = \sum_{k=0}^{\infty} \frac{p^{k(k-1)/2} x^k}{(p; p)_k}. \tag{A.8}$$

Then $E_p(-x)^{-1}$ also satisfies (A.6) and by comparing constant coefficients again we deduce $e_p(x) = E_p(-x)^{-1}$. By evaluating (A.2) at $x = 1$, we obtain $E_p(-x) = e_{p^{-1}}(p^{-1}x)$ and hence

$$e_p(x) = e_{p^{-1}}(p^{-1}x)^{-1} \in \mathbb{C}[[x]]. \tag{A.9}$$

Deformed exponentials in x and y satisfy various useful identities in particular deformations of the commutative algebra $\mathbb{C}[[x, y]]$. For instance, in any algebra generated by the symbols x and y such that $yx = \gamma xy$ for $\gamma \in \mathbb{C}$, the definition implies the following identity:

$$ye_p(x) = e_p(\gamma x)y \tag{A.10}$$

which we will use repeatedly. For a survey of product formulas analogous to $\exp(x)\exp(y) = \exp(x+y)$, see [Ko97]. We will need the following result.

Lemma A.1. *Let x, y be elements of an algebra such that $yx = px y$. The following identities hold as formal power series in x, y :*

$$e_p(x)e_p(y) = e_p(x+y), \tag{A.11}$$

$$\begin{aligned} e_p(y)e_p(x) &= e_p(x(1-y))e_p(y) \\ &= e_p(x)e_p(-xy)e_p(y) = e_p(x)e_p((1-x)y). \end{aligned} \tag{A.12}$$

Proof. (A.11) is a direct consequence of the well-known q-binomial formula, see e.g. [GR90, Ex. 1.35]. For (A.12) see [Ko97, Prop. 3.2]. \square

A.3. Deformed exponentials as linear maps. Let V be a \mathbb{C} -linear space. Call an operator f on V *locally nilpotent* if for all $v \in V$ there exists $o(v) \in \mathbb{Z}_{\geq 0}$ such that $f^{o(v)}(v) = 0$ (note that nilpotent operators are locally nilpotent and if V is finite-dimensional the converse is true). If f is nilpotent, the deformed exponential $e_p(f)$ defines an invertible map on V . If additionally y is an indeterminate then $e_p(yf)$ is a well-defined invertible element of $\text{End}(V)[[y]]$.

In the case $V = W \otimes W$ the following commutation relations for linear-operator valued formal series are satisfied, expressed in terms of the linear operators $a, a^\dagger, \bar{a}^\dagger, f(D)$ ($f \in \mathcal{F}$) on W defined in Sect. 4.2.

Lemma A.2. *Let y be a formal variable. In $\text{End}(W \otimes W)[[y]]$ the following identities hold:*

$$[e_p(ya_1\bar{a}_2^\dagger), f(D_1 + D_2)] = [e_p(ya_1\bar{a}_2^\dagger), a_1] = [e_p(ya_1\bar{a}_2^\dagger), \bar{a}_2^\dagger] = 0 \tag{A.13}$$

for all $f \in \mathcal{F}$ and

$$[e_p(ya_1\bar{a}_2^\dagger), a_1^\dagger] = yp^{D_1}\bar{a}_2^\dagger e_p(ya_1\bar{a}_2^\dagger), \tag{A.14}$$

$$[e_p(ya_1\bar{a}_2^\dagger), p^{-D_1}a_2] = ye_p(ya_1\bar{a}_2^\dagger)a_1p^{-D_1}. \tag{A.15}$$

Proof. Note that (A.13) follows directly from the definition of the deformed exponential. A straightforward inductive argument using (4.4) yields

$$[a^{k+1}, a^\dagger] = (1 - p^{k+1})p^D a^k, \tag{A.16}$$

$$[(\bar{a}^\dagger)^{k+1}, a]_{p^{k+1}} = (1 - p^{k+1})(\bar{a}^\dagger)^k, \tag{A.17}$$

for all $k \in \mathbb{Z}_{\geq 0}$, which imply (A.14) and (A.15), respectively. \square

B. Explicit Expressions for R-Operators

In this appendix we derive explicit formulas for $\mathcal{R}_{q\bar{q}}(z)$ and $\mathcal{R}_{v\phi}(z)$, defined by (5.10) as images of the universal R-matrix \mathcal{R} fixing $w_0 \otimes w_0$. We expect that these will be useful in further studies of Baxter’s Q-operators for the open XXZ spin chain; for now they will allow us to give a proof of the boundary factorization identity which does not rely on the universal K-matrix formalism. First we note that, by the second part of Theorem 2.4, $\mathcal{R}_{q\bar{q}}(z)$ and $\mathcal{R}_{v\phi}(z)$ lie in the centralizer

$$\mathcal{A}_0^{(2)} := \left\{ X \in \mathcal{A}^{(2)} \mid [X, q^{D_1+D_2}] = 0 \right\}. \tag{B.1}$$

One straightforwardly verifies that $\mathcal{A}_0^{(2)}$ is generated by elements of the form

$$\sum_{k \geq 0} (\bar{a}_2^\dagger)^k f_k(D_1, D_2) a_1^k, \quad \sum_{k \geq 0} (a_1^\dagger)^k f_k(D_1, D_2) a_2^k, \quad f_k \in \mathcal{F}^{(2)}. \tag{B.2}$$

Hence, elements of $\mathcal{A}_0^{(2)}$ in fact commute with all elements of the form $f(D_1 + D_2)$ ($f \in \mathcal{F}$).

B.1. Explicit expression for $\mathcal{R}_{v\phi}(z)$. We first state and prove an explicit formula for $\mathcal{R}_{v\phi}(z)$. We keep using the shorthand notation $p = q^2$.

Theorem B.1. *For all $z \in \mathbb{C}$ we have*

$$\mathcal{R}_{v\phi}(z) = e_p(z a_1^\dagger a_2) q^{(\mu-1)(D_2-D_1)-2D_1(D_2+1)}. \tag{B.3}$$

Proof. From Proposition 2.6 we deduce that $\mathcal{R}_{v\phi}(z)$ is a solution of the linear relation

$$X(v_z \otimes \phi^-)(\Delta(u)) = (v_z \otimes \phi^-)(\Delta^{\text{op}}(u))X \quad \text{for all } u \in U_q(\widehat{\mathfrak{b}}^-). \tag{B.4}$$

First of all, note that the element in the right-hand side of (B.3) satisfies (B.4) with $u \in \{k_0, k_1\}$ and so it suffices to prove that the vector space

$$\mathcal{X} = \left\{ X \in \mathcal{A}_0^{(2)} \mid X \text{ satisfies (B.4) for } u \in \{f_0, f_1\} \right\} \tag{B.5}$$

is spanned by $e_p(z^2 a_1^\dagger a_2) q^{(\mu-1)(D_2-D_1)-2D_1(D_2+1)}$. Using the explicit formulas (2.2), (4.8) and (4.16), we obtain that (B.4) is equivalent to the system

$$\begin{aligned} & X \left(z^{-1} a_1 (q^{-\mu} - q^{\mu-2D_1}) q^{-\mu-2D_2-1} + q^{-1} a_2 \right) \\ &= \left(z^{-1} a_1 (q^{-\mu} - q^{\mu-2D_1}) + q^{\mu-2(D_1+1)} a_2 \right) X, \\ & X a_1^\dagger q^{\mu+1+2D_2} = a_1^\dagger X. \end{aligned}$$

Without loss of generality we may write $X = \tilde{X}q^{(\mu-1)(D_2-D_1)-2D_1(D_2+1)}$ with $\tilde{X} \in \mathcal{A}_0^{(2)}$. Hence (B.4) is equivalent to

$$z^{-1}[\tilde{X}, a_1(1 - p^{\mu-D_1})] = p^{\mu-D_1-1}a_2\tilde{X} - \tilde{X}p^{D_1}a_2, \quad [\tilde{X}, a_1^\dagger] = 0. \quad (\text{B.6})$$

It is straightforward to check that the centralizer in $\mathcal{A}_0^{(2)}$ of a_1^\dagger is the subalgebra generated by elements of the form $\sum_{k \geq 0} (a_1^\dagger)^k f_k(D_2)a_2^k$ with $f_k \in \mathcal{F}$. It follows that \tilde{X} is of this form. Therefore (B.4) is equivalent to the single equation

$$\begin{aligned} & \sum_{k \geq 0} [(a_1^\dagger)^k, a_1(1 - p^{\mu-D_1})] f_k(D_2)a_2^k = \\ & = z \sum_{k \geq 0} (a_1^\dagger)^k (p^{\mu-D_1-k-1} f_k(D_2+1) - p^{D_1} f_k(D_2)) a_2^{k+1}. \end{aligned}$$

The commutator vanishes if $k = 0$ so in the left-hand side we replace k by $k + 1$. For $k \geq 0$ we have

$$[(a^\dagger)^{k+1}, a(1 - p^{\mu-D})] = (a^\dagger)^k (1 - p^{k+1})(p^{\mu-D-k-1} - p^D).$$

Hence (B.4) is equivalent to the recurrence relation

$$(1 - p^{k+1})(p^{\mu-D_1-k-1} - p^{D_1}) f_{k+1}(D_2) = z(p^{\mu-D_1-k-1} f_k(D_2+1) - p^{D_1} f_k(D_2)).$$

Viewing $\mathcal{F}^{(2)}(D_1, D_2)$ as an $\mathcal{F}(D_2)$ -module, the elements $p^{\pm D_1}$ are linearly independent. Hence the above recurrence relation is equivalent to the system

$$(1 - p^{k+1}) f_{k+1}(D) = z f_k(D+1), \quad f_k(D+1) = f_k(D).$$

This is in turn equivalent to $f_k(D) \in (p; p)_k^{-1} z^k \mathbb{C}$ for $k \in \mathbb{Z}_{>0}$, as required. \square

B.2. The automorphism χ and the q -oscillator subalgebra $\tilde{\mathcal{A}}$. To obtain an expression for $R_{\nu\phi}(z)$ in terms of deformed exponentials, it is very convenient to point out an additional automorphism χ . It cannot be defined on all of \mathcal{A} so we will consider a subalgebra $\tilde{\mathcal{A}}$. First, consider the subalgebra $\tilde{\mathcal{F}}(D) \subset \mathcal{F}(D)$ generated by

$$p^{\pm D(D+1)/2}, \quad \gamma^D, \quad (p\tilde{\gamma}; p)_D^{\pm 1}, \quad (p\gamma z^2; p)_D, \quad (-\gamma z^2)^{-D} (p\gamma^{-1} z^{-2}; p)_D^{-1}$$

for all $\gamma \in \mathbb{C}^\times$ and $\tilde{\gamma} \in \mathbb{C}^\times \setminus p^\mathbb{Z}$.

For elements of $\tilde{\mathcal{F}}(D)$, unlike general elements of $\mathcal{F}(D)$, the symbol D can be formally evaluated at negative integers. Accordingly, we define an involutive automorphism χ of $\tilde{\mathcal{F}}(D)$ accomplishing the formal replacement $D \mapsto -D - 1$. To be more precise, we set

$$\begin{aligned} \chi(p^{\pm D(D+1)/2}) &= p^{\pm D(D+1)/2}, & \chi(\gamma^D) &= \gamma^{-D-1}, \\ \chi((p\tilde{\gamma}; p)_D^{\pm 1}) &= (1 - \tilde{\gamma})^{\mp 1} p^{\pm D(D+1)/2} (-\tilde{\gamma})^{\mp D} (p\tilde{\gamma}^{-1}; p)_D^{\mp 1}, \\ \chi((p\gamma z^2; p)_D) &= (1 - \gamma z^2)^{-1} p^{D(D+1)/2} (-\gamma z^2)^{-D} (p\gamma^{-1} z^{-2}; p)_D^{-1}, \\ \chi((-\gamma z^2)^{-D} (p\gamma^{-1} z^{-2}; p)_D^{-1}) &= (1 - \gamma z^2) p^{-D(D+1)/2} (p\gamma z^2; p)_D. \end{aligned} \quad (\text{B.7})$$

We denote the subalgebra of $\text{End}(W)$ generated by a^\dagger , a and $\tilde{\mathcal{F}}(D)$ by $\tilde{\mathcal{A}}$. It is straightforward to check that χ extends to a (non-involutive) algebra automorphism of $\tilde{\mathcal{A}}$ by means of the assignments

$$\chi(a) = \tilde{a}^\dagger, \quad \chi(a^\dagger) = a. \tag{B.8}$$

We can formulate a completion of the tensor product $\tilde{\mathcal{A}} \otimes \tilde{\mathcal{A}}$ in a similar way as for $\mathcal{A} \otimes \mathcal{A}$. More precisely, we consider the subalgebra $\tilde{\mathcal{F}}^{(2)}$ of $\mathcal{F}^{(2)}$ generated by the subsets $\tilde{\mathcal{F}}(D_1)$, $\tilde{\mathcal{F}}(D_2)$ and the special elements $p^{\pm D_1(D_2+1)}$. The completed tensorial square of $\tilde{\mathcal{A}}$ is defined to be the subalgebra $\tilde{\mathcal{A}}^{(2)}$ of $\text{End}(W \otimes W)$ generated by the elements (4.7) with $g_{k,\ell}$, $h_{k,\ell} \in \tilde{\mathcal{F}}^{(2)}$. Note that the boundary factorization identity (8.2) is an identity in the subalgebra $\tilde{\mathcal{A}}^{(2)} \subset \text{End}(W \otimes W)[[z]]$.

The automorphism

$$\chi^{(2)} := \sigma \circ (\chi \otimes \chi^{-1}) \tag{B.9}$$

of $\tilde{\mathcal{A}} \otimes \tilde{\mathcal{A}}$ naturally extends to an automorphism of $\tilde{\mathcal{A}}^{(2)}$, fixing $p^{\pm D_1(D_2+1)}$ and acting termwise on power series in locally nilpotent operators.

Remark B.2. The map χ can be seen as an infinite-dimensional version of conjugation by anti-diagonal matrices; certain $U_q(\mathfrak{b}^+)$ -representations are naturally related this way. For instance, for the 2-dimensional representation Π , note that $\text{Ad}(J) \circ \Pi = \Pi \circ \Phi$ where Ad denotes ‘conjugation by’ and $J = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$. In the same way, χ relates the prefundamental representations ϱ and $\bar{\varrho}$ up to a twist by the diagram automorphism Φ : $\chi \circ \varrho = \bar{\varrho} \circ \Phi$. Hence, the condition (2.19) and the 1-dimensionality of the solution space of the relevant linear equation implies $(\text{Ad}(J) \otimes \chi)(\mathcal{L}_\varrho(z)) = \mathcal{L}_{\bar{\varrho}}(z)$. At the same time, a suitable scalar multiple of $\mathcal{R}_{\Pi \Pi}(z)$, i.e. the R-matrix for the XXZ chain, is fixed by $\text{Ad}(J \otimes J)$ and we will see in Sect. B.3 that the same statement is true for $\mathcal{R}_{\varrho \bar{\varrho}}(z)$ and $\chi^{(2)}$.

From (3.5) it follows that $\Phi(U_q(\mathfrak{k})) = U_q(\mathfrak{k})|_{\xi \mapsto \xi^{-1}}$. Hence, the boundary counterparts of the above relations also involve inversion of the free parameter ξ :

$$\text{Ad}(J)(\mathcal{K}_\Pi(z))|_{\xi \mapsto \xi^{-1}} = -\xi \mathcal{K}_\Pi(z), \quad \chi(\mathcal{K}_\varrho(z))|_{\xi \mapsto \xi^{-1}} = q^{-1}(z^2 - \xi^{-1})^{-1} \mathcal{K}_{\bar{\varrho}}(z).$$

In fact, applying $\chi \otimes \text{Ad}(J)$ to the reflection equation (6.4) with $\pi = \varrho$ and inverting ξ we see that

$$\mathcal{K}_\varrho(z) \mapsto \chi(\mathcal{K}_\varrho(z))|_{\xi \mapsto \xi^{-1}}$$

defines a bijection: $\text{RE}_\varrho \rightarrow \text{RE}_{\bar{\varrho}}$ of the solution spaces defined in (6.7). ∅

We can use the map $\chi^{(2)}$ to generate further relations similar to those in Lemma A.2.

Lemma B.3. *Let y be a formal parameter. In $\text{End}(W \otimes W)[[y]]$ the following identities hold:*

$$[\tilde{a}_2^\dagger, e_p(ya_1^\dagger a_2)] = ye_p(ya_1^\dagger a_2)a_1^\dagger p^{-D_2-1}, \tag{B.10}$$

$$[\tilde{a}_1^\dagger a_2, e_p(ya_1 \tilde{a}_2^\dagger)] = y(e_p(ya_1 \tilde{a}_2^\dagger)p^{-D_1-1} - p^{-D_2-1}e_p(ya_1 \tilde{a}_2^\dagger)). \tag{B.11}$$

Proof. In this proof we view the algebra \mathcal{A} as a subalgebra of $\text{End}(W)[[y]]$, and similarly for $\mathcal{A}^{(2)}$. To prove (B.10), first we apply $\chi^{(2)}$ to (A.14), obtaining

$$[e_p(ya_1\bar{a}_2^\dagger), a_2] = ya_1p^{-D_2-1}e_p(ya_1\bar{a}_2^\dagger). \tag{B.12}$$

Now consider the unique involutive algebra anti-automorphism $\eta : \mathcal{A} \rightarrow \mathcal{A}$ which exchanges a and a^\dagger and fixes $f(D)$ for all $f \in \mathcal{F}$ and the unique involutive algebra anti-automorphism $\bar{\eta} : \mathcal{A} \rightarrow \mathcal{A}$ which exchanges a and \bar{a}^\dagger and fixes $f(D)$ for all $f \in \mathcal{F}$. Then $\eta^{(2)} := \eta \otimes \bar{\eta}$ is an algebra antiautomorphism of $\mathcal{A} \otimes \mathcal{A}$. It extends in a natural way to an algebra antiautomorphism of $\mathcal{A}^{(2)}$. By applying $\eta^{(2)}$ to (B.12) we obtain (B.10).

Finally, to prove (B.11), upon right-multiplying (A.15) by $p^{D_1+D_2+1}$ we obtain

$$[e_p(ya_1\bar{a}_2^\dagger), a_1p^{D_2}] = ye_p(ya_1\bar{a}_2^\dagger)a_1p^{D_2}. \tag{B.13}$$

From (A.14) and (B.13) it follows that

$$\begin{aligned} [e_p(ya_1\bar{a}_2^\dagger), a_1^\dagger a_2 p^{D_2}] &= y(\bar{a}_2^\dagger p^{D_1} e_p(ya_1\bar{a}_2^\dagger) a_2 + a_1^\dagger e_p(ya_1\bar{a}_2^\dagger) a_1) p^{D_2} \\ &= y(p^{D_1} e_p(ya_1\bar{a}_2^\dagger)(p^{D_2} - 1) + (1 - p^{D_1}) e_p(ya_1\bar{a}_2^\dagger) p^{D_2}) \\ &= y(e_p(ya_1\bar{a}_2^\dagger) p^{D_2} - p^{D_1} e_p(ya_1\bar{a}_2^\dagger)). \end{aligned} \tag{B.14}$$

Now (B.11) follows as the $\chi^{(2)}$ -image of (B.14). □

B.3. Explicit expression for $\mathcal{R}_{\varrho\bar{\varrho}}(z)$. To aid the computation of $\mathcal{R}_{\varrho\bar{\varrho}}(z)$, consider the subalgebra $\tilde{\mathcal{A}}_0^{(2)} = \tilde{\mathcal{A}}^{(2)} \cap \mathcal{A}_0^{(2)}$, which is preserved by $\chi^{(2)}$.

Lemma B.4. $\mathcal{R}_{\varrho\bar{\varrho}}(z)$ is a $\tilde{\mathcal{A}}_0^{(2)}$ -valued formal power series whose coefficients are fixed by $\chi^{(2)}$.

Proof. It is clear from (4.10) and (4.16) that $\varrho \otimes \bar{\varrho}^-$ takes values in $\tilde{\mathcal{A}} \otimes \tilde{\mathcal{A}} \subset \tilde{\mathcal{A}}^{(2)}$. Now recall (2.20) and note that the factor κ acts as $p^{D_1(D_2+1)}$. Furthermore, noting the form of $(\Sigma_z \otimes \text{id})(\Theta)$ given by (2.26) with the components Θ_λ lying in $U_q(\hat{\mathfrak{n}}^+)_\lambda \otimes U_q(\hat{\mathfrak{n}}^-)_{-\lambda}$ ($\lambda \in \hat{Q}^+$), we obtain that the action of $\mathcal{R}(z)$ on $(\varrho \otimes \bar{\varrho}^-, W \otimes W)$ is by an element of $\tilde{\mathcal{A}}_0^{(2)}$. For the second part, note that

$$\chi^{(2)} \circ (\varrho \otimes \bar{\varrho}^-) = (\chi^{-1} \otimes \chi) \circ (\bar{\varrho}^- \otimes \varrho) \circ \sigma = (\varrho \otimes \bar{\varrho}^-) \circ (\omega \otimes \omega) \circ \sigma.$$

Applying this to $\mathcal{R}(z)$, making use of (2.27), (2.24) and (2.18), we obtain $\chi^{(2)}(\mathcal{R}_{\varrho\bar{\varrho}}(z)) = \mathcal{R}_{\varrho\bar{\varrho}}(z)$. □

In the derivation of the formula for $\mathcal{R}_{\varrho\bar{\varrho}}(z)$, we rely on the following result.

Lemma B.5. The centralizer of the subset $\{a_1^\dagger, \bar{a}_2^\dagger\}$ in $\mathcal{A}^{(2)}$ is equal to $\mathbb{C}[[z]]$.

Proof. This centralizer is the intersection of the centralizer of a_1^\dagger and the centralizer of \bar{a}_2^\dagger , which are easily found to be equal to

$$\left\{ \sum_{k,\ell \geq 0} (a_1^\dagger)^k f_{k,\ell}(D_2) a_2^\ell \mid f_{k,\ell} \in \mathcal{F} \right\}, \quad \left\{ \sum_{k,\ell \geq 0} (\bar{a}_2^\dagger)^k g_{k,\ell}(D_1) a_1^\ell \mid g_{k,\ell} \in \mathcal{F} \right\},$$

respectively. Clearly their intersection is trivial. □

Now we are ready to state and prove a formula for $\mathcal{R}_{\varrho\bar{\varrho}}(z)$ in terms of deformed exponentials.

Theorem B.6. *For all z we have*

$$\mathcal{R}_{\varrho\bar{\varrho}}(z) = e_{q^2}(q^3 z a_1 \bar{a}_2^\dagger) e_{q^2}(q^{-1} z a_1^\dagger a_2) q^{-2D_1(D_2+1)}. \tag{B.15}$$

Proof. Clearly, $w_0 \otimes w_0$ is fixed by the expression on the right-hand side of (B.15). In the following we initially work over the ring $\mathbb{C}[[z, z_2]]$ for some new indeterminate z_2 and write $z_1 = z z_2$. By applying $\varrho_{z_1} \otimes \Pi_1 \otimes \bar{\varrho}_{z_2}^-$ to (2.17) and left and right-multiplying by $\mathcal{L}_{\bar{\varrho}, 23}^-(z_2^{-1})^{-1}$ we obtain

$$\mathcal{R}_{\varrho\bar{\varrho}}(z)_{12} \mathcal{L}_\varrho(z_1)_{13} \mathcal{L}_{\bar{\varrho}}^-(z_2^{-1})_{32}^{-1} = \mathcal{L}_{\bar{\varrho}}^-(z_2^{-1})_{32}^{-1} \mathcal{L}_\varrho(z_1)_{13} \mathcal{R}_{\varrho\bar{\varrho}}(z)_{12} \tag{B.16}$$

an equation in $(\tilde{\mathcal{A}}^{(2)} \otimes \text{End}(\mathbb{C}^2))[[z_2]]$. By a direct computation we obtain

$$\mathcal{L}_{\bar{\varrho}}^-(z_2^{-1})^{-1} = \frac{1}{z_2^2 - 1} \begin{pmatrix} q^{-D-1} z_2^2 & \bar{a}^\dagger q^{-D-1} z_2 \\ a q^{D-1} z_2 & q^{D+1} z_2^2 - q^{-D-1} \end{pmatrix} \in \text{End}(\mathbb{C}^2) \otimes \tilde{\mathcal{A}}. \tag{B.17}$$

Now we consider the equation

$$(z_2^2 - 1) X_{12} \mathcal{L}_\varrho(z_1)_{13} \mathcal{L}_{\bar{\varrho}}^-(z_2^{-1})_{32}^{-1} = (z_2^2 - 1) \mathcal{L}_{\bar{\varrho}}^-(z_2^{-1})_{32}^{-1} \mathcal{L}_\varrho(z_1)_{13} X_{12} \tag{B.18}$$

in $(\tilde{\mathcal{A}}^{(2)} \otimes \text{End}(\mathbb{C}^2))[[z_2]]$, for some $X \in \tilde{\mathcal{A}}_0^{(2)}$ such that $\chi^{(2)}(X) = X$. It suffices to prove that

$$\mathcal{X} = \left\{ X \in \tilde{\mathcal{A}}_0^{(2)} \mid X \text{ satisfies (B.18) and is fixed by } \chi^{(2)} \right\}, \tag{B.19}$$

which by Lemma B.4 contains $(\varrho_z \otimes \bar{\varrho}^-)(\mathcal{R})$, is spanned by the element given in the right-hand side of (B.15).

By considering explicit expressions for $(z_2^2 - 1) \mathcal{L}_\varrho(z_1)_{13} \mathcal{L}_{\bar{\varrho}}^-(z_2^{-1})_{32}^{-1}$ and $(z_2^2 - 1) \mathcal{L}_{\bar{\varrho}}^-(z_2^{-1})_{32}^{-1} \mathcal{L}_\varrho(z_1)_{13}$, we obtain that (B.18) amounts to the system

$$\begin{aligned} X(q^{D_1-D_2-1} - a_1^\dagger a_2 q^{-D_1+D_2-2} z) &= (q^{D_1-D_2-1} - a_1 \bar{a}_2^\dagger q^{D_1-D_2} z) X, \\ X\left(\bar{a}_2^\dagger q^{D_1-D_2-1} + a_1^\dagger q^{-D_1-D_2-2} z\right) - a_1^\dagger q^{-D_1+D_2} z z_2^2 &= \\ &= \left(\bar{a}_2^\dagger q^{-D_1-D_2-1} - (a_1^\dagger q^{-D_1-D_2-2} + \bar{a}_2^\dagger q^{D_1-D_2+1} z) z z_2^2\right) X, \\ X\left(a_2 q^{-D_1+D_2-1} - (a_1 q^{D_1-D_2} + a_2 q^{D_1+D_2+1} z) z z_2^2\right) &= \\ &= \left((a_2 q^{D_1+D_2-1} + a_1 q^{D_1-D_2} z) - a_1 q^{D_1+D_2+2} z z_2^2\right) X, \\ X\left(q^{-D_1+D_2+1} + q^{D_1-D_2+1} z^2 - a_1 \bar{a}_2^\dagger q^{D_1-D_2} z\right) &= \\ &= \left(q^{-D_1+D_2+1} + q^{D_1-D_2+1} z^2 - a_1^\dagger a_2 q^{-D_1+D_2-2} z\right) X \end{aligned}$$

for $X \in \tilde{\mathcal{A}}_0^{(2)}$ fixed by $\chi^{(2)}$. Since $\mathbb{C}[[z, z_2]] \cong (\mathbb{C}[[z]])[[z_2]]$, considering expansion coefficients with respect to z_2 , we can use $[X, q^{D_1+D_2}] = 0$ to deduce that the above system is equivalent to

$$\begin{aligned} Xa_2q^{-2D_1} &= (a_2 + a_1q^{-2D_2+1}z)X, & a_1X &= X(a_1q^{-2(D_2+1)} + q^{-1}a_2z), \\ Xa_1^\dagger q^{2(D_2+1)} &= (a_1^\dagger + \bar{a}_2^\dagger q^{2D_1+3}z)X, & \bar{a}_2^\dagger X &= X(\bar{a}_2^\dagger q^{2D_1} + q^{-1}a_1^\dagger z), \end{aligned} \quad (\text{B.20})$$

$$\begin{aligned} [X, q^{2D_1}] &= (Xa_1^\dagger a_2 q^{2D_2-1} - a_1 \bar{a}_2^\dagger q^{2D_1+1} X)z, \\ [X, q^{2D_2} + q^{2D_1} z^2] &= (Xa_1 \bar{a}_2^\dagger q^{2D_1-1} - a_1^\dagger a_2 q^{2D_2-3} X)z. \end{aligned} \quad (\text{B.21})$$

Note that $q^{-2D_1(D_2+1)} \in \tilde{\mathcal{A}}_0^{(2)}$ is fixed by $\chi^{(2)}$. Hence without loss of generality we may write

$$X = \tilde{X}q^{-2D_1(D_2+1)}, \quad (\text{B.22})$$

for some $\tilde{X} \in \tilde{\mathcal{A}}_0^{(2)}$ fixed by $\chi^{(2)}$. The system (B.20–B.21) is equivalent to

$$[\tilde{X}, a_2] = q^{-2D_2+1}a_1\tilde{X}z, \quad [a_1, \tilde{X}] = \tilde{X}q^{2D_1-1}a_2z, \quad (\text{B.23})$$

$$[\tilde{X}, a_1^\dagger] = \bar{a}_2^\dagger q^{2D_1+3}\tilde{X}z, \quad [\bar{a}_2^\dagger, \tilde{X}] = \tilde{X}a_1^\dagger q^{-2D_2-3}z, \quad (\text{B.24})$$

$$[\tilde{X}, q^{2D_1}] = (\tilde{X}a_1^\dagger a_2 q^{2D_1-1} - a_1 \bar{a}_2^\dagger q^{2D_1+1}\tilde{X})z, \quad (\text{B.25})$$

$$[\tilde{X}, q^{2D_2} + q^{2D_1} z^2] = (\tilde{X}a_1 \bar{a}_2^\dagger q^{2D_2+3} - a_1^\dagger a_2 q^{2D_2-3}\tilde{X})z. \quad (\text{B.26})$$

Since $\chi^{(2)}$ fixes \tilde{X} , the equations in (B.23) and the equations in (B.24) are pairwise equivalent. At the same time, the system (B.23–B.24) implies (B.25) and (B.26). To show this, since $[\tilde{X}, q^{2D_1}] = [a_1^\dagger a_1, \tilde{X}]$ from (B.23–B.24) we obtain

$$\begin{aligned} [\tilde{X}, q^{2D_1}] + a_1 \bar{a}_2^\dagger q^{2D_1+1}\tilde{X}z - \tilde{X}a_1^\dagger a_2 q^{2D_1-1}z &= \\ = a_1 \bar{a}_2^\dagger q^{2D_1+1}\tilde{X}z - [\tilde{X}, a_2^\dagger]a_1 + a_1^\dagger [a_1, \tilde{X}] - \tilde{X}a_1^\dagger a_2 q^{2D_1-1}z &= \\ = (\bar{a}_2^\dagger q^{2D_1+3}[a_1, \tilde{X}] - [\tilde{X}, a_1^\dagger]a_2 q^{2D_1-1})z, \end{aligned}$$

which vanishes, thereby recovering (B.25). Applying $\chi^{(2)}$ to (B.25) we obtain $[\tilde{X}, q^{-2D_2}] = (\tilde{X}a_1^\dagger a_2 q^{-2D_2-1} - a_1 \bar{a}_2^\dagger q^{-2D_2+1}\tilde{X})z$. Left-and-right multiplying this by q^{2D_2} and using (B.23–B.24) to rewrite the result we obtain

$$[\tilde{X}, q^{2D_2}] = (\bar{a}_2^\dagger \tilde{X}a_1 q^{2D_2+3} - q^{2D_2-1}a_1^\dagger \tilde{X}a_2)z. \quad (\text{B.27})$$

Finally, using (B.27) and again (B.23–B.24), we derive that

$$\begin{aligned} [\tilde{X}, q^{2D_2} + q^{2D_1} z^2] - \tilde{X}a_1 \bar{a}_2^\dagger q^{2D_2+3}z + a_1^\dagger a_2 q^{2D_2-3}\tilde{X}z &= \\ = \bar{a}_2^\dagger \tilde{X}a_1 q^{2D_2+3}z - q^{2D_2-1}a_1^\dagger \tilde{X}a_2 z + [\tilde{X}, q^{2D_1}]z^2 + &= \\ - (\bar{a}_2^\dagger \tilde{X} - \tilde{X}a_1^\dagger q^{-2D_2-3}z)a_1 q^{2D_2+3}z + a_1^\dagger q^{2D_2-1}(\tilde{X}a_2 - a_1 q^{-2D_2+1}\tilde{X}z)z &= \\ = (\tilde{X}a_1^\dagger a_1 + a_1^\dagger a_1 \tilde{X} + [\tilde{X}, 1 - a_1^\dagger a_1])z^2 \end{aligned}$$

which vanishes, thereby proving (B.26) as well.

We have obtained that the system (B.23–B.26) is equivalent to the system (B.24). Writing $p = q^2$, without loss of generality we set

$$\tilde{X} = Y e_p(q^3 z a_1 \bar{a}_2^\dagger) e_p(q^{-1} z a_1^\dagger a_2)$$

for some $Y \in \tilde{\mathcal{A}}_0^{(2)}$ fixed by $\chi^{(2)}$, noting that $e_p(q^3 z a_1 \bar{a}_2^\dagger)$ and $e_p(q^{-1} z a_1^\dagger a_2)$ lie in $\tilde{\mathcal{A}}_0^{(2)}$ and are fixed by $\chi^{(2)}$. The theorem now follows from the following claim.

Claim: (B.24) is satisfied if and only if $Y \in \mathbb{C}[[z]]$.

In the special case $Y = 1$, (B.24) is indeed satisfied:

$$\begin{aligned} [\tilde{X}, a_1^\dagger] - \bar{a}_2^\dagger q^{2D_1+3} z \tilde{X} &= \\ &= \left([e_p(q^3 z a_1 \bar{a}_2^\dagger), a_1^\dagger] - \bar{a}_2^\dagger q^{2D_1+3} z e_p(q^3 z a_1 \bar{a}_2^\dagger) \right) e_p(q^{-1} z a_1^\dagger a_2), \\ [\bar{a}_2^\dagger, \tilde{X}] - \tilde{X} a_1^\dagger q^{-2D_2-3} z &= \\ &= e_p(q^3 z a_1 \bar{a}_2^\dagger) \left([\bar{a}_2^\dagger, e_p(q^{-1} z a_1^\dagger a_2)] - e_p(q^{-1} z a_1^\dagger a_2) a_1^\dagger q^{-2D_2-3} z \right), \end{aligned}$$

with the expressions in parentheses vanishing by virtue of (A.14) and (B.10) (with $y = q^{-1}z$).

For general Y we therefore have

$$\begin{aligned} [\tilde{X}, a_1^\dagger] - \bar{a}_2^\dagger q^{2D_1+3} z \tilde{X} &= [Y, a_1^\dagger] e_p(q^3 z a_1 \bar{a}_2^\dagger) e_p(q^{-1} z a_1^\dagger a_2), \\ [\bar{a}_2^\dagger, \tilde{X}] - \tilde{X} a_1^\dagger q^{-2D_2-3} z &= [\bar{a}_2^\dagger, Y] e_p(q^3 z a_1 \bar{a}_2^\dagger) e_p(q^{-1} z a_1^\dagger a_2). \end{aligned}$$

Both right-hand sides vanish, i.e. (B.24) is indeed satisfied, if and only if Y lies in the centralizer in $\tilde{\mathcal{A}}_0^{(2)}$ of $\{a_1^\dagger, \bar{a}_2^\dagger\}$, which is trivial by Lemma B.5. This proves the claim. \square

C. An Alternative Proof of the Main Theorem

In this part of the appendix we give a proof of the boundary factorization identity (8.2) independent of the universal K-matrix formalism, instead relying on the explicit expressions obtained in Appendix B. Before we state and prove a key lemma, note that expressions of the form $e_p(\gamma^D y)$ where $\gamma \in \mathbb{C}^\times$ and y is an indeterminate give rise to well-defined $\text{End}(W)$ -valued formal power series, sending w_j to $e_p(\gamma^j y) w_j$.

Lemma C.1. *Let y be a formal parameter and let p be a nonzero complex number, not a root of unity. In $\text{End}(W \otimes W)[[y]]$ we have the identities*

$$\begin{aligned} e_p(pa_1 \bar{a}_2^\dagger)(y; p)_{D_1} &= (y; p)_{D_1} e_p(-a_1 \bar{a}_2^\dagger p^{D_1} y) e_p(pa_1 \bar{a}_2^\dagger) \quad (\text{C.1}) \\ e_p(pa_1 \bar{a}_2^\dagger)(p^{1-D_1} y; p)_{D_1}^{-1} e_p(py \bar{a}_1^\dagger a_2) &= e_p(py \bar{a}_1^\dagger a_2)(p^{1-D_2} y; p)_{D_2}^{-1} e_p(pa_1 \bar{a}_2^\dagger). \quad (\text{C.2}) \end{aligned}$$

Proof. Note that

$$W \otimes W = \bigoplus_{m \in \mathbb{Z}_{\geq 0}} (W \otimes W)_m, \quad (W \otimes W)_m := \bigoplus_{\substack{j, k \geq 0 \\ j+k=m}} \mathbb{C} w_j \otimes w_k.$$

Because each factor in (C.1–C.2) preserves each finite-dimensional subspace $(W \otimes W)_m$, it suffices to prove the restrictions of (C.1–C.2) to $(W \otimes W)_m$, where $m \in \mathbb{Z}_{\geq 0}$ is fixed but arbitrary. Note that on $(W \otimes W)_m$ the operators appearing as arguments of the deformed exponentials are nilpotent. Therefore the operators on the left- and right-hand side of the restricted equations depend rationally on p and hence it suffices to prove them with p restricted to an open subset of \mathbb{C} .

We will prove the restriction of (C.1) to $(W \otimes W)_m$ for all $p \in \mathbb{C}$ such that $|p| < 1$. Combining (A.4) and (A.7) we obtain $(y; p)_D = \frac{e_p(p^D y)}{e_p(y)}$; as a consequence, (C.1) is equivalent to

$$e_p(pa_1\bar{a}_2^\dagger)e_p(p^{D_1}y) = e_p(p^{D_1}y)e_p(-a_1\bar{a}_2^\dagger p^{D_1}y)e_p(pa_1\bar{a}_2^\dagger). \tag{C.3}$$

But this equation follows directly from (A.12) and the observation $(a_1\bar{a}_2^\dagger)(p^{D_1}y) = p(p^{D_1}y)(a_1\bar{a}_2^\dagger)$.

On the other hand,¹³ we will prove the restricted version of (C.2) for all $p \in \mathbb{C}^\times$ such that $|p| > 1$. In this case, for all $j \in \mathbb{Z}_{\geq 0}$ we have

$$(p^{1-j}y; p)_j^{-1} = (y; p^{-1})_j^{-1} = \frac{(p^{-j}y; p^{-1})_\infty}{(y; p^{-1})_\infty} \in \mathbb{C}[[y]].$$

From (A.7) and (A.9) we deduce the identity

$$(p^{-j}y; p^{-1})_\infty = e_{p^{-1}}(p^{-j}y)^{-1} = e_p(p^{1-j}y) \in \mathbb{C}[[y]].$$

Hence $(p^{1-D}y; p)_D^{-1} = (y; p^{-1})_\infty^{-1}e_p(p^{1-D}y)$ in $\text{End}(W)[[y]]$ and (C.2) is equivalent to

$$e_p(pa_1\bar{a}_2^\dagger)e_p(p^{1-D_1}y)e_p(py\bar{a}_1^\dagger a_2) = e_p(py\bar{a}_1^\dagger a_2)e_p(p^{1-D_2}y)e_p(pa_1\bar{a}_2^\dagger). \tag{C.4}$$

To prove (C.4), note that (B.11) can be evaluated at $y = p$, and the result can be rewritten as

$$e_p(pa_1\bar{a}_2^\dagger)(p^{-D_1} + \bar{a}_1^\dagger a_2) = (p^{-D_2} + \bar{a}_1^\dagger a_2)e_p(pa_1\bar{a}_2^\dagger).$$

By iteration we obtain

$$e_p(pa_1\bar{a}_2^\dagger)e_p(p^{1-D_1}y + py\bar{a}_1^\dagger a_2) = e_p(p^{1-D_2}y + py\bar{a}_1^\dagger a_2)e_p(pa_1\bar{a}_2^\dagger). \tag{C.5}$$

Note that $(\bar{a}_1^\dagger a_2)p^{1-D_1} = p p^{1-D_1}(\bar{a}_1^\dagger a_2)$ and $p^{1-D_2}(\bar{a}_1^\dagger a_2) = p(\bar{a}_1^\dagger a_2)p^{1-D_2}$. Applying (A.11), we obtain (C.4), as required. \square

Alternative proof of Theorem 8.1. By virtue of (4.12), the desired identity, viz.

$$\mathcal{K}_v(z)_1 \mathcal{R}_{v\phi}(z^2) \mathcal{K}_\phi(z)_2 \mathcal{O} = \mathcal{O} \mathcal{K}_\varrho(q^{-\mu/2}z)_1 \mathcal{R}_{\varrho\bar{\varrho}}(z^2) \mathcal{K}_{\bar{\varrho}}(q^{\mu/2}z)_2 \tag{C.6}$$

for arbitrary $\mu \in \mathbb{C}$ and $q, \xi \in \mathbb{C}^\times$ such that q is not a root of unity, is equivalent to

$$\begin{aligned} e_p(pa_1\bar{a}_2^\dagger)\mathcal{K}_v(z)_1 \mathcal{R}_{v\phi}(z^2) \mathcal{K}_\phi(z)_2 e_p(pa_1\bar{a}_2^\dagger)^{-1} &= \\ = q^{\mu(D_1-D_2)/2} \mathcal{K}_\varrho(q^{-\mu/2}z)_1 \mathcal{R}_{\varrho\bar{\varrho}}(z^2) \mathcal{K}_{\bar{\varrho}}(q^{\mu/2}z)_2 q^{\mu(D_2-D_1)/2} &\tag{C.7} \end{aligned}$$

where $p = q^2$. The strategy of the proof is as follows. We move various simple factors in $\mathcal{F}^{(2)}(D_1, D_2)$ to the right in both sides of (C.7), thus bringing them to a similar form. Then more advanced product formulas involving q-exponentials and finite q-Pochhammer symbols yield the desired equality.

More precisely, we set $\gamma = pq^{-\mu}\xi^{-1} \in \mathbb{C}^\times$ and from (A.2) deduce

$$(\gamma z^{-2}; p)_j^{-1} = p^{j(1-j)/2} (-\gamma^{-1}z^2)^j (p^{1-j}\gamma^{-1}z^2; p)_j^{-1}$$

for all $j \in \mathbb{Z}_{\geq 0}$. Using the identities $q^{-D^2}a^\dagger = -q\bar{a}^\dagger q^{-D^2}$ and $q^{-D^2}a = aq^{2D-1}q^{-D^2}$, we obtain, for the left-hand side of (C.7),

¹³ We will need (C.2) with $|p| < 1$, but we are not aware of a direct proof of this.

$$\begin{aligned}
 & e_p(pa_1\bar{a}_2^\dagger)\mathcal{K}_v(z)_1\mathcal{R}_{v\phi}(z^2)\mathcal{K}_\phi(z)_2e_p(pa_1\bar{a}_2^\dagger)^{-1} = \\
 & = e_p(pa_1\bar{a}_2^\dagger)(-\gamma^{-1}q^{1-D_1})^{D_1}(\gamma z^2; p)_{D_1}(p^{1-D_1}\gamma^{-1}z^2; p)_{D_1}^{-1}e_p(z^2a_1^\dagger a_2) \cdot \\
 & \quad \cdot q^{(2\mu-1)D_1-2D_2-2D_1D_2-D_2^2}(-\xi)^{D_2}e_p(pa_1\bar{a}_2^\dagger)^{-1} \\
 & = e_p(pa_1\bar{a}_2^\dagger)(\gamma z^2; p)_{D_1}(p^{1-D_1}\gamma^{-1}z^2; p)_{D_1}^{-1} \cdot \\
 & \quad \cdot e_p(p\gamma^{-1}z^2\bar{a}_1^\dagger a_2)(-q^{-D_1-D_2-2\xi})^{D_1+D_2}e_p(pa_1\bar{a}_2^\dagger)^{-1} \\
 & = e_p(pa_1\bar{a}_2^\dagger)(\gamma z^2; p)_{D_1}(p^{1-D_1}\gamma^{-1}z^2; p)_{D_1}^{-1} \cdot \\
 & \quad \cdot e_p(p\gamma^{-1}z^2\bar{a}_1^\dagger a_2)e_p(pa_1\bar{a}_2^\dagger)^{-1}(-q^{-D_1-D_2-2\xi})^{D_1+D_2}. \tag{C.8}
 \end{aligned}$$

Similarly, for the right-hand side of (C.7) we obtain

$$\begin{aligned}
 & q^{\mu(D_1-D_2)/2}\mathcal{K}_\rho(q^{-\mu/2}z)_1\mathcal{R}_{\rho\bar{q}}(z^2)\mathcal{K}_{\bar{q}}(q^{\mu/2}z)_2q^{\mu(D_2-D_1)/2} = \\
 & = (\gamma z^2; p)_{D_1}q^{\mu(D_1-D_2)/2-D_1^2}(-\xi)^{D_1}e_p(q^3z^2a_1\bar{a}_2^\dagger)e_p(q^{-1}z^2a_1^\dagger a_2) \cdot \\
 & \quad \cdot (p^{1-D_2}\gamma^{-1}z^2; p)_{D_2}^{-1}q^{\mu(D_2-D_1)/2-2(D_1+D_2)-2D_1D_2-D_2^2}(-\xi)^{D_2} \\
 & = (\gamma z^2; p)_{D_1}e_p(-a_1\bar{a}_2^\dagger q^{2D_1}\gamma z^2) \\
 & \quad e_p(p\gamma^{-1}z^2\bar{a}_1^\dagger a_2)(p^{1-D_2}\gamma^{-1}z^2; p)_{D_2}^{-1}(-q^{-D_1-D_2-2\xi})^{D_1+D_2}. \tag{C.9}
 \end{aligned}$$

Therefore (C.7) is equivalent to

$$\begin{aligned}
 & e_p(pa_1\bar{a}_2^\dagger)(\gamma z^2; p)_{D_1}(p^{1-D_1}\gamma^{-1}z^2; p)_{D_1}^{-1}e_p(p\gamma^{-1}z^2\bar{a}_1^\dagger a_2)e_p(pa_1\bar{a}_2^\dagger)^{-1} = \\
 & = (\gamma z^2; p)_{D_1}e_p(-a_1\bar{a}_2^\dagger p^{D_1}\gamma z^2)e_p(p\gamma^{-1}z^2\bar{a}_1^\dagger a_2)(p^{1-D_2}\gamma^{-1}z^2; p)_{D_2}^{-1}. \tag{C.10}
 \end{aligned}$$

Applying (C.1) with $y = \gamma z^2$ and (C.2) with $y = \gamma^{-1}z^2$, we deduce (C.10), as required. □

References

- [AV22a] Appell, A., Vlaar, B.: Universal K-matrices for quantum Kac–Moody algebras. *Represent. Theory* **26**, 764–824 (2022)
- [AV22b] Appell, A., Vlaar, B.: Trigonometric K-matrices for finite-dimensional representations of quantum affine algebras (2022). Preprint at [arXiv:2203.16503](https://arxiv.org/abs/2203.16503)
- [Ba72] Baxter, R.: Partition function of the eight-vertex lattice model. *Ann. Phys.* **70**(1), 193–228 (1972)
- [Ba73] Baxter, R.: Eight-vertex model in lattice statistics and one-dimensional anisotropic Heisenberg chain. I. Some fundamental eigenvectors. *Ann. Phys.* **76**(1), 1–24 (1973)
- [BB13] Baseilhac, P., Belliard, S.: The half-infinite XXZ chain in Onsager’s approach. *Nucl. Phys. B* **873**(3), 550–584 (2013)
- [BGKNR10] Boos, H., Göhmann, F., Klümper, A., Nirov, K., Razumov, A.: Exercises with the universal R-matrix. *J. Phys. A Math. Theor.* **43**(41), 415208 (2010)
- [BGKNR13] Boos, H., Göhmann, F., Klümper, A., Nirov, K., Razumov, A.: Universal integrability objects. *Theor. Math. Phys.* **174**(1), 21–39 (2013)
- [BGKNR14] Boos, H., Göhmann, F., Klümper, A., Nirov, K., Razumov, A.: Universal R-matrix and functional relations. *Rev. Math. Phys.* **26**(6), 1430005 (2014)
- [BJMST09] Boos, H., Jimbo, M., Miwa, T., Smirnov, F., Takeyama, Y.: Hidden Grassmann structure in the XXZ model II: creation operators. *Commun. Math. Phys.* **286**(3), 875–932 (2009)

- [BK19] Balagović, M., Kolb, S.: Universal K-matrix for quantum symmetric pairs. *Journal für die reine und angewandte Mathematik (Crelles Journal)* (2019)
- [BLMS10] Bazhanov, V., Łukowski, T., Meneghelli, C., Staudacher, M.: A Shortcut to the Q-operator. *J. Stat. Mech. Theory Exp.* **11**, P11002 (2010)
- [BLZ96] Bazhanov, V., Lukyanov, S., Zamolodchikov, A.: Integrable structure of conformal field theory, quantum KdV theory and thermodynamic Bethe ansatz. *Commun. Math. Phys.* **177**(2), 381–398 (1996)
- [BLZ97] Bazhanov, V., Lukyanov, S., Zamolodchikov, A.: Integrable structure of conformal field theory II. Q-operator and DDV equation. *Commun. Math. Phys.* **190**, 247–278 (1997)
- [BLZ99] Bazhanov, V., Lukyanov, S., Zamolodchikov, A.: Integrable structure of conformal field theory III. The Yang–Baxter relation. *Commun. Math. Phys.* **200**(2), 297–324 (1999)
- [BS90] Bazhanov, V., Stroganov, Yu.: Chiral Potts model as a descendant of the six-vertex model. *J. Stat. Phys.* **59**(3), 799–817 (1990)
- [BT18] Baseilhac, P., Tsuboi, Z.: Asymptotic representations of augmented q -Onsager algebra and boundary K-operators related to Baxter Q-operators. *Nucl. Phys. B* **929**, 397–437 (2018)
- [BW18] Bao, H., Wang, W.: A new approach to Kazhdan-Lusztig theory of type B via quantum symmetric pairs. *Société mathématique de France, Paris* (2018)
- [Ch02] Chari, V.: Braid group actions and tensor products. *Int. Math. Res. Not.* **7**, 357–382 (2002)
- [Ch84] Cherednik, I.: Factorizing particles on a half-line and root systems. *Theor. Math. Phys.* **61**(1), 977–983 (1984)
- [CP94] Chari, V., Pressley, A.: Quantum affine algebras and their representations. Representation theory of groups (Banff, AB, 1994), CMS Conference Proceedings, vol. 16, pp. 59–78. American Mathematical Society, Providence (1994)
- [CP95] Chari, V., Pressley, A.: A guide to quantum groups. Cambridge University Press, Cambridge (1995). Corrected reprint of the 1994 original
- [De05] Derkachov, S.: Factorization of R-matrix and Baxter's Q-operator (2005). Preprint at [arXiv:math/0507252](https://arxiv.org/abs/math/0507252)
- [De07] Derkachov, S.: Factorization of the R-matrix. *I. J. Math. Sci.* **143**(1), 2773–2790 (2007)
- [DG02] Delius, G., George, A.: Quantum affine reflection algebras of type $d_n^{(1)}$ and reflection matrices. *Lett. Math. Phys.* **62**, 211–217 (2002)
- [DKK06] Derkachov, S., Karakhanyan, D., Kirschner, R.: Baxter Q-operators of the XXZ chain and R-matrix factorization. *Nucl. Phys. B* **738**(3), 368–390 (2006)
- [DM03] Delius, G., Mackay, N.: Quantum group symmetry in sine-Gordon and affine Toda field theories on the half-line. *Commun. Math. Phys.* **233**, 173–190 (2003)
- [Dr85] Drinfeld, V.: Hopf algebras and the quantum Yang–Baxter equation. *Soviet Math. Dokl.* **32**, 254–258 (1985)
- [Dr86] Drinfeld, V.: Quantum groups. In: Gleason, A.M. (ed.) *Proceedings of the International Congress of Mathematicians, Berkeley*, pp. 798–820. American Mathematical Society, Providence (1986)
- [EM03] Etingof, P., Moura, A.: Elliptic central characters and blocks of finite-dimensional representations of quantum affine algebras. *Represent. Theory* **7**, 346–373 (2003)
- [FH15] Frenkel, E., Hernandez, D.: Baxter's relations and spectra of quantum integrable models. *Duke Math. J.* **164**(12), 2407–2460 (2015)
- [FR92] Frenkel, I., Reshetikhin, N.: Quantum affine algebras and holonomic difference equations. *Commun. Math. Phys.* **146**(1), 1–60 (1992)
- [FR99] Frenkel, E., Reshetikhin, N.: The q -characters of representations of quantum affine algebras and deformations of W -algebras, in *Recent Developments in Quantum Affine Algebras and related topics*. *Contemp. Math.* **248**, 163–205 (1999)
- [FS15] Frassek, R., Szécsényi, I.: Q-operators for the open Heisenberg spin chain. *Nucl. Phys. B* **901**, 229–248 (2015)
- [GR90] Gasper, G., Rahman, M.: Basic hypergeometric series, *Encyclopedia of mathematics and its applications*, vol. 35. Cambridge University Press, Cambridge (1990)
- [He19] Hernandez, D.: Avancées concernant les R-matrices et leurs applications [d'après Maulik-Okounkov, Kang-Kashiwara-Kim-Oh, ...] (2019), no. 407, *Séminaire Bourbaki*, 69^{ème} année, Vol. 2016–2017, no. 1129, pp. 297–331
- [HJ12] Hernandez, D., Jimbo, M.: Asymptotic representations and Drinfeld rational fractions. *Compos. Math.* **148**(5), 1593–1623 (2012)
- [Ja96] Jantzen, J. C.: Lectures on quantum groups. *Grad. Stud. Math.*, vol. 6. American Mathematical Society, Providence (1996)
- [Ji86a] Jimbo, M.: A q -analogue of $U(\mathfrak{gl}(N+1))$, Hecke algebra, and the Yang–Baxter equation. *Lett. Math. Phys.* **11**(3), 247–252 (1986)

- [Ji86b] Jimbo, M.: Quantum R matrix for the generalized Toda system. *Commun. Math. Phys.* **102**, 537–547 (1986)
- [Ka90] Kac, V.: *Infinite-Dimensional Lie Algebras*, 3rd edn. Cambridge University Press, Cambridge (1990)
- [Ko14] Kolb, S.: Quantum symmetric Kac–Moody pairs. *Adv. Math.* **267**, 395–469 (2014)
- [Ko97] Koornwinder, T.: Special functions and q-commuting variables. In: *Special Functions, q-Series and Related Topics*, Fields Institute Communications 14, pp. 131–166. American Mathematical Society, Providence (1997)
- [KS95] Kazhdan, D., Soibelman, Y.: Representations of quantum affine algebras. *Sel. Math. (N.S.)* **1**(3), 537–595 (1995)
- [KT14] Khoroshkin, S., Tsuboi, Z.: The universal R-matrix and factorization of the L-operators related to the Baxter Q-operators. *J. Phys. A Math. Theor.* **47**(19), 192003 (2014)
- [Lu94] Lusztig, G.: *Introduction to Quantum Groups*. Birkhäuser, Boston (1994)
- [MN98] Mezincescu, L., Nepomechie, R.I.: Fractional-spin integrals of motion for the boundary Sine-Gordon model at the free fermion point. *Int. J. Mod. Phys. A* **13**, 2747–2764 (1998)
- [RSV15] Reshetikhin, N., Stokman, J., Vlaar, B.: Boundary quantum Knizhnik–Zamolodchikov equations and fusion. *Annales Henri Poincaré* 1–41 (2015)
- [RV16] Regelskis, V., Vlaar, B.: Reflection matrices, coideal subalgebras and generalized Satake diagrams of affine type
- [Sk88] Sklyanin, E.: Boundary conditions for integrable quantum systems. *J. Phys. A Math. Gen.* **21**, 2375–2389 (1988)
- [KT92] Tolstoy, V., Khoroshkin, S.: The universal R-matrix for quantum untwisted affine Lie algebras. *Funct. Anal. Appl.* **26**, 69–71 (1992)
- [Ts21] Tsuboi, Z.: Universal Baxter TQ-relations for open boundary quantum integrable systems. *Nucl. Phys. B* **963**, 115286 (2021)
- [VW20] Vlaar, B., Weston, R.: A Q-operator for open spin chains I: Baxter’s TQ relation. *J. Phys. A Math. Theor.* **53**(24), 245202 (2020)
- [YNZ06] Yang, W.-L., Nepomechie, R.I., Zhang, Y.-Z.: Q-operator and T-Q relation from the fusion hierarchy. *Phys. Lett. B* **633**(4–5), 664–670 (2006)
- [YZ06] Yang, W.-L., Zhang, Y.-L.: *T-Q* relation and exact solution for the XYZ chain with general nondiagonal boundary terms. *Nucl. Phys. B* **744**, 312–329 (2006)