

Geometric formulation of generalized root- $T\bar{T}$ deformations

H. Babaei-Aghbolagh,^{2,*} Song He,^{1,3,†} Tommaso Morone,^{4,‡} Hao Ouyang,^{1,§} and Roberto Tateo^{4,¶}

¹Center for Theoretical Physics and College of Physics, Jilin University, Changchun 130012, China

²Department of Physics, University of Mohaghegh Ardabili, P.O. Box 179, Ardabil, Iran

³Max Planck Institute for Gravitational Physics (Albert Einstein Institute), Am Mühlenberg 1, 14476 Golm, Germany

⁴Dipartimento di Fisica, Università di Torino, and INFN Sezione di Torino, Via P. Giuria 1, 10125, Torino, Italy

We develop a generic geometric formalism that incorporates both $T\bar{T}$ -like and root- $T\bar{T}$ -like deformations in arbitrary dimensions. This framework applies to a wide family of stress-energy tensor perturbations and encompasses various well-known field theories. Building upon the recently proposed correspondence between Ricci-based gravity and $T\bar{T}$ -like deformations, we further extend this duality to include root- $T\bar{T}$ -like perturbations. This refinement extends the potential applications of our approach and contributes to a deeper exploration of the interplay between stress tensor perturbations and gravitational dynamics. Among the various original outcomes detailed in this article, we have also obtained a deformation of the flat Jackiw-Teitelboim gravity action.

INTRODUCTION

Recent studies concerning deformations of classical and quantum field theories have revealed rich connections between geometry and field dynamics. A prime example is that of $T\bar{T}$ deformations [1, 2] of two-dimensional theories, driven by the irrelevant composite operator [3]

$$O_{T\bar{T}} = -\det(T^\mu{}_\nu) = \frac{1}{2}(T^{\mu\nu}T_{\mu\nu} - T^\mu{}_\mu T^\nu{}_\nu). \quad (1)$$

Despite being irrelevant, two-dimensional $T\bar{T}$ deformations remain well-controlled and even solvable at the quantum level. In the deformed theory, various quantities can be computed exactly from their counterparts in the original model. These include the finite-volume spectrum, the S-matrix [1, 2], the classical Lagrangian [4–6], and the torus partition function [7–10]. $T\bar{T}$ deformations connect with different topics in theoretical physics, such as string theory [11–14], holography [15–25], and quantum gravity [26–31]. We refer the reader to [32] for a pedagogical review on the subject.

Furthermore, the $T\bar{T}$ deformation lends itself to a number of geometric interpretations. It was proposed in [7] that $T\bar{T}$ -perturbing a theory is equivalent to coupling the original theory to a random geometry. $T\bar{T}$ deformations can also be interpreted as coupling the original theory to a flat space Jackiw-Teitelboim-like gravity [26, 27], or equivalently, a topological gravity [28, 33, 34].

Another interesting deformation of two-dimensional field theories, driven by the so-called root- $T\bar{T}$ operator [35–37]

$$R = \sqrt{\frac{1}{2}T^{\mu\nu}T_{\mu\nu} - \frac{1}{4}T^\mu{}_\mu T^\nu{}_\nu}, \quad (2)$$

has recently attracted growing attention. While its quantum-mechanical definition remains uncertain, the root- $T\bar{T}$ perturbation displays some surprising properties at the classical level. Notably, it commutes with the $T\bar{T}$, allowing for their simultaneous activation, and for

some integrable field theories, it preserves classical integrability [38]. Finally, the connection between the root- $T\bar{T}$ deformation and the modified boundary conditions in the holographic dictionary was studied in [39]. These results were later employed to explore the modular properties of deformed holographic conformal field theories in [40].

In higher space-time dimensions, stress-energy tensor perturbations give rise to many interesting field theory models [4, 7, 35]. Extensive research has focused on $T\bar{T}$ -like and root- $T\bar{T}$ -like deformations of four-dimensional Maxwell’s theory, exploring the relationship between electromagnetic duality invariance and stress tensor deformations [41–43]. The massive gravity formulation of duality-invariant non-linear electrodynamics was studied in [44] and, in three dimensions, it was shown that Born-Infeld theory displays a classical $T\bar{T}$ -like flow, connected to free Maxwell theory [45]. Furthermore, recent studies have explored nonlinear chiral two-form gauge theories in six dimensions as $T\bar{T}$ -like deformations [46].

This paper introduces a generic geometric approach to encompass a broader class of stress-energy tensor perturbations. We show that a two-dimensional theory deformed by both $T\bar{T}$ and root- $T\bar{T}$ operators is dynamically equivalent to the undeformed theory coupled to a novel gravity action, at least at a classical level. We further generalize the geometric formulation to accommodate various deformations in higher dimensions. While prior studies have investigated geometric formulations of $T\bar{T}$ -like deformations in higher dimensions within the metric approach [35], our formulation is based on the description in terms of eigenvalues of the product of the vielbein. This approach allows us to study stress-energy tensor-related flows within a simple and elegant setup.

The recent work [47] has emphasized that a $T\bar{T}$ -type deformed matter action coupled with the standard Einstein-Hilbert action is equivalent to an undeformed matter theory coupled with a Ricci-based gravity theory [48]. Adopting this perspective, we incorporate this logic into our geometric formulations and introduce Ricci-

based gravity actions linked with root- $T\bar{T}$ -like deformations. We develop a unified framework for $T\bar{T}$ and root- $T\bar{T}$ perturbations in field theories across various space-time dimensions, which may extend the class of exact-solvability preserving deformations and deepen our understanding of the fundamental principles of quantum gravity and string theory.

UNIFIED GEOMETRIC FORMULATION OF $T\bar{T}$ AND ROOT- $T\bar{T}$ DEFORMATIONS IN $d = 2$

We denote by $S_0[\phi, e_\mu^a]$ an arbitrary undeformed action, where ϕ indicates a generic collection of matter fields and e_ν^a denotes an auxiliary dynamical zweibein. The associated auxiliary metric is $g_{\mu\nu} = \eta_{ab} e_\mu^a e_\nu^b$. We couple the auxiliary zweibein to a second zweibein f_μ^a , and the metric tensor $h_{\mu\nu} = \eta_{ab} f_\mu^a f_\nu^b$ associated to f_μ^a will eventually emerge as the metric of the manifold on which the deformed theory lives. It is convenient to define two Lorentz invariant variables

$$\begin{aligned} y_1 &= \text{tr}(e^{-1}f) = f_\mu^a e_\mu^a, \\ y_2 &= \text{tr}[(e^{-1}f)^2] = f_\mu^a e_\nu^b f_\nu^c e_\mu^d. \end{aligned} \quad (3)$$

We now show that the combination of $T\bar{T}$ and root- $T\bar{T}$ deformations can be generated from the action:

$$S_{\gamma,\lambda}[\phi, e_\mu^a, f_\mu^a] = S_0[\phi, e_\mu^a] + S_{\text{grav}}[e_\mu^a, f_\mu^a], \quad (4)$$

where gravity action S_{grav} is

$$\begin{aligned} S_{\text{grav}}[e_\mu^a, f_\mu^a] &= \frac{1}{2\lambda} \int d^2x \det e \\ &\times \left(2 + y_1^2 - y_2 - 2y_1 \cosh \frac{\gamma}{2} + 2\sqrt{2y_2 - y_1^2} \sinh \frac{\gamma}{2} \right). \end{aligned} \quad (5)$$

The parameters λ and γ represent the $T\bar{T}$ and the root- $T\bar{T}$ perturbing parameters, respectively. When $\gamma = 0$, S_{grav} reduces to the topological gravity action associated to the $T\bar{T}$ deformation [28]:

$$S_{\text{grav}}[e_\mu^a, f_\mu^a] = \frac{1}{2\lambda} \int d^2x \epsilon^{\mu\nu} \epsilon_{ab} (e_\mu^a - f_\mu^a)(e_\nu^b - f_\nu^b), \quad (6)$$

where ϵ is the Levi-Civita symbol. Our analysis will be carried out using the Euclidean signature, and the generalization to the Lorentzian signature is straightforward.

The deformed action can be obtained by extremizing (4) with respect to the auxiliary zweibein e_μ^a : performing the variation of (4) with respect to e_μ^a , we have

$$\det e (T^{[0]})^\mu{}_\nu \equiv \frac{\delta S_0}{\delta e_\mu^a} e_\nu^a = -\frac{\delta S_{\text{grav}}}{\delta e_\mu^a} e_\nu^a, \quad (7)$$

where $(T^{[0]})^\mu{}_\nu$ is the stress-energy tensor of the undeformed theory, computed with respect to e_μ^a . We denote

the solution of the equation of motion by e_μ^{*a} . Note that (7) may admit multiple solutions e_μ^{*a} related to the choice of the branch for the square root of the root- $T\bar{T}$ operator: in this work, we ignore such branch ambiguities. However, in the quantum theory, we expect one should sum over contributions from all branches in the path integral. The deformed field theory is obtained substituting e_μ^{*a} back into (4):

$$S_{\text{deformed}}[\phi, f_\mu^a] = S_{\gamma,\lambda}[\phi, e_\mu^{*a}, f_\mu^a]. \quad (8)$$

The stress-energy tensor of the deformed theory can be computed as

$$T^\mu{}_\nu \equiv \frac{1}{\det f} \frac{\delta S_{\gamma,\lambda}}{\delta f_\mu^a} f_\nu^a = \frac{1}{\det f} \frac{\delta S_{\text{grav}}}{\delta f_\mu^a} f_\nu^a \Big|_{e=e^*}, \quad (9)$$

where we have used the on-shell condition (7) for e_μ^a , so that $S_{\gamma,\lambda}[\phi, e_\mu^{*a}, f_\mu^a]$ explicitly depends on f_μ^a alone. To simplify notation, we will not distinguish between e_ν^a and its on-shell value e_μ^{*a} , unless necessary. One can verify that the total action (4) obeys the following flow equations:

$$\frac{\partial S_{\gamma,\lambda}}{\partial \lambda} = - \int d^2x \det f \det(T^\nu{}_\mu), \quad (10)$$

$$\frac{\partial S_{\gamma,\lambda}}{\partial \gamma} = \int d^2x \det f \sqrt{\frac{1}{2} T^\mu{}_\nu T^\nu{}_\mu - \frac{1}{4} (T^\nu{}_\nu)^2}. \quad (11)$$

Therefore, the action (4) provides a geometric description of the combined $T\bar{T}$ and root- $T\bar{T}$ deformations. Since $S_{\gamma,\lambda}$ is defined as independent of the flow path, and since the operators do not have explicit λ and γ dependence, the two types of deformations commute with each other. As in [28], one can translate the vielbein formulation to the metric formulation by choosing a gauge such that

$$e^{-1}f = \sqrt{g^{-1}h}, \quad (12)$$

where we have omitted indices to simplify the notation. The validity of the flow equations (10) and (11) can also be verified in the metric formulation, and the details are shown in the appendix.

We now illustrate our methodology, starting from the simple undeformed action of a free scalar:

$$S_0[\phi, e_\mu^a] = \int d^2x \det e \left(\frac{1}{2} \eta^{ab} e_\mu^a e_\nu^b \partial_\mu \phi \partial_\nu \phi \right). \quad (13)$$

The solution of the equation of motion for e_μ^a is

$$\begin{aligned} e_\mu^{*a} &= \frac{1}{2} e^{\pm\frac{\gamma}{2}} \left(\frac{1}{\sqrt{1 - 4\lambda e^{\pm\gamma} X}} + 1 \right) f_\mu^a \\ &\mp \left(\frac{\sinh \frac{\gamma}{2} \pm 2\lambda e^{\pm\frac{\gamma}{2}} X}{X \sqrt{1 - 4\lambda e^{\pm\gamma} X}} + \frac{\sinh \frac{\gamma}{2}}{2X} \right) \eta^{ab} f_b^\nu \partial_\nu \phi \partial_\mu \phi, \end{aligned} \quad (14)$$

where $X = \frac{1}{2} \eta^{ab} f_\mu^a f_\nu^b \partial_\mu \phi \partial_\nu \phi$. Substituting the solution e_μ^{*a} into the action, we get

$$S_{\gamma,\lambda}[\phi, e_\mu^{*a}, f_\mu^a] = \int d^2x \frac{1 - \sqrt{1 - 4e^{\pm\gamma} \lambda X}}{2\lambda}, \quad (15)$$

which reproduces the result obtained in [37].

UPLIFT TO HIGHER DIMENSIONS

In this section, we uplift the geometric description to a family of deformations induced by functionals of the stress-energy tensor in higher dimensions. In d space-time dimensions, we consider the following general form for the gravity action:

$$S_{\text{grav}}[e_{\mu}^a, f_{\mu}^a] = \int d^d x \det e B(e^{-1} f), \quad (16)$$

where B is a Lorentz invariant function of $(e^{-1} f)_{\nu}^{\mu}$. Therefore, B depends only on the Lorentz invariant variables $y_n = \text{tr}[(e^{-1} f)^n]$, $n = 1, \dots, d$. For $n > d$, the y_n are not independent quantities. Since one can express B in terms of the variables y_n , the stress-energy tensor can be computed as

$$T \equiv \frac{1}{\det f} \frac{\delta S_{\text{grav}}}{\delta f} f = \sum_{n=1}^d \frac{n}{\det(e^{-1} f)} (e^{-1} f)^n \partial_{y_n} B, \quad (17)$$

where T denotes the matrix T_{ν}^{μ} . To construct higher-dimensional deforming operators, we need to compute Lorentz invariant functionals of the stress-energy tensor (17). Although we can express each invariant $\text{tr}(T^k)$ in terms of the y -variables, this approach is quite inefficient in arbitrary dimensions, since there is no simple general formula for y_n when $n > d$.

However, assuming $e^{-1} f$ can be diagonalized by means of some matrix U as $e^{-1} f = U \text{diag}(\alpha_1, \dots, \alpha_d) U^{-1}$, the function B can be expressed in terms of the eigenvalues α_i , and each y_n reduces to a power sum symmetric polynomial of the α_i . For this reason, working with the eigenvalues of $e^{-1} f$ proves to be a far more convenient strategy. The stress-energy tensor can be expressed as

$$T = \left(\prod_{k=1}^d \alpha_k \right)^{-1} U \text{diag}(\alpha_1 \partial_{\alpha_1} B, \dots, \alpha_d \partial_{\alpha_d} B) U^{-1}, \quad (18)$$

and

$$\text{tr}(T^k) = \left(\prod_{j=1}^d \alpha_j \right)^{-k} \sum_{i=1}^d (\alpha_i \partial_{\alpha_i} B)^k. \quad (19)$$

Expressing y_1 and y_2 in terms of eigenvalues of $e^{-1} f$, the two-dimensional gravity action (5) can be significantly simplified:

$$S_{\text{grav}}[e_{\mu}^a, f_{\mu}^a] = \frac{1}{\lambda} \int d^2 x \det e (\alpha_1 - e^{\frac{\gamma}{2}})(\alpha_2 - e^{-\frac{\gamma}{2}}). \quad (20)$$

Note that S_{grav} is not a symmetric function of the eigenvalues because of the non-analyticity of the root- $T\bar{T}$ operator. Exchanging two eigenvalues is equivalent to crossing a branch cut.

Motivated by the expression (20), we propose a generalization in arbitrary d space-time dimensions:

$$B = \frac{1}{\lambda^{\Sigma-1}} \prod_{k=1}^d (\alpha_k^{p_k} - \beta_k^{p_k})^{1/p_k}, \quad (21)$$

where λ and β_k are perturbing parameters, p_k are numbers characterizing the deformation, and $\Sigma = \sum_{k=1}^d p_k^{-1}$. When $d = 2$ and $p_k = 1$, the action (21) reduces to (20) if we identify $\beta_1 = e^{\frac{\gamma}{2}}$ and $\beta_2 = e^{-\frac{\gamma}{2}}$. We will show that the parameters λ and $\log \beta_k$ emerge as higher-dimensional analogs of the two-dimensional $T\bar{T}$ and root- $T\bar{T}$ deformation parameters, respectively. With the ansatz (21), the eigenvalues of the stress-energy tensor can be computed as

$$\tau_i = \alpha_i \partial_{\alpha_i} B \prod_{j=1}^d \alpha_j^{-1} = \frac{\alpha_i^{p_i}}{\alpha_i^{p_i} - \beta_i^{p_i}} B \prod_{j=1}^d \alpha_j^{-1}. \quad (22)$$

We also find

$$\prod_{k=1}^d \tau_k^{1/p_k} = \frac{1}{\lambda^{\Sigma-1}} B^{\Sigma-1} \left(\prod_{k=1}^d \alpha_k \right)^{1-\Sigma}. \quad (23)$$

Therefore, the flow equation for λ is

$$\frac{\partial S_{\text{grav}}}{\partial \lambda} = -(\Sigma - 1) \int d^d x \det f \left(\prod_{k=1}^d \tau_k^{1/p_k} \right)^{\frac{1}{\Sigma-1}}. \quad (24)$$

The operator on the right-hand side of the equation (24) is non-analytic and not symmetric in terms of the stress-energy tensor eigenvalues τ_i . Particularly, when all p_k are equal to p , we obtain a $(\det T)^{\frac{1}{d-p}}$ deformation [4, 7]. When $\Sigma = 2$, the deformation is of order $O(T^2)$. Let us now consider the flow equation for the β -parameters. We have:

$$\beta_i \partial_{\beta_i} B - \beta_j \partial_{\beta_j} B = -(\tau_i - \tau_j) \prod_{k=1}^d \alpha_k. \quad (25)$$

Equation (25) suggests that the flow should be confined to the surface defined by $\prod_{k=1}^d \beta_k = 1$. Otherwise, the perturbing operator would explicitly depend on λ and β_k . The resulting flow equation is:

$$\sum_{k=1}^d v^k \frac{\partial S_{\beta, \lambda}}{\partial \log \beta_k} = - \int d^d x \det f \left(\sum_{k=1}^d v^k \tau_k \right), \quad (26)$$

where v^k are constants satisfying $\sum_{k=1}^d v^k = 1$. Varying the β -parameters on the surface $\prod_{k=1}^d \beta_k = 1$ leads to non-analytic marginal deformations that commute with the $\prod_{k=1}^d \tau_k^{1/p_k}$ deformation. In two dimensions, the root- $T\bar{T}$ operator can be understood as the difference $\tau_1 - \tau_2$. However, explicitly expressing the difference between the τ_k in terms of $\text{tr}(T^j)$ is more difficult in higher

dimensions. Let us now examine the initial conditions of the flow equations. When integrating out the auxiliary vielbein e_μ^a , one needs the equations of motion of e_μ^a :

$$B(e^{-1}f)\delta_\nu^\mu - (e^{-1}f)_\alpha^\mu \frac{\partial B}{\partial (e^{-1}f)_\alpha^\nu} = -\frac{1}{\det e} \frac{\delta S_0}{\delta e_\mu^a} e_\nu^a. \quad (27)$$

The right-hand side is finite in the limit $\lambda \rightarrow 0$. Denoting the eigenvalues of $(T^{[0]})^\mu_\nu$ as $\tau_k^{[0]}$, the solution is

$$\alpha_j = \beta_j \left(\lambda (\tau_j^{[0]})^{-1} \left(\prod_{k=1}^d \tau_k^{[0]1/p_k} \right)^{\frac{1}{\Sigma-1}} + 1 \right)^{1/p_j}, \quad (28)$$

which implies that $\alpha_k = \beta_k + O(\lambda)$ when $\lambda \rightarrow 0$. When $\beta_k = 1$, we have $e_\mu^a \rightarrow f_\mu^a$ and the total action $S_{\beta,\lambda} = S_0[\phi, e_\mu^a] + S_{\text{grav}}[e_\mu^a, f_\mu^a]$ reduces to the original action $S_0[\phi, f_\mu^a]$. Equation (28) can be interpreted as the deformed boundary conditions in holography, formulated in terms of eigenvalue variables. In the appendix, we reproduce the root- $T\bar{T}$ deformed boundary conditions proposed in [39].

EXAMPLES

Several deformed field theories can be explored within this framework. A notable example is the ModMax theory [49] and its Born-Infeld-like (MMBI) extension [50]. The ModMax theory is a non-linear conformal- and duality-invariant modification of Maxwell's theory. The MMBI extension maintains the duality invariance, and the action satisfies two commuting flow-equations [5, 51, 52]:

$$\frac{\partial \mathcal{L}_{\text{MMBI}}}{\partial \tilde{\lambda}} = \frac{1}{8} \left(T_{\mu\nu} T^{\mu\nu} - \frac{1}{2} T_\mu{}^\mu T_\nu{}^\nu \right), \quad (29)$$

$$\frac{\partial \mathcal{L}_{\text{MMBI}}}{\partial \tilde{\gamma}} = \frac{1}{2} \sqrt{T_{\mu\nu} T^{\mu\nu} - \frac{1}{4} T_\mu{}^\mu T_\nu{}^\nu}. \quad (30)$$

In the MMBI theory, the stress-energy tensor admits two degenerate eigenvalues τ_1 and τ_2 , each of multiplicity 2. Therefore, the following relations hold:

$$\text{tr}(T^2) - \frac{1}{2}(\text{tr} T)^2 = -4\sqrt{\det T} = -4\tau_1\tau_2, \quad (31)$$

$$\sqrt{\text{tr}(T^2) - \frac{1}{4}(\text{tr} T)^2} = \tau_1 - \tau_2. \quad (32)$$

Turning off the irrelevant deformation momentarily, one can notice that the flow (24) is satisfied in $d = 4$ by fixing $p_k = 2$ for each k , up to rescaling the irrelevant flow parameter. On the other hand, setting $\beta_1 = \beta_2 = e^\gamma = \beta_3^{-1} = \beta_4^{-1}$, the flow equation (26) can be identified with (30), up to a rescaling of γ . This shows that the MMBI flows (29) and (30) can be realized by coupling

Maxwell's theory to the gravity action (21) with $d = 4$ and $p_k = 2$. In this case, (21) simply reduces to

$$S_{\text{grav}}[e_\mu^a, f_\mu^a] = \int d^4x \det e \left[\frac{1}{\lambda} \prod_{k=1}^4 (\alpha_k^2 - \beta_k^2)^{1/2} \right]. \quad (33)$$

Note that the quantities α_k^2 represent the eigenvalues of $g^{\mu\rho}h_{\rho\nu}$: if we switch off the deformation induced by the β_k 's, the corresponding action can be expressed explicitly in terms of the metrics:

$$S_{\text{grav}}[h_{\mu\nu}, g_{\mu\nu}] = \frac{1}{\lambda} \int d^4x \sqrt{\det(h_{\mu\nu} - g_{\mu\nu})}. \quad (34)$$

$T\bar{T}$ -like flows of six-dimensional two-form chiral theories were recently studied in [46]. In these models, T admits two degenerate eigenvalues of multiplicity 3 (throughout the flow), implying that our geometric construction can be straightforwardly implemented. Another example is the higher-dimensional generalized Nambu-Goto action of a self-interacting scalar field in d dimensions:

$$S_\lambda = \int d^d x \left[\frac{1 - \sqrt{1 - 2\lambda(1 - \lambda V)\partial^\mu\phi\partial_\mu\phi}}{\lambda(1 - \lambda V)} - \frac{2V}{1 - \lambda V} \right]. \quad (35)$$

The action (35) satisfies the flow equation

$$\begin{aligned} \frac{\partial S_\lambda}{\partial \lambda} = & \int d^d x \left(\frac{1}{2d} \text{tr}(T^2) - \frac{1}{d^2} (\text{tr} T)^2 \right. \\ & \left. - \frac{d-2}{2\sqrt{d-1}d^{3/2}} \text{tr}(T) \sqrt{\text{tr}(T^2) - \frac{1}{d}(\text{tr} T)^2} \right). \end{aligned} \quad (36)$$

The stress-energy tensor has a non-degenerate eigenvalue τ_1 and a degenerate eigenvalue τ_2 of multiplicity $d-1$. In terms of eigenvalues, the deforming operator can be written as $-\frac{1}{2}\tau_1\tau_2$. Therefore, such deformation can be achieved by setting $\beta_j = 1$, $p_1 = 1$ and $p_{k>2} = d-1$ in (24). It was shown in [45] that the three-dimensional Born-Infeld theory also satisfies the flow equation (36), and one can show that T has a non-degenerate eigenvalue τ_1 and a degenerate eigenvalue τ_2 of multiplicity 2, allowing for a similar description of the flow. Finally, alternative geometric formulations can be constructed when the theory's stress-energy tensor has two distinct degenerate eigenvalues, as described in the appendix.

INCLUSION OF DYNAMICAL GRAVITY

In [47], it was pointed out that a $T\bar{T}$ deformed matter action coupled to the Einstein-Hilbert action is equivalent to an undeformed matter theory coupled to a Ricci-based gravity. Continuing along the same line of thought, we now make the metric h dynamical and include the Einstein-Hilbert term within the first-order Palatini for-

malism. The total action is:

$$S[h, g, \Gamma, \phi] = \frac{1}{2\kappa} \int d^d x \sqrt{\det h} h^{\mu\nu} R_{\mu\nu}(\Gamma) + \int d^d x \sqrt{\det g} B(g^{-1}h) + S_0[g, \phi], \quad (37)$$

where the Ricci curvature tensor is a functional of the connection

$$R_{\mu\nu}(\Gamma) = \partial_\alpha \Gamma_{\nu\mu}^\alpha - \partial_\nu \Gamma_{\alpha\mu}^\alpha + \Gamma_{\alpha\beta}^\alpha \Gamma_{\nu\mu}^\beta - \Gamma_{\nu\beta}^\alpha \Gamma_{\alpha\mu}^\beta. \quad (38)$$

The equations of motion for the connection $\Gamma_{\mu\nu}^\lambda$ lead to the compatibility conditions

$$\Gamma_{\mu\nu}^\lambda = \frac{1}{2} (h^{-1})^{\lambda\alpha} (\partial_\nu h_{\mu\alpha} + \partial_\mu h_{\alpha\nu} - \partial_\alpha h_{\mu\nu}). \quad (39)$$

Integrating out g in the action (37) we get

$$S[h, \Gamma, \phi] = \frac{1}{2\kappa} \int d^d x \sqrt{\det h} h^{\mu\nu} R_{\mu\nu}(\Gamma) + S_{\text{deformed}}[h, \phi], \quad (40)$$

which can be viewed as a deformed matter action S_{deformed} coupled to the standard Einstein-Hilbert action.

To obtain the Ricci-based gravity description, one can integrate out h in the action (37) and obtain,

$$S[g, \Gamma, \phi] = \int d^d x \sqrt{\det g} \mathcal{L}(g^{-1}R) + S_0[g, \phi], \quad (41)$$

$$\mathcal{L} = \left[B(g^{-1}h) - \frac{1}{d-2} g^{\mu\alpha} h_{\alpha\nu} \frac{\partial B}{\partial (g^{\mu\beta} h_{\beta\nu})} \right] \Big|_{h=h^*(g)}, \quad (42)$$

which can be interpreted as an undeformed matter action coupled to a Ricci-based gravity theory $\mathcal{L}(g^{-1}R)$. This procedure yields a dynamical equivalence between an undeformed matter theory coupled to a Ricci-based gravity and a deformed theory coupled to standard general relativity.

It is, however, difficult to obtain an explicit expression for the Lagrangian $\mathcal{L}(g^{-1}R)$ associated with the B function given by (21) because the equations of motion of h are in general very complicated. In the appendix, we derive a flow equation for $\mathcal{L}(g^{-1}R)$:

$$\frac{\partial \mathcal{L}(\rho)}{\partial \lambda} = -(\Sigma - 1) \kappa^{-\frac{\Sigma}{\Sigma-1}} \left(\prod_{i=1}^d \alpha_i \right) \times \left(\prod_{k=1}^d (\alpha_k^{-2} \rho_k - \frac{1}{2} \sum_{j=1}^d \alpha_j^{-2} \rho_j)^{1/p_k} \right)^{\frac{1}{\Sigma-1}}, \quad (43)$$

where we express \mathcal{L} as a function of the eigenvalues ρ_k of $g^{-1}R$, and α_k can be determined through

$$\frac{\partial \mathcal{L}(\rho)}{\partial \rho_k} = \frac{1}{2\kappa \alpha_k^2} \prod_{i=1}^d \alpha_i. \quad (44)$$

The flow equation (43) allows computing the small λ expansion of $\mathcal{L}(g^{-1}R)$ (see the appendix).

Finally, in two dimensions, one can couple (5) to a Jackiw-Teitelboim-like gravity. We find that an undeformed matter theory coupled with a deformed Jackiw-Teitelboim-like gravity is dynamically equivalent to a deformed theory coupled to a Jackiw-Teitelboim-like gravity. The details are given in the appendix.

INCLUDING MARGINAL FLOWS IN GENERAL DEFORMATIONS

One can consider Ricci-based gravity theories associated with more general deformations. For instance, the stress tensor deformation originating from Eddington-inspired Born-Infeld gravity [53] plays a role in $d = 4$ $T\bar{T}$ -like deformations of Abelian gauge theories [47]. For a stress tensor flow driven by an arbitrary operator $f(\tau_i)$ with parameter λ , the associated B function satisfies the flow equation:

$$\frac{\partial B}{\partial \lambda} = f(\tau_i) \prod_{k=1}^d \alpha_k, \quad \tau_i = \alpha_i \partial_{\alpha_i} B \prod_{j=1}^d \alpha_j^{-1}. \quad (45)$$

One can also include marginal flows by replacing $\alpha_i \rightarrow \alpha_i/\beta_i$ with $\prod_{k=1}^d \beta_k = 1$ in the B function. The eigenvalues τ_i are modified as $\tau_i(\alpha_j) \rightarrow \tau_i(\alpha_j/\beta_j)$, and the form of the flow equation (45) remains unchanged. The flow equations with respect to the β -parameters are,

$$\beta_i \frac{\partial B}{\partial \beta_i} - \beta_j \frac{\partial B}{\partial \beta_j} = \alpha_i \frac{\partial B}{\partial \alpha_i} + \alpha_j \frac{\partial B}{\partial \alpha_j} = -(\tau_i - \tau_j) \prod_{k=1}^d \alpha_k. \quad (46)$$

Therefore, it is possible to incorporate commutative marginal flows for any stress tensor deformation that admits a geometric realization. It follows from (43) that the associated Ricci-based gravity action should be modified as $\mathcal{L}(\rho_j) \rightarrow \mathcal{L}(\beta_j^{-2} \rho_j)$.

CONCLUSIONS

This work introduces a geometric formulation for the combination of $T\bar{T}$ and root- $T\bar{T}$ deformations in $d = 2$. We demonstrate that these deformations can be classically formulated by coupling the undeformed theory with a massive gravity action. Additionally, we extend the geometric framework to encompass various stress-energy tensor deformations in higher dimensions. These deformations are related to several well-known theories, including ModMax and its Born-Infeld-like extension. Furthermore, we study the Ricci-based gravities associated with such deformations. These findings might have broad implications in key areas of string theory and holography,

improving our understanding of the effects of stress tensor deformations.

There are several compelling avenues for future exploration stemming from our current work. A natural question is whether our formulation allows for the study of root- $T\bar{T}$ or more general deformations at the quantum level. The quantization of root- $T\bar{T}$ deformed theory poses a complex challenge, and we anticipate that our formulation could offer insights into this intricate issue. One promising approach could involve utilizing the functional renormalization group, as demonstrated in the context of the $T\bar{T}$ deformation [54]. Additionally, it would be intriguing to extend the geometric formulation to encompass more general stress-energy tensor deformations. Another avenue worth exploring is investigating the holographic dictionary of these deformations.

Further, exploring the corresponding realization in celestial holography [55] would be valuable, as proposed in [56], which offers a potential avenue for constructing UV-complete gravity theories. The link between stress-energy flows and classical string or D-brane actions can provide insights into the UV completeness of deformed theories. Consequently, one can envisage constructing counterparts [56, 57] in the framework of celestial holography to investigate their role in UV-complete theories.

ACKNOWLEDGMENTS

The authors thank D. P. Sorokin for his valuable discussions and comments. H. B-A. expresses his gratitude to the Department of Physics and Astronomy "Galileo Galilei" at the University of Padova for their generous hospitality during the concluding phase of this project. S. H. acknowledges financial support from the Max Planck Partner Group, the Fundamental Research Funds for the Central Universities, and the Natural Science Foundation of China Grants No. 12075101 and No. 12235016. H. O. is supported by the National Natural Science Foundation of China, Grant No. 12205115, and by the Science and Technology Development Plan Project of Jilin Province of China, Grant No. 20240101326JC. T. M. and R. T. received partial support from the INFN project SFT and the PRIN Project No. 2022ABPB EY, with the title "Understanding quantum field theory through its deformations".

* h.babaei@uma.ac.ir

† hesong@jlu.edu.cn

‡ tommaso.morone@unito.it

§ haoouyang@jlu.edu.cn

¶ roberto.tateo@unito.it

- [1] A. Cavaglià, S. Negro, I. M. Szécsényi and R. Tateo, " $T\bar{T}$ -deformed 2D Quantum Field Theories," *JHEP* **10**, 112 (2016) [arXiv:1608.05534 [hep-th]].
- [2] F. A. Smirnov and A. B. Zamolodchikov, "On space of integrable quantum field theories," *Nucl. Phys. B* **915**, 363-383 (2017) [arXiv:1608.05499 [hep-th]].
- [3] A. B. Zamolodchikov, "Expectation value of composite field $T\bar{T}$ in two-dimensional quantum field theory," [arXiv:hep-th/0401146 [hep-th]].
- [4] G. Bonelli, N. Doroud and M. Zhu, " $T\bar{T}$ -deformations in closed form," *JHEP* **06**, 149 (2018) [arXiv:1804.10967 [hep-th]].
- [5] R. Conti, L. Iannella, S. Negro and R. Tateo, "Generalised Born-Infeld models, Lax operators and the $T\bar{T}$ perturbation," *JHEP* **11**, 007 (2018) [arXiv:1806.11515 [hep-th]].
- [6] R. Conti, S. Negro and R. Tateo, "The $T\bar{T}$ perturbation and its geometric interpretation," *JHEP* **02**, 085 (2019) [arXiv:1809.09593 [hep-th]].
- [7] J. Cardy, "The $T\bar{T}$ deformation of quantum field theory as random geometry," *JHEP* **10**, 186 (2018) [arXiv:1801.06895 [hep-th]].
- [8] S. Datta and Y. Jiang, " $T\bar{T}$ deformed partition functions," *JHEP* **08**, 106 (2018) [arXiv:1806.07426 [hep-th]].
- [9] O. Aharony, S. Datta, A. Giveon, Y. Jiang and D. Kutasov, "Modular invariance and uniqueness of $T\bar{T}$ deformed CFT," *JHEP* **01**, 086 (2019) [arXiv:1808.02492 [hep-th]].
- [10] S. He, Y. Sun and Y. X. Zhang, " $T\bar{T}$ -flow effects on torus partition functions," *JHEP* **09**, 061 (2021) [arXiv:2011.02902 [hep-th]].
- [11] M. Baggio and A. Sfondrini, "Strings on NS-NS Backgrounds as Integrable Deformations," *Phys. Rev. D* **98**, no.2, 021902 (2018) [arXiv:1804.01998 [hep-th]].
- [12] A. Dei and A. Sfondrini, "Integrable S matrix, mirror TBA and spectrum for the stringy $\text{AdS}_3 \times \text{S}^3 \times \text{S}^3 \times \text{S}^1$ WZW model," *JHEP* **02**, 072 (2019) [arXiv:1812.08195 [hep-th]].
- [13] S. Chakraborty, A. Giveon and D. Kutasov, " $T\bar{T}$, $J\bar{T}$, $T\bar{J}$ and String Theory," *J. Phys. A* **52**, no.38, 384003 (2019) [arXiv:1905.00051 [hep-th]].
- [14] N. Callebaut, J. Kruthoff and H. Verlinde, " $T\bar{T}$ deformed CFT as a non-critical string," *JHEP* **04**, 084 (2020) [arXiv:1910.13578 [hep-th]].
- [15] L. McGough, M. Mezei and H. Verlinde, "Moving the CFT into the bulk with $T\bar{T}$," *JHEP* **04**, 010 (2018) [arXiv:1611.03470 [hep-th]].
- [16] P. Kraus, J. Liu and D. Marolf, "Cutoff AdS_3 versus the $T\bar{T}$ deformation," *JHEP* **07**, 027 (2018) [arXiv:1801.02714 [hep-th]].
- [17] W. Cottrell and A. Hashimoto, "Comments on $T\bar{T}$ double trace deformations and boundary conditions," *Phys. Lett. B* **789**, 251-255 (2019) [arXiv:1801.09708 [hep-th]].
- [18] M. Taylor, " $T\bar{T}$ deformations in general dimensions," *Adv. Theor. Math. Phys.* **27**, no.1, 37-63 (2023) [arXiv:1805.10287 [hep-th]].
- [19] T. Hartman, J. Kruthoff, E. Shaghoulian and A. Tajdini, "Holography at finite cutoff with a T^2 deformation," *JHEP* **03**, 004 (2019) [arXiv:1807.11401 [hep-th]].
- [20] P. Caputa, S. Datta and V. Shyam, "Sphere partition functions & cut-off AdS," *JHEP* **05**, 112 (2019) [arXiv:1902.10893 [hep-th]].
- [21] M. Guica and R. Monten, " $T\bar{T}$ and the mirage of a bulk cutoff," *SciPost Phys.* **10**, no.2, 024 (2021)

- [arXiv:1906.11251 [hep-th]].
- [22] D. J. Gross, J. Kruthoff, A. Rolph and E. Shaghoulian, “ $T\bar{T}$ in AdS₂ and Quantum Mechanics,” *Phys. Rev. D* **101**, no.2, 026011 (2020) [arXiv:1907.04873 [hep-th]].
- [23] Y. Li and Y. Zhou, “Cutoff AdS₃ versus $T\bar{T}$ CFT₂ in the large central charge sector: correlators of energy-momentum tensor,” *JHEP* **12**, 168 (2020) [arXiv:2005.01693 [hep-th]].
- [24] S. He, Y. Li, Y. Z. Li and Y. Zhang, “Holographic torus correlators of stress tensor in AdS₃/CFT₂,” *JHEP* **06**, 116 (2023) [arXiv:2303.13280 [hep-th]].
- [25] S. He, Y. Z. Li and Y. Zhang, “Holographic torus correlators in AdS₃ gravity coupled to scalar field,” [arXiv:2311.09636 [hep-th]].
- [26] S. Dubovsky, V. Gorbenko and M. Mirbabayi, “Asymptotic fragility, near AdS₂ holography and $T\bar{T}$,” *JHEP* **09**, 136 (2017) [arXiv:1706.06604 [hep-th]].
- [27] S. Dubovsky, V. Gorbenko and G. Hernández-Chifflet, “ $T\bar{T}$ partition function from topological gravity,” *JHEP* **09**, 158 (2018) [arXiv:1805.07386 [hep-th]].
- [28] A. J. Tolley, “ $T\bar{T}$ deformations, massive gravity and non-critical strings,” *JHEP* **06**, 050 (2020) [arXiv:1911.06142 [hep-th]].
- [29] L. V. Iliesiu, J. Kruthoff, G. J. Turiaci and H. Verlinde, “JT gravity at finite cutoff,” *SciPost Phys.* **9**, 023 (2020) [arXiv:2004.07242 [hep-th]].
- [30] S. Okumura and K. Yoshida, “ $T\bar{T}$ -deformation and Liouville gravity,” *Nucl. Phys. B* **957**, 115083 (2020) [arXiv:2003.14148 [hep-th]].
- [31] S. Ebert, C. Ferko, H. Y. Sun and Z. Sun, “ $T\bar{T}$ in JT Gravity and BF Gauge Theory,” *SciPost Phys.* **13**, no.4, 096 (2022) [arXiv:2205.07817 [hep-th]].
- [32] Y. Jiang, “A pedagogical review on solvable irrelevant deformations of 2D quantum field theory,” *Commun. Theor. Phys.* **73**, no.5, 057201 (2021) [arXiv:1904.13376 [hep-th]].
- [33] P. Caputa, P. Caputa, S. Datta, S. Datta, Y. Jiang, Y. Jiang, P. Kraus and P. Kraus, “Geometrizing $T\bar{T}$,” *JHEP* **03**, 140 (2021) [erratum: *JHEP* **09**, 110 (2022)] [arXiv:2011.04664 [hep-th]].
- [34] S. Dubovsky, S. Negro and M. Porrati, “Topological gauging and double current deformations,” *JHEP* **05**, 240 (2023) [arXiv:2302.01654 [hep-th]].
- [35] R. Conti, J. Romano and R. Tateo, “Metric approach to a $T\bar{T}$ -like deformation in arbitrary dimensions,” *JHEP* **09**, 085 (2022) [arXiv:2206.03415 [hep-th]].
- [36] H. Babaei-Aghbolagh, K. Babaei Velni, D. Mahdavian Yekta and H. Mohammadzadeh, “Marginal $T\bar{T}$ -like deformation and modified Maxwell theories in two dimensions,” *Phys. Rev. D* **106**, no.8, 086022 (2022) [arXiv:2206.12677 [hep-th]].
- [37] C. Ferko, A. Sfondrini, L. Smith and G. Tartaglino-Mazzucchelli, “Root- $T\bar{T}$ Deformations in Two-Dimensional Quantum Field Theories,” *Phys. Rev. Lett.* **129**, no.20, 201604 (2022) [arXiv:2206.10515 [hep-th]].
- [38] R. Borsato, C. Ferko and A. Sfondrini, “Classical integrability of root- $T\bar{T}$ flows,” *Phys. Rev. D* **107**, no.8, 086011 (2023) [arXiv:2209.14274 [hep-th]].
- [39] S. Ebert, C. Ferko and Z. Sun, “Root- $T\bar{T}$ deformed boundary conditions in holography,” *Phys. Rev. D* **107**, no.12, 12 (2023) [arXiv:2304.08723 [hep-th]].
- [40] J. Tian, T. Lai and F. Omid, “Modular transformations of on-shell actions of (root-) $T\bar{T}$ deformed holographic CFTs,” [arXiv:2404.16354 [hep-th]].
- [41] H. Babaei-Aghbolagh, K. Babaei Velni, D. M. Yekta and H. Mohammadzadeh, “ $T\bar{T}$ -like flows in non-linear electrodynamic theories and S-duality,” *JHEP* **04**, 187 (2021) [arXiv:2012.13636 [hep-th]].
- [42] H. Babaei-Aghbolagh, K. Babaei Velni, D. M. Yekta and H. Mohammadzadeh, “Manifestly SL(2, R) Duality-Symmetric Forms in ModMax Theory,” *JHEP* **12**, 147 (2022) [arXiv:2210.13196 [hep-th]].
- [43] C. Ferko, S. M. Kuzenko, L. Smith and G. Tartaglino-Mazzucchelli, “Duality-invariant nonlinear electro-dynamics and stress tensor flows,” *Phys. Rev. D* **108**, no.10, 106021 (2023) [arXiv:2309.04253 [hep-th]].
- [44] T. Flöss, D. Roest and T. Westerdijk, “Non-linear electro-dynamics from massive gravity,” *JHEP* **02**, 194 (2024) [arXiv:2308.04349 [hep-th]].
- [45] C. Ferko, Y. Hu, Z. Huang, K. Koutrolikos and G. Tartaglino-Mazzucchelli, “ $T\bar{T}$ -like flows and $3d$ non-linear supersymmetry,” *SciPost Phys.* **16**, no.1, 038 (2024) [arXiv:2302.10410 [hep-th]].
- [46] C. Ferko, S. M. Kuzenko, K. Lechner, D. P. Sorokin and G. Tartaglino-Mazzucchelli, “Interacting Chiral Form Field Theories and $T\bar{T}$ -like Flows in Six and Higher Dimensions,” [arXiv:2402.06947 [hep-th]].
- [47] T. Morone, S. Negro and R. Tateo, “Gravity and $T\bar{T}$ flows in higher dimensions,” [arXiv:2401.16400 [hep-th]].
- [48] G. J. Olmo and D. Rubiera-Garcia, “Some recent results on Ricci-based gravity theories,” *Int. J. Mod. Phys. D* **31**, no.11, 2240012 (2022) [arXiv:2203.04116 [gr-qc]].
- [49] I. Bandos, K. Lechner, D. Sorokin and P. K. Townsend, “A non-linear duality-invariant conformal extension of Maxwell’s equations,” *Phys. Rev. D* **102**, 121703 (2020) [arXiv:2007.09092 [hep-th]].
- [50] I. Bandos, K. Lechner, D. Sorokin and P. K. Townsend, “On p-form gauge theories and their conformal limits,” *JHEP* **03**, 022 (2021) [arXiv:2012.09286 [hep-th]].
- [51] H. Babaei-Aghbolagh, K. B. Velni, D. M. Yekta and H. Mohammadzadeh, “Emergence of non-linear electrodynamic theories from $T\bar{T}$ -like deformations,” *Phys. Lett. B* **829**, 137079 (2022) [arXiv:2202.11156 [hep-th]].
- [52] C. Ferko, L. Smith and G. Tartaglino-Mazzucchelli, “On Current-Squared Flows and ModMax Theories,” *SciPost Phys.* **13**, no.2, 012 (2022) [arXiv:2203.01085 [hep-th]].
- [53] J. Beltran Jimenez, L. Heisenberg, G. J. Olmo and D. Rubiera-Garcia, “Born-Infeld inspired modifications of gravity,” *Phys. Rept.* **727**, 1-129 (2018) [arXiv:1704.03351 [gr-qc]].
- [54] J. Liu, J. Haruna and M. Yamada, “Nonperturbative aspects of two-dimensional $T\bar{T}$ -deformed scalar theory from functional renormalization group,” *Phys. Rev. D* **109**, no.6, 6 (2024) [arXiv:2309.15584 [hep-th]].
- [55] S. Pasterski, S. H. Shao and A. Strominger, “Flat Space Amplitudes and Conformal Symmetry of the Celestial Sphere,” *Phys. Rev. D* **96**, no.6, 065026 (2017) [arXiv:1701.00049 [hep-th]].
- [56] S. He, P. Mao and X. C. Mao, “ $T\bar{T}$ deformed soft theorem,” *Phys. Rev. D* **107**, no.10, L101901 (2023) [arXiv:2209.01953 [hep-th]].
- [57] S. He, P. Mao and X. C. Mao, “Loop corrections versus marginal deformation in celestial holography,” [arXiv:2307.02743 [hep-th]].

Appendix

Metric formulation

This section presents the geometric formulation of the combination of $T\bar{T}$ and root- $T\bar{T}$ deformation in the metric approach. One can choose a gauge such that

$$e^{-1}f = \sqrt{g^{-1}h}. \quad (47)$$

The gravity action S_{grav} can be expressed in terms of the metrics $g_{\mu\nu}$ and $h_{\mu\nu}$ by using the relation

$$\det(\sqrt{g^{-1}h}) = \sqrt{(\det g)^{-1} \det(h)} = \frac{1}{2} \left[\text{tr}(\sqrt{g^{-1}h}) \right]^2 - \frac{1}{2} \text{tr}(g^{-1}h). \quad (48)$$

We have

$$y_1 = \text{tr}(\sqrt{g^{-1}h}) = \sqrt{z_1 + \sqrt{2z_1^2 - 2z_2}}, \quad (49)$$

$$y_2 = \text{tr}(g^{-1}h) = z_1, \quad (50)$$

where we denote $z_n = \text{tr}[(g^{-1}h)^n]$. Then the gravity action S_{grav} can be written as

$$S_{\text{grav}}[g, h] = \frac{1}{2\lambda} \int d^2x \sqrt{\det g} \left(2\sqrt{z_1 - \sqrt{2z_1^2 - 2z_2}} \sinh \frac{\gamma}{2} - 2\sqrt{\sqrt{2z_1^2 - 2z_2} + z_1} \cosh \frac{\gamma}{2} + \sqrt{2z_1^2 - 2z_2} + 2 \right). \quad (51)$$

One can also verify the flow equations in the metric formulation. The stress-energy tensor and related quantities are

$$T_{\nu}^{\mu} = 2 \frac{1}{\sqrt{\det h}} \frac{\delta S_{\gamma, \lambda}}{\delta h_{\mu\alpha}} h_{\alpha\nu} = 2 \frac{1}{\sqrt{\det h}} \frac{\delta S_{\text{grav}}}{\delta h_{\mu\alpha}} h_{\alpha\nu}, \quad (52)$$

$$T_{\nu}^{\nu} = \frac{\sqrt{2}\sqrt{z_1 - \sqrt{2z_1^2 - 2z_2}} \sinh \frac{\gamma}{2}}{\lambda\sqrt{z_1^2 - z_2}} - \frac{\sqrt{2}\sqrt{\sqrt{2z_1^2 - 2z_2} + z_1} \cosh \frac{\gamma}{2}}{\lambda\sqrt{z_1^2 - z_2}} + \frac{2}{\lambda}, \quad (53)$$

$$T_{\nu}^{\mu} T_{\mu}^{\nu} = \frac{2\sqrt{2}\sqrt{z_1 - \sqrt{2z_1^2 - 2z_2}} \sinh \frac{\gamma}{2}}{\lambda^2\sqrt{z_1^2 - z_2}} - \frac{2\sqrt{2}\sqrt{\sqrt{2z_1^2 - 2z_2} + z_1} \cosh \frac{\gamma}{2}}{\lambda^2\sqrt{z_1^2 - z_2}} \quad (54)$$

$$+ \frac{-2 \left(\sqrt{2z_2 - z_1^2} \sinh \gamma + z_1 (\cosh \gamma + z_1) - z_2 \right)}{\lambda^2 (z_1^2 - z_2)}. \quad (55)$$

The action satisfies the flow equations

$$\frac{\partial S_{\gamma, \lambda}}{\partial \lambda} = \int d^2x \sqrt{\det h} \left(\frac{1}{2} T_{\nu}^{\mu} T_{\mu}^{\nu} - \frac{1}{2} (T_{\mu}^{\mu})^2 \right), \quad (56)$$

$$\frac{\partial S_{\gamma, \lambda}}{\partial \gamma} = \int d^2x \sqrt{\det h} \sqrt{\frac{1}{2} T_{\nu}^{\mu} T_{\mu}^{\nu} - \frac{1}{4} (T_{\mu}^{\mu})^2}. \quad (57)$$

Holographic boundary conditions

When e_{μ}^a is on-shell, one can express α_j and τ_j in terms of $\tau_k^{[0]}$ as:

$$\alpha_j = \beta_j \left(\lambda (\tau_j^{[0]})^{-1} \left(\prod_{k=1}^d \tau_k^{[0]1/p_k} \right)^{\frac{1}{\Sigma-1}} + 1 \right)^{1/p_j}, \quad (58)$$

$$\tau_j = \left(\lambda \left(\prod_{k=1}^d \tau_k^{[0]1/p_k} \right)^{\frac{1}{\Sigma-1}} + \tau_j^{[0]} \right) \prod_{k=1}^d \alpha_k^{-1}. \quad (59)$$

To derive the root- $T\bar{T}$ deformed boundary conditions, we take $d = 2$, $p_k = 1$, $\beta_1 = e^{\frac{\gamma}{2}}$, $\beta_2 = e^{-\frac{\gamma}{2}}$ and $\lambda \rightarrow 0$. We get, $\alpha_j = \beta_j$ and $\tau_j = \tau_j^{[0]}$. We write $T^{[0]\mu}_\nu$ explicitly:

$$T^{[0]\mu}_\nu = \begin{pmatrix} a & b \\ c & d \end{pmatrix}, \quad (60)$$

which can be diagonalized by

$$U = \begin{pmatrix} \sqrt{a^2 - 2ad + 4bc + d^2} + a - d & -\sqrt{a^2 - 2ad + 4bc + d^2} + a - d \\ 2c & 2c \end{pmatrix}. \quad (61)$$

Since U also diagonalises $g^{-1}h$, we get

$$g^{-1}h = U \text{diag}(e^\gamma, e^{-\gamma})U^{-1} = \begin{pmatrix} \frac{(a-d)\sinh(\gamma)}{\sqrt{(a-d)^2 + 4bc}} + \cosh(\gamma) & \frac{2b\sinh(\gamma)}{\sqrt{(a-d)^2 + 4bc}} \\ \frac{2c\sinh(\gamma)}{\sqrt{(a-d)^2 + 4bc}} & \frac{(d-a)\sinh(\gamma)}{\sqrt{(a-d)^2 + 4bc}} + \cosh(\gamma) \end{pmatrix}, \quad (62)$$

$$\Rightarrow h_{\mu\nu} = g_{\mu\nu} \cosh \gamma + \frac{\sinh \gamma}{\sqrt{\frac{1}{2}T^{[0]\mu}_\nu T^{[0]\nu}_\mu - \frac{1}{4}(T^{[0]\mu}_\mu)^2}} \tilde{T}^{[0]}_{\mu\nu}, \quad (63)$$

where $\tilde{T}^{[0]}$ is the traceless part of $T^{[0]}$. The deformed boundary conditions for the stress-energy tensors are $T^\mu_\nu = T^{[0]\mu}_\nu$.

On-shell geometric formulation of MMBI and generalized Nambu-Goto

When the stress-energy tensor of the undeformed theory has two degenerate eigenvalues $\tau_1^{[0]}$ of multiplicity d_1 and $\tau_2^{[0]}$ of multiplicity d_2 , we propose an alternative gravity action:

$$S_{\text{grav}} = \frac{1}{\lambda^{\Sigma-1}} \int d^d x \left[\left(\frac{e^{\frac{\gamma}{d_2}} \left(d_2 y_1 \mp \sqrt{d_1 d_2 ((d_1 + d_2) y_2 - y_1^2)} \right)}{d_2 (d_1 + d_2)} \right)^{p_2} - 1 \right]^{\frac{d_2}{p_2}} \times \left[\left(\frac{e^{-\frac{\gamma}{d_1}} \left(d_1 y_1 \pm \sqrt{d_1 d_2 ((d_1 + d_2) y_2 - y_1^2)} \right)}{d_1 (d_1 + d_2)} \right)^{p_1} - 1 \right]^{\frac{d_1}{p_1}}. \quad (64)$$

When e^a_μ is on-shell, the eigenvalues α_k and τ_k have the same degenerate structure as $\tau_k^{[0]}$. One finds

$$\frac{\partial S_{\text{grav}}}{\partial \lambda} = -(\Sigma - 1) \int d^d x \det f(\tau_1^{d_1/p_1} \tau_2^{d_2/p_2})^{\frac{1}{\Sigma-1}}, \quad (65)$$

$$\frac{\partial S_{\text{grav}}}{\partial \gamma} = \int d^d x \det f(\tau_2 - \tau_1). \quad (66)$$

For the MMBI case, we take $d_k = p_k = 2$ and (64) becomes

$$S_{\text{grav}} = \frac{1}{\lambda} \int d^4 x \det e \left[1 - \frac{y_2}{2} \cosh \gamma \pm \frac{y_1}{4} \sqrt{4y_2 - y_1^2} \sinh \gamma + \left(\frac{y_1^2}{8} - \frac{y_2}{4} \right)^2 \right]. \quad (67)$$

For the generalized Nambu-Goto case, we take $d_1 = p_1 = 1$, $d_2 = p_2 = d - 1$, and (64) becomes

$$S_{\text{grav}} = \frac{1}{\lambda} \int d^3 x \det e \left[\left(\frac{(d-1)y_1 \mp \sqrt{(d-1)(dy_2 - y_1^2)}}{(d-1)d} \right)^{d-1} - 1 \right] \left(\pm \frac{\sqrt{(d-1)(dy_2 - y_1^2)} + y_1}{d} - 1 \right). \quad (68)$$

Flow equations of $\mathcal{L}(g^{-1}R)$

In this section, we derive flow equations of $\mathcal{L}(g^{-1}R)$, which enable us to compute the perturbative expansion of $\mathcal{L}(g^{-1}R)$ in small deformation parameters. When total action is extremized with respect to $h_{\mu\nu}$, only explicit dependencies must be considered when differentiating the Lagrangian with respect to fields or parameters. We have

$$\frac{\partial}{\partial R_{\mu\nu}} \left(\sqrt{\det g} \mathcal{L}(g^{-1}R) \right) = \frac{\partial}{\partial R_{\mu\nu}} \left(\frac{1}{2\kappa} \sqrt{\det h} h^{\alpha\beta} R_{\alpha\beta} \right), \quad (69)$$

$$\frac{\partial}{\partial \lambda} \left(\sqrt{\det g} \mathcal{L}(g^{-1}R) \right) = \frac{\partial}{\partial \lambda} \left(\sqrt{\det g} B(g^{-1}h) \right). \quad (70)$$

It follows from the equations of motion of $h_{\mu\nu}$ that

$$g^{\mu\alpha} R_{\alpha\nu} = \kappa g^{\mu\alpha} h_{\alpha\beta} T^\beta_\nu - \frac{\kappa}{d-2} g^{\mu\alpha} h_{\alpha\nu} T^\beta_\beta, \quad (71)$$

and therefore $g^{\mu\alpha} R_{\alpha\nu}$ and $g^{\mu\alpha} h_{\alpha\nu}$ can be diagonalized simultaneously. Denoting ρ_k as the eigenvalues of $g^{\mu\alpha} R_{\alpha\nu}$ and writing \mathcal{L} as a function of ρ_k , equation (71) can be written as

$$\kappa \tau_k = \alpha_k^{-2} \rho_k - \frac{1}{2} \sum_{j=1}^d \alpha_j^{-2} \rho_j, \quad (72)$$

and equation (69) leads to

$$\frac{\partial \mathcal{L}(\rho)}{\partial \rho_k} = \frac{1}{2\kappa \alpha_k^2} \prod_{i=1}^d \alpha_i \Rightarrow \alpha_i = (2\kappa)^{\frac{1}{d-2}} \left(\frac{\partial \mathcal{L}(\rho)}{\partial \rho_k} \right)^{-1} \left(\prod_{i=1}^d \frac{\partial \mathcal{L}(\rho)}{\partial \rho_k} \right)^{\frac{1}{2d-4}}. \quad (73)$$

Using (24), (70) and (72), we get the flow equation of $\mathcal{L}(\rho)$ with respect to λ :

$$\frac{\partial \mathcal{L}(\rho)}{\partial \lambda} = -(\Sigma - 1) \kappa^{-\frac{\Sigma}{\Sigma-1}} \left(\prod_{i=1}^d \alpha_i \right) \left(\prod_{k=1}^d (\alpha_k^{-2} \rho_k - \frac{1}{2} \sum_{j=1}^d \alpha_j^{-2} \rho_j)^{1/p_k} \right)^{\frac{1}{\Sigma-1}}, \quad (74)$$

where one should also substitute (73) into the right-hand site. The initial condition can be obtained using the limit $\lambda \rightarrow 0$. We have:

$$\alpha_k = \beta_k + O(\lambda) \quad (75)$$

$$\mathcal{L}(\rho) = \frac{1}{2\kappa} \sum_{k=1}^d \beta_k^{-2} \rho_k + O(\lambda). \quad (76)$$

Using the flow equation and the initial condition, we can recursively solve the λ expansion of $\mathcal{L}(\rho)$. Up to order λ we find

$$\mathcal{L}(\rho) = \frac{1}{2\kappa} \sum_{k=1}^d \beta_k^{-2} \rho_k - \lambda (\Sigma - 1) \kappa^{-\frac{\Sigma}{\Sigma-1}} \left(\prod_{k=1}^d (\beta_k^{-2} \rho_k - \frac{1}{2} \sum_{j=1}^d \beta_j^{-2} \rho_j)^{1/p_k} \right)^{\frac{1}{\Sigma-1}} + O(\lambda^2). \quad (77)$$

For pure $(\det T)^{\frac{1}{d-p}}$ deformations with $\beta_k = 1$ and $p_k = p$, the expression reduces to

$$\begin{aligned} \mathcal{L} &= \frac{1}{2\kappa} \sum_{k=1}^d \rho_k - \lambda (d/p - 1) \kappa^{-\frac{d}{d-p}} \prod_{k=1}^d \left(\rho_k - \frac{1}{2} \sum_{j=1}^d \rho_j \right)^{\frac{1}{d-p}} + O(\lambda^2) \\ &= \frac{1}{2\kappa} \text{tr}(g^{-1}R) - \lambda (d/p - 1) \kappa^{-\frac{d}{d-p}} \det(g^{-1}R - \frac{1}{2} \text{tr}(g^{-1}R))^{1/p} + O(\lambda^2). \end{aligned} \quad (78)$$

However, for more general deformations, it is difficult to express the eigenvalues ρ_k in terms of $\text{tr}[(g^{-1}R)^k]$ explicitly.

Coupling to flat Jackiw-Teitelboim-like gravity action in two dimensions

In two dimensions, we couple the action (4) to a flat space Jackiw-Teitelboim-like gravity action in the first-order formalism for the zweibein f_μ^a and a vacuum energy term:

$$S = S_{JT} - \Lambda \int d^2x \det f + \int d^2x \det e B(e^{-1}f) + S_0[\phi, e_\mu^a], \quad (79)$$

$$S_{JT} = \frac{1}{\kappa} \int d^2x \epsilon^{\alpha\beta} (\epsilon_{ac} \sigma^c (\partial_\alpha f_\beta^a - \epsilon^a_b \omega_\alpha f_\beta^b) + \varphi \partial_\alpha \omega_\beta). \quad (80)$$

The equation of motion for f gives

$$\frac{1}{\kappa} \epsilon^{\alpha\beta} \epsilon_{ab} u_\beta^b - \Lambda \epsilon^{\alpha\beta} \epsilon_{ab} f_\beta^b + \det e \frac{\partial B}{\partial f_\alpha^a} = 0, \quad (81)$$

where we defined $u_\alpha^a = \partial_\alpha \sigma^a - \epsilon^a_c \sigma^c \omega_\alpha$. The solution is

$$f^* = \frac{\sqrt{2w_2 - w_1^2} w_1 \sinh \frac{\gamma}{2} + (w_1^2 - 2w_2) \cosh \frac{\gamma}{2}}{(w_1^2 - 2w_2)(1 - \lambda\Lambda)} e - \frac{\lambda - \frac{2\kappa \sinh \frac{\gamma}{2}}{\sqrt{2w_2 - w_1^2}}}{\kappa - \kappa\lambda\Lambda} u, \quad (82)$$

where $w_n = \text{tr}[(e^{-1}u)^n]$. In terms of y_n , we find

$$y_1^* = \frac{2\kappa \cosh \frac{\gamma}{2} - \lambda w_1}{\kappa - \kappa\lambda\Lambda}, \quad y_2^* = \frac{2\kappa^2 \cosh \gamma + \lambda \left(-2\kappa \sqrt{2w_2 - w_1^2} \sinh \frac{\gamma}{2} - 2\kappa w_1 \cosh \frac{\gamma}{2} + \lambda w_2 \right)}{\kappa^2 (\lambda\Lambda - 1)^2}. \quad (83)$$

Substituting back into (79), we get

$$S = \int d^2x \det e \left(\frac{2\kappa^2 \lambda\Lambda + 4\kappa^2 \cosh \gamma - 4\kappa^2 + 2\kappa \lambda \sqrt{2w_2 - w_1^2} \sinh \frac{\gamma}{2} - 2\kappa \lambda w_1 \cosh \frac{\gamma}{2} + \lambda^2 w_1^2 - \lambda^2 w_2}{2\kappa^2 \lambda (\lambda\Lambda - 1)} \right) + \int d^2x \epsilon^{\alpha\beta} \varphi \partial_\alpha \omega_\beta + S_0. \quad (84)$$

Denoting the eigenvalues of $e^{-1}u$ as ν_k , the action can be simplified as

$$S = \int d^2x \det e \left(\frac{\kappa^2 \lambda\Lambda + \kappa^2 e^{-\gamma} + \kappa^2 e^\gamma - 2\kappa^2 - \kappa \lambda e^{\gamma/2} \nu_1 - \kappa \lambda e^{-\gamma/2} \nu_2 + \lambda^2 \nu_1 \nu_2}{\kappa^2 \lambda (\lambda\Lambda - 1)} \right) + \int d^2x \epsilon^{\alpha\beta} \varphi \partial_\alpha \omega_\beta + S_0, \quad (85)$$

which can be interpreted as a matter theory S_0 coupled to a deformed Jackiw-Teitelboim-like gravity. Alternatively, integrating out the vielbein e_μ^a in the action (79) results in a matter theory that is deformed by both $T\bar{T}$ and root- $T\bar{T}$, coupled to Jackiw-Teitelboim-like gravity. Consequently, the dynamics of a matter theory S_0 coupled to a deformed Jackiw-Teitelboim-like gravity is equivalent to that of a matter theory subjected to $T\bar{T}$ and root- $T\bar{T}$ deformations coupled to Jackiw-Teitelboim-like gravity.