



Data Article

Quantitative variation of male and female-specific compounds in 99 drosophilid flies



Mohammed A. Khallaf^{a,b,1,*}, Melissa Diaz-Morales^{a,c},
Bill S. Hansson^a, Markus Knaden^{a,*}

^a Department of Evolutionary Neuroethology, Max Planck Institute for Chemical Ecology, Jena, Germany

^b Department of Zoology and Entomology, Faculty of Science, Assiut University, Assiut, Egypt

^c Lankester Botanical Garden, University of Costa Rica, Cartago, Costa Rica

ARTICLE INFO

Article history:

Received 13 July 2024

Revised 15 August 2024

Accepted 19 August 2024

Available online 3 September 2024

Dataset link: [Quantitative variation of male and female-specific compounds in 99 drosophilid flies \(Original data\)](#)

Keywords:

Pheromone

Drosophila

Male-specific compound

Female-specific compound

Thermal desorption–gas chromatography–mass spectrometry

Courtship

Female receptivity

ABSTRACT

Variation in sex pheromones is regarded as one of the causes of reproductive isolation and speciation. We recently identified 51 male- and female-specific compounds – many of which function as sex pheromones – in 99 drosophilid species [1]. Here, we report that despite many of these compounds being shared between species, their quantities differ significantly. For example, although 34 drosophilid species share the male-specific compound cis-vaccenyl acetate (cVA), which plays a critical role in regulating various social and sexual behaviors, the amount of cVA can differ by up to 600-fold between different species. Additionally, we found 7-tricosene, the cuticular hydrocarbon pheromone, present in 35 *Drosophila* species. Our findings indicate that 7-tricosene is equally present in both sexes of 14 species, more abundant in males of 14 species, and more abundant in females of 7 species. We provide raw data on the concentration of potential pheromone components in the 99 drosophilids, which can provide important insights for further research on the behavior and evolution of these species. Quantitative varia-

* Corresponding authors at: Department of Evolutionary Neuroethology, Max Planck Institute for Chemical Ecology, Jena, Germany.

E-mail addresses: Mohammed.khallaf@mdc-berlin.de (M.A. Khallaf), Mknaden@ice.mpg.de (M. Knaden).

Social media: [@khalaf13](#) (M.A. Khallaf), [@billhansson](#) (B.S. Hansson), [@mknaden1](#) (M. Knaden)

¹ Present address: Department of Neuroscience, Max Delbrück Center for Molecular Medicine, Berlin D-13122, Germany.

tions highlight species-specific patterns, suggesting an additional mechanism for reproductive isolation built on specific combinations of compounds at set concentrations.

© 2024 The Author(s). Published by Elsevier Inc.

This is an open access article under the CC BY license

(<http://creativecommons.org/licenses/by/4.0/>)

Specifications Table

Subject	Biological sciences
Specific subject area	Sex pheromones in <i>Drosophila</i>
Type of data	Raw data and figures
How the data were acquired	Thermal desorption-gas chromatography-mass spectrometry (TD-GC-MS); (Agilent GC 7890A fitted with an MS 5975C inert XL MSD unit; www.agilent.com) equipped with an HP5-MS UI column (19091S-433UI; Agilent Technologies). Analysis: Enhanced ChemsStation (MSD ChemStation F.01.03.2357). Library: NIST MS Search 2.2
Data format	Raw, Analyzed.
Description of data collection	TD-GC-MS: HP5-MS UI column (19091S-433UI; Agilent Technologies) No solvent Thermal desorption: Temperature 250 °C for 3 min. Trap: Temperature -50 °C using liquid nitrogen Vaporizer injector: Ramp to 270 °C (12 °C/s) and held for 5 min. The oven program: Initial temperature 50 °C for 3 min, ramp to 250 °C (15 °C/min) and held for 3 min, and then to 280 °C (20 °C/min) and held for 30 min. For MS, the transfer line, source, and quad: Temperature 260 °C, 230 °C, and 150 °C, respectively. Ion source: Electron ionization (EI) operating at 70 eV energy. Mass spectra: m/z 33 to 500.
Data source location	Institution: Max Planck institute for chemical ecology City/Town/Region: Jena Country: Germany
Data accessibility	The data are available within this article and in supplementary Table 1.
Related research article	Author's name: Mohammed A. Khallaf, Rongfeng Cui, Jerrit Weißflog, Maide Erdogmus, Aleš Svatoš, Hany K. M. Dweck, Dario Riccardo Valenzano, Bill S. Hansson & Markus Knaden Title: Large-scale characterization of sex pheromone communication systems in <i>Drosophila</i> Journal: Nature Communications DOI: 10.1038/s41467-021-24395-z

1. Value of the Data

- This dataset represents a thorough analysis of sex-specific compounds in 99 species within the Dipteran family Drosophilidae. It includes 42 compounds in males, 9 in females, and quantifies the presence of 7-tricosene in 35 species.
- These findings will be useful to researchers studying the evolution of sex pheromones and communication systems as well as to *Drosophila* and chemical ecology experts.
- The variation in presence and concentration of the male- and female-specific compounds, although being shared by various *Drosophila* species, advances our understanding of evolutionary mechanisms that might cause a divergence in sexual communication and reproductive isolation between closely related species.
- The comprehensiveness of this dataset will pave the road for numerous further investigations on mating systems in different *Drosophila* species and will open the door to investigate genetic and neural correlates linked to the evolution of sex pheromones.

2. Background

The diversification of sex-pheromone communication is driven by diverse factors and influenced by multiple pressures, including genetic constraints and environmental signals. Until recently, the enormous diversity of sex pheromones in *Drosophila* flies, along with their evolutionary diversification and detection, had not been comprehensively described. We recently characterized the sex pheromone communications systems for 99 species of drosophilid flies, identifying up to 43 male-specific and 9 female-specific compounds [1]. Male-specific compounds spanned various chemical classes and were often transferred to females during mating, whereas female-specific compounds were not transferred to males. Mapping these compounds onto the phylogenetic tree showed that some male-specific compounds are widely shared across distant species, while a few are species-specific. This study highlighted how species-specific olfactory signals can reinforce sexual isolation barriers between species. However, data quantifying the abundance of these compounds for each species was previously unavailable.

3. Data Description

To quantify the male- and female-specific compounds, we analyzed the chemical profiles of 99 species and compared the chromatograms of both sexes within each species. Out of 99, 81 species exhibited sexually dimorphic cuticular chemicals. Remarkably, all 81 of these species showcased the presence of male-specific compounds, which amounted to a total of 42 unique compounds. In contrast, only 15 species displayed female-specific compounds, amounting to 9 compounds in total (see Sheet 2 in supplementary Table 1) (Figs. 1 and 2).

4. Experimental Design, Materials and Methods

4.1. Fly stocks

Wild-type flies used in this study were obtained from the National *Drosophila* Species Stock Centre (NDSSC; <http://blogs.cornell.edu/drosophila/>) and Kyoto stock center (Kyoto DGGR; <https://kyotofly.kit.jp/cgi-bin/stocks/index.cgi>). All flies were reared at 25 °C, 12 h Light:12 h Dark and 50 % relative humidity. Stock numbers and breeding diets are listed in [1].

4.2. Thermal desorption–gas chromatography–mass spectrometry (TD–GC–MS)

Individual headless virgin male and female flies in different mating status were prepared for chemical profile collection as described previously [2,3], with some modifications. Briefly, the GC–MS device (Agilent GC 7890A fitted with an MS 5975C inert XL MSD unit; www.agilent.com) was equipped with an HP5-MS UI column (19091S-433UI; Agilent Technologies). After desorption at 250 °C for 3 min, the volatiles were trapped at –50 °C using liquid nitrogen for cooling. In order to transfer the components to the GC column, the vaporizer injector was heated gradually to 270 °C (12 °C/s) and held for 5 min. The temperature of the GC oven was held at 50 °C for 3 min, gradually increased (15 °C/min) to 250 °C and held for 3 min, and then to 280 °C (20 °C/min) and held for 30 min. For MS, the transfer line, source, and quad were held at 260 °C, 230 °C, and 150 °C, respectively. Eluted compounds were ionized in electron ionization (EI) source using electron beam operating at 70 eV energy and their mass spectra were recorded in positive ion mode in the range from m/z 33 to 500. All gas-chromatography data were collected and analyzed by MSD Chemstation software (F.01.03.2357).




Fig. 1. Quantification of the male-specific compound cis-vaccenyl acetate (cVA) in the 99 species. cVA was identified in 34 species, with the highest concentration observed in *D. funebris* and lowest in *D. serrata*. Each species underwent analysis with five or more replicates. Species names are ranked based on their relationships [1]. cVA is among the 42 male-specific compounds detected in the 81 dimorphic species (see Sheets 1 and 2 in supplementary Table 1).




Fig. 2. Quantification of the cuticular hydrocarbon pheromone, 7-tricosene, in 35 drosophilids. Box plots illustrate 7-tricosene abundance across five or more replicates in males (blue) and females (pink). 7-tricosene is equally present in both sexes of 14 species, more abundant in females of 7 species, and more abundant in males of 14 species (see Sheet 3 in supplementary Table 1). Notably, our findings indicate that 7-tricosene is a male-specific compound in four drosophilids: *D. virilis*, *D. americana*, *D. erecta* and *D. sucinea*. Pairwise comparisons between sexes within each species were conducted using the Mann-Whitney test. Ns $p > 0.05$; * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$.

Limitations

Not applicable.

Ethics Statement

Authors have read and follow the ethical requirements for publication in Data in Brief and confirming that the current work does not involve human subjects, or any data collected from social media platforms.

CRediT Author Statement

Mohammed A. Khallaf: Conceptualization, Methodology, Investigation, Software, Data curation, Visualization, Writing- Original draft preparation, **Melissa Diaz-Morales:** Data curation, Visualization, Writing- Reviewing and Editing, **Bill Hansson:** Conceptualization, Validation, Funding acquisition, Writing- Reviewing and Editing, **Markus Knaden:** Conceptualization, Validation, Writing- Reviewing and Editing.

Data Availability

[Quantitive variation of male and female-specific compounds in 99 drosophilid flies \(Original data\)](#) (Edmond – the Open Research Data Repository of the Max Planck Society).

Acknowledgments

We thank Ibrahim Alali for fly rearing. Wild-type flies were obtained from the San Diego Drosophila Species Stock Center (now The National Drosophila Species Stock Center, Cornell University) and KYOTO Stock Center. This research was supported through funding by the Max Planck Society.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships which have, or could be perceived to have, influenced the work reported in this article.

Supplementary Materials

Supplementary material associated with this article can be found, in the online version, at [doi:10.1016/j.dib.2024.110871](https://doi.org/10.1016/j.dib.2024.110871).

References

- [1] M.A. Khallaf, et al., Large-scale characterization of sex pheromone communication systems in Drosophila, Nat. Commun. 12 (2021) 4165, doi:[10.1038/s41467-021-24395-z](https://doi.org/10.1038/s41467-021-24395-z).
- [2] H.K.M. Dweck, et al., Pheromones mediating copulation and attraction in Drosophila, Proc. Natl. Acad. Sci. U.S.A. 112 (2015) E2829–E2835, doi:[10.1073/pnas.1504527112](https://doi.org/10.1073/pnas.1504527112).
- [3] M.A. Khallaf, et al., Mate discrimination among subspecies through a conserved olfactory pathway, Sci. Adv. 6 (2020) eaba5279, doi:[10.1126/sciadv.aba5279](https://doi.org/10.1126/sciadv.aba5279).