

Systematic integral evaluation for spin-resummed binary dynamics

Gang Chen,^{1,*} Jung-Wook Kim (김정욱),^{2,†} and Tianheng Wang^{3,‡}

¹*Niels Bohr International Academy, Niels Bohr Institute, University of Copenhagen, Blegdamsvej 17, DK-2100 Copenhagen Ø, Denmark*

²*Max Planck Institute for Gravitational Physics (Albert Einstein Institute), Am Mühlenberg 1, D-14476 Potsdam, Germany*

³*Center for Theoretical Physics, Seoul National University, 1 Gwanak-ro, Gwanak-gu, 08826, Seoul, South Korea*

Computation of spin-resummed observables in post-Minkowskian dynamics typically involve evaluation of Feynman integrals deformed by an exponential factor, where the exponent is a linear sum of the momenta being integrated. Such integrals can be viewed as tensor integral generating functions, which provide alternative approaches to tensor reduction of Feynman integrals. We develop a systematic method to evaluate tensor integral generating functions using *conventional* multiloop integration techniques. The spin-resummed aligned-spin eikonal at second post-Minkowskian order is considered as a phenomenologically relevant example where evaluation of tensor integral generating functions is necessary.

INTRODUCTION

Inspired by quantum field theoretic approaches to gravity [1–8], modern methods to compute scattering amplitude have been widely applied to the study of classical dynamics of binary black hole scattering, from conservative dynamics [9–31] and gravitational bremsstrahlung [32–45] to effects of spin [46–79] (see also refs. [80–98] for related approaches), motivated in part by the detection of gravitational waves by the LIGO-Virgo collaboration [99]. The relevant perturbative expansion is the post-Minkowskian (PM) expansion, where general relativistic corrections are added to the unperturbed special relativistic kinematics. The study of PM dynamics is more than a theoretical interest, as they can be used in building waveform models for gravitational wave detection [100].

The typical integrals encountered in the study of PM dynamics can be formulated as Feynman integrals, which can be handled efficiently by modern amplitude techniques such as the integration-by-parts (IBP) reduction [101–103] and the method of differential equations [104–106]. The function space of the master integrals and their associated graph topologies were studied for the conservative scattering dynamics at 4PM and 5PM orders [25, 28, 83, 107–110]. The function space remains practically the same when spin effects are incorporated perturbatively into the non-spinning dynamics.

However, the function space becomes completely different when *exact* spin dependence—or spin resummation—is considered. The motivation for studying spin resummation in binary Kerr dynamics is that gravitational wave physics is becoming a precision science; one potential obstruction for high-precision gravitational wave physics is inaccurate modelling of spin effects in gravitational waveform models [111], which needs to be improved with better understanding of spin effects in grav-

itational two-body dynamics.

The spin-induced multipole moments of black holes can be generated by the Newman-Janis shift [51, 112], where the position of the black hole is complexified and shifted in the imaginary spin direction. The translation operator in the imaginary direction is $\exp(i\alpha \cdot \nabla) = \exp(\alpha \cdot K)$, and the same factor appears when the Newman-Janis shift is promoted to a dynamical statement [113]. In particular, the typical Feynman integrals encountered in the context of PM dynamics are modified by exponential factors corresponding to translations along the imaginary spin direction.

Such integrals can be considered in a more general setting as *tensor integral generating functions* (TIGFs), where a typical Feynman integrand is deformed by an exponential factor of integration variables. The TIGFs provide an alternative route to tensor reduction of Feynman integrals [114]. More importantly, consideration of such integrals allows us to treat the impact parameter space Fourier transform on an equal footing with loop momenta integration, and leads us to new insights on PM dynamics presented in impact parameter space [45]. For example, the spurious poles from the intermediate loop integral reduction, and computation of the terms that contribute to the loop integrals but vanish under the impact parameter space transform, can be avoided by performing IBP reduction at the full integrand level.

Unfortunately, known multiloop integration techniques are not directly applicable to evaluation of such modified Feynman integrals. We introduce a method to tame the exponential factor and convert it into an extra delta constraint, rendering the integrand suitable for application of *conventional* multiloop integration techniques. Interestingly, the extra delta constraint leads to significantly different function space of the master integrals when compared to the original master integrals with the same topology.




FIG. 1: $G_1(x_1)G_1(x_2)$ (left) and $G_2(x_1, x_2)$ (right)

where $\ell_2 = q - \ell_1$. M_{a_2} and M_{a_1} are the three-point and four-point HEFT amplitudes provided in the appendix.

The key ingredient is the HEFT Compton amplitude [115, 116], which—contrary to the minimal coupling Compton amplitude [47]—is free of unphysical singularities. However, the amplitude has spurious singularities entering through the entire functions

$$G_1(x_1)G_1(x_2) = \frac{\sinh(x_1)}{x_1} \frac{\sinh(x_2)}{x_2}, \quad (9)$$

$$G_2(x_1, x_2) = \frac{1}{x_2} \left(\frac{\sinh(x_{12})}{x_{12}} - \cosh(x_2) \frac{\sinh(x_1)}{x_1} \right),$$

where $x_{12} = x_1 + x_2$ and $x_i = a_1 \cdot \ell_i$. As can be seen from the graph of the two functions in fig. 1, the apparent singularities of (9) do not correspond to actual singularities. Nevertheless, the spurious singularities can be problematic in the evaluation of the integrals, especially when the integrations are performed numerically as the singularities spoil numerical stability. We therefore introduce their integral representations

$$G_1(x_1)G_1(x_2) = \int_0^1 d\sigma_1 d\sigma_2 \cosh(\sigma_1 x_1) \cosh(\sigma_2 x_2),$$

$$G_2(x_1, x_2) = \int_0^1 d\sigma_1 d\sigma_2 \left(\sigma_1 \sinh(\sigma_1 x_1 + \sigma_1 \sigma_2 x_2) - \sinh(\sigma_2 x_2) \cosh(\sigma_1 x_1) \right), \quad (10)$$

which removes spurious denominators at the price of auxiliary integrations over σ_i . Analogous integral representations are used for G_1 functions appearing in the three-point amplitudes when constructing the integrand (8).

The eikonal χ is obtained by transforming the HEFT amplitude to impact parameter space

$$\chi = \int \frac{d^D q}{(2\pi)^{D-2}} \frac{e^{iq \cdot b} \delta(q \cdot v_1) \delta(q \cdot v_2)}{4m_1 m_2} \quad (11)$$

$$\times M_{\text{HEFT}}(q, v_1, v_2, a_1, a_2).$$

We focus on a slight generalisation of the aligned spin configuration of a binary Kerr system; $a^\mu \equiv a_1^\mu = \xi a_2^\mu$ and $v_1 \cdot a_2 = v_2 \cdot a_1 = 0$ with $a \cdot b \neq 0$.

Exchanging the order of integration to pull out the σ_i integrals, the eikonal can be schematically written as

$$\chi = \int_0^1 \prod_{j=1}^4 d\sigma_j \sum_{\alpha} \mathcal{I}^{(\alpha)}[\mathbf{y}], \quad (12)$$




FIG. 2: The graph topologies of master integrals, edges labelled by λ_r defined in (22). The dashed line is the delta constraint originating from the exponential factor. The thick lines are remaining delta constraints, while the thin lines are physical propagators.

where

$$y_1 = v_1 \cdot v_2, \quad y_2 = \frac{a \cdot a}{-b \cdot b}, \quad y_3 = b \cdot b, \quad y_4 = \frac{b \cdot a}{-b \cdot b} \quad (13)$$

are the Lorentz scalars the eikonal depends on. The integral representation (10) reorganises the original integrand as a sum over integrands of the form (1) with additional numerator factors, which we classify into distinct *sectors* (α) defined by the exponential factor $\exp(iq \cdot \hat{b} + i\ell_1 \cdot \hat{a})$.

$$(\alpha) \equiv (\hat{b}, \hat{a}) \quad (14)$$

$$\equiv (b + c'_1(\sigma) \tilde{a}_1 + c'_2(\sigma) \tilde{a}_2, c'_3(\sigma) \tilde{a}_1 + c'_4(\sigma) \tilde{a}_2),$$

$$(\mathbf{y}) = (y_1, \hat{y}_2, \hat{y}_3, \hat{y}_4, \sigma_1, \sigma_2, \sigma_3, \sigma_4),$$

$$(K_1, K_2) = (q, \ell_1).$$

The Lorentz scalars \hat{y}_i are defined similar to (13) with hatted variables \hat{b}, \hat{a} . The tilded spin vectors $\tilde{a}_j^\mu = ia_j^\mu$ are analytic continuation of the real spin(-length) vectors of the black holes a_j^μ . After evaluation, the master integrals can be analytically continued back to real spin.

After employing the methods developed in the previous section, a typical integrand for a given sector (α) takes the form

$$\mathcal{I}^{(\alpha)}[\mathbf{y}] = \int_{-\infty}^{\infty} dt e^{it} \int \frac{d^D q d^D \ell_1}{(2\pi)^{2D-3}} \delta(q \cdot \hat{b} + \ell_1 \cdot \hat{a} - t) \quad (15)$$

$$\times \frac{\delta(q \cdot v_1) \delta(q \cdot v_2) \delta(\ell_1 \cdot v_2)}{8m_1 m_2^2 \ell_1^2 (q - \ell_1)^2} \left(\sum_{r=0}^2 \frac{N^{(r)}(\mathbf{y})}{(\ell_1 \cdot v_1)^r} + \frac{N^{(3)}(\mathbf{y})}{q^2} \right),$$

where denominators of the integrand were kept explicit. The relevant topologies are given in fig. 2. Each sector can be reduced to master integrals by applying conventional multiloop integration techniques such as IBP reduction.

All master integrals satisfy the factorisation condition (6). Moreover, solving the differential equations in t yields power-law t -dependence of the master integrals. All terms resulting from reducing (15) have t -dependence of the form

$$\frac{t^{j-4\epsilon}}{\epsilon}, \quad j \in [0, 6]. \quad (16)$$

As remarked, the contributions from the t Fourier transform can be treated as a constant factor. It is remarked that the divergent $\frac{1}{\epsilon}$ term in dimensional regularisation vanishes under the t Fourier transform, therefore the constant factor is finite.

An exemplary term in the $N^{(3)}$ category is

$$\mathcal{I}_{ex}[\mathbf{y}] = \int \frac{d^D q d^D \ell_1}{(2\pi)^{2D-3}} e^{iq \cdot b} \frac{\delta(q \cdot v_1) \delta(q \cdot v_2) \delta(\ell_1 \cdot v_2)}{\ell_1^2 (q - \ell_1)^2 q^2} \quad (17)$$

$$\times \frac{(16\pi G_N)^2 m_1 m_2^2 (\ell_1 \cdot v_1)^2 \cosh((1 - \xi) a_1 \cdot q + 2\xi a_1 \cdot \ell_1)}{16}.$$

The cosh function splits the integrand into two sectors; sector (1) = $(\hat{a}_1, \hat{b}_1) = (2\tilde{a}_1 \xi, \tilde{a}_1(1 - \xi) + b)$ and sector (2) = $(\hat{a}_2, \hat{b}_2) = (-2\tilde{a}_1 \xi, \tilde{a}_1(\xi - 1) + b)$. Each sector reduces to three master integrals by IBP reduction,

$$\mathcal{I}_{ex}^{(\alpha)}[\mathbf{y}] = -(16\pi G_N)^2 \frac{\pi m_1 m_2^2 (y_1^2 - 1)}{2 \cdot 32 \hat{y}_2 \hat{y}_3} \quad (18)$$

$$\times \left(2\hat{y}_3 (\hat{y}_2 + 2\hat{y}_4 - 1) \widehat{\mathcal{I}}_1^{(\alpha)}[\mathbf{y}] + 2\widehat{\mathcal{I}}_2^{(\alpha)}[\mathbf{y}] + \widehat{\mathcal{I}}_3^{(\alpha)}[\mathbf{y}] \right),$$

where the master integrals are defined by (21) and the superscript (α) denotes the sectors. (17) becomes

$$\mathcal{I}_{ex}[\mathbf{y}] = \mathcal{I}_{ex}^{(1)}[\mathbf{y}] + \mathcal{I}_{ex}^{(2)}[\mathbf{y}], \quad (19)$$

where each $\mathcal{I}_{ex}^{(\alpha)}[\mathbf{y}]$ is complex but the sum $\mathcal{I}_{ex}[\mathbf{y}]$ is real.

The integrals in (12) reduce to 320 sectors. Each sector has five master integrals. However, only three master integrals per sector appear in the final result,

$$\chi = \int_0^1 \prod_{j=1}^4 d\sigma_j \sum_{\alpha=1}^{320} \left(c_{1,\alpha}(\mathbf{y}) \widehat{\mathcal{I}}_1^{(\alpha)}[\mathbf{y}] + c_{2,\alpha}(\mathbf{y}) \widehat{\mathcal{I}}_2^{(\alpha)}[\mathbf{y}] \right. \\ \left. + c_{3,\alpha}(\mathbf{y}) \widehat{\mathcal{I}}_3^{(\alpha)}[\mathbf{y}] \right), \quad (20)$$

where the master integrals are

$$\widehat{\mathcal{I}}_1^{(\alpha)}[\mathbf{y}] = \frac{D-4}{t^{2D-8}} J_{1,1,1,1,1,1,0,0}^{(\alpha)}, \\ \widehat{\mathcal{I}}_2^{(\alpha)}[\mathbf{y}] = \frac{1}{t^{2D-10}} J_{2,1,1,1,1,1,0,0}^{(\alpha)}, \\ \widehat{\mathcal{I}}_3^{(\alpha)}[\mathbf{y}] = \frac{1}{t^{2D-10}} J_{1,1,1,1,1,1,0,1}^{(\alpha)}, \quad (21)$$

defined as

$$J_{\lambda_1, \lambda_2, \lambda_3, \lambda_4, \lambda_5, \lambda_6, \lambda_7, \lambda_8}^{(\alpha)} \equiv \int \frac{d^D q d^D \ell_1}{(2\pi)^{2D-3}} \delta^{\lambda_6-1}(\hat{a} \cdot \ell_1 + \hat{b} \cdot q - t) \\ \times \frac{\delta^{\lambda_3-1}(\ell_1 \cdot v_2) \delta^{\lambda_4-1}(q \cdot v_1) \delta^{\lambda_5-1}(q \cdot v_2)}{(\ell_1^2)^{\lambda_1} ((q - \ell_1)^2)^{\lambda_2} (\ell_1 \cdot v_1)^{\lambda_7} (q^2)^{\lambda_8}}. \quad (22)$$

The numerators $N^{(0,1,2)}$ of (12) reduce to $\widehat{\mathcal{I}}_{1,2}^{(\alpha)}[\mathbf{y}]$, while the numerators $N^{(3)}$ of (12) reduce to $\widehat{\mathcal{I}}_{1,2,3}^{(\alpha)}[\mathbf{y}]$. The integrals $\widehat{\mathcal{I}}_{1,2}^{(\alpha)}[\mathbf{y}]$ correspond to the left topology of fig. 2,

while the integral $\widehat{\mathcal{I}}_3^{(\alpha)}[\mathbf{y}]$ corresponds to the right topology of fig. 2. The apparent t -dependence in (21) makes the master integrals $\widehat{\mathcal{I}}_{1,2,3}^{(\alpha)}[\mathbf{y}]$ independent of t . As mentioned previously, the other two master integrals, $J_{1,1,1,2,1,1,0,0}^{(\alpha)}$ and $J_{1,1,1,1,1,1,1,0}^{(\alpha)}$, never appear and their evaluation is unnecessary for evaluating the eikonal. It is also remarked that both $c_{j,\alpha}(\mathbf{y})$ and $\widehat{\mathcal{I}}_j^{(\alpha)}[\mathbf{y}]$ are finite in dimensional regularisation.

The final result for the eikonal phase is available in the public repository [KerrEikonal2pm](#) [125]. We emphasise that the result contains all-orders-in-spin contributions for the generalised aligned spin configuration with $a \cdot b \neq 0$.

EVALUATION OF THE MASTER INTEGRALS

We evaluate the master integrals using the method of differential equations [104–106]. The \hat{y}_3 -dependence of the master integrals can be determined from dimensional analysis, since the only dimensionful variable is \hat{y}_3 . Moreover, the differential operator ∂_{y_1} is diagonal in the system of master integrals, and the y_1 -dependence of the master integrals can be solved separately.

The nontrivial systems of differential equations are given by $\hat{y}_{2,4}$. The two systems are degenerate for the master integrals $\widehat{\mathcal{I}}_{1,2}$ and they only depend on the combination $\hat{y}'_2 = \hat{y}_2 + 2\hat{y}_4$. The differential equations over \hat{y}'_2 for $\widehat{\mathcal{I}}_{1,2}$ are solved by complete elliptic integrals [126],

$$\widehat{\mathcal{I}}_1[\mathbf{y}] = \frac{C}{\sqrt{-\hat{y}_3} \sqrt{y_1^2 - 1}} \frac{K(\hat{y}'_2)}{\pi}, \\ \widehat{\mathcal{I}}_2[\mathbf{y}] = -\frac{C \sqrt{-\hat{y}_3}}{\sqrt{y_1^2 - 1}} \frac{(\hat{y}'_2 - 1)K(\hat{y}'_2) + E(\hat{y}'_2)}{\pi}, \quad (23)$$

where the integration constant $C = -\frac{1}{16\pi^2}$ is fixed from the spinless limit $\hat{y}'_2 \rightarrow 0$.

The differential equations satisfied by $\widehat{\mathcal{I}}_3$ are

$$\partial_{\hat{y}'_2} \widehat{\mathcal{I}}_3[\mathbf{y}] = \frac{-1}{2(\hat{y}'_2 - 2\hat{y}_4)(\hat{y}'_2 + \hat{y}_4^2 - 2\hat{y}_4)} \times \\ \left(\hat{y}_4^2 \widehat{\mathcal{I}}_3[\mathbf{y}] - \frac{C(\sqrt{-\hat{y}_3})}{\pi \sqrt{y_1^2 - 1}} (\hat{y}'_2 + 2\hat{y}_4)(\hat{y}_4 - 1) E(\hat{y}'_2) \right. \\ \left. + \frac{C(\sqrt{-\hat{y}_3})}{\pi \sqrt{y_1^2 - 1}} (1 - \hat{y}_4)(\hat{y}'_2 - 2\hat{y}_4) K(\hat{y}'_2) \right), \quad (24)$$

$$\partial_{\hat{y}_4} \widehat{\mathcal{I}}_3[\mathbf{y}] = \frac{1}{(\hat{y}'_2 - 2\hat{y}_4)(\hat{y}'_2 + \hat{y}_4^2 - 2\hat{y}_4)} \left(\hat{y}_4(\hat{y}'_2 - \hat{y}_4) \widehat{\mathcal{I}}_3[\mathbf{y}] \right. \\ \left. - C \frac{(\sqrt{-\hat{y}_3})}{\sqrt{y_1^2 - 1}} (\hat{y}'_2 - 1)(\hat{y}'_2 - 2\hat{y}_4) K(\hat{y}'_2) \right. \\ \left. - C \frac{(\sqrt{-\hat{y}_3})}{\sqrt{y_1^2 - 1}} (\hat{y}_4 \hat{y}'_2 + \hat{y}'_2 - 2\hat{y}_4) E(\hat{y}'_2) \right). \quad (25)$$

The common denominator factor in the two equations implies that the master integral should be understood as

an integral over the elliptic curve

$$\mathcal{Y}^2 = (\hat{y}'_2 - 2z)(z^2 - 2z + \hat{y}'_2), \quad (26)$$

where z is the integration variable to be identified as $-\hat{y}_4$ after integration. $\widehat{\mathcal{I}}_3[\mathbf{y}]$ is computed by solving (24) at $\hat{y}_4 = 0$, which yields the boundary value for the remaining integration over \hat{y}_4 . Carrying out the \hat{y}_4 integration, we have

$$\begin{aligned} \widehat{\mathcal{I}}_3[\mathbf{y}] = & \frac{C(\sqrt{-\hat{y}_3})}{\sqrt{y_1^2 - 1}} \frac{\sqrt{\hat{y}'_2 + \hat{y}_4(\hat{y}_4 - 2)}}{\sqrt{\hat{y}'_2 - 2\hat{y}_4}} \left[\frac{2E(\hat{y}'_2) + \pi}{2\pi} \right. \\ & + \int_0^{\hat{y}_4} dz \left(\frac{(1 - \hat{y}'_2)(\hat{y}'_2 - 2z)K(\hat{y}'_2)}{\pi \mathcal{Y}(z^2 - 2z + \hat{y}'_2)} \right. \\ & \left. \left. - \frac{(\hat{y}'_2 z - 2z + \hat{y}'_2)E(\hat{y}'_2)}{\pi \mathcal{Y}(z^2 - 2z + \hat{y}'_2)} \right) \right]. \end{aligned} \quad (27)$$

The z -integral can be reduced by the IBP relations following from the exact differentials

$$d\left(\frac{\hat{y}'_2 - 2z}{\mathcal{Y}}\right), \quad d\left(\frac{(\hat{y}'_2 - 2z)^2}{\mathcal{Y}}\right). \quad (28)$$

Eliminating integrals with undesired denominators, we arrive at

$$\begin{aligned} \widehat{\mathcal{I}}_3[\mathbf{y}] = & \frac{C\sqrt{-\hat{y}_3}}{\sqrt{y_1^2 - 1}} \left[\frac{\sqrt{\hat{y}'_2 + \hat{y}_4^2 - 2\hat{y}_4}}{\pi \sqrt{\hat{y}'_2 - 2\hat{y}_4}} \left(\frac{\pi - 2K(\hat{y}'_2)}{2} \right. \right. \\ & - (E(\hat{y}'_2) - K(\hat{y}'_2)) \int_0^{\hat{y}_4} \frac{dz}{\mathcal{Y}} - K(\hat{y}'_2) \int_0^{\hat{y}_4} \frac{z dz}{\mathcal{Y}} \left. \right) \\ & \left. + \frac{E(\hat{y}'_2) + (1 - \hat{y}_4)K(\hat{y}'_2)}{\pi} \right], \end{aligned} \quad (29)$$

where the integrals can be converted to incomplete elliptic integrals, which can be evaluated to arbitrary numerical precision. The integration constants were fixed by requiring regularity of the spin expansion around $\hat{a}^\mu = 0$. Since $\hat{a}^\mu \rightarrow 0$ is a singular limit of the underlying curve (26), we expand (29) around \hat{y}_4 first and then expand in \hat{y}_2 to obtain the series expansion of $\widehat{\mathcal{I}}_3$. The master integrals $\widehat{\mathcal{I}}_{1,2,3}$ were checked against brute-force calculations based on the formulas in appendix B of ref. [113], and found to agree up to $\mathcal{O}(\hat{a}^{20})$.

RESULTS

The eikonal phase (20) is exact to all orders in spin, and can be expanded in spin before the σ_j -integration for a check against perturbative-in-spin calculations. The spin-expanded σ_j integrand turns out to be polynomial in σ_j , and the auxiliary σ_j integrals can be evaluated exactly. We provide files for evaluating the spin-expanded σ_j integrand in the public repository




FIG. 3: The eikonal (20) as a function of $\frac{|a|}{|b|}$ and $\cos(\theta) = \frac{-a \cdot b}{|a||b|}$ with numerical errors $\sim 10^{-15}$.

[KerrEikonal2pm](#) [125]. We have checked that (20) expanded to $\mathcal{O}(a^8)$ is consistent with the results of ref. [124].

The merit of the eikonal phase (20) is that we can study its full spin dependence quantitatively, since the σ_j integrations can be numerically evaluated with high precision. As an example of numerically studying the eikonal (20), we consider the scattering of a test Kerr black hole on a Schwarzschild background, corresponding to the substitution

$$\chi_{a_2=0} = \chi|_{\xi \rightarrow 0}, \quad (30)$$

which is presented in fig. 3. The parameters $y_1 = \frac{12}{10}$ and $y_3 = -1$ are chosen for the ranges $|\sqrt{-y_2}| < 0.75$ and $\cos(\theta) \in [0, 0.9]$, where θ denotes the angle between a^μ and b^μ such that $\sqrt{-y_2} \cos(\theta) = y_4$.

The numerical calculations can be tested against analytic predictions. For example, the 2PM aligned-spin ($\cos(\theta) = 0$) eikonal is expected to have the singularity $\propto (|b|^2 - |a|^2)^{-3/2}$ based on analytic studies [71, 96, 113]. Fig. 4 shows numerical evaluation of (20) in the aligned-spin configuration plotted against the reference curve $\frac{\beta}{(|b|^2 - |a|^2)^{3/2}}$, where the constant β was fit to (20) at $|b| = 1.02|a|$ and other unspecified parameters are the same as in fig. 3. The two lines on fig. 4 show a good agreement near the singular point $|b| = |a|$, having $\lesssim 5\%$ relative difference for $|b| \leq 1.04|a|$. This singularity can be interpreted as the dynamics detecting the ring singularity from spin resummation, since the eikonal has a singularity only at $|b| = 0$ when spin is treated perturbatively.

CONCLUSIONS AND OUTLOOK

In this letter, a systematic method to reduce and evaluate TIGFs was introduced and applied to the scattering dynamics of a binary Kerr system in the generalised aligned-spin configuration. The corresponding




FIG. 4: Aligned-spin χ near the singular point $|b| = |a|$. The reference $\frac{\beta}{(|b|^2 - |a|^2)^{3/2}}$ intersects at $|b| = 1.02|a|$.

2PM eikonal phase (20) was presented in a closed form, which can be studied analytically by expanding to arbitrary orders in spin, or can be studied numerically through numerical integrations for exact spin dependence.

An important application of TIGFs is generation of tensor integrals, directly providing an alternative method of performing tensor reduction in Feynman integrals [114]. The approach may prove useful for reduction of irreducible numerators, which in many cases becomes a bottleneck when evaluation of multiloop Feynman integrals are involved. In this regard, understanding the criteria for the factorisation of the t -dependence (e.g. from intersection theory [45, 127–129]) will be important, as the factorisation played a crucial role in obtaining closed-form expressions for the TIGFs considered.

Another future direction to explore would be the study of function space complexity for TIGFs. We have seen that elliptic integrals appear in effective two-loop (2PM) topologies for TIGFs (see also ref. [113] for electromagnetism). It is reasonable to expect that more complex functions (e.g. elliptic multi-polylogarithms [130], integrals over Calabi-Yau manifolds [131]) will appear in effective three-loop (3PM) graph topologies when integrals are deformed into TIGFs. Moreover, we can attempt to quantify the increase in transcendentality of the function space when typical Feynman integrals are deformed into TIGFs. This may even lead to an exploration of new geometries that have not yet been associated with Feynman integrals.

Special cases of TIGFs appear ubiquitously in quantum-field-theory-inspired approaches to classical gravitating systems, especially in the calculation of scattering waveforms [33–45]. It would also be interesting to apply the developed methods to these problems.

Coming back to the initial motivation for the study of

TIGFs, it would be interesting to apply the developed methods to 3PM all-orders-in-spin dynamics and study spin resummation. Whether we have the exact three-graviton-Kerr five-point amplitude—which is necessary for constructing the 3PM integrand—is less relevant, as long as the conjectured five-point amplitude correctly captures features of the dynamical Newman-Janis shift; we expect the singularity structures of the binary Kerr dynamics to be governed by the dynamical Newman-Janis shift, and correct singularity structures are the most important when we attempt to resum the perturbative two-body dynamics [113]. Compared to the 2PM dynamics studied in this letter, the spin-resummed 3PM dynamics is qualitatively different in that it is the first order where next-to-leading-order effects in the mass-ratio expansion enters into the dynamics [93], thereby including the first beyond background-probe limit effects. The insights gained from studying singularity structures of spin-resummed binary Kerr dynamics may motivate new resummation schemes for spinning binary dynamics, providing a more accurate incorporation of spin effects in waveform models used by gravitational wave observatories, potentially having far-reaching consequences for astrophysics and multi-messenger astronomy.

ACKNOWLEDGEMENTS

It is a pleasure to thank Emil Bjerrum-Bohr, Andreas Brandhuber, Graham Brown, Marcos Skowronek, Zhengwen Liu, Roger Morales, Gabriele Travaglini for interesting conversations. JWK would like to thank Stefano De Angelis and Fei Teng for stimulating discussions. The authors would like to thank Bo Feng, Andres Luna, and Roger Morales for comments on the draft. GC has received funding from the European Union’s Horizon 2020 research and innovation program under the Marie Skłodowska-Curie grant agreement No. 847523 “INTERACTIONS”. TW is supported by the NRF grant 2021R1A2C2012350.

* gang.chen@nbi.ku.dk

† jung-wook.kim@aei.mpg.de

‡ tianhengwang@snu.ac.kr

- [1] Y. Iwasaki, Prog. Theor. Phys. **46**, 1587 (1971).
- [2] J. F. Donoghue, Phys. Rev. D **50**, 3874 (1994), arXiv:gr-qc/9405057.
- [3] N. E. J. Bjerrum-Bohr, J. F. Donoghue, and B. R. Holstein, Phys. Rev. D **67**, 084033 (2003), [Erratum: Phys.Rev.D 71, 069903 (2005)], arXiv:hep-th/0211072.
- [4] B. R. Holstein and J. F. Donoghue, Phys. Rev. Lett. **93**, 201602 (2004), arXiv:hep-th/0405239 [hep-th].
- [5] B. R. Holstein and A. Ross, (2008), arXiv:0802.0716 [hep-ph].

- [6] D. Neill and I. Z. Rothstein, Nucl. Phys. **B877**, 177 (2013), arXiv:1304.7263 [hep-th].
- [7] N. E. J. Bjerrum-Bohr, J. F. Donoghue, and P. Vanhove, JHEP **02**, 111 (2014), arXiv:1309.0804 [hep-th].
- [8] T. Damour, Phys. Rev. **D97**, 044038 (2018), arXiv:1710.10599 [gr-qc].
- [9] N. E. J. Bjerrum-Bohr, P. H. Damgaard, G. Festuccia, L. Plante, and P. Vanhove, Phys. Rev. Lett. **121**, 171601 (2018), arXiv:1806.04920 [hep-th].
- [10] C. Cheung, I. Z. Rothstein, and M. P. Solon, Phys. Rev. Lett. **121**, 251101 (2018), arXiv:1808.02489 [hep-th].
- [11] A. Cristofoli, N. E. J. Bjerrum-Bohr, P. H. Damgaard, and P. Vanhove, Phys. Rev. D **100**, 084040 (2019), arXiv:1906.01579 [hep-th].
- [12] Z. Bern, C. Cheung, R. Roiban, C.-H. Shen, M. P. Solon, and M. Zeng, Phys. Rev. Lett. **122**, 201603 (2019), arXiv:1901.04424 [hep-th].
- [13] Z. Bern, C. Cheung, R. Roiban, C.-H. Shen, M. P. Solon, and M. Zeng, JHEP **10**, 206 (2019), arXiv:1908.01493 [hep-th].
- [14] J. Parra-Martinez, M. S. Ruf, and M. Zeng, JHEP **11**, 023 (2020), arXiv:2005.04236 [hep-th].
- [15] P. Di Vecchia, C. Heissenberg, R. Russo, and G. Veneziano, Phys. Lett. B **811**, 135924 (2020), arXiv:2008.12743 [hep-th].
- [16] P. Di Vecchia, C. Heissenberg, R. Russo, and G. Veneziano, Phys. Lett. B **818**, 136379 (2021), arXiv:2101.05772 [hep-th].
- [17] P. Di Vecchia, C. Heissenberg, R. Russo, and G. Veneziano, JHEP **07**, 169 (2021), arXiv:2104.03256 [hep-th].
- [18] E. Herrmann, J. Parra-Martinez, M. S. Ruf, and M. Zeng, JHEP **10**, 148 (2021), arXiv:2104.03957 [hep-th].
- [19] N. E. J. Bjerrum-Bohr, P. H. Damgaard, L. Planté, and P. Vanhove, Phys. Rev. D **104**, 026009 (2021), arXiv:2104.04510 [hep-th].
- [20] N. E. J. Bjerrum-Bohr, P. H. Damgaard, L. Planté, and P. Vanhove, (2021), arXiv:2105.05218 [hep-th].
- [21] P. H. Damgaard, L. Plante, and P. Vanhove, JHEP **11**, 213 (2021), arXiv:2107.12891 [hep-th].
- [22] A. Brandhuber, G. Chen, G. Travaglini, and C. Wen, JHEP **10**, 118 (2021), arXiv:2108.04216 [hep-th].
- [23] N. E. J. Bjerrum-Bohr, L. Planté, and P. Vanhove, JHEP **03**, 071 (2022), arXiv:2111.02976 [hep-th].
- [24] N. E. J. Bjerrum-Bohr, L. Planté, and P. Vanhove, (2022), arXiv:2212.08957 [hep-th].
- [25] Z. Bern, J. Parra-Martinez, R. Roiban, M. S. Ruf, C.-H. Shen, M. P. Solon, and M. Zeng, Phys. Rev. Lett. **128**, 161103 (2022), arXiv:2112.10750 [hep-th].
- [26] Z. Bern, J. Parra-Martinez, R. Roiban, M. S. Ruf, C.-H. Shen, M. P. Solon, and M. Zeng, PoS **LL2022**, 051 (2022).
- [27] Z. Bern, J. Parra-Martinez, R. Roiban, M. S. Ruf, C.-H. Shen, M. P. Solon, and M. Zeng, Phys. Rev. Lett. **126**, 171601 (2021), arXiv:2101.07254 [hep-th].
- [28] P. H. Damgaard, E. R. Hansen, L. Planté, and P. Vanhove, (2023), arXiv:2307.04746 [hep-th].
- [29] P. Di Vecchia, C. Heissenberg, R. Russo, and G. Veneziano, (2023), arXiv:2306.16488 [hep-th].
- [30] C. Heissenberg, (2023), arXiv:2308.11470 [hep-th].
- [31] T. Adamo and R. Gonzo, JHEP **05**, 088 (2023), arXiv:2212.13269 [hep-th].
- [32] D. A. Kosower, B. Maybee, and D. O’Connell, JHEP **02**, 137 (2019), arXiv:1811.10950 [hep-th].
- [33] A. Brandhuber, G. R. Brown, G. Chen, S. De Angelis, J. Gowdy, and G. Travaglini, JHEP **06**, 048 (2023), arXiv:2303.06111 [hep-th].
- [34] A. Herderschee, R. Roiban, and F. Teng, JHEP **06**, 004 (2023), arXiv:2303.06112 [hep-th].
- [35] A. Elkhidir, D. O’Connell, M. Sergola, and I. A. Vazquez-Holm, (2023), arXiv:2303.06211 [hep-th].
- [36] A. Georgoudis, C. Heissenberg, and I. Vazquez-Holm, JHEP **06**, 126 (2023), arXiv:2303.07006 [hep-th].
- [37] S. Caron-Huot, M. Giroux, H. S. Hannesdottir, and S. Mizera, (2023), arXiv:2308.02125 [hep-th].
- [38] L. Bohnenblust, H. Ita, M. Kraus, and J. Schlenk, (2023), arXiv:2312.14859 [hep-th].
- [39] D. Bini, T. Damour, and A. Geralico, Phys. Rev. D **108**, 124052 (2023), arXiv:2309.14925 [gr-qc].
- [40] A. Georgoudis, C. Heissenberg, and R. Russo, JHEP **03**, 089 (2024), arXiv:2312.07452 [hep-th].
- [41] T. Adamo, R. Gonzo, and A. Ilderton, JHEP **05**, 034 (2024), arXiv:2402.00124 [hep-th].
- [42] D. Bini, T. Damour, S. De Angelis, A. Geralico, A. Herderschee, R. Roiban, and F. Teng, (2024), arXiv:2402.06604 [hep-th].
- [43] F. Alessio and P. Di Vecchia, (2024), arXiv:2402.06533 [hep-th].
- [44] A. Georgoudis, C. Heissenberg, and R. Russo, Phys. Rev. D **109**, 106020 (2024), arXiv:2402.06361 [hep-th].
- [45] G. Brunello and S. De Angelis, (2024), arXiv:2403.08009 [hep-th].
- [46] A. Guevara, JHEP **04**, 033 (2019), arXiv:1706.02314 [hep-th].
- [47] N. Arkani-Hamed, T.-C. Huang, and Y.-t. Huang, JHEP **11**, 070 (2021), arXiv:1709.04891 [hep-th].
- [48] A. Guevara, A. Ochirov, and J. Vines, JHEP **09**, 056 (2019), arXiv:1812.06895 [hep-th].
- [49] M.-Z. Chung, Y.-T. Huang, J.-W. Kim, and S. Lee, JHEP **04**, 156 (2019), arXiv:1812.08752 [hep-th].
- [50] A. Guevara, A. Ochirov, and J. Vines, Phys. Rev. **D100**, 104024 (2019), arXiv:1906.10071 [hep-th].
- [51] N. Arkani-Hamed, Y.-t. Huang, and D. O’Connell, JHEP **01**, 046 (2020), arXiv:1906.10100 [hep-th].
- [52] R. Aoude, K. Haddad, and A. Helset, JHEP **05**, 051 (2020), arXiv:2001.09164 [hep-th].
- [53] M.-Z. Chung, Y.-t. Huang, J.-W. Kim, and S. Lee, JHEP **05**, 105 (2020), arXiv:2003.06600 [hep-th].
- [54] A. Guevara, B. Maybee, A. Ochirov, D. O’Connell, and J. Vines, JHEP **03**, 201 (2021), arXiv:2012.11570 [hep-th].
- [55] W.-M. Chen, M.-Z. Chung, Y.-t. Huang, and J.-W. Kim, JHEP **08**, 148 (2022), arXiv:2111.13639 [hep-th].
- [56] D. Kosmopoulos and A. Luna, JHEP **07**, 037 (2021), arXiv:2102.10137 [hep-th].
- [57] M. Chiodaroli, H. Johansson, and P. Pichini, JHEP **02**, 156 (2022), arXiv:2107.14779 [hep-th].
- [58] Y. F. Bautista, A. Guevara, C. Kavanagh, and J. Vines, (2021), arXiv:2107.10179 [hep-th].
- [59] L. Cangemi, M. Chiodaroli, H. Johansson, A. Ochirov, P. Pichini, and E. Skvortsov, (2022), arXiv:2212.06120 [hep-th].
- [60] A. Ochirov and E. Skvortsov, Phys. Rev. Lett. **129**, 241601 (2022), arXiv:2207.14597 [hep-th].
- [61] P. H. Damgaard, K. Haddad, and A. Helset, JHEP **11**, 070 (2019), arXiv:1908.10308 [hep-ph].

- [62] Z. Bern, A. Luna, R. Roiban, C.-H. Shen, and M. Zeng, *Phys. Rev. D* **104**, 065014 (2021), arXiv:2005.03071 [hep-th].
- [63] F. Comberiati and L. de la Cruz, (2022), arXiv:2212.09259 [hep-th].
- [64] B. Maybee, D. O’Connell, and J. Vines, *JHEP* **12**, 156 (2019), arXiv:1906.09260 [hep-th].
- [65] K. Haddad, *Phys. Rev. D* **105**, 026004 (2022), arXiv:2109.04427 [hep-th].
- [66] W.-M. Chen, M.-Z. Chung, Y.-t. Huang, and J.-W. Kim, *JHEP* **12**, 058 (2022), arXiv:2205.07305 [hep-th].
- [67] G. Menezes and M. Sergola, *JHEP* **10**, 105 (2022), arXiv:2205.11701 [hep-th].
- [68] F. Febres Cordero, M. Kraus, G. Lin, M. S. Ruf, and M. Zeng, *Phys. Rev. Lett.* **130**, 021601 (2023), arXiv:2205.07357 [hep-th].
- [69] F. Alessio and P. Di Vecchia, *Phys. Lett. B* **832**, 137258 (2022), arXiv:2203.13272 [hep-th].
- [70] Z. Bern, D. Kosmopoulos, A. Luna, R. Roiban, and F. Teng, (2022), arXiv:2203.06202 [hep-th].
- [71] R. Aoude, K. Haddad, and A. Helset, *Phys. Rev. Lett.* **129**, 141102 (2022), arXiv:2205.02809 [hep-th].
- [72] R. Aoude, K. Haddad, and A. Helset, *JHEP* **07**, 072 (2022), arXiv:2203.06197 [hep-th].
- [73] R. Aoude, K. Haddad, and A. Helset, *Phys. Rev. D* **108**, 024050 (2023), arXiv:2304.13740 [hep-th].
- [74] Y. F. Bautista, (2023), arXiv:2304.04287 [hep-th].
- [75] S. De Angelis, R. Gonzo, and P. P. Novichkov, (2023), arXiv:2309.17429 [hep-th].
- [76] A. Brandhuber, G. R. Brown, G. Chen, J. Gowdy, and G. Travaglini, *JHEP* **02**, 026 (2024), arXiv:2310.04405 [hep-th].
- [77] R. Aoude, K. Haddad, C. Heissenberg, and A. Helset, *Phys. Rev. D* **109**, 036007 (2024), arXiv:2310.05832 [hep-th].
- [78] M. M. Riva, F. Vernizzi, and L. K. Wong, *Phys. Rev. D* **106**, 044013 (2022), arXiv:2205.15295 [hep-th].
- [79] A. Luna, N. Moynihan, D. O’Connell, and A. Ross, (2023), arXiv:2312.09960 [hep-th].
- [80] A. Antonelli, A. Buonanno, J. Steinhoff, M. van de Meent, and J. Vines, *Phys. Rev. D* **99**, 104004 (2019), arXiv:1901.07102 [gr-qc].
- [81] T. Damour, *Phys. Rev. D* **102**, 124008 (2020), arXiv:2010.01641 [gr-qc].
- [82] G. Kälin, Z. Liu, and R. A. Porto, *Phys. Rev. Lett.* **125**, 261103 (2020), arXiv:2007.04977 [hep-th].
- [83] C. Dlapa, G. Kälin, Z. Liu, and R. A. Porto, *Phys. Lett. B* **831**, 137203 (2022), arXiv:2106.08276 [hep-th].
- [84] C. Dlapa, G. Kälin, Z. Liu, and R. A. Porto, *Phys. Rev. Lett.* **128**, 161104 (2022), arXiv:2112.11296 [hep-th].
- [85] C. Dlapa, G. Kälin, Z. Liu, J. Neef, and R. A. Porto, (2022), arXiv:2210.05541 [hep-th].
- [86] G. U. Jakobsen, G. Mogull, J. Plefka, B. Sauer, and Y. Xu, (2023), arXiv:2306.01714 [hep-th].
- [87] G. U. Jakobsen, G. Mogull, J. Plefka, and J. Steinhoff, *Phys. Rev. Lett.* **126**, 201103 (2021), arXiv:2101.12688 [gr-qc].
- [88] S. Mougiakakos, M. M. Riva, and F. Vernizzi, *Phys. Rev. D* **104**, 024041 (2021), arXiv:2102.08339 [gr-qc].
- [89] G. U. Jakobsen, G. Mogull, J. Plefka, and J. Steinhoff, (2021), arXiv:2106.10256 [hep-th].
- [90] G. Mogull, J. Plefka, and J. Steinhoff, *JHEP* **02**, 048 (2021), arXiv:2010.02865 [hep-th].
- [91] G. U. Jakobsen, G. Mogull, J. Plefka, and J. Steinhoff, *JHEP* **01**, 027 (2022), arXiv:2109.04465 [hep-th].
- [92] J. Vines, *Class. Quant. Grav.* **35**, 084002 (2018), arXiv:1709.06016 [gr-qc].
- [93] J. Vines, J. Steinhoff, and A. Buonanno, *Phys. Rev. D* **99**, 064054 (2019), arXiv:1812.00956 [gr-qc].
- [94] Z. Liu, R. A. Porto, and Z. Yang, *JHEP* **06**, 012 (2021), arXiv:2102.10059 [hep-th].
- [95] G. U. Jakobsen and G. Mogull, *Phys. Rev. Lett.* **128**, 141102 (2022), arXiv:2201.07778 [hep-th].
- [96] P. H. Damgaard, J. Hoogeveen, A. Luna, and J. Vines, *Phys. Rev. D* **106**, 124030 (2022), arXiv:2208.11028 [hep-th].
- [97] M. Bianchi, C. Gambino, and F. Riccioli, *JHEP* **08**, 188 (2023), arXiv:2306.08969 [hep-th].
- [98] R. Gonzo and C. Shi, (2024), arXiv:2405.09687 [hep-th].
- [99] B. P. Abbott *et al.* (LIGO Scientific, Virgo), *Phys. Rev. Lett.* **116**, 061102 (2016), arXiv:1602.03837 [gr-qc].
- [100] A. Buonanno, G. Mogull, R. Patil, and L. Pompili, (2024), arXiv:2405.19181 [gr-qc].
- [101] A. Smirnov, *JHEP* **10**, 107 (2008), arXiv:0807.3243 [hep-ph].
- [102] R. N. Lee, *JHEP* **04**, 108 (2015), arXiv:1411.0911 [hep-ph].
- [103] K. J. Larsen and Y. Zhang, *Phys. Rev. D* **93**, 041701 (2016), arXiv:1511.01071 [hep-th].
- [104] T. Gehrmann and E. Remiddi, *Nuclear Physics B* **580**, 485 (2000).
- [105] A. V. Kotikov, *Phys. Lett. B* **254**, 158 (1991).
- [106] J. M. Henn, *Phys. Rev. Lett.* **110**, 251601 (2013), arXiv:1304.1806 [hep-th].
- [107] H. Frellesvig, R. Morales, and M. Wilhelm, *Phys. Rev. Lett.* **132**, 201602 (2024), arXiv:2312.11371 [hep-th].
- [108] M. Driesse, G. U. Jakobsen, G. Mogull, J. Plefka, B. Sauer, and J. Usovitsch, (2024), arXiv:2403.07781 [hep-th].
- [109] A. Klemm, C. Nega, B. Sauer, and J. Plefka, (2024), arXiv:2401.07899 [hep-th].
- [110] H. Frellesvig, R. Morales, and M. Wilhelm, (2024), arXiv:2405.17255 [hep-th].
- [111] A. Dhani, S. Völkel, A. Buonanno, H. Estelles, J. Gair, H. P. Pfeiffer, L. Pompili, and A. Toubiana, (2024), arXiv:2404.05811 [gr-qc].
- [112] E. T. Newman and A. I. Janis, *J. Math. Phys.* **6**, 915 (1965).
- [113] J.-H. Kim, J.-W. Kim, and S. Lee, (2024), arXiv:2405.17056 [hep-th].
- [114] B. Feng, *Commun. Theor. Phys.* **75**, 025203 (2023), arXiv:2209.09517 [hep-ph].
- [115] N. E. J. Bjerrum-Bohr, G. Chen, and M. Skowronek, *Phys. Rev. Lett.* **132**, 191603 (2024), arXiv:2309.11249 [hep-th].
- [116] N. E. J. Bjerrum-Bohr, G. Chen, and M. Skowronek, *JHEP* **06**, 170 (2023), arXiv:2302.00498 [hep-th].
- [117] Z. Bern, L. J. Dixon, D. C. Dunbar, and D. A. Kosower, *Nucl. Phys.* **B425**, 217 (1994), arXiv:hep-ph/9403226 [hep-ph].
- [118] Z. Bern, L. J. Dixon, D. C. Dunbar, and D. A. Kosower, *Nucl. Phys.* **B435**, 59 (1995), arXiv:hep-ph/9409265 [hep-ph].
- [119] Although generating functions are usually defined by real exponents, imaginary exponents were used to avoid factors of i when solving for the integrals. Such an ana-

lytic continuation is known in the probability literature as characteristic functions. This analytic continuation is allowed since Feynman integrals should be understood as generalised functions which may not be well-defined by conventional definitions of integrals.

- [120] The delta constraints can be converted to denominators through the distributional identity $2\pi i \delta(x) = (x - i0^+)^{-1} - (x + i0^+)^{-1}$ and its derivatives, therefore they can be considered as propagator factors.
- [121] B. Feng, C. Hu, J. Shen, and Y. Zhang, (2024), arXiv:2403.16040 [hep-ph].
- [122] T. Li, Y. Song, and L. Zhang, (2024), arXiv:2404.04644 [hep-ph].
- [123] G. Brunello, G. Crisanti, M. Giroux, P. Mastrolia, and S. Smith, Phys. Rev. D **109**, 094047 (2024), arXiv:2311.14432 [hep-th].
- [124] G. Chen and T. Wang, (2024), arXiv:2406.09086 [hep-th].
- [125] G. Chen, J.-W. Kim, and T. Wang, “<https://github.com/amplitudegravity/kerreikon2pm>,” (2024).
- [126] We adopt *Mathematica*’s definition for the elliptic integrals.
- [127] K. Cho and K. Matsumoto, Nagoya Mathematical Journal **139**, 67 (1995).
- [128] P. Mastrolia and S. Mizera, JHEP **02**, 139 (2019), arXiv:1810.03818 [hep-th].
- [129] H. Frellesvig, F. Gasparotto, S. Laporta, M. K. Mandal, P. Mastrolia, L. Mattiazzi, and S. Mizera, JHEP **05**, 153 (2019), arXiv:1901.11510 [hep-ph].
- [130] J. Broedel, C. Duhr, F. Dulat, and L. Tancredi, JHEP **05**, 093 (2018), arXiv:1712.07089 [hep-th].
- [131] J. L. Bourjaily *et al.*, in *Snowmass 2021* (2022) arXiv:2203.07088 [hep-ph].
- [132] H. Johansson and A. Ochirov, JHEP **09**, 040 (2019), arXiv:1906.12292 [hep-th].
- [133] Z. Bern, J. J. M. Carrasco, and H. Johansson, Phys. Rev. D **78**, 085011 (2008), arXiv:0805.3993 [hep-ph].
- [134] Z. Bern, J. J. M. Carrasco, and H. Johansson, Phys. Rev. Lett. **105**, 061602 (2010), arXiv:1004.0476 [hep-th].
- [135] A. Brandhuber, G. Chen, G. Travaglini, and C. Wen, JHEP **07**, 047 (2021), arXiv:2104.11206 [hep-th].
- [136] A. Brandhuber, G. Chen, H. Johansson, G. Travaglini, and C. Wen, Phys. Rev. Lett. **128**, 121601 (2022), arXiv:2111.15649 [hep-th].
- [137] A. Brandhuber, G. R. Brown, G. Chen, J. Gowdy, G. Travaglini, and C. Wen, JHEP **12**, 101 (2022), arXiv:2208.05886 [hep-th].
- [138] L. Cangemi, M. Chiodaroli, H. Johansson, A. Ochirov, P. Pichini, and E. Skvortsov, (2023), arXiv:2312.14913 [hep-th].
- [139] Y. F. Bautista, G. Bonelli, C. Iossa, A. Tanzini, and Z. Zhou, (2023), arXiv:2312.05965 [hep-th].

Appendix: The Compton amplitude

The three point amplitude with one graviton momentum p_1^μ and classical black hole spin a^μ [48, 49, 51, 52, 116, 132] is

$$M_3(p_1 \cdot a) = \sqrt{32\pi G_N} (\bar{p} \cdot \varepsilon_1)(w_1 \cdot \varepsilon_1), \quad (31)$$

where $\bar{p}^\mu = mv^\mu$ denotes massive particle's momentum,

$$w_1^\mu := \cosh(p_1 \cdot a) \bar{p}^\mu - i \frac{\sinh(p_1 \cdot a)}{p_1 \cdot a} (p_1 \cdot S)^\mu, \quad (32)$$

and $(p_1 \cdot S)^\mu = p_{1\nu} S^{\nu\mu}$. The $S^{\mu\nu}$ denotes the spin tensor $S^{\mu\nu} = -\epsilon^{\mu\nu\rho\sigma} \bar{p}_\rho a_\sigma$.

The Compton amplitude with graviton momenta $p_{1,2}$ is constructed from the double copy [133, 134], kinematic Hopf algebra [135–137], and classical spin bootstrap [115, 116],

$$\begin{aligned} \frac{M_4}{32\pi G_N} = & - \left(\frac{\bar{p} \cdot F_1 \cdot F_2 \cdot \bar{p}}{p_2 \cdot \bar{p}} \right) \left(\frac{w_1 \cdot F_1 \cdot F_2 \cdot w_2}{2(p_1 \cdot p_2)(p_1 \cdot \bar{p})} - \frac{p_1 \cdot \bar{p} - p_2 \cdot \bar{p}}{4(p_1 \cdot p_2)(p_1 \cdot \bar{p})} \left(iG_2(x_1, x_2) (a \cdot F_1 \cdot F_2 \cdot S \cdot p_2) + iG_2(x_1, x_2) (a \cdot F_2 \cdot F_1 \cdot S \cdot p_1) \right. \right. \\ & \left. \left. + iG_1(x_{12}) \text{tr}(F_1 \cdot S \cdot F_2) + G_1(x_1)G_1(x_2) \left((a \cdot F_1 \cdot \bar{p})(a \cdot F_2 \cdot p_1) - (a \cdot F_1 \cdot p_2)(a \cdot F_2 \cdot \bar{p}) - \frac{p_2 \cdot \bar{p} - p_1 \cdot \bar{p}}{2} (a \cdot F_1 \cdot F_2 \cdot a) \right) \right) \right) \\ & + \frac{\left((\partial_{x_1} - \partial_{x_2}) G_1(x_1) G_1(x_2) \right)}{4(\bar{p} \cdot p_1)(\bar{p} \cdot p_2)} \left(\bar{p} \cdot p_2 (\bar{p}^2 (a \cdot F_1 \cdot F_2 \cdot a)(a \cdot F_2 \cdot F_1 \cdot \bar{p}) + a^2 (\bar{p}_4 \cdot F_1 \cdot F_2 \cdot \bar{p})(a \cdot F_1 \cdot F_2 \cdot \bar{p})) - (1 \leftrightarrow 2) \right) \\ & + \left(\frac{i(\partial_{x_1} - \partial_{x_2}) G_2(x_1, x_2)}{4(\bar{p} \cdot p_1)(\bar{p} \cdot p_2)} \right) \left((\bar{p} \cdot p_2) (a \cdot F_2 \cdot F_1 \cdot \bar{p}) \left((a \cdot F_2 \cdot \bar{p})(a \cdot \tilde{F}_1 \cdot \bar{p}) - (a \cdot F_1 \cdot \bar{p})(a \cdot \tilde{F}_2 \cdot \bar{p}) \right) + (1 \leftrightarrow 2) \right) \\ & + \left(\frac{(\partial_{x_1} - \partial_{x_2})^2}{2!} G_1(x_1) G_1(x_2) \right) \left((a \cdot F_1 \cdot \bar{p})(a \cdot F_2 \cdot \bar{p})(a \cdot F_1 \cdot F_2 \cdot a) - \frac{a^2}{2} \left((a \cdot F_1 \cdot F_2 \cdot p)(a \cdot F_2 \cdot F_1 \cdot \bar{p}) - (a \cdot F_1 \cdot F_2 \cdot a)(\bar{p} \cdot F_1 \cdot F_2 \cdot \bar{p}) \right) \right) \\ & + \left(\frac{i(\partial_{x_1} - \partial_{x_2})^2}{2!} G_2(x_1, x_2) \right) \left(-\frac{1}{2} \left((a \cdot F_1 \cdot F_2 \cdot a)(a \cdot F_2 \cdot \bar{p})(a \cdot \tilde{F}_1 \cdot \bar{p}) - (1 \leftrightarrow 2) \right) \right), \quad (33) \end{aligned}$$

where $x_i = a \cdot p_i$, $F_i^{\mu\nu} = p_i^\mu \varepsilon_i^\nu - \varepsilon_i^\mu p_i^\nu$, and $\tilde{F}^{\mu\nu} = \frac{1}{2} \epsilon^{\mu\nu\rho\sigma} F_{\rho\sigma}$. The differences between this Compton amplitude and other proposals [138, 139] stem from the differences in their respective formalisms and assumptions [115, 116], which are inherited by the one-loop amplitude [124]. The recursion relation for the integral representation of the G functions is

$$G_1(x_1) = \int_0^1 d\sigma_1 \cosh(\sigma_1 x_1), \quad (34)$$

$$G_2(x_1, x_2) = \int_0^1 d\sigma_2 \left[\partial_{x_1} G_1(x_1 + \sigma_2 x_2) - \sinh(\sigma_2 x_2) \cosh(\sigma_1 x_1) \right], \quad (35)$$

$$G_r(x_1, \dots, x_r) = \int_0^1 d\sigma_r \left[\partial_{x_1} G_{r-1}(x_1 + \sigma_r x_r, x_2, \dots, x_{r-1}) - G_{r-1}(x_1, \dots, x_{r-1}) \sinh(\sigma_r x_r) \right]. \quad (36)$$