

Erratum

Symbiotic Bacteria Protect Wasp Larvae from Fungal Infestation

Martin Kaltenpoth,^{1,2} Wolfgang Göttler,¹ Gudrun Herzner,¹ and Erhard Strohm^{1,*}

¹Department of Animal Ecology and Tropical Biology

University of Würzburg

Am Hubland 97074 Würzburg

Germany

²Department of Behavioral Physiology and Sociobiology

University of Würzburg

Am Hubland 97074 Würzburg

Germany

*Correspondence: martin.kaltenpoth@biozentrum.uni-wuerzburg.de

As a result of a printing error in the March 8, 2005 issue of *Current Biology*, pp. 475–479, the data in [Figure 2](#) in this paper were compromised. A reprinted figure appears below. *Current Biology* regrets the error.




Figure 2. Phylogenetic Position of Beewolf Endosymbionts within the Actinomycetes. First of three equally parsimonious trees from a full heuristic search with TBR branch swapping and random addition sequence (100 replicates). Analysis is based on 1324 bp of 16S rDNA, with 219 characters being parsimony informative. *A. globiformis* was defined as the outgroup. Values at the nodes represent bootstrap values from 1000 replicates. GenBank accession numbers are given behind the species names. *P. triangulum* specimens were collected at three different locations: in Schweinfurt (Germany SW), in Würzburg (Germany WÜ), and in the Ukraine.