

SHORT REPORT

Assessing the impact of *FOXP1* mutations on developmental verbal dyspraxia

Sonja C Vernes¹, Kay D MacDermot², Anthony P Monaco¹ and Simon E Fisher^{*,1}

¹Wellcome Trust Centre for Human Genetics, University of Oxford, Roosevelt Drive, Oxford, UK; ²North West Thames Regional Genetics Service (Kennedy-Galton Centre), North West London Hospitals NHS Trust, Harrow, UK

Neurodevelopmental disorders that disturb speech and language are highly heritable. Isolation of the underlying genetic risk factors has been hampered by complexity of the phenotype and potentially large number of contributing genes. One exception is the identification of rare heterozygous mutations of the *FOXP2* gene in a monogenic syndrome characterised by impaired sequencing of articulatory gestures, disrupting speech (developmental verbal dyspraxia, DVD), as well as multiple deficits in expressive and receptive language. The protein encoded by *FOXP2* belongs to a divergent subgroup of forkhead-box transcription factors, with a distinctive DNA-binding domain and motifs that mediate hetero- and homodimerisation. *FOXP1*, the most closely related member of this subgroup, can directly interact with *FOXP2* and is co-expressed in neural structures relevant to speech and language disorders. Moreover, investigations of songbird orthologues indicate that combinatorial actions of the two proteins may play important roles in vocal learning, leading to the suggestion that human *FOXP1* should be considered a strong candidate for involvement in DVD. Thus, in this study, we screened the entire coding region of *FOXP1* (exons and flanking intronic sequence) for nucleotide changes in a panel of probands used earlier to detect novel mutations in *FOXP2*. A non-synonymous coding change was identified in a single proband, yielding a proline-to-alanine change (P215A). However, this was also found in a random control sample. Analyses of non-coding SNP changes did not find any correlation with affection status. We conclude that *FOXP1* mutations are unlikely to represent a major cause of DVD.

European Journal of Human Genetics (2009) 17, 1354–1358; doi:10.1038/ejhg.2009.43; published online 8 April 2009

Keywords: speech and language; developmental verbal dyspraxia; transcription factor; *FOXP2*; *FOXP1*; heterodimerisation

Introduction

Developmental speech and language disorders are highly heritable, but the identification of genetic risk factors through classical mapping or association studies is hampered by genotypic and phenotypic complexity.¹ The implication of *FOXP2* in a rare monogenic form of

disorder² provides novel entry points into the critical molecular pathways.³ A heterozygous missense mutation of *FOXP2* co-segregates with speech and language disorder in the well-studied multigenerational KE family,² disrupting the function of the encoded protein.^{4,5} People carrying this mutation have problems sequencing mouth movements underlying speech (developmental verbal dyspraxia, DVD; MIM: 602081), along with impaired expressive and receptive language skills whether oral or written. Further cases of *FOXP2* disruptions causing verbal dyspraxia include a heterozygous nonsense mutation co-segregating with impairment in a small pedigree⁶ and several gross chromosomal rearrangements.^{7–9} However,

*Correspondence: Dr SE Fisher, Wellcome Trust Centre for Human Genetics, University of Oxford, Roosevelt Drive, Headington, Oxford, OX3 7BN, UK.

Tel: +44 1865 287647; Fax: +44 1865 287533;

E-mail: simon.fisher@well.ox.ac.uk

Received 23 December 2008; accepted 13 February 2009; published online 8 April 2009

aetiological point mutations of *FOXP2* likely account for only ~2% of children with DVD;⁶ other key genetic risk factors remain to be discovered. Importantly, data from functional studies of *FOXP2* can identify related genes acting in the same pathways to be considered as candidates for involvement in disorder.

FOXP2 encodes a member of the FOX group of transcription factors, featuring a characteristic forkhead-box DNA-binding domain. It belongs to a divergent subgroup (*FOXP1–4*) displaying a number of distinctive characteristics. Co-immunoprecipitation studies of murine orthologues showed that *Foxp1*, *Foxp2* and *Foxp4* form homo- and heterodimers, which are thought to be necessary for their efficient binding to target DNA.¹⁰ In contrast, most other forkhead proteins act as monomers.¹⁰ Furthermore, a *FOXP3* mutation that disturbs dimerisation results in immunodysregulation, polyendocrinopathy, enteropathy, X-linked syndrome.¹¹ Overall, for this subgroup of forkhead transcription factors, activity and/or specificity may be determined by the relative levels and combinations of different FOX proteins at a given time point, developmental stage or tissue/cell type.

Among the FOX subfamily, *FOXP1* is the most likely protein to influence language-related pathways in concert with *FOXP2*. The two proteins show a particularly high sequence homology and can cooperatively regulate downstream targets through heterodimerisation. Studies of *FoxP1* in diverse vertebrates, including humans, mice and songbirds, indicate several key sites of neural expression, displaying a significant overlap with *FoxP2* expression.¹² In the developing mouse lung (another site of co-expression), *Foxp1* and *Foxp2* have been shown to bind the promoter of a shared target gene (*T1 α*), acting together to modulate its expression.¹³ Moreover, although mice with heterozygous *Foxp1* disruption develop normally, and those with homozygous *Foxp2* loss show a subtly altered postnatal alveolarisation, combining both genotypes (*Foxp1* +/-, *Foxp2* -/-) yields much more severe lung defects, consistent with cooperative roles.¹³

Songbird studies further suggest that coordinated effects of *FOXP2* and *FOXP1* are relevant for spoken language. Like humans, songbirds display both innate vocalisations and vocal learning.¹⁴ In zebrafinches, only males learn and modify courtship song, doing so through the song system – neuronal networks spanning the cortical mantle, striatum and thalamus.¹² A striatal nucleus, area X, is necessary for song development and vocalisation, and is only present in the male brain. *FoxP2* and *FoxP1* are well conserved in zebrafinches, and are expressed in several parts of the song system, including high levels in area X and surrounding striatum.^{12,15} Knockdown of area-X *FoxP2* expression during song development results in inaccurate and incomplete imitation of tutor songs,¹⁴ suggesting important postnatal roles in auditory-guided motor learning. However, *FoxP2* does not show sexually dimorphic expression; there is no consistent differentiation between its

expression in area X of males and that in the corresponding region in females.¹² This suggests either that females have an unrealised potential for vocal learning or that this process is influenced by a sexually dimorphic co-regulator of *FoxP2*. Intriguingly, *FoxP1* expression in area X of male zebrafinches displays a sexual dimorphism closely resembling the pattern observed for the song circuit itself.¹² Thus, in songbirds, dimerisation with *FoxP1* may confer sexually dimorphic activity on *FoxP2* during vocal learning.

Integrating the above findings, it is plausible that development/function of language-related circuits in the human brain could be disturbed by imbalances in relative functional dosage of *FOXP2* and *FOXP1*. Thus, as explicitly proposed by Teramitsu *et al*,¹² *FOXP1* represents a strong functional candidate for involvement in speech disorders. In this study, we directly tested this hypothesis by mutation screening of all coding *FOXP1* exons and flanking splice sites in children with DVD.

Materials and methods

Children with DVD

A panel of 49 probands was assembled based on a primary clinical diagnosis of DVD (see Supplementary methods). Earlier analyses of this panel successfully identified novel mutations in the *FOXP2* gene.⁶

Genomic organisation of *FOXP1*

The intron/exon structure of *FOXP1* was determined by aligning GenBank entry AF146696 (*FOXP1* mRNA) with human genomic sequence through the UCSC Genome Browser (hg17 assembly, NCBI Build 35).

Denaturing high-performance liquid chromatography screening

Primers were designed to amplify the 16 coding exons of *FOXP1* (Supplementary Table S1), each with a fragment size optimal for denaturing high-performance liquid chromatography (DHPLC) (200–600 bp). After amplification using touchdown PCR, fragments were analysed through the Transgenomics WAVE DHPLC system followed by direct sequencing (Supplementary methods).

Results

FOXP1 consists of 16 coding exons spanning ~586 kb on 3p14.1 (Figure 1a). We screened these exons and flanking intronic sequence in 49 probands with clinically diagnosed DVD plus their siblings (59 individuals in total). DHPLC analysis, followed by sequencing of variants, revealed one exonic (coding) change and four intronic (non-coding) polymorphisms (Table 1). The coding change, identified in a singleton proband, was located in exon 5. This change constituted a heterozygous C-to-G transversion, yielding a proline-to-alanine substitution at position 215 of the




Figure 1 Mutation screening of *FOXP1* in verbal dyspraxia probands. (a) Genomic organisation of *FOXP1*. *FOXP1* is located on chromosome 3p14.1 and consists of 16 coding exons spanning 586 kb. Exons are represented by filled bars whose width is proportional to the length of the exon. Numbering scheme is based on alignment with the published mouse *Foxp1* structure.²⁰ Initiation codon is indicated by ‘atg’ and stop codon by ‘tga’. The Zinc-Finger domain spans exons 7 and 8, the Leucine-Zipper domain spans exons 8 and 9 and the forkhead-box motif is encoded by exons 10–13. Locations of base changes summarised in Table 1 are indicated on figure by bars, with base change and frequency in probands. (b) Direct sequencing confirmed the presence of base changes in probands displaying aberrant DHPLC elution patterns (see also Table 1). The exon 5 polymorphism is shown for the proband and HRC samples that carried the C-to-G transversion, aligned with a proband that did not carry the change for comparison.

Table 1 Base changes identified through DHPLC mutation screening and direct sequencing

Exon	Change	DbSNP reference	Intronic/exonic	Position ^a	No. of individuals (N= 49)
2	G-to-A	—	Intronic	–28 bp	1
5	C-to-G	—	Exonic	+132 bp	1
9	C-to-T	—	Intronic	+13 bp	1
15	C-to-A	rs7638391	Intronic	+19 bp	10
16	G-to-T	—	Intronic	–14 bp	1

^aPosition is given as upstream (–) or downstream (+) of exon/intron boundary.

encoded protein. Alignments of FOXP proteins from different species indicated that this was one of the few highly conserved residues in a region of low homology (Figure 2). To determine if this change was exclusive to DVD, 384 control chromosomes from Human Random Control (HRC) panels were screened using identical methods to the clinical samples. The same heterozygous C-to-G change was identified in a single chromosome within the controls (Figure 1b).

A heterozygous C-to-A transversion in the intron downstream of exon 15 (Figure 1, Table 1) was found in 10 of the 49 probands (an SNP frequency of ~10%). We tested whether this non-coding SNP might be in linkage disequilibrium with an undiscovered functional variant. DHPLC screening of 146 HRC samples identified the SNP in 20 controls (an SNP frequency of ~7%). As the SNP frequency is not significantly different between DVD cases and controls (χ^2 *P*-value > 0.25), it is unlikely to be associated with the disorder. The same heterozygous SNP was reported on dbSNP (<http://www.ncbi.nlm.nih.gov/SNP/>) at ~12.5% in populations of European descent. Again, the frequency did not significantly differ from that of our DVD cases (χ^2 *P*-value > 0.5).

Discussion

A range of *FOXP2* mutations has been identified in DVD, clearly showing the aetiological importance of this gene.^{2,6–9} Given the high heritability of speech/language impairments it is highly likely that other genes contribute. Here, we investigated the *FOXP1* functional candidate by mutation screening in a panel of cases that had earlier enabled us to identify *FOXP2* coding changes.⁶

A heterozygous *FOXP1* coding change was identified in a proband with no affected siblings. The resulting

substitution (P215A) replaced a proline with an alanine in the region encoded by exon 5. The region lies outside known functional domains and its structure is unknown. However, alignments of FOXP orthologues from diverse species (human to *Drosophila melanogaster*) indicate the high conservation of this proline (Figure 2). Mutations affecting proline residues commonly affect protein structure/stability.¹⁶ Owing to their rigid conformation, prolines frequently induce bends or kinks. Although a proline may sometimes enhance structural stability (eg, as the first residue of an α -helix), it often disrupts the secondary structure (eg, when located internally within an α -helix or a β -sheet).¹⁷ Thus, P215 could be important for maintaining local structure. Nonetheless, in the absence of a thorough characterisation of the structure of full-length FOXP1, it is difficult to determine the structural impact of the P215A substitution.

It is worth noting that a heterozygous P215A change was also identified in one of 384 control chromosomes from HRC panels (compared with a frequency of 1-in-98 chromosomes in the DVD panel). The possibility remains that this change produces a functional effect, particularly as developmental language disorders are observed in up to 7% of school-age children.¹⁸ Given that information is unavailable regarding phenotypes of HRC individuals, we cannot exclude the possibility that the control individual carrying the P215A change was affected with a mild speech/language disorder.

Despite evidence from human, mouse and songbirds suggesting the importance of FoxP1–FoxP2 neural interactions in pathways mediating speech and language, our study found no indication of clear correlations between *FOXP1* variants and DVD risk. Nevertheless, given emerging roles for this gene in nervous system development,¹⁹ it continues to represent a candidate for involvement in

Human	FOXP1 (Q9H334)	206	TIQPG--QPAL	PLQPLAQG	MIPTELQQLWKEVTS	SAHTAEETTGN	---HSSLDLTTTCVS	260
Mouse	Foxp1 (P58462)	234	TIQPG--QPAL	PLQPLAQG	MIPTELQQLWKEVTS	SAHTAEETTSS	---HSSLDLTSTCVS	288
Chicken	FOXP1 (Q58NQ4)	215	TIQPG--QPTL	PLQPLAQG	MIPTELQQLWKEVTS	SHTAEEAASNN	---HSSLDLSTTCVS	269
Xenopus	FoxP1 (Q5W1J5)	108	SIQPG--QPTL	PLQSLAQG	MIPAEELQQLWKEVT	GSHTADDVVCNN	---HSTLDLSTTCVS	162
Zebrafish	FoxP1 (Q2LE08)	193	SIQPN--Q-TL	PLHTLPQG	MIPAEELQQLWKEVT	NSHVKEENSVTN	NGHRGLDLS---	245
Human	FOXP2 (O15409)	241	SIPPG--QAAL	PVQSLPQAG	SPAELQQLWKEVT	GVHSMED---	NGIKHGGLDLTTNNS	297
Chimpanzee	FoxP2 (Q8MJA0)	244	SIPPG--QAAL	PVQSLPQAG	SPAELQQLWKEVT	GVHSMED---	NGIKHGGLDLTTNNS	298
Mouse	Foxp2 (P58463)	242	SIPPG--QAAL	PVQSLPQAG	SPAELQQLWKEVT	GVHSMED---	NGIKHGGLDLTTNNS	296
Zebra finch	FoxP2 (Q0QM04)	239	SIPPG--QSAL	PVQSLPQAG	SPAELQQLWKEVT	GVHSMED---	NGIKHGGLDLTTNNS	293
Xenopus	FoxP2 (Q4VYS1)	234	SIPPS--QSAL	PVQSLPQAG	SPAELQQLWKEVT	GVHSMED---	NGIKHGGLDLTTNIS	288
Zebrafish	FoxP2 (Q4JNX5)	203	SMPPGPGQPTL	PLGQTLPPAG	SPAELQQLWKDVT	TASHTMED---	NGMKHSGLDLSTNNNT	259
Human	Foxp4 (Q8IVH2)	204	SLQPN--QASG	PLQTLPQAA	CPTDLPQLWKGE	GAPGQPAE---	DSVKQEGDLTGTAAT	258
Mouse	Foxp4 (Q9DBY0)	210	SLQPS--QASG	PLQALPQ-AA	CPTDLPQLWKGE	GAPGQPAE---	DSGRQEGDLASTAVT	263
Xenopus	FoxP4 (Q4VYR7)	189	GLQSG--QGVP	PMQSLPQ--	VSPSDLHQLLKEM	SS---SQE---	ESSKQDVTDLMTSIT-	237
Human	Foxp3 (Q9BZS1)	111	AHARTPVLQVH	PLESPAMIS	TPPTATGVSFLK	KARPLPPG---	-----	152
Mouse	Foxp3 (Q99JB6)	110	AHAQT PVLQVR	PLDNPAMIS	LPPPSAATGVF	SLKARPLPPG---	-----	151
Drosophila	FoxP (Q4V6X1)	100	APVPDLGFYNV	PEFFISEQEK	MFSDAERFLR	SKDNEVCNND---	-----	140

*

Figure 2 Proline-215 is conserved within the FOXP subfamily. Amino acids 206–260 encoded by exon 5 of *FOXP1* were aligned (using CLUSTALW) with a range of sequences from FoxP1, FoxP2 and FoxP4 plus the closest drosophila homologue to the FOXP subgroup. Swiss-Prot/TrEMBL accession numbers are shown in parentheses. Conserved amino acids are shaded in black and similar amino acids are shaded in grey. Proline-215 is indicated by an asterisk and is completely conserved across all available FOXP sequences.

neurodevelopmental disorders. Further investigations of *FoxP1/FoxP2* in model systems promise greater insights into their coordinated effects on brain function.

Acknowledgements

We thank Nuala Sykes and Thomas Scerri for assistance in setting up the DHPLC mutation analysis and Thomas Bajorek for assistance during screening. The project was funded by the Wellcome Trust and Autism Speaks. SCV was supported by a Christopher Welch Biological Sciences Scholarship from the University of Oxford. SEF is a Royal Society Research Fellow.

References

- 1 Fisher SE, Lai CS, Monaco AP: Deciphering the genetic basis of speech and language disorders. *Annu Rev Neurosci* 2003; **26**: 57–80.
- 2 Lai CS, Fisher SE, Hurst JA, Vargha-Khadem F, Monaco AP: A forkhead-domain gene is mutated in a severe speech and language disorder. *Nature* 2001; **413**: 519–523.
- 3 Vernes SC, Spiteri E, Nicod J *et al*: High-throughput analysis of promoter occupancy reveals direct neural targets of FOXP2, a gene mutated in speech and language disorders. *Am J Hum Genet* 2007; **81**: 1232–1250.
- 4 Groszer M, Keays DA, Deacon RM *et al*: Impaired synaptic plasticity and motor learning in mice with a point mutation implicated in human speech deficits. *Curr Biol* 2008; **18**: 354–362.
- 5 Vernes SC, Nicod J, Elahi FM *et al*: Functional genetic analysis of mutations implicated in a human speech and language disorder. *Hum Mol Genet* 2006; **15**: 3154–3167.
- 6 MacDermot KD, Bonora E, Sykes N *et al*: Identification of FOXP2 truncation as a novel cause of developmental speech and language deficits. *Am J Hum Genet* 2005; **76**: 1074–1080.
- 7 Feuk L, Kalervo A, Lipsanen-Nyman M *et al*: Absence of a paternally inherited FOXP2 gene in developmental verbal dyspraxia. *Am J Hum Genet* 2006; **79**: 965–972.
- 8 Lennon PA, Cooper ML, Peiffer DA *et al*: Deletion of 7q31.1 supports involvement of FOXP2 in language impairment: clinical report and review. *Am J Med Genet A* 2007; **143**: 791–798.
- 9 Shriberg LD, Ballard KJ, Tomblin JB, Duffy JR, Odell KH, Williams CA: Speech, prosody, and voice characteristics of a mother and daughter with a 7;13 translocation affecting FOXP2. *J Speech Lang Hear Res* 2006; **49**: 500–525.
- 10 Li S, Weidenfeld J, Morrisey EE: Transcriptional and DNA binding activity of the Foxp1/2/4 family is modulated by heterotypic and homotypic protein interactions. *Mol Cell Biol* 2004; **24**: 809–822.
- 11 Chatila TA, Blaaser F, Ho N *et al*: JM2, encoding a fork head-related protein, is mutated in X-linked autoimmunity-allergic dysregulation syndrome. *J Clin Invest* 2000; **106**: R75–R81.
- 12 Teramitsu I, Kudo LC, London SE, Geschwind DH, White SA: Parallel FoxP1 and FoxP2 expression in songbird and human brain predicts functional interaction. *J Neurosci* 2004; **24**: 3152–3163.
- 13 Shu W, Lu MM, Zhang Y, Tucker PW, Zhou D, Morrisey EE: Foxp2 and Foxp1 cooperatively regulate lung and esophagus development. *Development* 2007; **134**: 1991–2000.
- 14 Haesler S, Rochefort C, Georgi B, Licznarski P, Osten P, Scharff C: Incomplete and inaccurate vocal imitation after knockdown of FoxP2 in songbird basal ganglia nucleus Area X. *PLoS Biol* 2007; **5**: e321.
- 15 Haesler S, Wada K, Nshdejan A *et al*: FoxP2 expression in avian vocal learners and non-learners. *J Neurosci* 2004; **24**: 3164–3175.
- 16 Bajaj K, Madhusudhan MS, Adkar BV *et al*: Stereochemical criteria for prediction of the effects of proline mutations on protein stability. *PLoS Comput Biol* 2007; **3**: e241.
- 17 Vanhoof G, Goossens F, De Meester I, Hendriks D, Scharpe S: Proline motifs in peptides and their biological processing. *FASEB J* 1995; **9**: 736–744.
- 18 Tomblin JB, Records NL, Buckwalter P, Zhang X, Smith E, O'Brien M: Prevalence of specific language impairment in kindergarten children. *J Speech Lang Hear Res* 1997; **40**: 1245–1260.
- 19 Dasen JS, De Camilli A, Wang B, Tucker PW, Jessell TM: Hox repertoires for motor neuron diversity and connectivity gated by a single accessory factor, FoxP1. *Cell* 2008; **134**: 304–316.
- 20 Wang B, Lin D, Li C, Tucker P: Multiple domains define the expression and regulatory properties of Foxp1 forkhead transcriptional repressors. *J Biol Chem* 2003; **278**: 24259–24268.



This work is licensed under the Creative Commons Attribution-NonCommercial-No Derivative Works 3.0 Licence. To view a copy of this licence, visit <http://creativecommons.org/licenses/by-nc-nd/3.0/>

Supplementary Information accompanies the paper on European Journal of Human Genetics website (<http://www.nature.com/ejhg>)