

Accurate Molecular Van Der Waals Interactions from Ground-State Electron Density and Free-Atom Reference Data

Alexandre Tkatchenko and Matthias Scheffler

Fritz-Haber-Institut der Max-Planck-Gesellschaft, Faradayweg 4-6, 14195, Berlin, Germany

(Received 3 November 2008; published 20 February 2009)

We present a parameter-free method for an accurate determination of long-range van der Waals interactions from mean-field electronic structure calculations. Our method relies on the summation of interatomic C_6 coefficients, derived from the electron density of a molecule or solid and accurate reference data for the free atoms. The mean absolute error in the C_6 coefficients is 5.5% when compared to accurate experimental values for 1225 intermolecular pairs, irrespective of the employed exchange-correlation functional. We show that the effective atomic C_6 coefficients depend strongly on the bonding environment of an atom in a molecule. Finally, we analyze the van der Waals radii and the damping function in the C_6R^{-6} correction method for density-functional theory calculations.

DOI: 10.1103/PhysRevLett.102.073005

PACS numbers: 31.15.eg, 71.15.Mb, 87.15.A-

Noncovalent forces, such as hydrogen bonding and van der Waals (vdW) interactions, are crucial for the formation, stability, and function of molecules and materials. At present, ubiquitous vdW interactions [1] can only be accounted for properly by high-level quantum-chemical wave function or by the Quantum Monte Carlo (QMC) method. In contrast, the correct long-range interaction tail, e.g., for separated molecules, is absent from all popular local-density or gradient corrected exchange-correlation (xc) functionals of density-functional theory (called DFT-XCA in what follows), as well as from the Hartree-Fock (HF) approximation [2,3].

Many encouraging concepts and methods have been proposed to include vdW interactions in DFT calculations [4–9]. Ultimately, a functional which is able to account for vdW interactions in a “seamless” manner is desirable. The Chalmers-Rutgers approach [4,10] is a step in that direction, but at this point, the performance is not certain and, for example, errors are as large as 70% for the binding energy of rare-gas dimers—prototypical vdW systems [10].

A popular remedy for the missing vdW interaction in present-day DFT consists of adding a pairwise interatomic C_6R^{-6} term (E_{vdW}) to the DFT energy [5,6,11–14],

$$E_{\text{vdW}} = -\frac{1}{2} \sum_{A,B} f_{\text{damp}}(R_{AB}, R_A^0, R_B^0) C_{6AB} R_{AB}^{-6}, \quad (1)$$

where R_{AB} is the distance between atoms A and B , C_{6AB} is the corresponding C_6 coefficient, R_A^0 and R_B^0 are the vdW radii. The R_{AB}^{-6} singularity at small distances is eliminated by the short-ranged damping function $f_{\text{damp}}(R_{AB}, R_A^0, R_B^0)$. Obviously, at the medium and short range, this approach can only work together with xc functionals that underestimate the binding energy. In particular, Grimme has proven such schemes to be accurate for a range of molecular systems [6,12]. A serious shortcoming of the C_6R^{-6} schemes is their empirical nature, since the parameters do not depend on the electronic structure, but are rather ob-

tained by fitting to experimental C_6 coefficients and/or post-Hartree-Fock binding energy data. Furthermore, we note that the damping function will also correct (or affect) other properties of the employed xc functional at short distances. Though a correction of present-day xc functionals is necessary, it is not satisfactorily handled by such an approach. Several methods exist to determine the C_6 coefficients either from ground-state orbitals [8,9,15] or time-dependent DFT (TDDFT) [16,17]. Unfortunately, the errors are quite large (15%–20% on average and maximum deviation of 40%–60%) [8,15,16]. The origin of such errors is related to the grossly overestimated polarizability in DFT-XCA [18,19].

In this Letter, we develop and assess a scheme to determine the C_6 coefficients and vdW radii from the mean-field ground-state electron density (DFT-XCA or HF) for molecules and solids. It is largely independent of the employed DFT-XCA approximation [tested here for local-density approximation (LDA), Perdew-Burke-Ernzerhof (PBE), and Becke-Lee-Yang-Parr (BLYP) functionals], and it shows a mean absolute error of 5.5% for intermolecular C_6 coefficients on a database of experimental dipole oscillator strength distribution (DOSD) data of Meath and co-workers for 1225 complexes (see, e.g., Refs. [5,8,20,21]). The DOSD for a given molecule is the (differential) dipole oscillator strength df/dE , as a function of excitation energy E , from the electronic absorption threshold E_0 to very high energies. Many important molecular dipole properties can be evaluated as integrals involving the DOSD [22]. The critical idea of our method is to use the electron density to compute the *relative* and not the absolute polarizability of an atom in a molecule. The method includes charge polarization effects in a transparent way, as shown for different atomic hybridization states and hydrogen bonding.

In order to develop our method, we start with the exact expression (Casimir-Polder integral) for the leading isotropic C_6 term describing the vdW interaction between two atoms or molecules A and B (Hartree atomic units used

throughout) [23],

$$C_{6AB} = \frac{3}{\pi} \int_0^\infty \alpha_A(i\omega)\alpha_B(i\omega)d\omega, \quad (2)$$

where $\alpha_{A/B}(i\omega)$ is the frequency-dependent polarizability of A and B evaluated at imaginary frequencies. We retain only the leading term in the Padé series [24] for $\alpha_{A/B}(i\omega)$, yielding for $\alpha_A^1(\omega)$

$$\alpha_A^1(\omega) = \alpha_A^0/[1 - (\omega/\eta_A)^2], \quad (3)$$

where α_A^0 is the static polarizability of A and η_A is an *effective frequency*. Substituting $\alpha^1(i\omega)$ of Eq. (3) for $\alpha(i\omega)$ in Eq. (2) yields the London formula [25],

$$C_{6AB} = \frac{3}{2}[\eta_A\eta_B/(\eta_A + \eta_B)]\alpha_A^0\alpha_B^0. \quad (4)$$

For $A = B$, we obtain

$$\eta_A = \frac{4}{3} \frac{C_{6AA}}{(\alpha_A^0)^2}, \quad (5)$$

with a corresponding expression for η_B . Combining Eqs. (4) and (5), we arrive at a formula for C_{6AB} which depends only on homonuclear parameters C_{6AA} , C_{6BB} , α_A^0 , and α_B^0 ,

$$C_{6AB} = \frac{2C_{6AA}C_{6BB}}{\left[\frac{\alpha_B^0}{\alpha_A^0}C_{6AA} + \frac{\alpha_A^0}{\alpha_B^0}C_{6BB}\right]}. \quad (6)$$

For the free-atom reference values of α_A^0 and C_{6AA} , we rely on the database of Chu and Dalgarno [16], which reports self-interaction corrected TDDFT values scaled to reproduce accurate all-order many-body calculations for rare gases, alkalis, and alkaline earth atoms. The accuracy of the free-atom results is presumed to be better than 3% for α_A^0 and C_{6AA} for nonmetallic elements (1% for rare gases, alkalis, and alkaline earth atoms). Using these homonuclear values along with Eq. (6), we obtain a mean absolute relative error (MARE) of just 2.7% on a database of 70 heteronuclear C_6 coefficients between light elements, rare gases, alkalis, and alkaline earth atoms from accurate many-body calculations [26–28]. The almost perfect correlation is shown in Fig. 1.

Let us now define the C_6 coefficient for an atom inside a molecule or a solid. This requires the definition of the *effective volume*, referenced to the free atom *in vacuo*. We take advantage of the direct relation between polarizability and volume [29], and employ the Hirshfeld partitioning of the electron density for the latter [8,30],

$$\frac{\kappa_A^{\text{eff}}}{\kappa_A^{\text{free}}} \frac{\alpha_A^{\text{eff}}}{\alpha_A^{\text{free}}} = \frac{V_A^{\text{eff}}}{V_A^{\text{free}}} = \left(\frac{\int r^3 w_A(\mathbf{r})n(\mathbf{r})d^3\mathbf{r}}{\int r^3 n_A^{\text{free}}(\mathbf{r})d^3\mathbf{r}} \right), \quad (7)$$

$$w_A(\mathbf{r}) = \frac{n_A^{\text{free}}(\mathbf{r})}{\sum_B n_B^{\text{free}}(\mathbf{r})}, \quad (8)$$

where κ_A^i is the proportionality constant between volume and polarizability for the free-atom and atom-in-a-molecule, $w_A(\mathbf{r})$ is the Hirshfeld atomic partitioning weight for the atom A , r^3 is the cube of the distance from




FIG. 1 (color online). Comparison of the C_6 coefficients for atom-atom interaction (At) and atom-molecule and molecule-molecule interaction (Mol). The reference results for atom-atom interaction are from accurate wave function calculations [26–28]. For molecules, DOSD results are taken as a reference [5,8,20,21]. Our results (only 211 values out of 1225 are shown) are compared to those of Chalmers-Rutgers collaboration [15] and Johnson-Becke [8]. The only outliers for our method are cases involving the H_2 molecule (20–44% deviation).

the nucleus of an atom A , $n(\mathbf{r})$ is the total electron density, $n_A^{\text{free}}(\mathbf{r})$ is the electron density of the free atom A , and the sum goes over all atoms in the system. Both $n(\mathbf{r})$ and $n_A^{\text{free}}(\mathbf{r})$ are calculated from DFT-XCA. The effective coefficient C_{6AA}^{eff} for an atom in a molecule is determined in the following way from Eqs. (4) and (5):

$$C_{6AA}^{\text{eff}} = \frac{\eta_A^{\text{eff}}}{\eta_A^{\text{free}}} \left(\frac{\kappa_A^{\text{free}}}{\kappa_A^{\text{eff}}} \right)^2 \left(\frac{V_A^{\text{eff}}}{V_A^{\text{free}}} \right)^2 C_{6AA}^{\text{free}}. \quad (9)$$

We assume the proportionality constant $\frac{\eta_A^{\text{eff}}}{\eta_A^{\text{free}}} \left(\frac{\kappa_A^{\text{free}}}{\kappa_A^{\text{eff}}} \right)^2$ to be unity and prove this choice to be remarkably good for a large variety of molecules. Indeed, this approximation breaks down only for the smallest H_2 molecule, with deviation of 44% for the C_6 coefficient. Already for molecules such as N_2 and CO_2 , the error is less than 10%. Since the static polarizability of a molecule cannot be expressed as a linear combination of atomic polarizabilities in general, and thus $\kappa_A^{\text{free}}/\kappa_A^{\text{eff}} \neq 1$, $\eta_A^{\text{eff}}/\eta_A^{\text{free}}$, and $(\kappa_A^{\text{free}}/\kappa_A^{\text{eff}})^2$ are inversely proportional.

Since the C_6 coefficients are additive [22,31], the intermolecular C_6 coefficient, C_6^{mol} , is given by the sum of all interatomic contributions

$$C_6^{\text{mol}} = \sum_{A \in M_1} \sum_{B \in M_2} C_{6AB}^{\text{eff}}, \quad (10)$$

where M_1 and M_2 refers to the first and the second molecule, respectively.

To assess the accuracy of our scheme, it was benchmarked on a database of 1225 intermolecular C_6 coefficients, derived from pseudo DOSD data of Meath and co-workers (see, e.g., Refs. [5,8,20,21]). The database con-

tains the C_6 coefficients for the interaction between 8 atoms and 42 molecules (organic and inorganic, from small dimers to C_8H_{18}). The geometry of every molecule was fully relaxed using the FHI-AIMS [32] code together with LDA, PBE, and BLYP functionals. The atomic C_6^{eff} coefficients were calculated from Eq. (9) using the DFT density and then the molecular C_6^{mol} coefficients were computed using Eq. (10). The correlation between the calculated C_6 coefficients and reference results is shown in Fig. 1 and compared with other methods for obtaining the C_6 coefficients from the ground-state density. With a MARE of 5.5%, we note that our scheme is a factor of 2–3 more accurate than existing methods [8,9,15]. Furthermore, if the H_2 molecule is excluded from the database, the MARE drops down to 4.5%. The C_6 coefficients vary only slightly for different exchange-correlation (xc) functionals. The MARE in the C_6 coefficients between LDA and PBE is 1.4% (maximum deviation of 3.8%), while for BLYP and PBE it is 0.68% (maximum deviation of 2.1%). Clearly, the difference in the electron density between xc functionals is compensated when computing the ratio in Eq. (7).

In Table I, we show the values of the C_6 coefficients for various atoms in the DOSD database. It is encouraging that the C_6 coefficients correspond very closely to those empirically fitted by Wu and Yang [5] for different hybridization states of C and O atoms. This further indicates that our scheme correctly accounts for different atomic environments without the need of empirical parameters. The largest variations occur for carbon and silicon, due to various possible hybridization states for these elements.

In order to further illustrate the change in the atomic C_6 coefficients as a function of molecular bonding and geometry, we show the case of the hydrogen-bonded water dimer in Fig. 2. As the H_2O molecules approach to form a hydrogen bond, the polarizabilities and the C_6 coefficients of all atoms are modified. The hydrogen involved in the hydrogen bond becomes significantly more polarizable along with the donor oxygen. On the other hand, the acceptor oxygen becomes less polarizable along with the attached hydrogens. The plot in Fig. 2 was done for a fixed water dimer geometry to illustrate that the change in the C_6 coefficients is a purely electronic effect. Relaxing the geometries for every O-O distance yields a similar plot for distances larger than the equilibrium one.

We carried out preliminary tests of our scheme for solids, calculating the C_6 coefficient of the carbon atom in a graphene sheet and in a diamond crystal at experimental geometries. For carbon in a graphene sheet, we get a value of 33.0, close to the expected value of 30.3 for the sp^2 hybridization in benzene. For diamond, we get a value of 38.6, significantly larger than 24.1 for the sp^3 hybridization in methane.

Our method can in principle treat all elements, including ions, on the same footing. Most problematic cases are those where the concept of *atoms-in-molecules* cannot be applied. Clearly, a pairwise summation of C_6R^{-6} interactions

TABLE I. Free-atom C_6 coefficients (hartree · bohr⁶) from Chu and Dalgarno [16] along with atom-in-a-molecule (minimum and maximum) C_6 coefficients for various atoms computed for molecules from the DOSD database. For CNOH atoms, these values are compared to empirical results of Wu and Yang [5] for different hybridization states.

	C_{6AA}^{free}	C_{6AA}^{min}	C_{6AA}^{max}	Wu-Yang		C_{6AA}^{free}	C_{6AA}^{min}	C_{6AA}^{max}
H	6.5	2.1	2.8	2.8	F	9.5	7.9	8.5
C	46.6	24.1	33.0	22.1–29.8	Si	305	146	158
N	24.2	17.1	19.6	19.3	S	134	113	134
O	15.6	11.7	14.7	11.6–13.0	Cl	94.6	88.8	89.4

can fail qualitatively for metallic low-dimensional systems due to nonadditive higher-order effects [33]. Our scheme can be further generalized to higher-order van der Waals coefficients (C_8 , etc.) since approximations similar to the London formula are known [22]. The Axilrod-Teller-Muto three-body term [22] can also be calculated since it involves an integral similar to the Casimir-Polder one.

Let us now briefly discuss the coupling of the above long-range scheme for correcting DFT-XCA calculations for shorter distances. For this, the damping function $f_{\text{damp}}(R_{AB}, R_A^0, R_B^0)$ in Eq. (1) must be defined. The vdW radii, $R_{A/B}^0$, are not experimental observables, unlike the C_6 coefficients. However, a rigorous theoretical definition does exist: the vdW radius corresponds to half of the distance between two atoms where the Pauli repulsion balances the London dispersion attraction [34]. Using the definition of the effective atomic volume in Eq. (7), the vdW radius of an atom in a molecule becomes

$$R_{\text{eff}}^0 = \left(\frac{V^{\text{eff}}}{V^{\text{free}}} \right)^{1/3} R_{\text{free}}^0. \quad (11)$$

According to the above definition, the free-atom vdW radii, R_{free}^0 , for rare-gas atoms correspond to the equilibrium distance of rare-gas dimers. Unfortunately, the vdW radii of Bondi [34] for other elements cannot be used since they correspond to atoms-in-molecules case. The simplest an-




FIG. 2 (color online). Dependence of the atomic C_6 coefficients on the O-O distance in the water dimer. Note the two different scales on both sides for H and O atoms.

satz for defining consistent free-atom vdW radii comes from the electron density for the (spherical) free atoms. The electron density contour value corresponding to the vdW radius can be determined for the rare-gas atoms and then used to define R_{free}^0 for other elements in the same row of the periodic table.

We used the above ansatz for the light elements (CNOH), to illustrate the performance of our nonempirical C_6 scheme with DFT. We use a Fermi-type damping function [5],

$$f_{\text{damp}}(R_{AB}, R_{AB}^0) = \frac{1}{1 + \exp[-d(\frac{R_{AB}}{s_R R_{AB}^0} - 1)]}, \quad (12)$$

where $R_{AB}^0 = R_A^0 + R_B^0$, d and s_R are free parameters. The d parameter adjusts the damping function steepness. Our analysis indicates that the results change negligibly for $12 < d < 45$. A small value of d affects covalently bonded systems, whereas a large value yields kinked binding energy curves. The choice of $d = 20$ turned out to satisfy both constraints as tested for binding energy curves of rare-gases and vdW-bonded organic molecule dimers (CH_4 and benzene), as also shown by Grimme in previous work [12]. Therefore, the scaling coefficient s_R remains as a single empirical parameter which determines the onset of the vdW correction for a particular xc functional in terms of the distance [14]. We use the S22 database of Jurecka *et al.* [35] to obtain the s_R parameter [36]. The database reports converged CCSD(T) binding energies for 22 different dimers with varying interaction strength, from a weakly vdW-bonded CH_4 dimer (23 meV) to a hydrogen-bonded uracil dimer (0.9 eV). The performance of our scheme, when coupled with the PBE functional, is significantly better than for highly empirical $C_6 R^{-6}$ approaches [6,12,14]. The mean absolute error (MAE) of our approach on the S22 database is 13 meV (20, 13, and 6 meV on hydrogen-bonded, vdW-bonded and mixed systems, respectively). This can be compared to 20 meV (29, 20, and 9 meV) when using empirical C_6 coefficients and vdW radii from Ref. [14]. It is especially encouraging that the overestimation of the hydrogen-bonded systems is significantly reduced due to larger effective vdW radii for atoms involved in hydrogen bonds. Since the atomic C_6 coefficients are functionals of the electron density [Eq. (9)], the potential due to the energy expression in Eq. (1) should enter the Kohn-Sham equations in DFT calculations. However, we do not expect a self-consistent treatment to give major changes as also noticed in Ref. [10]. This aspect will be addressed in future work.

In summary, we have presented an accurate nonempirical method to obtain molecular C_6 coefficients from ground-state electron density and reference values for the free atoms. Our scheme can also be used to improve the description of weakly bonded systems in DFT for a range of xc functionals.

A. T. acknowledges the Alexander von Humboldt (AvH) foundation for funding.

- [1] V. A. Parsegian, *Van der Waals Forces: A Handbook for Biologists, Chemists, Engineers and Physicists* (Cambridge University Press, Cambridge, England, 2005).
- [2] S. Kristyán and P. Pulay, *Chem. Phys. Lett.* **229**, 175 (1994).
- [3] J. M. Pérez-Jordá and A. D. Becke, *Chem. Phys. Lett.* **233**, 134 (1995).
- [4] M. Dion *et al.*, *Phys. Rev. Lett.* **92**, 246401 (2004).
- [5] Q. Wu and W. Yang, *J. Chem. Phys.* **116**, 515 (2002).
- [6] S. Grimme, *J. Comput. Chem.* **25**, 1463 (2004).
- [7] O. A. von Lilienfeld, I. Tavernelli, U. Rothlisberger, and D. Sebastiani, *Phys. Rev. Lett.* **93**, 153004 (2004).
- [8] E. R. Johnson and A. D. Becke, *J. Chem. Phys.* **123**, 024101 (2005).
- [9] P. L. Silvestrelli, *Phys. Rev. Lett.* **100**, 053002 (2008).
- [10] T. Thonhauser *et al.*, *Phys. Rev. B* **76**, 125112 (2007).
- [11] U. Zimmerli, M. Parrinello, and P. Koumoutsakos, *J. Chem. Phys.* **120**, 2693 (2004).
- [12] S. Grimme, *J. Comput. Chem.* **27**, 1787 (2006).
- [13] F. Ortman, F. Bechstedt, and W. G. Schmidt, *Phys. Rev. B* **73**, 205101 (2006).
- [14] P. Jurecka, J. Cerny, P. Hobza, and D. R. Salahub, *J. Comput. Chem.* **28**, 555 (2007).
- [15] Y. Andersson and H. Rydberg, *Phys. Scr.* **60**, 211 (1999).
- [16] X. Chu and A. Dalgarno, *J. Chem. Phys.* **121**, 4083 (2004).
- [17] S. Botti, A. Castro, X. Andrade, A. Rubio, and M. A. L. Marques, *Phys. Rev. B* **78**, 035333 (2008).
- [18] S. A. C. McDowell, R. D. Amos, and N. C. Handy, *Chem. Phys. Lett.* **235**, 1 (1995).
- [19] S. J. A. van Gisbergen *et al.*, *Phys. Rev. Lett.* **83**, 694 (1999).
- [20] D. J. Margoliash and W. J. Meath, *J. Chem. Phys.* **68**, 1426 (1978).
- [21] G. D. Zeiss and W. J. Meath, *Mol. Phys.* **33**, 1155 (1977).
- [22] A. J. Stone, *The Theory of Intermolecular Forces* (Oxford University Press, New York, 1996).
- [23] H. B. G. Casimir and B. Polder, *Phys. Rev.* **73**, 360 (1948).
- [24] K. T. Tang and M. Karplus, *Phys. Rev.* **171**, 70 (1968).
- [25] K. T. Tang, *Phys. Rev.* **177**, 108 (1969).
- [26] F. Maeder and W. Kutzelnigg, *Chem. Phys.* **42**, 95 (1979).
- [27] A. Derevianko, W. R. Johnson, M. S. Safronova, and J. F. Babb, *Phys. Rev. Lett.* **82**, 3589 (1999).
- [28] J. Mitroy and J.-Y. Zhang, *Phys. Rev. A* **76**, 032706 (2007).
- [29] T. Brink, J. S. Murray, and P. Politzer, *J. Chem. Phys.* **98**, 4305 (1993).
- [30] A. Olsz *et al.*, *J. Chem. Phys.* **127**, 224105 (2007).
- [31] I. G. Kaplan, *Intermolecular Interactions* (John Wiley & Sons, England, 2006).
- [32] V. Blum *et al.*, *Comput. Phys. Commun.* (to be published).
- [33] J. F. Dobson, A. White, and A. Rubio, *Phys. Rev. Lett.* **96**, 073201 (2006).
- [34] A. Bondi, *J. Phys. Chem.* **68**, 441 (1964).
- [35] P. Jurecka, J. Sponer, J. Cerny, and P. Hobza, *Phys. Chem. Chem. Phys.* **8**, 1985 (2006).
- [36] The determined free-atom vdW radii are 1.64, 1.90, 1.77, and 1.66 Å for H, C, N, and O atoms, respectively. The s_R coefficient for PBE is 0.94.