

TOPICAL REVIEW

Visual pathways and psychophysical channels in the primate

Barry B. Lee

SUNY College of Optometry, New York, NY, USA and Max Planck Institute for Biophysical Chemistry, Göttingen, Germany

The main cell systems of the retina that provide input to the striate cortex are now well described, although certain aspects of their anatomy and physiology remain contentious. Under simple stimulus conditions and in a threshold context psychophysical performance can often be assigned to one or other of these systems, and an identification of psychophysical channels with afferent pathways is justifiable. However, results from psychophysical studies using more complex stimulus conditions are more difficult to relate to ‘front end’ channels, and it is more difficult to separate the physiological contributions of afferent pathways from those of cortical mechanisms, in particular the separation of dorsal and ventral streams.

(Received 5 May 2010; accepted after revision 16 August 2010; first published online 19 August 2010)

Corresponding author B. B. Lee: SUNY Optometry, 33 W. 42nd St, New York, NY 10036, USA. Email: blee@sunyopt.edu

Introduction

General features of the anatomy and physiology of the visual pathways of the primate have become well established in past decades (see Lee & Dacey, 1997; Dacey, 1999, 2000; Lee, 2004 for reviews). This overview first briefly summarises standard views of the retinal anatomy of these afferent pathways, and some of their physiological properties. It mentions issues that are still unclear. In the second section these pathways are discussed in a psychophysical context. There is much classical evidence for the presence of psychophysical channels for colour and luminance under threshold conditions, which may map closely onto the different pathways originating in the retina. However, psychophysical performance in more complex visual contexts, such as natural scenes, now receives much attention and in such contexts these pathways relation to function is much less clear; it may be difficult to distinguish between properties of afferent pathways and those of central mechanisms.

Basic retinal structure in the primate


Figure 1 summarises Old-World primate retinal anatomy; the standard pattern of connectivity of receptor to bipolar cell to ganglion cell within the eye is outlined in Fig. 1A, and in the Nissl-stained section in Fig. 1B the laminar retinal structure is evident. Figure 1C provides a cartoon of retinal wiring, focusing on the three main cell systems that project to cortex through the lateral geniculate nucleus. A

more detailed description of primate retinal anatomy can be found elsewhere (Lee *et al.* 2010).

The cell systems providing the main input to the LGN are (from the top) the magnocellular (MC) pathway, which begins in parasol ganglion cells and projects to the striate cortex through the magnocellular layers of the lateral geniculate nucleus (LGN). There exist on- and off-centre cell types, each receiving input from various classes of diffuse bipolar. These diffuse bipolars almost entirely avoid input from short-wavelength (S) cones (Lee & Grünert, 2007). Input from medium- and long-wavelength (M and L) cones is thought to be derived at random from the underlying cone matrix. These cells have high achromatic contrast sensitivity, and very transient responses.

The second system is the parvocellular (PC) pathway beginning in the midget system. Midget ganglion cell

Barry Lee's main interest is the anatomy and physiology of primate retina and the relationship of retinal neurobiology to visual perception of colour and shape. After graduate work at University College in the UK and postdoctoral work at the University of Rochester, New York, he led a laboratory in the Department of Neurobiology at the Max Planck Institute for Biophysical Chemistry in Göttingen, concentrating on the primate visual system. Since 2000 he has been Professor of Biological Sciences at the College of Optometry, State University of New York.



anatomy strongly suggests that a single cone provides input to the centre through the one-to-one connectivity by which a single cone (in and near the fovea) provides input to a single midget bipolar cell which in turn contacts a single midget ganglion cell (Boycott & Dowling, 1969). Again, on- and off-centre varieties are present. Midget ganglion cells project to the PC layers of the LGN and thence to striate cortex. They display either M–L or L–M cone opponency, being excited by short and inhibited by long wavelengths or vice versa; they respond strongly to red–green chromatic modulation in a sustained manner and weakly to achromatic targets. In the sketch of Fig. 1C, these neurones are portrayed with red and green stripes; this reflects the fact that neighbouring ganglion cells may receive centre input from either an M- or L-cone and thus have opposite chromatic properties. This mixture of function within a single ganglion cell class of the same mosaic is unusual.

Lastly, S-cones provide input to several ganglion cell classes, of which the best described is the small bistratified cell (Dacey & Lee, 1994; Crook *et al.* 2009). Excitatory

(on) input is provided from S-cones by the S-cone bipolar and off input from the M- and L-cones mostly through diffuse bipolars, giving +blue–yellow (B–Y) responses. There are other cells with S-cone input, and at least one has inhibitory input from the S-cones. Their retinal connectivity remains uncertain but the class is well established from LGN recordings (Derrington *et al.* 1984; Valberg *et al.* 1986; Szmajda *et al.* 2006). These cell classes project to the koniocellular layers of the LGN (Tailby *et al.* 2008). Responses to blue–yellow modulation are strong and sustained with little response to achromatic modulation.

These three cell groups may relate to luminance and chromatic channels demonstrated psychophysically. How far this assumption is justified, is summarized in a subsequent section.

Issues of interest

The previous section presents a textbook view, but several unresolved issues have important functional implications.




Figure 1. A standard view of primate retinal structure

A, a view of the eye; the sketch of primate retinal wiring in C appears complex but is made up of standard retinal elements: receptors, bipolar and ganglion cells as indicated. The lamina structure of the primate retina is clearly delineated in B (scale bar 50 μm). C shows a sketch of primate retinal wiring. Each pathway conforms to the standard pattern (A), with receptors connected to bipolar cells to ganglion cells. The three main pathways sketched (parasol, midget and blue–yellow) are those providing the major input to striate cortex. Photomicrograph kindly provided by U. Grünert.

This section briefly points out some that have received recent attention.

Parasol ganglion cells. Dendritic trees cover a field six to eight cones across in the central retina; this is smaller than often assumed in the psychophysical literature. Measured centre sizes for midget and parasol ganglion cells are quite similar (Lee, 2004). Central parasol cells in the macaque are estimated to respond to 20–30 cycles deg^{-1} (cpd), close to the resolution limit for this species.

Parasol cells show a response minimum with equal luminance modulation (see below) but there is a residual response at twice the modulation frequency (Lee *et al.* 1989). This may be related to the effects of chromatic motion on luminance motion mechanisms (Cavanagh & Favreau, 1985; Mullen *et al.* 2003; Lee & Sun, 2009).

Recent recordings from arrays of parasol cells have indicated that there may be some degree of correlation of their activities (Shlens *et al.* 2006). Since parasol ganglion cells are implicated in the hyperacuity (Lee *et al.* 1994; Rüttiger *et al.* 2002), how these correlations may affect fine spatial processing is an interesting unresolved question.

Midget ganglion cells. The anatomy suggests that the centres of midget ganglion cells near the fovea are derived from a single cone. In most ganglion cells, centre diameter is defined by the spatial extent of the dendritic tree (Wässle *et al.* 1981), but midget ganglion cells are thus an exception to this rule. Measured receptive field centre sizes are much larger than expected on the basis of a single cone centre in the central retina. The reasons for this are probably partly optical (since the point spread function is larger than a single cone), but some measurements using interference fringes (McMahon *et al.* 2000) and adaptive optics (Sincich *et al.* 2009) have indicated a more complex picture. If this is so, current views of midget connectivity may have to be revised.

A second issue concerns the degree of specificity of connectivity in this pathway. Trichromatic colour vision is found only in primates among mammals. It evolved close to the beginning of primate evolution (Mollon, 1991). Although the blue–yellow system is phylogenetically ancient, the new red–green system presumably took over some existing cell class (Shapley & Perry, 1986; Wässle & Boycott, 1991). A centre derived from a single cone automatically confers cone specificity on the centre and some degree of red–green cone opponency, but whether or not surrounds are cone selective has generated much debate, with no clear conclusion (Lennie *et al.* 1991; Reid & Shapley, 1992, 2002). On evolution of this pathway, presumably surrounds were non-specific, but any selectivity would increase the strength of the chromatic signal (Lee, 2008), so some intermediate degree of specificity is plausible. A related question is cone specific

connectivity in the retinal periphery. Midget morphology changes in peripheral retina. Although the one-to-one connection between cones and midget bipolar cells is maintained to high eccentricity, convergence of midget bipolar cells onto midget ganglion cells becomes pronounced (Boycott & Dowling, 1969). Nevertheless, strongly cone opponent responses are maintained in a substantial percentage of cells in the mid-periphery (Martin *et al.* 2001; Solomon *et al.* 2005), implying some selectivity in this convergence; there is no known anatomical substrate for this.

The blue–yellow pathway. Physiology of the small bistratified cell is now well described (Crook *et al.* 2009). Other cell classes are less well defined; in particular the blue-off/yellow-on cell is less well understood (Dacey *et al.* 2002). There is also an additional blue-on/yellow-off class (Dacey & Packer, 2003), and the functional relationship between these two classes with apparently similar chromatic properties is obscure. Lastly, the photosensitive, melanopsin-containing ganglion cell also receives S-cone input (Dacey *et al.* 2005).

Visual pathways and channels

Beginning with Barlow and others (Barlow & Levick, 1969; Barlow, 1972), there have been attempts to rigorously link retinal physiology and psychophysical performance. In primate vision, there was early evidence for independent luminance and chromatic channels with different temporal properties (Kelly & Norren, 1977). A luminance channel was thought to underlie psychophysical performance on photometric tasks, such as heterochromatic flicker photometry (HFP) and the minimally distinct border (MDB). There is strong evidence that the MC pathway provides a physiological substrate for these tasks (Lee *et al.* 1988; Kaiser *et al.* 1990). The recent suggestion (Chatterjee & Callaway, 2002) that the MC pathway has significant S-cone input (which does not contribute to luminance) has been refuted (Sun *et al.* 2006). Also, psychophysical detection of luminance modulation (flicker) closely matches the properties of this pathway (Lee *et al.* 1990, 2007). On the other hand, detection of chromatic changes in a S-, M- and L-cone space can be partitioned into detection by S-cone or |M-L| cone mechanisms (Cole *et al.* 1993) and it is plausible that these map onto the S-cone-driven ganglion cells and the |M-L| cells of the PC pathway.

It is remarkable that psychophysical performance at threshold is constrained at a retinal level. There are a lot of synapses between the retina and a behavioural response, yet links between retinal physiology and psychophysics can be robust; the ‘nothing mucks it up’ principle (Teller, 1980).

However, there are a number of complications to this simple picture. Firstly, grating acuity (Pokorny *et al.* 1968) and the hyperacuities (Morgan & Aiba, 1985; Rüttiger & Lee, 2000) demonstrate a luminance-like spectral sensitivity. It has been proposed that the |M-L| cone opponent pathways does 'double duty' and, though underlying red-green vision at low spatial frequencies, at high spatial frequencies it underlies achromatic spatial vision (Ingling & Martinez-Uriegas, 1983; Lennie & D'Zmura, 1988). This view originated with the observations of Wiesel and Hubel (1966) that PC pathway cells in the LGN displayed centre-surround structure, and so large-field stimuli evoked |M-L| cone opponent responses while opponency is less with small targets. The alternative is that the MC pathway supports an achromatic channel of spatial vision, a view at first discounted since MC pathway cells were thought to have large centres and thus poor acuity (de Monasterio & Gouras, 1975). As noted above, however, recent data have shown that the difference in centre size is much smaller than first thought. This issue is beyond the scope of this brief review.

A peculiarity in the link between ganglion cell physiology and psychophysical performance is that chromatic pathways (S-cone and |M-L|) give responses to stimuli that are not perceived. Figure 2 shows three conditions for which this is the case. In Fig. 2A the temporal frequency tuning of |M-L| ganglion cells to red-green chromatic modulation is compared to psychophysical sensitivity to similar stimuli. Psychophysical sensitivity falls off above 4 Hz and above 10–12 Hz the chromatic

alternation cannot be seen. Yet ganglion cells respond vigorously to at least 30 Hz. A similar result holds for blue-yellow modulation and S-cone cells. Figure 2B shows another example; chromatic sensitivity to red-green modulation decreases rapidly with eccentricity, even with appropriately scaled stimuli, well before the one-to-one midget connectivity is lost. Midget ganglion cells at equivalent eccentricities have red-green chromatic sensitivity comparable to cells near the fovea. Lastly, grating visual resolution is higher with luminance gratings than with red-green equiluminant gratings (Mullen, 1985). However, resolution of PC pathways is similar to both grating types; independent of grating type, visual resolution is largely determined by centre size (Peichl & Wässle, 1979). Figure 2C is an example from the parafovea in which responses to both types of grating cut off near 10 cpd. This is much higher than parafoveal chromatic grating acuity. Again, it appears that central sites do not utilize a PC pathway signal. All these effects indicate that some aspects of the retinal signal are not utilized cortically, which substantially complicates the task of linking retinal physiology with psychophysical performance.

Distribution of function between the PC and MC pathways is often considered in terms of their spatial and temporal properties complementing one another to cover a broader range of frequencies than either alone (Kulikowski & Tolhurst, 1973; Silveira, 1996). Part of this distinction rests on the fact that PC pathway cells (and S-cone cells) show very sustained responses. This can provide information about brightness and chromaticity




Figure 2. Three examples in which retinal signals are not perceptually utilized

A, temporal frequency tuning for red-green modulation in psychophysics and PC pathway retinal ganglion cells, replotted from Lee *et al.* (2007 with permission from ARVO as the copyright holder). Psychophysical sensitivity (derived from Swanson *et al.* (1987)) and converted to cone contrast sensitivity. Physiological data are based on cell firing rates required to reach a response criterion. B, psychophysical and PC pathway sensitivity as a function of retinal eccentricity. Replotted from Martin *et al.* (2001). C, PC cell (+M-L cell, 4.8 deg eccentricity) responsivity to luminance (40% contrast) and red-green chromatic (100% modulation contrast) gratings (2 Hz drift rate) as a function of spatial frequency. Visual resolution is similar for both conditions.

of object surfaces; it is possible to derive psychological scaling of colour and brightness based on the activities of these neurons (Valberg & Seim, 1991). This differs somewhat from the textbook view of ganglion cell receptive fields as contrast or edge detectors, based on their centre-surround structure. A simple view of how PC and MC pathways deliver spatially distinct signals with complex patterns (in this case, colour Mondrians) was provided by Nothdurft and Lee (1982). Object boundaries derive from both luminance and colour, but shadows and shading tend to make chromatic properties more reliable indicators of material boundaries than luminance changes (Switkes *et al.* 1988; Kingdom *et al.* 2004). Algorithms for visual segmentation have found determination of surface characteristics (including chromatic properties) a useful approach (Li & Lennie, 2001; Maxwell *et al.* 2008).

There is now abundant evidence that colour in complex scenes provides a wealth of information about texture, structure and depth (see Shevell & Kingdom (2008) for a review from a psychophysical perspective). These data are sometimes difficult to relate to the apparent segregation of chromatic and luminance signals in the afferent pathways and in simple psychophysical detection tasks. Neurophysiologically, beginning in area 17 (Lennie *et al.* 1990), neurons are found which show chromatic properties intermediate between the characteristic chromatic signatures of the PC and KC pathways; also many neurons receive combined PC and MC input (Johnson *et al.* 2004). Given certain assumptions, linear combination of different pathways in cortex need not invalidate the 'nothing mucks it up' principle cited earlier, where psychophysical thresholds to simple stimuli map closely onto the properties of retinal neurons (or onto the cardinal axes of colour space; Krauskopf *et al.* 1982; Derrington *et al.* 1984). However, complex stimulus configurations appear to reveal 'higher order' chromatic mechanisms that might be generated at a cortical level (Krauskopf *et al.* 1986; Li & Lennie, 1997) but interpretation of these data is not straightforward (Eskew, 2009).

Nevertheless, complex spatial processing such as texture segmentation and pop-out can be based on chromatic cues; as pointed out by Shevell and Kingdom (2008), congruent (and incongruent) changes in luminance and chromaticity are an implicit aspect of normal viewing. For example, binocular vision, as a hyperacuity based on disparity, may be impaired with isoluminant patterns (Livingstone & Hubel, 1987), but binocular depth can readily be achieved with chromatic cues (Kingdom, 2003; Zaidi & Li, 2006). Various aspects of complex spatial processing may differentially rely on afferent pathways, but also psychophysical performance is likely to be heavily dependent on the separation of information in different cortical streams (see below). There have been numerous recent attempts to define cortical cells' properties in these complex contexts (e.g. Nothdurft *et al.* 1999; Roe *et al.*

2005), although links between physiology and behaviour tend to be less robust in the cortex than at earlier stages in the visual pathway.

Concluding remarks

The identification of psychophysical luminance and chromatic channels with afferent visual pathways may remain valid for simple detection tasks, but in more complex spatial contexts the way in which central pathways handle afferent input make such correlations less clear. In particular, dorsal (parietal) and ventral (temporal) processing streams with different functional properties receive differential input, with dorsal pathways dominated by the MC pathway and the ventral pathway receiving both luminance and chromatic input (Merigan & Maunsell, 1993). This ventral pathway processing must involve mechanisms that both assess surface chromatic properties of objects as well as using such information in spatial contexts; whether there are different mechanisms for these two functions, or whether they are two aspects of a unitary mechanism is obscure. Relating psychophysical performance to retinal mechanisms will be complicated by physiological properties of different cortical areas. For example, recent functional magnetic resonance imaging (fMRI) measurements suggest the V4 complex may have a sluggish temporal response (Liu & Wandell, 2005), perhaps related to the filtering of high temporal frequency chromatic information mentioned above. On the other hand, higher temporal frequency signals in the dorsal, motion systems appear better preserved. Lastly, the spectral sensitivities of chromatic mechanisms in detection tasks map closely onto that expected of retinal mechanism, but are inconsistent with psychophysical unique hues. This has been termed 'one of the central mysteries of colour science' (Mollon & Jordan, 1997) and is a further indication that cortical processing may modify afferent input in unexpected ways.

References

- Barlow HB (1972). Single units and sensation: a neuron doctrine for perceptual psychology? *Perception* **1**, 371–394.
- Barlow HB & Levick WR (1969). Three factors limiting the reliable detection of light by retinal ganglion cells of the cat. *J Physiol* **200**, 1–24.
- Boycott BB & Dowling JE (1969). Organization of the primate retina: light microscopy. *Philos Trans R Soc Lond B Biol Sci* **255**, 109–184.
- Cavanagh P & Favreau O (1985). Color and luminance share a common motion pathway. *Vision Res* **25**, 1595–1601.
- Chatterjee S & Callaway EM (2002). S cone contributions to the magnocellular visual pathway in macaque monkey. *Neuron* **35**, 1135–1146.

- Cole GR, Hine T & McIlhagga W (1993). Detection mechanisms in L-, M- and S-cone contrast space. *J Opt Soc Am A* **10**, 38–51.
- Crook JD, Davenport CM, Peterson BB, Packer OS, Detwiler PB & Dacey DM (2009). Parallel ON and OFF cone bipolar inputs establish spatially coextensive receptive field structure of blue-yellow ganglion cells in primate retina. *J Neurosci* **29**, 8372–8387.
- Dacey DM (1999). Primate retina: cell types, circuits and color opponency. *Prog Retin Eye Res* **18**, 737–763.
- Dacey DM (2000). Parallel pathways for spectral coding in primate retina. *Ann Rev Neurosci* **23**, 743–775.
- Dacey DM & Lee BB (1994). The blue-ON opponent pathway in primate retina originates from a distinct bistratified ganglion cell type. *Nature* **367**, 731–735.
- Dacey DM, Liao HW, Peterson BB, Robinson FR, Smith VC, Pokorny J, Yau KW & Gamlin PD (2005). Melanopsin-expressing ganglion cells in primate retina signal colour and irradiance and project to the LGN. *Nature* **433**, 749–754.
- Dacey DM & Packer OS (2003). Colour coding in primate retina: diverse cell types and cone specific circuitry. *Curr Opin Neurobiol* **13**, 421–427.
- Dacey DM, Peterson BB & Robinson FR (2002). Identification of an S-cone opponent OFF pathway in the macaque retina: morphology, physiology and possible circuitry. *Inv Ophthalmol Vis Sci* **43**, E-Abstract 2983.
- de Monasterio FM & Gouras P (1975). Functional properties of ganglion cells of the rhesus monkey retina. *J Physiol* **251**, 167–195.
- Derrington AM, Krauskopf J & Lennie P (1984). Chromatic mechanisms in lateral geniculate nucleus of macaque. *J Physiol* **357**, 241–265.
- Eskew RH (2009). Higher order color mechanisms: a critical review. *Vision Res* **49**, 2686–2704.
- Ingling CR & Martinez-Uriegas E (1983). The spatio-chromatic signal of the r-g channel. In *Colour Vision; Physiology and Psychophysics*, ed. Mollon J & Sharpe LT. Academic Press, London.
- Johnson EN, Hawken MJ & Shapley R (2004). Cone inputs in macaque primary visual cortex. *J Neurophysiol* **91**, 2501–2514.
- Kaiser PK, Lee BB, Martin PR & Valberg A (1990). The physiological basis of the minimally distinct border demonstrated in the ganglion cells of the macaque retina. *J Physiol* **422**, 153–183.
- Kelly DH & Norren Dv (1977). Two-band model of heterochromatic flicker. *J Opt Soc Am* **67**, 1081–1091.
- Kingdom FAA (2003). Color brings relief to human vision. *Nat Neurosci* **6**, 641–644.
- Kingdom FAA, Beauce C & Hunter L (2004). Colour vision brings clarity to shadows. *Perception* **33**, 907–914.
- Krauskopf J, Williams DR & Heeley DW (1982). Cardinal directions in color space. *Vision Res* **22**, 1123–1131.
- Krauskopf J, Williams DR, Mandler MB & Brown AM (1986). Higher order color mechanisms. *Vision Res* **26**, 23–32.
- Kulikowski JJ & Tolhurst DJ (1973). Psychophysical evidence for sustained and transient detectors in human vision. *J Physiol* **232**, 149–162.
- Lee BB (2004). Paths to color in the retina. *Clin Exptl Optom* **87**, 239–248.
- Lee BB (2008). Neural models and physiological reality. *Vis Neurosci* **25**, 231–241.
- Lee BB & Dacey DM (1997). Structure and function in primate retina. In *Color Vision Deficiencies XIII*, ed. Cavonius CR, pp. 107–118. Kluwer, Dordrecht, Holland.
- Lee BB, Martin PR & Grünert U (2010). Retinal connectivity and vision. *Progress in Retinal Research* In Press.
- Lee BB, Martin PR & Valberg A (1988). The physiological basis of heterochromatic flicker photometry demonstrated in the ganglion cells of the macaque retina. *J Physiol* **404**, 323–347.
- Lee BB, Martin PR & Valberg A (1989). Nonlinear summation of M- and L-cone inputs to phasic retinal ganglion cells of the macaque. *J Neurosci* **9**, 1433–1442.
- Lee BB, Pokorny J, Smith VC, Martin PR & Valberg A (1990). Luminance and chromatic modulation sensitivity of macaque ganglion cells and human observers. *J Opt Soc Am A* **7**, 2223–2236.
- Lee BB & Sun H (2009). The chromatic input to cells of the magnocellular pathway of primates. *J Vision* **9**, 151–158.
- Lee BB, Sun H & Zucchini W (2007). The temporal properties of the response of macaque ganglion cells and central mechanisms of flicker detection. *J Vision* **7**, 11–16.
- Lee BB, Wehrhahn C, Westheimer G & Kremers J (1994). The spatial precision of macaque ganglion cell responses and Vernier acuity. *Inv Ophthalmol Vis Sci Suppl* **35**, 2064.
- Lee SC & Grünert U (2007). Connections of diffuse bipolar cells in primate retina are biased against S-cones. *J Comp Neurol* **502**, 126–140.
- Lennie P & D’Zmura MD (1988). Mechanisms of color vision. <http://www.ncbi.nlm.nih.gov/pubmed/3048707> *Crit Rev Neurobiol* **3**, 333–400.
- Lennie P, Haake PW & Williams DR (1991). The design of chromatically opponent receptive fields. In *Computational Models of Visual Processing*, ed. Landy MS & Movshon JA, pp. 71–82. MIT Press, Cambridge, MA, USA.
- Lennie P, Krauskopf J & Sclar G (1990). Chromatic mechanisms in striate cortex of macaque. *J Neurosci* **10**, 649–669.
- Li A & Lennie P (1997). Mechanisms underlying segmentation of colored textures. *Vision Res* **37**, 83–97.
- Li A & Lennie P (2001). Importance of color in the segmentation of variegated surfaces. *J Opt Soc Am A* **18**, 1240–1251.
- Liu J & Wandell BA (2005). Specializations for chromatic and temporal signals in human visual cortex. *J Neurosci* **25**, 3459–3468.
- Livingstone MS & Hubel DH (1987). Psychophysical evidence for separate channels for the perception of form, color, motion and depth. *J Neurosci* **7**, 3416–3468.
- Martin PR, Lee BB, White AJ, Solomon SG & Rüttiger L (2001). Chromatic sensitivity of ganglion cells in peripheral primate retina. *Nature* **410**, 933–936.
- McMahon MJ, Lankheet MJ, Lennie P & Williams DR (2000). Fine structure of parvocellular receptive fields in the primate fovea revealed by laser interferometry. *J Neurosci* **20**, 2043–2053.

- Maxwell BA, Friedhoff RM & Smith CA (2008). A bi-illuminant dichromatic reflection model for understanding images. *IEEE Conference on Computer Vision and Pattern Recognition* 1–8.
- Merigan WH & Maunsell JHR (1993). How parallel are the primate visual pathways? *Ann Rev Neurosci* **16**, 369–402.
- Mollon JD (1991). Uses and evolutionary origins of primate color vision. In *Evolution of the Eye and Visual System*, ed. Cronly-Dillon JR & Gregory RL, pp. 306–319. MacMillan, London.
- Mollon JD & Jordan G (1997). On the nature of unique hues. In *John Dalton's Colour Vision Legacy*, ed. Dickinson C, Murray I & Carden D, pp. 381–392. Taylor & Francis, London, UK.
- Morgan MJ & Aiba TS (1985). Positional acuity with chromatic stimuli. *Vision Res* **25**, 689–695.
- Mullen KT (1985). The contrast sensitivity of human colour vision to red-green and blue-yellow chromatic gratings. *J Physiol* **359**, 381.
- Mullen KT, Yoshizawa T & Baker CL Jr (2003). Luminance mechanisms mediate the motion of red-green isoluminant gratings: the role of “temporal chromatic aberration”. *Vision Res* **43**, 1235–1247.
- Nothdurft HC, Gallant JL & Van Essen JC (1999). Response modulation by texture surround in primate area V1: correlates of “popout” under anesthesia. *Vis Neurosci* **16**, 15–34.
- Nothdurft HC & Lee BB (1982). Responses to coloured patterns in the macaque lateral geniculate nucleus; pattern processing in single neurones. *Exp Brain Res* **48**, 43–54.
- Peichl L & Wässle H (1979). Size, scatter and coverage of ganglion cell receptive field centres in the cat retina. *J Physiol* **291**, 117–141.
- Pokorny J, Graham CH & Lanson RN (1968). Effect of wavelength on foveal grating acuity. *J Opt Soc Am* **58**, 1410–1414.
- Reid RC & Shapley RM (1992). Spatial structure of cone inputs to receptive fields in primate lateral geniculate nucleus. *Nature* **356**, 716–718.
- Reid RC & Shapley RM (2002). Space and time maps of cone photoreceptor signals in macaque lateral geniculate nucleus. *J Neurosci* **22**, 6158–6175.
- Roe AW, Lu HD & Hung CP (2005). Cortical processing of a brightness illusion. *Proc Natl Acad Sci U S A* **102**, 3869–3874.
- Rüttiger L & Lee BB (2000). Chromatic and luminance contributions to a hyperacuity task. *Vision Res* **40**, 817–832.
- Rüttiger L, Lee BB & Sun H (2002). Transient cells can be neurometrically sustained; the positional accuracy of retinal signals to moving targets. *J Vision* **2**, 232–242.
- Shapley R & Perry VH (1986). Cat and monkey retinal ganglion cells and their visual functional roles. *Trends Neurosci* **9**, 229–235.
- Shevell SK & Kingdom FAA (2008). Color in complex scenes. *Ann Rev Psychol* **59**, 143–166.
- Shlens J, Field GD, Gauthier JL, Grivich MI, Petrusca D, Sher A, Litke AM & Chichilnisky EJ (2006). The structure of multi-neuron firing patterns in primate retina. *J Neurosci* **26**, 8254–8266.
- Silveira LCL (1996). Joint entropy loci of M and P cells: a hypothesis for parallel processing in the primate visual system. *Braz J Biol* **56**, 345–367.
- Sincich LC, Zhang Y, Tiruveedhula P, Horton JC & Roorda A (2009). Resolving single cone inputs to visual receptive fields. *Nat Neurosci* **12**, 967–969.
- Solomon SG, Lee BB, White AJ, Rüttiger L & Martin PR (2005). Chromatic organization of ganglion cell receptive fields in the peripheral retina. *J Neurosci* **25**, 4527–4539.
- Sun H, Smithson H, Zaidi Q & Lee BB (2006). Specificity of cone inputs to macaque ganglion cells. *J Neurophysiol* **95**, 837–849.
- Swanson WH, Ueno T, Smith VC & Pokorny J (1987). Temporal modulation sensitivity and pulse detection thresholds for chromatic and luminance perturbations. *J Opt Soc Am A* **4**, 1992–2005.
- Switkes E, Bradley A & DeValois KK (1988). Contrast dependence and mechanisms of making interactions among chromatic and luminance gratings. *J Opt Soc Am A* **5**, 1149–1162.
- Szmajda BA, Buzas P, Fitzgibbon T & Martin PR (2006). Geniculocortical relay of blue-off signals in the primate visual system. *Proc Natl Acad Sci U S A* **103**, 19512–19517.
- Tailby C, Szmajda BA, Buzas P, Lee BB & Martin PR (2008). Transmission of blue (S) cone signals through the primate lateral geniculate nucleus. *J Physiol* **586**, 5947–5967.
- Teller DY (1980). Locus questions in visual science. In *Visual Coding and Adaptability*, ed. Harris CS, pp. 151–176. Lawrence Erlbaum Associates, Hillsdale, New Jersey.
- Valberg A, Lee BB & Tigwell DA (1986). Neurons with strong inhibitory S-cone inputs in the macaque lateral geniculate nucleus. *Vision Res* **26**, 1061–1064.
- Valberg A & Seim T (1991). On the physiological basis of higher color metrics. In *From Pigments to Perception*, ed. Valberg A & Lee BB, pp. 425–436. Plenum, New York and London.
- Wässle H & Boycott BB (1991). Functional architecture of the mammalian retina. *Physiol Rev* **71**, 447–480.
- Wässle H, Boycott BB & Illing R-B (1981). Morphology and mosaic of on- and off-beta-cells of the cat retina and some functional considerations. *Proc R Soc Lond B Biol Sci* **212**, 177–195.
- Wiesel T & Hubel DH (1966). Spatial and chromatic interactions in the lateral geniculate body of the rhesus monkey. *J Neurophysiol* **29**, 1115–1156.
- Zaidi Q & Li A (2006). Three-dimensional shape perception from chromatic orientation flows. *Vis Neurosci* **23**, 323–330.

Acknowledgements

I thank Christoph Nothdurft and Qasim Zaidi for discussion of some of these issues. Preparation of this paper was supported by National Institutes of Health Grant EY 13112.