

Transcranial Magnetic Stimulation Intensities in Cognitive Paradigms

Jakob A. Kaminski^{1,2,3}, Franziska M. Korb^{1,3,3}, Arno Villringer^{1,2}, Derek V. M. Ott^{1,4*}

1 Department of Cognitive Neurology, Max Planck Institute for Human Cognitive and Brain Sciences, Leipzig, Germany, **2** Faculty of Medicine, University of Leipzig, Leipzig, Germany, **3** Center for Cognitive Neuroscience, Duke University, Durham, North Carolina, United States of America, **4** Department of Neurology, University Clinic Leipzig, Leipzig, Germany

Abstract

Background: Transcranial magnetic stimulation (TMS) has become an important experimental tool for exploring the brain's functional anatomy. As TMS interferes with neural activity, the hypothetical function of the stimulated area can thus be tested. One unresolved methodological issue in TMS experiments is the question of how to adequately calibrate stimulation intensities. The motor threshold (MT) is often taken as a reference for individually adapted stimulation intensities in TMS experiments, even if they do not involve the motor system. The aim of the present study was to evaluate whether it is reasonable to adjust stimulation intensities in each subject to the individual MT if prefrontal regions are stimulated prior to the performance of a cognitive paradigm.

Methods and Findings: Repetitive TMS (rTMS) was applied prior to a working memory task, either at the 'fixed' intensity of 40% maximum stimulator output (MSO), or individually adapted at 90% of the subject's MT. Stimulation was applied to a target region in the left posterior middle frontal gyrus (pmFG), as indicated by a functional magnetic resonance imaging (fMRI) localizer acquired beforehand, or to a control site (vertex). Results show that MT predicted the effect size after stimulating subjects with the fixed intensity (i.e., subjects with a low MT showed a greater behavioral effect). Nevertheless, the individual adaptation of intensities did not lead to stable effects.

Conclusion: Therefore, we suggest assessing MT and account for it as a measure for general cortical TMS susceptibility, even if TMS is applied outside the motor domain.

Citation: Kaminski JA, Korb FM, Villringer A, Ott DVM (2011) Transcranial Magnetic Stimulation Intensities in Cognitive Paradigms. PLoS ONE 6(9): e24836. doi:10.1371/journal.pone.0024836

Editor: Joseph Najbauer, City of Hope National Medical Center and Beckman Research Institute, United States of America

Received: April 20, 2011; **Accepted:** August 22, 2011; **Published:** September 29, 2011

Copyright: © 2011 Kaminski et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This research was funded by the Max Planck Institute for Human Cognitive and Brain Sciences. JAK's work was additionally supported with two travel grants from the German National Academic Foundation. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: ott@cps.mpg.de

These authors contributed equally to this work.

Introduction

Transcranial magnetic stimulation (TMS) has become increasingly important in brain research and is widely used for noninvasive evaluation and modulation of cortical function. Especially in neuropsychology, TMS gives insight into the function of a circumscribed brain area [1]. The temporary manipulation of neural activity impedes subjects' performance, as measured in prolonged reaction time (RT) and/or higher error rates, even beyond the duration of the stimulation [2]. Behavioral improvements have been reported after the application of certain TMS protocols as well [3]. It therefore allows the establishment of a causal relationship between structure and function. Nevertheless, individuals respond very differently to magnetic stimulation, and it is still unknown how a comparable biological TMS effect can be induced across subjects and how TMS intensity should be ideally gauged. Several factors have been shown to influence subjects' sensitivity to the magnetic pulse like, for example, the individual anatomy with varying distance from the coil to the underlying cortex [4]. Additionally, the trajectories of white matter fibers [5]

seem to have an impact on the TMS effect. Moreover, excitability also seems to depend on neuromodulator balance [6] and the functional state of the target area during stimulation [7].

From a technical point of view, the impact of TMS on a cluster of neurons depends on coil type, the geometry and direction of the induced electric field, which is determined by coil orientation [8–10], as well as on the pulse waveform defined by the stimulator model [11]. The assessment of these parameters was predominantly performed in the motor cortex; here, the responsiveness of cortical neurons can be measured in an objective way by registering observable muscle twitches or electromyographically monitored motor-evoked potentials (MEPs), often recorded from the first dorsal interosseus (FDI) muscle of the contralateral hand. The stimulator output which is needed to produce a reliable electromyographic response is commonly defined as the motor threshold (MT) [12,13]. Likewise, cortical excitability can be measured in the visual system by evoking phosphene [14], brief visual phenomena that are not caused by retinal stimulation. As with the MT, the cortical susceptibility to TMS in the occipital lobe can be determined by measuring the phosphene threshold

(PT), with the PT then being the minimum stimulator output necessary for a reliable induction of phosphenes. There are, however, some important shortcomings in the assessment of the PT as compared to MT, because the investigator has to rely on subjective reports, meaning that the effect cannot be measured in an objective way. Additionally, subjects need to be trained to recognize phosphenes and not everyone can perceive them.

In summary, cortical excitability can be measured in both the primary motor and visual cortex. But can regional excitability as expressed in MT and PT be used as a reference for stimulation in non-primary cortical areas? In other words, is there a global excitability level for each individual? Is it therefore valid to calibrate intensities according to those measures if other brain areas are stimulated? In order to elucidate these questions, several studies focused on whether and how MT and PT might correlate. Stewart et al. [15], amongst others [16–18], claim that there is no significant correlation between MT and PT. As a consequence, it would not be plausible that the excitability of any other brain region should relate to one of the two thresholds. Hence, adjusting stimulator output according to motor threshold would not reduce the variability of the TMS effect. Under this assumption, a predetermined, invariable intensity across all subjects has been applied in various studies [19–26]. On the other hand, one recent study by Deblieck et al. [27] does report a correlation between MT and PT and claimed that methodological shortcomings in the above-mentioned studies might be responsible for the discrepancies. If this was the case, stimulation intensity should be adjusted according to individual MT in order to reduce variability in the effect size, like it has also been practiced previously [28–35]. At the moment, the problem of generalizability of thresholds can be regarded as being unsolved and it appears to be more a matter of tradition and intuition whether experimental TMS in other cortical areas is referenced to motor (or phosphene) threshold or not.

Here, we address the question of how to calibrate stimulation intensity to prefrontal areas when performing cognitive experiments, in this case an n-back working memory task. Working memory is frequently defined as a system for the temporary storage and manipulation of remembered information [36]. The well-established n-back task requires this kind of cognitive processing and is known to evoke pronounced activation in the dorsolateral prefrontal cortex (DLPFC) or, more precisely, in the posterior part of the left middle frontal gyrus (pMFG; for a review see [37,38]). This is in line with cytoarchitectural and functional studies that suggest an important role of the posterior part of the middle frontal gyrus (Brodmann Area 9/46) in working memory processes [39], and due to its location near the cortical surface there is good access to this area using TMS. We located the pMFG as the TMS target region of interest (ROI) in individual functional magnetic resonance imaging (fMRI) maps of 2-back task activation. Because it has been shown that TMS affects local field potentials (LFPs) and BOLD-responses [40,41], we included a supplementary fMRI scanning session with a subgroup one week after the TMS experiments had ended, as an additional validation of consistency of the localizer, to ensure that the task-related BOLD-response in the target region had not shifted or decreased significantly. The TMS experiments consisted of four separate sessions, in which we applied repetitive TMS (rTMS) with either a predetermined ‘fixed’ intensity, identical for all subjects, or an individually adapted MT-related intensity prior to the working memory task. Both stimulation intensities were also applied to a control region not related to the task (vertex) in order to control for unspecific TMS effects.

If the excitability of the prefrontal cortex relates to the motor system’s excitability, we expect most consistent results (i.e., lowest

between-subject variability of the TMS-effect) in sessions with individually adapted intensities; at the same time, subjects should show differential effects in response to the pre-determined, fixed intensity, depending on whether they receive a relatively high or low stimulation intensity as compared to their individual sensitivity. In contrast, if there is no relationship between excitabilities, effects should be most pronounced when subjects are stimulated with the highest maximum stimulator output (MSO) (which we apply to the subjects with the highest MT in the individually adapted stimulation sessions); the pre-determined MSO, however, should then result in rather comparable effects across subjects.

Materials and Methods

Ethics Statement

All subjects gave their written informed consent, corresponding to the Code of Ethics of the World Medical Association (Declaration of Helsinki, sixth revision in 2008). The study was approved by the ethics committee of the University of Leipzig.

Subjects

Fifteen healthy subjects (mean age: 24.7 years, SD: 2.8 years) participated in the study. In order to reduce inter-subject variability, we chose only right-handed male participants (handedness was assessed with the Edinburgh inventory [42]) which are known to be a homogenous and stable group concerning their cortical excitability [43]. They had no history of neurological or psychiatric illness or knew of any cases of epilepsy amongst first-degree relatives. This was assessed with a questionnaire similar to the one established by Keel and others [44].

Experimental Design

All sessions (training, fMRI and four TMS sessions; see Figure 1) were one week apart. In the first week, subjects trained the n-back task in order to reduce training effects during the subsequent sessions. In the second week, the task was performed in an fMRI environment in order to acquire an anatomical and functional localizer for the target region, i.e. the activation peak in the left pMFG. During the next four weeks, we conducted the TMS sessions consisting of off-line stimulation followed by the task.

Overall, subjects were stimulated twice over the target region (pMFG) and twice over the control region (vertex), each time with either a predetermined fixed intensity (i.e., same for all participants) or an individually adapted intensity. The order of stimulation intensity and target site in the four sessions was pseudo-randomized and counterbalanced across participants. To assess stability of the functional localizer after this relatively high number of n-back task runs, a subgroup of six subjects underwent another fMRI scan in order to verify that the activation of the pMFG had not shifted or notably decreased.

Behavioral Task

For the n-back task, a series of letters had to be monitored with the instruction to respond whenever a letter appeared that was identical to the one presented n before (see Figure 2). Usually, n is a predefined integer as 1, 2 or 3. While a 1-back task does not reliably engage working memory, the interpretability of the results of a 3-back task are questioned because the performance decreases extremely and, additionally, there is a supposed capacity constraint [45]. Therefore we used a 2-back condition which still requires constant working memory updating and online-monitoring of the stored information. The so-called 0-back task served as standard experimental baseline condition (e.g., [46]). This is a simple target

Session type	MT measurement	Training n-back	fMRI for localization	rTMS	rTMS	rTMS	rTMS	fMRI consistency check
Stimulation site				pMFG (target)	pMFG (target)	vertex (control)	vertex (control)	
Intensity				Individual	Fixed	Individual	Fixed	




Figure 1. Overview of the experimental procedure. Note that the order of stimulation site and intensity is exemplary for one subject. The second fMRI experiment was only conducted with a subset of the sample (6 out of 15). doi:10.1371/journal.pone.0024836.g001

detection test, which does not need updating of information within working memory because subjects are required to respond whenever a predetermined, i.e., instructed, target letter occurs.

Each experimental block consisted of 25 stimuli out of which six were targets and 19 were non-target letters. In order to control for the possible strategy to simply respond to repeating stimuli in any of the conditions, two of the 19 non-target letters served as catch-trials (two 2-backs in the 0-back condition, one 1-back and one 3-back in the 2-back-condition). Each weekly session consisted of five blocks per condition (presented alternately and randomly beginning with either condition), so that 60 correct responses could optimally be obtained ($\frac{6 \text{ targets}}{\text{block}} \times 2[0\text{-back} + 2\text{-back}] \times 5\text{blocks} = 60 \text{ targets}$). In order to maximize a left lateralization of the activation, we used verbal stimuli (20 consonant letters; cf. [37]). Stimuli occurred in a pseudo-randomized order, with a stimulus duration of 500 ms at an inter-stimulus interval of 3000 ms. As a measure of accuracy, the

discrimination rate P_r was calculated. P_r is the difference between hit rate (i.e., correctly detected targets as indicated by a button press) and false alarm rate (i.e., erroneous responses to catch and foil trials), which is a recommended accuracy measure in target recognition tasks like the one described here [47].

fMRI-Localizer

For individual functional localization, subjects performed the paradigm in a 3T scanner (Medspec S300, Bruker, Ettlingen, Germany). Twenty axial slices (19.2 cm field of view, 64×64 matrix, 4 mm thickness, 1 mm spacing) parallel to the AC-PC plane and covering the whole brain were acquired using a single shot, gradient recalled echo-planar imaging (EPI) sequence (TR 2 s, TE 30 ms, 90° flip angle).

Preprocessing and Co-Registration. Data processing was performed using the software package LIPSIA [48], which contains




Figure 2. Exemplary extract of experimental block n-back conditions. Participants were instructed to respond to the stimuli labeled as 'target' with a right-hand index finger button press. doi:10.1371/journal.pone.0024836.g002

tools for preprocessing, co-registration, statistical evaluation, and visualization of fMRI data. Distortion correction was performed using a fieldmap scan that was acquired prior to the functional scan. Data were then corrected for motion using a matching metric, based on linear correlation. To correct for the temporal offset between the slices acquired in one scan, a cubic-spline-interpolation was applied.

To align the functional data slices with a 3D stereotactic coordinate reference system, a rigid linear registration with six degrees of freedom (three rotational, three translational) was performed. The rotational and translational parameters were determined on an optimal match between modified driven equilibrium Fourier transform (MDEFT; [49,50]), EPI-T1 scans that were run prior the functional data acquisition and an individual 3D reference data set, which had been measured for each subject during a previous scanning session. The high-resolution 3D reference data set with 160 slices and 1 mm slice thickness was standardized to the Talairach stereotactic space [51]. The rotational and translational parameters were subsequently transformed by linear scaling to standard space. The resulting parameters were then used to transform the functional slices using trilinear interpolation, so that the resulting functional slices were aligned with the stereotactic coordinate system, and functional data was re-sampled to $2 \times 2 \times 2$ mm. A temporal high-pass filter with a cut-off frequency at 1/100 Hz was used for baseline correction of the signal and a 3D spatial smoothing was applied with a Gaussian filter of 5.65 mm full width at half maximum (FWHM), which is equivalent to $\sigma = 1.2$.

Statistical Analysis. The statistical evaluation was based on a least-squares estimation using the general linear model for serially auto-correlated observations [52–55]. Data were modeled as mixed design including block-related information about the 0-back and 2-back conditions as well as pause/instruction periods between these task blocks, and event-related information about correctly detected 0-back and 2-back trials. The design matrix was generated with a synthetic hemodynamic response function [56,57] and its first derivative (for events only). The model equation, including the observation data, the design matrix and the error term, was convolved with a Gaussian kernel of dispersion of 4 s FWHM to deal with the temporal auto-correlation [55].

ROI Definition. In order to optimally determine the individual localization of the region of interest (ROI), i.e., the target of stimulation, 2-back vs. 0-back block-related activity were contrasted resulting in z-maps indicating activation differences between these conditions for each subject. The identification of the target region was based on functional and anatomical criteria (i.e., the highest z-value (Z_{target}) within the pMFG of the individual brain was classified as the ROI; see Table 1). For these single-subject data, no threshold or cluster size criteria were applied.

However, in order to compare our fMRI activation patterns with the existing ones in the n-back literature (for a review see: [37]), average statistical parametric maps based on the individual contrast images of the 2-back vs. the 0-back condition were computed.

The individual z-maps were converted to NIFTI-format and transferred to a PowerMac G5 (Apple Inc., Cupertino, USA) and used for localization with Brainsight TMS Software Version 1.7.8 (Rogue Research Inc., Montreal, Canada). The connected infrared stereo camera (Polaris Optical Tracking System, Northern Digital Inc., Waterloo, Canada) can detect both the coil and the participant by means of attached reflective trackers. This procedure provides the best precision in coil guidance [58] and furthermore allows reproduction of trajectories (i.e., the exact position and angle of the coil) in repeated sessions.

Table 1. Coordinates and Z value maximum of individual pMFG target regions.

Subject	Coordinates (mm)			Z_{target}
	x	y	z	
1	-34	8	38	1.61
2	-35	5	47	2.03
3	-35	4	52	2.09
4	-27	1	44	3.10
5	-45	33	38	1.60
6	-40	0	35	2.13
7	-46	38	17	3.21
8	-52	-1	34	3.41
9	-37	16	42	1.52
10	-36	-6	33	2.40
11	-34	13	37	1.32
12	-47	-4	44	1.99
13	-31	-2	55	1.96
14	-38	12	37	2.96
15	-38	14	32	2.33

Note that images were not normalized prior to fitting to the Talairach coordinate system provided by Brainsight; compatibility between subjects is therefore limited. Reference point for the origin of the underlying coordinate system is the anterior commissure (AC: 0 0 0) with the axial plane through anterior and posterior commissures (AC-PC) and the z-axis orthogonal to this plane. Values are like in the standard coordinate systems in mm with -x - left, +x - right, -y - posterior, +y - anterior, -z - inferior, +z - superior.
doi:10.1371/journal.pone.0024836.t001

TMS

Threshold Measurement. Stimulation was performed with a biphasic magstim Rapid² stimulator (The MAGSTIM Company Ltd, Whitland, UK) and a 70 mm figure-of-eight-shaped coil. Motor-threshold assessment followed recommendations by Rossini and Rothwell [12,13]: The coil was held tangentially to the skull at a 45° angle to the corresponding parasagittal line. MEPs were recorded from the right FDI using self-adhesive Ag-AgCl gel electrodes in a standard belly-tendon fashion. A Digitimer 360 (Digitimer Limited, Welwyn Garden City, UK) amplifier and the Spike Software Package version 4.02 (Cambridge Electronic Design Ltd, Cambridge, UK) were used to detect electromyographic responses. In order to find the MT hot-spot, we started with 55% MSO, which was sufficient in most participants to evoke MEPs; otherwise, intensity was increased in 5% steps. We started the stimulation 4 cm lateral and 1 cm anterior to the vertex. We systematically shifted the coil in steps of approximately 1 cm in anterior, posterior, lateral and medial direction, respectively, and repeated repositioning if higher MEPs were elicited from the new position. If coil replacement did not result in higher MEPs, we reduced the power at first in two then in one percent MSO decrements.

MT was both quantified at rest (rMT) as well as during activation (aMT). The rMT was defined at the percentage of MSO needed to evoke 50 μ V MEPs peak to peak in five out of ten consecutive trials. The aMT was assessed while the subjects had to contract the FDI with 20% of maximum force, which was measured with an analog dynamometer. The aMT was defined as the minimum MSO that would elicit MEPs exceeding contraction activity by 200 μ V in at least five out of ten consecutive stimulations. The MTs were measured in independent sessions prior to the actual experiment.

rTMS. Using an air-cooled figure-of-eight coil, we applied a continuous 1 Hz rTMS stimulation protocol for 15 min (total of 900 pulses) prior to task performance. The coil was held tangentially to the skull with the handle pointing upwards, and position was adjusted so that the electric current in the center of the coil would run perpendicular to the course of the inferior frontal sulcus [9,59]. As control site, the vertex was stimulated with the induced current running from posterior to anterior along the interhemispheric fissure. Stimulation intensity was adjusted to 90% of the individual rMT or set at a fixed intensity of 40% MSO. Ideally a fixed intensity should be set to the group average MT. However, the fixed intensity was chosen at 40% MSO because it corresponded to the rMT of the subject with the lowest thresholds and could thus be considered safe for all participants [60].

Scalp-Cortex Distance Measurement. MT has been shown to be influenced by the distance between the scalp and the cortex within the primary motor cortex (M1). Scalp-cortex distance (SCD) has been claimed to account for 25% [61], 29% [62] or even 45% [4] of interindividual MT variance. Therefore, we measured the SCD using the built-inBrainsight TMS Software metering rule in the target stimulation area and M1. In the high-resolution 3D-data set, we measured the distance from the closest voxel within the scalp surface to the target cortical site (pMFG) in all three axes in mm [62]. As we did not use neuronavigation in MT assessment, we measured the SCD in the neuroanatomical location of the hand-area (M1) above the hand-knob.

Results

Motor Threshold

rMT varied between 40 and 72% of MSO (mean rMT = 52.8; SD = 9.4), whereas aMT showed a variance between 37 and 68% of MSO (mean rMT = 47.4; SD = 8.0) (see Table 2). A two-tailed Pearson-test revealed a significant correlation of rMT and aMT ($r = 0.893$; $p < 0.01$).

Table 2. Motor thresholds and SCD.

Subject	aMT	rMT	SCD in mm	
			pMFG	M1
1	58	67	9	12.1
2	45	48	13.1	17.3
3	50	50	16.3	18.9
4	40	44	9.7	10
5	43	48	12.2	11.4
6	45	53	8.9	10.7
7	54	70	11.3	14.2
8	51	55	14	18.5
9	37	49	9	14.1
10	42	44	10	11.6
11	39	40	9.5	11.9
12	47	50	14.3	15
13	48	50	11.3	13.6
14	44	48	10.5	19.1
15	68	72	14.6	18.6

Motor threshold is expressed as % of MSO; note that SCD in pMFG was measured at the target stimulation site and the SCD in the M1 was measured on the anatomical area of the representation of the hand (handknob).

doi:10.1371/journal.pone.0024836.t002

Scalp-Cortex Distances

A one-tailed Pearson's correlation showed that SCD in the pMFG and the hand-knob did correlate ($r = 0.717$; $p < 0.05$). In M1, a one-tailed Pearson's correlation of SCD with aMT revealed a non-significant trend ($r = 0.431$; $p = 0.055$), whereas rMT did not show any significant correlation with the SCD in the hand area ($r = 0.264$; $p = 0.171$) (see Table 2).

fMRI Localizer

In the fMRI-contrasts (2-back vs. 0-back), we found activation in the posterior part of the left middle frontal gyrus (pMFG) in all participants. In the single subject contrasts, the z-value-threshold was gradually lowered until one blob within the pMFG with only one local maximum could be identified unambiguously. The peak voxel of this blob was then marked as the target stimulation site. Activation strength of the target, as indicated by the z-value of the local maximum (Z_{target}), varied between subjects (range: 1.32–3.41; mean 2.24; see Table 1) and was therefore included as regressor in later analysis in order to rule out an influence on effects of the stimulation.

The averaged contrast images of the 2-back vs. the 0-back condition revealed significant activation in the pMFG (Peak voxel Talairach coordinates $-45\ 26\ 24$, threshold at $z > 2.58$, i.e., $p < 0.005$, see Figure 3A). For the subsample of six participants undergoing a second post-TMS fMRI scanning session, the same average z-map was computed. Subsequently, a conjunction analysis of these subjects' pre and post TMS 2-back related activity was conducted, resulting in the same activation pattern in the pMFG (see Figure 3B).

Behavioral Data

General Effects. Calculating paired sample t-tests (2-back vs. 0-back), we found a significantly different performance (P_r -values) after stimulating the DLPFC both with a single intensity ($P_r = 99.8$ vs. 97.4; $t_{(14)} = 4.150$, $p < 0.001$) and with an individual intensity ($P_r = 99.9$ vs. 97.8; $t_{(14)} = 2.989$, $p < 0.01$). Performance in the 2-back task did not differ significantly from 0-back performance in the training or scanning sessions (see Figure 4).

The mean RT included only trials that were defined as targets (i.e., 0- and 2-back trials) in the respective block and were detected correctly by subjects. There was a significant difference between the two conditions, indicating slower RTs on 2-back trials (454.6 ms vs. 379.4 ms; $t_{(14)} = -2.812$, $p < 0.05$) in the very first session (i.e., the training session). All other sessions did not show any significant difference in RTs between conditions.

TMS-Specific Results. A 2×2 repeated measures ANOVA with the factors stimulation site (pMFG vs. vertex) and n-back condition (2-back vs. 0-back) was calculated separately for the sessions with a single intensity and the sessions with an individual intensity. Stimulating with a single intensity did not show any systematic effects, i.e. P_r and RT did not differ significantly between the target and control sessions.

In order to test whether the fixed intensity had differentiated effects on participants' performance based on their individual MT, we introduced a covariate which expresses 40% MSO as a percentage of individual MT ($\frac{40\%MSO}{rMT} \times 100$). As a consequence, we found a significant interaction between stimulation site and n-back condition ($F_{(1,13)} = 5.311$, $p < 0.05$) concerning task performance (P_r -values; see Figure 5A). In a subsequent post-hoc analysis, we correlated the rMT with the decrease in task attainment. The rMT correlated with 2-back performance ($r_{Pearson} = 0.488$; $p < 0.05$), indicating that a lower rMT lead to a larger decrease in task performance (see Figure 5B), whereas the 0-back condition did not show a significant correlation.




Figure 3. fMRI Results. (A) Averaged ($n = 15$) block-related 2-back vs. 0-back activity resulting in pMFG activation (peak voxel Talairach coordinates -45 26 24). Displayed are only voxels with $z > 2.58$ (i.e., $p < 0.005$). (B) fMRI check for consistency. Conjunction of averaged ($n = 6$) 2-back vs. 0-back block-related activity derived from pre- and post-TMS scans (peak voxel Talairach coordinates -45 26 24). doi:10.1371/journal.pone.0024836.g003

As rMT and aMT were highly correlated, we performed the same analysis with aMT as covariate. Again, the ANOVA revealed a significant interaction between stimulation site and n-back condition ($F_{(1,13)} = 5.004$, $p < 0.05$), with a corresponding correlation ($r_{\text{Pearson}} = 0.487$; $p < 0.05$; see Figure 5C).

When calculating the 2×2 ANOVA for the individually adapted stimulation intensity, no significant interaction was found. However, to test the possibility that aMT might be the better reference for overall excitability, we recalculated post-hoc the individual stimulation intensity applied (90% rMT), so it would be expressed as a percentage of aMT ($\frac{90\% \text{ rMT}}{\text{aMT}} \times 100$). As a consequence, we found a non-TMS-specific effect in the RT difference between 2- and 0-back condition ($F_{(1,13)} = 6.981$, $p < 0.05$) (see Figure 6A). The decrease in RT n-back effect correlated with the stimulation intensity relative to the aMT ($= \frac{90\% \text{ rMT}}{\text{aMT}} \times 100$) ($r_{\text{Pearson}} = 0.583$; $p < 0.05$) (see Figure 6B). Further analysis revealed that neither the

0-back nor the 2-back block alone accounted for this, but rather the interaction between an increase in 2-back RTs and a decrease of 0-back RTs (see Figure 6C). Neither SCD nor Z_{target} explained any variance when included as covariates to the ANOVA.

Discussion

The Problem

When applied in non-motor regions for neuropsychological studies, the calibration of intensity is one of the major problems in TMS studies [63]. This is due to the high interindividual variance in cortical sensitivity and the uncertain intraindividual transferability of measurable thresholds. Ideally, the effect of the stimulation should be comparable between subjects. In order to achieve small intersubject variance, individually adapted stimulation intensity is desirable. Some researchers try to achieve this by




Figure 4. General effects. Task performance on the two experimental conditions (2-back vs. 0-back) as measured by P_r . Paired-sample t-tests only revealed a significant difference when subjects were stimulated in the target region both with a single intensity (single int.: $P_r = 99.8$ vs. 97.4 ; $t_{(14)} = 4.150$, $p < 0.001$) and with an individually adapted intensity (individual int.: $P_r = 99.9$ vs. 97.8 ; $t_{(14)} = 2.989$, $p < 0.01$). doi:10.1371/journal.pone.0024836.g004




Figure 5. TMS-specific results. (A) Interaction between stimulation site and block type after stimulation with a fixed intensity (40% MSO). A sufficient level of statistical significance of that interaction was only reached when the repeated measures ANOVA was informed with the rMT as a covariate ($F_{(1,13)} = 5.311, p < 0.05$). (B) Correlation of resting motor threshold and costs in task accuracy, as revealed by the difference between control and target stimulation P_r value (rMT: $r_{Pearson} = 0.488; p < 0.05$). (C) Correlation of active motor threshold and costs in task accuracy, as revealed by the difference between control and target stimulation P_r value (aMT: $r_{Pearson} = 0.487; p < 0.05$). doi:10.1371/journal.pone.0024836.g005

adapting the stimulation intensities according to the MT. There are, however, good reasons to distrust MT as a valid measure for the sensitivity in non-motor cortical regions. As for any two given target areas, the cytoarchitecture of the cortex, the gyrification, the

homological structure, and the anatomy of the skull are different in prefrontal areas and the primary motor cortex. Therefore, the effect of TMS and the sensitivity to stimulation might be very different.




Figure 6. Non-specific influence of rTMS on task performance. (A) Statistically significant difference between 2-back and 0-back condition after stimulation with individually adapted intensity when informing the model with the aMT as a covariate, or the z-score of percent rMT in relation to aMT, respectively ($F_{(1,13)} = 6.981$, $p < 0.05$) (B) Correlation analysis indicating higher RT costs (i.e., 2-back minus 0-back RT) correlating with higher stimulation intensity in relation to aMT ($r_{\text{Pearson}} = 0.583$; $p < 0.05$) (C) The higher RT costs could not be attributed to an increase in 2-back RTs or decrease in 0-back RTs alone, but rather a combination of both. doi:10.1371/journal.pone.0024836.g006

Stimulation Intensities

Our results demonstrate that there is a link between cortical TMS excitability in the two regions examined (motor and prefrontal cortex). When all participants were stimulated with the same intensity (40% MSO), changes in performance (P_T) related to rMT: impact was strongest on subjects with a lower

rMT (e.g., 40% MSO), who were stimulated at comparatively high intensities ($\approx 100\%$ rMT); in contrast, for subjects with a high rMT (e.g., 72% MSO), the 40% MSO fixed intensity stimulation was far below their threshold and did not have a significant effect (see Figure 5B). Absolute stimulation intensity (in % MSO) alone had no significant influence either; otherwise, subjects with higher

thresholds should have shown greater deterioration in the sessions with individually adapted stimulation, because here they received the strongest stimulation of all subjects. However, if excitability of the non-motor cortices is related to rMT, why did individually adapted stimulation at 90% rMT yield no significant effects?

The most plausible explanation is that 90% rMT is not the most effective intensity for the present target site within the prefrontal cortex. Possibly, higher threshold-adapted intensities would yield stronger effects. However, it is known that higher stimulation intensities are not automatically more efficient [64] and the DLPFC might have a much higher or lower (TMS-)sensitivity than the motor cortex. Both an over- and an under-stimulation of any cortical area might lead to no, different, or opposite effects, depending on the neuronal populations predominately activated at any point. In principle, there might be any kind of non-linear yet stable relation between MT and the “DLPFC threshold”. Therefore, we cannot tell how to ideally adjust stimulation intensity. Based on our findings, we suggest that MT serves as a suitable measure of varying excitability between individuals. However, it does not provide an easy-to-use reference of how to stimulate any other part of cortex within the same brain, especially if basic features (cortical cytoarchitecture, etc.) differ.

After correcting individual stimulation intensity for aMT ($\frac{90\% \text{ of } rMT}{aMT} \times 100$), we found an impact of stimulation on response times over the whole period of the task, but surprisingly the effect was not specific for the stimulation site (pMFG vs. vertex). This indicates that the control stimulation to the vertex might not be the ideal option and possibly stimulates some midline cortical areas as the supplementary motor area, which could interfere with context-dependent motor planning. Choosing an adequate control site is yet another methodological issue that may be addressed in different ways: Given that stimulation to the DLPFC can result in disturbing jaw and facial muscle twitches, control stimulation should ideally elicit similar side effects, which is not true for either sham (tilted coil) or vertex stimulation [65]. Therefore, contralateral stimulation might be the best choice with regards to side-effects. But in the given working memory task, lateralization is often only partial, i.e., the contralateral site is involved in a cognitive process, and contralateral stimulation might have effects of its own. Therefore, we think that vertex stimulation is still the better choice, eliciting local somatosensory skin sensations, albeit not comparable to target stimulation.

Recommendation Based on Our Results

One of our main results is that there was an influence of individual MT on stimulation effects when all participants were stimulated at the same MSO, as we could show by a regression analysis. However, individually adapted stimulation yielded no significant TMS effects.

Based on these findings, it is not possible to say how the optimum DLPFC stimulation intensity would have to be adjusted. Therefore a whole range of intensities would have to be tested in order to evaluate a possible threshold in a specific brain area, which requires extremely meticulous and tedious studies, especially when involving cognitive processes. In most experiments, this would not be feasible if the determination of the threshold is not the overall aim of the study. Thus we suggest to use a fixed stimulation intensity across subjects, but include the individual MT in regression analyses, as an estimate of between-subject variability in cortical excitability. This idea would be in line with Deblicq et al. [27] who also suggested an intersubject comparability of different cortex areas concerning TMS excitability. Additionally, we could show that neither SCD nor differences in the fMRI data, depicted as Z_{target} , had a significant impact on the measured

TMS-Effects (i.e., change in P_r -value). Hence, we conclude that MT is the best marker for interindividual differences in TMS susceptibility.

TMS-Limitations

In any TMS experiment, one should consider sources of variance lying in the method itself: as discussed above TMS efficacy might be modulated to a variable degree by to different SCDs [61]. The fact that SCD in M1 and prefrontal regions correlate can be corrected e.g., by algorithms as the one introduced by Stokes et al. [62]. The adjustment of stimulation proposed in their publication, however, cannot take differences in tissue composition and conductivity into account [63]. In our subset of participants, we found a nonsignificant trend for the correlation of aMT and SCD in M1. However, as mentioned above, this finding could only partially explain a relationship between those two areas. The finding that rMT did not correlate with SCD in M1 might be due to the fact that rMT is not as precise as aMT as a measure of excitability.

Secondly, the induced effect is supposed to be a circumscribed cortical disturbance leading to measurable behavioral changes. It is, however, known that whole functional networks can be affected [25,66]. Possible long-distance effects are often ignored and some networks might prove functionally more resistant to TMS than others. Another interesting point to consider is the state dependency of TMS [7]. Adapted to our experimental design, one could speculate that any random cognitive operation performed by the subject during the stimulation period might possibly have an impact on stimulation efficiency, and thus subsequent behavioral changes, due to a different state of neuronal excitation.

Another finding noteworthy to be pointed out is the effect that rhythmic TMS patterns like the one applied here can have an effect on brain oscillations, some of which have been linked to cognitive tasks (for a review, see [67]). Although we cannot rule out that our TMS protocol caused such an entrainment of natural oscillations [68], we do not have to assume that it influenced the results. While the oscillations have been related to various TMS frequencies, regardless of intensity, we compared different intensities at the same frequency.

Concerning effects of the repetitive performance of the n-back task, a conjunction analysis was conducted for a subsample (i.e., $n=6$) for their pre- and post-TMS fMRI scans. We found precisely the same activation pattern in the pMFG that was evident in the averaged pre-TMS fMRI scans of the whole sample (i.e., $n=15$). Hence, we conclude that neither the excessive performance of the n-back task nor the repeated application of rTMS over a period of four weeks result in notably weaker recruitment of the same cortical network or the use of entirely different brain regions. Therefore, we may interpret all of the TMS effects as being exclusively attributable to the manipulation of stimulation intensity and site.

Conclusion

We found a link between cortical sensitivity in the motor cortex, and the left dorsolateral prefrontal cortex. However, the exact magnitude of this correlation remains unclear. As long as there is no strong hypothesis concerning the absolute excitability of the DLPFC, we suggest stimulating all subjects with the same stimulation intensity, and subsequently correcting for individual thresholds in a regression analysis. We also believe that this procedure is feasible for other non-primary cortical areas than the DLPFC. However, future research is necessary to prove if this is actually true.

Additionally, we showed the stability of an n-back fMRI-localizer over a total of six sessions, two of them including TMS to the peak of activation in the left pMFG; this consistency, which is crucial to any experiment including fMRI and TMS in different sessions, is implicitly assumed but, as to our knowledge, hadn't been shown previously.

Acknowledgments

We would like to thank Ramona Menger for her help with subject acquisition. We are equally grateful to Mandy Jochemko, Anke Kummer,

Annett Wiedemann and Simone Wipper for their assistance at the MRI scanner. We thank Rosie Wallis for her support with proof reading and two anonymous reviewers for their helpful comments.

Author Contributions

Conceived and designed the experiments: JAK FMK AV DVMO. Performed the experiments: JAK FMK DVMO. Analyzed the data: JAK FMK DVMO. Wrote the paper: JAK FMK AV DVMO.

References

- Walsh V, Rushworth M (1999) A primer of magnetic stimulation as a tool for neuropsychology. *Neuropsychologia* 37: 125–35.
- Pascual-Leone A, Walsh V, Rothwell J (2000) Transcranial magnetic stimulation in cognitive neuroscience—virtual lesion, chronometry, and functional connectivity. *Curr Opin Neurobiol* 10: 232–7.
- Guse B, Falkai P, Wobrock T (2010) Cognitive effects of high-frequency repetitive transcranial magnetic stimulation: a systematic review. *J Neural Transm* 117: 105–22.
- McConnell KA, Nahas Z, Shastri A, Lorberbaum JP, Kozel FA, et al. (2001) The transcranial magnetic stimulation motor threshold depends on the distance from coil to underlying cortex: a replication in healthy adults comparing two methods of assessing the distance to cortex. *Biol Psychiatry* 49: 454–9.
- Herbsman T, Forster L, Molnar C, Dougherty R, Christie D, et al. (2009) Motor threshold in transcranial magnetic stimulation: the impact of white matter fiber orientation and skull-to-cortex distance. *Hum Brain Mapp* 30: 2044–55.
- Ziemann U (2004) TMS and drugs. *Clin Neurophysiol* 115: 1717–29.
- Silvanto J, Muggleton N, Walsh V (2008) State-dependency in brain stimulation studies of perception and cognition. *Trends Cogn Sci* 12: 447–54.
- Amassian VE, Eberle L, Maccabee PJ, Cracco RQ (1992) Modelling magnetic coil excitation of human cerebral cortex with a peripheral nerve immersed in a brain-shaped volume conductor: the significance of fiber bending in excitation. *Electroencephalogr Clin Neurophysiol* 85: 291–301.
- Brasil-Neto JP, Cohen LG, Panizza M, Nilsson J, Roth BJ, et al. (1992) Optimal focal transcranial magnetic activation of the human motor cortex: effects of coil orientation, shape of the induced current pulse, and stimulus intensity. *J Clin Neurophysiol* 9: 132–6.
- Hill AC, Davey NJ, Kennard C (2000) Current orientation induced by magnetic stimulation influences a cognitive task. *Neuroreport* 11: 3257–3259.
- Kammer T, Beck S, Thielscher A, Laubis-Herrmann U, Topka H (2001) Motor thresholds in humans: a transcranial magnetic stimulation study comparing different pulse waveforms, current directions and stimulator types. *Clin Neurophysiol* 112: 250–8.
- Rothwell JC, Hallett M, Berardelli A, Eisen A, Rossini P, et al. (1999) Magnetic stimulation: motor evoked potentials. The International Federation of Clinical Neurophysiology. *Electroencephalogr Clin Neurophysiol Suppl* 52: 97–103.
- Rossini PM, Barker AT, Berardelli A, Caramia MD, Caruso G, et al. (1994) Non-invasive electrical and magnetic stimulation of the brain, spinal cord and roots: basic principles and procedures for routine clinical application. Report of an IFCN committee. *Electroencephalogr Clin Neurophysiol* 91: 79–92.
- Barker AT, Freeston IL, Jalilous R, Merton PA, Morton HB (1985) Magnetic stimulation of the human brain. *Journal of Physiology* 369(Suppl.): 3P.
- Stewart LM, Walsh V, Rothwell JC (2001) Motor and phosphene thresholds: a transcranial magnetic stimulation correlation study. *Neuropsychologia* 39: 415–9.
- Antal A, Nitsche MA, Kincses TZ, Lampe C, Paulus W (2004) No correlation between moving phosphene and motor thresholds: a transcranial magnetic stimulation study. *Neuroreport* 15: 297–302.
- Borojerd B, Meister IG, Foltys H, Sparing R, Cohen LG, et al. (2002) Visual and motor cortex excitability: a transcranial magnetic stimulation study. *Clin Neurophysiol* 113: 1501–4.
- Gervig M, Kastrup O, Meyer BU, Niehaus L (2003) Evaluation of cortical excitability by motor and phosphene thresholds in transcranial magnetic stimulation. *J Neurol Sci* 215: 75–8.
- Silvanto J, Cattaneo Z, Battelli L, Pascual-Leone A (2008) Baseline cortical excitability determines whether TMS disrupts or facilitates behavior. *J Neurophysiol* 99: 2725–30.
- Pitcher D, Walsh V, Yovel G, Duchaine B (2007) TMS evidence for the involvement of the right occipital face area in early face processing. *Curr Biol* 17: 1568–73.
- Buelte D, Meister IG, Staedtgen M, Dambek N, Sparing R, et al. (2008) The role of the anterior intraparietal sulcus in crossmodal processing of object features in humans: an rTMS study. *Brain Res* 1217: 110–8.
- Muggleton NG, Postma P, Moutsopoulou K, Nimmo-Smith I, Marcel A, et al. (2006) TMS over right posterior parietal cortex induces neglect in a scene-based frame of reference. *Neuropsychologia* 44: 1222–9.
- Beck DM, Muggleton N, Walsh V, Lavie N (2006) Right parietal cortex plays a critical role in change blindness. *Cereb Cortex* 16: 712–7.
- O'Shea J, Muggleton NG, Cowey A, Walsh V (2004) Timing of target discrimination in human frontal eye fields. *J Cogn Neurosci* 16: 1060–7.
- Ott DVM, Ullsperger M, Jocham G, Neumann J, Klein TA (2011) Continuous theta-burst stimulation (cTBS) over the lateral prefrontal cortex alters reinforcement learning bias. *Neuroimage* 57: 617–23.
- Dormal V, Andres M, Pesenti M (2011) Contribution of the right intraparietal sulcus to numerosity and length processing: an fMRI-guided TMS study. *Cortex*. doi:10.1016/j.cortex.2011.05.019.
- Deblieck C, Thompson B, Iacoboni M, Wu AD (2008) Correlation between motor and phosphene thresholds: a transcranial magnetic stimulation study. *Hum Brain Mapp* 29: 662–70.
- Rizzo S, Sandrini M, Papagno C (2007) The dorsolateral prefrontal cortex in idiom interpretation: an rTMS study. *Brain Res Bull* 71: 523–8.
- Mottaghy FM, Gangitano M, Sparing R, Krause BJ, Pascual-Leone A (2002) Segregation of areas related to visual working memory in the prefrontal cortex revealed by rTMS. *Cereb Cortex* 12: 369–75.
- Viggiano MP, Giovannelli F, Borgheresi A, Feurra M, Berardi N, et al. (2008) Disruption of the prefrontal cortex function by rTMS produces a category-specific enhancement of the reaction times during visual object identification. *Neuropsychologia* 46: 2725–31.
- Fitzgerald PB, Hoy K, Daskalakis ZJ, Kulkarni J (2009) A randomized trial of the anti-depressant effects of low- and high-frequency transcranial magnetic stimulation in treatment-resistant depression. *Depress Anxiety* 26: 229–34.
- Stern WM, Tormos JM, Press DZ, Pearlman C, Pascual-Leone A (2007) Antidepressant effects of high and low frequency repetitive transcranial magnetic stimulation to the dorsolateral prefrontal cortex: a double-blind, randomized, placebo-controlled trial. *J Neuropsychiatry Clin Neurosci* 19: 179–86.
- Dell'Osso B, Mundo E, D'Urso N, Pozzoli S, Buoli M, et al. (2009) Augmentative repetitive navigated transcranial magnetic stimulation (rTMS) in drug-resistant bipolar depression. *Bipolar Disord* 11: 76–81.
- Speer AM, Repella JD, Figueras S, Demian NK, Kimbrell TA, et al. (2001) Lack of adverse cognitive effects of 1 Hz and 20 Hz repetitive transcranial magnetic stimulation at 100% of motor threshold over left prefrontal cortex in depression. *J Ect* 17: 259–63.
- Uddin LQ, Molnar-Szakacs I, Zaidel E, Iacoboni M (2006) rTMS to the right inferior parietal lobule disrupts self-other discrimination. *Soc Cogn Affect Neurosci* 1: 65–71.
- Baddeley AD (1986) Working Memory. Oxford: Oxford University Press. pp 289.
- Owen AM, McMillan KM, Laird AR, Bullmore E (2005) N-back working memory paradigm: a meta-analysis of normative functional neuroimaging studies. *Hum Brain Mapp* 25: 46–59.
- Wager TD, Smith EE (2003) Neuroimaging studies of working memory: a meta-analysis. *Cogn Affect Behav Neurosci* 3: 255–74.
- Petrides M (2005) Lateral prefrontal cortex: architectonic and functional organization. *Philos Trans R Soc Lond B Biol Sci* 360: 781–95.
- Allen EA, Pasley BN, Duong T, Freeman RD (2007) Transcranial magnetic stimulation elicits coupled neural and hemodynamic consequences. *Science* 317: 1918–21.
- Sarfell AS, Diekhoff S, Wang LE, Liuzzi G, Uludağ K, et al. (2011) Convergence of human brain mapping tools: Neuronavigated TMS parameters and fMRI activity in the hand motor area. *Hum Brain Mapp*. doi: 10.1002/hbm.21272.
- Oldfield RC (1971) The assessment and analysis of handedness: the edinburgh inventory. *Neuropsychologia* 9: 97–113.
- Smith AE, Sale MV, Higgins RD, Wittert GA, Pitcher JB (2011) Male human motor cortex stimulus-response characteristics are not altered by aging. *J Appl Physiol* 110: 206–12.
- Keel JC, Smith MJ, Wassermann EM (2001) A safety screening questionnaire for transcranial magnetic stimulation. *Clin Neurophysiol* 112: 720.
- Callicott JH, Mattay VS, Bertolino A, Finn K, Coppola R, et al. (1999) Physiological characteristics of capacity constraints in working memory as revealed by functional MRI. *Cereb Cortex* 9: 20–6.
- Ragland JD, Turetsky BI, Gur RC, Gunning-Dixon F, Turner T, et al. (2002) Working memory for complex figures: an fMRI comparison of letter and fractal n-back tasks. *Neuropsychology* 16: 370–9.

47. Snodgrass JG, Corwin J (1988) Pragmatics of measuring recognition memory: applications to dementia and amnesia. *J Exp Psychol Gen* 117: 34–50.
48. Lohmann G, Muller K, Bosch V, Mentzel H, Hessler S, et al. (2001) LIPSIA – a new software system for the evaluation of functional magnetic resonance images of the human brain. *Comput Med Imaging Graph* 25: 449–57.
49. Ugurbil K, Garwood M, Ellermann J, Hendrich K, Hinke R, et al. (1993) Imaging at high magnetic fields: initial experiences at 4 T. *Magn Reson Q* 9: 259–77.
50. Norris DG (2000) Reduced power multislice MDEFT imaging. *J Magn Reson Imaging* 11: 445–51.
51. Talairach J, Tournoux P (1988) Co-planar stereotaxic atlas of the human brain. New York: Thieme. pp 122.
52. Friston K (1994) Statistical parametric mapping. San Diego: Academic Press. pp 79–93.
53. Friston K, Holmes A, Worsley K, Poline J, Frith C, et al. (1995) Statistical parametric maps in functional imaging: a general linear approach. *Hum Brain Mapping* 2: 189–210.
54. Friston KJ, Holmes AP, Poline JB, Grasby PJ, Williams SC, et al. (1995) Analysis of fMRI time-series revisited. *Neuroimage* 2: 45–53.
55. Worsley KJ, Friston KJ (1995) Analysis of fMRI time-series revisited—again. *Neuroimage* 2: 173–81.
56. Friston KJ, Fletcher P, Josephs O, Holmes A, Rugg MD, et al. (1998) Event-related fMRI: characterizing differential responses. *Neuroimage* 7: 30–40.
57. Josephs O, Turner R, Friston KJ (1997) Event-related fMRI. *Human Brain Mapping* 5: 243–248.
58. Sack AT, Kadosh RC, Schuhmann T, Moerel M, Walsh V, et al. (2009) Optimizing functional accuracy of TMS in cognitive studies: A comparison of methods. *J Cogn Neurosci* 21: 207–21.
59. Mills KR, Boniface SJ, Schubert M (1992) Magnetic brain stimulation with a double coil: the importance of coil orientation. *Electroencephalogr Clin Neurophysiol* 85: 17–21.
60. Wassermann EM (1998) Risk and safety of repetitive transcranial magnetic stimulation: report and suggested guidelines from the international workshop on the safety of repetitive transcranial magnetic stimulation, June 5–7, 1996. *Electroencephalogr Clin Neurophysiol* 108: 1–16.
61. Kozel FA, Nahas Z, deBrux C, Molloy M, Lorberbaum JP, et al. (2000) How coil-cortex distance relates to age, motor threshold, and antidepressant response to repetitive transcranial magnetic stimulation. *J Neuropsychiatry Clin Neurosci* 12: 376–84.
62. Stokes MG, Chambers CD, Gould IC, Henderson TR, Janko NE, et al. (2005) Simple metric for scaling motor threshold based on scalp-cortex distance: application to studies using transcranial magnetic stimulation. *J Neurophysiol* 94: 4520–4527.
63. Rossi S, Hallett M, Rossini PM, Pascual-Leone A, Safety of TMS Consensus Group (2009) Safety, ethical considerations, and application guidelines for the use of transcranial magnetic stimulation in clinical practice and research. *Clin Neurophysiol* 120: 2008–39.
64. Munchau A, Bloem BR, Irlbacher K, Trimble MR, Rothwell JC (2002) Functional connectivity of human premotor and motor cortex explored with repetitive transcranial magnetic stimulation. *J Neurosci* 22: 554–61.
65. Loo CK, Taylor JL, Gandevia SC, McDermont BN, Mitchell PB, et al. (2000) Transcranial magnetic stimulation (TMS) in controlled treatment studies: are some “sham” forms active? *Biol Psychiatry* 47: 325–31.
66. Knoch D, Gianotti LR, Pascual-Leone A, Treyer V, Regard M, et al. (2006) Disruption of right prefrontal cortex by low-frequency repetitive transcranial magnetic stimulation induces risk-taking behavior. *J Neurosci* 26: 6469–72.
67. Thut G, Miniussi C (2009) New insights into rhythmic brain activity from TMS-EEG studies. *Trends Cogn Sci* 13: 182–9.
68. Thut G, Veniero D, Romei V, Miniussi C, Schyns P, et al. (2011) Rhythmic TMS causes local entrainment of natural oscillatory signatures. *Curr Biol* 21: 1176–85.