

Classical GR as a topological theory with linear constraints

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2011 J. Phys.: Conf. Ser. 314 012044

(<http://iopscience.iop.org/1742-6596/314/1/012044>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 194.94.224.254

The article was downloaded on 15/11/2011 at 09:12

Please note that [terms and conditions apply](#).

Classical GR as a topological theory with linear constraints¹

Steffen Gielen

DAMTP, University of Cambridge, Centre for Mathematical Sciences, Wilberforce Road,
Cambridge CB3 0WA, U.K.

Max Planck Institute for Gravitational Physics (Albert Einstein Institute), Am Mühlenberg 1,
D-14476 Golm, Germany

E-mail: sg452@cam.ac.uk, gielen@aei.mpg.de

Abstract. We investigate a formulation of continuum 4d gravity in terms of a constrained topological (BF) theory, in the spirit of the Plebanski formulation, but involving only linear constraints, of the type used recently in the spin foam approach to quantum gravity. We identify both the continuum version of the linear simplicity constraints used in the quantum discrete context and a linear version of the quadratic volume constraints that are necessary to complete the reduction from the topological theory to gravity. We illustrate and discuss also the discrete counterpart of the same continuum linear constraints. Moreover, we show under which additional conditions the discrete volume constraints follow from the simplicity constraints, thus playing the role of secondary constraints. Our analysis clarifies how the discrete constructions of spin foam models are related to a continuum theory with an action principle that is equivalent to general relativity.

1. Motivation

As is well known, general relativity in three dimensions is topological, as there are no local degrees of freedom. Its first order formulation in terms of a connection ω^{AB} and a triad E^A is given by

$$S = \frac{1}{16\pi G} \int_{\Sigma} \epsilon_{ABC} E^A \wedge R^{BC}[\omega]. \quad (1)$$

The quantisation of such a theory is rather well understood and different possible quantisation procedures are known. This is of course in stark contrast to general relativity in four dimensions, whose quantisation is to a large extent still an open problem. One may try to exploit what is known in 3D to write general relativity in 4D as a *constrained* topological theory with action

$$S = \int_{\Sigma} B^{AB} \wedge R_{AB}[\omega] + \lambda^{\alpha} C_{\alpha}[B]; \quad (2)$$

one then needs to add constraints (the “simplicity constraints”) which enforce $B^{AB} = \frac{1}{16\pi G} \epsilon^{AB}{}_{CD} E^C \wedge E^D$ (for some set of 1-forms E^A) to recover general relativity. The main obstacle in quantisation is then the identification and implementation of such constraints, which has indeed been a topic of much recent debate in the discrete setting of spin foam models [2].

¹ Based on a talk given at the Spanish Relativity Meeting (ERE2010), Granada, on 6 September 2010 [1].

If, for example as the starting point for loop quantum gravity, one wants to add a Holst term [3] to the first order connection form for general relativity, one instead needs to impose $\Sigma^{AB} = \frac{1}{8\pi\gamma G} E^A \wedge E^B$ on the two-form field Σ^{AB} , defined in terms of B^{AB} by

$$\Sigma^{AB} = \frac{1}{1 \pm \gamma^2} \left(B^{AB} - \frac{\gamma}{2} \epsilon^{AB}{}_{CD} B^{CD} \right) \quad (3)$$

(note that this is just a linear redefinition). We will focus on this case in the following and try to determine an appropriate set of constraints for Σ^{AB} .

In the traditional (Plebanski) formulation [4], one uses *quadratic constraints* for Σ^{AB} ,

$$\epsilon_{ABCD} \Sigma_{ab}^{AB} \Sigma_{cd}^{CD} = V \epsilon_{abcd}, \quad (4)$$

which have two *separate* sectors of solutions²:

$$\text{either } \Sigma^{AB} = \pm e^A \wedge e^B \quad \text{or } \Sigma^{AB} = \pm \frac{1}{2} \epsilon^{AB}{}_{CD} e^C \wedge e^D \quad (5)$$

for some set of 1-forms e^A . Classically, one can consistently remain within the ‘‘GR’’ sector [5]; quantum mechanically, the situation is less clear.

The discrete analogue of this construction has been used in spin foam models of quantum gravity: One introduces a triangulation of spacetime, integrates Σ^{AB} over triangles

$$\Sigma_{ab}^{AB}(x) \Rightarrow \Sigma_{\Delta}^{AB} \equiv \int_{\Delta} \Sigma^{AB} \in \mathfrak{so}(4) \simeq \Lambda^2 \mathbb{R}^4, \quad (6)$$

and imposes the constraint $\epsilon_{ABCD} \Sigma_{\Delta}^{AB} \Sigma_{\Delta'}^{CD} = 0$ if $\Delta = \Delta'$ or Δ and Δ' share an edge. It can be shown [6] that the remaining ‘‘volume’’ constraints can be replaced by the ‘‘closure constraint’’

$$\sum_{\Delta \subset \mathbb{A}} \Sigma_{\Delta}^{AB} = 0. \quad (7)$$

The quantum-mechanical implementation of these constraints led to the Barrett-Crane model [7]. The recently proposed new spin foam models [2], on the other hand, rely on the replacement of quadratic constraints on Σ_{Δ}^{AB} by *linear* constraints:

$$n_A(\mathbb{A}) \Sigma^{AB}(\Delta) = 0 \quad \forall \Delta \subset \mathbb{A}, \quad (8)$$

where $n_A(\mathbb{A})$ is normal to the tetrahedron \mathbb{A} . These are stronger than the quadratic constraints; they restrict Σ^{AB} to the discrete analogue of $\Sigma^{AB} = \pm E^A \wedge E^B$. Here we present an extension of this construction, which was done in the discrete setting, to the *classical continuum theory*: We give a set of constraints whose solution implies that $\Sigma^{AB} = \pm E^A \wedge E^B$ for some E^A (ignoring the factor $8\pi\gamma G$ in the following). We then discretise these constraints and identify some which are sufficient at the discrete level. For details and proofs of our claims, see [8].

² This is under the additional assumption that $V \neq 0$! $V = 0$ configurations are not geometric at all, leading to additional potential difficulties in quantisation [5].

2. The construction

2.1. Continuum construction

Clearly, to find a continuum analogue of (8) that is derivable from an action, one has to introduce dynamical fields that play the analogue of the normal vectors n^A in (8). We introduce a basis of 1-forms e_a^A (i.e. assuming $\det(e_a^A) \neq 0$), which is dual to a basis of 3-forms by

$$n_{Adef} \equiv \epsilon_{ADEF} e_d^D e_e^E e_f^F, \quad e_A^c \sim \epsilon^{cdef} n_{Adef}. \quad (9)$$

Claim 1. For a basis of 3-forms n_A , the general solution to

$$n_{Adef} \Sigma_{ab}^{AB} = 0 \quad \forall \{a, b\} \subset \{d, e, f\} \quad (10)$$

is

$$\Sigma_{ab}^{AB} = G_{ab} e_a^{[A} e_b^{B]}, \quad (11)$$

where e^A is defined in terms of n_A as above. Note that $G_{ab} = G_{ab}(x)$ are functions on spacetime. One could try a linear redefinition $e_a^A = \lambda_a E_a^A$ to identify this general solution

$$\Sigma_{ab}^{AB} = G_{ab} e_a^{[A} e_b^{B]} \quad (12)$$

with $\Sigma^{AB} = \pm E^A \wedge E^B$, but this is not possible in general; one needs additional conditions.

In [8], we show that imposing the additional three constraints

$$\sum_b \sum_{\{a,f\} \not\subset \{b,e\}} n_{Abef} \Sigma_{ab}^{AB} = 0, \quad e \in \{0, 1, 2\} \text{ fixed}, \quad (13)$$

implies that $G_{ab}(x)$ is the same function $c(x)$ for all a, b , which can be absorbed by a pointwise rescaling $E^A = \sqrt{|c|} e^A$ to complete the identification $\Sigma^{AB} = \pm E^A \wedge E^B$. Note that not the field e^A we have introduced to define the constraints (10), but the rescaled E^A will play the role of an orthonormal tetrad determining physical lengths and angles. In particular, a completely degenerate triad $E^A = 0$ is possible since $c(x) = 0$ at some points is not excluded.

2.2. Discrete construction

The translation of the constraints into discretised variables is straightforward. Again one introduces a triangulation of spacetime and integrates Σ^{AB} over triangles and n_A over tetrahedra:

$$\Sigma_{ab}^{AB}, n_{Adef} \Rightarrow \Sigma^{AB}(\triangle), n_A(\triangle). \quad (14)$$

Then the discrete analogue of the set of constraints (10) is (essentially by construction) the set of linear constraints used in the EPR(L)/FK models, $n_A(\triangle) \Sigma^{AB}(\triangle) = 0 \quad \forall \triangle \subset \triangle$. Perhaps more interestingly, the discrete version of the remaining constraints (13) would take the form

$$\sum_{\{i,j\} \not\supset \mathbf{A}} n_A(\triangle_i) \Sigma^{AB}(\triangle_{\mathbf{A}j}) = 0, \quad (15)$$

where we label the tetrahedra in a 4-simplex by $\mathbf{A}, \mathbf{B}, \mathbf{C}, \mathbf{D}, \mathbf{E}$; there are five of these constraints per simplex (where \mathbf{A} is replaced by $\mathbf{B}, \mathbf{C}, \mathbf{D}, \mathbf{E}$ respectively). These constraints play the role of the ‘‘volume’’ constraints which could be, in the case of quadratic constraints, replaced by a closure constraint relating different triangles in a tetrahedron. The situation here is completely analogous:

Result: The constraints (15) follow from the EPR(L)/FK linear constraints, the closure constraint on $\Sigma^{AB}(\Delta)$, plus an analogous “4D closure constraint”

$$n_A(\Delta_{\mathbf{A}}) + n_A(\Delta_{\mathbf{B}}) + n_A(\Delta_{\mathbf{C}}) + n_A(\Delta_{\mathbf{D}}) + n_A(\Delta_{\mathbf{E}}) = 0, \quad (16)$$

which can be given a clear geometric motivation, just as closure on Σ^{AB} : While the closure constraint (7) can be interpreted as the requirement for the triangles specified by Σ_{Δ}^{AB} to close up to form a tetrahedron, or alternatively as the discrete version of the field equation $\nabla_{[a}\Sigma_{bc]}^{AB} = 0$ integrated over an infinitesimal 3-ball, the four-dimensional constraint (16) can be interpreted as demanding that the five tetrahedra specified by $n^A(\Delta)$ close up to form a 4-simplex. The derivation of (16) from a continuum field equation is more subtle since one has to assume both $\nabla_{[a}\Sigma_{bc]}^{AB} = 0$ and the identification of Σ^{AB} with e^A to obtain the equation $\nabla_{[a}n_{bcd]}^A = 0$.

3. Summary

We have established a formulation of classical general relativity as a topological BF theory plus linear constraints which relied on the introduction of a basis of 3-forms at each point in addition to the two-form field Σ^{AB} and the connection ω^{AB} . This new continuum action is more closely related to current spin foam models, such as the EPR(L) and FK models, as discretising it leads to variables $\Sigma^{AB}(\Delta)$, $n_A(\Delta)$ which appear in these spin foam models, and are subject to the constraints used in these models, plus closure constraints on both Σ^{AB} and n_A . The latter is a new constraint which has been suggested before, but not received much attention in the literature so far. It suggests a new formulation for spin foam models in which the normals n_A carry information about the 3-volume and thus are given a fully geometric role, other than in the current models where one can set them to a fixed value. Other than searching for a modification of spin foam models in this direction, one should perform a detailed canonical analysis of the constraints used in our formulation. There are also indications for a close relation of our constructions both to current research in group field theories, as well as to the “edge simplicity” constraints introduced in [9]. We leave all this to future work.

Acknowledgments

This work was supported by EPSRC and Trinity College, Cambridge, partly through a Rouse Ball Travelling Studentship in Mathematics.

References

- [1] Slides available as <http://www.iaa.es/ere2010/website/modules/tinyd0/content/talks/gielen.pdf>.
- [2] Engle J, Pereira R and Rovelli C 2008 Flipped spinfoam vertex and loop gravity *Nuclear Physics B* **798** 251–290; Freidel L and Krasnov K 2008 A new spin foam model for 4D gravity *Classical and Quantum Gravity* **25** 125018
- [3] Holst S 1996 Barbero’s Hamiltonian derived from a generalized Hilbert-Palatini action *Physical Review D* **53** 5966–5969
- [4] Plebański J F 1977 On the separation of Einsteinian substructures *Journal of Mathematical Physics* **18** 2511–2520
- [5] Reisenberger M P 1999 Classical Euclidean general relativity from ‘left-handed area = right-handed area’ *Classical and Quantum Gravity* **16** 1357–1371
- [6] Livine E R and Speziale S 2008 Solving the simplicity constraints for spinfoam quantum gravity *Europhysics Letters* **81** 50004
- [7] Barrett J W and Crane L 1998 Relativistic spin networks and quantum gravity *Journal of Mathematical Physics* **39** 3296–3302
- [8] Gielen S and Oriti D 2010 Classical general relativity as BF-Plebanski theory with linear constraints *Classical and Quantum Gravity* **27** 185017
- [9] Dittrich B and Ryan J P 2008 Phase space descriptions for simplicial 4d geometries *Preprint* 0807.2806