

Global Pointwise Decay Estimates for Defocusing Radial Nonlinear Wave Equations

ROGER BIELI AND NIKODEM SZPAK

Max-Planck-Institut für Gravitationsphysik, Albert-Einstein-Institut,
Golm, Germany

We prove global pointwise decay estimates for a class of defocusing semilinear wave equations in $n = 3$ dimensions restricted to spherical symmetry. The technique is based on a conformal transformation and a suitable choice of the mapping adjusted to the nonlinearity. As a result we obtain a pointwise bound on the solutions for arbitrarily large Cauchy data, provided the solutions exist globally. The decay rates are identical with those for small data and hence seem to be optimal. A generalization beyond the spherical symmetry is suggested.

Keywords Decay estimates; Large data; Nonlinear wave equations; Pointwise decay; Spherical symmetry.

Mathematics Subject Classification 35L05; 35L15; 35L17.

1. Introduction

We consider a class of nonlinear wave equations

$$\partial_t^2 \phi - \Delta \phi = f(\phi) \quad (1)$$

in $n = 3$ spatial dimensions where the nonlinearity $f(\phi)$ is of a defocusing type. It means that the nonlinear term has a repulsive action on the waves and focusing of waves is suppressed by an energy condition. The sign of the nonlinear term which is chosen is essential such that the following conserved energy is positive definite

$$E = \int_{\mathbb{R}^3} \left(\frac{1}{2} |\partial_t \phi|^2 + \frac{1}{2} |\nabla \phi|^2 + F(\phi) \right) d^3x \quad (2)$$

with $f = -F'$. Equation (1) has been intensively studied in the literature over a few decades, in particular in the case of a pure power nonlinearity $f(\phi) = -|\phi|^{p-1}\phi$.

Received March 18, 2009; Accepted March 28, 2010

Address correspondence to Roger Bieli, Max-Planck-Institut für Gravitationsphysik, Albert-Einstein-Institut, Am Mühlenberg 1, Golm 14476, Germany; E-mail: bieli@aei.mpg.de

Let us collect the most important results defining the context for this work. The global existence of C^2 solutions to the Cauchy problem has been shown by Jörrens [11] in the energy subcritical case $1 < p < 5$ and later by Grillakis [8] in the critical case $p = 5$ (see also [17] for the spherically symmetric $p = 5$ case) while not much is known about the supercritical case $p > 5$. For more references related to global existence we recommend the book of Sogge [13]. Uniform boundedness of solutions for $2 < p < 5$ has been proved by Pecher [12]. Uniform decay $1/t^{1-\epsilon}$ and scattering have been proved for $3 \leq p < 5$ by Strauss [14]. Bahouri and Shatah [2] have shown that finite energy solutions decay to zero for $p = 5$ and Hidano [9] has shown scattering and decay to zero for $2.5 < p \leq 3$. Ginibre and Velo [7] have shown scattering in the energy space for nonlinearities near $p = 1 + 4/(n - 2)$ in dimensions $n \geq 3$ but for $n = 3$ only with smallness assumption. Scattering results imply, in some sense, that the solutions behave asymptotically like solutions of the linear equation. (For more scattering results we refer to the monograph of Strauss [15].)

Here, we go further and study the pointwise behavior of solutions for $3 \leq p < 5$. Such results in dimension $n = 3$ exist only for small data [1, 16, 18, 19] and are based on perturbation techniques which cannot be generalized to large data. We extend the technique of conformal compactification developed by Choquet-Bruhat, Christodoulou and others in [3, 5, 6]. The novelty is that we use a suitably chosen conformal transformation, adjusted to the nonlinearity. In this setting, we first show uniform boundedness of the transformed solutions in the precompact region and then use the inverse transformation to get an extra decaying (conformal) factor and thus a pointwise decay estimate for the original solution. We claim that this estimate is optimal (for generic initial data) as it is identical with the one obtained for small data where it has been shown to be optimal [18, 19]. Numerical simulations done by Bizoń et al. [4] also support this picture.

The Main Result and the Method

In the following we study the Cauchy problem for the radial semilinear wave equation

$$\partial_t^2 \phi - \Delta \phi = -|\phi|^{p-1} \phi \quad (3)$$

with the initial data of compact support $r \in [0, \alpha_p[$, where $\alpha_p > 0$ will be specified later and chosen at time $t = 1$

$$\phi(1, r) = \phi_0, \quad \partial_t \phi(1, r) = \phi_1 \quad (4)$$

and prove that the solutions satisfy for all $t \geq 1$ and $0 \leq r \leq t$

$$|\phi(t, r)| \leq \frac{C}{(1+t+r)(1+t-r)^{p-2}} \quad (5)$$

with some constant C depending only on ϕ_0 , ϕ_1 and p . (Symmetries of the equation (3) allow for mapping any compactly supported initial data onto the interval $r \in [0, \alpha_p[$ at $t = 1$.)

Briefly, our method is based on a conformal mapping $(u, v) \rightarrow (\tilde{u}, \tilde{v})$ defined in the double-null coordinates $u = t + r, v = t - r$,

$$\tilde{u} := -\frac{1}{u^{p-2}}, \quad \tilde{v} := -\frac{1}{v^{p-2}}. \quad (6)$$

The conformal factor $\Omega := \tilde{r}/r$ multiplies the transformed solution $\tilde{\phi}(\tilde{u}, \tilde{v}) := \Omega^{-1}\phi(u, v)$ which satisfies the transformed wave equation

$$\tilde{\square}\tilde{\phi} + \frac{1}{(p-2)^2} \left[\frac{\tilde{u} - \tilde{v}}{(-\tilde{v})^{1/(p-2)} - (-\tilde{u})^{1/(p-2)}} \right]^{p-1} \tilde{\phi}|\tilde{\phi}|^{p-1} = 0 \quad (7)$$

in a precompact region of spacetime. There, we are able to show boundedness of some pseudo-energy flux what we further use to show the uniform boundedness of $|\tilde{\phi}(\tilde{u}, \tilde{v})| \leq \tilde{C}$. Finally, the inverse transformation provides the desired estimate

$$|\phi(u, v)| \leq \tilde{C} \cdot \Omega = \tilde{C} \cdot \frac{1}{(uv)^{p-2}} \cdot \frac{u^{p-2} - v^{p-2}}{u - v} \leq \frac{C}{u \cdot v^{p-2}}.$$

The power $p - 2$ in the mapping (6) cannot be increased, what would potentially lead to a stronger pointwise decay of the solutions, because then the factor multiplying the nonlinearity in (7) becomes singular and our boundedness theorems cannot be applied.

The estimate (5) is also optimal in the sense of compatibility with the small data case. For small initial data it has been shown in [19] that the solutions behave asymptotically, for large t and fixed r , like

$$\phi(t, r) = \frac{C}{t^{p-1}} + \mathcal{O}(t^{-p})$$

where the constant C can be expressed explicitly via ϕ_0 and ϕ_1 . Hence, the estimate (5) applied to the small data case gives the optimal decay rate.

The assumption of spherical symmetry is essential for the method to yield optimal decay for $p > 3$. On $\mathbb{R} \times \mathbb{R}^3$ the obvious analogue of the map $(u, v) \rightarrow (\tilde{u}, \tilde{v})$ is no longer conformal, unless $p = 3$, of course. The reason is that radial and angular parts of the wave operator transform diversely and the resulting equation is no longer semilinear. Applying the transformation corresponding to $p = 3$, which is conformal, to an equation with nonlinearity of power $p > 3$ still gives a global $1/(uv)$ -decay result, being however not optimal and independent of the actual power p .

2. Conformal Transformation

In the case $p = 3$, the radial wave equation (3), having the explicit form

$$\partial_t^2 \phi - \partial_r^2 \phi - \frac{2}{r} \partial_r \phi + \phi^3 = 0,$$

is not only invariant under time translation, $\phi(t, r) \mapsto \phi(t + a, r)$, scaling, $\phi(t, r) \mapsto \lambda\phi(\lambda t, \lambda r)$, and reflection, $\phi \mapsto -\phi$, but also under a *conformal inversion* which is

given by

$$\phi(t, r) \mapsto \frac{1}{t^2 - r^2} \phi\left(\frac{-t}{t^2 - r^2}, \frac{r}{t^2 - r^2}\right). \quad (8)$$

This inversion maps solutions on the interior of the forward light-cone $K^+ := \{(t, r) \mid 0 \leq r < t\}$ to solutions on the interior of the backward light-cone $K^- := \{(t, r) \mid 0 \leq r < -t\}$ of the origin and vice versa. Thus, establishing boundedness of a solution on K^- towards the future immediately implies pointwise decay estimates for the transformed solution on K^+ . In order for this to yield optimal decay in the case of more general nonlinearities it is necessary to also consider more general conformal transformations which is what will be discussed in the following. For this, the standard conformal method (see e.g., [10, Section 6.7]) is applied with a different choice of the conformal factor.

On $(t, r) \in \mathbb{R} \times \mathbb{R}_+$ consider null coordinates $u := t + r$ and $v := t - r$. Then u and v are positive on K^+ and negative on K^- . For any $p > 2$ define, with respect to the (u, v) -coordinate system, the map

$$\Phi : K^+ \rightarrow K^-, \quad (u, v) \mapsto \left(-\frac{1}{u^{p-2}}, -\frac{1}{v^{p-2}}\right).$$

It is analytic with analytic inverse

$$\Phi^{-1} : K^- \rightarrow K^+, \quad (u, v) \mapsto ((-u)^{-\frac{1}{p-2}}, (-v)^{-\frac{1}{p-2}}).$$

Furthermore, there exists a positive analytic function $\Omega > 0$ on K^+ with the property that

$$r\Omega = r \circ \Phi \quad (9)$$

holds on K^+ . If $\eta = dt^2 - dr^2$ denotes the Minkowski metric on $\mathbb{R} \times \mathbb{R}_+$ then its pullback by Φ satisfies

$$\Phi^*\eta = \frac{(p-2)^2}{(t^2 - r^2)^{p-1}} \eta \quad \text{on } K^+,$$

which shows that mapping by Φ is conformal. In the case $p = 3$ the map $\phi \mapsto \Omega \Phi^* \phi = \Omega \cdot (\phi \circ \Phi)$ corresponds to the conformal inversion (8) since then $\Omega = 1/(t^2 - r^2)$. This conformal transformation is illustrated in Figure 1. Note that with

$$\alpha_p := 1 - \frac{1}{2^{\frac{1}{p-2}}} \in]0, 1[,$$

for any compact interval I contained in $\{1\} \times [0, \alpha_p[$ the interval J is compactly contained in $\{-1\} \times [0, 1[$, where J is the intersection of the future of the curve $H := \Phi(I)$ with the line $\{t = -1\}$. Moreover, for $p \geq 3$ one has $\alpha_p \leq 1/2$,




Figure 1. The function Φ maps the interior of the forward light-cone K^+ to the interior of the backward light-cone K^- conformally and bianalytically. An interval I of the line $\{t = 1\}$, containing the support of the original Cauchy data, is mapped to a curve H . The future of the interval I is mapped into the future of the curve H . Initial data on H is evolved in the region S to yield new initial data on the interval J of the line $\{t = -1\}$.

Suppose that $h \in C^2(K^-)$ is a twice continuously differentiable function on K^- . Then $\Omega\Phi^*h$ is a function on K^+ of class C^2 for which

$$\square(\Omega\Phi^*h) = (\square\Omega)\Phi^*h + \Omega \frac{(p-2)^2}{(t^2-r^2)^{p-1}} \Phi^*\square h,$$

where $\square := \partial_t^2 - \partial_r^2 - (2/r)\partial_r$ on $\mathbb{R} \times \mathbb{R}_+$. With the special choice of $h = 1/r$, this identity together with relation (9) implies that $(\square\Omega)\Phi^*(1/r) = \square[\Omega\Phi^*(1/r)] = 0$ since $\square(1/r) = 0$ and $\Omega\Phi^*(1/r) = 1/r$. Therefore, $\square\Omega = 0$, such that in general

$$\square(\Omega\Phi^*h) = (p-2)^2(uv)^{-(p-1)}\Omega\Phi^*\square h. \tag{10}$$

Hence, if $p > 2$ and $\phi \in C^2(K^+)$ is a classical solution of the radial wave equation

$$\square\phi + \phi|\phi|^{p-1} = 0 \tag{11}$$

on K^+ , its conformal transformation $\psi := \Phi_*(\Omega^{-1}\phi) \in C^2(K^-)$ satisfies

$$\square\psi + \frac{1}{(p-2)^2}[\Phi_*(\Omega uv)^{p-1}]\psi|\psi|^{p-1} = 0, \quad (12)$$

where Φ_* denotes the push-forward by the diffeomorphism Φ . On the other hand, it also follows directly from (10) that if $\psi \in C^2(K^-)$ is a classical solution of the transformed equation (12) on K^- , the function $\phi := \Omega\Phi^*\psi \in C^2(K^+)$ solves the original equation (11) on K^+ .

3. Evolution of the Pseudo-Energy

As mentioned introductorily, for equations of the form (1) there is a *conserved* energy (2) that in particular controls the spatial L^{p+1} -norm of the solution. One can no longer expect to find such a quantity for the conformally transformed equation (12) or, more generally, for an equation

$$\square\psi + c(t, r)\psi|\psi|^{p-1} = 0 \quad (13)$$

with a non-negative function c of class C^2 . However, an analogue of the energy expression, although not conserved exactly, will turn out to be sufficient to prove boundedness of ψ on the relevant region of K^- , provided the function c is monotonically decreasing in t there. This pseudo-energy will now be defined.

Let $\psi \in C^2(K^-)$ be a classical solution of (13) on K^- for $p > 2$. Then the vector field \mathcal{E} given by

$$\mathcal{E} := r^2 \left[\frac{1}{2}(\partial_t \psi)^2 + \frac{1}{2}(\partial_r \psi)^2 + \frac{1}{p+1}c|\psi|^{p+1} \right] \partial_t - r^2(\partial_t \psi \partial_r \psi) \partial_r,$$

is continuously differentiable and

$$\operatorname{div} \mathcal{E} = r^2 [\square\psi + c\psi|\psi|^{p-1}] \partial_t \psi + \frac{1}{p+1}r^2(\partial_r c)|\psi|^{p+1} = \frac{1}{p+1}r^2(\partial_r c)|\psi|^{p+1}$$

holds on K^- . Assume furthermore that $\partial_r c \leq 0$ is non-positive such that the same is true for $\operatorname{div} \mathcal{E}$. This implies, recalling Figure 1, together with the assumption that the support I of the initial data is compactly contained in $\{1\} \times [0, \alpha_p]$, that

$$\begin{aligned} 0 &\geq \int_S (\operatorname{div} \mathcal{E}) dt \wedge dr = \int_{\partial S} i_{\mathcal{E}}(dt \wedge dr) = \int_{J \cup H} [(i_{\mathcal{E}} dt) dr - (i_{\mathcal{E}} dr) dt] \\ &= \int_J r^2 \left[\frac{1}{2}(\partial_t \psi)^2 + \frac{1}{2}(\partial_r \psi)^2 + \frac{1}{p+1}c|\psi|^{p+1} \right] dr \\ &\quad - \frac{p-2}{2} \int_I (i_{\mathcal{E}} dt) \circ \Phi(1+r, 1-r) \cdot \left[\frac{1}{(1-r)^{p-1}} + \frac{1}{(1+r)^{p-1}} \right] dr \\ &\quad - \frac{p-2}{2} \int_I (i_{\mathcal{E}} dr) \circ \Phi(1+r, 1-r) \cdot \left[\frac{1}{(1-r)^{p-1}} - \frac{1}{(1+r)^{p-1}} \right] dr, \end{aligned}$$

where $i_{\mathcal{E}}$ denotes the interior multiplication with the vector field \mathcal{E} . The volume form $dt \wedge dr$ is chosen such that the interval J carries the standard orientation by dr as

does the interval I . The last two integrals over the interval I depend only on the initial data ϕ_0 and ϕ_1 on I and are certainly finite. Thus

$$E_0 := \int_J r^2 \left[\frac{1}{2} (\partial_t \psi)^2 + \frac{1}{2} (\partial_r \psi)^2 + \frac{1}{p+1} c |\psi|^{p+1} \right] dr \leq C \quad (14)$$

for a constant C .

The pseudo-energy E_0 now controls light-cone integrals of the quantity $c|\psi|^{p+1}$ according to the following Proposition 1, a fact that will be essential for proving boundedness of ψ on the region $K := K^- \cap \{-1 \leq t < 0\}$.

Proposition 1. *Let $\psi \in C^2(K)$ be a solution of (13) with initial data supported on J satisfying the estimate (14). If $\partial_t c \leq 0$ on K then*

$$\frac{1}{p+1} \int_{-1}^{t_0} [r^2 c |\psi|^{p+1}](s, t_0 - s) ds \leq E_0$$

holds for any $-1 \leq t_0 < 0$.

Proof. It is useful to consider the functions

$$E_{t_0}(t) := \int_0^{t_0-t} (i_{\mathcal{E}} dt)(t, r) dr$$

for any $-1 \leq t \leq t_0 < 0$. Then, for a fixed $-1 < t_0 < 0$, E_{t_0} is continuously differentiable and

$$\begin{aligned} E'_{t_0}(t) &= -(i_{\mathcal{E}} dt)(t, t_0 - t) - \int_0^{t_0-t} [\partial_r (i_{\mathcal{E}} dr)](t, r) dr + \int_0^{t_0-t} (\operatorname{div} \mathcal{E})(t, r) dr \\ &= -(t_0 - t)^2 \left[\frac{1}{2} (\partial_t \psi - \partial_r \psi)^2 + \frac{1}{p+1} c |\psi|^{p+1} \right](t, t_0 - t) + \int_0^{t_0-t} (\operatorname{div} \mathcal{E})(t, r) dr \end{aligned}$$

holds for all $-1 \leq t \leq t_0$. Hence, it follows that

$$\begin{aligned} \operatorname{Flux}(-1, t_0) &:= \int_{-1}^{t_0} i_{\mathcal{E}}(dt + dr)(s, t_0 - s) ds \\ &= \int_{-1}^{t_0} (t_0 - s)^2 \left[\frac{1}{2} (\partial_t \psi - \partial_r \psi)^2 + \frac{1}{p+1} c |\psi|^{p+1} \right](s, t_0 - s) ds \\ &= - \int_{-1}^{t_0} E'_{t_0}(s) ds + \int_{-1}^{t_0} \int_0^{t_0-t} (\operatorname{div} \mathcal{E})(s, r) dr ds \\ &= E_{t_0}(-1) + \int_{-1}^{t_0} \int_0^{t_0-t} (\operatorname{div} \mathcal{E})(s, r) dr ds \leq E_0, \end{aligned}$$

since $E_{t_0}(t_0) = 0$, $E_{t_0}(-1) \leq E_0$ and $\operatorname{div} \mathcal{E} \leq 0$. But this implies

$$\frac{1}{p+1} \int_{-1}^{t_0} (t_0 - s)^2 [c |\psi|^{p+1}](s, t_0 - s) ds \leq \operatorname{Flux}(-1, t_0) \leq E_0,$$

and hence the claim. \square

4. Boundedness

Using the light-cone estimate established in Proposition 1 the boundedness of ψ in the region $K = K^- \cap \{-1 \leq t < 0\}$ can be shown by a simple argument going back to Pecher [12].

Theorem 1. *Let $\psi \in C^2(K)$ be a solution of (13) with $2 < p < 5$ and initial data supported on J satisfying the estimate (14). If $c \geq 0$ is uniformly bounded and $\partial_t c \leq 0$ on K then ψ is uniformly bounded.*

Proof. Let $\psi_0 \in C^2(K)$ be a classical solution of the homogeneous equation $\square\psi_0 = 0$ with the same initial data as ψ . Then, for a fixed $(t, r) \in K$, it holds that

$$|\psi - \psi_0|(t, r) \leq \frac{1}{2r} \int_{-1}^t \int_{|t-s-r|}^{t-s+r} \gamma c(s, y) |\psi|^p(s, y) dy ds.$$

Due to the fact that $2 < p < 5$ there exists a q with $3/2 < q < (p + 1)/(p - 1)$. Changing variables and applying Hölder’s inequality yields

$$\begin{aligned} & \frac{1}{2r} \int_{-1}^t \int_{|t-s-r|}^{t-s+r} \gamma c(s, y) |\psi|^p(s, y) dy ds \\ &= \frac{1}{2r} \int_a^{t+r} \int_{-1}^b (t_0 - s) c(s, t_0 - s) |\psi|^p(s, t_0 - s) ds dt_0 \\ &\leq \frac{1}{2r} \left[\int_a^{t+r} \int_{-1}^b (t_0 - s)^2 c^q(s, t_0 - s) |\psi|^{pq}(s, t_0 - s) ds dt_0 \right]^{\frac{1}{q}} \\ &\quad \times \left[\int_a^{t+r} \int_{-1}^b (t_0 - s)^{\frac{q-2}{q-1}} ds dt_0 \right]^{\frac{q-1}{q}}, \end{aligned} \tag{15}$$

where the abbreviations $a := \max\{t - r, r - t - 2\}$ and $b := [t_0 + (t - r)]/2$ were introduced. Consider the first integral. Since $q > 1$ and c is bounded, so is c^{q-1} . Furthermore, $0 < pq - (p + 1) < q$, so that

$$\begin{aligned} & \int_a^{t+r} \int_{-1}^b (t_0 - s)^2 c^q(s, t_0 - s) |\psi|^{pq}(s, t_0 - s) ds dt_0 \\ &\leq C \left(\sup_{-1 \leq \tau \leq t} \|\psi(\tau, \cdot)\|_{L^\infty} \right)^{\gamma q} \int_a^{t+r} \int_{-1}^b (t_0 - s)^2 c(s, t_0 - s) |\psi|^{p+1}(s, t_0 - s) ds dt_0, \end{aligned}$$

where $0 < \gamma < 1$ is such that $\gamma q = pq - (p + 1)$. The integration variable t_0 takes values $t_0 \geq a$ which ensures $-1 \leq b \leq t_0$. Thus, the integral in s can be estimated by virtue of Proposition 1 to yield

$$\int_a^{t+r} \int_{-1}^b (t_0 - s)^2 c(s, t_0 - s) |\psi|^{p+1}(s, t_0 - s) ds dt_0 \leq \int_{t-r}^{t+r} (p + 1) E_0 dt_0 \leq 2rCE_0.$$

The second integral in (15) can be calculated directly to give

$$\int_a^{t+r} \int_{-1}^b (t_0 - s)^{\frac{q-2}{q-1}} ds dt_0 \leq \frac{q - 1}{2q - 3} \int_{t-r}^{t+r} (1 + t_0)^{\frac{2q-3}{q-1}} dt_0 \leq 2rC$$

because $q > 3/2$. To sum up, the estimate

$$|\psi - \psi_0|(t, r) \leq \frac{C}{2r} \left(\sup_{-1 \leq \tau \leq t} \|\psi(\tau, \cdot)\|_{L^\infty} \right)^\gamma (2rE_0)^{\frac{1}{q}} (2r)^{\frac{q-1}{q}} = CE_0^{\frac{1}{q}} \left(\sup_{-1 \leq \tau \leq t} \|\psi(\tau, \cdot)\|_{L^\infty} \right)^\gamma$$

holds true for any $(t, r) \in K$. But since the solution ψ_0 of the homogeneous equation is clearly bounded and $0 < \gamma < 1$ this estimate implies the boundedness of ψ itself uniformly on K , for the function $t \mapsto \sup_{-1 \leq \tau \leq t} \|\psi(\tau, \cdot)\|_{L^\infty}$ is continuous and finite at $t = -1$. \square

With the function ψ bounded on K the decay of ϕ in the future of I follows immediately.

Corollary 1. *Let ϕ be a classical solution of the wave equation (11) for $3 \leq p < 5$ with initial data $\phi_0 \in C^2(\mathbb{R}_+)$ and $\phi_1 \in C^1(\mathbb{R}_+)$ given at $t = 1$ and which exists globally towards the future. Assume that the support of ϕ_0 and ϕ_1 is contained in the compact interval $I \subset [0, \alpha_p[$. Then there is a constant $C \in \mathbb{R}_+$ such that ϕ satisfies the decay estimate*

$$|\phi(t, r)| \leq \frac{C}{(1+t+r)(1+t-r)^{p-2}} \quad (16)$$

for all $t \geq 1$ and $0 \leq r \leq t$.

Proof. Given such a solution ϕ of class C^2 , it was shown in section 2 that its conformal transformation $\psi = \Phi_*(\Omega^{-1}\phi)$ is a classical solution of the wave equation (13) on the future of the curve H in K^- , cf. Figure 1, with

$$c = \frac{1}{(p-2)^2} \Phi_*(\Omega uv)^{p-1}$$

according to equation (12). Now, for $v < u < 0$

$$\Phi_*(\Omega uv) = \frac{u-v}{(-v)^{\frac{1}{p-2}} - (-u)^{\frac{1}{p-2}}},$$

so that c is bounded on the future of H in K^- owing to $p \geq 3$. Furthermore,

$$\partial_t [\Phi_*(\Omega uv)] = \frac{1}{p-2} \frac{\Phi_*(\Omega uv)^2}{2r} [(-v)^{-\frac{p-3}{p-2}} - (-u)^{-\frac{p-3}{p-2}}],$$

which implies $\partial_t c \leq 0$ on K^- again by reason of $p \geq 3$. Moreover, as detailed in section 3, the assumptions on the support of ϕ_0 and ϕ_1 guarantee that J is compactly contained in $\{-1\} \times [0, 1[$ and that the estimate (14) holds. Thus, Theorem 1 applies and yields boundedness of ψ on K . Since ψ is also bounded on the compact region S it follows that ψ is bounded on the whole future of H in K^- , say $|\psi| \leq C$. But then

$$|\phi(t, r)| = \Omega(t, r) |\Phi^* \psi(t, r)| \leq C \Omega(t, r)$$

on the whole future of I in K^+ , while

$$\Omega(t, r) = \frac{1}{2rv^{p-2}} \left[1 - \left(\frac{v}{u} \right)^{p-2} \right] \leq \frac{p-2}{2rv^{p-2}} \left(1 - \frac{v}{u} \right) = \frac{p-2}{uv^{p-2}}.$$

Since in this region $u \geq 1$ and $v \geq 1 - \alpha_p$ and outside, for $v < 1 - \alpha_p$, the solution vanishes identically, the bound for $\phi(t, r)$ can be written in the regularized form (16). \square

5. Discussion

As already noted in the introduction, the conformal transformations which we consider for the radial problem are not conformal for the full problem, beyond the spherical symmetry, unless $p = 3$. However, for the full problem with $3 < p < 5$ we can still apply the conformal ($p = 3$) transformation and obtain a global pointwise decay $1/(uv)$. The decay rate is then not optimal but can act as a prerequisite for a more refined asymptotic analysis. Indeed, that decay rate is sufficient for the solution to become small at large times in the sense which allows a perturbative analysis with methods similar to those developed in [18]. We want to address this issue in a forthcoming publication.

It seems, at least as far as the spherical symmetry is concerned, that our method can be applied in higher than $n = 3$ odd dimensions.

Also the linear wave equations with strong positive potentials

$$\partial_t^2 \phi - \Delta \phi + V(x)\phi = 0$$

having prescribed decay at spatial infinity $V(x) \sim 1/|x|^k$ still lack a sharp pointwise decay estimate and seem to be treatable with our method after some modifications. Such equations have a positive definite energy, too, and can be conformally transformed to a form analogous to (13) however with a weakly singular positive function c . This idea shall be addressed in another publication.

Acknowledgments

We would like to thank the referee for his valuable comments improving the manuscript. One of the authors (NS) wants to express his gratitude to the Mittag-Leffler Institute and to the organizers of the workshop “Geometry, Analysis, and General Relativity” (Fall 2008) for hospitality and fantastic, creative and relaxed atmosphere during the work which resulted in this article.

References

- [1] Asakura, F. (1986). Existence of a global solution to a semi-linear wave equation with slowly decreasing initial data in three space dimensions. *Comm. Part. Diff. Eqs.* 13:1459–1487.
- [2] Bahouri, H., Shatah, J. (1998). Decay estimates for the critical semilinear wave equation. *Ann. Inst. Henri Poincaré* 15:783–789.
- [3] Baez, J.C., Segal, I.E., Zhou, Z.-F. (1990). The global Goursat problem and scattering for nonlinear wave equations. *J. Func. Anal.* 93:239–269.

- [4] Bizoń, P. (2008). Private communication.
- [5] Choquet-Bruhat, Y., Paneitz, S.M., Segal, I.E. (1983). The Yang-Mills equations on the universal cosmos. *J. Func. Anal.* 53:112–150.
- [6] Christodoulou, D. (1986). Global solutions of nonlinear hyperbolic equations for small initial data. *Comm. Pure Appl. Math.* 39:267–282.
- [7] Ginibre, J., Velo, G. (1989). Scattering theory in the energy space for a class of non-linear wave equations. *Comm. Math. Phys.* 123:535–573.
- [8] Grillakis, M.G. (1990). Regularity and asymptotic behavior of the wave equation with a critical nonlinearity. *Ann. Math.* 132:485–509.
- [9] Hidano, K. (2001). Scattering problem for the nonlinear wave equation in the finite energy and conformal charge space. *J. Func. Anal.* 187:274–307.
- [10] Hörmander, L. (1997). *Lectures on Nonlinear Hyperbolic Differential Equations*. Berlin: Springer-Verlag.
- [11] Jörgens, K. (1961). Das Anfangswertproblem im Grossen für eine Klasse nichtlinearer Wellengleichungen [The initial boundary problem in the large for a class of nonlinear wave equations]. *Math. Z.* 77:295–307.
- [12] Pecher, H. (1974). Das Verhalten globaler Lösungen nichtlinearer Wellengleichungen für grosse Zeiten [The behavior of global solutions on nonlinear wave equations for large times]. *Math. Z.* 136:67–92.
- [13] Sogge, C.D. (1995). *Lectures on Nonlinear Wave Equations*. Monographs in Analysis, Vol. II. Boston: International Press Inc.
- [14] Strauss, W. (1968). Decay and asymptotics for $u = F(u)$. *J. Func. Anal.* 2:409–457.
- [15] Strauss, W. (1989). *Nonlinear Wave Equations*. Providence, RI: American Mathematical Society.
- [16] Strauss, W., Tsutaya, K. (1997). Existence and blow up of small amplitude nonlinear waves with a negative potential. *Discr. Cont. Dyn. Sys.* 3:175–188.
- [17] Struwe, M. (1988). Globally regular solutions to the u^5 Klein-Gordon equation. *Ann. Sci. Norm. Sup. Pisa* 15:495–513.
- [18] Szpak, N. (2008). Linear and nonlinear tails I: general results and perturbation theory. *J. Hyp. Diff. Eqs.* 5:741–765.
- [19] Szpak, N., Bizoń, P., Chmaj, T., Rostworowski, A. (2009). Linear and nonlinear tails II: Spherical symmetry. *J. Hyp. Diff. Eqs.* 6:107–125.