

Future non-linear stability for solutions of the Einstein-Vlasov system of Bianchi types II and VI₀

Ernesto Nungesser

Albert-Einstein-Institute Max-Planck-Institute for Gravitational Physics, Am Muehlenberg 1, Golm 14476, Germany

(Received 31 August 2012; accepted 28 September 2012; published online 18 October 2012)

In a recent paper [E. Nungesser, “Future non-linear stability for reflection symmetric solutions of the Einstein-Vlasov system of Bianchi types II and VI₀,” *Annales Henri Poincaré* (2012)], we have treated the future nonlinear stability for reflection symmetric solutions of the Einstein-Vlasov system of Bianchi types II and VI₀. We have been able now to remove the reflection symmetry assumption, thus treating the non-diagonal case. Apart from the increasing complexity, the methods have been essentially the same as in the diagonal case, showing that they are thus quite powerful. Here, the challenge was to put the equations in a form that permits the use of the previous results. We are able to conclude that after a possible basis change, the future of the non-diagonal spacetimes in consideration is asymptotically diagonal. © 2012 American Institute of Physics. [<http://dx.doi.org/10.1063/1.4758930>]

I. THE EINSTEIN-VLASOV SYSTEM

A cosmological model represents a universe at a certain averaging scale. It is described via a Lorentzian metric $g_{\alpha\beta}$ (we will use signature $- + + +$) on a manifold M and a family of fundamental observers. The metric is assumed to be time-orientable, which means that at each point of M the two halves of the light cone can be labelled past and future in a way, which varies continuously from point to point. This enables to distinguish between future-pointing and past-pointing timelike vectors. This is a physically reasonable assumption from both a macroscopic point of view, e.g., the increase of entropy and also from a microscopic point of view, e.g., the kaon decay. One has also to specify the matter model and this we will do in the following. The interaction between the geometry and the matter is described by the Einstein field equations (we use geometrized units, i.e., the gravitational constant G and the speed of light in vacuum c are set equal to one):

$$G_{\alpha\beta} = 8\pi T_{\alpha\beta},$$

where $G_{\alpha\beta}$ is the Einstein tensor and $T_{\alpha\beta}$ is the energy-momentum tensor. For the matter model, we will take the point of view of kinetic theory.¹² The sign conventions of Ref. 10 and the Einstein summation convention that repeated indices are to be summed over are used. Latin indices run from one to three and Greek ones from zero to three.

We will consider from now on that all the particles have *equal* mass m . We will choose units such that $m = 1$, which means that a distinction between velocities and momenta is not necessary. The collection of particles (galaxies or clusters of galaxies) will be described (statistically) by a non-negative real valued distribution function $f(x^\alpha, p^\alpha)$ on the mass shell. This function represents the density of particles at a given spacetime point with given four-momentum. Using the geodesic equations, the restriction of the Liouville operator to the mass shell has the following form:

$$L = p^\alpha \frac{\partial}{\partial x^\alpha} - \Gamma_{\beta\gamma}^a p^\beta p^\gamma \frac{\partial}{\partial p^a},$$

where $\Gamma_{\beta\gamma}^{\alpha}$ are the components of the metric connection. We will consider the collisionless case, which is described via the Vlasov equation

$$L(f) = 0.$$

The unknowns of our system are a 4-manifold M , a Lorentz metric $g_{\alpha\beta}$ on this manifold and the distribution function f on the mass shell defined by the metric. We have the Vlasov equation defined by the metric for the distribution function and the Einstein equations. It remains to define the energy-momentum tensor $T_{\alpha\beta}$ in terms of the distribution and the metric. Before that, we need a Lorentz invariant volume element on the mass shell. A point of the tangent space has the volume element $|g^{(4)}|^{\frac{1}{2}} dp^0 dp^1 dp^2 dp^3$ ($g^{(4)}$ is the determinant of the spacetime metric), which is Lorentz invariant. Now considering p^0 as a dependent variable, the induced (Riemannian) volume of the mass shell considered as a hypersurface in the tangent space at that point is

$$\varpi = 2H(p^\alpha)\delta(p_\alpha p^\alpha + m^2)|g^{(4)}|^{\frac{1}{2}} dp^0 dp^1 dp^2 dp^3,$$

where δ is the Dirac distribution function and $H(p^\alpha)$ is defined to be one if p^α is future directed and zero otherwise. We can write this explicitly as

$$\varpi = |p_0|^{-1}|g^{(4)}|^{\frac{1}{2}} dp^1 dp^2 dp^3.$$

Now we define the energy momentum tensor as follows:

$$T_{\alpha\beta} = \int f(x^\alpha, p^\alpha) p_\alpha p_\beta \varpi.$$

One can show that $T_{\alpha\beta}$ is divergence-free and thus it is compatible with the Einstein equations. For collisionless matter, all the energy conditions hold. The Vlasov equation in a fixed spacetime can be solved by the method of characteristics

$$\frac{dX^a}{ds} = P^a; \quad \frac{dP^a}{ds} = -\Gamma_{\beta\gamma}^a P^\beta P^\gamma.$$

Let $X^a(s, x^\alpha, p^a)$, $P^a(s, x^\alpha, p^a)$ be the unique solution of that equation with initial conditions $X^a(t, x^\alpha, p^a) = x^a$ and $P^a(t, x^\alpha, p^a) = p^a$. Then the solution of the Vlasov equation can be written as

$$f(x^\alpha, p^a) = f_0(X^a(0, x^\alpha, p^a), P^a(0, x^\alpha, p^a)),$$

where f_0 is the restriction of f to the hypersurface $t = 0$. It follows that if f_0 is bounded the same is true for f . We will assume that f has compact support in momentum space for each fixed t . This property holds if the initial datum f_0 has compact support and if each hypersurface $t = t_0$ is a Cauchy hypersurface. Before coming to our symmetry assumption, we want to briefly introduce the initial value problem for the Einstein-Vlasov system. In general, the initial data for the Einstein-matter equations consist of a metric g_{ab} on the initial hypersurface, the second fundamental form k_{ab} on that hypersurface, and some matter data. Thus, we have a Riemannian metric g_{ab} , a symmetric tensor k_{ab} and some matter fields defined on an abstract 3-dimensional manifold S .

Solving the initial value problem means embedding S into a 4-dimensional M on which are defined a Lorentzian metric $g_{\alpha\beta}$ and matter fields such that g_{ab} and k_{ab} are the pullbacks to S of the induced metric and second fundamental form of the image of the embedding of S while f is the pullback of the matter fields. Finally, $g^{\alpha\beta}$ and f have to satisfy the Einstein-matter equations.

For the Einstein-Vlasov system, it has been shown that given an initial data set, there exists a corresponding solution of the Einstein-Vlasov system and that this solution is locally unique up to diffeomorphism. The extension to a global theorem has not been achieved yet. However, if one assumes that the initial data have certain symmetry, this symmetry is inherited by the corresponding solutions. In particular, for the case we will deal with, i.e., expanding Bianchi models (except type IX) coupled to dust or to collisionless matter, the spacetime is future complete (Theorem 2.1 of Ref. 9).

II. BIANCHI SPACETIMES

The basis for the classification of homogeneous spacetimes is the work of Bianchi, which was introduced to cosmology by Taub. Here, we will use the modern terminology and we define Bianchi spacetimes as follows:

Definition 1: A Bianchi spacetime is defined to be a spatially homogeneous spacetime whose isometry group possesses a three-dimensional subgroup G that acts simply transitively on the spacelike orbits.

Our results concern only a special class of the Bianchi spacetimes, namely that of class A.

Definition 2: A Bianchi A spacetime is a Bianchi spacetime whose three-dimensional Lie algebra has traceless structure constants, i.e., $C_{ba}^a = 0$.

We will study II and VI₀. For Bianchi II, the only non-vanishing structure constants are

$$C_{23}^1 = 1 = -C_{32}^1 \quad (1)$$

and in the case of Bianchi VI₀, these are

$$C_{31}^2 = 1 = -C_{13}^2, \quad C_{21}^3 = 1 = -C_{12}^3. \quad (2)$$

We will use the metric approach. If \mathbf{W}^a denote the 1-forms dual to the frame vectors \mathbf{E}_a , the metric of a Bianchi spacetime takes the form

$${}^4g = -dt^2 + g_{ab}(t)\mathbf{W}^a\mathbf{W}^b, \quad (3)$$

where g_{ab} (and all other tensors) on G will be described in terms of the frame components of a left invariant frame. A dot above a letter will denote a derivative with respect to the cosmological time t . We will use the 3+1 decomposition of the Einstein equations as made in Ref. 10. Comparing our metric (3) with (2.28) of Ref. 10, we have that $\alpha = 1$ and $\beta^a = 0$, which means that the lapse function is the identity and the shift vector vanishes. There, the abstract index notation is used. We can interpret the quantities as being frame components. There are different projections of the energy momentum tensor, which are important

$$\begin{aligned} \rho &= T^{00}, \\ j_a &= T_a^0, \\ S_{ab} &= T_{ab}, \end{aligned}$$

where ρ is the energy density and j_a is the matter current.

The second fundamental form k_{ab} can be expressed as

$$\dot{g}_{ab} = -2k_{ab}. \quad (4)$$

The Einstein equations

$$\dot{k}_{ab} = R_{ab} + kk_{ab} - 2k_{ac}k_b^c - 8\pi(S_{ab} - \frac{1}{2}g_{ab}S) - 4\pi\rho g_{ab}, \quad (5)$$

where we have used the notations $S = g^{ab}S_{ab}$, $k = g^{ab}k_{ab}$, and R_{ab} is the Ricci tensor of the three-dimensional metric. The evolution equation for the mixed version of the second fundamental form is (2.35) of Ref. 10

$$\dot{k}_b^a = R_b^a + kk_b^a - 8\pi S_b^a + 4\pi\delta_b^a(S - \rho). \quad (6)$$

From the constraint equations since k only depends on the time variable, we have that

$$R - k_{ab}k^{ab} + k^2 = 16\pi\rho, \quad (7)$$

$$\nabla^a k_{ab} = 8\pi j_b, \quad (8)$$

where R is the Ricci scalar curvature.

Another useful relation concerns the determinant g of the induced metric ((2.30) of Ref. 10)

$$\frac{d}{dt}(\log g) = -2k. \quad (9)$$

Taking the trace of (6)

$$\dot{k} = R + k^2 + 4\pi S - 12\pi\rho. \quad (10)$$

With (7), one can eliminate the energy density and (10) reads

$$\dot{k} = \frac{1}{4}(k^2 + R + 3k_{ab}k^{ab}) + 4\pi S. \quad (11)$$

Finally, if one substitutes for the Ricci scalar with (7)

$$\dot{k} = k_{ab}k^{ab} + 4\pi(S + \rho). \quad (12)$$

Now with the 3+1 formulation, our initial data are $(g_{ij}(t_0), k_{ij}(t_0), f(t_0))$, i.e., a Riemannian metric, a second fundamental form, and the distribution function of the Vlasov equation, respectively, on a three-dimensional manifold $S(t_0)$. This is the initial data set at $t = t_0$ for the Einstein-Vlasov system.

We assume that $k < 0$ for all time following Ref. 8 (see comments below Lemma 2.2 of Ref. 8). This enables us to set without the loss of generality $t_0 = -2/k(t_0)$. The reason for this choice will become clear later and is of technical nature.

We will now introduce several new variables in order to use the ones, which are common in Bianchi cosmologies and to be able to compare results. We can decompose the second fundamental form introducing σ_{ab} as the trace-free part

$$k_{ab} = \sigma_{ab} - Hg_{ab}, \quad (13)$$

$$k_{ab}k^{ab} = \sigma_{ab}\sigma^{ab} + 3H^2. \quad (14)$$

Using the Hubble parameter,

$$H = -\frac{1}{3}k,$$

we define

$$\Sigma_a^b = \frac{\sigma_a^b}{H} \quad (15)$$

and

$$\Sigma_+ = -\frac{1}{2}(\Sigma_2^2 + \Sigma_3^3), \quad (16)$$

$$\Sigma_- = -\frac{1}{2\sqrt{3}}(\Sigma_2^2 - \Sigma_3^3). \quad (17)$$

Thus,

$$\Sigma_a^b = \begin{pmatrix} 2\Sigma_+ & \Sigma_2^1 & \Sigma_3^1 \\ \Sigma_1^2 & -\Sigma_+ - \sqrt{3}\Sigma_- & \Sigma_3^2 \\ \Sigma_1^3 & \Sigma_2^3 & -\Sigma_+ + \sqrt{3}\Sigma_- \end{pmatrix}.$$

The reason for using the variables Σ_+ and Σ_- is that the diagonal case has been very important to understand the non-diagonal case. Define also

$$\Omega = 8\pi\rho/3H^2, \quad (18)$$

$$q = -1 - \frac{\dot{H}}{H^2}, \quad (19)$$

$$\frac{d\tau}{dt} = H. \quad (20)$$

The time variable τ is dimensionless and sometimes very useful. From (7), we obtain the constraint equation

$$\frac{1}{6H^2}(R - \sigma_{ab}\sigma^{ab}) = \Omega - 1,$$

and from (11), the evolution equation for the Hubble variable

$$\partial_t(H^{-1}) = \frac{3}{2} + \frac{1}{12}\left(\frac{R}{H^2} + \frac{3}{H^2}\sigma_{ab}\sigma^{ab}\right) + \frac{4\pi S}{3H^2}. \quad (21)$$

Combining the last two equations with (6), we obtain the evolution equations for Σ_- and Σ_+

$$\dot{\Sigma}_+ = H\left[\frac{2R - 3(R_2^2 + R_3^2)}{6H^2} - \Sigma_+(3 + \frac{\dot{H}}{H^2}) + \frac{4\pi}{3H^2}(3S_2^2 + 3S_3^3 - 2S)\right], \quad (22)$$

$$\dot{\Sigma}_- = H\left[\frac{R_3^3 - R_2^2}{2\sqrt{3}H^2} - (3 + \frac{\dot{H}}{H^2})\Sigma_- + \frac{4\pi(S_2^2 - S_3^3)}{\sqrt{3}H^2}\right]. \quad (23)$$

Since we use a left-invariant frame, f will not depend on x^a and the Vlasov equation takes the form

$$p^0 \frac{\partial f}{\partial t} - \Gamma_{\beta\gamma}^a p^\beta p^\gamma \frac{\partial f}{\partial p^a} = 0.$$

It turns out that the equation simplifies if we express f in terms of p_i instead of p^i what we can do due to the mass shell relation

$$p^0 \frac{\partial f}{\partial t} - \Gamma_{a\beta\gamma} p^\beta p^\gamma \frac{\partial f}{\partial p_a} = 0.$$

Because of our special choice of frame, the metric has the simple form (3). Due to the fact that we are contracting and the antisymmetry of the structure constants, we finally arrive at

$$\frac{\partial f}{\partial t} + (p^0)^{-1} C_{ba}^d p^b p_d \frac{\partial f}{\partial p_a} = 0. \quad (24)$$

From (24), it is also possible to define the characteristic curve V_a

$$\frac{dV_a}{dt} = (V^0)^{-1} C_{ba}^d V^b V_d, \quad (25)$$

for each $V_i(\bar{t}) = \bar{v}_i$ given \bar{t} . Note that if we define

$$V = g^{ij} V_i V_j \quad (26)$$

due to the antisymmetry of the structure constants, we have with (25)

$$\frac{dV}{dt} = \frac{d}{dt}(g^{ij})V_i V_j. \quad (27)$$

Let us also write down the components of the energy momentum tensor in our frame

$$T_{00} = \int f(t, p^a) p^0 \sqrt{g} dp^1 dp^2 dp^3, \quad (28)$$

$$T_{0j} = - \int f(t, p^a) p_j \sqrt{g} dp^1 dp^2 dp^3, \quad (29)$$

$$T_{ij} = \int f(t, p^a) p_i p_j (p^0)^{-1} \sqrt{g} dp^1 dp^2 dp^3. \quad (30)$$

III. THE ASYMPTOTICS OF BIANCHI II AND VI₀

Before coming to the non-diagonal case, we have a look at the tilted fluid models, since they are non-diagonal as well and they may help us to understand the non-diagonal case with collisionless matter. For the tilted Bianchi II, we use the corresponding equations of Ref. 5 and for Bianchi VI₀ the equations of Ref. 4, in both cases with $\gamma = 1$. We will not go into the details for this we refer to the mentioned work. The point is that looking at the linearization we see that the variables, which did not appear in the diagonal case have decay rates, which are between the ones considered previously. This is a good sign. Also in Ref. 2, the stability of the Ellis-MacCallum solution, in fact the stability of the Collins solution, was already considered within the Einstein-Euler system.

A. Equations of the non-diagonal case

Using (15), we arrive with (6) for $a \neq b$ to

$$\dot{\Sigma}_a^b = H \left[\frac{R_a^b}{H^2} - \Sigma_a^b \left(3 + \frac{\dot{H}}{H^2} \right) - \frac{8\pi S_a^b}{H^2} \right]; \quad a \neq b,$$

which together with (22)–(23), i.e.,

$$\begin{aligned} \dot{\Sigma}_+ &= H \left[\frac{2R - 3(R_2^2 + R_3^2)}{6H^2} - \Sigma_+ \left(3 + \frac{\dot{H}}{H^2} \right) + \frac{4\pi}{3H^2} (3S_2^2 + 3S_3^2 - 2S) \right], \\ \dot{\Sigma}_- &= H \left[\frac{R_3^3 - R_2^2}{2\sqrt{3}H^2} - \left(3 + \frac{\dot{H}}{H^2} \right) \Sigma_- + \frac{4\pi(S_2^2 - S_3^2)}{\sqrt{3}H^2} \right] \end{aligned}$$

describe the evolution of Σ_a^b . The expression for the Ricci tensor is

$$R_{ij} = -\frac{1}{2} C_{ki}^l (C_{lj}^k + g_{lm} g^{kn} C_{nj}^m) - \frac{1}{4} C_{nk}^m C_{ql}^p g_{jm} g_{ip} g^{kq} g^{ln}, \quad (31)$$

and

$$R_i^j = R_{ib} g^{bj} = -\frac{1}{2} C_{ki}^l g^{bj} (C_{lb}^k + g_{lm} g^{kn} C_{nb}^m) - \frac{1}{4} C_{nk}^j C_{ql}^p g_{ip} g^{kq} g^{ln}. \quad (32)$$

We will now derive some expression concerning the derivative of (31)

$$\begin{aligned} \dot{R}_{ij} &= C_{ki}^l C_{nj}^m (k_{lm} g^{kn} - g_{lm} k^{kn}) + \\ &\quad \frac{1}{2} C_{nk}^m C_{ql}^p (k_{jm} g_{ip} g^{kq} g^{ln} + g_{jm} k_{ip} g^{kq} g^{ln} - g_{jm} g_{ip} k^{kq} g^{ln} - g_{jm} g_{ip} g^{kq} k^{ln}). \end{aligned}$$

Thus,

$$\begin{aligned} g^{jr} \dot{R}_{ij} &= g^{jr} C_{ki}^l C_{nj}^m (k_{lm} g^{kn} - g_{lm} k^{kn}) + \\ &\quad \frac{1}{2} C_{ql}^p [C_{nk}^m k_{ip} g^{kq} g^{ln} + C_{nk}^r (k_{ip} g^{kq} g^{ln} - g_{ip} (k^{kq} g^{ln} + g^{kq} k^{ln}))]. \end{aligned}$$

For $r = i$ and relabelling, the m with i for the terms with the prefactor $\frac{1}{2}$,

$$g^{ji} \dot{R}_{ij} = g^{ji} C_{ki}^l C_{nj}^m (k_{lm} g^{kn} - g_{lm} k^{kn}) + \frac{1}{2} C_{ql}^p C_{nk}^i [2k_{ip} g^{kq} g^{ln} - g_{ip} (k^{kq} g^{ln} + g^{kq} k^{ln})].$$

Rearranging terms

$$g^{ji} \dot{R}_{ij} = C_{ki}^l C_{nj}^m (k_{lm} g^{kn} g^{ji} - g_{lm} k^{kn} g^{ji}) + C_{ql}^p C_{nk}^i [k_{ip} g^{kq} g^{ln} - g_{ip} k^{kq} g^{ln}].$$

We see that the first with the third and the second with the fourth term cancel each other, hence,

$$g^{ji} \dot{R}_{ij} = 0. \quad (33)$$

The evolution equation for the Ricci scalar due to (33) is

$$\dot{R} = 2R^i_j k_i^j = 2H(-R + R^i_j \Sigma_i^j).$$

Define

$$N_i^j = \frac{R_i^j}{H^2}.$$

The derivative of this expression is

$$\dot{N}_i^j = \frac{g^{pj} \dot{R}_{pi}}{H^2} + 2H(N_i^p \Sigma_p^j - (1 + \frac{\dot{H}}{H^2})N_i^j).$$

Consider the quantity $N = R/H^2$. Its evolution equation is

$$\dot{N} = 2H[qN + N_i^j \Sigma_i^j]. \quad (34)$$

B. Curvature expressions

For bookkeeping reasons, we define the following quantities where we use from now on g for the determinant of the metric:

$$\begin{aligned} A &= g^{22}g^{33} - (g^{23})^2 = \frac{g_{11}}{g}; & B &= g^{13}g^{23} - g^{12}g^{33} = \frac{g_{12}}{g}, \\ C &= g^{12}g^{23} - g^{13}g^{22} = \frac{g_{13}}{g}; & D &= g^{12}g^{13} - g^{11}g^{23} = \frac{g_{23}}{g}, \\ E &= g^{11}g^{33} - (g^{13})^2 = \frac{g_{22}}{g}; & F &= g^{11}g^{22} - (g^{12})^2 = \frac{g_{33}}{g}. \end{aligned}$$

Let us denote the quantities divided by H^2 with small letters, i.e., $a = \frac{A}{H^2}$.

1. Curvature expressions for Bianchi II

Using (32) for Bianchi II,

$$R_i^j = \frac{1}{2}g_{11}[C_{2i}^1(g^{23}g^{2j} - g^{22}g^{3j}) + C_{i3}^1(g^{23}g^{3j} - g^{33}g^{2j})] + \frac{1}{2}g_{i1}C_{23}^j A.$$

We obtain

$$R = -\frac{1}{2}g_{11}A = -\frac{1}{2}\frac{(g_{11})^2}{g}$$

and as in the diagonal case,

$$\begin{aligned} R_1^1 &= -R = -R_2^2 = -R_3^3, \\ R_1^2 &= R_1^3 = R_2^3 = R_3^2 = 0. \end{aligned}$$

However, in the non-diagonal case, we have

$$\begin{aligned} R_2^1 &= -2\frac{g_{12}}{g_{11}}R, \\ R_3^1 &= -2\frac{g_{13}}{g_{11}}R. \end{aligned}$$

Thus,

$$\dot{N} = -2H[(1 + \frac{\dot{H}}{H^2} + 4\Sigma_+)N - W_{II}],$$

where $W_{II} = N_2^1 \Sigma_1^2 + N_3^1 \Sigma_1^3$. In order to calculate the derivative of N_2^1 , we need the following expression:

$$R \frac{d}{dt}(-2\frac{g_{12}}{g_{11}}) = 2H[2\Sigma_2^1 R + (3\Sigma_+ + \sqrt{3}\Sigma_-)R_2^1 - \frac{1}{2R}((R_2^1)^2 \Sigma_1^2 + R_3^1 R_2^1 \Sigma_1^3)].$$

Hence,

$$\begin{aligned}\dot{N}_2^1 &= H[4N\Sigma_2^1 - 2(\Sigma_+ + 1 - \sqrt{3}\Sigma_- + \frac{\dot{H}}{H^2})N_2^1 + W_2^1], \\ \dot{N}_3^1 &= H[4N\Sigma_3^1 - 2(\Sigma_+ + 1 + \sqrt{3}\Sigma_- + \frac{\dot{H}}{H^2})N_3^1 + W_3^1],\end{aligned}$$

where

$$\begin{aligned}W_2^1 &= -2\Sigma_2^3 N_3^1 + N_2^1 N^{-1}(\Sigma_1^2 N_2^1 + \Sigma_1^3 N_3^1), \\ W_3^1 &= -2\Sigma_3^2 N_2^1 + N_3^1 N^{-1}(\Sigma_1^2 N_2^1 + \Sigma_1^3 N_3^1).\end{aligned}$$

2. Curvature expressions for Bianchi VI₀

With (32), we obtain

$$\begin{aligned}-2R_i^j &= g^{1j}(C_{2i}^3 + C_{3i}^2) + g_{i2}(C_{13}^j E - C_{12}^j D) + g_{i3}(-C_{13}^j D + C_{12}^j F) \\ &+ g_{22}[C_{1i}^2(-g^{3j}g^{11} + g^{1j}g^{13}) + C_{3i}^2(-g^{3j}g^{31} + g^{1j}g^{33})] \\ &+ g_{33}[C_{1i}^3(-g^{2j}g^{11} + g^{1j}g^{12}) + C_{2i}^3(-g^{2j}g^{21} + g^{1j}g^{22})] \\ &+ g_{23}[C_{1i}^2(g^{1j}g^{12} - g^{2j}g^{11}) + C_{3i}^2(g^{1j}g^{23} - g^{2j}g^{13}) \\ &+ C_{1i}^3(g^{1j}g^{13} - g^{3j}g^{11}) + C_{2i}^3(g^{1j}g^{23} - g^{3j}g^{21})].\end{aligned}$$

In particular,

$$\begin{aligned}R &= -\frac{1}{2}[(\sqrt{g_{22}E} + \sqrt{g_{33}F})^2 - 4g_{23}D] = -\frac{1}{2g}[(g_{22} + g_{33})^2 - 4g_{23}^2], \\ R_2^2 &= \frac{1}{2}(g_{22}E - g_{33}F) = \frac{1}{2g}[(g_{22})^2 - (g_{33})^2],\end{aligned}$$

and like in the diagonal case,

$$\begin{aligned}R &= R_1^1, \\ R_2^2 &= -R_3^3, \\ R_2^1 &= R_3^1 = 0.\end{aligned}$$

However, we have

$$\begin{aligned}N_2^3 &= -N_3^2 = g_{23}(f - e) = -N_{23}(N_3 + N_2), \\ N_1^2 &= -2\frac{g^{12}}{H^2} + g_{12}(e - f) = N_{12}(N_2 - N_3) - 2N_{13}N_{23}, \\ N_1^3 &= -2\frac{g^{13}}{H^2} + g_{13}(f - e) = N_{13}(N_2 - N_3) - 2N_{12}N_{23},\end{aligned}$$

where N_{ij} is defined as

$$N_{ij} = \frac{g_{ij}}{\sqrt{g}H},$$

and

$$\begin{aligned}N_2 &= N_{22}, \\ N_3 &= -N_{33},\end{aligned}$$

which means that $N_2^2 = R_2^2/H^2$

$$N_2^2 = \frac{1}{2}((N_2)^2 - (N_3)^2).$$

Recalling that

$$\frac{\dot{g}}{g} = 6H,$$

we can compute the derivatives of N_{ij} using the following formula:

$$\dot{N}_{ij} = H[qN_{ij} - 2\Sigma_i^l N_{lj}].$$

Hence,

$$\dot{N}_{12} = H[(q - 4\Sigma_+)N_{12} - 2\Sigma_1^2 N_2 - 2\Sigma_1^3 N_{23}], \quad (35)$$

$$\dot{N}_{13} = H[(q - 4\Sigma_+)N_{13} - 2\Sigma_1^2 N_{23} + 2\Sigma_1^3 N_3], \quad (36)$$

$$\dot{N}_{23} = H[(2\Sigma_+ + 2\sqrt{3}\Sigma_- + q)N_{23} + 2\Sigma_2^3 N_3 - 2\Sigma_2^1 N_{13}], \quad (37)$$

$$\dot{N}_2 = H[(2\Sigma_+ + 2\sqrt{3}\Sigma_- + q)N_2 - 2\Sigma_2^1 N_{12} - 2\Sigma_2^3 N_{23}], \quad (38)$$

$$\dot{N}_3 = H[(2\Sigma_+ - 2\sqrt{3}\Sigma_- + q)N_3 + 2\Sigma_3^1 N_{13} + 2\Sigma_3^2 N_{23}]. \quad (39)$$

From (34), we obtain

$$\dot{N} = 2H[(2\Sigma_+ + q)N - 2\sqrt{3}\Sigma_- N_2^2 + N_3^2 \Sigma_2^3 + N_2^3 \Sigma_3^2 + N_1^2 \Sigma_2^1 + N_1^3 \Sigma_3^1]. \quad (40)$$

The evolution equation for N_2^2

$$\begin{aligned} \dot{N}_2^2 = & H[2(2\Sigma_+ + q)N_2^2 + 2\sqrt{3}\Sigma_-((N_3)^2 + (N_2)^2) - 2(\Sigma_2^1 N_{12} N_2 \\ & + \Sigma_3^1 N_{13} N_3 + \Sigma_2^3 N_{23} N_2 + \Sigma_3^2 N_{23} N_3)]. \end{aligned} \quad (41)$$

C. The non-diagonal asymptotics of Bianchi II and VI₀

We will now discuss the asymptotics of the non-diagonal case. The structure of the analysis is very similar to the diagonal case. We start with a bootstrap argument and end with applying Arzela-Ascoli. Next we will collect the bootstrap assumptions. The prefactors denoted by A and some index are small constants.

1. Bootstrap assumptions for Bianchi II

$$|\Sigma_+ - \frac{1}{8}| \leq A_+(1+t)^{-\frac{3}{8}},$$

$$|N + \frac{9}{32}| \leq A_c(1+t)^{-\frac{3}{8}},$$

$$|\Sigma_3^2| \leq A_{23}(1+t)^{-\frac{3}{8}},$$

$$|\Sigma_2^3| \leq A_{32}(1+t)^{-\frac{3}{8}},$$

$$|\Sigma_2^1| \leq A_{12},$$

$$|\Sigma_3^1| \leq A_{13},$$

$$|N_2^1| \leq A_{c12},$$

$$|N_3^1| \leq A_{c13},$$

$$P \leq A_m(1+t)^{-\frac{1}{3}},$$

$$|\Sigma_-| \leq A_-(1+t)^{-\frac{3}{4}},$$

$$|\Sigma_1^2| \leq A_{21}(1+t)^{-\frac{3}{4}},$$

$$|\Sigma_1^3| \leq A_{31}(1+t)^{-\frac{3}{4}}.$$

2. Bootstrap assumptions for Bianchi VI₀

$$|\Sigma_+ + \frac{1}{4}| \leq A_+(1+t)^{-\frac{3}{8}},$$

$$|\Sigma_-| \leq A_-(1+t)^{-\frac{3}{8}},$$

$$|N + \frac{9}{8}| \leq A_{c1}(1+t)^{-\frac{3}{8}},$$

$$|N_2^2| \leq A_{c2}(1+t)^{-\frac{3}{8}},$$

$$|N_{12}| \leq C_1,$$

$$|N_{13}| \leq C_2,$$

$$|N_{23}| \leq A_{c23},$$

$$P \leq A_m(1+t)^{-\frac{1}{3}},$$

$$|\Sigma_3^2| \leq A_{23}(1+t)^{-\frac{3}{4}},$$

$$|\Sigma_2^3| \leq A_{32}(1+t)^{-\frac{3}{4}},$$

$$|\Sigma_2^1| \leq A_{12}(1+t)^{-\frac{3}{4}},$$

$$|\Sigma_3^1| \leq A_{13}(1+t)^{-\frac{3}{4}},$$

$$|\Sigma_1^3| \leq C_3,$$

$$|\Sigma_1^2| \leq C_4.$$

3. Mean curvature

Concerning the estimate of H , there is no difference with respect to the diagonal case. The reason is that the estimate of D

$$D = \frac{1}{12}(N + \frac{3}{H^2}\sigma_{ab}\sigma^{ab}) + \frac{4\pi S}{3H^2}$$

is the same. Thus, as in the diagonal case, it follows from (21) that

$$\partial_t(H^{-1}) = \frac{3}{2} + O(\epsilon t^{-\frac{3}{8}}),$$

and following the steps made for the diagonal case, we arrive at.

$$H = \frac{2}{3}t^{-1}(1 + O(\epsilon t^{-\frac{3}{8}}))$$

will hold.

4. Estimate of the metric and P

For a matrix A , its norm can be defined as

$$\|A\| = \sup\{|Ax|/|x| : x \neq 0\}.$$

Let B and C be $n \times n$ symmetric matrices with C positive definite. It is possible to define a *relative norm* by

$$\|B\|_C = \sup\{|Bx|/|Cx| : x \neq 0\}.$$

Clearly,

$$\|B\| \leq \|B\|_C \|C\|.$$

It also true that

$$\|B\|_C \leq \sqrt{\text{tr}(C^{-1}BC^{-1}B)}. \tag{42}$$

This can be shown as follows. Consider the common eigenbasis b_i of B and C . Then there exist α_i such that $Bb_i = \alpha_i Cb_i$ for each i . Then (42) is equivalent to the statement that the maximum modulus of any α_i is smaller than $\Sigma_i \alpha_i^2$. Using (42), we obtain in the sense of *quadratic forms*

$$\sigma^{ab} \leq (\sigma_{cd}\sigma^{cd})^{\frac{1}{2}} g^{ab}. \tag{43}$$

Define

$$\bar{g}^{ab} = t^{\frac{p}{q}} g^{ab}.$$

Then

$$\frac{d}{dt}(t^{-\gamma} \bar{g}^{ab}) = t^{-\gamma-1} \bar{g}^{ab} (-\gamma + \frac{p}{q}) + 2t^{-\gamma+\frac{p}{q}} (\sigma^{ab} - H g^{ab}),$$

where we have introduced for technical reasons a small positive parameter γ . Using now the inequality (43)

$$\frac{d}{dt}(t^{-\gamma} \bar{g}^{ab}) \leq t^{-\gamma-1} \bar{g}^{ab} [-\gamma + \frac{p}{q} + 2tH((H^{-2}\sigma_{cd}\sigma^{cd})^{\frac{1}{2}} - 1)]. \tag{44}$$

Using the equation (44) and the estimate of H

$$\frac{d}{dt}(t^{-\gamma} \bar{g}^{ab}) \leq t^{-\gamma-1} \bar{g}^{ab} [-\gamma + \frac{p}{q} + \frac{4}{3}(1 + O(\epsilon t^{-\frac{3}{8}}))((H^{-2}\sigma_{cd}\sigma^{cd})^{\frac{1}{2}} - 1)].$$

We obtain decay for the metric (in the sense of quadratic forms) provided that $(H^{-2}\sigma_{cd}\sigma^{cd})^{\frac{1}{2}} \leq 1$. This holds for Bianchi II and VI₀ with for instance $\frac{p}{q} = 0.4$. Thus, we have

$$g^{ab} \leq t^{-\frac{p}{q}} t_0^{\frac{p}{q}} g^{ab}(t_0).$$

This implies that the components of the metric are also bounded by some constant $C(t_0)$, which depends on the terms of $g^{ab}(t_0)$. Consider now

$$\dot{g}^{bf} = 2H(\Sigma_a^b - \delta_a^b)g^{af}.$$

Since the metric components are bounded, the non-diagonal terms will contribute only with an ϵ . Thus, we have for every component g^{ij} (no summation over the indices in the following equation):

$$\dot{g}^{ij} = 2H(\Sigma_i^i - 1 + \epsilon)g^{ij} \leq 2H(\max(\Sigma_i^i) - 1 + \epsilon)g^{ij} = 2H(-\frac{3}{4} + \epsilon)g^{ij}.$$

Using now the estimate of H

$$\dot{g}^{ij} \leq t^{-1}(-1 + \epsilon)g^{ij}. \quad (45)$$

One can conclude that

$$\|g^{-1}\| \leq O(t^{-1+\epsilon}).$$

From (45),

$$\dot{V} = \dot{g}^{bf} V_b V_f \leq t^{-1}(-1 + \epsilon)V,$$

which means that

$$V = O(t^{-1+\epsilon}),$$

which gives us the same decay for P as in the diagonal case

$$P = O(t^{-\frac{1}{2}+\epsilon}).$$

5. Closing the bootstrap argument for Bianchi II

It follows immediately by the same arguments as in the diagonal case

$$\Sigma_- = O(t^{-1+\epsilon})$$

$$\Sigma_1^2 = O(t^{-1+\epsilon}),$$

$$\Sigma_1^3 = O(t^{-1+\epsilon}),$$

$$\Sigma_2^3 = O(t^{-1+\epsilon}),$$

$$\Sigma_3^2 = O(t^{-1+\epsilon}).$$

Defining $(N_1)^2 = -2N$, we arrive at

$$\dot{\Sigma}_+ = H\left[\frac{(N_1)^2}{3} - \Sigma_+(3 + \frac{\dot{H}}{H^2}) + \frac{4\pi}{3H^2}(3S_2^2 + 3S_3^3 - 2S)\right],$$

$$\dot{\Sigma}_- = H\left[-(3 + \frac{\dot{H}}{H^2})\Sigma_- + \frac{4\pi(S_2^2 - S_3^3)}{\sqrt{3}H^2}\right],$$

$$\dot{N}_1 = H\left[(1 + \frac{\dot{H}}{H^2} + 4\Sigma_+)N_1 + 2\frac{W_{II}}{N_1}\right].$$

Since $2\frac{W_{LL}}{N_1}$ decays like $t^{-1+\epsilon}$, we see that we can apply the same arguments as in the diagonal case to obtain an improvement of the bootstrap assumptions

$$\begin{aligned} \Sigma_+ - \frac{1}{8} &= O(t^{-\frac{1}{2}+\epsilon}), \\ \Sigma_- &= O(t^{-\frac{1}{2}+\epsilon}), \\ N_1 - \frac{3}{4} &= O(t^{-\frac{1}{2}+\epsilon}). \end{aligned}$$

The system which remains using the time variable τ is the following:

$$\begin{aligned} (\Sigma_2^1)' &= \Sigma_2^1(q - 2) + N_2^1 - \frac{8\pi S_2^1}{H^2}, \\ (\Sigma_3^1)' &= \Sigma_3^1(q - 2) + N_3^1 - \frac{8\pi S_3^1}{H^2}, \\ (N_2^1)' &= -2(N_1)^2 \Sigma_2^1 - 2(\Sigma_+ - q - \sqrt{3}\Sigma_-)N_2^1 + W_2^1, \\ (N_3^1)' &= -2(N_1)^2 \Sigma_3^1 - 2(\Sigma_+ - q + \sqrt{3}\Sigma_-)N_3^1 + W_3^1. \end{aligned}$$

Let us focus on the $\Sigma_2^1 - N_2^1$ -system. Using the estimates obtained, we arrive at

$$\begin{pmatrix} \Sigma_2^1 \\ N_2^1 \end{pmatrix}' = \begin{pmatrix} -\frac{3}{2} & 1 \\ -\frac{9}{8} & \frac{3}{4} \end{pmatrix} \begin{pmatrix} \Sigma_2^1 \\ N_2^1 \end{pmatrix} + O(\epsilon e^{(-\frac{3}{4}+\epsilon)\tau}) \begin{pmatrix} 1 \\ 1 \end{pmatrix}.$$

Let us go to the basis of eigenvectors of the linear system via the linear transformation

$$\begin{pmatrix} \check{\Sigma}_2^1 \\ \check{N}_2^1 \end{pmatrix} = \begin{pmatrix} \frac{3}{2} & -1 \\ -\frac{3}{2} & 2 \end{pmatrix} \begin{pmatrix} \Sigma_2^1 \\ N_2^1 \end{pmatrix}.$$

Thus, we arrive at

$$\begin{pmatrix} \check{\Sigma}_2^1 \\ \check{N}_2^1 \end{pmatrix}' = \begin{pmatrix} -\frac{3}{4} & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} \check{\Sigma}_2^1 \\ \check{N}_2^1 \end{pmatrix} + O(\epsilon e^{(-\frac{3}{4}+\epsilon)\tau}) \begin{pmatrix} 1 \\ 1 \end{pmatrix}.$$

Using the bootstrap assumptions for Σ_2^1 and N_2^1 , we have an assumption for $\check{\Sigma}_2^1$. By the usual contradiction argument, we arrive at

$$\check{\Sigma}_2^1 = \check{\Sigma}_2^1(\tau_0)e^{(-\frac{3}{4}+\epsilon)\tau}.$$

Integrating the equation for \check{N}_2^1 , we arrive at

$$\check{N}_2^1 = \check{N}_2^1(\tau_0) + O(\epsilon).$$

Going back to the variables Σ_2^1 and N_2^1 via

$$\begin{pmatrix} \Sigma_2^1 \\ N_2^1 \end{pmatrix} = \frac{1}{3} \begin{pmatrix} 4 & 2 \\ 3 & 3 \end{pmatrix} \begin{pmatrix} \check{\Sigma}_2^1 \\ \check{N}_2^1 \end{pmatrix},$$

$$\Sigma_2^1(\tau) = [2\Sigma_2^1(\tau_0) - \frac{4}{3}N_2^1(\tau_0)]e^{(-\frac{3}{4}+\epsilon)\tau} + \frac{4}{3}N_2^1(\tau_0) - \Sigma_2^1(\tau_0) + O(\epsilon),$$

$$N_2^1(\tau) = [\frac{3}{2}\Sigma_2^1(\tau_0) - N_2^1(\tau_0)]e^{(-\frac{3}{4}+\epsilon)\tau} + 2N_2^1(\tau_0) - \frac{3}{2}\Sigma_2^1(\tau_0) + O(\epsilon).$$

Changing back to the time variable t

$$\Sigma_2^1(t) = C(t_0)[2\Sigma_2^1(t_0) - \frac{4}{3}N_2^1(t_0)]t^{-\frac{1}{2}+\epsilon} + \frac{4}{3}N_2^1(t_0) - \Sigma_2^1(t_0) + O(\epsilon),$$

$$N_2^1(t) = C(t_0)[\frac{3}{2}\Sigma_2^1(t_0) - N_2^1(t_0)]t^{-\frac{1}{2}+\epsilon} + 2N_2^1(t_0) - \frac{3}{2}\Sigma_2^1(t_0) + O(\epsilon),$$

where C is a constant, in particular $C(t_0) = t_0^{\frac{1}{3}} e^{-\frac{3}{4}\tau_0}$. The only term which could prevent us from improving the estimates is the ϵ coming from the bootstrap assumptions of Σ_2^1 , but note that it comes in combination with Σ_1^2 as a product of both, thus the last term $O(\epsilon)$ on the right hand side of the last two equations does not prevent us from improving our estimates. Thus, if we wait long time enough and choose $N_2^1(t_0)$ and $\Sigma_2^1(t_0)$ small enough, we will have an improvement for N_2^1 and Σ_2^1 since we can choose them independently and smaller than A_{12} and A_{c12} . There is no difference in the procedure for N_3^1 and Σ_3^1 .

6. Arzela-Ascoli for Bianchi II

Since all estimates have been improved, we can apply Arzela-Ascoli and we arrive for Σ_2^1 and N_2^1 to

$$\begin{aligned} \Sigma_2^1(t = \infty) &= \frac{4}{3}N_2^1(t_0) - \Sigma_2^1(t_0), \\ N_2^1(t = \infty) &= 2N_2^1(t_0) - \frac{3}{2}\Sigma_2^1(t_0). \end{aligned}$$

Consider now the following transformation of the basis vector

$$\begin{aligned} \tilde{e}_1 &= e_1, \\ \tilde{e}_2 &= e_2 + ae_1, \\ \tilde{e}_3 &= e_3 + be_1. \end{aligned}$$

It preserves the Lie-algebra, i.e., the Bianchi type. The following relation holds between the variables Σ_2^1 and Σ_3^1 in the different basis:

$$\begin{aligned} \begin{pmatrix} \tilde{\Sigma}_1^1 & \tilde{\Sigma}_1^2 & \tilde{\Sigma}_1^3 \\ \tilde{\Sigma}_2^1 & \tilde{\Sigma}_2^2 & \tilde{\Sigma}_2^3 \\ \tilde{\Sigma}_3^1 & \tilde{\Sigma}_3^2 & \tilde{\Sigma}_3^3 \end{pmatrix} &= \begin{pmatrix} 1 & 0 & 0 \\ a & 1 & 0 \\ b & 0 & 1 \end{pmatrix} \begin{pmatrix} \Sigma_1^1 & 0 & 0 \\ \Sigma_2^1 & \Sigma_2^2 & 0 \\ \Sigma_3^1 & 0 & \Sigma_3^3 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ -a & 1 & 0 \\ -b & 0 & 1 \end{pmatrix} \\ &= \begin{pmatrix} \Sigma_1^1 & 0 & 0 \\ \Sigma_2^1 + a(\Sigma_1^1 - \Sigma_2^2) & \Sigma_2^2 & 0 \\ \Sigma_3^1 + b(\Sigma_1^1 - \Sigma_3^3) & 0 & \Sigma_3^3 \end{pmatrix} \\ &= \begin{pmatrix} \Sigma_1^1 & 0 & 0 \\ \Sigma_2^1 + a(3\Sigma_+ + \sqrt{3}\Sigma_-) & \Sigma_2^2 & 0 \\ \Sigma_3^1 + b(3\Sigma_+ - \sqrt{3}\Sigma_-) & 0 & \Sigma_3^3 \end{pmatrix}. \end{aligned}$$

We see that choosing $a = -\frac{8}{3}\Sigma_2^1(\infty)$ and $b = -\frac{8}{3}\Sigma_3^1(\infty)$ the transformed variables $\tilde{\Sigma}_2^1, \tilde{\Sigma}_3^1$ are zero asymptotically. By direct calculation, one can see that the same is true for the transformed variables \tilde{N}_2^1 and \tilde{N}_3^1 . Thus, we obtain the same asymptotics as in the diagonal case and we can conclude:

Theorem 1: Consider any C^∞ solution of the Einstein-Vlasov system with Bianchi II symmetry and with C^∞ initial data. Assume that $|\Sigma_+(t_0) - \frac{1}{8}|, |\Sigma_-(t_0)|, |\Sigma_2^1(t_0)|, |\Sigma_3^1(t_0)|, |\Sigma_3^2(t_0)|, |\Sigma_2^3(t_0)|, |\Sigma_1^2(t_0)|, |\Sigma_1^3(t_0)|, |N_1(t_0) - \frac{3}{4}|, |N_2^1(t_0)|, |N_3^1(t_0)|$, and $P(t_0)$ are sufficiently small. Then at late times,

after possibly a basis change, the following estimates hold:

$$\begin{aligned}
 H(t) &= \frac{2}{3}t^{-1}(1 + O(t^{-\frac{1}{2}})), \\
 \Sigma_+ - \frac{1}{8} &= O(t^{-\frac{1}{2}}), \\
 \Sigma_- &= O(t^{-1}), \\
 \Sigma_2^1 &= O(t^{-\frac{1}{2}}), \\
 \Sigma_3^1 &= O(t^{-\frac{1}{2}}), \\
 \Sigma_3^2 &= O(t^{-1}), \\
 \Sigma_2^3 &= O(t^{-1}), \\
 \Sigma_1^2 &= O(t^{-1}), \\
 \Sigma_1^3 &= O(t^{-1}), \\
 N_1 - \frac{3}{4} &= O(t^{-\frac{1}{2}}), \\
 N_2^1 &= O(t^{-\frac{1}{2}}), \\
 N_3^1 &= O(t^{-\frac{1}{2}}), \\
 P(t) &= O(t^{-\frac{1}{2}}).
 \end{aligned}$$

7. Closing the bootstrap argument of Bianchi VI₀

It follows immediately by the same arguments as in the diagonal case:

$$|\Sigma_2^1| = O(t^{-1+\epsilon}), \tag{46}$$

$$|\Sigma_3^1| = O(t^{-1+\epsilon}), \tag{47}$$

$$|\Sigma_2^3 + \Sigma_3^2| = O(t^{-1+\epsilon}). \tag{48}$$

Now consider the $\Sigma_2^3 N_{23}$ system. Using the fact that $N_2^3 = -N_{23}(N_3 + N_2)$, we obtain

$$\begin{aligned}
 \dot{\Sigma}_2^3 &= H[-N_{23}(N_3 + N_2) - \Sigma_2^3(3 + \frac{\dot{H}}{H^2}) - \frac{8\pi S_2^3}{H^2}], \\
 \dot{N}_{23} &= H[(2\Sigma_+ + 2\sqrt{3}\Sigma_- + q)N_{23} + 2\Sigma_2^3 N_3 - 2\Sigma_2^1 N_{13}].
 \end{aligned}$$

Using the bootstrap assumptions, the estimates obtained and the variable τ

$$\begin{pmatrix} \Sigma_2^3 \\ N_{23} \end{pmatrix}' = \begin{pmatrix} -\frac{3}{2} + \epsilon_1 & \epsilon_2 \\ -\frac{3}{2} + \epsilon_3 & \epsilon_1 \end{pmatrix} \begin{pmatrix} \Sigma_2^3 \\ N_{23} \end{pmatrix} + O(\epsilon e^{(-\frac{3}{2}+\epsilon)\tau}) \begin{pmatrix} 1 \\ 1 \end{pmatrix},$$

where $\epsilon_1, \epsilon_2,$ and ϵ_3 have the following origin. The quantity ϵ_1 is determined essentially by the error in N and Σ_+ and note that Σ_2^3 comes in combination with Σ_3^2 , thus this term can be chosen as small as we want. The quantity ϵ_2 comes from $N_2 + N_3$ and can be determined by the error of N_2^2 and finally the quantity ϵ_3 which comes from N_3 depends on the error of $N, N_2^2,$ and N_{23}^2 . Note in the last term that the quantity is squared, thus it is negligible. Having a look at the linearization and going

to the eigenbasis via

$$\begin{pmatrix} \check{\Sigma}_2^3 \\ \check{N}_{23} \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} \Sigma_2^3 \\ N_{23} \end{pmatrix},$$

we come to the system

$$\begin{pmatrix} \check{\Sigma}_2^3 \\ \check{N}_{23} \end{pmatrix}' = \begin{pmatrix} -\frac{3}{2} + \epsilon_1 + \epsilon_2 & \epsilon_2 \\ \epsilon_3 - \epsilon_2 & \epsilon_1 - \epsilon_2 \end{pmatrix} \begin{pmatrix} \check{\Sigma}_2^3 \\ \check{N}_{23} \end{pmatrix} + O(\epsilon e^{(-\frac{3}{2} + \epsilon)\tau}) \begin{pmatrix} 1 \\ 1 \end{pmatrix}.$$

From which follows:

$$\begin{aligned} \check{\Sigma}_2^3 &= \check{\Sigma}_2^3(\tau_0) e^{(-\frac{3}{2} + \epsilon)\tau}, \\ \check{N}_{23} &= \check{N}_{23}(\tau_0) + O(\epsilon), \end{aligned}$$

and going back

$$\begin{aligned} \Sigma_2^3 &= \Sigma_2^3(\tau_0) e^{(-\frac{3}{2} + \epsilon)\tau}, \\ N_{23} &= N_{23}(\tau_0) - \Sigma_2^3(\tau_0) + O(\epsilon). \end{aligned}$$

We see that we have improved N_{23} , Σ_2^3 and with that also Σ_3^2 .

Using these estimates and the bootstrap assumptions, let us focus now on the following system:

$$\begin{aligned} \dot{\Sigma}_+ &= H\left[\frac{N}{3} + \Sigma_+(q - 2) + O(t^{-1+\epsilon})\right], \\ \dot{\Sigma}_- &= H\left[-\frac{N^2}{\sqrt{3}} + (q - 2)\Sigma_- + O(t^{-1+\epsilon})\right], \\ \dot{N} &= H[2(2\Sigma_+ + q)N - 4\sqrt{3}\Sigma_-N^2 + O(t^{-1+\epsilon})], \\ \dot{N}_2^2 &= H[2(2\Sigma_+ + q)N^2 + (\frac{9}{4}\sqrt{3} + O(\epsilon))\Sigma_- + O(t^{-1+\epsilon})], \end{aligned}$$

where in the last equation, $(N_2)^2 + (N_3)^2$ was estimated with N_2^2 , N , and N_{23} . The $O(\epsilon)$ -term will not play a role since it can be absorbed in the ϵ of the estimate. Let us look at the linearization using the variables $\check{\Sigma}_+ = \Sigma_+ + \frac{1}{4}$, $\check{\Sigma}_- = \Sigma_-$, $\check{N} = N + \frac{9}{8}$, $\check{N}_2^2 = N_2^2$ and the variable τ

$$\begin{pmatrix} \check{\Sigma}_+ \\ \check{N} \\ \check{\Sigma}_- \\ \check{N}_2^2 \end{pmatrix}' = \begin{pmatrix} -\frac{21}{16} & +\frac{5}{16} & 0 & 0 \\ -\frac{45}{16} & -\frac{3}{16} & 0 & 0 \\ 0 & 0 & -\frac{3}{2} & -\frac{\sqrt{3}}{3} \\ 0 & 0 & \frac{9}{4}\sqrt{3} & 0 \end{pmatrix} \begin{pmatrix} \check{\Sigma}_+ \\ \check{N} \\ \check{\Sigma}_- \\ \check{N}_2^2 \end{pmatrix}.$$

The eigenvalues are

$$\begin{aligned} \lambda_{1/2} &= -\frac{3}{4} \pm \frac{3}{4}i\sqrt{3}, \\ \lambda_{3/4} &= -\frac{3}{4} \pm \frac{3}{4}i. \end{aligned}$$

These eigenvalues are the same, which appeared in the reflection symmetric case. Using the same arguments, we arrive at

$$\begin{aligned}
 |\Sigma_+ + \frac{1}{4}| &\leq A_+(1+t)^{-\frac{1}{2}+\epsilon}, \\
 |\Sigma_-| &\leq A_-(1+t)^{-\frac{1}{2}+\epsilon}, \\
 |N + \frac{9}{8}| &\leq A_{c1}(1+t)^{-\frac{1}{2}+\epsilon}, \\
 |N_2^2| &\leq A_{c2}(1+t)^{-\frac{1}{2}+\epsilon}.
 \end{aligned}$$

Finally, only $\Sigma_1^2(\tau_0)$, $\Sigma_1^3(\tau_0)$, $N_{12}(\tau_0)$, and $N_{13}(\tau_0)$ have to be improved. Let us look at the Σ_1^2, N_{12} system. There is no difference between this system and the $\Sigma_1^3-N_{13}$ system.

$$\begin{pmatrix} \Sigma_1^2 \\ N_{12} \end{pmatrix}' = \begin{pmatrix} -\frac{3}{2} + \epsilon_1 & \frac{3}{2} + \epsilon_2 \\ -\frac{3}{2} + \epsilon_3 & \frac{3}{2} + \epsilon_1 \end{pmatrix} \begin{pmatrix} \Sigma_1^2 \\ N_{12} \end{pmatrix} + O(\epsilon) \begin{pmatrix} 1 \\ 1 \end{pmatrix},$$

$$\begin{aligned}
 \lambda_1 &= \frac{1}{2}(2\epsilon_1 - \sqrt{2\epsilon_2\epsilon_3 - 3\epsilon_2 + 3\epsilon_3}), \\
 \lambda_2 &= \frac{1}{2}(2\epsilon_1 + \sqrt{2\epsilon_2\epsilon_3 - 3\epsilon_2 + 3\epsilon_3}).
 \end{aligned}$$

Now choosing the error of N bigger than $\Sigma^{ab}\Sigma_{ab} \epsilon_1$ will be negative. ϵ_2 and ϵ_3 can be chosen in such a way that the square root of the term is positive but in total smaller than ϵ_1 , such that we have two small and different eigenvalues.

8. Arzela-Ascoli for Bianchi VI₀

For Bianchi VI₀, we can apply Arzela-Ascoli as well. We see that N_{23} will be zero and $\Sigma_1^2(\tau_0)$, $\Sigma_1^3(\tau_0)$, $N_{12}(\tau_0)$, and $N_{13}(\tau_0)$ will be come constants. This time we can make the following basis change, which preserves the Lie-algebra to obtain that the mentioned variables tend to zero

$$\begin{aligned}
 \tilde{e}_1 &= e_1 + ae_2 + be_3, \\
 \tilde{e}_2 &= e_2, \\
 \tilde{e}_3 &= e_3.
 \end{aligned}$$

We can conclude

Theorem 2: Consider any C^∞ solution of the Einstein-Vlasov system with Bianchi VI₀ symmetry and with C^∞ initial data. Assume that $|\Sigma_+(t_0) + \frac{1}{4}|$, $|\Sigma_-(t_0)|$, $|\Sigma_2^1(t_0)|$, $|\Sigma_3^1(t_0)|$, $|\Sigma_3^2(t_0)|$, $|\Sigma_2^3(t_0)|$, $|\Sigma_1^2(t_0)|$, $|\Sigma_1^3(t_0)|$, $|N(t_0) + \frac{9}{8}|$, $|N_2^2(t_0)|$, $|\Sigma_1^2(t_0)|$, $|N_{12}(t_0)|$, $|N_{13}(t_0)|$, and $P(t_0)$ are sufficiently small.

Then at late times, after possibly a basis change, the following estimates hold:

$$\begin{aligned}
 H(t) &= \frac{2}{3}t^{-1}(1 + O(t^{-\frac{1}{2}})), \\
 \Sigma_+ - \frac{1}{4} &= O(t^{-\frac{1}{2}}), \\
 \Sigma_- &= O(t^{-\frac{1}{2}}), \\
 \Sigma_2^1 &= O(t^{-1}), \\
 \Sigma_3^1 &= O(t^{-1}), \\
 \Sigma_3^2 &= O(t^{-1}), \\
 \Sigma_2^3 &= O(t^{-1}), \\
 \Sigma_1^3 &= O(t^{-\frac{1}{2}}), \\
 \Sigma_1^2 &= O(t^{-\frac{1}{2}}), \\
 N_{12} &= O(t^{-\frac{1}{2}}), \\
 N_{13} &= O(t^{-\frac{1}{2}}), \\
 N_{23} &= O(t^{-\frac{1}{2}}), \\
 N_2^2 &= O(t^{-\frac{1}{2}}), \\
 N + \frac{9}{8} &= O(t^{-\frac{1}{2}}), \\
 P(t) &= O(t^{-\frac{1}{2}}).
 \end{aligned}$$

IV. CONCLUSIONS

As mentioned in the abstract, the challenge here was to put the equations in a form such that the results of the diagonal case can be used. This can be seen especially in the curvature variables. For Bianchi II, it was sufficient to use the new variables $N_j^i = \frac{R_j^i}{H^2}$. For Bianchi VI₀, we had to introduce in addition to that the new variables $N_{ij} = \frac{g_{ij}}{g\sqrt{H}}$. The notation might be a little bit confusing, but in both cases these variables have a connection to the curvature variables N_1, N_2 , and N_3 of the diagonal case and this is the reason for the notation. In contrast to the diagonal case where the treatment of Bianchi II and VI₀ was almost identical, here the latter case was more difficult. One reason could be the obvious increase in complexity. In the Bianchi II case, it was sufficient to deal with N instead of N_1 and look at the differences. In the case of Bianchi VI₀, N had to be used to start the bootstrap argument. Then also N_2^2 and N_{23} . This last variable made the correspondence to the diagonal case more difficult. As can be seen in the chapter where the bootstrap argument was closed for Bianchi VI₀, we had to look more carefully on the dependence of the different ϵ . Note also that we did not use exactly the linearization in our last improvement of the estimates. We would have obtained that zero is a multiple eigenvalue and we would have not obtained decay, but logarithmic growth. This would have been sufficient to close the bootstrap argument with corresponding suitable bootstrap assumptions, but there would exist difficulties to apply the Arzela-Ascoli theorem and to obtain that the non-diagonal components become constant. Another difference to the diagonal case is the use of a basis change in the end. In general, the non-diagonal components will become constants and thus not relevant. However, to obtain “diagonal” asymptotics, a basis change will in general be necessary.

It would be interesting to investigate whether the work on homogeneous Ricci solitons³ can help to understand the similarities and differences between Bianchi II and VI₀ (in Thurston's classification Nil and Sol).

We have discussed the future asymptotics of some Bianchi models; however, what about the higher types? The case of Bianchi VII₀ will probably be quite different. For instance in Ref. 13, it was discovered that the Bianchi VII₀ spacetimes with a non-tilted fluid are not asymptotically self-similar in the future and that some oscillations take place. It is shown that dynamics are dominated by the Weyl curvature. However for dust, a bifurcation of the Weyl curvature takes place (Theorem 2.4 of Ref. 13 and comments below). For this reason, it is likely to expect difficulties when applying our techniques to this case. Something similar, but even more complicated happens in the case of Bianchi VIII spacetimes with a non-tilted fluid.⁶

What about inhomogeneous models? Some direction to generalize our results could be to analyze the Gowdy model, which is the simplest inhomogeneous case. In Ref. 11, different links between Bianchi and (twisted) Gowdy spacetimes are considered, in particular for Bianchi I, II, VI₀, and VII₀. The analysis of perturbations is another interesting approach towards the understanding of inhomogeneous models (see Ref. 1 for recent developments).

- ¹ P. T. Allen and A. D. Rendall, "Asymptotics of linearized cosmological perturbations," *J. Hyp. Differ. Equations* **7**, 255–277 (2010).
- ² J. D. Barrow and S. Hervik, "The future of tilted Bianchi universes," *Class. Quantum Grav.* **20**, 2841–2854 (2003).
- ³ C. Guenther, J. Isenberg, and D. Knopf, "Linear stability of homogeneous Ricci solitons," *Int. Math. Res. Notices* **2006**, 96253.
- ⁴ S. Hervik, "The asymptotic behaviour of tilted Bianchi type VI₀ universes," *Class. Quantum Grav.* **21**, 2301 (2004).
- ⁵ C. G. Hewitt, R. Bridson, and J. Wainwright, "The asymptotic regimes of tilted Bianchi II cosmologies," *Gen. Relativ. Gravit.* **33**, 65–94 (2001).
- ⁶ J. T. Horwood, M. J. Hancock, D. The, and J. Wainwright, "Late-time asymptotic dynamics of Bianchi VIII cosmologies," *Class. Quantum Grav.* **20**, 1757–1778 (2003).
- ⁷ E. Nungesser, "Future non-linear stability for reflection symmetric solutions of the Einstein-Vlasov system of Bianchi types II and VI₀," *Annales Henri Poincaré* (2012).
- ⁸ A. D. Rendall, "Cosmic censorship for some spatially homogeneous cosmological models," *Ann. Phys.* **233**, 82–96 (1994).
- ⁹ A. D. Rendall, "Global properties of locally spatially homogeneous cosmological models with matter," *Math. Proc. Cambridge Philos. Soc.* **118**, 511–526 (1995).
- ¹⁰ A. D. Rendall, *Partial Differential Equations in General Relativity* (Oxford University Press, Oxford, 2008).
- ¹¹ A. D. Rendall, "Dynamics of solutions of the Einstein equations with twisted Gowdy symmetry," *J. Geom. Phys.* **62**, 569–577 (2012).
- ¹² J. M. Stewart, *Non-equilibrium Relativistic Kinetic Theory*, Lecture Notes in Physics Vol. 10 (Springer, Berlin, 1971).
- ¹³ J. Wainwright, M. J. Hancock, and C. Uggla, "Asymptotic self-similarity breaking at late times in cosmology," *Class. Quantum Grav.* **16**, 2577–2598 (1999).